RESEARCH ARTICLE

Title: Restoration and resilience to sea level rise of a salt marsh affected by dieback events in

Charleston, SC

Authors: Rolando, JL[a]; Hodges, M[b]; Garcia, KD[a]; Krueger, G[a]; Williams, N[c]; Carr Jr., J[d];

Robinson; J[c,e]; George, A[f]; Morris; J[g]; Kostka, JE[a,h,i]

Affiliations:

[a]Georgia Institute of Technology, School of Biological Sciences, Atlanta, GA 30332, USA

[b]South Carolina Department of Natural Resources, Charleston, SC 29412, USA

[c]Robinson Design Engineers, Charleston, SC 29407, USA

[d]Ashleyville Neighborhood Association, Charleston, SC 29407, USA

[e]College of Charleston, Marine Biology Graduate Program, Charleston, SC 29424, USA

[f]South Carolina Aquarium, Conservation Department, Charleston, SC 29401, USA

[g]University of South Carolina, Belle Baruch Institute for Marine & Coastal Sciences, Columbia,

SC 29208, USA

[h]Georgia Institute of Technology, School of Earth and Atmospheric Sciences, Atlanta, GA 30332,

USA

[i]Center for Microbial Dynamics and Infection, Georgia Institute of Technology, Atlanta, GA

30332, USA

Corresponding author:

Kostka, JE. e-mail: joel.kostka@biology.gatech.edu

Abstract

The frequency of salt marsh dieback events has increased over the last 25 years with unknown consequences to the resilience of the ecosystem to accelerated sea level rise (SLR). Salt marsh ecosystems impacted by sudden vegetation dieback events were previously thought to recover naturally within a few months to years. In this study, we provide evidence that approximately 14% of total marsh area has not revegetated 10-years after a dieback event in Charleston, SC. Dieback onset coincided with a severe drought in 2012, and a second dieback event occurred in 2016 after a historic flood influenced by Hurricane Joaquin in October of 2015, with unvegetated zones reaching nearly 30% of total marsh area in 2017. Most affected areas were associated with lower elevation zones in the interior of the marsh (midmarsh). During the 2013 dieback event, we estimate that unvegetated midmarsh area expanded by 300%. Grass planting was shown to be an effective restoration practice, with restored plants having greater aboveground biomass than relict sites after two years of transplanting. A positive restoration outcome indicated that the stressors that caused the initial dieback are no longer present. Despite that, many dieback areas have not recovered naturally even as they are located within the typical elevation range of a healthy vegetated marsh. A mechanistic modelling approach was used to assess the effects of vegetation dieback on salt marsh resilience to SLR. Predictions indicate that a highly productive restored marsh (2000 g m^{-2} y^{-1}) would persist at a moderate SLR rate of 60 cm 100 y^{-1}, whereas a non-restored mudflat would lose all of its elevation capital after 100 years. Thus, rapid restoration of marsh dieback is critical to avoid further degradation. Also, failure to incorporate the increasing frequency and intensity of extreme climatic events which trigger irreversible marsh diebacks underestimates salt marsh vulnerability to climate change. At an elevated SLR rate of 122 cm 100 y^{-1}, most likely an extreme climate change scenario, even highly productive ecosystems augmented

47 by sediment placement would not keep pace with SLR. Thus, climate change mitigation actions

48 are also urgently needed to preserve present-day marsh ecosystems.

49 Keywords: salt marsh, dieback, drought, elevation, resilience, sea level rise, urban ecosystem

1. Introduction:

The resilience of salt marsh ecosystems to climate change has been a source of intense scientific

debate, often focused on the response of marshes to accelerated sea level rise (SLR) (Donnelly and

Bertness, 2001; Kirwan et al., 2010; Kirwan et al., 2016a; Kirwan and Megonigal, 2013; Morris

et al., 2002; Langston et al., 2021; Raposa et al., 2017; Rogers et al., 2019; Schuerch et al., 2018;

Weston, 2014). Ultimately, the capacity of present-day marshes to persist under accelerated SLR

will depend on the ability to increase in surface elevation at a rate that at least matches that of SLR

(Morris et al., 2002; Reed, 1995). Salt marsh surface elevation is determined by the balance

between sediment deposition and erosion, organic matter sequestration, and land subsidence

(Kirwan and Megonigal, 2013; Morris et al., 2016; Mudd et al., 2009). Plant productivity is a key

variable modulating marsh vertical accretion not only by accruing organic matter from

belowground biomass, but also by facilitating the deposition of suspended matter during tidal

inundation (Baustian et al., 2012; Mudd et al., 2009, 2010). Moreover, relative marsh elevation or

elevation capital (Cahoon and Guntenspergen, 2010), the distance of the wetland surface to the

lowest elevation at which plants can survive, largely determines the vulnerability to SLR (Morris

et al., 2021).

In the last 25 years, sudden salt marsh dieback events in which extensive areas of vegetation die-

off in short periods of time have been increasing in frequency with little understanding of the

implications for the long-term resilience of marshes to SLR (Alber et al., 2008; McKee et al.,

2004). Sudden salt marsh dieback events have been associated with drought, and suggested to be

caused by the interaction of multiple physical, chemical, and biological stressors in the Southeast

and Gulf of Mexico US (Alber et al., 2008). Specifically, elevated porewater salinity, acidification

of air-exposed sediment, fungal infections, and increased snail herbivory have all been reported in

previous events (Alber et al., 2008; Hughes et al., 2012; McKee et al., 2004; Silliman et al., 2005).

In many cases, areas affected by sudden salt marsh dieback have naturally recovered within a few

months to years (Alber et al., 2008; Lindstedt et al., 2006; Linthurst and Seneca, 1980; Marsh et

al., 2016). However, there are also reports of marshes that have not recovered to date (Alber et al.,

2008; Baustian et al., 2012; Marsh et al., 2016), with most of these cases representing low elevation

zones that experienced an irreversible regime shift to ponds, requiring active restoration for

recovery (Baustian et al., 2012; McKee et al., 2004; Ogburn and Alber, 2006; Schepers et al.,

2020). In contrast, little is known about the causes and fate of dieback events causing long-term

vegetation loss in areas that transitioned from salt marsh to mudflat, as well as their vulnerability

to accelerated SLR. Furthermore, to the best of our knowledge there are limited studies

characterizing the trajectory and recovery of salt marsh dieback events at decadal timescales (e.g.,

Marsh et al., 2016).

At a global scale, the area of marshland that is projected to drown as a consequence of accelerated

SLR has been suggested to be offset by salt marsh migration onto higher land (Kirwan et al.,

2016a). Nevertheless, even if that holds true, urban salt marshes are highly vulnerable to

accelerated SLR because built infrastructure associated with coastal cities acts as a physical barrier

preventing marsh migration (Kirwan and Gedan, 2019; Schuerch et al., 2018). If no actions are

taken to conserve present-day urban salt marshes, the ecosystem services they provide will be lost.

For example, salt marshes provide a natural defense to storm surge for coastal populations and

cities (Barbier et al., 2011; Temmerman et al., 2013). Up to 41% of the world's population lives

on the coastline, with 630 million people living below projected flood levels for 2100, and lower

income populations from underrepresented minority communities are more likely to live in flood

prone zones (Kulp and Strauss, 2019; Martínez et al., 2006; Chakraborty et al., 2014; Eisenman et

al., 2007). Conservation of urban marshes represents an effective climate change adaptation

strategy with increasing relevance over time, as the frequency and intensity of tropical storms and

hurricanes are already increasing (Brown et al., 2019; Paerl et al., 2019). To the best of our

knowledge, little information is available on the vulnerability or resilience of urban salt marshes

to accelerated SLR. Thus, there is great need to investigate the potential response of urban marshes

to various climate change, ecological, and management scenarios in order to support adaptive

management of the coastline.

This study was initiated to assess the vulnerability and restoration potential of salt marshes in

Charleston, SC, motivated by members of the Ashleyville Neighborhood Association in West

Ashley, a densely populated low-lying area that borders on urban marshes. Salt marshes in the area

serve as essential habitat for many fish and shellfish species (e.g., red drum, sharks, oysters), and

the National Fish and Wildlife Foundation (NFWF)-funded Coastal Resilience Assessment

prioritized the West Ashley area for restoration because its habitats are generally in relatively poor

condition, wildlife is exposed to high stress, and human assets show a very high vulnerability to

storms along with SLR (Crist et al. 2019). Thus, we used this case study to expand the current

knowledge on the trajectory of salt marsh dieback events and their effects on salt marsh resilience

to climate change-induced accelerated rate of SLR. Objectives of this study were to: i. characterize

the onset, dynamic, and trajectory of salt marsh dieback events in Charleston, SC ii. correlate the

onset of dieback with climatological and salt marsh geophysical properties, iii. evaluate the

effectiveness of salt marsh restoration by grass planting in an unvegetated marsh affected by a

dieback event, and finally iv. model the resilience of dieback affected salt marshes to SLR under

contrasting restoration scenarios.

2. Materials and methods

2.1 <u>Study Area</u>

The study was conducted in *Spartina alterniflora*-dominated salt marshes located in West Ashley,

Charleston, SC. The watershed west of the Ashley River has great cultural and historical relevance

since it contains Maryville and Charles Towne Landing. Maryville and Charles Towne Landing

are the sites of a Gullah Geechee community established after the American Civil War and first

permanent English settlement in the Carolinas in 1670, respectively. The dieback area was first

discovered by members of the local Gullah Geechee community in a salt marsh adjacent to

Maryville (Fig. S1).

In order to relate the expansion of dieback areas to drought and extreme precipitation, we retrieved

monthly Palmer drought stress index (PDSI) values from the South Carolina Southern Division

(January 2009 to August 2021) using NOAA´s National Centers for Environmental Information

database (https://www.ncdc.noaa.gov/temp-and-precip/drought/historical-palmers/), and monthly

precipitation values from the Charleston area, SC (ThreadEx) using NOAA's Online Weather Data

(NOWData, https://www.weather.gov/wrh/Climate?wfo=chs). The PDSI is a widely used drought

index based on the supply and demand of a water balance equation of precipitation,

evapotranspiration, soil moisture loss and recharge, and runoff, which has been used to study the

effects of water limitation on agricultural and ecological systems (Palmer, 1965).

2.2 <u>Characterization of marsh dieback trajectories using multispectral high-resolution imagery</u>

Multispectral imagery was acquired from three sources, satellite imagery from the PlanetScope

constellation (Planet Team, 2017), and MAXAR's WorldView-3 satellite, and aerial imagery from

US Department of Agriculture's (USDA) National Agriculture Imagery Program (NAIP). The

NAIP dataset consists of 4-band [blue, green, red, and near infrared (NIR)], 1-meter resolution

imagery   (http://www.fsa.usda.gov/programs-and-services/aerial-photography/imagery-

programs/naip-imagery). The PlanetScope constellation acquires daily, 3-meter resolution

imagery, using a 4-band (blue, green, red, NIR) sensor. We used the Planet's Surface Reflectance

product, processed to top of atmosphere reflectance, and atmospherically corrected to surface

reflectance (Planet Team, 2019). PlanetScope images were acquired during low tide, and on

cloudless days. The MAXAR WorldView-3 acquires 4-band (blue, green, red, NIR), 30-cm

resolution images. The used View-Ready WorldView-3 image was retrieved during low tide on a

cloudless day, was radiometrically and sensor corrected, and atmospherically corrected to surface

reflectance (MAXAR, 2021). The complete imagery used in this study consisted of 9 images

retrieved from 2009 to 2021 (further details on Table S1).

Two vegetation indices were calculated from all acquired images: the normalized difference

vegetation index (NDVI), and the modified soil-adjusted vegetation index 2 (MSAVI2):

153
$$NDVI = \frac{(NIR-Red)}{(NIR+Red)} \qquad (1)$$

154
$$MSAVI2 = \frac{2NIR+1-\sqrt{(2NIR+1)^2-8(NIR-Red)}}{2} \qquad (2)$$

Total unvegetated area was estimated for all NAIP images, and the 2016 WorldView-3, and 2021

PlanetScope satellite images. We visually inspected the coordinates of vegetated and unvegetated

points (median observations per year: 218) using high-resolution images at a scale of 1:500 to

1:1000 in QGIS 3.16.5 (QGIS.org, 2021). Raster values of the four bands and NDVI were extracted

from the pixels containing the vegetated and unvegetated points. Random forest models were used

to estimate the probability that a pixel was unvegetated using the four bands and NDVI as

explanatory variables. Out-of-bag estimate of error rate was lower than 3.5% for all models. Pixels

with a greater than 75% probability of being unvegetated were defined as such, while pixels with

a lower than 25% likelihood of being unvegetated were defined as vegetated areas. This analysis

was performed using the raster, and randomForest R packages in the R v. 4.1.0 environment

(Hijmans and van Etten, 2012; Liaw and Wiener, 2002; R Core Team, 2021).

In order to investigate if unvegetated areas were preferentially associated with creek bank or

midmarsh zones, we drew all creek bank edges from the study area (Fig. S1). Edges were drawn

using the 2019 NAIP aerial image and the corrected 2017 LIDAR derived DEM at a scale of 1:250

to 1:500 in QGIS 3.16.5 (QGIS.org, 2021). We generated a raster measuring the distance of each

pixel to the closest creek bank edge using the "Join attributes by nearest" algorithm in QGIS with

a resolution of 1 m. Based on the well-known natural zonation of *S. alterniflora* (Mendelssohn and

Morris, 2002), with taller plants found closer to tidal creeks, areas within 7.5 m of a tidal creek

were defined as "creek bank marsh", whereas areas > 7.5 m distal to creeks were defined as

"midmarsh" (Valiela et al., 1978). The 7.5 m threshold characterizing the ecosystem state change

was defined by visually inspecting an aboveground biomass vs. distance from tidal creek biplot

(Fig. S2, method for aboveground biomass estimation below).

2.3 <u>Field sampling</u>

Field sampling was performed during spring and summer of 2021. Two field campaigns were

performed on 27-28 May, and 20-26 July of 2021. We sampled 60, and 119 locations, in May and

July, respectively (total = 179 points). Field sampling was performed non-randomly by

establishing transects across gradients of *S. alterniflora* biomass from the creek bank to the interior

of the marsh (Fig. S1). The dieback area adjacent to Maryville was oversampled due to our interest

as a potential restoration site. At all locations (n = 179), a high-accuracy real time kinematic (RTK)

survey was conducted using a Trimble GNSS GPS receiver (Trimble, Sunnyvale, CA, USA).

Elevation was expressed in meters in the North American Vertical Datum of 1988 (NAVD88)

GEOID 18. Instrument accuracy is of 2 cm root mean square error (RMSE), horizontal and vertical.

During the July field sampling, *S. alterniflora* shoot density and height were measured using a

0.25 m^2 quadrat at 82 points sampled in parallel for RTK surface elevation. Shoot density was

measured by counting all *S. alterniflora* stems inside the 0.25 m^2 quadrat. Shoot height was

measured randomly from 6 stems per quadrat. In order to estimate aboveground biomass, 71 *S.*

alterniflora shoots of known height were harvested from 37 sampling points across contrasting *S.*

alterniflora growth zones. Harvested shoots were oven-dried at 60 ºC for one week and their shoot

height and dry weight were used to construct a biomass allometric equation. For all sampling

points, biomass was calculated for individual shoots using the constructed allometric equation, and

mean shoot biomass (n = 6) multiplied by shoot density to calculate aboveground biomass at the

plot level (g m^{-2}).

2.4 Acquisition of LIDAR derived DEM

A light detection and ranging (LIDAR) derived bare-earth digital elevation model (DEM)

constructed from LIDAR aerial acquisition conducted from February 25, 2017 to March 09, 2017

in Charleston County was used in this study (South Carolina DNR, 2018). LIDAR acquisition

involved Axis GeoSpatial (Easton, MD, USA) operating a Cessna 206H outfitted with a Riegl

LMS-Q1560 dual channel laser scanner system at 2043 m above ground level. Scanner Pulse Rate

was 800 kHz with unlimited number of returns per pulse. The DEM vertical datum used was

NAVD88 Geoid 12B and was projected to the NAD83 (2011) South Carolina State Plane at a pixel

resolution of 1 m. System parameters for LIDAR acquisition are described in Table S2 (further

details can be found in South Carolina DNR, 2018).

2.5 Correction of the LIDAR derived DEM

High-accuracy RTK elevations were used to correct the bare-earth LIDAR derived DEM. Error of

the LIDAR derived DEM was defined as: Z_{LIDAR} - Z_{RTK} for each RTK observation, where Z is

NAVD88 elevation. We used a multispectral PlanetScope satellite image retrieved 5 days before

LIDAR acquisition (02/20/2017) to correct the DEM based on the known relation between LIDAR

error and *S. alterniflora* height (Hladik and Alber, 2012). The 179 RTK observations were

randomly split for model training and testing in a 7 to 3 ratio, respectively. Error of the LIDAR

derived DEM was fitted to a multiple lineal regression model using automated model selection

based on Akaike Information Criterion (AIC) as implemented in the glmulti R function (method =

"h") (Calcagno and de Mazancourt, 2010). We used the four bands of the PlanetScope image (Red,

Green, Blue, and NIR), two vegetation indices (NDVI, and MSAVI2), and the original LIDAR

derived DEM elevations as potential explanatory variables. The DEM was corrected by solving

the multiple linear regression for the RTK observed elevation. Performance of the best model was

tested based on root mean square error (RMSE) calculated as: $RMSE =$

$\sqrt{\sum(Observed_i - Predicted_i)^2/n}$, where *Observed$_i$* is the ith observation of the RTK elevation,

and *Predicted$_i$* is the ith observation of the LIDAR derived DEM (Maune et al., 2007). We

calculated the RMSE for the test and train datasets of both the corrected and uncorrected LIDAR

derived DEM.

2.6 Estimation of 2021 aboveground biomass

Aboveground biomass was estimated for the total study area based on *in situ* biomass observations

made during the July 2021 field campaign. Aboveground biomass (g m^{-2}) estimated from 82

sampling points (section 2.3) were randomly split into train and test datasets (80% and 20%,

respectively). A random forest model was constructed using the corrected 2017 LIDAR derived

DEM, MSAVI2, NDVI, and the green, red and NIR bands from a PlanetScope image (acquired in

08/01/2021) as explanatory variables. The model grew 5000 trees; and in each split, 2 variables

were randomly sampled as candidates. The performance of the model was assessed by calculating

the RMSE of the train and test datasets (as in section 2.5). A raster containing the predicted

aboveground biomass for the whole study area was constructed using the random forest model.

Aboveground biomass was set to 0 in all unvegetated areas (as inferred in section 2.2). The model

was built using the raster and randomForest R packages in the R v. 4.1.0 environment (Hijmans

and van Etten, 2012; Liaw and Wiener, 2002; R Core Team, 2021).

2.7 Pilot grass planting restoration

A pilot restoration project led by the South Carolina Department of Natural Resources (SC-DNR)

was initiated in the Maryville marsh in 2019 in order to test the effectiveness of marsh restoration

in long-term affected ecosystems impacted by dieback events. The restoration effort consisted of

S. alterniflora planting using seedlings grown from local sources. Botanical seeds were annually

collected in October thru December each year (2019, 2020, and 2021) from 11 unique sites, all

located in Charleston County. After collection, seeds were kept at 4ºC for eight weeks, after which

they were germinated in planting trays filled with fresh water substrate. After 6-7 months,

seedlings ~55.0 cm tall were transplanted into the dieback area from May to August. When

transplanting, a spacing of 30.5 cm was used. Restoration was conducted in 6 individual patches

with dimensions ranging from 48.6 m^2 to 183.3 m^2. Plots from transplants performed in 2019,

2020 and 2021 were studied. In order to assess the effectiveness of the pilot restoration project,

aboveground biomass was estimated during the July field campaign in 5 and 12 quadrats (0.25-

m^2) from restored plots planted in 2020 and 2021, respectively. In order to have a complete and

balanced statistical design at the end of the growing season, a second field assessment was

performed in September 22nd 2021. In this assessment, 10 quadrats per plot were sampled in plots

planted in 2019, 2020, and 2021, and two patches of adjacent relict marsh. Aboveground biomass

was calculated based on shoot height and density as previously described (section 2.3). A one-way

ANOVA with the restored/natural plots as a fixed factor was used to explain differences in *S.*

alterniflora aboveground biomass for both the July and September assessments. In case of finding

statistical significance ($p < 0.05$), a Tukey's post-hoc test using the *emmeans* R package was

performed (Lenth, 2016).

2.8 Modeling of marsh elevation under primary production, management, and sea level rise

scenarios

The Marsh Equilibrium Model (MEM) v. 9.01 (Morris et al., 2002, 2021) was employed to assess

the resilience of the Charleston salt marsh to accelerated SLR under different scenarios of (i) *S.*

alterniflora primary productivity, (ii) human intervention (none or thin layer placement), and (iii)

rates of SLR (regional NOAA's intermediate low of 60 cm in 100 y^{-1} and intermediate of 122 cm

in 100 y^{-1}). The rationale to utilize the MEM was to assess the long-term effect (100 y) of marsh

vegetation loss with and without restoration in Charleston, SC. The MEM is a one-dimensional

mechanistic model that projects change in marsh elevation with SLR as a function of primary

production and sediment accretion. It was modified to simulate thin-layer placement (TLP) of

mineral sediment by allowing the user to specify periodic sediment additions to a specified

thickness, net of compaction. The model was coded in VBA behind an Excel user interface.

We simulated two rates of SLR, based on regional NOAA's intermediate low (60 cm in 100 years),

and intermediate (122 cm in 100 years) projections by 2100 (Sweet et al., 2017). The intermediate

low scenario has a 73% and 96% probability of being exceeded in 2100 under RCP4.5 and RCP8.5

scenarios, respectively; while the intermediate scenario has a 3% and 17% probability of being

exceeded by 2100 under RCP4.5 and RCP8.5 scenarios, respectively (Sweet et al., 2017). Starting

sea-level rise rate was set as 3.4 mm y^{-1}, as calculated for the Charleston Harbor by Morris and

Renken (2020). The elevation range of marsh vegetation was assumed to be within 25 cm below

mean sea level (MSL), and 30 cm above mean high water (MHW), as characterized in South

Carolina salt marshes (Morris et al., 2013). Tidal range, MSL and MHW during the study period

(Jan 2009 to August 2021) was 76 cm, 4 cm NAVD88, and 80 cm NAVD88, respectively. Optimal

elevation was determined as 5 cm above MSL based on a biplot of elevation and aboveground

biomass from the study area (Fig. 4f). Four different optimal values of aboveground primary

productivity were modelled: 0, 500, 1000, and 2000 g m^{-2} y^{-1} in order to assess the effects of plant

productivity and restoration by grass planting on marsh resilience to SLR. The median elevation

of all dieback pixels in 2021 was set as the initial elevation: 0.478 m NAVD88. Total suspended

matter (TSM) was estimated as 23 mg L^{-1} from remotely sensed GlobColour OLCIA imagery

(http://hermes.acri.fr/) as the average of monthly values from 2016 to 2021. TSM was extracted

from the closest pixels to the southern border of our study area in the Charleston Harbor. Finally,

a treatment with 5 cm of thin layer placement (TLP) every 25 years was included into the study to

simulate a more aggressive restoration scenario. Thin layer placement consists of spreading

sediment onto the marsh platform to increase surface elevation (Ford et al., 1999). After TLP,

marsh vegetation was simulated to be completely lost, and fully recovered 10 years after

disturbance.

3. Results

3.1 Salt marsh dieback trajectory

We observed an expansion of large unvegetated areas in 2013 that coincided with a severe drought in 2011-2012 (Fig. 1). Charleston, SC, experienced its most severe drought over the last 100 years in 2011-2012, concurrent with an over 200 % increase of unvegetated marsh area (64 ha, Fig. 2a). The South Carolina Southern NOAA division, which contains the city of Charleston, experienced an extreme drought (PDSI < -4.0) from June of 2011 to April of 2012 (Fig. 1). A second large expansion of unvegetated area was observed in 2016-2017 which was not drought-induced. The onset of the 2016-2017 dieback event coincided with the historic flood of the city of Charleston caused by record precipitation associated with the tropical cyclone Joaquin in October 2015 (Fig. 1). In 2019, the vegetated area of West Ashley had recovered up to 2013 levels which have persisted since. Based on field inspection and aerial imagery, we found that the affected salt marsh ecosystem transitioned from vegetated salt marsh to mudflat after both dieback events.

The dieback of 2013 caused a 300% expansion of unvegetated area in the interior of the marsh (Fig. 2b). In the 2016 event, the midmarsh and creekbank experienced a 50% and 33% increase of unvegetated area, respectively (Fig. 2b,c). Moreover, from 2011 to 2017 there was a constant loss of about 50% of creekbank salt marsh every two years (Fig. 2c). A LIDAR derived DEM acquired on 2017 was corrected for the studied area, improving the error estimation by approximately 30% on both training and testing datasets (RMSE after correction: less or equal to 8 cm in both train/test datasets, more detailed information in Supplementary Text S1, and Fig. S3). Using the corrected 2017 LIDAR derived DEM, we observed that both creek bank and midmarsh unvegetated areas were associated with lower elevation terrain (Fig. 3). The difference in median elevations between vegetated and unvegetated areas ranged from 13.3 cm to 27.4 cm, and 6.3 cm to 14.3 cm in creek

bank and midmarsh zones across years, respectively. Since the difference in elevation could have

been caused by land subsidence after the dieback event, we repeated the analysis with an

uncalibrated 2009 LIDAR derived DEM (South Carolina DNR, 2009), finding similar results (Fig.

S4). In the 2009 DEM, difference between median elevation from vegetated and unvegetated areas

ranged from 4.4 to 23.0 cm, and 2.4 to 16.3 cm in creek bank and midmarsh zones, respectively.

The results suggest that lower elevation persisted in the unvegetated areas prior to the dieback

event.

3.2 *Spartina alterniflora* primary production

A site-specific allometric equation was constructed to quantify the primary production of the

Charleston salt marsh during July of 2021 (Fig. 4a,c). The random forest model used to estimate

plant aboveground biomass was based on the 2017 calibrated LIDAR derived DEM, vegetation

indices, and satellite bands from an August 2021 color near infrared PlanetScope image. The

RMSE$_{test}$ of the model was 268.1 g m^{-2}. Median predicted aboveground biomass was 623.1 g m^{-2},

ranging from 0 to 1944 g m^{-2} (Fig. 4b). Biomass was within the expected range for Charleston's

latitude (predicted end of season productivity based on Kirwan et al., 2009: 574.8 g m^{-2} y^{-1}).

Aboveground biomass was closely associated with elevation and distance to tidal creeks (Fig. 4e,f,

Fig. S2), with greater aboveground biomass observed in lower elevation areas within closer

proximity to tidal creeks (Fig. 4e,f).

3.3 Grass transplanting is a successful restoration method in West Ashley, SC

Guided by the South Carolina Department of Natural Resources, seedlings from botanical seeds

of *S. alterniflora* were collected in Charleston, SC, propagated under greenhouse conditions, and

transplanted by community volunteers in July of 2019, 2020, and 2021 in a West Ashley dieback

area that has been unvegetated since 2013 (Fig. 5, S1). Aboveground biomass was determined

during the July field trip for the 2020 transplanted plot compared to reference plots of natural

midmarsh grass in the study area (Fig. 5). Aboveground biomass of the naturally vegetated

midmarsh and the 2020 transplanted plot were non-statistically different and had an average $\pm$ S.E.

of 527 ± 43 g m^{-2}, and 700 ± 185 g m^{-2}, respectively. Repeated measurements conducted in

September showed that the restored plots planted in 2019 had greater aboveground biomass in

comparison to reference relict marshes adjacent to the restoration site (Fig. 5). The plot

transplanted in 2020 was non-statistically different from reference relict marshes (Fig. 5). Average

$\pm$ S.E. aboveground biomass was 1726 ± 207 g m^{-2}, and 1290 ± 168 g m^{-2} for the 2019 and 2020

transplanted plots, respectively.

3.4 <u>Salt marsh resilience to sea level rise under contrasting restoration scenarios.</u>

The model output indicated that mudflat areas (*S. alterniflora* primary productivity = 0 g m^{-2} y^{-1})

rapidly lose elevation capital at a SLR rate of 60 cm in 100 y; decreasing in elevation by a total of

38 cm after the 100 years (Fig. 6). Restored salt marsh ecosystems with an optimal primary

productivity of 500 g m^{-2} y^{-1}, 1000 g m^{-2} y^{-1}, and 2000 g m^{-2} y^{-1} would persist under the 60 cm in

100 y SLR scenario, but losing 37.4 cm, 33.0 cm, and 24.5 cm of elevation capital after 100 years,

respectively (Fig. 6). At a SLR rate of 122 cm in 100 y, even a healthy restored salt marsh with an

optimal primary productivity of 2000 g m^{-2} y^{-1} would collapse in less than 75 years (Fig. 6).

Simulating an external supply of sediment (i.e., thin layer placement) at a rate of 5 cm of inorganic

sediment every 25 years would maintain the elevation capital of all simulated salt marsh

ecosystems in the 60 cm in 100 y SLR scenario. However, thin layer placement of a highly

productive salt marsh (2000 g m^{-2} y^{-1}) under the extreme 122 cm in 100 y SLR scenario would

result in the persistence of the ecosystem for only 100 years (Fig. 6).

4. Discussion

4.1 The resilience of salt marshes to accelerated sea level rise and extreme climatic events.

Extreme drought events have been linked to sudden vegetation diebacks in several salt marshes in the USA Southeast and Gulf of Mexico over the last 25 years (Alber et al., 2008; Lindstedt et al., 2006; McKee et al., 2004). The drought-induced mechanism of sudden marsh dieback has been associated with the interaction and cascading effects of multiple stressors, such as extreme porewater salinity, sediment acidification due to the oxidation of air-exposed sulfur minerals into sulfuric acid, infection by fungal pathogens, and an exponential increase of periwinkle snail herbivory (Alber et al., 2008; Hughes et al., 2012; McKee et al., 2004; Silliman et al., 2005). More recently, extreme precipitation from catastrophic tropical cyclones was suggested to trigger marsh vegetation diebacks by either intrusion of salt water into brackish/freshwater marshes and/ or by extending periods of tidal inundation which in turn exceed the physiological tolerance of marsh plants (Marsh et al., 2016; Ramsey et al., 2012; Stagg et al., 2021). The co-occurrence of biannual vegetation loss with extreme drought and flooding events in our study provides substantial evidence that marsh vegetation dieback is attributed to extreme climatic events in the USA Southeast.

Salt marsh dieback events are often described as transient phenomena (Alber et al., 2008). In contrast, we show that marsh dieback triggered by extreme climatic events can be long-lasting, with an approximate 14% loss of total vegetated area after 10 years. To the best of our knowledge, this is the first time a decadal timeframe is used to characterize a long-term salt marsh dieback event induced by an extreme climatic event. It is intriguing that the surface elevation of the dieback areas, or elevation capital, remains within the range of a hypothetically healthy marsh (Morris et

al., 2002). Furthermore, we provide evidence that the marsh may be restored by grass planting,

suggesting that the stressors that caused dieback may be overcome, despite the fact that the

majority of dieback areas have not revegetated naturally. Potential explanations for slow or no

recovery require further research and may be associated with limited seed dispersion and viability,

poor vegetative reproduction, as well as disturbed marsh hydrology and geomorphology (i.e., tidal

creek erosion and sedimentation limiting previously normal tidal flow).

The increasing frequency and intensity of severe climatic events due to climate change is likely to

further develop marsh diebacks similar to those we observed in Charleston (Brown et al., 2019;

Dai, 2013; Paerl et al., 2019). Long-term vegetation loss, as shown in this study and previous work,

increases the vulnerability of salt marsh ecosystems to SLR (Baustian et al., 2012; Mudd et al.,

2009; 2010; Fig. 6). Thus, the effects of extreme climatic events on marsh vegetation must be

incorporated into assessments of the resilience of the ecosystem to climate change. Current models

designed to project marsh resilience to accelerated SLR often assume that the ecosystem collapses

due to the progressive loss of elevation relative to mean sea level (Morris et al, 2002; Kirwan et

al., 2010, 2016b). In other words, the marsh drowns with the loss of elevation capital. Failing to

incorporate the increasing frequency and intensity of extreme climatic events which trigger

irreversible marsh diebacks will lead to an underestimation of the vulnerability of salt marshes to

climate change. For example, our modelling approach forecasted that after 100 years under a

moderate climate change scenario, the elevation of a highly productive marsh would exceed that

of an unvegetated dieback marsh by 17.5 cm (see below for further discussion).

4.2 Call for restoration of urban marshes

Urban salt marsh ecosystems are more vulnerable than non-urban marshes due to their limited

migration space (Schuerch et al., 2018). At the same time, the value of the ecosystem services they

provide are very high due to their close proximity to heavily populated areas. Thus, their

conservation and restoration have been proposed as an effective adaptation strategy to climate

change, with special emphasis on their ability to attenuate storm surge and coastal erosion

(Temmerman et al., 2013; Geedicke et al., 2018). Hybrid solutions in which engineered structures

are combined with coastal wetlands to protect built infrastructure from storm surge were shown to

be more effective than engineered defenses alone (Temmerman et al., 2013; Zhu et al., 2020).

However, despite the enhanced value and vulnerability of urban salt marshes, these ecosystems

remain understudied in comparison to their non-urban counterparts (Alldred et al., 2020).

Our results corroborate evidence from non-urban marshes to show that marsh grass planting is an

effective restoration practice in an urban marsh that experienced dieback (Baustian et al., 2012;

Linthurst and Seneca, 1980; Ogburn and Alber, 2006). The multiple stressors that lead to sudden

vegetation dieback have yet to be completely revealed (Evans et al., 2021). The ability of

unvegetated marshes to respond positively to grass planting is likely dependent on the strength of

the marsh platform and whether it is located within the elevation range of a stable marsh domain

(*sensu* Morris et al., 2002), along with sufficient sediment supply from the surrounding estuary

(Baustian et al., 2012; Liu et al., 2021; Webb et al., 1995; Wilsey et al., 1992). As shown in our

modelling approach and by previous work, plant primary production is a key variable modulating

marsh accretion (Morris et al., 2002; Mudd et al., 2009, 2010; Silliman et al., 2019). Thus, rapid

restoration of marsh dieback is crucial in order to avoid further degradation, especially in stressed

urban salt marshes. The urgency increases when taking into consideration that dieback areas in

West Ashley and other studied locations are already associated with lower elevation terrain (Alber

et al., 2008; Marsh et al., 2016). Moreover, unvegetated marshes are prone to fill adjacent tidal

channels and to experience marsh platform erosion, which is exacerbated during low tide rainfall

events (Ganju et al., 2017; Temmerman et al., 2012; Torres et al., 2004). Thus, rapid restoration

by grass planting in marshes that have experienced long-term dieback would not only reinstate the

marsh's ability to accrete sediment, but also prevent further disturbance of marsh hydrology. The

effectiveness of any restoration effort over the long term must include careful consideration of

elevation capital and hydrology. Recent studies show that drainage density and distance from a

channel network increase marsh resilience to future drought-induced dieback events (Liu et al.,

2020).

4.3 Adaptive management of salt marsh ecosystems under climate change

Grass transplanting is an effective restoration practice that bolsters high rates of sediment accretion

under conservative SLR scenarios (Silliman et al., 2019; Temmink et al., 2020). However, our

model results indicate that even highly productive marshes will likely drown under more extreme

climate change scenarios. Our results are supported by long-term modelling performed in Georgia,

USA which predicts an 88 % loss of initial marsh area by 2160 (Langston et al., 2021). Field

verification of remotely sensed data and site-specific monitoring are critical for the assessment of

salt marsh resilience to accelerated SLR in urban environments. If grass planting alone is not

sufficient to maintain marsh elevation under accelerated SLR, natural accretion could be

augmented by the direct application of sediment. This can be performed by either beneficial use

of dredging material, or by sediment redistribution within the same marsh system from edge marsh

erosion, or other processes (Ford et al., 1999; Hopkinson et al., 2018). However, even periodic

placement of dredged material may not be sufficient under extreme climate change scenarios (Fig

6). Thus, constant monitoring, rapid restoration, and climate change mitigation actions are all

urgently needed to preserve highly valuable present-day marsh ecosystems close to urban

environments.

5. **Acknowledgements**

This research was funded by grant 66115 (to J.E.K.) from the National Fish and Wildlife

Foundation under the National Coastal Resilience Fund program. The views and conclusions

contained in this document are those of the authors and should not be interpreted as representing

the opinions or policies of the U.S. Government or the National Fish and Wildlife Foundation and

its funding sources. Mention of trade names or commercial products does not constitute their

endorsement by the U.S. Government, or the National Fish and Wildlife Foundation or its funding

sources. Grass planting in Maryville, South Carolina was implemented through funding received

from NOAA National Marine Fisheries Service (to M.H.). This research was also funded by grant

1654853 (to J.T.M.) from the National Science Foundation's Division of Environmental Biology

6. **Conflict of Interest**

The authors declare that they have no conflict of interest.

7. **Contributions**

J.LR., J.E.K., M.H., N.W., J.R., A.G., J.C., and J.T.M. designed the research. J.L.R., M.H., K.D.G.,

G.K., N.W., J.R., and J.E.K performed field work, J.L.R., and M.H., conducted laboratory work,

J.L.R. analyzed the data with input from J.T.M., and J.E.K. J.L.R. wrote the first draft of the

manuscript with inputs from J.E.K. and all authors equally contributed to revisions and gave

approval for publication

8. **Data availability statement**

Remote sensed imagery was retrieved from publicly available repositories (https://earthexplorer.usgs.gov/ and https://planet.com/). All code and field data are publicly available at a Zenodo repository: https://doi.org/10.5281/zenodo.6961467

9. References

1. Alber, M., Swenson, E.M., Adamowicz, S.C., Mendelssohn, I.A., 2008. Salt marsh dieback: an overview of recent events in the US. Estuarine, Coastal and Shelf Science, 80:1-11.
2. Alldred, M., Borrelli, J.J., Hoellein, T., Bruesewitz, D., Zarnoch, C., 2020. Marsh plants enhance coastal marsh resilience by changing sediment oxygen and sulfide concentrations in an urban, eutrophic estuary. Estuaries and Coasts, 43:801-813.
3. Barbier, E.B., Hacker, S.D., Kennedy, C., Koch, E.W., Stier, A.C., Silliman, B.R., 2011. The value of estuarine and coastal ecosystem services. Ecological Monographs, 81:169-193.
4. Baustian, J.J., Mendelssohn, I.A., & Hester, M.W., 2012. Vegetation's importance in regulating surface elevation in a coastal salt marsh facing elevated rates of sea level rise. Global Change Biology, 18:3377-3382.
5. Brown, V.M., Keim, B.D., Black, A.W., 2019. Climatology and trends in hourly precipitation for the southeast United States. Journal of Hydrometeorology, 20:1737-1755.
6. Cahoon, D.R., Guntenspergen, G.R., 2010. Climate change, sea-level rise, and coastal wetlands. National Wetlands Newsletter, 32:8-12.
7. Calcagno, V., de Mazancourt, C., 2010. glmulti: An R package for easy automated model selection with (generalized) linear models. Journal of Statistical Software, 34:1–29.
8. Chakraborty, J., Collins, T.W., Montgomery, M.C., Grineski, S.E., 2014. Social and spatial inequities in exposure to flood risk in Miami, Florida. Natural Hazards Review, 15:04014006.
9. Crist, P.J., White, R., Chesnutt, M., Scott, C., Sutter, R., Cutter, P., Dobson, G., 2019. Coastal Resilience Assessment of the Charleston Harbor Watershed. National Fish and Wildlife Foundation
10. Dai, A., 2013. Increasing drought under global warming in observations and models. Nature Climate Change, 3:52-58.
11. Donnelly, J.P., Bertness, M.D., 2001. Rapid shoreward encroachment of salt marsh cordgrass in response to accelerated sea-level rise. Proceedings of the National Academy of Sciences, 98:14218-14223.
12. Eisenman, D.P., Cordasco, K.M., Asch, S., Golden, J.F., Glik, D., 2007. Disaster planning and risk communication with vulnerable communities: lessons from Hurricane Katrina. American Journal of Public Health, 97:S109-S115.
13. Evans, B.R., Möller, I., Spencer, T., 2021. Topological and morphological controls on morphodynamics of salt marsh interiors. Journal of Marine Science and Engineering, 9:311.
14. Ford, M.A., Cahoon, D.R., Lynch, J.C., 1999. Restoring marsh elevation in a rapidly subsiding salt marsh by thin-layer deposition of dredged material. Ecological Engineering, 12:189-205.

15. Ganju, N.K., Defne, Z., Kirwan, M.L., Fagherazzi, S., D'Alpaos, A., Carniello, L., 2017. Spatially integrative metrics reveal hidden vulnerability of microtidal salt marshes. Nature communications, 8:1-7.

16. Geedicke, I., Oldeland, J., Leishman, M.R., 2018. Urban stormwater run-off promotes compression of saltmarshes by freshwater plants and mangrove forests. Science of The Total Environment, 637:137-144.

17. Hijmans, R.J., van Etten, J., 2012. raster: Geographic analysis and modeling with raster data. R package version 3.4-13. http://CRAN.R-project.org/package=raster

18. Hladik, C., Alber, M., 2012. Accuracy assessment and correction of a LIDAR-derived salt marsh digital elevation model. Remote Sensing of Environment, 121:224-235.

19. Hopkinson, C.S., Morris, J.T., Fagherazzi, S., Wollheim, W.M., Raymond, P.A., 2018. Lateral marsh edge erosion as a source of sediments for vertical marsh accretion. Journal of Geophysical Research: Biogeosciences, 123:2444-2465.

20. Hughes, A.L., Wilson, A.M., Morris, J.T., 2012. Hydrologic variability in a salt marsh: Assessing the links between drought and acute marsh dieback. Estuarine, Coastal and Shelf Science, 111:95-106.

21. Kirwan, M.L., Guntenspergen, G.R., Morris, J.T., 2009. Latitudinal trends in *Spartina alterniflora* productivity and the response of coastal marshes to global change. Global Change Biology, 15:1982-1989.

22. Kirwan, M.L., Guntenspergen, G.R., D'Alpaos, A., Morris, J.T., Mudd, S.M., Temmerman, S., 2010. Limits on the adaptability of coastal marshes to rising sea level. Geophysical Research Letters, 37:L23401.

23. Kirwan, M.L., Megonigal, J.P., 2013. Tidal wetland stability in the face of human impacts and sea-level rise. Nature, 504:53-60.

24. Kirwan, M.L., Temmerman, S., Skeehan, E.E., Guntenspergen, G.R., Fagherazzi, S., 2016a. Overestimation of marsh vulnerability to sea level rise. Nature Climate Change, 6:253-260.

25. Kirwan, M.L., Walters, D.C., Reay, W.G. and Carr, J.A., 2016b. Sea level driven marsh expansion in a coupled model of marsh erosion and migration. Geophysical Research Letters, 43:4366-4373.

26. Kirwan, M.L., Gedan, K.B., 2019. Sea-level driven land conversion and the formation of ghost forests. Nature Climate Change, 9:450-457.

27. Kulp, S.A., Strauss, B.H., 2019. New elevation data triple estimates of global vulnerability to sea-level rise and coastal flooding. Nature communications, 10:1-12.

28. Langston, A.K., Alexander, C.R., Alber, M., & Kirwan, M.L., 2021. Beyond 2100: Elevation capital disguises salt marsh vulnerability to sea-level rise in Georgia, USA. Estuarine, Coastal and Shelf Science, 249:107093.

29. Lenth, R.V., 2016. Least-squares means: The R package lsmeans. Journal of Statistical Software, 69:1–33.

30. Liaw, A., Wiener, M., 2002. Classification and Regression by randomForest. R News 2, 18-22.

31. Lindstedt, D.M., Swenson, E.M., Reed, D., Twilley, R., Mendelssohn, I.A., 2006. The case of the dying marsh grass. Report submitted to the Louisiana Department of Natural Resources, Baton Rouge, LA.

32. Linthurst, R.A., Seneca, E.D., 1980. Dieback of salt-water cordgrass (*Spartina alterniflora*
Loisel.) in the lower Cape Fear estuary of North Carolina: an experimental approach to re-
establishment. Environmental Conservation, 7:59-66.
33. Liu, Z., Fagherazzi, S., She, X., Ma, X., Xie, C., Cui, B., 2020. Efficient tidal channel networks
alleviate the drought-induced die-off of salt marshes: Implications for coastal restoration and
management. Science of The Total Environment, 749:141493.
34. Liu, Z., Fagherazzi, S., Cui, B., 2021. Success of coastal wetlands restoration is driven by
sediment availability. Communications Earth & Environment, 2:1-9.
35. Marsh, A., Blum, L.K., Christian, R.R., Ramsey, E., Rangoonwala, A., 2016. Response and
resilience of *Spartina alterniflora* to sudden dieback. Journal of Coastal Conservation, 20:335-
350.
36. Martínez, M.L., Intralawan, A., Vázquez, G., Pérez-Maqueo, O., Sutton, P., & Landgrave, R.,
2007. The coasts of our world: Ecological, economic and social importance. Ecological
Economics, 63:254-272.
37. Maune, D.F., Maitra, J.B., McKay, E.J., 2007. Accuracy standards and guidelines. In D. Maune
(Ed.), Digital elevation model technologies and applications: The DEM user's manual (pp. 65–
97). (2nd Edition). Bethesda, Maryland: American Society for Photogrammetry and Remote
Sensing.
38. MAXAR, 2021. Satellite imagery. Product guide. v. 2.6
39. McKee, K.L., Mendelssohn, I.A., Materne, M.D., 2004. Acute salt marsh dieback in the
Mississippi River deltaic plain: a drought-induced phenomenon? Global Ecology and
Biogeography, 13:65-73.
40. Mendelssohn I.A., Morris J.T., 2002. Eco-Physiological Controls on the Productivity of
Spartina Alterniflora Loisel. In: Weinstein M.P., Kreeger D.A. (eds) Concepts and
Controversies in Tidal Marsh Ecology. Springer, Dordrecht.
41. Morris, J.T., Sundareshwar, P.V., Nietch, C.T., Kjerfve, B., Cahoon, D.R., 2002. Responses
of coastal wetlands to rising sea level. Ecology, 83:2869-2877.
42. Morris, J.T., Sundberg, K., Hopkinson, C.S., 2013, Salt marsh primary production and its
responses to relative sea level and nutrients in estuaries at Plum Island, Massachusetts, and
North Inlet, South Carolina, USA. Oceanography 26:78-84.
43. Morris, J.T., Barber, D.C., Callaway, J.C., Chambers, R., Hagen, S.C., Hopkinson, C.S., et al.,
2016. Contributions of organic and inorganic matter to sediment volume and accretion in tidal
wetlands at steady state. Earth's Future, 4:110-121.
44. Morris, J.T., Renken, K.A., 2020. Past, present, and future nuisance flooding on the Charleston
peninsula. Plos one, 15:e0238770.
45. Morris, J. T., Cahoon, D. R., Callaway, J. C., Craft, C., Neubauer, S. C., Weston, N. B., 2021.
Marsh Equilibrium Theory: Implications for Responses to Rising Sea Level. In: FitzGerald, D.
M. and Hughes, Z. J. (eds) Salt Marshes: Function, Dynamics, and Stresses. Cambridge:
Cambridge University Press, pp. 157–177.
46. Mudd, S.M., Howell, S.M., Morris, J.T., 2009. Impact of dynamic feedbacks between
sedimentation, sea-level rise, and biomass production on near-surface marsh stratigraphy and
carbon accumulation. Estuarine, Coastal and Shelf Science, 82:377-389.

47. Mudd, S.M., D'Alpaos, A., Morris, J.T., 2010. How does vegetation affect sedimentation on tidal marshes? Investigating particle capture and hydrodynamic controls on biologically mediated sedimentation. Journal of Geophysical Research: Earth Surface, 115:F03029.

48. Ogburn, M.B., Alber, M., 2006. An investigation of salt marsh dieback in Georgia using field transplants. Estuaries and Coasts, 29:54-62.

49. Paerl, H.W., Hall, N.S., Hounshell, A.G., Luettich, R.A., Rossignol, K.L., Osburn, C.L., Bales, J., 2019. Recent increase in catastrophic tropical cyclone flooding in coastal North Carolina, USA: Long-term observations suggest a regime shift. Scientific reports, 9:1-9.

50. Palmer, W. C., 1965: Meteorological drought. U.S. Weather Bureau Research Paper 45, 58 pp. Washington, DC

51. Planet Team, 2019. Planet Imagery Product Specifications. San Francisco, CA. https://assets.planet.com/docs/Planet_Combined_Imagery_Product_Specs_letter_screen.pdf

52. QGIS.org, 2021. QGIS Geographic Information System. QGIS Association. http://www.qgis.org

53. R Core Team, 2021. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

54. Ramsey III, E., Werle, D., Suzuoki, Y., Rangoonwala, A., Lu, Z., 2012. Limitations and potential of satellite imagery to monitor environmental response to coastal flooding. Journal of Coastal Research, 28:457-476.

55. Raposa, K.B., Ekberg, M.L.C., Burdick, D.M., Ernst, N.T., Adamowicz, S.C., 2017. Elevation change and the vulnerability of Rhode Island (USA) salt marshes to sea-level rise. Regional Environmental Change, 17:389-397.

56. Reed, D.J., 1995. The response of coastal marshes to sea-level rise: Survival or submergence? Earth Surface Processes and Landforms, 20:39-48.

57. Rogers, K., Kelleway, J.J., Saintilan, N., Megonigal, J.P., Adams, J.B., Holmquist, J.R., et al., 2019. Wetland carbon storage controlled by millennial-scale variation in relative sea-level rise. Nature, 567:91-95.

58. Schepers, L., Brennand, P., Kirwan, M.L., Guntenspergen, G.R., & Temmerman, S., 2020. Coastal marsh degradation into ponds induces irreversible elevation loss relative to sea level in a microtidal system. Geophysical Research Letters, 47:e2020GL089121.

59. Schuerch, M., Spencer, T., Temmerman, S., Kirwan, M.L., Wolff, C., Lincke, D., et al., 2018. Future response of global coastal wetlands to sea-level rise. Nature, 561:231-234.

60. Silliman, B.R., Van De Koppel, J., Bertness, M.D., Stanton, L.E. and Mendelsohn, I.A., 2005. Drought, snails, and large-scale die-off of southern US salt marshes. Science, 310:1803-1806.

61. Silliman, B.R., He, Q., Angelini, C., Smith, C.S., Kirwan, M.L., Daleo, P., Renzi, J.J., Butler, J., Osborne, T.Z., Nifong, J.C., van de Koppel, J., 2019. Field experiments and meta-analysis reveal wetland vegetation as a crucial element in the coastal protection paradigm. Current Biology, 29:1800-1806.

62. South Carolina DNR, 2018. Report of the South Carolina Charleston County Lidar. https://prd-tnm.s3.amazonaws.com/StagedProducts/Elevation/metadata/SC_CoastalCounties_2017_A17/SC_Charleston_2016/reports/South_Carolina_Lidar_Charleston_Report_180720.pdf

63. South Carolina DNR, 2009. Report of the South Carolina Charleston County Lidar. https://www.fisheries.noaa.gov/inport/item/49978

64. Stagg, C.L., Osland, M.J., Moon, J.A., Feher, L.C., Laurenzano, C., Lane, T.C., Jones, W.R., Hartley, S.B., 2021. Extreme Precipitation and Flooding Contribute to Sudden Vegetation Dieback in a Coastal Salt Marsh. Plants, 10:1841.

65. Sweet, W.V., Kopp, R.E., Weaver, C.P., Obeysekera, J., Horton, R.M., Thieler, E.R., et al., 2017. Global and Regional Sea Level Rise Scenarios for the United States. NOAA Technical Report NOSCO-OPS 086; 44pp.

66. Temmerman, S., Moonen, P., Schoelynck, J., Govers, G., Bouma, T.J., 2012. Impact of vegetation die-off on spatial flow patterns over a tidal marsh. Geophysical Research Letters, 39:L03406.

67. Temmerman, S., Meire, P., Bouma, T.J., Herman, P.M., Ysebaert, T., De Vriend, H.J., 2013. Ecosystem-based coastal defence in the face of global change. Nature, 504:79-83.

68. Temmink, R.J., Christianen, M.J., Fivash, G.S., Angelini, C., Boström, C., Didderen, K., Engel, S.M., Esteban, N., Gaeckle, J.L., Gagnon, K., Govers, L.L., 2020. Mimicry of emergent traits amplifies coastal restoration success. Nature communications, 11:1-9.

69. Torres, R., Goni, M.A., Voulgaris, G., Lovell, C.R., Morris, J.T., 2004. Effects of low tide rainfall on intertidal zone material cycling. Coastal and Estuarine Studies, 59:93-114.

70. Valiela, I., Teal, J.M., Deuser, W.G., 1978. The nature of growth forms in the salt marsh grass *Spartina alterniflora*. The American Naturalist, 112:461–470

71. Webb, E.C., Mendelssohn, I.A., Wilsey, B.J., 1995. Causes for vegetation dieback in a Louisiana salt marsh: a bioassay approach. Aquatic Botany, 51:281-289.

72. Weston, N.B., 2014. Declining sediments and rising seas: an unfortunate convergence for tidal wetlands. Estuaries and Coasts, 37:1-23.

73. Wilsey, B. J., McKee, K.L., & Mendelssohn, I.A., 1992. Effects of increased elevation and macro-and micronutrient additions on *Spartina alterniflora* transplant success in salt-marsh dieback areas in Louisiana. Environmental Management, 16:505-511.

74. Zhu, Z., Vuik, V., Visser, P.J., Soens, T., van Wesenbeeck, B., van de Koppel, J., Jonkman, S.N., Temmerman, S., Bouma, T.J., 2020. Historic storms and the hidden value of coastal wetlands for nature-based flood defence. Nature Sustainability, 3:853-862.

Figure legends:

Figure 1: Dieback trajectory of the saltmarsh located in West Ashley, Charleston, SC. **a** Percentage of unvegetated marsh area over time. Total area is partitioned into creek bank or midmarsh categories based on areas closer or further than 7.5 m to its closest creek bank, respectively. **b** Palmer drought severity index (PDSI) from the South Carolina Southern Division the over time. Values lower than -2 represent moderate drought (dotted line), while values lower than -4 extreme drought conditions (dashed line). **c** Monthly precipitation from the Charleston, SC area retrieved from NOAA's Online Weather Data (NOWData). **d** Maps showing the probability of an area being unvegetated. Based on individual random forest models per image.

Figure 2: Dieback events were associated with an expansion of unvegetated area in the midmarsh zone. Barplots showing the percent increase in unvegetated areas since the last evaluation (1 or 2 years). Analysis performed for the total marsh, midmarsh, and creekbank areas.

Figure 3: Characterization of unvegetated and vegetated areas within the West Ashley, SC salt marsh. **a** Median ± interquartile range of the field-corrected 2017 light detection and ranging (LIDAR) derived digital elevation model (DEM) of vegetated and unvegetated areas over time. **b** Median ± interquartile range of the distance from the closest tidal creek of vegetated and unvegetated areas over time.

Figure 4: Aboveground biomass (AGB) estimation of the West Ashley, SC salt marsh during July 2021. **a** Map of AGB based on a random forest model using the 2017 elevation digital elevation model, vegetation indices and bands from a PlanetScope 4-band image. **b** Density plot of the predicted AGB values. **c** Allometric biomass equation based on shoot biomass. **d** Goodness of fit of the random forest model in the testing and training datasets. Relation between predicted AGB and distance from tidal creek **(e)**, and the 2017 elevation model **(f)**. Points and error bars represent AGB median and interquartile range on equidistant increments, respectively.

Figure 5 Grass planting is an effective restoration practice in Maryville, SC. **a** Drone image from the restoration site. Plots sampled are labelled on top in white. Photo acquired by the South Carolina Department of Natural Resources Shellfish Research Section on 21 July of 2021. **b** Boxplot representing the interquartile range of aboveground biomass from all sampled creek bank, midmarsh, and 2020 and 2021 restored plot during the July 2021 field campaign. **c** Boxplot representing the interquartile range of aboveground biomass from 10 quadrats sampled from the two reference sites, and the 2019, 2020, and 2021 restored plot during September of 2021. Different letters indicate statistically significant differences ($p < 0.05$) between treatments according to a Tukey test using *emmeans* R package (Lenth, 2016).

Figure 6 Modelled salt marsh resilience under different scenarios of sea level rise, human intervention, and plant primary production using the Marsh Equilibrium Model (MEM) (Morris et al., 2002). Modelled optimal biomass values ranged from 0 g m^{-2} y^{-1} (mudflat) to highly productive salt marshes of 2000 g m^{-2} y^{-1}. Thin Layer Placement was assumed to be performed every 25 years by applying 5 cm of inorganic sediment. Vegetation was assumed to fully recover after 10 years of disturbance. The two sea level rise scenarios modelled were based on regional NOAA´s intermediate low, and intermediate of 60 cm and 122 cm in 100 y, respectively (Sweet et al., 2017).

716   Marsh elevation (cm) over time projected to NAVD88 (**a**), and mean sea level (MSL) (**b**). **c**
717   Modelled salt marsh aboveground biomass over time.

Figure 1: Dieback trajectory of the saltmarsh located in West Ashley, Charleston, SC. **a** Percentage of unvegetated marsh area over time. Total area is partitioned into creek bank or midmarsh categories based on areas closer or further than 7.5 m to its closest creek bank, respectively. **b** Palmer drought severity index (PDSI) from the South Carolina Southern Division the over time. Values lower than -2 represent moderate drought (dotted line), while values lower than -4 extreme drought conditions (dashed line). **c** Monthly precipitation from the Charleston, SC area retrieved from NOAA's Online Weather Data (NOWData). **d** Maps showing the probability of an area being unvegetated. Based on individual random forest models per image.

Figure 2: Dieback events were associated with an expansion of unvegetated area in the midmarsh zone. Barplots showing the percent increase in unvegetated areas since the last evaluation (1 or 2 years). Analysis performed for the total marsh (a), midmarsh (b), and creekbank (c) areas.

765 **Figure 3:** Characterization of unvegetated and vegetated areas within the West Ashley, SC salt
766 marsh. **a** Median ± interquartile range of the field-corrected 2017 light detection and ranging
767 (LIDAR) derived digital elevation model (DEM) of vegetated and unvegetated areas over time. **b**
768 Median ± interquartile range of the distance from the closest tidal creek of vegetated and
769 unvegetated areas over time.

Figure 4: Aboveground biomass (AGB) estimation of the West Ashley, SC salt marsh during July
2021. **a** Map of AGB based on a random forest model using the 2017 elevation digital elevation
model, vegetation indices and bands from a PlanetScope 4-band image. **b** Density plot of the
predicted AGB values. **c** Allometric biomass equation based on shoot biomass. **d** Goodness of fit
of the random forest model in the testing and training datasets. Relation between predicted AGB
and distance from tidal creek **(e)**, and the 2017 elevation model **(f)**. Points and error bars represent
AGB median and interquartile range on equidistant increments, respectively.

 Figure 5 Grass planting is an effective restoration practice in Maryville, SC. **a** Drone image from
 the restoration site. Plots sampled are labelled on top in white. Photo acquired by the South
 Carolina Department of Natural Resources Shellfish Research Section on 21 July of 2021. **b**
 Boxplot representing the interquartile range of aboveground biomass from all sampled creek bank,
 midmarsh, and 2020 and 2021 restored plot during the July 2021 field campaign. **c** Boxplot
 representing the interquartile range of aboveground biomass from 10 quadrats sampled from the
 two reference sites, and the 2019, 2020, and 2021 restored plot during September of 2021.
 Different letters indicate statistically significant differences ($p < 0.05$) between treatments
 according to a Tukey test using *emmeans* R package (Lenth, 2016).

Figure 6 Modelled salt marsh resilience under different scenarios of sea level rise, human intervention, and plant primary production using the Marsh Equilibrium Model (MEM) (Morris et al., 2002). Modelled optimal biomass values ranged from 0 g m⁻² y⁻¹ (mudflat) to highly productive salt marshes of 2000 g m⁻² y⁻¹. Thin Layer Placement was assumed to be performed every 25 years by applying 5 cm of inorganic sediment. Vegetation was assumed to fully recover after 10 years of disturbance. The two sea level rise scenarios modelled were based on regional NOAA´s intermediate low, and intermediate of 60 cm and 122 cm in 100 y, respectively (Sweet et al., 2017). Marsh elevation (cm) over time projected to NAVD88 (**a**), and mean sea level (MSL) (**b**). **c** Modelled salt marsh aboveground biomass over time.

Supplementary Information

Supplementary Text S1

<u>Correction of LIDAR derived DEM.</u>

Error correction of the 2017 light detection and ranging (LIDAR) derived digital elevation model

(DEM) was constructed based on a multiple regression model (further detail in Materials and

Methods). Error was determined based on the difference between real-time kinematic (RTK)

survey points and values from the LIDAR derived DEM raster. Inputs for the model selection were

NDVI, MSAVI2, and independent bands from a PlanetScope satellite image taken days before the

LIDAR acquisition. Model selection determined the following regression model as the best fit:

$\ln(Z_{DEM} - Z_{RTK} + 1) = 0.226\ Z_{DEM} + 105.967\ MSAVI2 + 177.408\ Red - 179.504\ NIR + 0.069$

Where Z is elevation in NAVD88 (m).

The adjusted R^2 in the training dataset was 41.17%

The median ($1^{st} - 3^{rd}$ quartile) error of the original and corrected DEM were -0.03 m (-0.06 – 0.002

m), and -0.01 m (-0.05 – 0.01 m), respectively. Model correction reduced the RMSE of both the

test and train datasets in 27.3% and 30.5%, respectively (Fig. S2). The RMSE of the corrected test

and train dataset were of 5.7 cm, and 8.0 cm, respectively. These values are withing the range of

other error corrected DEMs. For instance, RMSE values from a corrected DEM in Sapelo Island,

GA ranged from 5 cm to 30 cm depending on the cover class (Hladik and Alber, 2012).

Supplementary Table S1: Technical details of aerial and satellite imagery utilized in the present study.

Imagery Source	Year	Acquisition Date (mm/dd/yyyy)	Resolution (m)	Sensor	Red (nm)	Green (nm)	Blue (nm)	NIR (nm)
NAIP	2009	04/21/2009	1	ADS-40	610-660	535-585	430-490	835-885
NAIP	2011	05/08/2011	1	ADS80-SH82	608-662	533-587	428-492	833-887
NAIP	2013	09/29/2013	1	ADS40-SH51/ADS40-SH81	610-660	535-585	430-490	835-885
NAIP	2015	06/12/2015	1	ADS-100	619-651	525-585	435-495	808-882
MAXAR	2016	10/11/2016	0.3	WorldView-3	630-690	510-580	450-510	770-895
NAIP	2017	09/19/2017	1	ADS-100	619-651	525-585	435-495	808-882
NAIP	2019	07/01/2019	1	ADS-100	619-651	525-585	435-495	808-882
Planet	2017	02/20/2017	3	PlanetScope	590-670	500-590	455-515	780-860
Planet	2021	08/01/2021	3	PlanetScope	590-670	500-590	455-515	780-860

845   **Supplementary Table S2**: Technical description of the light detection and ranging (LIDAR)

846   acquisition. Information retrieved from South Carolina DNR (2018).

Item	Parameter
System	Riegl LMS-Q1560
Altitude (AGL meters)	2043
Approx. Flight Speed (knots)	150
Scanner Pulse Rate (kHz)	800
Scan Frequency (hz)	169
Pulse Duration of the Scanner (nanoseconds)	3
Pulse Width of the Scanner (m)	0.9
Swath width (m)	2289
Central Wavelength of the Sensor Laser (nanometers)	1064
Did the Sensor Operate with Multiple Pulses in The Air? (yes/no)	Yes
Beam Divergence (milliradians)	0.25 mrad
Nominal Swath Width on the Ground (m)	2289
Swath Overlap (%)	30
Total Sensor Scan Angle (degree)	58.52
Computed Down Track spacing (m) per beam	0.68
Computed Cross Track Spacing (m) per beam	0.68
Nominal Pulse Spacing (single swath), (m)	0.7
Nominal Pulse Density (single swath) (ppsm), (m)	2
Aggregate NPS (m) (if ANPS was designed to be met through single coverage, ANPS and NPS will be equal)	0.7
Aggregate NPD (m) (if ANPD was designed to be met through single coverage, ANPD and NPD will be equal)	2
Maximum Number of Returns per Pulse	unlimited

Source: South Carolina DNR, 2018

Supplementary Figure S1. Study area and sampling locations of the present study in West Ashley, Charleston, SC. **a** Map of the study
area. Blue lines represent tidal creeks, orange dots sampled location in the field trips of May and July of 2021. **b** Picture of the now
dieback Maryville marsh circa 1970. Image provided by Mr. John Carr Jr. **c** Drone image of the Maryville marsh taken on 21 July of
2021. Drone image acquired by the South Carolina Department of Natural Resources Shellfish Research Section.

Supplementary Figure S2. Relationship between *in situ* aboveground biomass and distance from tidal creek. The vertical dotted line
represents the tipping point (7.5 m distance from the closest tidal creek) defining the state change between creek bank and midmarsh
Spartina alterniflora population. Vegetation data was generated during the field trip of July of 2021.

Supplementary Figure S3: Error correction of the 2017 light detection and ranging (LIDAR) derived digital elevation model (DEM).
Biplot of the observed elevation from a real time-kinematic (RTK) survey and elevation from the uncorrected (**a**) and corrected (**d**) 2017
LIDAR derived DEM. Frequency profile of the DEM error (LIDAR elevation – RTK elevation) of the uncorrected (**b**) and corrected (**e**)
dataset. Frequency profile of the uncorrected (**c**) and corrected (**f**) elevation from the 2017 LIDAR derived DEM.

 Supplementary Figure S4. Median ± interquartile range of an uncorrected 2009 light detection and ranging (LIDAR) derived digital
 elevation model (DEM) of vegetated and unvegetated areas over time (South Carolina DNR, 2009).

References cited in the Supplementary Information

1. Hladik, C., Alber, M., 2012. Accuracy assessment and correction of a LIDAR-derived salt
marsh digital elevation model. Remote Sensing of Environment, 121:224-235.
2. South Carolina DNR, 2018. Report of the South Carolina Charleston County Lidar. https://prd-
tnm.s3.amazonaws.com/StagedProducts/Elevation/metadata/SC_CoastalCounties_2017_A17
/SC_Charleston_2016/reports/South_Carolina_Lidar_Charleston_Report_180720.pdf
3. South Carolina DNR, 2009. Report of the South Carolina Charleston County Lidar.
https://www.fisheries.noaa.gov/inport/item/49978