Title:

Biofilms preserve the transmissibility of a multi-drug resistance plasmid

Authors:

Genevieve A. Metzger[1,2,3], Benjamin J. Ridenhour[2,3,4], Michael France[2,3], Karol Gliniewicz[1,2], Jack

Millstein[1,2], Matthew L. Settles[2], Larry J. Forney[1,2,3], Thibault Stalder[1,2], Eva M. Top[1,2,3] *

Authors Affiliations:

[1] Department of Biological Sciences, University of Idaho, Moscow, Idaho, USA

[2] Institute for Interdisciplinary Data Sciences, University of Idaho, Moscow, Idaho, USA

[3] Bioinformatics and Computational Biology Graduate Program (BCB), Moscow, Idaho, USA

[4] Department of Mathematics and Statistical Science, University of Idaho, Moscow, Idaho, USA

<u>* Corresponding Author:</u>

Department of Biological Sciences

Life Sciences South 258

University of Idaho

Moscow, ID 83844-3051

Phone: +01 (208) 885-5015

email: evatop@uidaho.edu

Keywords: in *Acinetobacter baumannii*, antibiotic resistance, horizontal gene transfer, conjugation,

plasmid

ABSTRACT:

Self-transmissible multidrug resistance (MDR) plasmids are a major health concern because they can spread antibiotic resistance to pathogens. Even though most pathogens form biofilms, little is known about how MDR plasmids persist and evolve in biofilms. We hypothesize that (i) biofilms act as refugia of MDR plasmids by retaining them in the absence of antibiotics longer than well-mixed planktonic populations, and that (ii) the evolutionary trajectories that account for the improvement of plasmid persistence over time differ between biofilms and planktonic populations. In this study, we evolved *Acinetobacter baumannii* with an MDR plasmid in biofilm and planktonic populations with and without antibiotic selection. In the absence of selection biofilm populations were better able to maintain the MDR plasmid than planktonic populations. In planktonic populations plasmid persistence improved rapidly but was accompanied by a loss of genes required for the horizonal transfer of plasmids. In contrast, in biofilms most plasmids retained their transfer genes, but on average plasmid persistence improved less over time. Our results showed that biofilms can act as refugia of MDR plasmids and favor the horizontal mode of plasmid transfer, which has important implications for the spread of MDR.

INTRODUCTION

Antibiotic resistance is widely recognized as one of the most serious problems facing healthcare today. Resistance to antibiotics, including those of last resort, often results from the acquisition of resistance genes by horizontal gene transfer (HGT) [1,2]. When present as part of mobile genetic elements such as plasmids, resistance genes can rapidly spread to various species or strains of bacteria [3,4]. This makes multidrug resistance (MDR) plasmids, particularly those with a broad host range, a major problem in fighting the spread of antibiotic resistance.

Understanding how these plasmids evolve in the presence and absence of antibiotics is critically important. In the presence of antibiotics, MDR plasmids are maintained in bacterial populations via positive selection [5], but when antibiotics are removed the fitness cost of plasmid carriage [6–8] is no longer offset by the benefit of antibiotic resistance [9]. Therefore, plasmids are expected to be lost from bacterial populations via purifying selection unless (i) partitioning and post-segregational mechanisms limit the formation of plasmid-free cells, (ii) reacquisition of plasmids counteracts plasmid loss and cost, (iii) co-residing plasmids and other bacteria in the community can prevent plasmid loss [10,11], or (iv) the cost of plasmid carriage is reduced, eliminated or reversed by evolution [12]. The rate at which plasmids are lost from a population probably depends on a combination of these factors and is inversely proportional to 'plasmid persistence' – the ability of a plasmid to maintain itself in the absence of known positive selection for the plasmid. Previous studies from our and other research groups have repeatedly shown that plasmid persistence can improve during growth in the presence or absence of antibiotics (for example, [13–24]). However, almost all these studies used planktonic populations. Much less is known about how plasmid persistence evolves in spatially structured populations such as biofilms, even though they represent the most common form of bacterial growth and are the causes of many recalcitrant infections [25–28].

Due to spatial structure, bacteria that grow in biofilms only compete locally with their neighbors, which protracts selective sweeps and leads to increased genotypic and phenotypic variation within a population [29–39]. Additionally, the gradients of nutrients and electron acceptors cause habitat heterogeneity that results in local adaptation of subpopulations. These biofilm features allow bacteria to access more fitness peaks in rugged adaptive landscapes. This can result in evolutionary outcomes that may not be observed in planktonic populations that routinely experience strong selective sweeps [40–42]. Previously we showed that under antibiotic selection, evolution in biofilms results in a higher diversity of plasmid persistence phenotypes as compared to planktonic populations [43]. In a separate study we have found that biofilms of plasmid-bearing cells maintain a greater diversity of mutations than planktonic populations. Among these diverse biofilm genotypes were clones that better retained their plasmid than any clone evolved in planktonic populations [44]. These seminal studies showed how biofilm growth can affect the evolution of plasmids persistence. Because many bacterial pathogens with antibiotic resistance plasmids [45,46] form biofilms, we need to study the evolution of their plasmids under these growth conditions.

The biofilm-forming bacterium *Acinetobacter baumannii* is an emerging threat in the United States and worldwide because it causes wound infections, ventilator-associated pneumonia, and sepsis [47,48]. Further, it is known to rapidly acquire new forms of antibiotic resistance by HGT [49–51] and pan-drug resistant strains have been documented [52]. Despite the medical importance of *A. baumannii* and the role of HGT in its acquisition of antibiotics resistance, little information is available on how this organism and recently acquired MDR plasmids co-evolve.

Previously we found that the persistence of a broad-host range MDR plasmids pB10 in *A. baumannii* increased over time in both biofilm and planktonic populations treated with antibiotics, but this phenotype showed more diversity after evolution in biofilms [43]. Building on this previous study we

sought to determine the effect of biofilm growth (i) on plasmid persistence in the absence of antibiotics,

and (ii) on the evolutionary trajectories of this plasmid in the presence and absence of antibiotics. Our

findings can be summarized as follows. First, in the absence of selection, biofilm populations

maintained the MDR plasmid longer than planktonic populations. Second, the evolution of plasmid

persistence in the absence of antibiotic selection was protracted in biofilms. Third, large deletions of the

plasmid conjugative transfer regions were a common evolutionary outcome in planktonic populations

but less so in biofilms, indicating that growth in biofilms conserved the plasmid's ability to transfer

horizontally. Finally, plasmid adaptation to one bacterial species can either promote or decrease its

persistence in other bacteria, including other opportunistic pathogens.

RESULTS

Experimental set-up of biofilm and planktonic populations.

To investigate the effects of biofilm growth and the presence of a plasmid-selective antibiotic on

the persistence and evolution of MDR plasmid pB10 in *A. baumannii,* we set up an experiment with a

full factorial design with two growth environments and two treatments. As shown in Fig. 1, the growth

environment was either a biofilm flow cell or serial batch culture; the treatments were the presence

(Tet+) or absence (Tet-) of tetracycline in the growth medium. In order to initiate all four

environment/treatment combinations with equivalent populations, replicate biofilm populations were

first established in flow cells for a period of four days ($t_{-4} – t_0$). To avoid plasmid loss during this initial

phase, the medium was supplemented with Tet for four days, regardless of the subsequent treatment

(Tet+ or Tet-). After this biofilm establishment phase at t_0, triplicate flow cells were harvested and used

to inoculate triplicate serial batch cultures per treatment. From here on we refer to these serial batch

cultures as planktonic populations. The remainder of the growth and sampling scheme is depicted in Fig.

1 (see Materials and Methods for more details).

Biofilms retain the plasmid in the absence of selection.

To assess the ability of bacterial biofilms to retain an MDR plasmid in the absence of antibiotics,

we compared the proportion of plasmid-containing cells present in biofilm and planktonic populations at

t_0 and after 14 days of growth in Tet-free media (t_{14}) (Table 1). This was done by plating samples of

these populations on selective and non-selective media. The plasmid was lost more slowly from biofilms

than from planktonic populations. After 14 days 12.3 ($\pm$ 3.5)% of the viable cells from the biofilms still

carried the plasmid. In contrast, plasmid-bearing cells could no longer be detected in two of the three

planktonic populations (i.e., their frequency in these populations was $<10^{-8}$), and were present in only

0.016% of the third. This observation demonstrates the importance of biofilms in the persistence of

plasmid-mediated antibiotic resistance.

Biofilms protracted the evolution of plasmid persistence in the absence of selection.

We previously showed that after 28 days of evolution in the presence of Tet, bacteria isolated

from biofilms showed a higher diversity in plasmid persistence dynamics than their counterparts evolved

in planktonic populations, and on average, a smaller increase in plasmid persistence [43]. In the current

study we compared the evolution of plasmid persistence in clones evolved over 14 days in the presence

and absence of Tet (Tet+ and Tet- populations). Because the plasmid was rapidly lost in Tet- planktonic

populations, we ended the entire Tet- experiment on day 14. In one of these three Tet- populations,

plasmid-containing clones could no longer be detected after day 6, and in another after day 10.

Therefore the plasmid-containing clones from Tet- planktonic populations were isolated from samples

archived on t_6, t_{10}, and t_{14}. In contrast, all the clones from Tet- biofilm populations were isolated on t_{14}.

For each of the triplicate growth environment/treatment combinations plasmid persistence was tested for

six clones. First, we verified that clones isolated at t_0 showed the same plasmid persistence profile as the

ancestor (Fig. 2A). Based on linear regression there was no significant change in the rate of plasmid

loss, meaning the plasmid persistence in the populations used to start the experiment did not change

during the 4-day biofilm establishment phase (t-test p-values for t_0, Tet+ was 0.7037 and for t_0, Tet- was

0.5195). Therefore, the t_0 clones were used for the comparisons of plasmid persistence described below.

134   After 6–14 days of evolution in the absence of antibiotics, plasmid pB10 showed improved

persistence in *A. baumannii* clones from planktonic populations but not in clones from biofilms (Fig.

2B; t-test p-values were 0.0007 and 0.2107, respectively). These results are in contrast to the clones

evolved in the same conditions but in the presence of Tet, where plasmid persistence improved in clones

isolated from both biofilm and planktonic populations after 14 days (Fig. 2C; t-test p-values were

0.0053 and <0.0001, respectively). The latter result is consistent with our previously published findings

after 28 days of biofilm and planktonic growth [43]. Thus on average, plasmid persistence always

improved in planktonic populations regardless of antibiotic treatment, but only improved in biofilms

grown with Tet.

Plasmid transfer genes were mostly retained in biofilms but completely lost in planktonic populations.

144   To determine the genotypic changes that occurred during the experiment, we sequenced all 18

clones per environment/treatment at each time point (6 per triplicate population, Fig. 1), for a total of

144 strains. Here we focus on genetic changes in the plasmid pB10. We identified three types of

mutations, as shown by a visual summary in Fig. 3 and a list in Supplementary Table 1. By far the most

common genetic changes were large deletions in the plasmid regions encoding conjugative transfer (*tra*),

mating pair formation (*trb*), and an intervening class 1 integron containing sulfonamide and amoxicillin

resistance genes. These large deletions also often included a few genes thought to be involved in

plasmid maintenance and central control, adjacent to the *tra* region. Conjugation assays with a

representative clone showed that this gene loss resulted in the inability of these plasmids to transfer by

conjugation. When the donor strain contained a deletion mutant of plasmid pB10 we could not detect

any transconjugants (Supplementary Table 2). In contrast, transconjugants were detected for the ancestor

and a randomly chosen evolved clone with intact plasmid, and the average counts (T) were significantly

different from 0 (t-test p-values were 0.03662 and 0.01467, respectively).

We also identified nine deletions smaller than 1,000 bp located in maintenance/control genes

kfrA, *klcB*, or one of the *trb* genes. Finally, there were four single nucleotide polymorphisms (SNPs):

two in *kfrA*, one in a *trb* gene, and one in an intergenic region near the *tetR* gene. In summary, large

deletions that remove virtually all the genes required for horizontal transfer were by far the most

common genetic changes.

The size of the large deleted plasmid regions varied drastically between clones, ranging from

21,755 bp to 34,594 bp. The number of genes deleted per plasmids ranged from 23 to 35, but a common

set of 19 genes (from *trbF* through *traC*) were deleted or truncated in every deletion variant (Fig. 3,

Supplementary Table 1). As plasmid pB10 has 65 complete coding sequences (CDS) in addition to a

few truncated transposases [53], all large deletions represented more than a third of the gene content of the

plasmid. Whereas the boundaries of these deletions were all different, most contained short flanking

direct repeats (typically 8-12 bp, with one being only 6 bp, Supplementary Table 1). One of these

repeats was always eliminated during deletion of the intervening region. Deletions between these direct

repeats are indicative of recombination events [54].

There were striking differences in the kinds of plasmid genotypes observed in the biofilm versus

planktonic populations (Fig. 3, Supplementary Table 1). Most of the clones from planktonic populations

showed the large deletions described above. Only 2 of the 54 clones had retained the ancestral pB10

sequence, both from the Tet- population sampled at t_6. In five of the six Tet+ planktonic populations

sampled at either 14 or 28 days, all six clones contained plasmids with identical large deletions.

Moreover, two of the three planktonic populations showed the same large deletion at both time points.

These results indicate strong selective sweeps of large deletion mutants in planktonic populations under

antibiotic selection for the plasmid. Even in the absence of Tet, all six clones from the population that

still retained plasmids by day 14 showed an identical large deletion. In stark contrast, all clones from the

three Tet+ biofilms at t_{14} still contained the ancestral plasmid genotype. Even two weeks later, one of the

three Tet+ biofilm populations still showed intact plasmids in 100% of the clones, and in the other two

populations around half the clones still had intact plasmids. In the Tet- biofilms the deletions were more

frequent at t_{14} but still only found in half or less than half of the clones per population (Supplementary

Table 1 and Fig. 3). Moreover, of the 54 evolved biofilm clones only two pairs showed an identical

deletion. These results clearly show that there were no sweeps of large deletion mutants in the biofilms.

In conclusion, biofilms still contained a large proportion of transmissible plasmids after 14 or 28 days,

whereas transmissible plasmids were no longer detected at these time points in planktonic populations.

Loss of plasmid transfer genes improved plasmid persistence.

To determine the effect of the large plasmid sequence deletions, we compared the persistence of

truncated and full-length plasmids in two ways. First, we compared the persistence of these plasmids in

their co-evolved hosts. On average, clones with truncated plasmids showed a significantly higher

plasmid persistence than clones with full-length plasmids (Fig. 4A, t-test p-value was < 0.0001).

However, a few biofilm clones with full-length plasmids identical in sequence to ancestral pB10, also

demonstrated high plasmid persistence (Fig. 4A), suggesting this change was caused by chromosomal

mutations. Next, to ensure that the observed improvement in persistence of the truncated plasmids could

entirely be explained by the plasmid deletions and not chromosomal changes, we transformed the

ancestral host with one full-length pB10 and several truncated plasmids from t_{28} of the Tet$^+$ treatments.

Plasmid persistence was higher for the truncated plasmids than for the full-length plasmid (t-test p-value

was < 0.0001) or ancestral pB10 (Fig. 4B; t-test p-value was 0.0151). Thus, the large deletions in

plasmid pB10 were a major driver of improved plasmid persistence in *A. baumannii*.

Persistence of evolved pB10 in other naive hosts.

To understand the broader effects of the large deletions in pB10 on plasmid persistence, we

tested one evolved truncated plasmid from a t_{28} clone in three bacterial strains: *Pseudomonas*

moraviensis R28, *Pseudomonas sp. nov.* H2, and *Stenotrophomonas maltophilia* P21 (see footnote 3 in

Supplementary Table 1). All three strains were previously shown to rapidly lose pB10 [55]. The evolved

plasmid was even less persistent than ancestral pB10 in the two *Pseudomonas* hosts, but it was more

persistent in *S. maltophilia* P21 (Fig. 5). Our data demonstrate that even though deletions that eliminate

conjugative transfer of the plasmid may be beneficial in some species, that benefit may not extend to all.

Some chromosomal mutations are also responsible for improved plasmid persistence.

While the mutations found in the plasmids explained most of the increased persistence in the

isolated clones, we identified a few clones where chromosomal mutations must be responsible for the

improved persistence (Figure 4). In the genome sequences of the 144 clones, we identified many

chromosomal mutations when compared to the ancestral sequence (Supplementary Table 3). However,

none of these mutations were in genes previously shown to be involved in plasmid persistence. Due to

the high number of unique mutations, mutation reconstruction in the ancestor would be required to test

their effects, which is outside the scope of this study.

DISCUSSION

In this study we showed that growth in biofilms can promote the retention of MDR plasmids, while preserving the horizontal mode of plasmid transmission better than planktonic populations. These findings have several important implications. First, they demonstrate the positive effect of biofilm growth on MDR plasmid persistence in the absence of antibiotics. This is significant given most bacteria in clinical and environmental habitats grow in biofilms [25]. Our findings are consistent with previous studies that showed that the deeper layers of biofilms can act as refugia for plasmids even in the absence of selection for the plasmid [56,57]. A plausible explanation is that bacteria in biofilm layers farthest away from the bulk medium grow slowly or not at all. This allows plasmids to be better retained in these subpopulations as plasmid loss requires cells to divide. Our results thus suggest that reduced use of antibiotics may help lower the relative abundance of resistance genes in biofilms but not necessarily eliminate them from a population. Second, evolution in biofilms conserved the horizontal mode of transmission of a MDR plasmid, while growth in well-mixed systems resulted in sweeps of much smaller, non-self-transmissible plasmids that persist better through vertical transmission. Finally, this study cautions against indiscriminately extrapolating results from experimental evolution in planktonic cultures to more complex environments such as bacterial biofilms.

Perhaps the most important finding of our study was the striking difference in evolutionary trajectories between the biofilm and planktonic populations. Independent of the presence of selection for the plasmid, virtually every tetracycline resistant *A. baumannii* clone analyzed from planktonic populations had lost a very large segment of plasmid pB10 that contained horizontal transfer and a few antibiotic resistance genes. The loss of these genes resulted in the inability of the plasmid to transfer by conjugation (Supplementary Table 2). While such gene loss may not be a general adaptive strategy for most conjugative plasmids, the findings are consistent with other studies that observed loss of plasmid

transmissibility due to deletion or mutations of transfer genes during evolution in planktonic populations

[21,44,58,59]. In contrast, the majority of plasmids in evolved biofilm clones remained intact and thus self-

transmissible. In only one of the nine evolved biofilm populations did plasmid genotypes with deletions

in transfer genes outnumber the ancestral genotype (a Tet+ biofilm at t_{28}). Biofilms can thus protract the

emergence of non-transmissible plasmid mutants.

There are a few possible explanations for these contrasting results between biofilms and

planktonic populations. The most parsimonious one is that the spatial structure of biofilms protracted

selective sweeps, which is in line with previous studies [29–33,44]. Although we did not compare plasmid

fitness cost, the higher persistence of the much smaller evolved plasmids can almost certainly be

explained by a reduced fitness cost on their host. First, loss of the two transfer regions would be

expected to reduce plasmid cost as horizontal transfer (conjugation) is known to be costly [21,58,59].

Moreover, in the class 1 integron region that was always deleted together with the transfer regions, the

integrase gene could have added to the cost of the plasmid [60,61]. Since this gene is not repressed by LexA

in *Acinetobacter* spp., a highly active integrase expressed from pB10 could have had a toxic effect on

Acinetobacter spp. [62]. Regardless of which genes conferred the highest cost, losing these large plasmid

segments allowed the hosts of truncated plasmids to rapidly sweep through the planktonic Tet resistant

populations. In contrast, due to the absence of global competition in biofilms, clones with wild-type

plasmids were not as readily outcompeted by clones with deletion variants. An argument could be made

that the bacteria in biofilms may have undergone fewer doublings than those in planktonic populations

[63,64], thus slowing evolution. Unfortunately, while the number of doublings in the planktonic populations

can readily be estimated, this is not the case for biofilm populations where the growth rate varies

according to depth and the availability of nutrients. Even if the number of generations differed between

the two environments, what matters is that within a certain time frame, such as a course of antibiotics,

some biofilm forming pathogens may lose the ability to transfer their MDR plasmids less rapidly than

expected based on results from well-mixed systems.

An additional hypothesis that can explain the increased retention of plasmid transfer genes in

bacteria evolving in biofilms is that a tradeoff exists between horizontal and vertical transmission [58]. It is

known that in biofilms close cell-to-cell contact and stabilized mating pair formation can improve the

efficiency of conjugative transfer to neighboring cells relative to planktonic populations [65–68]. This

makes horizontal transfer a more important mechanism of plasmid 'survival' in biofilms [69]. It would

then follow that selection on plasmid transferability may be stronger in biofilms than in axenic

planktonic populations, where it would present a cost rather than a benefit. This is especially true for

IncP-1 plasmids like pB10, which do not transfer efficiently in shaken liquid cultures due to their short

rigid sex pili [70,71]. In such a well-mixed environment, smaller non-self-transmissible plasmids with a

lower fitness cost could thus outcompete transmissible ones and propagate through vertical transmission

only.

Another plausible explanation for higher persistence of self-transmissible plasmids in biofilms is

the positive selection for retention of the sex pilus genes in this environment, which are part of the

transfer gene region *trb* that was deleted in the planktonic clones. Previous studies have shown that

conjugative plasmids can stabilize biofilms during the early stages of biofilm formation due to the

formation of sex pili [65,72]. If pilus expression was favorable for biofilm formation in our flow cells, loss

of the pilus could have been selected against in this environment.

Overall, our results suggest that in planktonic populations, the cost of maintaining the transfer

regions and integron is apparently higher than the benefit of retaining the plasmid through horizontal

transfer. The stark difference in plasmid evolutionary trajectories between planktonic and biofilm

populations and the higher diversity in plasmid persistence phenotypes reported by us previously [43,44]

suggests that it is critical to study bacteria-plasmid coevolution in biofilms to fully understand plasmid

population biology.

The deletions of the large plasmid segments in *A. baumannii* may result from homology-

facilitated illegitimate recombination due to short direct repeats that flanked the boundaries[54].

Illegitimate recombination due to short repeats is well understood in *E. coli,* and has also been reported

in *Acinetobacter baylyi* [73]. The same plasmid pB10 was previously shown by us to lose its tetracycline

resistance operon in *E. coli* through recombination between much larger direct repeats [74]. Interestingly

Porse *et al.* (2016) [21] identified a similar pattern of large deletions of a plasmid's conjugative transfer

region in an IncN plasmid from *Klebsiella pneumoniae* that was evolved in strains of *Escherichia coli*.

In that study the observed deletions were very similar to one another and corresponded to the location of

IS26, whereas the pB10 deletions in our study were not adjacent to any IS element. Given the very short

length of the flanking repeats on plasmid pB10 and the very large deleted fragments, it would be

interesting in future studies to determine the rate at which these deletions occur in this *A. baumannii*

strain.

In our previous studies we never observed loss of the entire plasmid transfer regions of pB10 or

other plasmids of the same IncP-1 group [13,17,18]. Therefore we examined the persistence of one of our

pB10 deletion variants in three other bacterial species. The deletion caused a large increase in plasmid

persistence in *S. maltophilia*, but not in two environmental *Pseudomonas* isolates. *S. maltophilia* is

increasingly found to be an important nosocomial pathogen[75]. Our findings thus show that deletion of

plasmid transfer genes could also contribute to the persistence of MDR plasmids in distantly related

bacterial pathogens.

The persistence of plasmids that remained in the planktonic populations had always improved

within 14 days (140 generations) or less, regardless of the presence or absence of Tet. This finding is

consistent with previous work that showed a reduction in plasmid cost or improvement of persistence

over time in the absence of plasmid-selective antibiotics[15,19,76]. Thus even when MDR plasmids are in

bacteria that are not exposed to antibiotics, they can undergo genetic changes that rapidly improve their

persistence. Upon later exposure to antibiotics, these deletion variants would sweep through the

population, as they can outcompete ancestral or plasmid-free clones. In contrast, in biofilms the average

plasmid persistence only improved significantly in the Tet+ biofilms. The observed discrepancy between

biofilm and planktonic populations may be due to the same variety of factors described above.

Our findings suggest that bacterial biofilms play an important role in the maintenance and spread

of MDR plasmids in natural bacterial populations in several ways. First, growth in biofilms can increase

MDR plasmid persistence in the absence of antibiotics. Second, evolution in biofilms may facilitate the

preservation of plasmid transferability. So far this has not been considered as a possible positive effect

of biofilm growth on the spread of MDR plasmids. These findings point out that while experimental

evolution of bacteria in planktonic populations have addressed important basic evolutionary questions,

these growth conditions are far removed from those of naturally occurring bacterial populations. It is

thus vital that future research on the ecology and evolution of MDR plasmids includes studies on

bacterial biofilms.

METHODS

Bacteria, plasmid, and growth media.

The ancestor used in our study was derived from *Acinetobacter baumannii* strain ATCC 17978

(Accession #CP000521) and is the same as used in our previous study [43]. Here we refer to this strain

simply as *A. baumannii*. The plasmid used was pB10, a 64.5-kbp broad-host-range IncP-1 plasmid from

a German wastewater treatment plant [53] that is poorly maintained in naïve *A. baumannii*. It encodes

resistance to tetracycline, streptomycin, amoxicillin, sulfonamides, and $HgCl_2$. Upon whole-genome

sequencing of the ancestral strain after electroporation with pB10 we observed the loss of a large DNA

segment (~140-kbp). This region was previously thought to be a chromosomal island [77] but more

recently shown to be part of a 150-kbp plasmid pAB3 in this strain [78]. We determined that upon

introduction of plasmid pB10 in ATCC 17978 by electroporation, our ancestor lost pAB3 and is thus a

derivative of ATCC 17978 without that mobile genetic element. In addition to pB10, the strain retained

the two small native plasmids, pAB1 and pAB2. Other strains used in this study were *Pseudomonas* sp.

nov. H2 [79], *Pseudomonas moraviensis* R28 [80], and *Stenotrophomonas maltophilia* P21 [55].

For information regarding the construction of the ancestral host, the MBMS medium with and

without tetracycline (Tet, 10 mg/l), and the culture conditions, we refer to Ridenhour et al. [43].

Experimental evolution protocol.

A timeline of our experimental plan is shown in Fig. 1. The evolution experiment consisted of a

full factorial design with two growth environments and two treatments. The growth environment was

either a biofilm flow cell or serial batch culture. In one treatment (Tet+) the plasmid-selective antibiotic

Tet was added during the entire length of the experiment, whereas in the Tet- treatment Tet was only

present during the 4-day biofilm establishment period ($t_{-4} - t_0$). This antibiotic selection phase was

necessary to avoid establishing biofilms that would consist of a mixture of plasmid-containing and

plasmid-free cells. In all treatments, time point t_0 represents the end of the 'biofilm establishment phase',

which occurred four days after the flow of MBMS-Tet through the flow cells was initiated following

strain inoculation. As the set-up and growth conditions of the Tet+ treatments have been described in

detail by us previously [43], we focus here on conditions that differed between treatments. The ancestral

strain, grown overnight in MBMS-Tet, was inoculated in multiples of three flow cells, as biofilm

sampling was destructive, and triplicate populations were harvested at each time point. This harvesting

was done at t_0 and after 14 and 28 days (t_{14}, t_{28}) for the Tet+ treatment, and at t_0 and t_{14} for the Tet-

treatment. The Tet- treatment was ended at t_{14} instead of t_{28}, due to rapid plasmid loss in the parallel Tet-

planktonic populations. The harvesting procedures are detailed in Ridenhour *et al.* [43].

The Tet+ and Tet- planktonic populations were initiated by inoculating three replicate test tubes

containing five ml of MBMS-Tet or MBMS with cell suspensions from the triplicate t_0 biofilms of the

Tet+ and Tet- treatments, respectively (Fig. 1). Each of these planktonic populations were then grown in

parallel to the remaining biofilm populations. We used this inoculation approach to control for the level

of diversity already present in the biofilms at t_0, due to evolution that had taken place during the biofilm

establishment phase. Following inoculation, the planktonic populations were serially passaged every 24

($\pm$ 1) hours to obtain approximately 10 generations per day, as described previously [43]. For the Tet+

treatment, the planktonic populations were archived at t_{14} and t_{28}. For the Tet- treatment, the planktonic

populations were archived daily because we did not know how long plasmid-containing (Tet-resistant)

cells would be maintained in these populations. Appropriate dilutions were plated at t_5, t_{10}, and t_{14} on

lysogeny broth agar (LBA) and LBA supplemented with tetracycline (10 mg/l) (LBA$_{Tet}$). When plasmid-

containing cells were no longer detected on LBA$_{Tet}$, the populations archived at previous time points

were plated until we could identify the last time point at which the population still contained plasmid-

bearing cells.

For both biofilm and planktonic populations at each sampling point (see Fig. 1), serial dilutions

of cell suspensions were plated on LBA$_{Tet}$ and plates were incubated overnight. To isolate evolved

clones for plasmid persistence tests and whole-genome sequencing, six resistant clones per replicate

population were purified by restreaking twice and archived. The procedures used for archiving clones

and for plasmid extractions were described by us previously [43].

To determine the effect of large plasmid deletions on plasmid persistence, we also transferred

plasmid DNA extracted from five resistant clones obtained from t_{28} of the Tet+ treatment into the

ancestral *A. baumannii* host. This was done by electroporation, essentially as described previously,

except for washing the cells in cold deionized water[81].

Plasmid fate in biofilms versus planktonic populations in the absence of antibiotics.

To determine the loss of the plasmid in the Tet- biofilms and planktonic populations between t_0

and t_{14}, cell suspensions were diluted and plated on LBA and LBA$_{Tet}$. From these colony-forming unit

(cfu) counts, we determined the total number of culturable cells and the proportion of plasmid-

containing cells. In addition to the number of culturable (=viable) cells, we also determined the total

number of cells in the Tet- biofilms using phase-contrast microscopy and a Petroff-Hausser counting

chamber.

Plasmid persistence assays.

To determine plasmid persistence, we performed triplicate plasmid persistence assays on six

clones per time point per treatment, as described in Ridenhour *et al.* [43]. Briefly, each clone was grown

overnight in MBMS-Tet and used to inoculate three test tubes with MBMS, which were incubated for 24

hours (+/- 1 hour) before passage into new media. This was repeated for a total of 8 days. To estimate

the fraction of plasmid-bearing cells on days 5 and 8 by real-time quantitative PCR (qPCR), cell pellets

obtained after centrifugation of population samples were frozen at -20°C. DNA extractions and qPCR

reactions were performed to determine the copy number of the pB10-encoded replication initiator gene

trfA relative to the number of 16S rRNA copies. These qPCR-based estimates of the

plasmid:chromosome ratio were used as a measure of the fraction of plasmid-bearing cells. This gene

ratio is a proxy for the fraction of plasmid-containing cells but not an exact representation, as there are

five copies of the 16 rRNA gene in *A. baumannii* strain ATCC 17978 and on average three copies of the

plasmid pB10 (Supplementary Figure 1). In addition, this ratio may be slightly affected by slight

variations in copy number (i.e., less than two-fold) of the pB10 and partial chromosomes replicating in

fast-growing cells. However, these variations were small, as Supplementary Fig. 1 shows very limited

variation of the plasmid:chromosome sequence read coverage among all sequenced clones. Given this,

the significant decline of plasmid:chromosome ratios in some populations during the plasmid persistence

assays was not due to changes in plasmid copy number per cell, but due to plasmid loss in these

populations. As described by us previously [43], we used the log-linear model to estimate the rate of

plasmid loss over time.

Plasmid persistence in other hosts.

We extracted ancestral pB10 and one evolved truncated pB10 variant from a clone obtained from

a t_{28} biofilm (Supplementary Table 1) and used electroporation to insert each into three additional

bacterial strains. Plasmid persistence assays for these clones were performed as described above, but for

a total of seven days. They were sampled at days 0, 4, 7, and plasmid presence was determined using

serial dilutions and plate counting on LBA and LBA$_{Tet}$ instead of qPCR.

Plasmid transfer assays

The ability to transfer their plasmid by conjugation was compared for evolved and ancestral

strains as described by us previously, with *E. coli* K12Nal as the recipient [17]. Briefly, 1 ml of donor and

recipient cultures – grown overnight at 37 °C were harvested by centrifugation and resuspended in 100

µl LB broth; 50 µ l of each cell type was then mixed together and spotted onto an LB agar plate for the

actual conjugation and spotted separately as negative controls. Mating plates were incubated at 37 °C.

After overnight incubation, the entire cell mass of each control and conjugation mixture was harvested

and suspended in 500 µl saline, and a dilution series were spread onto the appropriate donor-, recipient-

and transconjugant-selective LB agar media.

Sequencing, and sequence analysis.

For DNA sequencing, total genomic DNA from each clone was extracted from 2mL of an

overnight culture using the GenElute Bacterial Genomic DNA kit (Sigma-Aldrich) and the quality and

integrity of the DNA was assessed on a 1% agarose gel and the concentrations were determined

fluorometrically using Quant-iT™ PicoGreen® dsDNA Reagent (ThermoFisher Scientific, Waltham,

MA, USA) with the TBS-380 Mini-Fluorometer (Turner BioSystems) (Molecular Devices, Sunnyvale,

CA, USA). Samples were submitted to the University of Idaho IBEST Genomic Resources Core facility

for library preparation and whole-genome sequencing using Illumina MiSeq (Illumina, San Diego, CA,

USA) and associated chemistry.

Following sequencing, the sequence data were screened to remove low-quality reads, sequencing

adapters, and duplicate read pairs using the software package htstream

(https://s4hts.github.io/HTStream/). To identify mutations in the genome, cleaned reads were mapped

against the reference sequence for pB10 (AJ564903.1) and *A. baumannii* ATCC 17978

(GCA_000015425.1) using *breseq* v0.35.7 [82]. Prior mapping discrepancy between the reference

sequence of *A. baumannii* ATCC 17978 and our ancestral seed stock were corrected using the gdtools

APPLY command from *breseq*. The command was applied iteratively until no more mutations were

detected between the ancestor reads and the reference sequence. Mutations between strains were

compared using the gdtools COMPARE from *breseq* and the presence of undetected deletions was

manually screened by searching for Missing Coverage Evidence and the presence of corresponding

New-Junction Evidence, indicating reads spanning the area where no reads were mapped. The plasmid

copy numbers shown in Supplementary Figure 1 were estimated using the predicted mean read-depth

coverage given by *breseq* for the plasmids and the chromosome (note that the predicted values are

corrected for the effects of deleted regions).

Statistical analysis.

Statistical analysis of the qPCR data was done using a linear mixed-effect model approach in R

with the nlme package as detailed in Ridenhour *et al.* [43]. Briefly, the log plasmid:chromosome ratio of

each group was predicted using the origin of the clones (i.e., t_0, biofilm, or planktonic population) as

fixed effect and the clone replicate measure as random effect. Each parameter was nested within the day

of the experiment (Days 0, 5, 8) as a continuous variable.

DATA AVAILABILITY

457 All sequencing data pertaining to this project have been made available at the National Center for

458 Biotechnology Information (SRA accession number PRJNA869051). All plasmid persistence data are

459 available at https://github.com/tstal/Metzger-et-al-2022.

CODE AVAILABILITY

462 The code used analyze the plasmid persistence data is available at https://github.com/tstal/Metzger-et-al-

463 2022.

ACKNOWLEDGEMENTS

This project was funded by grant DOD-DM110149. G.A.M. was also in part supported by the

Bioinformatics and Computational Biology Program at the University of Idaho in partnership with the

Institute for Bioinformatics and Evolutionary Studies (now Institute for Interdisciplinary Data Sciences,

IIDS). T.S. and E.M.T. were also partially supported by the National Institute of Allergy and Infectious

Diseases at the National Institute of Health (grant number R01 AI084918). The genome sequencing was

carried out by the IIDS Genomic and Bioinformatic Resources Core, and made possible thanks to NIH

National Institute of General Medical Sciences Award No. P30 GM103324. We would like to thank

Wesley Loftie-Eaton, and Silvia Smith for helpful discussion on the design of this project, and Rachana

Regmi, Andrew Avery, Sean West and Taylor Wilkinson for their roles in data collection.

COMPETING INTERESTS: The authors declare no competing interests.

AUTHOR CONTRIBUTIONS

E.M.T. and L.J. F., conceived the project; G.A.M., E.M.T., L.J. F., B.J.R., and M.L.S., designed the

study; G.A.M., M.F., K.G., and J.M. performed experiments and collected the data; G.A.M. and T.S.

performed the genomic analysis; G.A.M. and B.J.R. performed the statistical analyses; G.A.M., T.S. and

E.M.T. wrote the manuscript; All authors helped revising the manuscript.

REFERENCES

1. Davies, J. Inactivation of antibiotics and the dissemination of resistance genes. *Science* **264**,

375–382 (1994).

2. Schwarz, S. & Johnson, A. P. Transferable resistance to colistin: a new but old threat. *J.*

Antimicrob. Chemother. **71**, 2066–2070 (2016).

3. Mazel, D. & Davies, J. Antibiotic resistance in microbes. *Cell Mol. Life Sci.* **56**, 742–754 (1999).

4. Mathers, A. J. *et al.* Molecular dissection of an outbreak of carbapenem-resistant

Enterobacteriaceae reveals intergenus KPC carbapenemase transmission through a

promiscuous plasmid. *mBio* **2**, e00204-11 (2011).

5. Dionisio, F., Conceição, I. C., Marques, A. C. R., Fernandes, L. & Gordo, I. The evolution of a

conjugative plasmid and its ability to increase bacterial fitness. *Biol. Lett.* **1**, 250–252 (2005).

6. Stewart, F. M. & Levin, B. R. The population biology of bacterial plasmids: a PRIORI conditions

for the existence of conjugationally transmitted factors. *Genetics* **87**, 209–228 (1977).

7. Bergstrom, C. T., Lipsitch, M. & Levin, B. R. Natural selection, infectious transfer and the

existence conditions for bacterial plasmids. *Genetics* **155**, 1505–1519 (2000).

8. Baltrus, D. A. Exploring the costs of horizontal gene transfer. *Trends. Ecol. Evol.* **28**, 489–495

(2013).

9. San Millan, A. & MacLean, R. C. Fitness costs of plasmids: a limit to plasmid transmission.

Microbiol Spectr. **5**, (2017).

10. Gama, J. A., Zilhão, R. & Dionisio, F. Plasmid interactions can improve plasmid persistence in

bacterial populations. *Front. Microbiol.* **11**, 2033 (2020).

11. Brockhurst, M. A. & Harrison, E. Ecological and evolutionary solutions to the plasmid paradox.

Trends Microbiol. S0966-842X(21)00267–5 (2021) doi:10.1016/j.tim.2021.11.001.

12. Bouma, J. E. & Lenski, R. E. Evolution of a bacteria/plasmid association. *Nature* **335**, 351–352

(1988).

13. De Gelder, L., Williams, J. J., Ponciano, J. M., Sota, M. & Top, E. M. Adaptive plasmid evolution

results in host-range expansion of a broad-host-range plasmid. *Genetics* **178**, 2179–2190

(2008).

14. Sota, M. *et al.* Shifts in the host range of a promiscuous plasmid through parallel evolution of

its replication initiation protein. *ISME J.* **4**, 1568–1580 (2010).

15. San Millan, A. *et al.* Positive selection and compensatory adaptation interact to stabilize non-

transmissible plasmids. *Nat. Commun.* **5**, 5208 (2014).

16. Harrison, E., Guymer, D., Spiers, A. J., Paterson, S. & Brockhurst, M. A. Parallel compensatory

evolution stabilizes plasmids across the parasitism-mutualism continuum. *Curr. Biol.* **25**,

2034–2039 (2015).

17. Loftie-Eaton, W. *et al.* Compensatory mutations improve general permissiveness to antibiotic

resistance plasmids. *Nat. Ecol. Evol.* **1**, 1354–1363 (2017).

18. Stalder, T. *et al.* Emerging patterns of plasmid-host coevolution that stabilize antibiotic

resistance. *Sci. Rep.* **7**, 4853 (2017).

19. Wein, T., Hülter, N. F., Mizrahi, I. & Dagan, T. Emergence of plasmid stability under non-

selective conditions maintains antibiotic resistance. *Nat. Commun.* **10**, 2595 (2019).

20. Lopatkin, A. J. *et al.* Persistence and reversal of plasmid-mediated antibiotic resistance. *Nat.*

Commun. **8**, 1689 (2017).

21. Porse, A., Schønning, K., Munck, C. & Sommer, M. O. A. Survival and evolution of a large

multidrug resistance plasmid in new clinical bacterial hosts. *Mol. Biol. Evol.* **33**, 2860–2873

(2016).

22. Hall, J. P. J. *et al.* Plasmid fitness costs are caused by specific genetic conflicts enabling

resolution by compensatory mutation. *PLoS Biol.* **19**, e3001225 (2021).

23. Kloos, J., Gama, J. A., Hegstad, J., Samuelsen, Ø. & Johnsen, P. J. Piggybacking on niche adaptation

improves the maintenance of multidrug-resistance plasmids. *Mol. Biol. Evol.* **38**, 3188–3201

(2021).

24. Kawano, H. *et al.* A novel small RNA on the *Pseudomonas putida* KT2440 chromosome is

involved in the fitness cost imposed by IncP-1 plasmid RP4. *Front. Microbiol.* **11**, (2020).

25. Vlamakis, H., Chai, Y., Beauregard, P., Losick, R. & Kolter, R. Sticking together: building a biofilm

the *Bacillus subtilis* way. *Nat. Rev. Microbiol.* **11**, 157–168 (2013).

26. Flemming, H.-C. & Wuertz, S. Bacteria and archaea on Earth and their abundance in biofilms.

Nat. Rev. Microbiol. **17**, 247–260 (2019).

27. Costerton, J. W., Stewart, P. S. & Greenberg, E. P. Bacterial biofilms: a common cause of

persistent infections. *Science* **284**, 1318–1322 (1999).

28. Parsek, M. R. & Singh, P. K. Bacterial biofilms: an emerging link to disease pathogenesis. *Annu.*

Rev. Microbiol. **57**, 677–701 (2003).

29. Boles, B. R., Thoendel, M. & Singh, P. K. Self-generated diversity produces "insurance effects" in

biofilm communities. *Proc. Natl. Acad. Sci. U.S.A.* **101**, 16630–16635 (2004).

30. Korona, R., Nakatsu, C. H., Forney, L. J. & Lenski, R. E. Evidence for multiple adaptive peaks

from populations of bacteria evolving in a structured habitat. *Proc. Natl. Acad. Sci. U.S.A.* **91**,

9037–9041 (1994).

31. Ponciano, J. M., La, H.-J., Joyce, P. & Forney, L. J. Evolution of diversity in spatially structured

Escherichia coli populations. *Appl. Environ. Microbiol.* **75**, 6047–6054 (2009).

32. Rainey, P. B. & Travisano, M. Adaptive radiation in a heterogeneous environment. *Nature* **394**,

69–72 (1998).

33. Tyerman, J. G., Ponciano, J. M., Joyce, P., Forney, L. J. & Harmon, L. J. The evolution of antibiotic

susceptibility and resistance during the formation of *Escherichia coli* biofilms in the absence of

antibiotics. *BMC Evol. Biol.* **13**, 22 (2013).

34. Steenackers, H. P., Parijs, I., Foster, K. R. & Vanderleyden, J. Experimental evolution in biofilm

populations. *FEMS Microbiol. Rev.* **40**, 373–397 (2016).

35. Santos-Lopez, A., Marshall, C. W., Scribner, M. R., Snyder, D. J. & Cooper, V. S. Evolutionary

pathways to antibiotic resistance are dependent upon environmental structure and bacterial

lifestyle. *eLife* **8**, e47612 (2019).

36. Martin, M., Hölscher, T., Dragoš, A., Cooper, V. S. & Kovács, Á. T. Laboratory evolution of

microbial interactions in bacterial biofilms. *J. Bacteriol.* **198**, 2564–2571 (2016).

37. France, M. T., Ridenhour, B. J. & Forney, L. J. Effects of spatial structure and reduced growth

rates on evolution in bacterial populations. in *Molecular Mechanisms of Microbial Evolution* (ed.

Rampelotto, P. H.) 175–197 (Springer International Publishing, 2018). doi:10.1007/978-3-

319-69078-0_8.

38. Gordo, I. & Campos, P. R. A. Adaptive evolution in a spatially structured asexual population.

Genetica **127**, 217 (2006).

39. Habets, M. G. J. L., Czárán, T., Hoekstra, R. F. & de Visser, J. A. G. M. Spatial structure inhibits the

rate of invasion of beneficial mutations in asexual populations. *Proc. R. Soc. B: Biol.* **274**, 2139–

2143 (2007).

40. Nahum, J. R. *et al.* A tortoise-hare pattern seen in adapting structured and unstructured

populations suggests a rugged fitness landscape in bacteria. *Proc. Natl. Acad. Sci. U.S.A.* **112**,

7530–7535 (2015).

41. France, M. T. & Forney, L. J. The relationship between spatial structure and the maintenance of

diversity in microbial populations. *Am. Nat* **193**, 503–513 (2019).

42. Coenye, T., Bové, M. & Bjarnsholt, T. Biofilm antimicrobial susceptibility through an

experimental evolutionary lens. *NPJ Biofilms Microbiomes* **8**, 82 (2022).

43. Ridenhour, B. J. *et al.* Persistence of antibiotic resistance plasmids in bacterial biofilms. *Evol.*

Appl. **10**, 640–647 (2017).

44. Stalder, T. *et al.* Evolving populations in biofilms contain more persistent plasmids. *Mol. Biol.*

Evol. **37**, 1563–1576 (2020).

45. Galimand, M. *et al.* Multidrug resistance in *Yersinia pestis* mediated by a transferable plasmid.

N. Engl. J. Med. **337**, 677–680 (1997).

46. Gay, K. *et al.* Plasmid-mediated quinolone resistance in non-Typhi serotypes of *Salmonella*

enterica. *Clin. Infect. Dis.* **43**, 297–304 (2006).

47. Richet, H. & Fournier, P. E. Nosocomial infections caused by *Acinetobacter baumannii*: a major

threat worldwide. *Infect. Control Hosp. Epidemiol.* **27**, 645–646 (2006).

48. Sunenshine, R. H. *et al.* Multidrug-resistant *Acinetobacter* infection mortality rate and length of

hospitalization. *Emerg. Infect. Dis.* **13**, 97–103 (2007).

49. Perez, S. *et al.* Increase in hospital-acquired carbapenem-resistant *Acinetobacter baumannii*

infection and colonization in an acute care hospital during a surge in COVID-19 Admissions —

New Jersey, February–July 2020. *MMWR Morb. Mortal Wkly Rep.* **69**, 1827–1831 (2020).

595 50. Ma, F. *et al.* Identification of a novel plasmid carrying *mcr-4.3* in an *Acinetobacter baumannii*

596  strain in China. *Antimicrob. Agents Chemother.* **63**, e00133-19 (2019).

597 51. Ibrahim, S., Al-Saryi, N., Al-Kadmy, I. M. S. & Aziz, S. N. Multidrug-resistant *Acinetobacter*

598  *baumannii* as an emerging concern in hospitals. *Mol. Biol. Rep.* **48**, 6987–6998 (2021).

599 52. Karakonstantis, S., Ioannou, P., Samonis, G. & Kofteridis, D. P. Systematic review of

600  antimicrobial combination options for pandrug-resistant *Acinetobacter baumannii*. *Antibiotics*

601  **10**, 1344 (2021).

602 53. Schlüter, A. *et al.* The 64 508 bp IncP-1beta antibiotic multiresistance plasmid pB10 isolated

603  from a waste-water treatment plant provides evidence for recombination between members

604  of different branches of the IncP-1beta group. *Microbiology* **149**, 3139–3153 (2003).

605 54. Albertini, A. M., Hofer, M., Calos, M. P. & Miller, J. H. On the formation of spontaneous deletions:

606  The importance of short sequence homologies in the generation of large deletions. *Cell* **29**,

607  319–328 (1982).

608 55. De Gelder, L., Ponciano, J. M., Joyce, P. & Top, E. M. Stability of a promiscuous plasmid in

609  different hosts: no guarantee for a long-term relationship. *Microbiology* **153**, 452–463 (2007).

610 56. Røder, H. L. *et al.* Biofilms can act as plasmid reserves in the absence of plasmid specific

611  selection. *npj Biofilms Microbiomes* **7**, 1–6 (2021).

612 57. Ma, H., Katzenmeyer, K. N. & Bryers, J. D. Non-invasive in situ monitoring and quantification of

613  TOL plasmid segregational loss within *Pseudomonas putida* biofilms. *Biotechnol. Bioeng.* **110**,

614  2949–2958 (2013).

615 58. Turner, P. E., Cooper, V. S. & Lenski, R. E. Tradeoff between horizontal and vertical modes of

616  transmission in bacterial plasmids. *Evolution* **52**, 315–329 (1998).

59. Dahlberg, C. & Chao, L. Amelioration of the cost of conjugative plasmid carriage in *Eschericha*

coli K12. *Genetics* **165**, 1641–1649 (2003).

60. Starikova, I. *et al.* A trade-off between the fitness cost of functional integrases and long-term

stability of integrons. *PLoS Pathog.* **8**, e1003043 (2012).

61. Lacotte, Y., Ploy, M.-C. & Raherison, S. Class 1 integrons are low-cost structures in *Escherichia*

coli. *ISME J.* **11**, 1535–1544 (2017).

62. Couvé-Deacon, E. *et al.* Class 1 integrons in *Acinetobacter baumannii*: a weak expression of

gene cassettes to counterbalance the lack of LexA-driven integrase repression. *Int. J.*

Antimicrob. Agents **53**, 491–499 (2019).

63. Poulsen, L. K., Ballard, G. & Stahl, D. A. Use of rRNA fluorescence in situ hybridization for

measuring the activity of single cells in young and established biofilms. *Appl. Environ.*

Microbiol. **59**, 1354–1360 (1993).

64. Wang, L., Fan, D., Chen, W. & Terentjev, E. M. Bacterial growth, detachment and cell size control

on polyethylene terephthalate surfaces. *Sci. Rep.* **5**, 15159 (2015).

65. Molin, S. & Tolker-Nielsen, T. Gene transfer occurs with enhanced efficiency in biofilms and

induces enhanced stabilisation of the biofilm structure. *Curr. Opin. Biotechnol.* **14**, 255–261

(2003).

66. Hausner, M. & Wuertz, S. High rates of conjugation in bacterial biofilms as determined by

quantitative *in situ* analysis. *Appl. Environ. Microbiol.* **65**, 3710–3713 (1999).

67. Krol, J. E. *et al.* Increased transfer of a multidrug resistance plasmid in *Escherichia coli* biofilms

at the air-liquid interface. *Appl. Environ. Microbiol.* **77**, 5079–5088 (2011).

68. Stalder, T. & Top, E. Plasmid transfer in biofilms: a perspective on limitations and

opportunities. *npj Biofilms Microbiomes* **2**, (2016).

69. Zwanzig, M. *et al.* Mobile compensatory mutations promote plasmid survival. *mSystems* **4**,

e00186-18 (2019).

70. Zhong, X., Krol, J. E., Top, E. M. & Krone, S. M. Accounting for mating pair formation in plasmid

population dynamics. *J. Theor. Biol.* **262**, 711–719 (2010).

71. Bradley, D. E., Taylor, D. E. & Cohen, D. R. Specification of surface mating systems among

conjugative drug resistance plasmids in *Escherichia coli* K-12. *J. Bacteriol.* **143**, 1466–1470

(1980).

72. Ghigo, J. M. Natural conjugative plasmids induce bacterial biofilm development. *Nature* **412**,

442–445 (2001).

73. Vries, J. de & Wackernagel, W. Integration of foreign DNA during natural transformation of

Acinetobacter sp. by homology-facilitated illegitimate recombination. *PNAS* **99**, 2094–2099

(2002).

74. De Gelder, L. *et al.* Combining mathematical models and statistical methods to understand and

predict the dynamics of antibiotic-sensitive mutants in a population of resistant bacteria

during experimental evolution. *Genetics* **168**, 1131–1144 (2004).

75. Senol, E. *Stenotrophomonas maltophilia*: the significance and role as a nosocomial pathogen. *J.*

Hosp. Infect. **57**, 1–7 (2004).

76. Hall, J. P. J., Wright, R. C. T., Guymer, D., Harrison, E. & Brockhurst, M. A. Y. 2020. Extremely fast

amelioration of plasmid fitness costs by multiple functionally diverse pathways. *Microbiology*

166, 56–62 (2019).

77. Smith, M. G. *et al.* New insights into *Acinetobacter baumannii* pathogenesis revealed by high-

density pyrosequencing and transposon mutagenesis. *Genes Dev.* **21**, 601–614 (2007).

78. Weber, B. S., Ly, P. M., Irwin, J. N., Pukatzki, S. & Feldman, M. F. A multidrug resistance plasmid

contains the molecular switch for type VI secretion in *Acinetobacter baumannii*. *Proc. Natl.*

Acad. Sci. U S A **112**, 9442–9447 (2015).

79. Loftie-Eaton, W. *et al.* Draft Genome Sequence of *Pseudomonas* sp. nov. H2. *Genome Announc.* **3**,

e00241-15 (2015).

80. Hunter, S. S. *et al.* Draft genome sequence of *Pseudomonas moraviensis* R28-S. *Genome Announc.*

2, (2014).

81. Yano, H., Deckert, G. E., Rogers, L. M. & Top, E. M. Roles of long and short replication initiation

proteins in the fate of IncP-1 plasmids. *J. Bacteriol.* **194**, 1533–1543 (2012).

82. Deatherage, D. E. & Barrick, J. E. Identification of mutations in laboratory-evolved microbes

from next-generation sequencing data using breseq. *Methods Mol. Biol.* **1151**, 165–188 (2014).

 FIGURE LEGENDS

Figure 1: Timeline of the evolution experiments. Red represents time periods when media contained

tetracycline (Tet+ treatment); blue represents the absence of antibiotics (Tet- treatment). Large ticks

represent inoculation and harvest/sampling events. Four days prior to initiation of the evolution

experiments (-4) the biofilm flow cells were inoculated with the ancestral strain. Four days later, at day

0 (t_0), the first set of randomly selected flow cells (n=3) were harvested. A subsample of the cell

suspensions from each of these replicates was used to inoculate each of the three planktonic populations

(grey curved arrow). Small tick marks indicate the daily serial passage of planktonic populations.

Asterisks on days 6 and 10 in the timeline of the Tet- planktonic populations indicate the last day at

which plasmid-containing cells were detected in two of the three populations due to rapid plasmid loss;

these time points were the final sampling points for these two populations (t_6 and t_{10}). The Tet-

experiment was terminated after 14 days and the Tet+ experiment after 28. For more details on the

actual protocol, see Materials and Methods and Ridenhour et al. [43].

Figure 2: Plasmid persistence shown as the estimated ratio of *trfA*/16S rRNA genes over a period of

eight days in liquid serial batch cultures in the absence of selection; (A) the ancestor used to inoculate

the flow cells at t_{-4} compared to clones isolated from all the biofilms harvested after four days of growth

with tetracycline, t_0, showing similar plasmid persistence dynamics (Tet+ and Tet-: populations treated

with and without tetracycline after t_0); (B) clones isolated from the Tet- populations, showing a drastic

increase in plasmid persistence after evolution in planktonic but not in biofilm populations (note that

two of the three planktonic populations no longer contained detectable plasmids after day 6 and 10,

respectively, requiring analysis on these days instead of t_{14} – the final sampling times are therefore

referred to as t_{final}); (C) clones isolated at t_0 and t_{14} of the Tet+ evolution experiment, showing an

increase in plasmid persistence over time for all populations. Lines are the output of the log-linear model

for each group. * denotes a significant difference (t-test p-value <0.05) between the groups pointed out

by the brackets: planktonic, t_{final} versus t_0 and biofilm t_{14} versus t_0. The plasmid persistence data from t_0

in the Tet+ experiment were reported by us previously [43].

Figure 3: Maps of evolved plasmids with major segment functions identified, showing all plasmid

mutations observed in each sequenced clone evolved under different treatments (Tet+ and Tet-) in

different environments (biofilm and planktonic populations). Evolved clones are grouped by treatment,

environment, and time point, as shown by different colors and shades and identified in the open wedge

of the circle. The plasmid genome of each clone is shown as a single band – at each time point, 18

clones were analyzed per treatment and environment (six clones from each triplicate population). Tet+,

Tet-: evolved with and without Tet. Black areas in the plasmid maps represent deletions (note that small

indels are shown as small spots) and the dots represent SNPs; see Supplementary Table 1 for details.

Note that the tetracycline resistance operon is located in the 'accessory region'.

Figure 4: Persistence of plasmids evolved in Tet+ biofilms and planktonic populations tested in (A) their

co-evolved hosts (92 full-length and 49 truncated plasmids), and (B) in the ancestral host (2 full-length

and 4 truncated plasmids). In both panels the ancestral plasmid was tested in the ancestral host. Plasmid

persistence is shown as the estimated ratio of $trfA$/16S rRNA genes over time. Lines are the results of

the log-linear model for each group. * denotes a significant difference (t-test p-value was <0.05)

between the groups indicated by the brackets: truncated plasmids versus ancestral plasmid and versus

full-length plasmids.

Figure 5: Plasmid persistence of ancestral plasmid pB10$_{Anc}$ and truncated plasmid pB10$_{ev}$ in three

additional hosts, as determined by plate counting after 8 days, showing that the plasmid deletion was

only beneficial to persistence in *Stenotrophomonas maltophilia*. (P= *Pseudomonas*). Boxes represent the

median and the first and third quartiles, and the whiskers the lower and upper 1.5 interquartile range.

Inoculate
N=6

Harvest
N=3

Harvest
N=3

Biofilm

Tet-

Inoculate
N=3

*

*

Sample
N=3

Planktonic

Inoculate
N=9

Harvest
N=3

Harvest
N=3

Harvest
N=3

Biofilm

Tet+

Inoculate
N=3

Sample
N=3

Sample
N=3

Planktonic

-4   0   14   28

Time (days)

Replication

Tet-

Tet+

Accessory region

Biofilm t14
Planktonic tfinal
t0
Biofilm t28
Biofilm t14
Planktonic t28
Planktonic t14
t0

Mating-Pair-Formation (trb)

Class 1 Integron

Plasmid Regulation & Inheritance

Conjugative Transfer (tra)

Table 1: Cell counts as well as absolute counts and fractions of plasmid-containing (P+) cells for each replicate biofilm or planktonic population of the Tet- treatment.

	Total cells (per mL)[1]	Total viable cells (per mL) [2]	P+ cells (per mL)	Fraction P+ cells [3]
_t_0				
Rep 1	9.6×10^9	6.1×10^6	5.9×10^6	0.98
Rep 2	3.9×10^9	4.7×10^6	3.8×10^6	0.82
Rep 3	3.5×10^9	3.4×10^6	3.4×10^6	1.00
t0 Avg	_5.6×10^9_	_4.7×10^6_	_4.4×10^6_	_0.93_
_t_14 Biofilm				
Rep 1	1.3×10^{11}	2.9×10^6	5.5×10^5	0.19
Rep 2	1.3×10^{10}	3.1×10^6	3.6×10^5	0.11
Rep 3	1.1×10^{10}	1.8×10^6	1.3×10^5	0.07
Biofilm Avg	_5.1×10^{10}_	_2.6×10^6_	_3.5×10^5_	_0.12_
_t_14 Planktonic				
Rep 1	n/a	2.0×10^8	3.2×10^4	0.00016
Rep 2	n/a	1.6×10^8	<1	0
Rep 3	n/a	2.0×10^8	<1	0
Planktonic Avg	_n/a_	_1.9×10^8_	_1.1×10^4_	_0.00005_

[1] Total cell counts were obtained by microscopy; [2] total viable cell counts correspond to colony forming units (cfu); [3] P+ fractions were calculated as the ratio of Tet^R over total cfu counts, obtained by plating on selective and non-selective media. Note that total cell counts estimated by microscopy were only determined for biofilms. n/a: count not performed.