Incorporating selection pressures into risk predictions of pesticide resistance evolution in arthropod pests

Joshua A. Thia [1,§], James Maino [1,2], Alicia Kelly [1], Ary A. Hoffmann [1], Paul A. Umina [1,2]

[1] Bio21 Institute, School of BioSciences, University of Melbourne, Parkville, VIC, Australia

[2] Cesar Australia, Brunswick, VIC, Australia

[§] Corresponding author: email, joshua.thia@unimelb.edu.au and josh.thia@live.com; phone, +61390358667

KEYWORDS

Bayesian inference; ecological traits; pesticide product registrations; predictive modelling; resistance management

PREPRINT

An earlier version of this manuscript has been previously posted on BioRXiv under a CC-BY-NC-ND 4.0 International license, DOI: 10.1101/2022.04.18.488707.

ABSTRACT

Introduction — Chemical resistance in pest organisms threatens global food security and human health, yet resistance issues are mostly dealt with reactively. Predictive models of resistance risk provide opportunity for field practitioners to implement proactive management but require knowledge of variables that drive the evolution of resistance. Despite the importance of chemical selection pressure on resistance evolution, poor availability of chemical usage data has limited the use of a general multi-species measure of selection pressure in predictive models.

Results — We demonstrate the use of pesticide product registrations as a predictor of resistance status and potential proxy of chemical selection pressure. Pesticide product registrations were obtained for 427 USA and 209 Australian agricultural arthropod pests, for 42 and 39 chemical Mode of Action groups, respectively. We constructed Bayesian logistic regressions of resistance status as a function of the number of pesticide product registrations and two ecological traits, phagy and voltinism. Our models were well-supported, and we demonstrated considerable power to discriminate between resistant and susceptible observation in both USA and Australian species sets using cross-validation. Importantly, we observed strong support for a positive association between pesticide products and resistance status.

Conclusions — Our results suggest that pesticide product registrations can be used to approximate pesticide usage and thus selection pressure for resistance in agro-ecosystems, which combined with ecological information can be used to predict the risk of resistance in agricultural pests. Our work expands the horizon for proactive management via modeling of specific species-by-chemical resistance risks using publicly available data, with broad applicability across agricultural pests, including arthropods, weeds and fungi.

BACKGROUND

Decades of research into the evolution of pesticide resistance and its molecular basis have contributed to a rich understanding of the dynamics of resistance development, which has benefited management strategies in agricultural and human health contexts (Crow 1957; Georghiou and Taylor 1986; McKenzie and Batterham 1994; Hawkins et al. 2019). But despite our understanding of pesticide resistance evolution, resistance remains a prevalent global issue for pest management and food security (Pimentel et al. 1992; Liu 2015; Sparks and Nauen 2015; Gould et al.

2018; Pu et al. 2020). Indeed, notwithstanding the best efforts of scientists, field practitioners and multi-national chemical companies, entire chemical Mode of Action (MoA) groups have become ineffective against certain pest groups due to widespread resistance (Nauen 2007; Bass et al. 2015; Horowitz et al. 2020).

A major challenge in managing pesticide resistance is the ongoing importance of chemical control, which places populations under repeated selective pressure to evolve resistance. Pest management strategies are often *reactive*, only being implemented once resistance has evolved and resistant genotypes have reached a detectable frequency in field populations (Tabashnik et al. 2014; Sparks and Nauen 2015; Umina et al. 2019). More *proactive* strategies would routinely assess resistance risks prior to control failures developing. Predictions about where and when a species might exhibit high risks of resistance evolution could be used to prioritize research investment and target in-field monitoring efforts. To achieve this, a comprehensive understanding of factors that increase the risk of resistance evolution is required.

One of the most important factors expected to drive rates of pesticide resistance evolution is selection pressure (Crow 1957; Georghiou and Taylor 1977; McKenzie and Batterham 1994). However, quantifying the magnitude of selection in pest populations is challenging in agro-ecosystems. Many countries lack policy for reporting and documenting pesticide usage, making regulation and risk assessment of pesticide usage difficult (Handford et al. 2015; Sharma et al. 2019). Studies investing in the co-collection of pesticide usage and phenotypic resistance data have provided valuable demonstrations of the link between on-farm practice and the evolution of pesticide resistance. For example, studies of weeds from the United Kingdom show that the spatial distribution of herbicide resistance is linked to historical application rates and frequency of herbicide applications (Evans et al. 2016; Hicks et al. 2018). Yet collecting such data requires considerable investment, especially across large spatial scales, and is limited in the scope of species and pesticides that can be assessed. Proactive management would therefore benefit from measures that can approximate pesticide usage (selection pressure) in agro-ecosystems across multiple pest species and chemical MoAs.

Selection alone is not the only factor influencing the rate of pesticide resistance evolution. Species-specific ecological traits or evolutionary constraints might modulate the response of selection of pest populations (Georghiou and Taylor 1977; Rosenheim and Tabashnik 1990). In arthropod pests (insects and mites), phagy and voltinism are two ecological traits with demonstrated importance in pesticide resistance evolution. Arthropods that are more polyphagous (broader diets) might be

preadapted with xenobiotic detoxification machinery that can be co-opted into pesticide detoxification (Hardy et al. 2018; Rane et al. 2019; Crossley et al. 2020). Greater voltinism (generations per year) may positively affect resistance evolution through population genetic or population growth effects (Robertson 1960; Georghiou and Taylor 1986; Li et al. 1987; Weber 1990; Baer et al. 2007). There may also be curvilinear effects of these traits, such that the effect of increasing diet breadth or generations per year is not simply additive (Rosenheim and Tabashnik 1991; Hardy et al. 2018; Crossley et al. 2020). There may also be evolutionary constraints that lead to significant phylogenetic differences among arthropod lineages in their capacity to evolve resistance (Brevik et al. 2018; Hardy et al. 2018). Accounting for ecology and evolutionary factors is therefore necessary in predicting the risk of resistance evolution given an inferred level of pesticide usage.

In this present work, we consider the need to derive inferences of selection pressure in agro-ecosystems in the absence of well-maintained records of pesticide usage. We focus on arthropod pests and their resistances to pesticides (insecticides and acaricides). We have previously demonstrated that pesticide product registrations are positively correlated with pesticide usage agro-ecosystems from the USA and Australia (Maino et al., in press). Here, we extend these findings by demonstrating a positive association between pesticide product registrations and pesticide resistance in agricultural arthropod pests from both these countries. Our results support the notion that pesticide product registrations might be a generalisable selection pressure in agro-ecosystems. These findings demonstrate how pesticide product registrations can be combined with ecological information to develop risk predictions of resistance evolution and guide proactive resistance management for specific species and chemical MoAs.

RESULTS

DATASET COMPILATION

We compiled two datasets comprising agricultural arthropod pests from the USA and Australia, their ecological traits (phagy and voltinism), the number of pesticide products registered against them for different chemical MoAs, and their resistance statuses to those chemical MoAs. Our USA species set included 427 species, 81 families, and 10 orders, and for the Australian species set, there were 209 species, 65 families, and 12 orders. The taxonomic composition of total species was

similar between countries (Figure 1). In the USA, Lepidoptera (32%), Hemiptera (29%), Coleoptera (19%), and Trombidiformes (9%) made up the bulk of the total pest species, which was very similar to the Australian species set (30%, 29%, 18% and 9%, respectively). The taxonomic composition of species with resistance (to at least one chemical MoA largely paralleled the total species set (Figure 1). In the USA, Lepidoptera (32%), Coleoptera (26%), Hemiptera (25%), and Trombidiformes (9%) made up the bulk of resistant pest species (n = 76). In Australia, resistant pests (n = 24) included species from Lepidoptera (46%), Hemiptera (21%), Trombidiformes (17%), Thysanoptera (8%), and Coleoptera (8%).

[FIGURE 1]

Values of phagy and voltinism were sourced from previous pest community studies (Rosenheim et al. 1996; Hardy et al. 2018; Crossley et al. 2020) or were sourced from a literature search. However, many species required imputation. Of the USA species, 30.9% (n = 132) had at least one ecological trait imputed from a literature search, whereas for Australian species, 56.0% (n = 117) of species had at least one ecological trait imputed. The USA species set contained 42 chemical MoAs, and the Australian species set contained 39 chemical MoAs. Our working datasets were structured with respect to species–MoA observations, where the number of pesticide product registrations for each unique chemical MoA was assigned to each pest species. There were n = 17,934 species–chemical MoA observations in the USA species set and n = 8,151 for the Australian species set.

PREDICTIVE MODELS OF RESISTANCE

Our 'full' model of resistance fitted resistance status of species to different chemical MoAs (a binary response) as a function of the fixed effects of pesticide product registrations ($\beta_{Products}$), the first and second polynomials of phagy (β_{Phagy^1}, β_{Phagy^2}) and voltinism ($\beta_{Voltinism^1}$, $\beta_{Voltinism^2}$), and the random effects of taxonomy and chemical MoA. We also fitted a 'products' model that only included the fixed effect of products and the random effects of taxonomy and chemical MoA, and a 'null' model that only fitted the random effects of taxonomy and chemical MoA. Our models were fitted using a Bayesian framework and the likelihood of the 'full' and 'products' models were tested by comparison to the 'null' model.

We tested whether ecological trait imputation affected the predictions of the 'full' model by comparing predictions models using imputed values of phagy and voltinism (all species) with models using non-imputed values (excluding species with missing data). Because predicted risk scores using the 'full' model using imputed and non-imputed ecological traits exhibited very high

correlations ($r > 0.98$; Figure S1), all our interpretations consider those models fitted to datasets with imputed ecological traits. Distributions for fixed effects are illustrated in Figure 2. Fitted terms for the 'full' models are tabulated in Table 1 and illustrated in Figure 2.

[FIGURE 2]

[TABLE 1]

We found a positive association between pesticide product registrations and resistance status (Figure 2 & 3). For a given chemical MoA, species with a larger number of pesticide product registrations were more likely to be resistant to that chemical MoA than those with fewer pesticide product registrations. This effect was consistent between species sets from the USA, $\beta_{Products} = 0.68$, and Australia, $\beta_{Products} = 1.09$, with overlapping Bayesian 95% credible intervals (Table 1; Figure 3). Consistent with other studies, we found a positive second order polynomial for phagy, and a negative second order polynomial for voltinism, suggesting that resistance evolution is minimal at intermediate levels of phagy, and maximised at intermediate levels of voltinism (Hardy et al. 2018; Crossley et al. 2020). We observed consistent estimates in the second order polynomials for both countries, with overlapping Bayesian 95% credible intervals: for USA species, $\beta_{Phagy^2} = 79.30$ and $\beta_{Voltinism^2} = -100.90$; and for Australian species, $\beta_{Phagy^2} = 199.58$ and $\beta_{Voltinism^2} = -179.64$ (Table 1; Figure 3). Diagnostic statistics indicated that we reliably sampled the posterior distributions for these fixed effect coefficients (Table 1): the $\hat{R}$ statistics were equal to 1.00, indicating convergence, and ESS values were >1,000, indicating many uncorrelated draws were obtained (Muth et al. 2018).

[FIGURE 3]

Our 'full' model including pesticide product registrations and ecological traits were substantially more likely (log-Bayes factor $\gg$ 1.1) to explain resistance status of a species to different chemical MoAs relative to 'null' models excluding these factors. For the USA species set, the 'full' model had an estimated log-Bayes factors of 88.24 and a Bayesian R^2 of 0.33. For the Australian species, the 'full' model had an estimated log-Bayes factors of 83.04 and a Bayesian R^2 of 0.45. Additionally, the 'products' model had a log-Bayes factor of 45.85 for the USA species set, with a Bayesian R^2 of 0.31. For the Australian species set, the 'products' model had a log-Bayes factor of 29.85 and a Bayesian R^2 of 0.34. Therefore, models fitting only pesticide products as a fixed effect were substantially more likely to explain resistance status of a species to different chemical MoAs relative to the 'null' models.

MODEL GENERALITY

K-fold cross-validation was used to test the predictive power of the 'full' and 'products' model using 100 training–testing species set partitions. In each partition, the training species were used to construct the model, predictions were made on the testing species, and an AUC was calculated. Predictive power was similar between models and countries (Figure 4). For the USA species set, the mean AUC scores were 0.81 (±0.004 SEM) for the 'full' model and 0.80 (±0.003 SEM) for the 'products' model. For the Australian species set, the mean AUC scores were 0.82 (±0.007 SEM) for the 'full' model and 0.79 (±0.006 SEM) for the 'products' model. AUC scores were obtained using only the estimated fixed effect coefficients and not the estimated random intercepts for taxonomy or chemical MoA group. Hence, these *K*-fold cross-validation analyses show that resistance can be predicted well without controlling for ecology, taxonomy, and chemical MoA group, because species with resistance to a chemical MoA group tend to have greater product registrations than those susceptible to that chemical MoA group.

[FIGURE 4]

DISCUSSION

Identifying key predictors of pesticide resistance and pinpointing high-risk species would unlock new possibilities for making proactive, data-driven risk management decisions. Despite the importance of selection for pesticide resistance, we are unaware of attempts to identify a multi-species measure of selection to different chemical MoAs. Here, we successfully used pesticide product registrations as a predictor of resistance in agricultural arthropod pests from the USA and Australia. In a parallel study, we have found that pesticide product registrations are positively associated with pesticide usage in both the USA and Australia (Maino et al., in press). We have expanded those results in this present work by further demonstrating that pesticide product registrations are also positively associated with resistance status of pests to specific chemical MoAs. These joint observations support the notion that pesticide product registrations may be a useful, general proxy for selection pressures in agro-ecosystems where pesticide usage is poorly recorded. Note that we do not imply to have estimated selection coefficients themselves, but instead have a defined a covariate for modelling purposes that approximates selection pressure.

We hypothesize that the positive association between pesticide product registrations and resistance status arises from the accumulation of processes that generally increase the selection pressure on pest populations. The risk of pesticide resistance in any pest is expected to be relative to the magnitude and frequency of selection (Georghiou and Taylor 1977; Evans et al. 2016; Hicks et al. 2018; Hawkins et al. 2019). Additionally, for a chemical MoA to be extensively used, it must be accessible, affordable, and in demand (Sparks and Nauen 2015; Umina et al. 2019). Chemical databases may therefore capture trends in both the availability and application intensity of chemicals in the field, which translate into greater selection pressures in the field, and therefore risk of resistance evolution. While other studies have demonstrated the importance of ecology and taxonomy in determining the number of resistances and the rate of pesticide resistance evolution across many diverse arthropod pests (Rosenheim et al. 1996; Brevik et al. 2018; Hardy et al. 2018; Crossley et al. 2020), ours is the first to implement a proxy for selection intensity for many species and chemical MoA combinations, at least to the best of our knowledge. To a degree, the use of a "severity index" (number of research papers) for pest species employed in other studies may partially capture the selection pressure experienced by pests (Hardy et al. 2018; Crossley et al. 2020). However, this severity index does not extend to specific chemical MoAs and more likely describes an inherent adaptive potential of a pest to evolve resistance, versus a proxy of selection pressure.

We see the application of pesticide product registrations to proactive management as a two-step procedure. First, as we have done here, a model fitting resistance status of species to chemical MoA groups needs to be constructed. Second, after fitting this model, species with high predicted risk scores for a chemical MoA, but with no documented resistances to the chemical MoA, can then be identified as potential candidates for further inquiry. We illustrate such predicted risks for the top ten resistant pests (those with the most resistances to different chemical MoAs) in the USA (Figure 5a) and Australian (Figure 5b) species sets. We highlight the three species–chemical MoA combinations with highest predicted risks from an Australian context:

1. *Tetranychus urticae*, the two-spotted mite, to avermectins (MoA group 6, $\hat{y}$ = 0.69). *Tetranychus urticae* has resistance to 12 different chemical MoAs in Australia, and there are only six chemical MoAs for which *T. urticae* is susceptible. Within Australia, avermectins are widely used to control *T. urticae* in grain, vegetable, cotton, and fruit crops (Australian Pesticides and Veterinary Medicines Authority 2021; Dodds and Fearnley 2021; Grundy et al. 2021). Avermectins are the third most registered chemical MoA for *T. urticae* (following

organophosphates and pyrethroids), so resistance to this chemical MoA would constitute a non-trivial loss of chemical control options. Field resistance of *T. urticae* to avermectins has already been reported in other parts of the world, involving metabolic mechanisms (Stumpf and Nauen 2002), and reported laboratory-evolved resistance has been underpinned by target-site mutations (Kwon et al. 2010; Dermauw et al. 2012). Our results indicate Australian *T. urticae* populations have a moderate risk of evolving resistance to avermectins, highlighting the value of screening field populations to assess the status of local resistance to this chemical group.

2. *Epiphyas postvittana*, light brown apple moth, to pyrethroids (MoA group 3A, $\hat{y}$ = 0.47). Reported resistances in this pest are to older chemical groups (*i.e.,* carbamates, organophosphates and organochlorines) from the 1960s through to the 1980s (Kerr 1964; Mota-Sanchez and Wise 2021). There are many chemical options for *E. postvittana* control, but pyrethroids are the third most registered chemical MoA for *E. postvittana* in Australia (following organophosphates and carbamates) and are used for control of this pest in fruit, vegetable, and ornamental crops (Australian Pesticides and Veterinary Medicines Authority 2021). Although non-chemical management options are available to control *E. postvittana* (Suckling and Brockerhoff 2010), the moderately high predicted risk to pyrethroids suggests judicious chemical rotations are an important part of management strategies implemented for this pest in Australia.

3. *Helicoverpa armigera*, corn earworm, to amitraz (MoA group 19, $\hat{y}$ = 0.46). Amitraz is primarily used in the Australian cotton industry to manage *H. armigera*, with up to four applications often recommended per growing season (Grundy et al. 2021). Amitraz is the only chemical MoA group currently registered against *H. armigera* for which no current resistance data is available in Australia (Grundy et al. 2021), although it was presumably still susceptible in the early 2000s (Gunning 2002). *Helicoverpa armigera* are already resistant to eight chemical MoA groups in Australia, many of which are widely distributed throughout the country (Thia et al. 2021). Although there are no reports of resistance to amitraz globally in *H. armigera* (Mota-Sanchez and Wise 2021), our results suggest Australian populations have a moderately high risk of evolving resistance. Baseline data on current amitraz susceptibilities in Australia would be valuable to develop a complete picture of chemical tolerances and evaluate the need for preemptive strategies to maintain amitraz susceptibilities.

[FIGURE 5]

Consistency in the effect of pesticide product registration in both the USA and Australian models of resistance status is a noteworthy outcome of our analyses. Such a finding might, in part, be due to shared characteristics of the cropping industries and agro-ecosystems of these countries. For example, the taxonomic composition of agricultural arthropod pests was similar between countries: the orders Lepidoptera and Hemiptera had the most registered pest species, followed by Coleoptera and Trombifidormes (Figure 1). However, at the species level, both countries have different pest communities. In both countries, cotton and grains are the highest contributing cropping commodities to total earnings from agricultural production. In the USA, corn, soybeans, wheat, and cotton make up 15%, 11%, 3% and 2% of the total earnings (respectively), and in Australia, wheat, barley, canola, and cotton make up 13%, 5%, 4%, 3% (respectively) (Howden and Zammit 2019). Whilst the effect size of pesticide product registrations on resistance status was similar between the USA and Australia, application of this predictor into other geographic regions will require construction of region-specific datasets and models of resistance status. In doing so, the framework developed here could be extended to other countries and potentially to other agricultural pests, such as weeds and fungi, by compiling similar datasets of specific resistances and product registrations to herbicides and fungicides.

Whilst pesticide usage is necessary to impose selection, the ecology of pests can play an important role in the rate at which resistance evolves. Our work demonstrated that resistance evolution in Australian agricultural pests exhibits similar curvilinear relationships with phagy and voltinism as that observed in USA (Rosenheim and Tabashnik 1991; Rosenheim et al. 1996; Hardy et al. 2018; Crossley et al. 2020). There was a striking consistency in both the direction and magnitude of effect in partial regression coefficients between the pest species sets from the USA and Australia (Figure 3). The positive second order effect of phagy suggests that pest species with an intermediate host plant breadth have the lower risk of resistance, and that narrow and wide host plant breadth is associated with greatest risk of resistance. For more specialized pests with narrow host plant breadth, a greater proportion of the population is constrained to treated crops and exposed to selection (Georghiou and Taylor 1986). For more generalist pests with wide host plant breadth, a greater arsenal of biochemical machinery used to metabolize plant phytochemicals are available to detoxify pesticides (Dermauw et al. 2018; Rane et al. 2019). The negative second order effect of voltinism suggests that pest species with an intermediate number of generations per year have the greatest risk of resistance, and that few and many generations per year is associated with lower risk of resistance. The curvilinear relationship between voltinism and resistance risk has a less clear interpretation. In some cases, voltinism has been described as a nuisance parameter (Crossley et al.

2020). However, the number of generations per year may modulate the effect of other operational, ecological, and population genetic factors on resistance evolution (Georghiou and Taylor 1977; Rosenheim and Tabashnik 1990). Regardless of the underlying mechanism, our analyses show that phagy and voltinism may be truly general predictors of resistance evolution risk in agro-ecosystems.

The predictive framework developed here is not without limitations, including (i) legacy effects captured in pesticide product registration and resistance status datasets; (ii) ephemeral resistances; (iii) lag between the evolution of resistance and the accumulation of pesticide product registrations; (iv) method of assigning pesticide product registrations for taxonomically ambiguous pest names and chemical mixtures; and (v) the coarse scale of our predictions.

(i)   Legacy effects pertain the analysis of temporally accumulated data. For pesticide registrations, we summed the cumulative number of products across years, and for resistance status, we scored pests as resistant to a chemical MoA if there was at least one record, irrespective of when it was reported. Some MoA groups are no longer used in many countries due to their negative health and ecotoxicology effects, such as organochlorines and DDTs, or because resistance is so commonplace they are no longer efficacious, such as pyrethroids for *Frankliniella occidentalis* (western flower thrips) control in Australia (Herron and Gullick 2001).

(ii)   Ephemeral, transient resistances also contribute toward legacy effects. Some species periodically exhibited high levels of resistance, presumably in response to intense short-term selection pressure, which later disappears. In Australia, two such examples include *Therioaphis trifolii* (spotted alfalfa aphid) resistance to carbamates and organophosphates (Holtkamp et al. 1992), and *Helicoverpa punctigera* (native budworm) resistance to pyrethroids (Gunning et al. 1997).

(iii)   Lag effects may occur if pests evolve resistance to a chemical MoA before pesticide product registrations accumulate to a notably "at risk" level. We attempted to account for this by fitting random intercepts for taxonomy and chemical MoA. But we also found that models including pesticide product registrations alone (no ecological predictors) could predict resistance status (Figure 4). Hence, despite being an approximate measure of pesticide usage, we believe that pesticide product registrations sufficiently covary with resistance status to be a useful predictor of resistance risk in agricultural pests that are currently susceptible.

(iv)   Assigning pesticide product registrations required decisions on how to assign taxonomically ambiguous pest names and chemical mixtures. In both the USA and Australian chemical databases, many pesticide products were registered against pest common names that could not be assigned to the species level (genus or higher). To overcome this issue, we assigned pesticide product registrations to all species nested within the relevant taxonomic rank (up to the family level). The effect of this treatment is to make all species within a taxonomic rank more similar. Chemical mixtures contributed counts to each unique chemical MoA present in the pesticide product. This treatment effectively assumes that chemical mixtures apply selection for all constituent chemical MoAs and that their effect is equivalent to pesticide products comprising a single chemical MoA. Treating pesticide product registrations in this way was necessary to exploit as much information as possible from chemical databases and avoid fitting complicated models. Both taxonomically aggregating and separate MoA group scoring of chemical mixtures would be expected to add noise to the data and reduce power.

(v)   Spatially and temporally patchy resistances are somewhat misaligned with our models, which have a coarse spatio-temporal resolution and consider all observations as contemporaneous. Whilst the risk predictions produced by our models are useful for identifying candidate species for proactive management at a region-wide level, they lack the fine-scale resolution necessary to guide action at local levels. Although localized models can be successfully developed, they are almost always constrained to single species (Feng et al. 2010; Ives et al. 2017; Maino et al. 2018; Ma et al. 2021). Research and data management infrastructures that capture fine-scale chemical usages patterns and spatio-temporal variability in pesticide tolerance phenotypes across many pest species are needed (Thia et al. 2021). Such data is necessary to generate flexible multi-species predictive models at scales relevant for localized management.

CONCLUSIONS

Developing analytical methods that delimit 'high-risk' versus 'low-risk' species can guide future resistance monitoring efforts and facilitate decisions on resource allocation. We detail the novel use of pesticide product registrations as a predictor of resistance in agricultural arthropod pests from the USA and Australia. We suggest that knowledge of available pesticide product registrations for

specific pest species can be used to pinpoint when the intensity and duration of chemical selection in the field warrants careful attention. Our work should aid the development of proactive pest management programs in agro-ecosystems.

METHODS

DATASET COMPILATION

Data on pesticide products, phagy and voltinism were collected for agricultural arthropod pests from the USA (n = 427 species) and Australia (n = 209 species). Our working dataset was compiled from publicly available databases and published studies. All data filtering was performed in R v4.1.2 (R Core Team 2022).

Pesticide products — In the absence of detailed information on chemical usage patterns against agricultural pests in most parts of the world, our study takes the approach of using pesticide product registration data as a potential proxy for chemical usage and selection pressure. Our rationale is that economically important arthropod pests will have many pesticide products registered against them and will be exposed to greater pesticide load in the field. Here, we strictly refer to "pesticide product registrations" as all unique product registrations associated with a specific chemical MoA group.

For USA pesticide products, we downloaded pesticide product registration data from the Environmental Protection Agency (EPA; https://www.epa.gov/; accessed February 11th, 2021). Likewise, registrations of Australian pesticide products were obtained from the Australian Pesticides and Veterinary Medicines Authority (APVMA; https://apvma.gov.au/; accessed February 9th, 2021). Each of these databases contain tens of thousands of product registrations across time that include agricultural and non-agricultural uses (EPA, n = 165,661; APVMA, n = 22,090). Keyword searches were used to isolate those registrations related to pesticide products for agricultural pests. For the EPA database, we filtered the "Site" variable for keywords like "FOLIAR TREATMENT" or "SEEDLING" (List S1). For the APVMA database, we filtered the "Category" variable for "AGRICULTURAL" products and then filtered the "Product_type" variable for the keywords: "INSECTICIDE", "MITICIDE", and "MOLLUSCICIDE".

Pest common names were assigned to species scientific names, and multiple common names per species were merged. For logistical reasons, we only considered the top 50% most registered species: first, the number of pesticide product registrations were pooled across MoAs within a pest common name to get the cumulative number of pesticide product registrations; second, pest common names were ranked from most to least cumulative pesticide product registrations; third, only those pest common names above the 50th percentile were selected for downstream analyses. This procedure removed unique pest common names with very few pesticide product registrations and eased manual curation of our dataset. Taxonomic assignment of pest common names was initially automated with the R package *taxize* (Chamberlain and Szöcs 2013; Chamberlain et al. 2020) and were manually assigned otherwise. Species lists were then compared against reference databases of species occurrence to remove any putatively incorrect species assignments. USA species were compared against a published inventory of USA pests (Hardy et al. 2018) and Australian species were compared against occurrence records in the Atlas of Living Australia (https://www.ala.org.au/; accessed July 7th, 2021). We also removed pests like cockroaches, invasive ants, or pests of ornamentals, conifers, and turf. We were left with 427 USA species (750 common names) and 209 Australian species (247 common names) that were agricultural pests (further details in the Supplementary Methods).

Chemical MoA groups were assigned to each product registration based on their listed active ingredients from the Insecticide Resistance Action Committee (IRAC) classifications (Table S1). Next, we summed the number of unique chemical registrations per species for each chemical MoA across relevant pest common names. In both chemical databases, some chemical registrations are listed at higher order taxonomic classifications (*i.e.,* genus or higher), for example: "aphids", "thrips", or "moths". We considered two alternate ways to deal with these ambiguous assignments. Firstly, all entries that cannot be assigned to species level could be removed. However, this would substantially decrease the number of entries in certain pests where many registrations are not at the species level (e.g., cereal aphids). Secondly, entries could be assigned to all species within that taxonomic rank. However, this would inflate the total number of entries for pests that are not targeted by a pesticide product but occur in the same taxonomic rank as a targeted species. This is expected to increase the similarity between targeted and non-targeted species with respect to the number of products registered against them, and hence reduce signals of selection pressure. To retain as much data as possible, we opted for the second approach, allocating pesticide products up to the family level, ignoring registrations at the order level or higher. Additionally, chemical mixtures provided a unique count for each chemical MoA present in the product. For example, a

product containing both a pyrethroid (group 3A) and a neonicotinoid (group 4A) would comprise a pesticide product for each of these chemical MoAs separately.

Phagy and voltinism — We collected data on two ecological traits previously associated with increased rates of pesticide resistance evolution: phagy (number of host plant families) and voltinism (number of generations per year). We leveraged the data published in Hardy et al. (Hardy et al. 2018), which expanded work presented in Rosenheim et al. (Rosenheim et al. 1996). For each candidate species, we extracted available values of phagy and voltinism. For those species not included in Hardy et al. (2018), we performed a literature search to acquire this information using Google Scholar (see Supplementary Methods).

Resistance status — For each species, for each chemical MoA group, we assigned resistance status (present or absent), as reported in the Arthropod Pesticide Resistance Database (APRD: Mota-Sanchez and Wise 2021) and from the literature. For Australian species, we combined APRD records with a literature search. For USA species, APRD records alone were used to assign resistance status. APRD records for both countries were filtered to only include those marked as "Field Evolved Resistances" to ensure we did not include any records of experimentally evolved resistance

STATISTICAL ANALYSES

Predictive models of resistance — We used Bayesian logistic regression to model resistance status in each species against different chemical MoA groups. Our 'full' model was constructed as follows:

$$\text{RESISTANCE} \sim Bernouilli(p)$$

$$logit(p) = \text{PRODUCTS} + \text{PHAGY} + \text{PHAGY}^2 + \text{VOLTINISM} + \text{VOLTINISM}^2 + \text{TAXONOMY} + \text{CHEM}$$

RESISTANCE was a Bernoulli response for resistance status, 0 for susceptible and 1 for resistant, for each species to a given. PRODUCTS (number of product registrations), PHAGY (number of host plant families), and VOLTINISM (number of generations per year), are continuous fixed effects. Each of these fixed effect predictors were log-scaled: where $\vec{x}$ was the vector of predictor values, we applied the transformation $\log_{10}(\vec{x} + 1)$ and then scaled to a mean of 0 and variance of 1. Because prior studies have indicated curvilinear relationships between resistance evolution and these two ecological traits (Rosenheim et al. 1996; Hardy et al. 2018; Crossley et al. 2020), we fitted first and second order orthogonal polynomials for PHAGY and VOLTINISM using R's *poly* package. TAXONOMY is a random effect accounting for taxonomic structure, fitting a random intercept for each genus, nested in family,

nested in order, as used elsewhere (Hardy et al. 2018; Crossley et al. 2020). CHEM is a random effect, fitting a random intercept for each chemical MoA group.

We fit our 'full' model to pest datasets with imputed and non-imputed ecological traits to evaluate the effect of this imputation on predictions. To evaluate the performance of our resistance predictors, we constructed a 'null' model that simply fitted the global mean and the random effects of taxonomy and chemical MoA group. A 'products' model was also fitted to test the predictive power of pesticide product registrations in isolation. All three models were fit separately on the USA and Australian pest species sets in a Bayesian framework with R's *brms* package (Bürkner 2017). All models were fitted using 10 chains, each with 8,000 iterations, for a total of 80,000 iterations. The first 50% of iterations were used as burn-in, leaving 40,000 iterations for estimating model coefficients. An adaptive delta value of 0.999 was used to set a high average target acceptance rate and prevent divergent transitions after the burn-in phase. The 'full' and 'products' models were compared against the 'null' model using the *bayestestR* R package (Makowski et al. 2019) to obtain Bayes factors. A Bayes factor >3, or a log-Bayes factor >1.1, indicates substantial support for a focal model relative to the null model (Wetzels et al. 2011).

Model generality — We used *K*-fold cross-validation to evaluate the generality and predictive power of our 'full' model and the 'products' model. We analyzed 100 training–testing partitions of both models used imputed datasets. For each partition, 85% of the resistant and susceptible species were randomly selected for the training subset, and the remaining 15% of resistant and susceptible species were allocated to the testing subset. Thus, proportion of resistant to susceptible species was retained in both the training and testing subsets. The 'full' and 'products' models were fitted to each training subset using the *brms* function with 10 chains, 1,000 iterations per chain, and a burn-in of 5,000 chains, with 5,000 iterations for coefficient estimation, and an adaptive delta of 0.999. Predictions on the testing subsets were made using the *posterior_predict* function from the *brms* package without inclusion of random effects. Random effects were excluded because random subsampling of species led to non-overlapping combinations of order, family, and genus between the training and testing subsets. Because predictions were made without consideration of random intercepts for taxonomy and chemical MoA group, our cross-validation analyses provided a more conservative assessment of the generality of our predictive models. For direct comparison, additive and polynomial models were also fitted to the same training–testing species subsets.

To quantify predictive power, we calculated AUC (area under the receiver–operator characteristic curve) scores for each testing subset. AUC scores describe the relative rate of true to false positives

for increasing predicted risk scores. AUC scores range from 0–1: as values approach 1, there is increasing likelihood that resistant species will be assigned higher predicted risk scores than susceptible species. For each testing subset, observed (0 or 1) and predicted (probability from 0 to 1) values were passed to the *auc* function from R's *Metrics* package to obtain an AUC score.

DECLARATIONS

Ethics approval and consent to participate — No ethics approval was required to conduct this study.

Consent for publications — No ethics approval was required to conduct this study.

Availability of data and materials — Data will be uploaded to Dryad, pending acceptance.

Competing interests — The authors declare no competing interests associated with manuscript.

Funding — This research was supported through funding provided by the Grains Research and Development Corporation and the University of Melbourne.

Authors' contributions — PU and AH conceived the project and secured funding. JT and JM designed the study and compiled the pesticide registration data. AK led the compilation of trait data. JT led the statistical analysis and drafted the original manuscript. All authors contributed revisions and approved the manuscript.

Acknowledgements — We thank David Mota-Sanchez and John Wise for their permission to use the Arthropod Pesticide Resistance Database records in our study. We acknowledge Kaya Moore, Candida Wong, Courtney Brown and Safieh Soleimannejad for helping curate taxonomic information and ecological trait data. Computational resources were provided by The University of Melbourne's Research Computing Services and the Petascale Campus Initiative.

REFERENCES

Australian Pesticides and Veterinary Medicines Authority (2021) Public Chemical Registration Information System Search. https://portal.apvma.gov.au/pubcris. Accessed 26 Apr 2021

Baer CF, Miyamoto MM, Denver DR (2007) Mutation rate variation in multicellular eukaryotes: Causes and consequences. Nat Rev Genet 8:619–631. https://doi.org/10.1038/nrg2158

Bass C, Denholm I, Williamson MS, Nauen R (2015) The global status of insect resistance to neonicotinoid insecticides. Pestic Biochem Physiol 121:78–87. https://doi.org/10.1016/j.pestbp.2015.04.004

Brevik K, Schoville SD, Mota-Sanchez D, Chen YH (2018) Pesticide durability and the evolution of resistance: A novel application of survival analysis. Pest Manag Sci 74:1953–1963. https://doi.org/10.1002/ps.4899

Bürkner P-C (2017) brms: An R package for Bayesian multilevel models using Stan. J Stat Softw 80:1–28

Chamberlain SA, Szöcs E (2013) taxize: taxonomic search and retrieval in R. F1000Research 2:

Chamberlain SA, Szoecs E, Foster Z, et al (2020) taxize: Taxonomic information from around the web

Crossley MS, Snyder WE, Hardy NB (2020) Insect–plant relationships predict the speed of insecticide adaptation. Evol Appl 1–7. https://doi.org/10.1111/eva.13089

Crow JF (1957) Genetics of Insect Resistance to Chemicals. Annu Rev Entomol 2:227–246. https://doi.org/10.1146/annurev.en.02.010157.001303

Dermauw W, Ilias A, Riga M, et al (2012) The cys-loop ligand-gated ion channel gene family of *Tetranychus urticae*: Implications for acaricide toxicology and a novel mutation associated with abamectin resistance. Insect Biochem Mol Biol 42:455–465. https://doi.org/10.1016/j.ibmb.2012.03.002

Dermauw W, Pym A, Bass C, Leeuwen T Van (2018) Does host plant adaptation lead to pesticide resistance in generalist herbivores? Curr Opin Insect Sci 26:25–33. https://doi.org/10.1016/j.cois.2018.01.001

Dodds K, Fearnley J (2021) Orchard plant protection guide for deciduous fruits in NSW

Evans JA, Tranel PJ, Hager AG, et al (2016) Managing the evolution of herbicide resistance. Pest Manag Sci 72:74–80. https://doi.org/10.1002/ps.4009

Feng H, Gould F, Huang Y, et al (2010) Modeling the population dynamics of cotton bollworm *Helicoverpa armigera* (Hübner) (Lepidoptera: Noctuidae) over a wide area in northern China. Ecol Modell 221:1819–1830. https://doi.org/10.1016/j.ecolmodel.2010.04.003

Georghiou GP, Taylor CE (1977) Operational influences in the evolution of insecticide resistance. J Econ Entomol 70:653–658. https://doi.org/10.1093/jee/70.5.653

Georghiou GP, Taylor CE (1986) Factors influencing the evolution of resistance. In: Pesticide Resistance: Strategies and Tactics for Management. National Academic Press, Washington, DC, USA, pp 157–169

Gould F, Brown ZS, Kuzma J (2018) Wicked evolution: Can we address the sociobiological dilemma of pesticide resistance? 360:728–732

Grundy P, Koetz E, Holman S, et al (2021) Cotton Pest Management Guide 2021–22

Gunning R (2002) *Heliothis* insecticide resistance continues to climb. Aust Cottongrower 23:73

Gunning R V., Moores GD, Devonshire AL (1997) Esterases and fenvalerate resistance in a field population of *Helicoverpa punctigera* (Lepidoptera: Noctuidae) in Australia. Pestic Biochem Physiol 58:155–162. https://doi.org/10.1006/pest.1997.2295

Handford CE, Elliott CT, Campbell K (2015) A review of the global pesticide legislation and the scale of challenge in reaching the global harmonization of food safety standards. Integr Environ Assess Manag 11:525–536. https://doi.org/10.1002/ieam.1635

Hardy NB, Peterson DA, Ross L, Rosenheim JA (2018) Does a plant-eating insect's diet govern the evolution of insecticide resistance? Comparative tests of the pre-adaptation hypothesis. Evol Appl 11:739–747. https://doi.org/10.1111/eva.12579

Hawkins NJ, Bass C, Dixon A, Neve P (2019) The evolutionary origins of pesticide resistance. Biol Rev 94:135–155. https://doi.org/10.1111/brv.12440

Herron G, Gullick G (2001) Insecticide resistance in Australian populations of *Frankliniella occidentalis* (Pergande) (Thysanoptera: Thripidae) causes the abandonment of Pyrethroid

chemicals for its control. Gen Appl Entomol 30:21–26

Hicks HL, Comont D, Coutts SR, et al (2018) The factors driving evolved herbicide resistance at a national scale. Nat Ecol Evol 2:529–536. https://doi.org/10.1038/s41559-018-0470-1

Holtkamp RH, Edge VE, Dominiak BC, Walters J (1992) Insecticide Resistance in *Therioaphis trifolii f. maculata* (Hemiptera: Aphididae) in Australia. J Econ Entomol. https://doi.org/10.1093/jee/85.5.1576

Horowitz AR, Ghanim M, Roditakis E, et al (2020) Insecticide resistance and its management in *Bemisia tabaci* species. J Pest Sci (2004) 93:893–910. https://doi.org/10.1007/s10340-020-01210-0

Howden M, Zammit K (2019) Analysis of United States and Australian agriculture – a comparison

Ives AR, Paull C, Hulthen A, et al (2017) Spatio-temporal variation in landscape composition may speed resistance evolution of pests to Bt crops. PLoS One 12:1–20. https://doi.org/10.1371/journal.pone.0169167

Kerr RW (1964) Notes on arthropod resistance to chemicals used in their control in Australia. J Aust Inst Agric Sci 30:

Kwon DH, Yoon KS, Clark JM, Lee SH (2010) A point mutation in a glutamate-gated chloride channel confers abamectin resistance in the two-spotted spider mite, *Tetranychus urticae* Koch. Insect Mol Biol 19:583–591. https://doi.org/10.1111/j.1365-2583.2010.01017.x

Li WH, Tanimura M, Sharp PM (1987) An evaluation of the molecular clock hypothesis using mammalian DNA sequences. J Mol Evol 25:330–342. https://doi.org/10.1007/BF02603118

Liu N (2015) Insecticide resistance in mosquitoes: Impact, mechanisms, and research directions. Annu Rev Entomol 60:537–559. https://doi.org/10.1146/annurev-ento-010814-020828

Ma C Sen, Zhang W, Peng Y, et al (2021) Climate warming promotes pesticide resistance through expanding overwintering range of a global pest. Nat Commun 12:1–10. https://doi.org/10.1038/s41467-021-25505-7

Maino JL, Thia J, Hoffmann AA, Umina PA (in press) Estimating rates of pesticide usage from trends in herbicide, insecticide, and fungicide product registrations. Crop Prot

Maino JL, Umina PA, Hoffmann AA (2018) Climate contributes to the evolution of pesticide

resistance. Glob Ecol Biogeogr 27:223–232. https://doi.org/10.1111/geb.12692

Makowski D, Ben-Shachar MS, Lüdecke D (2019) bayestestR: Describing effects and their uncertainty, existence and significance within the Bayesian framework. J Open Source Softw 4:1541

McKenzie JA, Batterham P (1994) The genetic, molecular and phenotypic consequences of selection for insecticide resistance. Trend Ecol Evol 9:166–169

Mota-Sanchez D, Wise JC (2021) The Arthropod Pesticide Resistance Database. http://www.pesticideresistance.org. Accessed 3 Feb 2021

Muth C, Oravecz Z, Gabry J (2018) User-friendly Bayesian regression modeling: A tutorial with rstanarm and shinystan. Quant Methods Psychol 14:99–119. https://doi.org/10.20982/tqmp.14.2.p099

Nauen R (2007) Insecticide resistance in disease vectors of public health importance. Pest Manag. Sci.

Pimentel D, Acquay H, Biltonen M, et al (1992) Environmental and Economic Costs of Pesticide Use. Bioscience 42:750–760. https://doi.org/10.2307/1311994

Pu J, Wang Z, Chung H (2020) Climate change and the genetics of insecticide resistance. Pest Manag Sci 79:846–852. https://doi.org/10.1002/ps.5700

R Core Team (2022) R: A language and environment for statistical computing (v4.1.3)

Rane R V, Ghodke AB, Hoffmann AA, et al (2019) Detoxifying enzyme complements and host use phenotypes in 160 insect species. Curr Opin Insect Sci 31:131–138. https://doi.org/10.1016/j.cois.2018.12.008

Robertson A (1960) A Theory of Limits in Artificial Selection. Proc R Soc B Biol Sci 153:234–249

Rosenheim JA, Johnson MW, Mau RFL, et al (1996) Biochemical preadaptations, founder events, and the evolution of resistance in arthropods. J Econ Entomol 89:263–273. https://doi.org/10.1093/jee/89.2.263

Rosenheim JA, Tabashnik BE (1990) Evolution of pesticide resistance: interactions between generation time and genetic, ecological, and operational factors. J Econ Entomol 83:1184–1193. https://doi.org/10.1093/jee/83.4.1184

Rosenheim JA, Tabashnik BE (1991) Influence of generation time on the rate of response to selection. Am Nat 137:527–541. https://doi.org/10.1086/285181

Sharma A, Kumar V, Shahzad B, et al (2019) Worldwide pesticide usage and its impacts on ecosystem. SN Appl Sci 1:1–16. https://doi.org/10.1007/s42452-019-1485-1

Sparks TC, Nauen R (2015) IRAC: Mode of action classification and insecticide resistance management. Pestic Biochem Physiol 121:122–128. https://doi.org/10.1016/j.pestbp.2014.11.014

Stumpf N, Nauen R (2002) Biochemical Markers Linked to Abamectin Resistance in *Tetranychus urticae* (Acari: Tetranychidae). Pestic Biochem Physiol 72:111–121. https://doi.org/10.1006/pest.2001.2583

Suckling DM, Brockerhoff EG (2010) Invasion biology, ecology, and management of the light brown apple moth (Tortricidae). Annu Rev Entomol 55:285–306. https://doi.org/10.1146/annurev-ento-112408-085311

Tabashnik BE, Mota-Sanchez D, Whalon ME, et al (2014) Defining terms for proactive management of resistance to Bt crops and pesticides. J Econ Entomol 107:496–507. https://doi.org/10.1603/EC13458

Thia JA, Hoffmann AA, Umina PA (2021) Empowering Australian insecticide resistance research with genetic information: the road ahead. Austral Entomol 60:147–162. https://doi.org/10.1111/aen.12512

Umina PA, McDonald G, Maino J, et al (2019) Escalating insecticide resistance in Australian grain pests: contributing factors, industry trends and management opportunities. Pest Manag Sci 75:1494–1506. https://doi.org/10.1002/ps.5285

Weber KE (1990) Increased selection response in larger populations. I. Selection for wing-tip height in *Drosophila melanogaster* at three population sizes. Genetics 125:579–584

Wetzels R, Matzke D, Lee MD, et al (2011) Statistical evidence in experimental psychology: An empirical comparison using 855 *t* tests. Perspect Psychol Sci 6:291–298. https://doi.org/10.1177/1745691611406923

FIGURES AND TABLES

Figure 1. Taxonomic distribution of analysed agricultural arthropod pests. Panels illustrate the proportion of species from different taxonomic groups with resistance (left column) or in the total species set (right column) in the USA (top panels) or in Australia (bottom panels). Taxonomic orders are on the *y*-axis, and the proportion of species is on the *x*-axis. The length of bars indicates the proportion of species, coloured by order (see legend), with numbers to the right of bars noting the actual number of species.

Figure 2. Resistance predictor distributions. Curves depict the kernel density estimate, with area under the curve equal to 1.00. Resistant species–chemical MoA observations are depicted by pink curves, and blue curves depict susceptible species–chemical MoA observations. Panels contain observations from the USA and the Australian species sets separately. (a) Products, the log-scaled number of product registrations. (b) Phagy, the log-scaled number of host plant families. (c) Voltinism, the log-scaled number of generations per year.

Figure 3. Intercept and partial regression coefficient estimates in models of resistance. Each panel represents a modelled term, with the *x*-axis representing the country, and the *y*-axis representing the estimate, with Bayesian 95% credible intervals. The dashed line demarcates an estimate of zero.

Figure 4. Comparisons of predictive power between models using AUC score using 100 training–testing partitions. Curves depict the kernel density estimate, with area under the curve equal to 1.00. Dark grey curves depict the distribution of AUC scores obtained from the 'full' model and light grey curves depicts scores obtained from the 'products' model. The dashed lines depict an AUC score of 0.50, that is, an equal probability that the model assigns a higher score to resistant or susceptible species–chemical MoA observation. AUC scores approaching 1.00 indicate an increasing probability that resistance species–chemical MoA observations will be assigned higher scores than susceptible observations. Panels contain results from the USA and the Australian species sets separately.

Figure 5. Predicted risks for the ten most resistant agricultural pest species in the USA and Australia. Closed circles represent species–MoA combinations that exhibit resistance, whereas open circles represent species–MoA combinations that are susceptible. Circles are coloured by the magnitude of predicted risk, with low risk in pale blue (0.00) to high risk in red (1.00). Only risks for chemical MoAs registered for each pest species are shown. (a) The USA top ten pest species with most cases of resistance (field evolved resistances reported in the APRD). (b) The Australian top ten pest species with most cases of resistance (field evolved resistances reported in the APRD and literature search).

Table 1. Posterior estimates of model terms and Bayesian diagnostic statistics for 'full' models of resistance.

Country	Term	β	CI$_{2.5\%}$	CI$_{97.5\%}$	$\hat{R}$	Bulk ESS	Tail ESS
USA	Intercept	−9.17	0.86	−10.97	−7.56	1.00	16,560
	Products	0.68	0.08	0.53	0.83	1.00	38,115
	Phagy[1]	82.26	26.52	31.91	135.73	1.00	24,326
	Phagy[2]	79.30	21.15	38.63	121.70	1.00	27,828
	Voltinism[1]	71.60	35.10	1.70	139.77	1.00	24,331
	Voltinism[2]	−100.90	25.44	−152.36	−52.28	1.00	30,365
Australia	Intercept	−14.31	−20.00	−10.12	1.00	11,925	17,263
	Products	1.09	0.82	1.37	1.00	27,153	29,236
	Phagy[1]	219.49	95.52	377.60	1.00	15,801	21,471
	Phagy[2]	199.58	94.87	326.51	1.00	17,930	23,804
	Voltinism[1]	103.71	−68.40	295.92	1.00	20,449	21,431
	Voltinism[2]	−179.64	−380.84	−25.06	1.00	16,824	21,061

Note: CIs are the Bayesian credible intervals; $\hat{R}$ is a convergence diagnostic statistic, with values of 1.00 indicating convergence among chains; Bulk and tail ESS (effective sample size) indicates the effective number of independent samples from the bulk and tail ends of sampling distribution, respectively, with values >1,000 being considered adequate.

SUPPLEMENTARY INFORMATION

LISTS OF KEYWORDS FOR DATABASE QUERIES AND LITERATURE SEARCHES

List S1. Key words used to filter EPA database of chemical registration in America.

- BAIT APPLICATION
- BARK TREATMENT
- DIP TREATMENT
- DORMANT APPLICATION
- FOLIAR TREATMENT
- FORAGE
- GRAIN
- GREENHOUSE
- NURSERY
- PASTURE
- POSTHARVEST
- ROOT
- SEED
- SEEDLING
- SEED CROP
- SEED TREATMENT
- SOIL FUMIGATION
- SOIL TREATMENT
- WATER TREATMENT

List S2. Key words used to search for phagy values in Google Scholar.

- food plant
- host
- hosts
- host plant
- host plants
- phagy

List S3. Key words used to search for voltinism values in Google Scholar.

- biology
- ecology
- generation per year
- generations per year
- life cycle
- life history
- multivoltine
- phenology
- reproductive rate
- univoltine
- voltinism

SUPPORTING TABLES AND FIGURES

Table S1. Chemical MoA groups and active ingredients included in this study.

MoA	Chemical class	Target	Active ingredients	USA	Australia
1A	Carbamates	Acetylcholinesterase inhibitors	Aldicarb, Bendiocarb, Carbaryl, Carbofuran, Methomyl, Promecarb, Propoxur, Thiodicarb, Methiocarb, Pirimicarb, Carbosulfan, Oxamyl	✓	✓
1B	Organophosphates	Acetylcholinesterase inhibitors	Fenitrothion, Malathion, Acephate, Azinphos-ethyl, Azinphos-methyl, Chlorpyrifos-methyl, Chlorpyrifos, Diazinon, Dichlorvos, Dimethoate, Disulfoton, Fenamiphos, Fenthion, Methamidophos, Methidathion, Naled, Omethoate, Parathion-methyl, Parathion, Phorate, Phosmet, Prothiofos, Sulprofos, Thiometon, Trichlorfon, Vamidothion, Ethion, Isofenphos, Monocrotophos, Cadusafos, Mevinphos, Profenofos, Ddvp, Ronnel, Oxydemeton-methyl, Phosphamidon, Carbophenothion, Dicrotophos, Phosalone, Terbufos, Dioxathion, Tetrachlorvinphos, Fonofos, Tetraethyl pyrophosphate, Pirimiphos-methyl, Chlorethoxyfos, Isazofos, Fosthiazate	✓	✓
2A	Cyclodiene organochlorines	GABA-gated chloride channel blocks	Chlordane, Dieldrin, Lindane, Endosulfan, Endrin, Dienochlor	✓	✓
2B	Phenypyrazoles	GABA-gated chloride channel blocks	Fipronil	✓	✓
3A	Pyrethroids	Sodium channel modulators	Deltamethrin, Bioallethrin, Bioresmethrin, Pyrethrins, Alpha-cypermethrin, Cyfluthrin, Cypermethrin, Tau-fluvalinate, Bifenthrin, Esfenvalerate, Zeta-cypermethrin, Lambda-cyhalothrin, Permethrin, Gamma-cyhalothrin, Imiprothrin, Resmethrin, Beta-cyfluthrin, Tetramethrin, Fenvalerate, Fluvalinate, Flucythrinate, Fenpropathrin, Phenothrin, Prallethrin, Etofenprox, Allethrin, Tralomethrin, Tefluthrin	✓	✓
3B	DDT	Sodium channel modulators	DDT, Methoxychlor	✓	✓

4A	Neonicotinoids	Nicotinic acetylcholine receptor competitive modulators	Imidacloprid, Thiamethoxam, Thiacloprid, Acetamiprid, Clothianidin, Dinotefuran	✓	✓
4C	Sulfoximines	Nicotinic acetylcholine receptor competitive modulators	Sulfoxaflor	✓	✓
4D	Butenolides	Nicotinic acetylcholine receptor competitive modulators	Flupyradifurone	✓	
5	Spinosyns	Nicotinic acetylcholine receptor allosteric modulators	Spinosad, Spinetoram	✓	✓
6	Avermectins and milbemycins	Glutamate-gated chloride channel allosteric modulators	Abamectin, Milbemectin, Emamectin benzoate	✓	✓
7A	Juvenile hormone analogues	Juvenile hormone mimics	Methoprene	✓	✓
7B	Fenoxycarb	Juvenile hormone mimics	Fenoxycarb	✓	✓
7C	Pyriproxyfen	Juvenile hormone mimics	Pyriproxyfen	✓	✓
8A	Alkyl halides	Miscellaneous non-specific inhibitors	Ethylene dibromide	✓	
8B	Chloropicrin	Miscellaneous non-specific inhibitors	Chloropicrin	✓	✓
8C	Fluorides	Miscellaneous non-specific inhibitors	Sulfuryl fluoride	✓	✓

9B	Pyridine azomethine derivatives	Chordontonal organ TRPV channel modulators	Pymetrozine	✓	✓
9D	Pyropenes	Chordontonal organ TRPV channel modulators	Afidopyropen	✓	✓
10A	Clofentezine, Diflovidazin, Hexythiazox	Mite growth inhibitors affecting chs1	Clofentezine, Hexythiazox	✓	✓
10B	Etoxazole	Mite growth inhibitors affecting chs1	Etoxazole	✓	✓
11	Bacillus products	Microbial disruptors of insect midgut membranes	Bacillus thuringiensis subsp. kurstaki, Bacillus thuringiensis subsp. israelensis	✓	✓
12A	Diafenthiuron	Inhibitors of mitochondrial ATP synthase	Diafenthiuron		✓
12B	Organotin miticides	Inhibitors of mitochondrial ATP synthase	Azocyclotin, Cyhexatin, Fenbutatin oxide	✓	✓
12C	Propargite	Inhibitors of mitochondrial ATP synthase	Propargite	✓	✓
12D	Tetradifon	Inhibitors of mitochondrial ATP synthase	Tetradifon	✓	✓
13	Pyrroles, dinitrophenols, sulfuramid	Oxidative phosphorylation uncouplers via proton gradient disruption	Chlorfenapyr, Dinocap	✓	✓

15	Benzoylureas	Inhibitors of chitin biosynthesis affecting chs1	Triflumuron, Diflubenzuron, Novaluron	✓	✓
16	Buprofezin	Inhibitors of chitin biosynthesis	Buprofezin	✓	✓
17	Cyromazine	Moulting disruptor, dipterans	Cyromazine	✓	✓
18	Diacylhydrazines	Ecdysone receptor agonists	Tebufenozide, Methoxyfenozide	✓	✓
19	Amitraz	Octopamine receptor agonists	Amitraz, Chlordimeform	✓	✓
20A	Hydramethylnon	Mitochondrial complex III electron transport inhibitors	Hydramethylnon	✓	
20B	Acequinocyl	Mitochondrial complex III electron transport inhibitors	Acequinocyl	✓	✓
20D	Bifenazate	Mitochondrial complex III electron transport inhibitors	Bifenazate	✓	✓
21A	METI acaricides and insecticides	Mitochondrial complex I electron transport inhibitors	Pyridaben, Tebufenpyrad, Fenpyroximate, Fenazaquin	✓	✓
21B	METI acaricides and insecticides	Mitochondrial complex I electron transport inhibitors	Rotenone	✓	✓
22A	Oxadiazines	Voltage-dependent sodium channel blocks	Indoxacarb	✓	✓

				USA	Australia
23	Tetronic and tetramic acid derivatives	Acetyl CoA carboxylase inhibitors	Spirotetramat (iso), Spirotetramat, Spiromesifen, Spirodiclofen	✓	✓
24A	Phosphides	Mitochondrial complex IV electron transport inhibitors	Phosphine	✓	✓
25A	Beta-ketonitrile derivatives	Mitochondrial complex II electron transport inhibitors	Cyflumetofen	✓	
28	Diamides	Ryanodine receptor modulators	Chlorantraniliprole, Flubendiamide, Cyantraniliprole, Cyclaniliprole	✓	✓
29	Flonicamid	Chordontonal organ modulators, undefine target site	Flonicamid	✓	✓

Note: The "Active ingredients" column contains the names of all chemical compounds associated with each chemical MoA observed in this study. Ticks in the "USA" and "Australia" column indicate that the associated chemical MoA was modelled in our analyses, but it does not indicate that all associated active ingredients listed in this table were present in that country.

Figure S1. Comparisons of predicted resistance risks from the 'full' model using datasets with imputed and non-imputed ecological traits. Points represent predicted risks for species–chemical MoA combinations, coloured by resistance status: pink for resistant, and blue for susceptible. The dashed line demarcates the 1:1 association. Panels contain predictions for the USA (left) and Australian (right) species sets.