An rRNA fragment in extracellular vesicles secreted by human airway epithelial cells

increases the fluoroquinolone sensitivity of *P. aeruginosa*

Katja Koeppen[1]*, Thomas H. Hampton[1], Scott A. Gerber[2], Young Ah Goo[3], Byoung-Kyu Cho[3],

Danielle M. Vermilyea[1], Deborah A. Hogan[1] and Bruce A. Stanton[1]

[1]Department of Microbiology and Immunology, Geisel School of Medicine at Dartmouth, Hanover,

New Hampshire, United States of America

[2]Norris Cotton Cancer Center, Geisel School of Medicine at Dartmouth, Lebanon, New

Hampshire, United States of America

[3]Mass Spectrometry Technology Access Center, McDonnell Genome Institute, Washington

University School of Medicine, St. Louis, Missouri, United States of America

*Corresponding author

E-mail: Koeppen.Katja@gmail.com

Abstract

Lung infection by antibiotic resistant strains of *Pseudomonas aeruginosa* is a well-known

concern for immunocompromised hosts including people with lung diseases such as cystic

fibrosis. We have previously demonstrated that extracellular vesicles (EVs) secreted by primary

human airway epithelial cells (AEC) deliver miRNA let-7b-5p to *P. aeruginosa* where it

suppresses biofilm formation and increases sensitivity to beta-lactam antibiotics. In this study

we used RNA-seq to characterize the small RNA (sRNA) content of EVs secreted by AEC and

demonstrate transfer of multiple distinct RNA fragments from EVs to *P. aeruginosa*.

Bioinformatic predictions reveal that several sRNAs may target all three subunits of the

fluoroquinolone efflux pump MexHI-OpmD, an effect predicted to increase antibiotic sensitivity

to fluoroquinolone antibiotics. Exposure of *P. aeruginosa* to EVs resulted in a significant

reduction in the protein levels of MexH (-48%), MexI (-50%) and OpmD (-35%). Moreover, EVs

reduced planktonic growth of *P. aeruginosa* in the presence of the fluoroquinolone antibiotic

ciprofloxacin by 20%. A *mexGHI-opmD* deletion mutant of *P. aeruginosa* phenocopied this

increased sensitivity to ciprofloxacin. Finally, we found that a fragment of an 18S rRNA external

transcribed spacer that was transferred to *P. aeruginosa* by EVs was sufficient to reduce

planktonic growth of *P. aeruginosa* in the presence of ciprofloxacin, to reduce the minimum

inhibitory concentration (MIC) of *P. aeruginosa* for ciprofloxacin by over 50%, and to significantly

reduce protein levels of MexH and OpmD. In conclusion, an rRNA fragment secreted by AEC in

EVs increases the ciprofloxacin sensitivity of *P. aeruginosa* by targeting and down-regulating

the fluoroquinolone efflux pump MexHI-OpmD. A combination of rRNA fragments and

ciprofloxacin packaged in nanoparticles or EVs may benefit patients with antibiotic-resistant *P.

aeruginosa* infections.

Author Summary

According to the World Health Organization and the U.S. Centers for Disease Control and Prevention the development of antibiotic resistant strains of bacteria, including *Pseudomonas aeruginosa*, are a significant global threat to human health. Thus, development of new approaches to eliminate antibiotic resistant infections is required. In this study, we report that lung epithelial cells secrete extracellular vesicles (EVs) that fuse with and deliver small rRNAs to *P. aeruginosa*, and that the rRNAs increase the sensitivity of *P. aeruginosa* to the antibiotic ciprofloxacin by reducing protein levels of the drug efflux pump MexHI-OpmD. We identified one rRNA fragment that by itself significantly reduced the protein levels of MexH and OpmD and increased the ability of ciprofloxacin to kill *P. aeruginosa*. We propose that developing synthetic vesicles containing a combination of the rRNA that inhibits antibiotic efflux pumps and ciprofloxacin would benefit patients with antibiotic resistant *P. aeruginosa* infections.

Introduction

Pseudomonas aeruginosa is an opportunistic bacterial pathogen that can cause chronic infections in immunocompromised hosts, leading to increased morbidity and mortality in individuals with cystic fibrosis (CF), chronic obstructive pulmonary disease (COPD), ventilator-associated pneumonia and burn wounds (1). *P. aeruginosa* is also emerging as a pathogen that commonly co-infects and exacerbates COVID-19 infections (2–5). The rise of difficult-to-eradicate, antibiotic resistant strains of *P. aeruginosa* is posing treatment challenges and increasingly becoming a concern in healthcare settings (6–9). Multidrug-resistant strains of *P.*

aeruginosa are estimated to have caused about 32,600 hospital-acquired infections and 2700

deaths in the US in 2017 and have been rated as a serious threat by the CDC in a 2019 report

(10). Thus, new approaches are needed to eradicate antibiotic resistant infections by *P.*

aeruginosa.

Bi-directional host-pathogen communication through short RNA (sRNA) in extracellular vesicles

(EVs) and the resulting regulation of the host immune response as well as bacterial phenotypes

such as antibiotic sensitivity have become a research focus in recent years (11–19). We have

previously shown that *P. aeruginosa* outer membrane vesicles deliver a transfer RNA (tRNA)

fragment to human airway epithelial cells (AEC), suppressing the host innate immune response

to infection, thereby facilitating chronic infections (11). Eukaryotic EVs have also been reported

to contain tRNA fragments (20–22) and tRNA fragments regulate gene expression in eukaryotes

and prokaryotes (23–27), and to play a role in various human diseases (28,29). Previous

studies have primarily focused on the regulatory effects of tRNA fragments from a given

organism on that same organism, rather than on inter-species crosstalk. Much less is known

about the role of tRNA fragments and other sRNAs in inter-kingdom signaling between

eukaryotic and prokaryotic organisms.

Besides tRNA fragments and messenger RNAs (mRNA), eukaryotic EVs contain a diverse

assortment of non-coding RNA species, including long non-coding RNAs (lncRNA), microRNAs

(miRNA), piwi-interacting RNAs (piRNA), Y RNAs, antisense RNAs, transfer RNAs (tRNA),

tRNA fragments, and ribosomal RNAs (rRNA) (20,22,30–33). EVs as well as their miRNA cargo

are involved in immune regulation and their secretion and RNA content have been shown to be

dysregulated in patients with airway diseases (34–38). For example, a recent study found that

increased levels of miR-223-3p or miR-451a and decreased levels of miR-27b-3p in sputum

supernatants correlated with exacerbation in children with CF and infection with *Aspergillus*,

Haemophilus, or *Pseudomonas*, respectively, suggesting the involvement of EV miRNAs in

pulmonary exacerbation (39). Two recent studies provide a comprehensive characterization of

the sRNA content of EVs secreted by AEC – one reporting changes in the sRNA content of EVs

secreted by the human airway epithelial cell line A549 in response to viral infection (40), and

another describing cigarette smoke induced changes in the RNA content of EVs secreted by

primary small airway epithelial cells (41). Prior studies of EVs secreted by airway epithelial cells

(AEC) have focused mainly on their miRNA cargo (19,42–44), whereas the function of other

sRNA species contained in AEC EVs has not yet been reported. While there is a growing body

of literature on the regulatory effects of EV sRNAs within a given organism, little is known about

the ability of sRNAs secreted in eukaryotic EVs to regulate gene expression in prokaryotes. For

example, eukaryotic miRNAs affect the growth of gut microbes through an unknown mechanism

(45–47). Recently, we provided the first direct evidence that a miRNA, let-7b-5p, in EVs

secreted by human airway epithelial cells, is transferred to *P. aeruginosa,* where it increases

beta-lactam antibiotic sensitivity and decreases biofilm formation by reducing the abundance of

the beta-lactamase AmpC and several proteins required for biofilm formation (19). Here, we

report that human EVs also reduce the abundance of all three subunits of the fluoroquinolone

efflux pump MexHI-OpmD and increase fluoroquinolone sensitivity, an effect mediated by an

rRNA fragment secreted in EVs.

Results

EVs increase the fluoroquinolone sensitivity of *P. aeruginosa*

In a previous study we demonstrated that EVs secreted by AEC increased the sensitivity of *P. aeruginosa* to beta lactam antibiotics by delivering the miRNA let-7b to *P. aeruginosa* (19). To test the hypothesis that EVs may also alter the sensitivity to other antibiotics typically used to

treat infections by *P. aeruginosa* (48), we examined the ability of EVs to alter the sensitivity of *P.*

aeruginosa to the fluoroquinolone antibiotic ciprofloxacin. To that end, the planktonic growth of

P. aeruginosa strain PA14 as well as the mucoid *P. aeruginosa* CF clinical isolate SMC1585

(49) were measured in the presence and absence of EVs and in the presence and absence of

eight different concentrations of ciprofloxacin to determine the minimal inhibitory concentration

(MIC). EVs alone had no significant effect on planktonic growth of PA14 in the absence of

ciprofloxacin (Fig 1A) but reduced the MIC of ciprofloxacin by 25% (Fig 1B). EVs reduced

planktonic growth of PA14 in the presence of 0.02 µg/ml ciprofloxacin (corresponding to about

one third of the MIC) by 26%, compared to 0.02 µg/ml ciprofloxacin alone (Fig 1C). Likewise,

EVs alone had no significant effect on planktonic growth of clinical isolate SMC1585 in the

absence of ciprofloxacin (Fig 1D) but reduced the MIC of ciprofloxacin by 48% (Fig 1E). EVs

reduced planktonic growth of SMC1585 in the presence of 0.02 µg/ml ciprofloxacin by 27%,

compared to 0.02 µg/ml ciprofloxacin alone (Fig 1F). Full 20-hour growth curves for EV-exposed

and vehicle exposed PA14 and SMC1585 in the presence and absence of 0.02 µg/ml

ciprofloxacin are shown in S1 Fig. In the presence of 0.02 µg/ml ciprofloxacin there was a

statistically significant difference in the growth curves of PA14 and SMC1585 with EVs

compared to controls for the last 4 hours of growth. Thus, the combination of EVs and

ciprofloxacin reduced the MIC of ciprofloxacin as well as the growth of PA14 and a clinical

isolate of *Pseudomonas* (SMC1585).

EVs increase the fluoroquinolone sensitivity of *P. aeruginosa*

by targeting MexHI-OpmD

To begin to elucidate the mechanism whereby EVs increased the sensitivity of *P. aeruginosa* to

ciprofloxacin we analyzed the proteome of *P. aeruginosa* that had been exposed to EVs or

vehicle control. EVs reduced the protein levels of all three subunits of the fluoroquinolone efflux

pump MexHI-OpmD (Table 1). Compared to control, EVs decreased protein levels of MexH by 48%, MexI by 50% and OpmD by 35%. Normalized peak intensities for each sample set as well as average log2 fold changes and p-values for all 1619 detected proteins in *P. aeruginosa* are provided as S1 Table. To determine whether a reduction of MexHI-OpmD alone increased fluoroquinolone sensitivity, planktonic growth experiments were performed with a *mexGHI-opmD* deletion strain of *P. aeruginosa* PAO1 and the parental PAO1 wild type strain. In the presence of ciprofloxacin EVs significantly decreased planktonic growth of wild type PAO1 (Fig 2). By contrast, in the presence of ciprofloxacin, EVs had no significant effect on the *mexGHI-opmD* deletion strain (Δ+EV). S2 Fig contains full 20-hour growth curves for the data shown in Fig 2. Taken together, the data suggest that EVs increase *P. aeruginosa* ciprofloxacin sensitivity by targeting and down-regulating the fluoroquinolone efflux pump MexHI-OpmD.

Table 1. EVs significantly reduce the protein levels of all three subunits of the fluoroquinolone efflux pump MexHI-OpmD.

Uniprot ID	Locus	Name	Product	log2 FC	% decrease	P-value
A0A0H2ZGA2	PA14_09530	MexH	RND efflux membrane fusion protein	-0.93	-48	0.0008
A0A0H2ZGB3	PA14_09520	MexI	RND efflux transporter	-1.01	-50	0.0003
A0A0H2ZF64	PA14_09500	OpmD	outer membrane protein	-0.62	-35	0.0161

Protein level changes in *P. aeruginosa* exposed to EVs compared to vehicle controls. FC = fold change, N = three biological replicates of EVs secreted by different AEC donors. Statistical significance was determined with QPROT.

Characterization of sRNA content of EVs

We previously found that EVs secreted by AEC deliver the miRNA let-7b-5p to *P. aeruginosa*, where it reduced biofilm formation and increased sensitivity to beta-lactam antibiotics by targeting key genes involved in biofilm formation and beta-lactam resistance (19). However, let-7b-5p alone did not reduce protein levels of MexHI-OpmD. To determine whether a different sRNA was responsible for the EV-mediated reduction in MexHI-OpmD protein levels and the resulting increased sensitivity to fluoroquinolone antibiotics, we re-analyzed our existing RNA-seq data (GSE174690) to identify sRNAs in EVs that are predicted to target MexHI-OpmD. The alignment-based characterization of EV sRNA content showed that tRNA and tRNA like fragments were the most abundant sequences in EVs, accounting for 65% of all aligned reads (Table 2). To gain a better understanding of the sRNA content of EVs at the sequence level, we generated count tables for unique sequences and filtered them for the most abundant unique sequences with at least 20 nucleotide length and at least 100 counts in each sample. The resulting 1346 sequences were aligned to homo sapiens using BLAST+ to obtain annotations for the 909 most abundant unique sequences with 100% identity and coverage (S2 Table). Collectively, these 909 unique sequences accounted for 60% of total reads detected in all 3 replicate samples. The sequences with the highest total count (listed at the top of S2 Table) include many tRNA and lncRNA fragments.

Table 2. Alignment-based characterization of EV sRNA content.

RNA type	% of total reads
tRNA	46.0
tRNA like	19.0
RefSeq anti	16.5
RefSeq	9.9

rfam	3.0
rRNA	1.8
other ncRNA	1.4
piRNA	1.1
lncRNA	0.7
miRNA	0.3
lncRNA anti	0.3
other ncRNA anti	0.1
CDBox	0.003

188   RNA types (column 1) and their relative abundance as a percentage of total reads (column 2).

189   EVs were derived from three individual AEC donors. tRNA = transfer RNA, RefSeq =

190   messenger RNA, anti = antisense, rfam = non-coding RNA annotated in Rfam database, rRNA

191   = ribosomal RNA, ncRNA = non-coding RNA, piRNA = Piwi-interacting RNA, lncRNA = long

192   non-coding RNA, miRNA = microRNA, CDBox = Small nucleolar RNA (snoRNA).

Transfer of EV sRNAs to *P. aeruginosa*

We previously described the transfer of miRNAs secreted in AEC EVs to *P. aeruginosa* (19). To

assess the transfer of other sRNAs besides miRNAs to *P. aeruginosa*, we re-analyzed RNA-seq

data of EV-exposed *P. aeruginosa* and unexposed controls (GSE174710). We generated count

tables of unique sequences from trimmed reads for individual samples - three unexposed

controls and three samples of *P. aeruginosa* exposed to EVs secreted by different AEC donors.

Reads were filtered to include only sequences with a length of at least 20 nucleotides that were

detected in all three EV-exposed *P. aeruginosa* samples and none of the unexposed control

samples. Filtering on these criteria yielded 1022 sequences, 15 of which were perfect matches

to human sequences in a BLAST search. The length requirement of at least 20 nucleotides was

chosen because shorter sequences tend to be less specific and map to many different regions

in the genome. The 15 human EV sRNA sequences listed in Table 3 were detected in EV-

exposed *P. aeruginosa*, but not in any of the unexposed control samples, suggesting that they

are delivered to *P. aeruginosa* by EVs. Subsequently, we used IntaRNA to predict target genes

of these 15 human EV sRNAs in *P. aeruginosa.*

Table 3. Human EV RNA fragments delivered to *P. aeruginosa*.

Sequence	Length	Description	Total Count	*mexH* Energy	*mexI* Energy	*opmD* Energy
TGGATTTTTGGAGCAGGGAG	20	CpG island DNA genomic Mse1 fragment	7	-13.19	-15.07	-17.44
AAATGGATTTTTGGAGCAGGGAG	**23**	**LHRI_LNC217 lncRNA**	**24**	**-13.18**	**-15.06**	**-17.43**
AAATGGATTTTTGGAGCAGGG	21	LHRI_LNC217 lncRNA	9	-8.68	-15.46	-16.97
GCATTGGTGGTTCAGTGGTAGAATTCTCGCCTG	**33**	**tRNA-Gly-GCC**	**11**	**-9.16**	**-14.77**	**-15.15**
GCGGCGTCCGGTGAGCTCTCGCTGGCC	**27**	**external transcribed spacer 18S ribosomal RNA**	**11**	**-10.52**	**-15.68**	**-14.77**
TGAGGTAGGAGGTTGTATAGTT	22	microRNA hsa-let-7e-5p (mature)	28	-10.01	-13.55	-14.6
TGAGGTAGTAGGTTGTATGGTT	22	microRNA let-7c-5p (mature)	21	-11.61	-13.67	-11.3

CAGGGATAACTGGCTTGTGGCGGCCAAG	28	external transcribed spacer 18S ribosomal RNA	4	-9.29	-13.95	-10.94
TCCCACATGGTCTAGCGGTTAGGATTCCTGG	31	tRNA-Glu-TTC	5	-10.26	-14.27	-10.92
AGTAAGGTCAGCTAAATAAGCTATCGGGCCCATACCCCGAAAATGTTGGTTATA	54	mitochondrial tRNA-fMet	6	-8.18	-10.98	-8.97
AGTAAGGTCAGCTAAATAAGCTATCGGGCCCATACCCCGAAAATGTTGGTTA	52	mitochondrial tRNA-fMet	8	-8.27	-10.96	-8.91
GAAAAAAAAAAAGGATGAGG	20	CKLF-like MARVEL transmembrane domain containing 6	7	-5.59	-9.04	-8.27
TGAGGTAGTAGATTGTATAGT	21	microRNA let-7f-5p, partial sequence (missing 3' T)	26	-7.59	-10.77	-7.83
CGCGACCTCAGATCAGACGTGGCG	24	external transcribed spacer 18S ribosomal RNA	6	-3.93	-5.8	-3.2
TCCCTGAGACCCTTTAACCTGT	22	miR-125a-5p, partial sequence	11	-8.78	-8.7	0

		(missing 3' GA)				

Listed are 15 human EV RNAs that were exclusively detected in EV-exposed *P. aeruginosa.*

Three sequences that were top candidates for follow-up experiments based on abundance and

targeting prediction for MexHI-OpmD are highlighted in bold font. Column 1 contains the sRNA

sequence, column 2 the sequence length in nucleotides (nt), column 3 the sequence identity as

determined with BLAST+, column 4 the total number of reads detected in *P. aeruginosa* after

EV exposure, and columns 5-7 the IntaRNA energy scores for *mexH*, *mexI* and *opmD*,

respectively. EVs were derived from three individual AEC donors.

220 **Human EV sRNAs are predicted to target *P. aeruginosa* efflux**

221 **pump *mexHI-opmD***

To predict which *P. aeruginosa* genes the 15 human sRNAs that were delivered to *P.*

aeruginosa by EVs may target, the RNA-RNA interaction tool IntaRNA (50,51) was used to

calculate binding energy scores. The more negative the energy score, the higher the likelihood

of an RNA-RNA interaction. Table 3 lists the IntaRNA energy scores for each of the 15 EV

sRNAs for predicted targeting of the fluoroquinolone efflux pump subunits *mexH*, *mexI*, and

opmD. Three sRNAs with a high count in PA14+EV samples and the best (i.e., lowest) energy

scores for targeting of *mexH*, *mexI*, and *opmD* were selected for follow-up validation

experiments: a 23-nt lncRNA fragment, a 33-nt tRNA-Gly fragment and a 27-nt fragment of an

18S rRNA external transcribed spacer (highlighted in bold font in Table 3). As described next, of

the three top candidate sRNAs based on bioinformatic predictions, the rRNA fragment was

validated in wet lab experiments to increase the fluoroquinolone sensitivity of *P. aeruginosa* and

to decrease MexHI-OpmD protein levels.

A human rRNA fragment mediates the increased fluoroquinolone sensitivity of *P. aeruginosa* by targeting MexHI-OpmD

To test whether the 18S rRNA spacer fragment in EVs mediates the increased ciprofloxacin

sensitivity of *P. aeruginosa* in response to EVs, we expressed the rRNA fragment (sequence

listed in Table 3) in *P. aeruginosa* using a plasmid with an arabinose-inducible promoter and

measured planktonic growth in the presence and absence of ciprofloxacin using the empty

vector as the control strain (ctrl). We observed that the rRNA fragment significantly decreased

planktonic growth of *P. aeruginosa* in the presence of ciprofloxacin (0.03 µg/ml, Fig 3A), while it

did not significantly alter planktonic growth in the absence of ciprofloxacin (Fig 3B).

Full 20-hour growth curves for the data in Fig 3A and 3B are provided in Fig S3. The rRNA

fragment also significantly decreased the MIC of *P. aeruginosa* for ciprofloxacin (Fig 3C), and

significantly reduced the number of live bacteria in the presence of ciprofloxacin (Fig 3D). In the

presence of the rRNA fragment planktonic growth of *P. aeruginosa* was reduced by 76%

compared to the empty vector control, the MIC of *P. aeruginosa* for ciprofloxacin was decreased

two-fold and the number of live bacteria was reduced by 73%.

To validate that the rRNA fragment increased the ciprofloxacin sensitivity of *P. aeruginosa* by

targeting the fluoroquinolone efflux pump MexHI-OpmD, we measured protein levels of MexHI-

OpmD in the presence and absence of the rRNA fragment and found that compared to the

empty vector control, the rRNA fragment decreased MexH by 53%, MexI by 26% and OpmD by

52% (Table 4). The decrease in the protein levels of MexH and OpmD was statistically

significant, and there was a trend for the rRNA fragment to decrease MexI protein levels that did

not quite reach statistical significance. Normalized peak intensities for each sample set as well

as average log2 fold changes and p-values for all 1911 detected proteins in *P. aeruginosa* are

259 provided as S3 Table. Taken together, these experiments demonstrate that the rRNA fragment

260 increased the ciprofloxacin sensitivity of *P. aeruginosa* by targeting the fluoroquinolone efflux

261 pump MexHI-OpmD.

263 **Table 4. A human rRNA fragment significantly reduces protein levels of MexH and OpmD.**

Uniprot ID	Locus	Name	Product	log2 FC	% decrease	p-value
A0A0H2ZGA2	PA14_09530	MexH	RND efflux membrane fusion protein	-1.09	-53	0.0003
A0A0H2ZGB3	PA14_09520	MexI	RND efflux transporter	-0.43	-26	0.1344
A0A0H2ZF64	PA14_09500	OpmD	outer membrane protein	-1.07	-52	0.0107

265 Protein level changes in a *P. aeruginosa* strain induced with 100 mM arabinose to express the

266 rRNA fragment compared to the empty vector control strain. FC = fold change, N = three

267 different clones of each strain. Statistical significance was determined with QPROT.

Discussion

We report that AECs secrete EVs containing lncRNA, tRNA and 18S rRNA external transcribed

spacer fragments that are transferred to *P. aeruginosa*, and that EVs increased the

fluoroquinolone sensitivity of *P. aeruginosa* by reducing protein levels of the drug efflux pump

MexHI-OpmD. An 18S rRNA external transcribed spacer fragment by itself significantly reduced

the protein levels of MexH and OpmD, thereby increasing antibiotic sensitivity to ciprofloxacin.

The finding that AEC EVs contain rRNA fragments is consistent with prior characterizations of

the sRNA content of EVs secreted by airway epithelial cells (40,41). It has been previously

described that rRNA fragments are produced in a controlled manner that is population-, sex-

and tissue-specific, suggesting they constitute another class of small non-coding regulatory

RNAs (52,53). Several reports predict and speculate about a regulatory function of rRNA

derived fragments (52,54,55), with some studies providing evidence for the regulation of various

biological functions by rRNA fragments. Chen et al. report RNAi knockdown of an endogenous

20-nt 28S rRNA fragment in the human non-small cell lung carcinoma cell line H1299 induced

apoptosis and inhibited cell proliferation (56), while Wei et al. found that endogenous rRNA

fragments detected in human and murine liver regulate glucose metabolism in the murine

hepatoma cell line Hepa 1-6 and are differentially expressed in the livers of diabetic mice (57).

Moreover, several studies show co-immunoprecipitation of eukaryotic rRNA fragments with

Argonaute proteins (54,57,58), further corroborating our conclusion that rRNA fragments are

involved in the regulation of gene expression. Most of the previously described rRNA fragments

are derived from 5.8S, 18S or 28S rRNA as well as the internal transcribed spacer and, in fact,

some sequences that had been previously classified as miRNAs or piRNAs are in fact rRNA-

derived (53). There is very limited knowledge about the existence and function of fragments

derived from the rRNA external transcribed spacer. Lee et al. describe upregulation of rRNA

external transcribed spacer fragments in *Neurospora crassa* in response to DNA damage (59),

while Gupta et al. found that fragments derived from an rRNA external transcribed spacer are

upregulated in the parasitic protist *Entamoeba histolytica* in response to growth stress,

suggesting that these fragments play a regulatory role in the stress response (60). However,

direct evidence for the regulation of gene or protein expression by rRNA external transcribed

spacer fragments has not been published. In addition, to our knowledge there are no studies

showing cross kingdom transfer of rRNA fragments from human EVs to a bacterial pathogen,

nor is anything known about the effect of human rRNA fragments on bacterial gene expression.

Thus, to our knowledge, our data represent the first report of an rRNA fragment secreted by a

eukaryotic cell that regulates protein expression and alters the phenotype of a prokaryotic

organism. The exact mechanism of action and potential protein binding partners of rRNA

fragments in prokaryotes are beyond the scope of this work and need to be elucidated in future

studies.

We found that some, but not all, of the most abundant sequences in EVs can be detected in *P.*

aeruginosa following EV exposure. However, some human EV sRNA sequences detected in

EV-exposed *P. aeruginosa* are not among the most abundant EV sequences, suggesting

differential stability in *P. aeruginosa*. While RNA contained within EVs is protected from

degradation by extracellular RNases, once the EV RNA cargo is delivered to *P. aeruginosa,*

RNA that is not protected by binding to protein or *P. aeruginosa* mRNA targets may be more

prone to degradation, explaining the discrepancy between abundance in EVs compared to the

abundance after delivery to *P. aeruginosa* for some of the sRNAs.

One limitation of our study is that we did not assess other AEC EV cargo such as DNA, protein,

lipids, or metabolites, which could contribute to EV induced sensitivity to ciprofloxacin. However,

our experiment expressing the rRNA fragment in *P. aeruginosa* demonstrates that the rRNA

fragment alone is sufficient to reduce the abundance of MexH and OpmD and to increase the

sensitivity to ciprofloxacin.

Another limitation is that due to redundancy of fluoroquinolone efflux pumps in *P. aeruginosa*,

the suppression of MexHI-OpmD only leads to a partial effect on fluoroquinolone sensitivity.

MexEF-OprN serves as another fluoroquinolone efflux pump in *P. aeruginosa* (61), and there

are several additional efflux pumps that can serve this function (62), which helps explain why

deletion of MexHI-OpmD does not result in a more drastic increase in fluoroquinolone sensitivity

(Fig 2). On the other hand, as shown in Fig 2, the downregulation of MexHI-OpmD by EVs is

enough to reduce fluoroquinolone sensitivity to a level achieved with a complete deletion of the

efflux pump. Moreover, we found that expressing the rRNA fragment by itself led to a larger

reduction in *P. aeruginosa* planktonic growth in the presence of ciprofloxacin than EVs, and that

while both EVs and the rRNA fragment alone led to a protein level reduction of the

fluoroquinolone efflux pump MexHI-Opm, the rRNA fragment significantly suppressed additional

proteins involved in mediating fluoroquinolone resistance, including MexA, GyrA , and ParC

(62,63). This may explain the increased ability of the rRNA fragment to reduce planktonic

growth in the presence of ciprofloxacin compared to EVs, which contain additional cargo that

may counteract the effect of the rRNA fragment.

P. aeruginosa in the lung is exposed to EVs secreted by other cell types in addition to those

secreted by AEC, most notably immune cells like neutrophils and macrophages, whose EV RNA

content may include the rRNA fragment in our study as well as other sRNAs that may affect

antibiotic resistance of *Pseudomonas*. Therefore, a final limitation is that our study cannot

elucidate the cumulative effect that EV sRNAs from multiple cell types would have on *P.*

aeruginosa in the lungs*.* However, re-analysis of publicly available data (GSE126051) from a

study characterizing sRNAs in human plasma (64) revealed that rRNA fragments that were

identical in length and sequence to the rRNA fragment described in this study were present in

plasma of healthy control subjects. Moreover, identical rRNA fragments were detected in EVs

isolated from human bronchoalveolar lavage fluid (BALF; Dr. Alix Ashare, personal

communication). These observations reveal the biological significance of our study since our

rRNA fragment was found in clinical samples and is not limited to our model system of primary

AECs. So far, there are no published studies that comprehensively characterize the sRNA

content of EVs isolated from human BALF, and, as noted in a recent review by Zareba et al.,

future in-depth RNA-seq profiling of distinct EV populations in BALF will elucidate their role in

disease pathogenesis and provide insight into treatment strategies (65). The important

questions of whether *P. aeruginosa* exposure alters the sRNA content of AEC EVs and whether

there is a difference in the sRNA cargo of EVs secreted by WT and CF AEC need to be

addressed in future studies.

To assess whether EV sRNA targeting of efflux pumps is limited to *P. aeruginosa* or extends to

efflux pump orthologs in other common lung pathogens we utilized IntaRNA to predict targeting

of *Burkholderia cenocepacia* (which has orthologs for *mexH*, *mexI* and *opmD*) and

Staphylococcus aureus (which has an ortholog for *mexI*). Our candidate rRNA spacer fragment

is predicted to target orthologs of *mexI* and *opmD* in *Burkholderia cenocepacia*, suggesting that

AEC EV sRNAs are likely to affect other lung pathogens beyond *P. aeruginosa* and warrant

future analysis beyond the scope of the current study.

In summary, our study provides the first evidence for transfer of lncRNA, tRNA and rRNA

fragments from eukaryotic EVs to a prokaryote and provides direct evidence for regulation of the

prokaryote by a eukaryotic rRNA fragment, resulting in subsequent phenotypic alterations such

as increased fluoroquinolone sensitivity. The development of new treatment approaches utilizing

a combination of human rRNA fragments and antibiotics in nanoparticles or EVs may benefit

individuals with chronic antibiotic-resistant *P. aeruginosa* infections.

Materials and Methods

Culture of airway epithelial cells and EV isolation

De-identified primary human AEC were obtained from Dr. Scott Randell (University of North

Carolina, Chapel Hill, NC) and cultured in BronchiaLife basal medium (Lifeline Cell Technology,

Frederick, MD, Cat. # LM-0007) supplemented with the BronchiaLife B/T LifeFactors Kit (Lifeline

Cell Technology, Cat. # LS-1047) as well as 10,000 U/ml Penicillin and 10,000 µg/ml

Streptomycin (Sigma-Aldrich, Cat. # P4333), as previously described (66). EVs were isolated

from AEC culture supernatants (passages 4-8) with the ExoQuick-TC EV isolation kit (System

Biosciences, Palo Alto, CA, Cat. # EXOTC50A-1) and characterized as previously described

(67).

Bacterial strains and culture

For all experiments, *P. aeruginosa* strains PA14, PAO1, and the mucoid CF clinical isolate SMC1585 (49) were cultured in Luria broth (LB, Thermo Fisher Scientific, Waltham, MA). A deletion mutant for the fluoroquinolone efflux pump *mexGHI*-opmD in the PAO1 background (strain DW101, PAO1ΔGHID), as well as the matching parental PAO1 wild type strain, were generously provided by Dr. Helen Zgurskaya from the University of Oklahoma (61). Deletion of *opmD* in the PAO1Δ*mexGHI*-opmD strain was confirmed by PCR. Briefly, RNA was isolated from PAO1Δ*mexGHI-opmD* and the parental PAO1 wild type strain using the miRNeasy Mini Kit (Qiagen, Germantown, MD, Cat. # 217004), cDNA was synthesized with the Invitrogen RETROscript Reverse Transcription Kit (Thermo Fisher Scientific Cat. # 10585595), and the presence of *opmD* was interrogated by PCR amplification using the following primers: forward 5'-CCTGGTGGAGTTTCTTCGAC-3' and reverse 5'- CGTCGTAGTCCAGCTGTTGT-3'.

P. aeruginosa strains expressing either the 18S rRNA spacer fragment under the control of an arabinose-inducible promoter or the empty vector control were produced by transforming PA14 with a pMQ70 expression vector (68) containing either the rRNA fragment sequence (listed in Table 3) as an insert, or no insert (control). Strains were cultured with 300 μg/ml carbenicillin (Sigma-Aldrich, Cat. # C1389) to select for bacteria containing the plasmid and 100 mM L-(+)-arabinose (Sigma-Aldrich, St. Louis, MO, Cat. # A3256) was added during experiments to induce expression of the rRNA fragment.

Bacterial growth curves and CFU count

The minimal inhibitory concentration (MIC) of the fluoroquinolone antibiotic ciprofloxacin was determined by incubation of each *P. aeruginosa* strain with 8 different concentrations of ciprofloxacin (0-0.25 μg/ml) in triplicate for 20 hours in a plate reader at 37°C. The OD600 was recorded every 15 minutes and the MIC for ciprofloxacin was calculated by fitting the growth

curve data with a Gompertz function, as previously described (69,70). At the beginning of the

20-hour growth curves, 5000 colony forming units (CFUs) of *P. aeruginosa* were exposed to

5×10^9 EV/ml, corresponding to a concentration of EVs measured in AEC culture supernatants

by us, as well as in human bronchoalveolar lavage fluid (42,71). During exposure of *P.*

aeruginosa to EVs, ciprofloxacin (Sigma-Aldrich, Cat. # 17850-5G-F) was present in the culture

medium at a concentration equivalent to one-third to one-half the MIC of a given strain, as

indicated in the figure legends. CFU/ml for viable planktonic bacteria were obtained by dilution

plating of supernatants.

Proteomics of *P. aeruginosa*

1.5×10^7 CFUs of PA14 were exposed to 1.5×10^{10} EV isolated from AEC from three donors or

vehicle controls for a PA14:EV ratio of 1:1000 and incubated for 15 h at 37°C at 225 rpm.

Samples were processed and TMT labeled as described previously (19). All six samples were

run in duplicate, peak intensities from duplicate runs were averaged and filtered to retain 1619

proteins that were detected in all replicate samples. Differentially abundant proteins were

identified using QPROT (72). The mass spectrometry proteomics data have been deposited to

the ProteomeXchange Consortium (73) via the MassIVE (74) partner repository with the dataset

identifier PXD033213.

In addition, following overnight cultures with 300 µg/ml carbenicillin to select for bacteria with the

plasmid containing the rRNA fragment or the empty plasmid, 3 different clones of the PA14

strains expressing either the rRNA fragment or the empty vector control were cultured with 100

mM arabinose for 15 h at 37°C at 225 rpm to induce expression of the rRNA fragment. Cell

pellets were washed with PBS, lysed and proteins (200 µg of each sample) were purified by

acetone/trichloroacetic acid precipitation overnight at -20°C. After washing the pellet with ice-

cold acetone, the resulting protein pellet was resuspended in 50 µL of 8 M urea in 400 mM

ammonium bicarbonate, pH 7.8, reduced with 4 mM dithiothreitol at 50°C for 30 minutes, and

cysteines were alkylated with 18 mM iodoacetamide for 30 minutes. The solution was then

diluted to <2 M urea and trypsin (Promega, Madison, WI) was added at a final trypsin/protein

ratio of 1:100 prior to overnight incubation at 37°C. The resulting peptides were desalted using

solid-phase extraction on a C18 Spin column and eluted in 240 µL of 80% acetonitrile in 0.1%

formic acid. All samples were analyzed in technical duplicates by LC-MS/MS using a nanoElute

coupled to a timsTOF Pro Mass Spectrometer (Bruker Daltonics). Two hundred nanogram of

each digested peptide sample was loaded on a capillary C18 column (25 cm length, 75 µm

inner diameter, 1.6 µm particle size, 120 Å pore size; IonOpticks). The flow rate was kept at 300

nL/min. Solvent A was 0.1% FA in water, and Solvent B was 0.1% FA in ACN. The peptides

were separated on a 100 min analytical gradient from 2% ACN/0.1% FA to 35% ACN/0.1% FA

for a total of 120 min gradient. The timsTOF Pro was operated in the PASEF mode. MS and

MS/MS spectra were acquired from 100-1700 m/z. The inverse reduced ion mobility $1/K_0$ was

set to $0.60-1.60$ V·s/cm^2 over a ramp time of 100 ms. Data-dependent acquisition was

performed using 10 PASEF MS/MS scans per cycle with a near 100% duty cycle. The resulting

protein tandem MS data was queried for protein identification and label-free quantification

against the Uniprot *Pseudomonas aeruginosa* strain UCBPP-PA14 database (proteome ID

UP000000653, retrieved on Feb 2, 2022) using MaxQuant v2.0.3.1. The following modifications

were set as search parameters: peptide mass tolerance at 20 ppm, trypsin digestion cleavage

after K or R (except when followed by P), 2 allowed missed cleavage sites,

carbamidomethylated cysteine (static modification), and oxidized methionine, deaminated

asparagine/glutamine, and protein N-term acetylation (variable modification). Search results

were validated with peptide and protein FDR both at 0.01. LFQ intensities were averaged

across technical replicates, filtered to retain 1911 proteins detected in at least two replicate

samples, and normalized prior to QPROT analysis (72). The mass spectrometry proteomics

data have been deposited to the ProteomeXchange Consortium (73) via the PRIDE (75) partner

repository with the dataset identifier PXD033165.

Characterization of EV sRNA content

Previously, we characterized the miRNA content of AEC EVs (19); here, we focus on the EV

sRNA content beyond miRNAs. The raw data as well as count tables of aligned reads provided

by System Biosciences can be accessed through NCBI's Gene Expression Omnibus (GEO

Series accession number GSE174690). In addition to the alignment-based characterization we

also performed a sequence-based characterization of EV sRNA content. Adaptor trimming of

raw reads was performed with fastp version 0.20.0 (76). Trimmed reads from EVs secreted by

three donors of AEC yielded close to 500,000 unique sequences, about 40,000 of which were

detected in all three biological replicates. Count tables with the number of reads per sample for

each unique sequence were generated using the Unix command "cat fastp.fastq.gz | gunzip |

awk '(NR%4==2)' | sort | uniq -c | sort -k 1 -n". To determine the origin of the most abundant

unique EV sRNA sequences, we ran command line BLAST+ (77) for 1346 sequences that had

a minimum of 100 counts per sample and a length of at least 20 nucleotides. Arguments to the

blastn function included -taxids 9606 (homo sapiens), -perc_identity 100 (requiring a perfect

match), -qcov_hsp_perc 100 (requiring 100% query coverage) and -task blastn-short (to adjust

for a short input sequence). Genbank accessions returned by BLAST were annotated using the

R package rentrez (78). Subsequently, RNA-seq samples from EV-exposed *P. aeruginosa* were

used to assess whether human sRNAs are transferred to *P. aeruginosa* by EVs.

RNA-seq to detect transfer of EV RNA to *P. aeruginosa*

Existing RNA-seq data of *P. aeruginosa* exposed to PBS vehicle control in triplicate or to EVs

from 3 AEC donors (19) was re-analyzed to identify EV-mediated transfer of sRNA other than

miRNAs to *P. aeruginosa.* The raw data can be accessed through NCBI's Gene Expression

Omnibus (GEO Series accession number GSE174710). For each sample, trimmed reads were

compiled into count tables of unique sequences and samples were filtered for unique

sequences of at least 20 nucleotides in length that were contained in all three EV-exposed *P.*

aeruginosa samples, but none of the control samples. The origin of 1022 unique sequences

matching these criteria was determined using BLAST+ (77), as described above, and included a

separate alignment to -taxids 287 (*P. aeruginosa*). Genbank accessions returned by BLAST

were annotated using the R package rentrez (78). For EV RNA sequences that were of human

origin, we generated IntaRNA targeting predictions (50,51) for *P. aeruginosa* genes, as

described below.

IntaRNA target predictions

We used the IntaRNA algorithm (50,51) to predict whether human sRNAs transferred to *P.*

aeruginosa by EVs might regulate *P. aeruginosa* gene expression by targeting bacterial

mRNAs. *P. aeruginosa* UCBPP-PA14 reference sequences and standard parameters were

used to obtain energy scores as a measure of the likelihood of RNA-RNA interaction based on

sequence similarity. The more negative the energy score, the higher the likelihood of interaction.

Three top candidate RNA fragments derived from a lncRNA, a tRNA, and an rRNA were

selected based on predicted targeting of MexHI-OpmD and abundance in *P. aeruginosa*

following EV exposure. To assess whether targeting predictions extend beyond *P. aeruginosa* to

other common lung pathogens, we ran IntaRNA to predict targeting of MexHI-OpmD orthologs

by our top candidate RNA fragments in *Burkholderia cenocepacia* J2315, a multidrug-resistant

CF clinical isolate, and *Staphylococcus aureus* COL, a MRSA clinical isolate.

Statistical Analysis

Data were analyzed with Prism 8 for macOS (version 9.3.1, Graphpad, San Diego, CA)

and the R software environment for statistical computing and graphics version 4.1.1 (79) using

appropriate statistical methods, as indicated in the figure legends. Statistically significant

differences between bacterial growth curves were determined with permutation tests using the

compareGrowthCurves function from the R package statmod (version 1.4.36) (80,81).

Acknowledgements

The authors would like to thank Dr. Helen Zgurskaya (University of Oklahoma) for providing the

PAO1 deletion mutant for the fluoroquinolone efflux pump *mexGHI*-opmD as well as the matching

parental PAO1 wild type strain. Mass Spectrometry analyses of the rRNA fragment expressing

and empty vector strain of *P. aeruginosa* were performed by the Mass Spectrometry Technology

Access Center at McDonnell Genome Institute (MTAC@MGI) at Washington University School

of Medicine.

Figures

Fig 1. EVs increase the fluoroquinolone sensitivity of *P. aeruginosa*.

EVs did not significantly affect planktonic growth (OD600) of PA14 in the absence of

ciprofloxacin (CIP) after a 20-hour incubation (A), but significantly reduced the MIC of

ciprofloxacin (B) as well as planktonic growth of PA14 in the presence of 0.02 µg/ml

ciprofloxacin (C). EVs did not significantly affect planktonic growth (OD600) of the clinical isolate

SMC1585 in the absence of ciprofloxacin after a 20-hour incubation (D), but significantly

reduced the MIC of ciprofloxacin (E) as well as planktonic growth in the presence of 0.02 µg/ml

ciprofloxacin (F). Statistical significance was determined using unpaired t-tests. EVs were

isolated from four AEC donors and each data point is the mean of three technical replicates

from each donor. ***$P < 0.001$, *$P < 0.05$.

Fig 2. EVs increase *P. aeruginosa* ciprofloxacin sensitivity by targeting the
fluoroquinolone efflux pump MexHI-OpmD. EVs decreased planktonic growth of wild type
(WT) PAO1 in the presence of ciprofloxacin (0.03 µg/ml). EVs in the presence of ciprofloxacin
(0.03 µg/ml) had no significant effect on the *mexHI-opmD* knockout strain (Δ+EV). Statistical
significance was determined with repeated measures one-way ANOVA followed by Dunnett's
multiple comparisons test. EVs were isolated from six AEC donors and each data point is the
mean of three technical replicates from each donor. ***$P < 0.001$ compared to WT ctrl, **$P <$
0.01 compared to WT ctrl, ns = not significant.

Fig 3. A human rRNA fragment increases the fluoroquinolone sensitivity of *P.

aeruginosa. Compared to the empty vector control (ctrl), the rRNA fragment significantly

reduced planktonic growth (OD600) of PA14 in the presence of 0.03 µg/ml ciprofloxacin (CIP)

and 100 mM arabinose after a 20-hour incubation (A) but did not significantly affect planktonic

growth in the absence of ciprofloxacin (B). The rRNA fragment also significantly reduced the

minimal inhibitory concentration (MIC) of ciprofloxacin (C) as well as the number of live bacteria

in the presence of 0.03 µg/ml ciprofloxacin (D). Statistical significance was determined using

unpaired t-tests for panels A-C with N = four independent experiments using different clones of

each strain and each data point is the mean of three technical replicates and a negative

binomial regression model for panel D with N = three independent experiments using different

clones of each strain. Panel D shows the batch-corrected estimates from the regression model.

CFU = colony forming units. ****P < 0.0001 compared to ctrl, *P < 0.05 compared to ctrl.

Supporting information

S1 Fig. EVs increase the fluoroquinolone sensitivity of *P. aeruginosa*.

(A) 20-hour planktonic growth curves (OD600) of PA14 exposed to EVs or vehicle control (ctrl)

in the absence of ciprofloxacin. There was no significant difference between the growth curves

for the last 4 hours of growth. (B) 20-hour planktonic growth curves (OD600) of PA14 exposed

to 0.02 µg/ml ciprofloxacin and EVs or 0.02 µg/ml ciprofloxacin and vehicle control (ctrl). There

was a significant difference between the growth curves for the last 4 hours of growth. (C) 20-

hour planktonic growth curves (OD600) of clinical isolate SMC1585 exposed to EVs or vehicle

control (ctrl) in the absence of ciprofloxacin. There was no significant difference between the

growth curves for the last 4 hours of growth. (D) 20-hour planktonic growth curves (OD600) of

clinical isolate SMC1585 exposed to 0.02 µg/ml ciprofloxacin and EVs or 0.02 µg/ml

569 ciprofloxacin and vehicle control (ctrl). There was a significant difference between the growth

570 curves for the last 4 hours of growth. Each data point shows the mean +/- SEM of EVs isolated

571 from four AEC donors (or matching controls), averaged across three technical replicates.

572 Statistical significance was determined using permutation tests. * P < 0.05.

S2 Fig. EVs increase *P. aeruginosa* ciprofloxacin sensitivity by targeting the

fluoroquinolone efflux pump MexHI-OpmD. 20-hour planktonic growth curves (OD600) of

PAO1 wild type (WT) and the PAO1 *mexHI-opmD* knockout strain (Δ) exposed to 0.03 µg/ml

ciprofloxacin (CIP) in the presence and absence of EVs. There was a significant difference

between WT ctrl+CIP and all other growth curves for the last hour of growth, while here was no

significant difference between the other three growth curves. Each data point shows the mean

+/- SEM of EVs isolated from six AEC donors (or matching controls), averaged across three

technical replicates. Statistical significance was determined using permutation tests. * P < 0.05.

S3 Fig. A human rRNA fragment increases the fluoroquinolone sensitivity of *P.*

aeruginosa. (A) 20-hour planktonic growth curves (OD600) of PA14 expressing the rRNA

fragment or empty vector control (ctrl) in the presence of 0.03 µg/ml ciprofloxacin (CIP) and 100

mM arabinose. There was a significant difference between the growth curves for the empty

vector control (ctrl) and the rRNA expressing strain. (B) There was no significant difference in

the 20-hour planktonic growth curves (OD600) of PA14 expressing the rRNA fragment or empty

vector control in the absence of ciprofloxacin. Each data point shows the mean +/- SEM from

four different clones of each strain averaged across three technical replicates. Statistical

significance was determined using permutation tests. * P < 0.05.

S1 Table. Proteomics data of EV-exposed *P. aeruginosa* and unexposed controls.

Normalized peak intensities for each sample as well as average log2 fold changes and p-values

for 1619 proteins that were detected in duplicate runs of all six samples. EVs were derived from

three individual AEC donors.

S2 Table. Origin of most abundant sRNAs from the sequence-based characterization of

EV sRNA content. 909 most abundant unique sequences with at least 100 counts per sample

accounted for 60% of total reads for sequences detected in all 3 replicate samples. Listed are

the nucleotide sequence (column 1), sequence length (column 2), number of reads in EVs

(column 3) and origin of sequence as determined with BLAST+ (column 4). EVs were derived

from three individual AEC donors.

S3 Table. Proteomics data of *P. aeruginosa* expressing the rRNA fragment compared to

empty vector controls. Normalized peak intensities for each sample as well as average log2

fold changes and p-values for 1911 proteins that were detected in at least two of three replicate

samples.

References

References

1. Behzadi P, Baráth Z, Gajdács M. It's Not Easy Being Green: A Narrative Review on the
Microbiology, Virulence and Therapeutic Prospects of Multidrug-Resistant Pseudomonas
aeruginosa. Antibiotics (Basel) [Internet]. 2021 Jan 4 [cited 2021 Apr 12];10(1). Available
from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7823828/

2. Chong WH, Saha BK, Ananthakrishnan Ramani, Chopra A. State-of-the-art review of
secondary pulmonary infections in patients with COVID-19 pneumonia. Infection. 2021
Mar 11;1–15.

3. Giacobbe DR, Battaglini D, Enrile EM, Dentone C, Vena A, Robba C, et al. Incidence and
Prognosis of Ventilator-Associated Pneumonia in Critically Ill Patients with COVID-19: A
Multicenter Study. J Clin Med [Internet]. 2021 Feb 3 [cited 2021 Apr 12];10(4). Available
from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7913191/

4. Qu J, Cai Z, Liu Y, Duan X, Han S, Liu J, et al. Persistent Bacterial Coinfection of a
COVID-19 Patient Caused by a Genetically Adapted Pseudomonas aeruginosa Chronic
Colonizer. Front Cell Infect Microbiol. 2021;11:641920.

5. Yang S, Hua M, Liu X, Du C, Pu L, Xiang P, et al. Bacterial and fungal co-infections
among COVID-19 patients in intensive care unit. Microbes Infect [Internet]. 2021 Mar 5
[cited 2021 Apr 12]; Available from:
https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7933791/

6. Bullington W, Hempstead S, Smyth AR, Drevinek P, Saiman L, Waters VJ, et al.
Antimicrobial resistance: Concerns of healthcare providers and people with CF. Journal of
Cystic Fibrosis [Internet]. 2020 Jun 17 [cited 2021 Apr 12]; Available from:
https://www.sciencedirect.com/science/article/pii/S1569199320301594

7. López-Causapé C, Rojo-Molinero E, Macià MD, Oliver A. The problems of antibiotic
resistance in cystic fibrosis and solutions. Expert Review of Respiratory Medicine. 2015
Jan 2;9(1):73–88.

8. Tchakal-Mesbahi A, Metref M, Singh VK, Almpani M, Rahme LG. Characterization of
antibiotic resistance profiles in Pseudomonas aeruginosa isolates from burn patients. Burns.
2021 Mar 17;

9. Baiou A, Elbuzidi AA, Bakdach D, Zaqout A, Alarbi KM, Bintaher AA, et al. Clinical
characteristics and risk factors for the isolation of multi-drug-resistant Gram-negative
bacteria from critically ill patients with COVID-19. J Hosp Infect. 2021 Apr;110:165–71.

10. CDC. Antibiotic Resistance Threats in the United States, 2019. 2019;150.

11. Koeppen K, Hampton TH, Jarek M, Scharfe M, Gerber SA, Mielcarz DW, et al. A Novel
Mechanism of Host-Pathogen Interaction through sRNA in Bacterial Outer Membrane
Vesicles. PLoS Pathog. 2016 Jun;12(6):e1005672.

12.   Blenkiron C, Simonov D, Muthukaruppan A, Tsai P, Dauros P, Green S, et al.
648   Uropathogenic Escherichia coli Releases Extracellular Vesicles That Are Associated with
649   RNA. PLOS ONE. 2016;16.

650   13.   Choi J-W, Kim S-C, Hong S-H, Lee H-J. Secretable Small RNAs via Outer Membrane
651   Vesicles in Periodontal Pathogens. J Dent Res. 2017 Apr 1;96(4):458–66.

652   14.   Lefebvre FA, Lécuyer E. Small Luggage for a Long Journey: Transfer of Vesicle-Enclosed
653   Small RNA in Interspecies Communication. Front Microbiol [Internet]. 2017 Mar 16 [cited
654   2021 Apr 12];8. Available from:
655   http://journal.frontiersin.org/article/10.3389/fmicb.2017.00377/full

656   15.   Lee H-J. Microbe-Host Communication by Small RNAs in Extracellular Vesicles: Vehicles
657   for Transkingdom RNA Transportation. Int J Mol Sci. 2019 Mar 25;20(6).

658   16.   Munhoz da Rocha IF, Amatuzzi RF, Lucena ACR, Faoro H, Alves LR. Cross-Kingdom
659   Extracellular Vesicles EV-RNA Communication as a Mechanism for Host–Pathogen
660   Interaction. Front Cell Infect Microbiol. 2020 Nov 18;10:593160.

661   17.   Lécrivain A-L, Beckmann BM. Bacterial RNA in extracellular vesicles: A new regulator of
662   host-pathogen interactions? Biochimica et Biophysica Acta (BBA) - Gene Regulatory
663   Mechanisms. 2020 Jul 1;1863(7):194519.

664   18.   Zhang H, Zhang Y, Song Z, Li R, Ruan H, Liu Q, et al. sncRNAs packaged by Helicobacter
665   pylori outer membrane vesicles attenuate IL-8 secretion in human cells. International
666   Journal of Medical Microbiology. 2020 Jan 1;310(1):151356.

667   19.   Koeppen K, Nymon A, Barnaby R, Bashor L, Li Z, Hampton TH, et al. Let-7b-5p in
668   vesicles secreted by human airway cells reduces biofilm formation and increases antibiotic
669   sensitivity of P. aeruginosa. Proc Natl Acad Sci U S A. 2021 Jul 13;118(28):e2105370118.

670   20.   Vojtech L, Woo S, Hughes S, Levy C, Ballweber L, Sauteraud RP, et al. Exosomes in
671   human semen carry a distinctive repertoire of small non-coding RNAs with potential
672   regulatory functions. Nucleic Acids Res. 2014 Jun;42(11):7290–304.

673   21.   Chiou N-T, Kageyama R, Ansel KM. Selective Export into Extracellular Vesicles and
674   Function of tRNA Fragments during T Cell Activation. Cell Reports. 2018
675   Dec;25(12):3356-3370.e4.

676   22.   Abramowicz A, Story MD. The Long and Short of It: The Emerging Roles of Non-Coding
677   RNA in Small Extracellular Vesicles. Cancers. 2020 Jun 2;12(6):1445.

678   23.   Garcia-Silva MR, Cabrera-Cabrera F, Güida MC, Cayota A. Hints of tRNA-Derived Small
679   RNAs Role in RNA Silencing Mechanisms. Genes (Basel). 2012 Oct 10;3(4):603–14.

680   24.   Lee YS, Shibata Y, Malhotra A, Dutta A. A novel class of small RNAs: tRNA-derived
681   RNA fragments (tRFs). Genes Dev. 2009 Nov 15;23(22):2639–49.

25. Kumar P, Anaya J, Mudunuri SB, Dutta A. Meta-analysis of tRNA derived RNA fragments
reveals that they are evolutionarily conserved and associate with AGO proteins to recognize
specific RNA targets. BMC Biol [Internet]. 2014 Oct 1 [cited 2021 Apr 13];12. Available
from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4203973/

26. Raina M, Ibba M. tRNAs as regulators of biological processes. Front Genet [Internet]. 2014
Jun 11 [cited 2021 Apr 13];5. Available from:
https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4052509/

27. Avcilar-Kucukgoze I, Kashina A. Hijacking tRNAs From Translation: Regulatory
Functions of tRNAs in Mammalian Cell Physiology. Front Mol Biosci. 2020;7:610617.

28. Anderson P, Ivanov P. tRNA fragments in human health and disease. FEBS Lett. 2014 Nov
28;588(23):4297–304.

29. Xie Y, Yao L, Yu X, Ruan Y, Li Z, Guo J. Action mechanisms and research methods of
tRNA-derived small RNAs. Signal Transduction and Targeted Therapy. 2020 Jun
30;5(1):1–9.

30. Ogawa Y, Taketomi Y, Murakami M, Tsujimoto M, Yanoshita R. Small RNA
Transcriptomes of Two Types of Exosomes in Human Whole Saliva Determined by Next
Generation Sequencing. Biological & Pharmaceutical Bulletin. 2013;36(1):66–75.

31. Fatima F, Nawaz M. Vesiculated Long Non-Coding RNAs: Offshore Packages Deciphering
Trans-Regulation between Cells, Cancer Progression and Resistance to Therapies. Non-
Coding RNA. 2017 Mar;3(1):10.

32. Li Z, Stanton BA. Transfer RNA-Derived Fragments, the Underappreciated Regulatory
Small RNAs in Microbial Pathogenesis. Front Microbiol. 2021;12:687632.

33. Stanton BA. Extracellular Vesicles and Host-Pathogen Interactions: A Review of Inter-
Kingdom Signaling by Small Noncoding RNA. Genes (Basel). 2021 Jun 30;12(7):1010.

34. Fujita Y, Kosaka N, Araya J, Kuwano K, Ochiya T. Extracellular vesicles in lung
microenvironment and pathogenesis. Trends in Molecular Medicine. 2015 Sep;21(9):533–
42.

35. Fujita Y, Kadota T, Araya J, Ochiya T, Kuwano K. Extracellular Vesicles: New Players in
Lung Immunity. Am J Respir Cell Mol Biol. 2018 May;58(5):560–5.

36. Hough KP, Chanda D, Duncan SR, Thannickal VJ, Deshane JS. Exosomes in
immunoregulation of chronic lung diseases. Allergy. 2017 Apr;72(4):534–44.

37. McVey MJ, Maishan M, Blokland KEC, Bartlett N, Kuebler WM. Extracellular vesicles in
lung health, disease, and therapy. Am J Physiol Lung Cell Mol Physiol. 2019 Jun
1;316(6):L977–89.

38.   He S, Chen D, Hu M, Zhang L, Liu C, Traini D, et al. Bronchial epithelial cell extracellular
vesicles ameliorate epithelial–mesenchymal transition in COPD pathogenesis by alleviating
M2 macrophage polarization. Nanomedicine: Nanotechnology, Biology and Medicine.
2019 Jun;18:259–71.

39.   Stachowiak Z, Wojsyk-Banaszak I, Jończyk-Potoczna K, Narożna B, Langwiński W,
Kycler Z, et al. MiRNA Expression Profile in the Airways Is Altered during Pulmonary
Exacerbation in Children with Cystic Fibrosis—A Preliminary Report. J Clin Med
[Internet]. 2020 Jun 16 [cited 2021 Apr 15];9(6). Available from:
https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7356328/

40.   Chahar HS, Corsello T, Kudlicki AS, Komaravelli N, Casola A. Respiratory Syncytial
Virus Infection Changes Cargo Composition of Exosome Released from Airway Epithelial
Cells. Sci Rep [Internet]. 2018 Jan 10 [cited 2021 Apr 14];8. Available from:
https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5762922/

41.   Corsello T, Kudlicki AS, Garofalo RP, Casola A. Cigarette Smoke Condensate Exposure
Changes RNA Content of Extracellular Vesicles Released from Small Airway Epithelial
Cells. Cells. 2019 Dec 17;8(12).

42.   Armstrong DA, Nymon AB, Ringelberg CS, Lesseur C, Hazlett HF, Howard L, et al.
Pulmonary microRNA profiling: implications in upper lobe predominant lung disease. Clin
Epigenetics [Internet]. 2017 May 30 [cited 2021 Apr 6];9. Available from:
https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5450072/

43.   Gon Y, Shimizu T, Mizumura K, Maruoka S, Hikichi M. Molecular techniques for
respiratory diseases: MicroRNA and extracellular vesicles. Respirology. 2020
Feb;25(2):149–60.

44.   Guiot J, Struman I, Louis E, Louis R, Malaise M, Njock M-S. Exosomal miRNAs in Lung
Diseases: From Biologic Function to Therapeutic Targets. J Clin Med [Internet]. 2019 Aug
29 [cited 2021 Apr 15];8(9). Available from:
https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6781233/

45.   Liu S, da Cunha AP, Rezende RM, Cialic R, Wei Z, Bry L, et al. The Host Shapes the Gut
Microbiota via Fecal MicroRNA. Cell Host & Microbe. 2016 Jan;19(1):32–43.

46.   Liu S, Rezende RM, Moreira TG, Tankou SK, Cox LM, Wu M, et al. Oral Administration
of miR-30d from Feces of MS Patients Suppresses MS-like Symptoms in Mice by
Expanding Akkermansia muciniphila. Cell Host Microbe. 2019 Dec 11;26(6):779-794.e8.

47.   Teng Y, Ren Y, Sayed M, Hu X, Lei C, Kumar A, et al. Plant-Derived Exosomal
MicroRNAs Shape the Gut Microbiota. Cell Host Microbe. 2018 Nov 14;24(5):637-652.e8.

48.   Heijerman H, Westerman E, Conway S, Touw D, Döring G, consensus working group.
Inhaled medication and inhalation devices for lung disease in patients with cystic fibrosis:
A European consensus. J Cyst Fibros. 2009 Sep;8(5):295–315.

49.   Moreau-Marquis S, Coutermarsh B, Stanton BA. Combination of hypothiocyanite and
754   lactoferrin (ALX-109) enhances the ability of tobramycin and aztreonam to eliminate
755   Pseudomonas aeruginosa biofilms growing on cystic fibrosis airway epithelial cells. J
756   Antimicrob Chemother. 2015 Jan;70(1):160–6.

757   50.   Busch A, Richter AS, Backofen R. IntaRNA: efficient prediction of bacterial sRNA targets
758   incorporating target site accessibility and seed regions. Bioinformatics. 2008 Dec
759   15;24(24):2849–56.

760   51.   Mann M, Wright PR, Backofen R. IntaRNA 2.0: enhanced and customizable prediction of
761   RNA–RNA interactions. Nucleic Acids Res. 2017 Jul 3;45(Web Server issue):W435–9.

762   52.   Cherlin T. Ribosomal RNA fragmentation into short RNAs (rRFs) is modulated in a sex-
763   and population of origin-specific manner. 2020;19.

764   53.   Lambert M, Benmoussa A, Provost P. Small Non-Coding RNAs Derived From Eukaryotic
765   Ribosomal RNA. ncRNA. 2019 Feb 4;5(1):16.

766   54.   Guan L, Grigoriev A. Age-Related Argonaute Loading of Ribosomal RNA Fragments.
767   MIRNA. 2020 Mar 4;9(2):142–52.

768   55.   Guan L, Grigoriev A. Computational meta-analysis of ribosomal RNA fragments: potential
769   targets and interaction mechanisms. Nucleic Acids Research. 2021 Apr 19;49(7):4085–103.

770   56.   Chen Z, Sun Y, Yang X, Wu Z, Guo K, Niu X, et al. Two featured series of rRNA-derived
771   RNA fragments (rRFs) constitute a novel class of small RNAs. Xue Y, editor. PLoS ONE.
772   2017 Apr 25;12(4):e0176458.

773   57.   Wei H, Zhou B, Zhang F, Tu Y, Hu Y, Zhang B, et al. Profiling and Identification of Small
774   rDNA-Derived RNAs and Their Potential Biological Functions. Müller M, editor. PLoS
775   ONE. 2013 Feb 13;8(2):e56842.

776   58.   Czech B, Malone CD, Zhou R, Stark A, Schlingeheyde C, Dus M, et al. An endogenous
777   small interfering RNA pathway in Drosophila. Nature. 2008 Jun 5;453(7196):798–802.

778   59.   Lee H-C, Chang S-S, Choudhary S, Aalto AP, Maiti M, Bamford DH, et al. qiRNA, a new
779   type of small interfering RNA induced by DNA damage. Nature. 2009 May
780   14;459(7244):274–7.

781   60.   Gupta AK, Panigrahi SK, Bhattacharya A, Bhattacharya S. Self-circularizing 5′-ETS RNAs
782   accumulate along with unprocessed pre ribosomal RNAs in growth-stressed Entamoeba
783   histolytica. Sci Rep. 2012 Dec;2(1):303.

784   61.   Wolloscheck D, Krishnamoorthy G, Nguyen J, Zgurskaya HI. Kinetic control of quorum
785   sensing in Pseudomonas aeruginosa by multidrug efflux pumps. ACS Infect Dis. 2018 Feb
786   9;4(2):185–95.

62. Lister PD, Wolter DJ, Hanson ND. Antibacterial-Resistant *Pseudomonas aeruginosa* :
Clinical Impact and Complex Regulation of Chromosomally Encoded Resistance
Mechanisms. Clin Microbiol Rev. 2009 Oct;22(4):582–610.

63. Rehman A, Patrick WM, Lamont IL. Mechanisms of ciprofloxacin resistance in
Pseudomonas aeruginosa: new approaches to an old problem. Journal of Medical
Microbiology. 2019 Jan 1;68(1):1–10.

64. Giraldez MD, Spengler RM, Etheridge A, Goicochea AJ, Tuck M, Choi SW, et al.
Phospho-RNA-seq: a modified small RNA-seq method that reveals circulating mRNA and
lncRNA fragments as potential biomarkers in human plasma. EMBO J [Internet]. 2019 Jun
3 [cited 2021 Apr 16];38(11). Available from:
https://onlinelibrary.wiley.com/doi/10.15252/embj.2019101695

65. Zareba L, Szymanski J, Homoncik Z, Czystowska-Kuzmicz M. EVs from BALF-Mediators
of Inflammation and Potential Biomarkers in Lung Diseases. Int J Mol Sci. 2021 Apr
1;22(7).

66. Fulcher ML, Randell SH. Human nasal and tracheo-bronchial respiratory epithelial cell
culture. Methods Mol Biol. 2013;945:109–21.

67. Koeppen K, Nymon AB, Barnaby R, Li Z, Hampton TH, Ashare A, et al. CF Monocyte
Derived Macrophages Have an Attenuated Response to Extracellular Vesicles Secreted by
Airway Epithelial Cells. Am J Physiol Lung Cell Mol Physiol. 2021 Jan 20;

68. Shanks RMQ, Caiazza NC, Hinsa SM, Toutain CM, O'Toole GA. Saccharomyces
cerevisiae-based molecular tool kit for manipulation of genes from gram-negative bacteria.
Appl Environ Microbiol. 2006 Jul;72(7):5027–36.

69. Lambert RJW, Pearson J. Susceptibility testing: accurate and reproducible minimum
inhibitory concentration (MIC) and non-inhibitory concentration (NIC) values. Journal of
Applied Microbiology. 2000;88(5):784–90.

70. Malek AA, Wargo MJ, Hogan DA. Absence of Membrane Phosphatidylcholine Does Not
Affect Virulence and Stress Tolerance Phenotypes in the Opportunistic Pathogen
Pseudomonas aeruginosa. PLoS One [Internet]. 2012 Feb 17 [cited 2021 Apr 6];7(2).
Available from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3281885/

71. Scheller N, Herold S, Kellner R, Bertrams W, Jung AL, Janga H, et al. Proviral
MicroRNAs Detected in Extracellular Vesicles From Bronchoalveolar Lavage Fluid of
Patients With Influenza Virus–Induced Acute Respiratory Distress Syndrome. The Journal
of Infectious Diseases. 2019 Jan 29;219(4):540–3.

72. Choi H, Kim S, Fermin D, Tsou C-C, Nesvizhskii AI. QPROT: statistical method for
testing differential expression using protein-level intensity data in label-free quantitative
proteomics. J Proteomics. 2015 Nov 3;129:121–6.

73.   Deutsch EW, Bandeira N, Sharma V, Perez-Riverol Y, Carver JJ, Kundu DJ, et al. The
ProteomeXchange consortium in 2020: enabling "big data" approaches in proteomics.
Nucleic Acids Res. 2020 Jan 8;48(D1):D1145–52.

74.   Wang M, Wang J, Carver J, Pullman BS, Cha SW, Bandeira N. Assembling the
Community-Scale Discoverable Human Proteome. Cell Syst. 2018 Oct 24;7(4):412-421.e5.

75.   Perez-Riverol Y, Bai J, Bandla C, García-Seisdedos D, Hewapathirana S, Kamatchinathan
S, et al. The PRIDE database resources in 2022: a hub for mass spectrometry-based
proteomics evidences. Nucleic Acids Res. 2022 Jan 7;50(D1):D543–52.

76.   Chen S, Zhou Y, Chen Y, Gu J. fastp: an ultra-fast all-in-one FASTQ preprocessor.
Bioinformatics. 2018 Sep 1;34(17):i884–90.

77.   Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, et al. BLAST+:
architecture and applications. BMC Bioinformatics. 2009 Dec 15;10:421.

78.   Winter DJ. rentrez: An R package for the NCBI eUtils API. The R Journal. 2017;9(2):520–
6.

79.   R Core Team. R: A language and environment for statistical computing. [Internet]. Vienna,
Austria: R Foundation for Statistical Computing; 2021. Available from: URL
https://www.R-project.org/

80.   Baldwin T, Sakthianandeswaren A, Curtis JM, Kumar B, Smyth GK, Foote SJ, et al.
Wound healing response is a major contributor to the severity of cutaneous leishmaniasis in
the ear model of infection. Parasite Immunology. 2007;29(10):501–13.

81.   Elso CM, Roberts LJ, Smyth GK, Thomson RJ, Baldwin TM, Foote SJ, et al. Leishmaniasis
host response loci (lmr1–3) modify disease severity through a Th1/Th2-independent
pathway. Genes Immun. 2004 Mar;5(2):93–100.