Cross-modal interaction of Alpha Activity does not reflect inhibition

of early sensory processing: A frequency tagging study using EEG

and MEG

Brickwedde, M.[1,2,3], Limachya, R.[3], Markiewicz, R. [3,4], Sutton, E. [3], Postzich, C.[5], Shapiro, K.

[3], Jensen, O[3]., and Mazaheri, A[3].

[1]Department of Child and Adolescent Psychiatry, Charité - Universitätsmedizin Berlin, Germany

[2]Physikalisch Technische Bundesanstalt, Berlin, Germany

[3]Centre for Human Brain Health, School of Psychology, University of Birmingham, Uk

[4]Istituto Italiano di Tecnologia, Genova, Italy

[5]Max-Planck Research Institute, Leipzig, Germany

Abstract

Selective attention involves prioritizing relevant sensory input while suppressing irrelevant

stimuli. It has been proposed that oscillatory alpha-band activity (~10 Hz) aids this process by

functionally inhibiting early sensory regions. However, recent studies have challenged this

notion. Our EEG and MEG studies aimed to investigate whether alpha oscillations serve as a

'gatekeeper' for downstream signal transmission. We first observed these effects in an EEG

study and then replicated them using MEG, which allowed us to localize the sources.

We employed a cross-modal paradigm where visual cues indicated whether upcoming targets

required visual or auditory discrimination. To assess inhibition, we utilized frequency-

tagging, simultaneously flickering the fixation cross at 36 Hz and playing amplitude-

modulated white noise at 40 Hz during the cue-to-target interval.

Consistent with prior research, we observed an increase in posterior alpha activity following

cues signalling auditory targets. However, remarkably, both visual and auditory frequency

tagged responses amplified in anticipation of auditory targets, correlating with alpha activity

amplitude. Our findings suggest that when attention shifts to auditory processing, the visual

stream remains responsive and is not hindered by occipital alpha activity. This implies that

alpha modulation does not solely regulate 'gain control' in early visual areas but rather

orchestrates signal transmission to later stages of the processing stream.

Introduction

In our daily life, we are often confronted with sensory information from many different sources, all at once. To operate effectively, we require selective attention, reconciling the tension between environmental inputs relevant for top-down goals and sensory information that may be perceptually salient but task-irrelevant. Previous research gave rise to the prominent alpha inhibition hypothesis, which suggests that oscillatory activity in the alpha range (~10 Hz) plays a mechanistic role in selective attention through functional inhibition of irrelevant cortices (see Fig. 1A; Foxe et al., 1998; Jensen & Mazaheri, 2010; Klimesch et al., 2007). Functional inhibition refers to an area of the cortex being actively hindered to process input, which is distinctly different from idling, where a part of the cortex is not actively involved. Evidence supporting this theory revealed an increase in alpha-power over task-irrelevant sensory cortices after the onset of cues indicating the spatial location (Kelly et al., 2006; Okazaki et al., 2014; Thut et al., 2006; Worden et al., 2000; Zumer et al., 2014) or specific modality of an upcoming target (Foxe et al., 1998; Fu et al., 2001; Mazaheri et al., 2014). Moreover, previous investigations have observed 'spontaneous fluctuations' of alpha power in sensory regions, particularly in the visual cortex, to be inversely related to discrimination ability (Ergenoglu et al., 2004; Van Dijk et al., 2008). Alpha inhibition is believed to be transmitted in a phasic manner, as phosphene perception as well as high-frequency and spiking activity vary in line with the alpha cycle (Dugué et al., 2011; Haegens et al., 2011; Spaak et al., 2012).

Recent evidence challenged a direct connection between alpha activity and visual information processing in early visual cortex. As such, both visual steady-state responses and alpha power were modulated by attention, but did not covary when investigating individual trials (Zhigalov & Jensen, 2020). Unfortunately, very few studies have investigated direct connections between alpha activity, attention and sensory signals, especially over trials. Furthermore,

results seem to depend on timing of alpha activity in relation to sensory responses as well as

stimulus type and outcome measure (Morrow et al., 2023).

Accordingly, the objective of the current study is to test the alpha inhibition hypothesis

compared to an alternative theory. Based on the alpha inhibition hypothesis, alpha modulation

is connected to 'gain control' in early visual areas through modulation of excitability (Foxe &

Snyder, 2011; Jensen & Mazaheri, 2010; Van Diepen et al., 2019). In contrast, we propose

that inhibitory effects of alpha modulation are exhibited at later stages of the processing

stream (Peylo et al., 2021; Yang et al., 2023; Zhigalov & Jensen, 2020; Zumer et al., 2014),

gating feedforward or feedback communication between sensory areas (see Fig. 1; Bauer et

al., 2020; Haegens et al., 2015; Uemura et al., 2021).

To this end, we applied frequency-tagging, the rhythmic presentation of sensory stimuli,

which elicits steady-state sensory evoked potentials or fields (SSEP/SSEF), consisting of

rhythmic neuronal activity in the frequency of stimulation (Brickwedde et al., 2020; Colon et

al., 2012; Dinse et al., 1998; Marzoll et al., 2018; Regan, 1982; Snyder, 1992; Stapells et al.,

1984; Tobimatsu et al., 1999). The magnitude of SSEPs is attention-dependent (de Jong et al.,

2010; Müller et al., 1998; Müller & Hillyard, 2000; Porcu et al., 2013; Saupe et al., 2009;

Toffanin et al., 2009) even for frequencies too fast to perceive consciously (Brickwedde et al.,

2022; Zhigalov et al., 2019). Visual SSEP/SSEFs are most strongly observable over occipital

areas, whereas auditory SSEP/SSEFs appear most strongly over temporal (MEG) or fronto-to-

central (EEG) areas (de Jong et al., 2010; Hari et al., 1989; Pantev et al., 1996; Regan, 1982).

Furthermore, when applying two different frequencies for two different sensory modalities,

their intermodulation frequency (f1-f2) has been suggested to reflect cross-modal integration

(Drijvers et al., 2021). Due to distinct responses, localisation and attention-dependence,

frequency-tagging provides an optimal tool to study sensory signal processing and integration

over time.

The aim of our study was to directly test the alpha inhibition hypothesis by investigating if

cue-induced modulation of alpha activity coincides with the suppression of frequency-tagging

responses in task-irrelevant modalities. Based on previous studies, we utilized a cross-modal

attention paradigm, in which symbolic visual cues signalled the target modality (visual or

auditory) of an upcoming discrimination task (e.g., van Diepen et al., 2015). Here, we

included an additional experimental manipulation in the form of frequency-tagging to assess

the involvement of the auditory and visual systems in the *cue-to-target interval*.

Figure 1. Illustration of the alpha inhibition hypothesis. The alpha inhibition hypothesis suggests that occipital
alpha inhibits visual information processing in a phasic manner. If alpha activity is high, it suggests that the
whole area is inhibited and thereby disengaged (Foxe et al., 1998; Jensen & Mazaheri, 2010; Klimesch et al.,
2007). We propose a revision of this hypothesis, whereby alpha activity exerts its phasic inhibition to regulate
downstream information transfer, creating enhanced signal packages of prioritised information (see also Yang
et al., 2023; Zhigalov & Jensen, 2020; Zumer et al., 2014).

In line with previous results, we hypothesized that signalling an upcoming auditory target

would lead to increased alpha activity over visual occipital regions as well as increased SSEP

responses to auditory and decreased SSEP to visual stimuli as indexed by frequency-tagging.

In brief, while we observed the expected cue-induced early-visual alpha modulation, the

amplitude of auditory and *visual* SSEP/SSEFs as well as their intermodulation frequency

increased just prior to the onset of the auditory target, contradicting the alpha inhibition

hypothesis. The difference between conditions of visual SSEP/SSEFs originated from sensory

integration areas and correlated with early sensory alpha activity on a trial-by-trial basis,

speaking to an effect of alpha modulation on signal transmission rather than inhibition of

early visual areas.

Results Study 1 – cross-modal EEG experiment

To assess audio-visual excitability in anticipation of either visual or auditory targets, we

displayed visual cues to signal the modality of the upcoming target (auditory, visual, or non-

specific). In a three-second cue-to-target interval, we frequency-tagged the fixation cross at 36

Hz and played 40 Hz amplitude modulating white noise. Immediately afterwards, participants

had to either discriminate between three different pitch sounds (auditory target) or three

different Gabor patch orientations (visual target). If the target modality was cued (e.g. not

non-specific), 50% of the trials were accompanied by a random distractor from the target pool

of the opposing sensory modality (see Fig. 2A).

118
119
Behavioural performance

We found that accuracy differed significantly between conditions ($F_{(5,105)} = 44.16$; $p < .001$).

Overall, participants were significantly less accurate in the auditory discrimination task

('overall auditory', $M = 79\%$ correct, $SD = 11.7$) than in the visual discrimination task

('overall visual', $M = 97\%$, $SD = 2.3$; see Fig. 2B), with the worst performance occurring

when auditory targets were paired with visual distractors (auditory +: $M = 74\%$ correct, $SD =$

13).

Reaction times yielded a similar pattern, with auditory reactions (overall auditory: $M = 662$

ms, $SD = 136$) being slower than for the visual task (overall visual: $M = 597$ ms, $SD = 130$;

main effect over all conditions: $F_{(5,105)} = 27.47$; $p < .001$; see Fig. 2C). This was mostly driven

by slow responses to auditory targets paired with distractors (auditory +: $M = 723$ ms).

Figure 2. A, Illustration of the cross-model discrimination task in Study 1. Trials were separated by a 4 s
interval, in which a fixation cross was displayed. A brief central presentation of the cue (100 ms) initiated the
trial, signalling the target modality (see figure above from left to right: auditory, non-specific, visual). In the
cue-to-target interval, the fixation cross was frequency-tagged at 36 Hz. At the same time, a sound was
displayed over headphones, which was frequency-tagged at 40 Hz. Both tones and fixation cross contained no
task-relevant information. The target, consisting either of a static Gabor patch or a tone, was presented for 25
ms. Participants had to differentiate via button presses between 3 different Gabor rotations or tone pitches,
respectively. In 50% of auditory and visually cued trials, a distractor in form of a random pitch or rotation of the
un-cued modality was presented alongside the target. **B-C, Analysis of task accuracy and reaction time**
indicates increased difficulty of auditory targets. B, Task accuracy compared between all 6 experimental
conditions reveals a drop in accuracy for responses to auditory targets. **C,** reaction times of correct trials
compared between all 6 experimental conditions. The slowest reaction times are observable following auditory
targets alongside visual distractors. N = 22; *** sig < .001; ** sig. < .01; * sig. < .05;
145

To ascertain that cues and distractors were functionally relevant, we calculated attentional

benefit (cued - non-specific condition) and distractor cost (cued condition with distractor –

cued condition without distractor). As expected, cues improved reaction times over all

conditions and distractors impaired the accuracy over all conditions and reaction times for

auditory targets. (see suppl Fig. 1). Interestingly, auditory distractors reduced reaction times

to visual targets, which could be explained by a generally faster processing of auditory targets

(Jain et al., 2015). As such, the auditory distractor possibly caused intersensory facilitation

(Nickerson., 1973), whereby reaction times to a target can be facilitated when accompanied

by stimuli of other sensory modalities, even if they are irrelevant or distracting.

Figure 3. Post-cue modality specific modulation of alpha power in anticipation of an auditory versus a visual
target A-B, the time course of post-cue alpha power. Cluster permutation analysis resulted in two condition
effects, both indicating heightened alpha activity when expecting an auditory compared to a visual target (A:
p < .01; B: p < .01). C-D, Time-frequency representation of power in the cue-to-target interval. A greater
increase in alpha power was observed when expecting an auditory target (average over significant electrodes
for the condition difference in A). note: Cluster electrodes are marked in white. Shading represents standard
error from the mean; Δ / Σ represents (a-b)/(a+b) normalization.

Cross-modal cues differentially modulated pre-target alpha activity

We conducted a time-frequency analysis of power in the cue-to-target interval and found a

stronger amplitude increase from baseline for auditory compared to visual target conditions

starting around 2 s before target onset (Fig. 3). We calculated the time course of alpha power

changes using the Hilbert-transformation (8 – 12 Hz, Fig. 3A-B). We then applied cluster

permutation analysis, whereby real condition differences were tested against coincidental

findings by randomly permutating the condition labels to the data and testing for condition

differences 1000 times (Maris & Oostenveld, 2007). Consistent with previous work (Mazaheri

et al., 2014; van Diepen & Mazaheri, 2017), analysing the last two seconds before target onset

revealed two significant clusters of difference in alpha power when expecting an auditory

compared to a visual target. These effects corresponded to clusters extending from -1.84 to –

0.64 s (p = .004) and -0.62 to 0s (p = .005). When presenting non-specific cues, alpha power

changes were not significant, but descriptively larger compared to visual target conditions and

lower compared to auditory target conditions (see suppl Fig. 2). However as significant alpha

modulation was a prerequisite to test our hypotheses, we excluded this condition from further

analysis.

Cross-modal cues increased the amplitude of the frequency-tagged responses across both
modalities

To assess the temporal development of frequency-tagging responses, steady-state potentials

were calculated using data band-pass filtered around the tagging frequency. Neuronal 40 Hz

auditory-steady state evoked potential (ASSEP) responses were strongest over central areas

(Fig. 4A), and 36 Hz auditory-steady state evoked potential (VSSEP) responses were

strongest over occipital areas (Fig. 4B).

To assess the differences between conditions, the Hilbert envelope of the steady-state

potentials was analysed using cluster permutation analyses. When expecting an auditory

target compared to a visual target, the 40 Hz ASSEP response was larger shortly before target

onset (see Fig. 4C; -0.15 to -0.08 s; p = .041).

Figure 4. Increase in amplitude of both visual and auditory frequency tagged responses when anticipating visual or auditory targets. Event-related potentials and scalp topographies reveal distinct modality specific responses at the tagged frequencies. A, auditory steady-state evoked potential (ASSEP) averaged over 6 central electrodes displaying the highest 40 Hz power (Cz, FC1, FC2, C1, C2, FCz; marked with white squares). B, visual steady-state evoked potential (VSSEP) averaged over 4 occipital electrodes displaying the highest 36 Hz power (POz, O1, O2, Oz; marked with white squares) C, the Hilbert-envelope of the 40 Hz ASSEP reveals an increase shortly before target onset when anticipating an auditory compared to a visual target (p = .041); D, the Hilbert-envelope of the 36 Hz VSSEP likewise reveals an increase shortly before target onset when anticipating an auditory compared to a visual target (p = .014). *note*: Cluster electrodes are marked in white. Shading represents standard error from the mean. Δ / ∑ represents (a-b)/(a+b) normalization.

Surprisingly, the 36 Hz VSSEP response was likewise increased shortly before expecting an auditory compared to a visual target (see Fig. 4D; -0.16 to -0.06 s; p = .014). As such, the observed frequency tagging responses might reflect effort, which affects the vigilance of the whole sensory system rather than sensory-specific allocations of attention. For both 36 Hz VSSEP and 40 Hz ASSEP responses, condition differences appeared strongest over mid-parietal regions in contrast to primary sensory cortices.

Alpha power was positively correlated with amplitude of frequency tagged responses

Following the observation of condition differences in alpha activity and frequency-tagging responses, we were further interested in exploring whether these responses were connected.

Accordingly, we conducted trial-by-trial correlations using alpha condition differences and

their electrode positions as seed, which was correlated with frequency-tagging signals over all

electrodes. Multiple comparison correction was applied by testing the correlation matrix

against a zero-correlation matrix with a cluster permutation approach. A positive correlation

was observed over right parietal-to-occipital areas between the late alpha cluster activity and

both 40 Hz ASSEP ($p = .009$) and 36 Hz VSSEP ($p = .004$) responses when expecting a visual

target (see Fig. 5.A-B). To estimate the robustness of these results, we additionally conducted

median split analyses between trials with high and low alpha power for each participant, as

well as averaged the correlation coefficient of each participant and calculated a one-sample t-

test against 0. For each analysis we provided the Bayes Factor, which estimates the strength of

support for or against the null hypothesis (BF > 3.2 is considered as substantial evidence and

BF > 10 is considered as strong evidence; Kass & Raftery, 1995).

The median split was highly significant for the 36 Hz VSSEP response (Fig. 5A, middle

columns, $p = .033$; $t_{(19)} = 2.29$; $BF_{(10)} = 1.91$) but did not reach significance for the 40 Hz

ASSEP response (Fig. 5B, middle column; $p = 0.20$; $t_{(19)} = 1.32$; $BF_{(10)} = 0.49$). Correlation

coefficients indicated strong correlations with alpha activity for both 40 Hz ASSEP (Fig. 5B,

right column; $p < .001$; $t_{(19)} = 4.95$; $BF_{(10)} = 306.93$) and 36 Hz VSSEP activity (Fig. 5A, right

column; $p < .01$; $t_{(19)} = 3.66$; $BF_{(10)} = 23.57$).

The same positive correlation with alpha activity was found when expecting an auditory target

for 36 Hz VSSEP activity ($p = .031$), but not for 40 Hz ASSEP activity (see Fig. 5D). A

median split between high and low alpha activity ($p = .005$; $t_{(19)} = 3.14$; $BF_{(10)} = 8.53$) and

correlation coefficients ($p = .002$; $t_{(19)} = 3.52$; $BF_{(10)} = 17.76$) provided moderate to strong

evidence for this effect. The likely origin of alpha modulations from early visual cortices may

explain a stronger connection to 36 Hz VSSPE responses compared to 40 Hz ASSEP

responses.

Additionally, we compared how correlation coefficients between alpha activity and

frequency-tagging differed when anticipating an auditory versus a visual target. Multiple

comparison correction was applied with cluster permutation analysis. Interestingly, an

interaction between the strength of the correlation associating alpha, 36 Hz VSSEP activity

and condition became apparent ($p = .044$; *see* Fig. 5E) and was observed most strongly over

right-central electrodes. Comparing the correlation coefficients of participants over this

cluster revealed a strong effect and even a change of direction in the correlation ($p < .001$;

$t_{(21)} = -4.76$; $BF_{(10)} = 259.97$). Particularly, when expecting a visual target, there was a

negative correlation between 36 Hz VSSEP and alpha activity, which turned positive when

expecting an auditory target. Both correlations also differed significantly from 0 (expecting a

visual target: $p < .001$; $t_{(21)} = -3.87$; $BF_{(10)} = 39.62$; expecting an auditory target: $p = .020$;

$t_{(21)} = 2.53$; $BF_{(10)} = 2.83$). In contrast to the previously observed positive correlation between

alpha activity and 36 Hz VSSEP activity, the significant electrode cluster was located more

ventrally. This effect could possibly hint at dynamic adaptability of oscillatory alpha effects

on later processing stages.

It is further noteworthy that the correlation between alpha activity and 36 Hz VSSEP response

when expecting a visual target, was also present when utilizing the early alpha cluster as seed

(~ 1200 to 400 ms before target onset, see Fig. 5C), in which case alpha modulation greatly

preceded the 36 Hz VSSEP modulation ($p = .016$). For this correlation, the median split

between high and low alpha trials did not reach significance ($p = .11$; $t_{(20)} = 1.69$;

$BF_{(10)} = 0.76$), however testing the correlation coefficients against 0 again revealed a

significant effect ($p = .003$; $t_{(20)} = 3.39$; $BF_{(10)} = 14.22$). This result may support the

directionality of alpha modulation affecting visual information processing (and not vice

versa).

Figure 5. Relationship between cue induced alpha modulation and amplitude of frequency tagged responses in study 1. Previously obtained alpha clusters (see Fig. 3) were correlated over trials with auditory 40 Hz and visual 36 Hz clusters (see Fig. 4), where alpha electrodes/sensors were applied as seeds. The analysis was performed using a cluster-permutation approach, testing a correlation model against a 0-correlation model. Clusters significantly diverging from the 0-correlation model are presented topographically. Additionally, median splits between high and low alpha trials as well as correlation coefficients of these clusters are displayed for all participants **A-B,** a positive correlation is visible between alpha activity in the last 400 ms and steady state potentials shortly before target onset when expecting a visual target (visual 36 HZ: *p* = .013; 40 Hz: *p* = .009). **D,** when expecting an auditory target, there is a positive correlation between alpha activity in the last 400 ms and visual 36 Hz activity shortly before target onset (*p* = .010). **E,** the correlation between alpha activity 400 ms and visual 36 Hz activity shortly before target onset changes its direction depending on whether an auditory or a visual target is expected (*p* = .037). **C,** a positive correlation is also visible between alpha activity as early as ~1200 ms to 400 ms and visual 36 Hz activity shortly before target onset when expecting a visual target (*p* = .016). N = 22; *** sig < .001; ** sig < .01; * sig < .05. + sig. < .1

Intermodulation frequency

Lastly, we analysed the steady-state response of the intermodulation frequency at 4 Hz. Increased intermodulation shortly before target onset could be observed when expecting an auditory compared to a visual target (-0.51 to -0.0620 s; *p* < .001). This effect was strongest over left fronto-to-central electrodes and right central-to-occipital electrodes (see Fig. 6A) and may suggest in increase in cross-modal integration.

With the goal to examine whether there are any behavioral effects of the condition difference,

trial by trial correlations with reaction times of each of the 2 auditory and visual conditions

were performed. Only in the easiest condition, when expecting a visual target that was not

accompanied by a distractor, a negative correlation with reaction time could be found,

strongest over right central electrodes (see Fig. 6B; $p = .046$). While a median split between

slow and fast trials did not reach significance ($p = .50$; $t_{(21)} = -0.69$; $BF_{(10)} = .28$), testing the

correlation coefficients against 0 revealed strong evidence for a correlation ($p = .004$;

$t_{(21)} = -3.28$; $BF_{(10)} = 11.98$). If the intermodulation frequency reflects cross-modal integration

(Drijvers et al., 2021), this effect would indicate faster reaction times for trials with stronger

sensory integration.

Figure 6. Steady-state response in the intermodulation frequency and its behavioural relevance. A, the Hilbert-envelope of the 4 Hz steady-state response reveals an increase shortly before target onset when anticipating an auditory compared to a visual target ($p < .01$). **B**, there is a trial-by-trial correlation between 4 Hz activity and reaction time when a visual target without distractor was presented. The correlation is further illustrated by a median split between fast and slow reaction time trials as well as by correlation coefficients for each participant. N = 22; ** sig. < .01;

Results Study 2 – cross-modal MEG experiment

To test the robustness of our results and to employ additional control analyses, we replicated our experiment using MEG (see Fig. 7A). While an increase in visual information processing parallel to an increase in alpha modulation already contradicts the notion of alpha inhibition exerting "gain control", affecting the whole visual cortex, our claim that alpha modulation instead affects visual information at later processing stages still required further validation. As such, our goal was to perform source analyses showing alpha modulation originating from primary visual areas affected visual information at later processing stages (e.g. not in primary visual cortex). Additionally, to exclude that the uncertainty over possible distractors affected our results, we employed a block design, where block 1 consisted only of trials without distractors and in block 2 targets were always accompanied by a distractor. Furthermore, we aligned the visual and auditory task to be more similar, both of them now featuring frequency-discrimination, which related to sound pitch (frequency) in the auditory condition and stripe-frequency of the Gabor patch in the visual condition. Lastly, to make sure our effects were driven by sensory modality-differences rather than task-difficulty differences, we included a short calibration phase. Prior to the experiment, difficulty of pitch sounds, and Gabor patch frequency were calibrated for each individual, ascertaining a success rate between 55% to 75%.

Figure 7. A, Illustration of the cross-model discrimination task in Study 2. Trials were separated by a 1-1.3 s interval, in which a fixation cross was displayed. A brief central presentation of the cue (100 ms) initiated the trial, signalizing the target modality (from left to right: auditory, visual). In the cue-to-target interval, the fixation cross was frequency-tagged at 36 Hz. At the same time, a sound was displayed over headphones, which was frequency-tagged at 40 Hz. Both tones and fixation cross contained no task-relevant information. The target, consisting either of a static Gabor patch or a tone, was presented for 25 ms. Participants had to differentiate via button presses between 3 different Gabor stripe frequencies or tone pitches, respectively. In block 2, a distractor in form of a random sound or Gabor patch of the un-cued modality was always presented alongside the target. B-C, analysis of task accuracy and reaction time indicates comparable difficulties in block 2. B, task accuracy differences were only observable in the first block without distractors. C, reaction times to visual targets in the first block were strongly increased compared to all other conditions. In the second block, no significant difference in reaction times was observable. N = 27; *** sig < .001; ** sig. < .01; * sig. < .05;

Behavioural performance

Our adjustments in the MEG-study streamlined performances to be more in line between auditory and visual conditions in the second block, however in the first block there were strong condition differences (Main effect condition: ($F_{(3,75)}$ = 10.26; $p < .01$). In block 1, participants were significantly more accurate in the auditory ('block 1 auditory', M = 84%

correct, $SD = 10.03$) compared to the visual task ('block 1 visual', $M = 70\%$, $SD = 10.61$; see

Fig. 7B). However, in block 2, there was no observable difference between visual and

auditory task accuracy ('block 2 auditory', $M = 77\%$ correct, $SD = 10.03$; ('block 2 visual', M

$= 78\%$, $SD = 13.15$). Reaction times were comparable between conditions (overall: $M = 661$

ms, $SD = 76$), with the exception of responses to visual targets in block 1 (block 1 visual:

$M = 726$ ms, $SD = 85$), which were significantly slower than all other reaction times (main

effect condition: $F_{(3,75)} = 18.13$; $p < .001$; see Fig. 7C).

Due to the block design, distractor cost could not be differentiated from learning effects. For

auditory targets, there was a significant decrease in accuracy between block 1 and 2, while for

visual targets, there was a significant increase. Similar to study 1, reaction times for auditory

targets stayed roughly the same, but became shorter for visual targets (see suppl Fig. 1).

Cross-modal cues differentially modulated pre-target alpha activity in early visual areas

In line with previous studies (van Diepen & Mazaheri, 2017), condition differences in alpha

activity were only significant in block 2, where distractors were always present. As alpha

modulation was a prerequisite to test our hypotheses, we performed the following analyses

solely with data from block 2 (see Fig. 8).

Replicating our results from study 1, cluster permutation on the Hilbert-transformed alpha

activity revealed a significant cluster between -1.47 to -1.18 s (p = .034), where alpha change

from baseline was higher when expecting an auditory compared to a visual target (see Fig.

8A). Applying roughly the cluster time-window (-1.5 and -1 s), we conducted a source

localization, contrasting the two conditions with cluster permutation analysis. We found that

condition differences in alpha activity originated from early visual areas, albeit with a

stronger effect on the right compared to the left hemisphere (p < .01; peak spm coordinates:

41 -82 -19 mm, in the right lingual gyrus see Fig. 8B). Additionally, we replicated this effect

with a virtual channel analysis in V1 (see SUPPL Fig. 12)

Figure 8. Post-cue modality specific early visual modulation of alpha power in anticipation of an auditory versus a visual target A, the time course of post-cue alpha power. Cluster permutation analysis resulted in a condition effects, indicating heightened alpha activity when expecting an auditory compared to a visual target (p = .034). **B,** source localization of the condition difference between expecting an auditory versus a visual target, revealing a significant cluster in early visual areas with stronger effects on the right hemisphere (p < .01). **C-D,** Time-frequency representation of power in the cue-to-target interval (average over electrodes that showed maximal condition difference in A). *note:* Cluster electrodes are marked in white. Shading represents standard error from the mean; Δ / ∑ represents (a-b)/(a+b) normalization.

Cross-modal cues increased the amplitude of the frequency-tagged responses across both modalities

In accordance with study 1, neuronal responses to 40 Hz auditory steady-state evoked field (ASSEF) responses were strongest over temporal areas (Fig. 9A) and 36 Hz visual steady-state evoked field (VSSEF) responses were strongest over occipital areas (Fig. 9C). As expected, the auditory tagging response originated from the right-hemispheric early auditory cortex (cluster significance: p < .01; peak spm coordinates: 69 -22 5 mm, in the right superior temporal gyrus; see Fig. 9B) and the visual tagging response originated from the early visual cortex (cluster significance: p < .001; peak spm coordinates: 19 -105- 11 mm, in the right lingual gyrus; see Fig. 9D). Additionally, we observed significantly reduced 40 Hz activity in the left-hemispheric visual-to-central cortex (cluster significance: p < .01).

386   anticipating auditory target ▬▬▬
        anticipating visual target ▬▬▬▬

Figure 9. Post-cue modality specific early visual modulation of alpha power in anticipation of an auditory versus a visual target. A, auditory steady-state evoked fields (ASSEF) averaged over 24 temporal sensors displaying the highest auditory 40 Hz power (12 right, 12 left; marked in white squares). **B,** ASSEF source localization revealed a significant positive cluster in the right-hemispheric early auditory cortex ($p < .001$). **C,** visual steady-state evoked fields (VSSEF) averaged over 10 occipital sensors, displaying the highest visual 36 Hz power (marked in white squares). **D,** VSSEF source localization revealed a significant positive cluster in the early visual cortex ($p < .001$). **E,** the Hilbert-envelope of the 40 Hz ASSEF reveals an increase shortly before target onset when anticipating an auditory compared to a visual target ($p = .043$); **F,** condition differences in the 40 Hz ASSEF response did not reach significance in sensor space. **G,** the Hilbert-envelope of the 36 Hz VSSEF likewise reveals an increase shortly before target onset when anticipating an auditory compared to a visual target ($p = .019$). **H,** condition differences in the 36 Hz VSSEF response were significant over several areas of the visual stream, including most strongly the medial occipital cortex, the calcarine fissure, and the precuneus ($p = .047$); *note*: Cluster electrodes are marked in white. Shading represents standard error from the mean. Δ / Σ represents (a-b)/(a+b) normalization.

Applying cluster permutation on the Hilbert-transformed frequency-tagging data over the last

500 ms before target onset replicated the results of study 1, where both 36 Hz VSSEF as well

as 40 Hz ASSEF signals were significantly enhanced when expecting an auditory compared to

a visual target (see Fig. 9E-G; auditory target: $p = .043$; visual target: $p = .019$).

Source localization confirmed the condition difference in 36 Hz VSSEF activity to originate

from later stages of the processing stream, encompassing a wide range of areas, most strongly

the medial occipital cortex and the precuneus (cluster significance: $p = .047$; peak spm

coordinates: 3 -66 44 mm, in the left and right precuneus; see Fig. 9H). In source space, the

effect was not significant for auditory 40 Hz activity (p = .11; peak spm coordinates: -9 -39 0

mm, in the left precuneus; see Fig. 9F). Furthermore, a virtual channel analysis in V1 and

Heschl's gyrus confirmed that there were no condition differences in primary visual and

auditory areas (see SUPPL Fig. 12).

Alpha power was positively correlated with amplitude of frequency tagged responses

The positive correlations between alpha power and frequency-tagging amplitude observed in

study 1 could be replicated. Alpha activity 500 ms before target onset correlated both with 36

Hz VSSEF and 40 Hz ASSEF activity when expecting a visual target (see Fig. 10A-B, 36 Hz

response: cluster significance: $p < .01$; median split: $p < .001$; $t_{(24)} = 4.33$; $BF_{(10)} = 127$; t-test:

$p < .001$; $t_{(24)} = 5.33$; $BF_{(10)} = 1272$; 40 Hz response: cluster significance: $p < .001$; median

split: $p < .001$; $t_{(24)} = 7.05$; $BF_{(10)} = 59443$; t-test: $p < .001$; $t_{(24)} = 6.75$; $BF_{(10)} = 30515$).

The same correlation was present when expecting an auditory target for both 36 Hz VSSEF

activity (see Fig. 10. D-E; cluster significance: $p < .001$; median split: $p = .001$; $t_{(23)} = 3.62$;

$BF_{(10)} = 25.51$; t-test: $p < .001$; $t_{(23)} = 4.60$; $BF_{(10)} = 216$) and 40 Hz ASSEF activity (cluster

significance: $p = .005$; median split: $p < .001$; $t_{(25)} = 3.75$; $BF_{(10)} = 36.40$; t-test: $p < .001$;

$t_{(25)} = 4.06$; $BF_{(10)} = 73.61$).

Figure 10. Relationship between cue induced alpha modulation and amplitude of frequency tagged responses in study 2. A-C, a positive correlation is visible between alpha activity in the last 500 ms as well as alpha activity in the last 1500ms–1000 ms and steady state potentials shortly before target onset when expecting a visual target (36 HZ late: p = .013; 40 Hz late: p = .009; 40 Hz early: p = 002). **D-F,** when expecting an auditory target, there is a positive correlation between alpha activity in the last 500 ms as well as alpha activity in the last 1500ms–1000 ms and steady state potentials shortly before target onset (36 HZ late: p < .001; 40 Hz late: p = .005; 36Hz early: p = 011). N = 27; *** sig < .001; ** sig. < .01; * sig. < .05. + sig. < .1

In accordance with the results of study 1, we then tested whether alpha activity preceding frequency-tagging activity showed similar correlations. Our data revealed that during the last 1 to 1.5 s before target onset alpha activity correlated with the 36 Hz VSSEF response during the last 500 ms prior to an auditory target (see Fig. 10F; cluster significance: p = .01; median split: p < .001; $t_{(24)}$ = 4.66; BF$_{(10)}$ = 271; t-test: p < .001; $t_{(24)}$ = 5.66; BF$_{(10)}$ = 2688). The same alpha activity correlated with 40 Hz ASSEF activity during the last 500 ms prior to a visual target (see Fig. 10C; cluster significance: p = .002; median split: p = .002; $t_{(23)}$ = 3.41; BF$_{(10)}$ = 16.38; t-test: p < .001; $t_{(23)}$ = 5.57; BF$_{(10)}$ = 1901).

Lastly, both alpha activity as well as 36 Hz VSSEF responses 500 ms before target onset

correlated negatively with reaction time on a trial-by-trial basis, indicating faster reaction

times in trials with higher pre-stimulus activity (alpha: $p = .037$; median split: $p = .013$;

$t_{(25)} = -2.67$; $BF_{(10)} = 3.78$; t-test: $p < .01$; $t_{(25)} = -3.34$; $BF_{(10)} = 14.84$; 36 Hz: $p = .002$; median

split: $p = .004$; $t_{(25)} = -3.20$; $BF_{(10)} = 8.98$; t-test: $p < .01$; $t_{(25)} = -3.46$; $BF_{(10)} = 19.12$. See suppl

Fig. 3-4).

Figure 11. Steady-state response in the intermodulation frequency A, similar to study 1, 4 Hz frequency-tagging activity was increased shortly before target onset when expecting an auditory compared to a visual target ($p = .006$). **D**, source localization showed activity over auditory sensory areas, but did not reach significance. N = 27;

467   The same increase in the intermodulatory frequency observed in study 1, could be observed in

468   our second study, during the last 500 ms prior to target onset (see Fig. 11A; $p = .006$). In

469   source space, a descriptive condition difference was visible in auditory and visual sensory

470   cortices, however this effect did not reach significance (p = .49; peak spm coordinates: 45 -83

471   -19 mm, in the right lingual gyrus; see Fig. 11B).

Discussion

The neuropsychological account of attention defines it as the selective facilitation (i.e.,

prioritization) of relevant sensory input and suppression of irrelevant sensory input.

Oscillatory activity in the alpha range (~10 Hz) has been suggested to play a mechanistic role

in attention through inhibition of irrelevant cortices, commonly referred to as the 'alpha

inhibition hypothesis' (Foxe et al., 1998; Jensen & Mazaheri, 2010; Klimesch et al., 2007). In

the current cross-modal attention study we directly tested this hypothesis by using frequency-

tagging to specifically examine how cues signalling the modality of an upcoming target

(either the auditory or visual modality) affected the responsiveness of the relevant and

irrelevant sensory cortices prior to target onset. In-line with previous work, we observed a

post-cue increase in posterior alpha power in anticipation of processing auditory targets.

However, contrary to prevalent theories proposing visual gain suppression when focusing on

the auditory modality, we observed that the amplitude of visual frequency-tagging responses

increased just prior to the onset of the auditory target. This suggests that responsiveness of the

visual stream was not inhibited when attention was directed to auditory processing and was

not inhibited by occipital alpha activity, which directly contradicts the alpha inhibition

hypothesis. Our results reconcile previously paradoxical results on audio-visual attention and

support the view that alpha activity gates downstream communication pathways.

Frequency-tagging

In the current experiment, we specifically chose to analyse the cue-to-target interval, where

both visual and auditory SSEPs/SSEFs present preparatory states for the upcoming task,

independent of task-related processing or performance. The magnitude of auditory

SSEPs/SSEFs was increased shortly before target onset when expecting an auditory target

compared to a visual target, very much in line with previous reports (e.g., Saupe et al., 2009).

In contrast to the results reported in Saupe et a., (2009), where visual SSEPs decreased when

attending the auditory modality, visual SSEPs/SSEFs increased shortly before target onset

when expecting an *auditory* target in our data. This is especially surprising as auditory targets

were frequently, or in case of our second study, always accompanied by visual distractors,

rendering it optimal for task success to completely ignore any visual input. As auditory targets

were significantly more difficult than visual targets in our first study and of comparable

difficulty in our second study, these results strongly speak to a vigilance increase of sensory

processing independent of modality and an inability to selectively disengage one sensory

modality in anticipation of a demanding task. This view is consistent with previous work in

which visual SSEPs elicited by irrelevant background stimulation increased with task load in

an auditory discrimination task (Jacoby et al., 2012). Furthermore, our results indicate that

task demand is a strong candidate to reconcile previously seemingly paradox results, as

splitting attention between the auditory and visual system seemed possible in simpler tasks

(Driver & Spence, 1998; Saupe et al., 2009) and impossible under high demand (de Jong et

al., 2010; Driver, 1996; Driver & Spence, 1998; Spence & Driver, 1996). An alternative

account for our findings stems from the evidence, that participants are more likely to only

perceive and react to the visual modality, when confronted with audio-visual stimuli

(Colavita, 1974; Spence, 2009). However, this effect was mostly limited to speeded modality

discrimination/target detection tasks (Sinnett et al., 2008; Spence, 2009). Furthermore, the

increased difficulty of the here-used auditory stimuli was confirmed in a previous block-

design study (van Diepen & Mazaheri, 2017) and in our second study, performances over the

visual and auditory tasks were comparable. Nevertheless, visual dominance could play a role

for auditory target difficulty as well as predictions over the reciprocity of the audio-visual

relationship.

A revision of the alpha inhibition hypothesis

Top-down cued changes in alpha power have now been widely viewed to play a functional role in directing attention: the processing of irrelevant information is attenuated by increasing alpha power in cortices involved with processing this information (Foxe, Simpson, & Ahlfors, 1998; Hanslmayr et al., 2007; Jensen & Mazaheri, 2010). However, recent evidence suggests that posterior alpha activity does not inhibit gain in early sensory processing stages (Antonov et al., 2020; Gundlach et al., 2020; Gutteling et al., 2022; Zhigalov & Jensen, 2020). To date there has been no direct investigation into the effect of alpha increases on later stages in the processing stream. In the current study, as expected, we observed a post-cue increase in occipital alpha activity in anticipation of an auditory target. However, we also observed an increase in the amplitude of visual SSEPs during the cue-target interval, which directly contradicts the widespread view of alpha activity exerting 'gain control' in early sensory areas by regulating excitability (Foxe & Snyder, 2011; Jensen & Mazaheri, 2010; Van Diepen et al., 2019). Here we propose that alpha activity, rather than modulating early primary sensory processing, exhibits its inhibitory effects at later stages of the processing stream (Antonov et al., 2020; Gundlach et al., 2020; Zhigalov & Jensen, 2020; Zumer et al., 2014), gating feedforward or feedback communication between sensory areas (Bauer et al., 2020; Haegens et al., 2015; Uemura et al., 2021). Our data provides evidence in favour of this view, as we can show that early sensory alpha activity covaries over trials with SSEP magnitude in higher order sensory areas. If alpha activity exerted gain control in early visual regions, increased alpha activity would have to lead to a decrease in SSEP responses. In contrast, we observe that increased alpha activity originating from early visual cortex is related to *enhanced* visual processing. Source localization confirmed that this enhancement was not originating from early visual areas, but from areas associated with later stages of the processing stream such as the precuneus, which has been connected to sensory integration (Al-Ramadhani et al., 2021; Xie et al., 2019). While we cannot completely rule out alternative explanations, it seems plausible to assume that inhibition of other task-irrelevant communication pathways leads to

prioritised and thereby enhanced processing over relevant pathways. In line with previous

literature (Morrow et al., 2023; Peylo et al., 2021; Zhigalov & Jensen, 2020), we therefore

suggest that alpha activity limits task-irrelevant feedforward communication, thereby

enhancing processing capabilities in relevant downstream areas (see Fig. 1A). The benefit of

alpha-modulation and its effect on visual information processing is underlined by reaction

times, which were faster both for trials with high pre-target alpha activity and high pre-target

visual SSVEF activity.

It should be noted, the comparison between modulation in alpha activity and in SSEP/SSEFs

is difficult, especially concerning timing. This is largely owed to differences in signal-to-noise

due to trial averaging in the frequency versus the time domain and temporal and frequency lag

in the estimation of alpha activity (Peylo et al., 2021). It is further noteworthy, that the

majority of evidence for the alpha inhibition hypothesis focused on the effect of pre-target

alpha modulation on behaviour and target-related potentials (Morrow et al., 2023). However,

in our data alpha modulation occurs clearly ahead of SSVEP/SSVEF modulation on a scale

that could not be simply explained by temporal or frequency smearing. Additionally,

significant trial-by-trial correlations, which occur in the frequency domain for both signal

types, underline the strong relationship between both measurements.

Interestingly, we could show that the magnitude of the correlation between alpha power and

visual information processing varied between conditions, suggesting a dynamic and adaptive

regime. This notion supports the view that alpha oscillations represent a mechanism rather

than a specific function, which can fulfil different roles depending on task demand and

network location, which has been confirmed in a recent study revealing functionally distinct

alpha networks (Clausner et al., 2024). As such, it is conceivable that alpha oscillations can in

some cases inhibit local processing, while in other cases, depending on network location,

connectivity and demand, alpha oscillation can facilitate signal transmission. In different

contexts, utilizing unimodal targets and distractors, spatial cueing, or covert attention,

different functional processes could be involved (Morrow et al., 2023). Future research should

intensify efforts to disentangle these effects, investigating localized alpha networks

intracranially or through combinations of fMRI, EEG and MEG, to clearly measure their

effects on sensory processing and behaviour.

It is known that the localisation of alpha activity reflects the retinotopic organisation of visual

spatial attention topographically over the parietooccipital cortex (Kelly et al., 2006; Popov et

al., 2019). Notably, recent studies provided evidence, that the same organisation can be

observed for auditory attention. Specifically, the localisation of visual alpha activity in the

parietooccipital cortex reflects the spatial direction of auditory attention (Klatt et al., 2021;

Popov et al., 2021). This observation can be explained through micro-saccades towards the

spatial location of sounds, which are irrevocably connected to alpha oscillations (Popov et al.,

2021). While we did not manipulate spatial attention, our results fit well to the notion of

visual alpha activity serving as a sensory orientation system, relaying visual information to

task-relevant downstream processing areas, and blocking communication to irrelevant

pathways.

The intermodulation frequency

Previous research showed that simultaneous frequency-tagging in multiple frequencies evokes

a response in the intermodulation frequency (f1 – f2). In multimodal settings, this frequency is

thought to reflect cross-modal integration (Drijvers et al., 2021). This is very well in line with

our findings, where increased vigilance of the sensory system arising from anticipation of a

difficult auditory target resulted in an increase in the intermodulation frequency. Likewise,

our data shows that visual signal enhancement was localized in precuneus, which has been

connected to sensory integration (Al-Ramadhani et al., 2021; Xie et al., 2019). Furthermore,

we could show that the intermodulation frequency covaries over trials with reaction time in

the easiest condition, where visual targets were presented without any distractors. A lack of

this connection in other conditions might reflect increasing interferences from higher task

difficulty, rather than a lack of the effect itself, but this remains to be tested. We cannot

exclude an alternative explanation, as theta oscillations are known to be involved in

movement preparation, it is possible that phase-resets could lead to time-locked appearance of

these oscillations (Lakatos et al., 2008; Tomassini et al., 2017).

Conclusion

Our results taken together suggest that under high attention, audio-visual excitability is

enhanced, reflecting an increase in vigilance for both visual as well as auditory information,

even if this increases processing of distracting information. We showed that this vigilance

shift, as reflected by SSEP/SSEF responses, is regulated by early visual alpha activity,

presumably through relaying of visual information over communication pathways, thereby

controlling the downstream flow of visual information.

Materials and Methods

Participants EEG-Study

In total, 24 healthy volunteers participated in this study (mean age: 19.1 ± 1.8 SD; 17

women). Due to technical difficulties, one participant could not finish the experiment and one

participant did not exceed chance level in the behavioural task (~33 %). Both were therefore

removed from any further analysis. All remaining participants reported normal or corrected-

to-normal vision, no history of psychiatric or neurological illness and provided written

informed consent. After completion of the experiment, participants received either monetary

compensation or certification of their participation for their university course program. The

study protocol was approved by the Ethics Committee of the School of Psychology at the

University of Birmingham and is in accordance with the Declaration of Helsinki.

Participants MEG-Study

In total, 28 healthy volunteers participated in this study (mean age: 23.4 ± 3.6 SD; 20

women). One participant was removed from further analysis, as they only responded to ~42%

of trials correctly in the second block, which related to 27/19 trials per condition respectively.

All remaining participants reported normal or corrected-to-normal vision, no history of

psychiatric or neurological illness and provided written informed consent. After completion of

the experiment, participants received either monetary compensation or certification of their

participation for their university course program. The study protocol was approved by the

Ethics Committee of the School of Psychology at the University of Birmingham and is in

accordance with the Declaration of Helsinki.

Cross-modal attention paradigm EEG-Study

Each trial was initiated by a brief presentation of a cue (100 ms) signalling the modality of the

upcoming discrimination task (v-shape: visual modality; inversed v-shape: auditory modality;

diamond-shape: non-specific). During the following cue-to-target interval, the fixation cross

was frequency-tagged at 36 Hz. At the same time, a 40 Hz frequency-tagged sound

(amplitude modulated white noise) was played over headphones (see Fig. 1). The volume of

tones was initially adjusted to a level that was clearly perceivable but not uncomfortable and

remained stable over participants. No task was connected to this interval and participants did

not need to pay attention to either the sound or the fixation cross. After three seconds, the

frequency-tagging stopped, and right after the cessation of stimuli, the target was presented

for a very brief moment (25 ms). It consisted either of a Gabor patch (visual modality) or a

sound (auditory modality). If the target modality was visual, participants had to use the three

arrow buttons on the keyboard to indicate whether the Gabor patch was tilted to the left (-10°;

left arrow button), vertical (0°; down arrow button), or tilted to the right (10°; right arrow

button). Additionally, in 50% of the trials, a random distractor from the pool of auditory

targets was presented simultaneously to the visual target over headphones. Similarly, if the

target modality was auditory, participants used the same buttons to indicate whether the pitch

of the tone was low (500 Hz), medium (1000 Hz) or high (2000 Hz). Again, in 50% of the

trials, a random distractor from the pool of visual targets was also presented. If the cue was

non-specific (diamond shape), either a visual (50% of non-specific trials) or an auditory target

was presented, never accompanied by any distractors. Experimental trials were separated by

an inter-trial interval of 4 seconds, to avoid carry-over effects from previous trials. The

resulting 6 conditions were randomly ordered and balanced out over the experiment.

During the experiment, participants were instructed to keep their gaze locked to a fixation

cross presented at the centre of the screen. Preceding data collection, participants performed

36 practice trials to get accustomed to the task and the target stimuli. The ensuing experiment

was split into 26 trial sequences, separated by self-chosen breaks, which together resulted in

468 trials and lasted between 80 and 90 minutes. The discrimination task was programmed

and presented with MATLAB® R2020b and Psychtoolbox-3 on an LCD-monitor featuring a

140 Hz refresh rate. The onset of the visual and auditory tagging frequencies (i.e steady state

stimuli)   were   tracked   using   the   Cedrus   Stimtracker

(https://cedrus.com/stimtracker/index.htm).

Adjustments to the attention paradigm in the MEG-Study

In our second study, we removed all ambiguity concerning targets and distractors and

therefore developed a blocked design, incorporating two blocks. The first block did not

display distractors and only cues which correctly predicted the target were presented. Cues in

the second block were likewise always correctly indicating the target modality, but this time,

each target was accompanied by a random distractor from the non-target modality.

Furthermore, the visual task was adjusted to be more in line with the auditory task. As such,
the Gabor patches now featured stripes in different frequencies (e.g. a low number of stripes,
a medium number of stripes and a high number of stripes. The participant's task was to
discriminate between these three Gabor patches. As auditory targets had been markedly more
difficult in our first study, we now included a brief difficulty calibration prior to the
experiment. First, we presented 21 Gabor patches with 3 different amounts of stripes
following a standard difficulty. If participants could discriminate them correctly 55 – 75% of
the time, this difficulty setting was chosen. Otherwise, depending on the performance, the
stripe-frequency of the Gabor patches was adjusted. There were maximally 3 sessions of 21
Gabor patches, after which we had enough data to calibrate the individual difficulty setting.

The same procedure was then performed with the difficulty of the tones, calibrating the pitch
frequency for each individual participant.

Lastly, visual frequency-tagging stimulation now followed a sinusoidal contrast-change rather
than an on-off stimulation, which was possible due to a high-resolution projector featuring a
refresh rate of 1440 Hz (PROPixx DLP LED projector ;VPixx Technologies Inc., Canada).

Eye-tracking

To confirm that participants had focused on the fixation cross during the cue-to-target
interval, we incorporated eye-tracking into our MEG-experiment (EyeLink 1000 Plus).
Correct trials of the second block were analysed for vertical and horizontal eye-movements.
To exclude blinks from this analysis, trials with very large eye-movements (> 10 degrees of
visual angle) were removed from the eye-tracking data (See suppl Fig. 5).

Behavioral analysis

We were interested in accuracy in the discrimination of visual targets and auditory targets, as
well as reaction times. Furthermore, we examined the distraction cost of having a target
presented with a distractor of a different modality as well as the reaction time to make the
target discrimination. The distraction cost was calculated as the reaction time difference
between cued targets with distractors (i.e. visual and auditory stimuli presented together) and
cued targets without distractors (either a visual or auditory stimulus presented alone). All
incorrect trials as well as trials with reaction times faster than 100 ms or exceeding 1500 ms
were removed from analysis (0.5 % too fast, 9.2% too slow).

 EEG data acquisition

 All EEG recordings were conducted using a WaveGuard Cap (ANTneuro), featuring 64

 Ag/AgCL electrodes (10-10 system; ground: Fz; reference: Cpz; EOG: left canthus).

 Electrodes positions were prepared with OneStep cleargel conductive paste and impedances

 were kept below 100 kΩ. The measured signal was transmitted using an ANTneuro

 EEGosports amplifier (low-pass filter: 150 Hz; high-pass filter: 0.5 Hz; sampling rate: 500

 Hz).

 MEG data acquisition

 Prior to the experiment, feducial positions and head-shape were recorded using a FASTRAK

 system (Polhemus, USA). The experiment took place in a dimly lit room, where participants

 were seated in a comfortable chair in the gantry of a 306-sensor TRIUX Elekta system with

 204 orthogonal planar gradiometers and 102 magnetometers (Elekta, Finland). The 71*40 cm

 screen was positioned at ~1.40 m distance from the participant.

 EEG Preprocessing

 Offline analyses were performed in MATLAB ® R2020b. The data was pruned from artifacts

 by visual inspection using the EEGLAB toolbox (Delorme & Makeig, 2004). Additionally,

 blinks and ocular artefacts were removed from the data using independent component analysis

 (ICA). EEG channels were re-referenced to an average of all channels (excluding EOG).

 MEG Preprocessing

 Offline analyses were performed in MATLAB ® R2020b and Python. Spatiotemporal Signal-

 Source-Separation (SSS) was applied to the raw data via MNE's inbuilt maxfilter function

 with a duration window of 10 s and a correlation value of .9. The data was pruned from

 artifacts by visual inspection using the Fieldtrip toolbox (Delorme & Makeig, 2004).

 Additionally, blinks and ocular artefacts were removed from the data using independent

 component analysis (ICA). In sensor space, planar gradiometers were combined for further

 analyses. In source space, all individual planar gradiometers were analysed.

 Amplitude of the evoked frequency-tagging response

 To investigate the temporal dynamics of amplitude of the frequency tagged responses after

 the onset of the attentional cues (also the precise onset of the frequency tagged stimuli) the

data was epoched into 6-second segments starting 1.5 seconds prior to cue onset. Next the

data were narrow-band filtered around the 36 Hz activity to capture the visual frequency-

tagging, 40 Hz activity in to capture the auditory frequency-tagging, and the intermodulation

frequency at 4 Hz, which can be derived by subtracting both frequency-tagging responses ($f_i =$

$f_{auditory} - f_{visual}$; see Drijvers, Spaak & Jensen, 2020). Here we used a Blackmann-windowed

sync filters adapted to a suitable ratio of temporal and frequency resolution for the specific

frequency of each of the tagged signals: filter order 116 for 35.5 to 36.5 Hz and 39.5 to 40.5,

filter order 344 for 3.5 to 4.5 Hz. The filtered data at each of the tagged frequencies as well

the intermodulated frequency were baseline corrected (interval between 700 and 200 ms

preceding cue onset) before calculating the average over trials to obtain steady-state evoked

potentials). The power envelope of the SSEPs of tagged frequencies was estimated using

Hilbert transformation. To confirm that 4 Hz is a sufficient distance between tagging

frequencies, we repeated to analysis for 43.5 to 44.5. We found no indication of frequency-

bleeding over, as the effects observed at 40 Hz, were not present at 44 Hz (see SUPPL Fig.

11).

Temporal dynamics of the induced EEG changes

In addition to looking at the cue evoked changes in the amplitude of the frequency tagged

signals, we investigated the induced changes in the EEG signal at the frequencies of the

tagged auditory and visual stimuli, alpha activity (9-11 Hz, filter order 276 for 7.5 to 12.5

Hz), as well as the intermodulation frequency (4 Hz; filter order 344 for 3.5 to 4.5 Hz). Here

rather than averaging the epoched data filtered at the specific frequency ranges, we performed

the Hilbert transform, and averaged the power-envelope of the specific frequencies across

trials. This approach is very much analogous to the standard time-frequency analysis using

convolutions (van Diepen & Mazaheri, 2017; Zhigalov et al., 2019), but affords more control

concerning temporal versus frequency resolution to examine the temporal dynamics of the

specific frequencies of interest. In our second study, the individual peak alpha frequency was

used for bandpass-filtering in contrast to a standardised band applied in the first study.

Time-frequency representations of power

In addition to the estimating frequency power envelopes, Time–frequency representations

(TFRs) of power of the EEG signal were estimated using the Fieldtrip toolbox (Oostenveld et

al., 2011). The power or frequencies between 5 and 20 Hz were calculated for each trial, using

a sliding time window (frequency steps: 0.5; time steps: 10 ms). The length of the window

was adjusted to a length of 3 cycles per frequency and tapered with a Hanning window. For

each trial, both datasets were normalised to display relative percent change from baseline

using the following formula: [(activity – baseline) / baseline], where baseline refers to the

interval between 700 and 200 ms before cue onset. To estimate the topographical distribution

of voltage differences between conditions, uncorrected power values were normalised

applying the following formula: $[\Delta/\sum = (a - b) / (a + b)]$, where a and b reflect the different

conditions.

*Source localization*Source localization was performed with a beamformer approach using the

Fieldtrip toolbox. Headmodels were created based on individual T1-scans fitted to fiducial

points and head shapes. These data were fit to a 5mm 3d sourcemodel and warped into MNI-

space. Two participants were missing individual T1-scans. In these cases, we applied a

standardized T1-scan using the Colin 27 Average Brain Model (Holmes et al., 1998).

Frequency-domain data was localized using the Dynamic Imaging of Coherent Sources (DICS

beamformer) method with a dpss taper of 2 Hz assuming fixed orientation. As condition

differences between frequency-tagging responses were better estimated in the time-domain,

we assessed them applying the Synthetic Aperture Magnetometry (SAM-beamformer) method

with optimal fixed rotation (Sekihara et al., 2004). Significant brain areas and peak

coordinates were related to brain areas using the Anatomical Automatic Labeling (AAL) atlas

for SPM8 (Tzourio-Mazoyer et al., 2002) . After statistical analysis, source localized data was

interpolated onto the Colin 27 Average Brain Model MRI. Cerebellar and brainstem

interpolations were excluded from the coordinate system.

Statistical analysis

Condition differences in the behavioural task were estimated with paired t-tests and repeated

measures ANOVAS, utilizing Tukey-Kramer Post-Hoc test. Power differences between

conditions, as well as source-space contrasts were analysed using cluster permutation analysis

(Maris & Oostenveld, 2007). In this procedure, condition labels were randomly shuffled 1000

times, creating pairs of surrogate conditions. To test for significance, paired t-tests were

conducted for each data point and each channel, resulting in one t- matrix for real conditions

and 1000 t-matrices for surrogate conditions. Significant t-values ($p < .05$) were defined as

clusters if there was at least one significant data point present at the same time and frequency

in at least two neighbouring channels. To correct for multiple comparisons, a condition

difference was only assumed, if the maximum sum of t-values in a real cluster exceeded the
same sum of 95% of the clusters found in the surrogate data. To replicate our results in the
second study, we applied the same statistics and averaged the previously found time-intervals
into windows of 500ms (e.g.: if we found an effect -0.51 to -0.0620 s prior to target onset we
tested this effect for a time window of -0.5 to 0 s prior to target onset).

The relationship between induced changes in alpha activity and frequency tagged responses
was assessed using trial by trial Spearman correlations. For each participant and each
electrode, a correlation coefficient was calculated between the average activity in a previously
identified cluster, which was used as seed (e.g., condition differences in alpha activity), and
the average activity of the electrophysiological correlate of interest (e.g., 36 Hz activity over
the previously identified time window). The correlation coefficients were z-transformed, and
the resulting channel by participant matrix was tested against null-correlation model using the
cluster permutation approach described above. Derived clusters were additionally tested and
visualised by comparing median split trials of high vs low activity. For this analysis, outliers
(values deviating more than 2 standard deviations from the mean) were excluded.
Furthermore, the average correlation coefficient of the cluster was tested against a 0-
correlation model for each participant using t-statistics. Lastly, an interaction between
electrophysiological correlations and conditions was performed using correlation coefficients
for each participant and electrode, testing them between conditions using the cluster
permutation approach. Perceptually uniform and universally readable colormaps were applied
to all visualisations (Crameri et al., 2020). All data are presented as mean ± standard error of
the mean (SEM).

Acknowledgements: This work was made possible by funding support from Facebook

Oculus and BBSRC (BB/R018723/1).

Competing interests: There are no competing interests

Data availability: Codes for analyses and figures will be made available on github and the
data will be uploaded to Dryad or OSF

References

Al-Ramadhani, R. R., Shivamurthy, V. K. N., Elkins, K., Gedela, S., Pedersen, N. P., & Kheder, A. (2021). The Precuneal Cortex: Anatomy and Seizure Semiology. *Epileptic Disorders : International Epilepsy Journal with Videotape*, *23*(2), 218. https://doi.org/10.1684/EPD.2021.1257

Antonov, P. A., Chakravarthi, R., & Andersen, S. K. (2020). Too little, too late, and in the wrong place: Alpha band activity does not reflect an active mechanism of selective attention. *NeuroImage*, *219*, 117006. https://doi.org/10.1016/J.NEUROIMAGE.2020.117006

Bauer, A. K. R., Debener, S., & Nobre, A. C. (2020). Synchronisation of Neural Oscillations and Cross-modal Influences. In *Trends in Cognitive Sciences* (Vol. 24, Issue 6, pp. 481–495). Elsevier Ltd. https://doi.org/10.1016/j.tics.2020.03.003

Brickwedde, M., Bezsudnova, Y., Kowalczyk, A., Jensen, O., & Zhigalov, A. (2022). Application of rapid invisible frequency tagging for brain computer interfaces. *Journal of Neuroscience Methods*, *382*, 109726. https://doi.org/10.1016/J.JNEUMETH.2022.109726

Brickwedde, M., Schmidt, M. D., Krüger, M. C., & Dinse, H. R. (2020). 20 Hz Steady-State Response in Somatosensory Cortex During Induction of Tactile Perceptual Learning Through LTP-Like Sensory Stimulation. *Frontiers in Human Neuroscience*. https://doi.org/10.3389/fnhum.2020.00257

Clausner, T., Marques, J., Scheeringa, R., & Bonnefond, M. (2024). Feature specific neuronal oscillations in cortical layers. *BioRxiv*, 2024.07.31.605816. https://doi.org/10.1101/2024.07.31.605816

Colavita, F. B. (1974). Human sensory dominance*. In *Perception & Psychophysics* (Vol. 16, Issue 2).

Colon, E., Legrain, V., & Mouraux, A. (2012). Steady-state evoked potentials to study the processing of tactile and nociceptive somatosensory input in the human brain. *Neurophysiologie Clinique/Clinical Neurophysiology*, *42*(5), 315–323. https://doi.org/10.1016/J.NEUCLI.2012.05.005

Crameri, F., Shephard, G. E., & Heron, P. J. (2020). The misuse of colour in science communication. *Nature Communications*, *11*(1). https://doi.org/10.1038/s41467-020-19160-7

de Jong, R., Toffanin, P., & Harbers, M. (2010). Dynamic crossmodal links revealed by steady-state responses in auditory–visual divided attention. *International Journal of Psychophysiology*, *75*(1), 3–15. https://doi.org/10.1016/J.IJPSYCHO.2009.09.013

Dinse, H. R., Ragert, P., Pleger, B., Schwenkreis, P., & Tegenthoff, M. (1998). Pharmacological Modulation of Perceptual Learning and Associated Cortical Reorganization. *SCIENCE*, *282*, 865.

Drijvers, L., Jensen, O., & Spaak, E. (2021). Rapid invisible frequency tagging reveals nonlinear integration of auditory and visual information. *Human Brain Mapping*, *42*(4), 1138–1152. https://doi.org/10.1002/hbm.25282

Driver, J. (1996). Enhancement of selective listening by illusory mislocation of speech sounds due to lip-reading. *NATURE ·*, *381*.

Driver, J., & Spence, C. (1998). Cross-modal links in spatial attention. *Philos Trans R Soc Lond B Biol Sci*, *353*, 1319–1331.

Dugué, L., Marque, P., & VanRullen, R. (2011). The phase of ongoing oscillations mediates the causal relation between brain excitation and visual perception. *Journal of Neuroscience*, *31*(33), 11889–11893. https://doi.org/10.1523/JNEUROSCI.1161-11.2011

Ergenoglu, T., Demiralp, T., Bayraktaroglu, Z., Ergen, M., Beydagi, H., & Uresin, Y. (2004). Alpha rhythm of the EEG modulates visual detection performance in humans. *Cognitive Brain Research*, *20*(3), 376–383. https://doi.org/10.1016/J.COGBRAINRES.2004.03.009

Foxe, J. J., Simpson, G. V, & Ahlfors, S. P. (1998). Parieto-occipital ~10Hz activity reflects anticipatory state of visual attention mechanisms. *NeuroReport*.

Foxe, J. J., & Snyder, A. C. (2011). The role of alpha-band brain oscillations as a sensory suppression mechanism during selective attention. *Frontiers in Psychology*, *2*(JUL). https://doi.org/10.3389/fpsyg.2011.00154

Fu, K. M. G., Foxe, J. J., Murray, M. M., Higgins, B. A., Javitt, D. C., & Schroeder, C. E. (2001). Attention-dependent suppression of distracter visual input can be cross-modally cued as indexed by anticipatory parieto–occipital alpha-band oscillations. *Cognitive Brain Research*, *12*(1), 145–152. https://doi.org/10.1016/S0926-6410(01)00034-9

Gundlach, C., Moratti, S., Forschack, N., & Müller, M. M. (2020). Spatial Attentional Selection Modulates Early Visual Stimulus Processing Independently of Visual Alpha Modulations. *Cerebral Cortex*, *30*(6), 3686–3703. https://doi.org/10.1093/cercor/bhz335

Gutteling, T. P., Sillekens, L., Lavie, N., & Jensen, O. (2022). Alpha oscillations reflect suppression of distractors with increased perceptual load. *Progress in Neurobiology*, *214*. https://doi.org/10.1016/J.PNEUROBIO.2022.102285

Haegens, S., Barczak, A., Musacchia, G., Lipton, M. L., Mehta, A. D., Lakatos, P., & Schroeder, C. E. (2015). Laminar profile and physiology of the α rhythm in primary visual, auditory, and somatosensory regions of neocortex. *Journal of Neuroscience*, *35*(42), 14341–14352. https://doi.org/10.1523/JNEUROSCI.0600-15.2015

Haegens, S., Nácher, V., Luna, R., Romo, R., & Jensen, O. (2011). α-Oscillations in the monkey sensorimotor network influence discrimination performance by rhythmical inhibition of neuronal spiking. *Proceedings of the National Academy of Sciences*, *108*(48), 19377–19382. https://doi.org/10.1073/PNAS.1117190108

Hari, R., Hämäläinen, M., & Joutsiniemi, S. L. (1989). Neuromagnetic steady-state responses to auditory stimuli. *Journal of the Acoustical Society of America*, *86*(3), 1033–1039. https://doi.org/10.1121/1.398093

Holmes, C. J., Hoge, R., Collins, L., Woods, R., Toga, A. W., & Evans, A. C. (1998).
Enhancement of MR images using registration for signal averaging. *Journal of Computer*
Assisted Tomography, *22*(2), 324–333. https://doi.org/10.1097/00004728-199803000-
00032

Jacoby, O., Hall, S. E., & Mattingley, J. B. (2012). A crossmodal crossover: Opposite effects
of visual and auditory perceptual load on steady-state evoked potentials to irrelevant
visual stimuli. *NeuroImage*, *61*(4), 1050–1058.
https://doi.org/10.1016/J.NEUROIMAGE.2012.03.040

Jensen, O., & Mazaheri, A. (2010). Shaping functional architecture by oscillatory alpha
activity: Gating by inhibition. *Frontiers in Human Neuroscience*, *4*.
https://doi.org/10.3389/fnhum.2010.00186

Kass, R. E., & Raftery, A. E. (1995). Bayes factors. *Journal of the American Statistical*
Association, *90*(430), 773–795. https://doi.org/10.1080/01621459.1995.10476572

Kelly, S. P., Lalor, E. C., Reilly, R. B., & Foxe, J. J. (2006). Increases in alpha oscillatory
power reflect an active retinotopic mechanism for distracter suppression during sustained
visuospatial attention. *Journal of Neurophysiology*, *95*(6), 3844–3851.
https://doi.org/10.1152/jn.01234.2005

Klatt, L.-I., Getzmann, S., & Schneider, D. (2021). Attentional Modulations of Alpha Power
Are Sensitive to the Task-relevance of Auditory Spatial Information. *BioRxiv*.
https://doi.org/10.1101/2021.02.12.430942

Klimesch, W., Sauseng, P., & Hanslmayr, S. (2007). EEG alpha oscillations: The inhibition–
timing hypothesis. *Brain Research Reviews*, *53*(1), 63–88.
https://doi.org/10.1016/J.BRAINRESREV.2006.06.003

Lakatos, P., Karmos, G., Mehta, A. D., Ulbert, I., & Schroeder, C. E. (2008). Entrainment of
neuronal oscillations as a mechanism of attentional selection. *Science (New York, N.Y.)*,
320(5872), 110–113. https://doi.org/10.1126/SCIENCE.1154735

Maris, E., & Oostenveld, R. (2007). Nonparametric statistical testing of EEG- and MEG-data.
Journal of Neuroscience Methods, *164*(1), 177–190.
https://doi.org/10.1016/J.JNEUMETH.2007.03.024

Marzoll, A., Saygi, T., & Dinse, H. R. (2018). The effect of LTP- and LTD-like visual
stimulation on modulation of human orientation discrimination. *Scientific Reports*, *8*(1).
https://doi.org/10.1038/s41598-018-34276-z

Mazaheri, A., van Schouwenburg, M. R., Dimitrijevic, A., Denys, D., Cools, R., & Jensen, O.
(2014). Region-specific modulations in oscillatory alpha activity serve to facilitate
processing in the visual and auditory modalities. *NeuroImage*, *87*, 356–362.
https://doi.org/10.1016/J.NEUROIMAGE.2013.10.052

Morrow, A., Elias, M., & Samaha, J. (2023). Evaluating the Evidence for the Functional
Inhibition Account of Alpha-band Oscillations during Preparatory Attention. *Journal of*
Cognitive Neuroscience, *35*(8), 1195–1211. https://doi.org/10.1162/JOCN_A_02009

Müller, M. M., & Hillyard, S. (2000). Concurrent recording of steady-state and transient
event-related potentials as indices of visual-spatial selective attention. *Clinical*
Neurophysiology, *111*(9), 1544–1552. https://doi.org/10.1016/S1388-2457(00)00371-0

Müller, M. M., Picton, T. W., Valdes-Sosa, P., Riera, J., Teder-Sälejärvi, W. A., & Hillyard,
S. A. (1998). Effects of spatial selective attention on the steady-state visual evoked
potential in the 20–28 Hz range. *Cognitive Brain Research*, *6*(4), 249–261.
https://doi.org/10.1016/S0926-6410(97)00036-0

Okazaki, Y. O., De Weerd, P., Haegens, S., & Jensen, O. (2014). Hemispheric lateralization
of posterior alpha reduces distracter interference during face matching. *Brain Research*,
1590(1), 56–64. https://doi.org/10.1016/J.BRAINRES.2014.09.058

Pantev, C., Roberts, L. E., Elbert, T., Roß, B., & Wienbruch, C. (1996). Tonotopic
organization of the sources of human auditory steady-state responses. *Hearing Research*,
101(1–2), 62–74. https://doi.org/10.1016/S0378-5955(96)00133-5

Peylo, C., Hilla, Y., & Sauseng, P. (2021). Cause or consequence? Alpha oscillations in
visuospatial attention. *Trends in Neurosciences*, *44*(9), 705–713.
https://doi.org/10.1016/J.TINS.2021.05.004/ASSET/69AE0787-5757-4CDC-B7D6-
968818664AAB/MAIN.ASSETS/GR2.JPG

Popov, T., Gips, B., Kastner, S., & Jensen, O. (2019). Spatial specificity of alpha oscillations
in the human visual system. *Human Brain Mapping*, *40*(15), 4432–4440.
https://doi.org/10.1002/hbm.24712

Popov, T., Gips, B., Weisz, N., & Jensen, O. (2021). Brain areas associated with visual spatial
attention display topographic organization during auditory spatial attention. *BioRxiv*.
https://doi.org/10.1101/2021.03.15.435371

Porcu, E., Keitel, C., & Müller, M. M. (2013). Concurrent visual and tactile steady-state
evoked potentials index allocation of inter-modal attention: A frequency-tagging study.
Neuroscience Letters, *556*, 113–117. https://doi.org/10.1016/J.NEULET.2013.09.068

Regan, D. (1982). COMPARISON OF TRANSIENT AND STEADY-STATE METHODS*.
Annals of the New York Academy of Sciences.

Saupe, K., Schröger, E., Andersen, S. K., & Müller, M. M. (2009). Neural mechanisms of
intermodal sustained selective attention with concurrently presented auditory and visual
stimuli. *Frontiers in Human Neuroscience*, *3*(NOV).
https://doi.org/10.3389/neuro.09.058.2009

Sekihara, K., Nagarajan, S. S., Poeppel, D., & Marantz, A. (2004). Asymptotic SNR of scalar
and vector minimum-variance beanformers for neuromagnetic source reconstruction.
IEEE Transactions on Biomedical Engineering, *51*(10), 1726–1734.
https://doi.org/10.1109/TBME.2004.827926

Sinnett, S., Soto-Faraco, S., & Spence, C. (2008). The co-occurrence of multisensory
competition and facilitation. *Acta Psychologica*, *128*(1), 153–161.
https://doi.org/10.1016/J.ACTPSY.2007.12.002

Snyder, A. (1992). Steady-state vibration evoked potentials: description of technique and
characterization of responses. *Electroencephalography and Clinical*

Neurophysiology/Evoked Potentials Section, *84*(3), 257–268.
https://doi.org/10.1016/0168-5597(92)90007-X

Spaak, E., Bonnefond, M., Maier, A., Leopold, D. A., & Jensen, O. (2012). Layer-Specific
Entrainment of Gamma-Band Neural Activity by the Alpha Rhythm in Monkey Visual
Cortex. *Current Biology*, *22*(24), 2313–2318.
https://doi.org/10.1016/J.CUB.2012.10.020

Spence, C. (2009). Explaining the Colavita visual dominance effect. In *Progress in Brain*
Research (Vol. 176, pp. 245–258). https://doi.org/10.1016/S0079-6123(09)17615-X

Spence, C., & Driver, J. (1996). Audiovisual Links in Endogenous Covert Spatial Attention.
Journal of Experimental Psychology: Human Perception and Performance, *22*(4), 1005–
1030.

Stapells, D. R., Linden, D., Braxton Suffield, J., Hamel, G., Picton, T. W., & Wp, M. T.
(1984). *Electrophysiologic Techniques in Audiology and Otology Human Auditory*
*Steady State Potentials** (Vol. 5, Issue 2).

Thut, G., Nietzel, A., Brandt, S. A., & Pascual-Leone, A. (2006).
Behavioral/Systems/Cognitive-Band Electroencephalographic Activity over Occipital
Cortex Indexes Visuospatial Attention Bias and Predicts Visual Target Detection.
https://doi.org/10.1523/JNEUROSCI.0875-06.2006

Tobimatsu, S., Zhang, Y. M., & Kato, M. (1999). Steady-state vibration somatosensory
evoked potentials: physiological characteristics and tuning function. *Clinical*
Neurophysiology, *110*(11), 1953–1958. https://doi.org/10.1016/S1388-2457(99)00146-7

Toffanin, P., de Jong, R., Johnson, A., & Martens, S. (2009). Using frequency tagging to
quantify attentional deployment in a visual divided attention task. *International Journal*
of Psychophysiology, *72*(3), 289–298. https://doi.org/10.1016/J.IJPSYCHO.2009.01.006

Tomassini, A., Ambrogioni, L., Medendorp, W. P., & Maris, E. (2017). Theta oscillations
locked to intended actions rhythmically modulate perception. *ELife*, *6*.
https://doi.org/10.7554/eLife.25618.001

Tzourio-Mazoyer, N., Landeau, B., Papathanassiou, D., Crivello, F., Etard, O., Delcroix, N.,
Mazoyer, B., & Joliot, M. (2002). Automated anatomical labeling of activations in SPM
using a macroscopic anatomical parcellation of the MNI MRI single-subject brain.
NeuroImage, *15*(1), 273–289. https://doi.org/10.1006/nimg.2001.0978

Uemura, J.-I., Hoshino, A., Go Igarashi, |, Matsui, Y., Chishima, M., Hoshiyama, | Minoru, &
Mazaheri, A. (2021). Pre-stimulus alpha oscillation and post-stimulus cortical activity
differ in localization between consciously perceived and missed near-threshold
somatosensory stimuli. *Eur J Neurosci*. https://doi.org/10.1111/ejn.15388

Van Diepen, R. M., Cohen, M. X., Denys, D., & Mazaheri, A. (2015). Attention and temporal
expectations modulate power, not phase, of ongoing alpha oscillations. *Journal of*
Cognitive Neuroscience, *27*(8), 1573–1586. https://doi.org/10.1162/jocn_a_00803

Van Diepen, R. M., Foxe, J. J., & Mazaheri, A. (2019). The functional role of alpha-band
activity in attentional processing: the current zeitgeist and future outlook. *Current*
Opinion in Psychology, *29*, 229–238. https://doi.org/10.1016/J.COPSYC.2019.03.015

van Diepen, R. M., & Mazaheri, A. (2017). Cross-sensory modulation of alpha oscillatory
activity: suppression, idling, and default resource allocation. *European Journal of*
Neuroscience, *45*(11), 1431–1438. https://doi.org/10.1111/ejn.13570

Van Dijk, H., Schoffelen, J. M., Oostenveld, R., & Jensen, O. (2008). Prestimulus oscillatory
activity in the alpha band predicts visual discrimination ability. *Journal of Neuroscience*,
28(8), 1816–1823. https://doi.org/10.1523/JNEUROSCI.1853-07.2008

Worden, M. S., Foxe, J. J., Wang, N., & Simpson, G. V. (2000). *Anticipatory Biasing of*
Visuospatial Attention Indexed by Retinotopically Specific-Band
Electroencephalography Increases over Occipital Cortex.
http://www.jneurosci.org/cgi/content/full/4016

Xie, Y. J., Li, Y. Y., Xie, B., Xu, Y. Y., & Peng, L. (2019). The neural basis of complex
audiovisual objects maintenances in working memory. *Neuropsychologia*, *133*, 107189.
https://doi.org/10.1016/J.NEUROPSYCHOLOGIA.2019.107189

Yang, X., Fiebelkorn, I. C., Jensen, O., Knight, R. T., & Kastner, S. (2023). Differential
neural mechanisms underlie cortical gating of visual spatial attention mediated by alpha-
band oscillations. *BioRxiv*, 2023.08.21.553303.
https://doi.org/10.1101/2023.08.21.553303

Zhigalov, A., Herring, J. D., Herpers, J., Bergmann, T. O., & Jensen, O. (2019). Probing
cortical excitability using rapid frequency tagging. *NeuroImage*, *195*, 59–66.
https://doi.org/10.1016/J.NEUROIMAGE.2019.03.056

Zhigalov, A., & Jensen, O. (2020). Alpha oscillations do not implement gain control in early
visual cortex but rather gating in parieto-occipital regions. *Human Brain Mapping*,
41(18), 5176–5186. https://doi.org/10.1002/HBM.25183

Zumer, J. M., Scheeringa, R., Schoffelen, J. M., Norris, D. G., & Jensen, O. (2014). Occipital
Alpha Activity during Stimulus Processing Gates the Information Flow to Object-
Selective Cortex. *PLoS Biology*, *12*(10). https://doi.org/10.1371/journal.pbio.1001965

1053

1054

1056

SUPPL Fig. 1 | Distractor cost and attentional benefit. A-B, illustration of distractor cost: mean performance
over trials with distractors was subtracted from mean performance over trials without distractors. Distractor
effects were observable for accuracy as well as reaction time; **A,** accuracy: auditory- − auditory +: M = 10.0 %;
SD = 7.3; p < .001; $t_{(21)}$ = 7.32; visual- − visual+: M = 1.5%; SD = 3.06; p = .02). The effect was stronger for
auditory than for visual target trials (p < .001; $t_{(21)}$ = 7.67). Reaction time: (auditory- - auditory+: M = -108.1 ms;
SD = 84.8; p < .001; $t_{(21)}$ = -5.98; visual- - visual+: M = 123.6 ms; SD = 76.3; p < .001; $t_{(21)}$ = 7.60). auditory
distracters decreased response time to visual targets (p < .001; $t_{(21)}$ = -11.99). **B,** (accuracy: auditory- − auditory
+: M = 7.2 %; SD = 7.5; p = .001; t(25) = 4.9; visual- − visual+: M = -7.6%; SD = 10.80; p < .01; t(25) = -3.59;
Reaction time: auditory- − auditory +: M = -20.64 ms; SD = 57.6; n.s.: p = .08; t(25) = -1.83; visual- − visual+: M =
60.1 ms ; SD = 58.52; p < .001; t(25) = 5.23). **C,** Illustration of attentional benefit: mean performance over non-
specific trials was subtracted from mean performance over modality-cued trials without distractor. attentional
benefit auditory: non-specifically cued auditory targets - informatively cued auditory targets = M = 81.2 ms; SD
= 54.9; p < .001; t(21) = 6.94; attentional benefit visual: non-specifically cued visual targets - informatively cued
visual targets -; M = 54.4 ms; SD = 41.1 ; p <.001; t(21) = 5.19). The magnitude of the effect on reaction time
also differed between conditions (p = .043; t(21) = 2.16), with stronger attentional benefit for auditory target
cues. Attentional cues did not affect response accuracy, neither in auditory nor visual target conditions
(auditory: p = 0.49; visual: p = 0.32). EEG Study: N = 22; MEG-Study: N = 27; *** sig < .001; ** sig. < .01; * sig. <
.05;

1075

1076

1077

SUPPL Fig. 2 | Time course of alpha activity and frequency-tagging responses for the non-specific compared
to the visually-cued condition. **A,** alpha activity compared between expecting a visual target and having
received a non-specific cue. **B,** 36 Hz frequency-tagging response between expecting a visual target and having
received a non-specific cue. **C,** 40 Hz frequency-tagging response between expecting a visual target and having
received a non-specific cue. **D,** alpha activity compared between expecting an auditory target and having
received a non-specific cue. **E,** 36 Hz frequency-tagging response between expecting an auditory target and
having received a non-specific cue. **F,** 40 Hz frequency-tagging response between expecting an auditory target
and having received a non-specific cue.
1086
1087
1088

1089

SUPPL Fig. 3 | Correlation of prestimulus alpha change from baseline with reaction time in the MEG study. A, the analysis was performed using a cluster-permutation approach, testing a correlation model against a 0-correlation model. Clusters significantly diverging from the 0-correlation model are presented topographically (p = .037). Additionally, median splits between fast and slow reaction time trials (p = .013; $t_{(25)}$ = -2.67) as well as correlation coefficients (p = .003; $t_{(25)}$ = -3.34) of these clusters are displayed for all participants. A negative correlation is visible between alpha modulation and reaction times in the last 500 ms before target onset when expecting a visual target. B, Correlation between alpha modulation and reaction time for each participant. Black diamonds represent trials from the first block (without distractor) and blue dots represent trials from the second block (with auditory distractor).

1099

SUPPL Fig. 4 | Correlation of 36 Hz change from baseline with reaction time in the MEG study. A, the analysis
was performed using a cluster-permutation approach, testing a correlation model against a 0-correlation
model. Clusters significantly diverging from the 0-correlation model are presented topographically (p = .040).
Additionally, median splits between fast and slow reaction time trials (p = .005; $t_{(25)}$ = -3.10) as well as
correlation coefficients (p = .002; $t_{(25)}$ = -3.46) of these clusters are displayed for all participants. A negative
correlation is visible between 36 Hz modulation and reaction times in the last 500 ms before target onset when
expecting a visual target. **B,** Correlation between 36 Hz modulation and reaction time for each participant.
Black diamonds represent trials from the first block (without distractor) and blue dots represent trials from the
second block (with auditory distractor).

1109

1110

1111

1112   **SUPPL Fig. 5 | Illustration of eye-tracking during the cue-to-target interval (2.5 − 0 s before target onset).** All
1113   datapoints of eye-positions during the cue-to-trial interval for all trials and all participants were plotted with 5%
1114   visibility on top of each other. Only 3% of datapoints showed eye-movement larger than 3 degrees of visual
1115   angle away from the fixation cross.

1116
1117   **SUPPL Fig. 6 | Individual ERP power spectra of the cue-to-target interval when anticipating an auditory target**
1118   **in the MEG-study.** Fast-fourier transformation was applied to the averaged trials using a dynamic hanning-
1119   tapered sliding time-window of 7 cycles per frequency. The Dotted line represents 40 Hz (auditory frequency-
1120   tagging).

1121

1122 **SUPPL Fig. 7 | Individual ERP power spectra of the cue-to-target interval when anticipating a visual target in**
1123 **the MEG-study.** Fast-fourier transformation was applied to the averaged trials using a dynamic hanning-
1124 tapered sliding time-window of 7 cycles per frequency. The Dotted line represents 36 Hz (auditory frequency-
1125 tagging).

1126

1127 **SUPPL Fig. 8 | Individual ERP power spectra of the cue-to-target interval when anticipating an auditory target**
1128 **in the EEG-study.** Fast-fourier transformation was applied to the averaged trials using a dynamic hanning-
1129 tapered sliding time-window of 7 cycles per frequency. The Dotted line represents 40 Hz (auditory frequency-
1130 tagging).

1131

1132 **SUPPL Fig. 9 | Individual ERP power spectra of the cue-to-target interval when anticipating a visual target in**
1133 **the EEG-study.** Fast-fourier transformation was applied to the averaged trials using a dynamic hanning-tapered
1134 sliding time-window of 7 cycles per frequency. The Dotted line represents 36 Hz (auditory frequency-tagging).
1135

1136

1137

1138

1139 **SUPPL Fig. 10 | Exemplary illustration of the correlation between alpha power (0.5 – 0 s before target onset)**
1140 **and 36 Hz steady-state response (0.5 – 0 s before target onset) for each participant in block 2 (distractors**
1141 **present).** Alpha activity was averaged over the significant group difference cluster for alpha condition
1142 differences (seed cluster). Frequency-tagging activity was averaged over the significant cluster in the
1143 correlation with the alpha seed activity.

1144
1145
1146
1147

anticipating auditory target
anticipating visual target

1148

1149 **SUPPL Fig. 11 | Illustration of bleeding over effects over a span of 4 Hz. A,** 40 Hz frequency-tagging data over
1150 the significant cluster differing between when expecting an auditory versus a visual target (identical to Fig. 9 in
1151 the manuscript). **B,** 44 Hz signal over the same cluster chosen for A. The analysis was identical with the analysis
1152 performed in A, apart from the frequency for the band-pass filter.

1153
1154

1155

1156

1157

SUPPL Figure 12 | Virtual channels for V1 and Helschl's gyrus. **A**, alpha power for the virtual channel created
in V1 (Calcerine_L and Calcerine_R from AAL atlas; Tzourio-Mazoyer et al., 2002, NeuroImage). A cluster
permutation analysis over time (between -2 and 0) revealed a significant condition difference between ~ -2 and
-1.7 s (*p* = 0.0449). **B**, 36 Hz frequency-tagging signal for the virtual channel created in V1 (equivalent to the
procedure in A). The same cluster permutation as performed in A revealed no significant condition differences.
C, 40 Hz frequency-tagging signal for the virtual channel created in Heschl's gryrus (Heschl_L and Heschl_R
from AAL atlas; Tzourio-Mazoyer et al., 2002, NeuroImage). The same cluster permutation as performed in A
revealed no significant condition differences.

1166

1167

1168

1169