Wolbachia inhibits ovarian formation and increases blood feeding rate in female *Aedes*

aegypti

Meng-Jia Lau[1*]; Perran A. Ross[1]; Nancy M. Endersby-Harshman[1]; Qiong Yang[1]; Ary A.

Hoffmann[1]

[1] Pest and Environmental Adaptation Research Group, Bio21 Institute and the School of

BioSciences, The University of Melbourne, Parkville, Victoria, Australia

* Corresponding author, email: lau.mengjia@gmail.com; ORCID: 0000-0003-2943-7022

Abstract

Wolbachia, a gram-negative endosymbiotic bacterium widespread in arthropods, is well-

known for changing the reproduction of its host in ways that increase its rate of spread, but

there are also costs to hosts that can reduce this. Here we investigated a novel reproductive

alteration of *Wolbachia w*AlbB on its mosquito host *Aedes aegypti*, based on studies of

mosquito life history traits, ovarian dissections and gene expression assays. We found that an

extended period of the larval stage as well as the egg stage, as previously shown, can increase

the proportion of *Wolbachia*-infected females that become infertile; the impact of *Wolbachia*

on infertility therefore accumulated before pupation. We found that ovarian formation was

blocked in infertile females, even though infected females had relatively lower *Wolbachia*

densities than fertile infected females. Infertile females also showed a higher frequency of

blood feeding following a prior blood meal, indicating that they do not enter a complete

gonotrophic cycle. Treatments leading to infertility decreased expression of genes related to

reproduction. Our results demonstrate effects related to the causes and consequences of

infertile *w*AlbB-infected *Ae. aegypti* females with implications for *Wolbachia* releases and

have evolutionary implications for *Wolbachia* infections in novel hosts.

Keywords: *Wolbachia*; reproductive alteration; *Aedes aegypti*; ovarian formation

Introduction

Wolbachia, a *Rickettsia*-like maternally inherited endosymbiotic bacterium widespread in

insects [1], can alter insect reproduction in multiple ways to enhance its transmission [2].

Specific alterations include parthenogenesis – infected females can produce female offspring

without mating; feminization – infected genetic males are transformed into functional

females and produce female offspring; male killing – infected immature males die while

females survive, and cytoplasmic incompatibility (CI) – females infected with the

predominant *Wolbachia* strain obtain a fertility advantage because males infected with

Wolbachia can cause female embryonic lethality when they mate with uninfected females or

females infected with a different *Wolbachia* strain [3, 4]. Aside from these alterations that

enhance the spread of *Wolbachia*, there can also be costs to the infection that slow its spread

[5-7]. These effects of *Wolbachia* are normally based on investigations of natural *Wolbachia*

strains that have existed in their hosts for many years, but are also frequently found in

artificial infections [8].

In the last decade, artificially introduced *Wolbachia* have been used to reduce the

transmission of arboviral diseases, including the *Wolbachia* strains *w*Mel [9] and *w*AlbB [10]

in *Aedes aegypti* mosquitoes. Studies on native hosts tend to indicate only a weak fitness cost

of *Wolbachia* [6], but *Wolbachia* can have diverse fitness costs in its newly introduced host

Ae. aegypti, such as life-shortening, heat sensitivity, reduced quiescent egg viability and

reduced blood feeding success [11-14]. The effects of *Wolbachia* in native and introduced

hosts point to the possibility of a mutualistic relationship between *Wolbachia* developing

over time as a consequence of co-evolution and adaptation. In fact, it is acknowledged that

sometimes *Wolbachia* can act as both parasite and mutualist [5, 15, 16]. As a result,

Wolbachia field release and establishment in new populations can be useful for studying

evolutionary changes in *Wolbachia* given that the invasion history of the population is known

[17-19].

In our previous study of the effects of *Wolbachia w*Mel and *w*AlbB on *Ae. aegypti* eggs, we

discovered a novel reproductive effect induced by *w*AlbB, in which infected females that

hatched from quiescent eggs lost their fertility despite successful mating, with the proportion

of female infertility being correlated with the duration of egg quiescence [20]. The *w*Mel

strain also causes female infertility following egg storage, but to a lesser extent than *w*AlbB

[20, 21]. Several other reproductive fitness costs of *Wolbachia*, including those on the

number and viability of eggs, can be attributed to the consequence of nutritional competition

between *Wolbachia* and its host [22] and represent quantitative effects. However, in our study

Wolbachia-infected females lost their fertility entirely, indicating a qualitative effect difficult

to attribute to nutritional competition [20], so we suspect the involvement of other

mechanisms.

In this paper, we investigate the causes and consequences of *Ae. aegypti* female infertility

induced by *Wolbachia w*AlbB infection after egg quiescence. Firstly, we tested if the effects

of *Wolbachia* on female infertility are permanent and dissected female ovaries. Secondly, we

performed a larval starvation experiment combined with real-time PCR assays to understand

the influence of *Wolbachia* on female development during different stages that could lead to

infertility effects. Finally, we tested whether the loss of ovaries in infertile females affected

the rate of female blood feeding. Our study highlights the novel nature of reproductive effects

altered by *Wolbachia w*AlbB in its new host *Ae. aegypti*, with implications for invasion of

Wolbachia into uninfected populations and the evolution of mutualism more generally.

Materials and Methods

Mosquito populations

We used uninfected and *Wolbachia w*AlbB-infected *Aedes aegypti*. The uninfected

population was derived from eggs collected in 2019 from regions in Cairns, Queensland,

Australia where *Ae*. *aegypti* were not infected with *Wolbachia*. The *w*AlbB-infected

population was generated by microinjection [23] and infected females were crossed to males

from the uninfected population regularly [24, 25] to maintain a similar genetic background

between populations. Mosquitoes were maintained in the laboratory following methods

described previously [26]. In the following experiments, uninfected females were hatched

from eggs stored at 26 ± 1 °C for less than one month. *w*AlbB-infected females from non-

stored eggs were hatched from freshly-laid eggs that had been dried and conditioned for one

week at 26 ± 1 °C. All lines tested in the same experiment had the same rearing and feeding

conditions.

Determination of female fertility status

To determine the fertility status of individual females in the experiments, we hatched a

mixture (approximately 1 : 1) of *w*AlbB-infected eggs that had been stored for two or four

months. Female mosquitoes were blood fed on the $4^{th} \pm 1$ day post emergence. 200 engorged

females were aspirated individually in 70 mL specimen cups with larval rearing water and

sandpaper strips to allow them to lay eggs. After a week, we separated females that laid

(fertile) or did not lay (infertile) eggs and grouped them separately in a 19.7-L BugDorm-1®

adult cage (MegaView Science Co., Ltd., Taichung City, Xitun District, Taiwan). The

process above is called "fertility separation". And to test the consistency of female infertility

status, we then provided fertile and infertile females with a second blood meal after the

"fertility separation", and isolated 20 of "fertile" and 20 of "infertile" females in 70 mL cups

individually for another week to observe if their fertility status was maintained. Next, we

dissected the remaining "fertile" and "infertile" females for their ovarian appearance: females

were killed by storing at -80 ℃ for 30 minutes then returning them to room temperature for

10 minutes before dissecting them in saline solution (0.9% sodium chloride) under a

compound light microscope (Motic B1 series, Australian Instrument Services Pty. Ltd.,

Australia).

Since we have confirmed the absence of ovaries from infertile females through dissection, we

further improved our method of identifying female fertility status by dissecting females

before "fertility separation". We did this dissection for 65 wAlbB-infected females that had

been stored as eggs for 11 weeks on the $4^{th} \pm 1$ day post-emergence (before their first blood

meal), and then used an NIS Elements BR imaging microscope (Nikon Instruments, Japan)

for photography.

Finally, to test the effect of wAlbB on the infertility of *Ae. aegypti* females with different

genetic background, we also dissected females from a recent wAlbB transinfection line that

had been repeatedly backcrossed to a Saudi Arabian background [27]. We dissected ovaries

from 113 wAlbB-infected mosquitoes that had been stored as eggs for 12 weeks, as well as

50 wAlbB-infected females stored as eggs for 1 week and 50 uninfected Saudi Arabian

females stored for 12 weeks (Supplementary material 1).

***Wolbachia* density**

After "fertility separation" in the previous section, we measured relative *Wolbachia* density

of fertile and infertile females for 16 individuals per group based on the $2^{\Delta Ct}$ method [28].

Screening based on real-time PCR assays followed methods described previously [29]. For

each replicate, all samples were set up in a 384-well white plate and measured by primers *aeg*

and *wMwA* [30]; two consistent replicates ($\Delta Ct < 1$) were obtained and their values were

averaged for the density analysis.

Impact of larval stage extension on mosquito infertility

To investigate whether the impact of *Wolbachia* infection on female infertility was only

mediated through the egg stage or whether the prepupal stage also had an accumulating

impact, we deprived 2^{nd} instar *w*AlbB-infected larvae of food for two weeks before feeding

them again until pupation and then followed this with a "fertility separation". In non-starved

controls, *w*AlbB-infected larvae were provided with food *ad libitum.* Each starvation

treatment was performed with both stored eggs (stored at 26 ± 1□ for 10 weeks) and non-

stored eggs (stored at 26 ± 1□ for 1 week) for a total of four treatments. For the group in

which infected females were neither stored nor starved, we had two replicates, while for the

other three experimental groups we had three replicates, with each replicate containing 30

individual females. Before the blood meal, 15 individual females from each treatment 4 ± 1

days post-emergence were screened for *Wolbachia* infection and *Wolbachia* density, using

the screening methods described in the previous section. After the fertile and infertile females

had been separated, females were also dissected to confirm their fertility status by checking

for the presence of ovaries. We also performed all four treatments with uninfected *Ae.*

aegypti (with similar egg storage and larval starvation conditions) and dissected 15

individuals 4 ± 1 days post-emergence from each replicate to check for presence of ovaries

(Supplementary material 1).

Gene expression assays

We selected three genes related to mosquito reproduction and tested their expression levels in

females at different developmental stages through a real-time PCR assay [29, 30]. One of the

genes tested was the *vitellogenin receptor* (*vgr*): following blood feeding, vitellogenin is

secreted from the fat body and internalized by ovaries through the receptor VgR [31-33]. The

other two genes were the *ecdysone receptor* (*ecr*) [34, 35] and the *eggshell organizing factor*

(*eof*); the vitellogenin transcript is regulated by *ecr* while *eof* is an essential gene encoding a

protein for eggshell formation at the late stage of egg production [36] (Supplementary

material 2). Gene expression level was quantified relative to a control gene, *RPS17* [37]. We

tested female pupae 1 ± 0.5 days post-pupation and female adults 1 ± 0.5 days post-

emergence in *w*AlbB-infected females that been stored as eggs for 14 weeks as well as both

*w*AlbB-infected and uninfected females that had not been stored, and considered the

uninfected mosquitoes as a control. For the *w*AlbB-infected females that had been stored as

eggs, we also tested gene expression levels in fertile and infertile females after testing female

fertility, and fertile females were treated as the control. Mosquito samples were given a

second blood meal and stored in RNA*later*® (Sigma Aldrich Cat No. R0901-100ML) on the

third day after feeding, before RNA extraction, reverse transcription and real-time PCR

assays were conducted. In all of the above comparisons, we screened 10 individuals from

each group; the details of RNA extraction and real-time PCR can be found in Supplementary

material 3.

Blood feeding rate of female mosquitoes

Normally, fully engorged fertile females are reluctant to feed within a gonotrophic cycle. In

this experiment, after determining female fertility, we measured blood feeding rate of female

mosquitoes by providing fertile and infertile females a third blood meal three days after their

second blood feeding, when engorged blood was almost digested. The same volunteer who

provided previous blood meals placed their forearm into a BugDorm cage for 15 minutes.

Fully engorged females were considered as having successfully fed after this feeding attempt.

Three replicates were completed with 20–30 individual females for each replicate. We also

weighed 20 random females that had fully fed and 20 that had not been provided with a blood

meal from all three replicates to estimate their blood meal weight using a Sartorius Analytical

balance BP 210 D (Sartorius, Gottigen, Germany, Readability: 0.01 mg).

Statistics

We used R v. 3.6.0 with R studio v. 1.1.453 to conduct data analyses and visualizations [38]

using the "car" library for ANOVA [39], the base library for other statistical analyses, and the

"ggplot2" library for visualization [40]. After real-time PCR measurement, we used the $2^{-\Delta\Delta Ct}$

method to compare gene expression levels [41], using either uninfected females from pupae

or young adult stage as controls for the same stage, or fertile females as the control group of

infertile females after blood feeding. The values of $2^{-\Delta\Delta Ct}$ were natural log-transformed for

ANOVA analysis. For *Wolbachia* density we used the $2^{\Delta Ct}$ method [28] to calculate relatively

density before log10 transformation for ANOVA analysis. We also used Tukey's honest

significant difference (HSD) tests [42] for further pairwise comparisons. For the proportion

data from larval starvation and blood feeding experiments, we analysed data with binomial

logistic regression models [43]. In the blood feeding rate test, student's t-tests were used to

test for changes of mosquito weight after blood feeding.

Results

Immature ovaries and lower *Wolbachia* densities in infertile *Aedes aegypti* females

We did a "fertility separation" for *w*AlbB-infected females and provided a second blood meal

to fertile and infertile females. We confirmed that all females we scored as fertile or infertile

after the first gonotrophic cycle maintained their phenotype in the second gonotrophic cycle.

We then dissected and compared the ovarian appearance of fertile and infertile females with

an Australian or Saudi genetic background and discovered that ovaries could not be observed

in infertile females under a microscope (Figure 1). Only occasionally (around one in ten) can

immature ovarian structure be seen, but with a narrow width similar to that of a Malpighian

tube (Supplementary material 4). We also found that fertile females had higher relative

densities of *Wolbachia* (Figure 2: ANOVA: $F_{1,30}$ = 8.632, P = 0.006, fertile: mean ± se = 5.15

± 0.906; infertile: mean ± se = 2.290 ± 0.673).

Figure 1. The appearance of internal organs in *w*AlbB-infected *Aedes aegypti* mosquitoes. (a)

Infertile females lack observable ovaries, while ovaries are present in (b) fertile females. The

background of the photographs was removed for clarity, with the original photographs

presented in Supplementary material 4.

Figure 2. Box plots of relative *Wolbachia* density of fertile and infertile *w*AlbB-infected

females after females were separated into fertile and infertile groups (one week after blood

meal). Densities are based on two consistent real-time PCR replicates of 16 females from

each group.

Impact of larval stage extension on mosquito infertility

We starved mosquito larvae to increase the duration of the larval stage from approximately

one week to three weeks to test if there is any impact of larval stage extension on infertility in

*w*AlbB-infected females. We found significant effects of both storage and starvation

treatment, with an interaction between these terms (Figure 3A: storage: $F_{1,7} = 256.304$, P <

0.001; starvation: $F_{1,7} = 32.412$, P < 0.001; interaction: $F_{1,7} = 21.515$, P = 0.002). Infertility

rates associated with *w*AlbB infection increased further with extension of the larval stage

over and above that seen with stored eggs by 15%, and the extension of the larval stage by

starvation also induced infertility of around 10% by itself. For uninfected *Ae. aegypti* treated

in the same manner, all females that were dissected contained ovaries, indicating the essential

role of *Wolbachia w*AlbB in inducing female infertility during larval starvation and egg

storage (Supplementary material 1).

Figure 3. (A) Boxplots showing the proportion of infertile females for *w*AlbB-infected

mosquitoes when stored or not stored as eggs and when larvae had been starved or not

starved. Values above boxplots represent corresponding averaged percentages based on two

replicates for control and three replicates for treatments with each replicate containing 30

individuals. (B) Boxplots of relative *Wolbachia* density in 4 ± 1 days post-emergence females

Each group based on two consistent real-time PCR replicates of 15 individuals with failure of

detection were excluded (Supplementary material 5).

We also screened for *Wolbachia* in all the treatments to confirm *Wolbachia* infection status.

We failed to detect an infection in five out of 60 individuals and these were excluded from

the density analysis (Supplementary material 5). *Wolbachia* density was significantly

influenced by larval starvation but not egg quiescence, with a significant interaction (Figure

3B: ANOVA: starvation: $F_{1,51} = 7.145$, P = 0.010; egg quiescence: $F_{1,51} = 2.475$, P = 0.122,

interaction: $F_{1,51} = 8.396$, P = 0.006). Egg quiescence decreased *Wolbachia* density, but

density increased with larval starvation, potentially cancelling out this effect [44].

Gene expression assays

We selected three genes related to reproductive development and tested their expression level

at different developmental stages. At the pupal stage, there are significant differences among

the three groups of mosquitoes, but no interaction with the genes (Figure 4A, ANOVA:

mosquito group: $F_{2,81} = 26.717$, p < 0.001; genes: $F_{2,81} = 1.044$, p = 0.357; interaction: $F_{4,81} =$

0.423, p = 0.792). Specifically, in Tukey's HSD test, *w*AlbB-infected females that been

stored as eggs had lower expression levels for all three genes than expression levels in

uninfected females, but only gene *ecr* had a lower expression level in *w*AlbB-infected

females that had not been stored.

For young adults 1 ± 0.5 days post-emergence, significant differences of gene expression

levels were found among mosquito groups, but there was no interaction with the genes

(Figure 4B, mosquito group: $F_{2,81} = 15.641$, $p < 0.001$; genes: $F_{2,81} = 0.383$, $p = 0.683$;

interaction: $F_{4,81} = 7.100$, $p = 0.792$). Specifically, only differences for *vgr* expression were

significant between *w*AlbB-infected females that been stored and the other two mosquito

groups, with some individuals from stored eggs having very low expression levels - these

females are probably infertile. The *w*AlbB infection may have increased expression of this

gene, indicating that there may also be some impact of *Wolbachia* on reproduction of non-

stored females (Figure 4B).

Figure 4. Boxplots of relative expression of reproduction-related genes *ecdysone receptor*

(*ecr*), *eggshell organizing factor* (*eof*) and *vitellogenin receptor* (*vgr*) in female (A) pupae

and (B) young adults (1 ± 0.5 days post-emergence). Values with the same letter are not

significantly different based on Tukey's HSD tests. Results are based on two consistent real-

time PCR replicates of ten individual females from each group.

After "fertility separation", female mosquitoes were provided a second blood meal and were

screened for gene expression levels three days later. Significant differences were found

between fertile and infertile females, and also between genes and their interaction (Figure 5,

mosquito group: $F_{1,54} = 235.696$, $p < 0.001$; genes: $F_{2,54} = 68.553$, $p < 0.001$; interaction: $F_{2,54}$

$= 68.553$, $p < 0.001$). All three genes showed significant differences between fertile and

infertile females for expression, especially in the case of *vgr* whose expression in infertile

mosquitoes was only around 0.01% of that seen in fertile mosquitoes, likely reflecting the

fact that infertile females were unable to produce eggs.

Figure 5. Boxplots of relative expression of reproduction-related genes *ecdysone receptor*

(*ecr*), *eggshell organizing factor* (*eof*) and *vitellogenin receptor* (*vgr*) in fertile and infertile

females one week after their second blood meal. Values with the same letter are not

significantly different according to Tukey's HSD tests. Results are based on two consistent

real-time PCR replicates of ten individual females from each group.

Blood feeding rate

After "fertility separation", we provided female mosquitoes a third blood meal on the third

day after they had become engorged with their second blood meal to test their blood feeding

behaviour. Infertile females had a higher proportion feeding compared to fertile and

uninfected females (Figure 6A, $F_{2,6} = 61.037$, $p < 0.001$). We also compared the weight of

fully fed and unfed females and noted significant differences between mosquito groups,

feeding status and their interaction (Figure 6B, ANOVA: mosquito group: $F_{1,73} = 29.06$, $p <$

0.001; feeding: $F_{1,73} = 322.89$, $p < 0.001$; interaction: $F_{2,73} = 15.26$, $p < 0.001$). Specifically,

significant weight increases were recorded for both *w*AlbB-infected fertile females ($t_{35} =$

9.627, $p < 0.001$) and *w*AlbB-infected infertile females ($t_{38} = 15.658$, $p < 0.001$), but the

weight of fully fed fertile females was significantly lower compared to infertile females ($t_{35} =$

-4.645, $p < 0.001$), while there was no significant difference between unfed fertile and

infertile females ($t_{38} = -1.096$, $p = 0.280$), suggesting that infertile females took in a larger

amount of blood during successive feeding.

Figure 6. Boxplots of (A) female proportion that blood fed three days after they had been

fully fed, in which three replicates were completed with 20–30 individual females for each

replicate; and (B) the weights of fed and unfed females. 20 females that had fully fed and 20

that had not been provided with a blood meal from all three replicates were weighed at

random.

Discussion

Wolbachia is a well-known bacterium that can alter the reproduction of its host to benefit its

spread. In our previous study, *Wolbachia w*AlbB was found to cause *Ae. aegypti* females to

become infertile following egg storage [20], inhibiting vertical transmission. In this paper, we

investigated this phenomenon and obtained the following results: 1) in infertile females, the

development of ovaries has been interrupted by *Wolbachia*; 2) the effect of *Wolbachia* on

female fertility accumulates across pre-pupal stages, in that the frequency of female infertility

depends on the duration before metamorphosis (including egg and larval stages); and 3)

infertile females maintain other female characteristics such as mating and blood feeding, but

they do not enter a gonotrophic cycle and blood feeding occurs more frequently. We,

therefore, confirmed that *Wolbachia w*AlbB altered the reproductive process in its new host,

Ae. aegypti.

The effect of *Wolbachia* on female infertility is different from other fitness impacts of

Wolbachia and the four typical reproductive alterations (parthenogenesis, feminization, male

killing and cytoplasmic incompatibility) in which *Wolbachia* alters host reproduction in ways

that increase its vertical transmission success [4, 45]. The infertility effect we describe here

converts functional females into non-functional females without male characteristics,

suggesting different *Wolbachia*-related mechanisms to those involved in other phenomena

like feminization [46, 47]. By testing the expression level of *vgr, ecr* and *eof* at the female

pupal, young adult before blood feeding and old adult after blood feeding stages , we found

substantial differences in gene expression between fertile and infertile groups. For pupae,

large variation in expression levels were found among individuals, especially for expression

of the gene *vgr*, probably reflecting the fact that its expression mainly occurs in a narrow

period during the pupal stage. *w*AlbB-infected *Ae. aegypti* females that had been stored long-

term as eggs had lower expression levels of the three reproductive genes compared with

uninfected females, especially for the gene *vgr* at the young adult stage, suggesting the

formation and development of oocytes was impacted. This is also supported by the very low

relative expression of *vgr* and *eof* in infertile females when compared with fertile females at

the third day after blood feeding, and the absence of mature ovaries in infertile females.

Moreover, we found much lower expression of *ecr* in *w*AlbB-infected female pupae hatched

from long-stored eggs, while this difference was not found in young adults. As *ecr* is

expressed in a variety of organs to encode the receptor of a hormone [48], the down-

regulation of *ecr* may indicate that egg quiescence has a substantial impact on the pupation

process of *w*AlbB-infected females. However, interactions between *Wolbachia* and specific

reproductive signals and mechanisms are poorly understood and require further investigation

for example, through hormone level tests, transcriptome analysis and pathway analysis [49,

50].

The novel reproductive alteration induced by *Wolbachia w*AlbB could slow the spread of the

infection in mosquito populations as females are less likely to produce offspring. This may

occur not only when the egg desiccation period is extended during dry and warm seasons [51,

52], but also when the larval period is extended due to poor provision of food [53-55]. These

effects can reduce the efficiency of *Wolbachia* invasion which aims to inhibit the

transmission of arboviruses [8, 25, 56]. While *w*AlbB has established successfully in

Malaysia where the warm and humid year-round climate precludes extended egg desiccation

periods and promotes fast development of larvae [10, 57], our discovery provides guidance

for future releases in other climates and highlights the importance of future monitoring [20].

Our observation that infertile females show an increased rate of blood feeding highlights a

potential risk of increased nuisance biting following a *Wolbachia* release program, which

may lead to community discontentment. Normal *Ae. aegypti* females ingest human blood to

obtain unique nutrients for egg development [58, 59]. However, while infertile females lack

ovaries, they still blood feed. Previous research found that multiple feeding within a

gonotrophic cycle is caused by nutritional reserve depletion or feeding interruption [60].

Nevertheless, we found a large proportion of infertile females fed again on the third day after

their second blood meal, indicating they do not follow a gonotrophic cycle. It is unclear if

and how the process of blood digestion is impacted by infertility. Infertile females also took

in a larger amount of blood, though this might be a compensatory reaction towards starvation

at egg and larval stages. This behavioural change of *Wolbachia*-infected infertile females has

significant implications for vector control population release strategies [8, 61]; in particular it

would be interesting to test whether the ability of these infertile females to carry and transmit

arbovirus has changed as the density of their *Wolbachia* infection may be lower than that of

fertile females [62, 63]. Similar concerns have been raised about increased biting frequencies

of irradiated female mosquitoes in sterile insect technique programs [64], however the

prevalence of infertile females is likely to be much higher in *Wolbachia*-infected mosquito

populations compared to irradiated females that are only released accidentally.

The reproductive alterations of *Wolbachia* discovered in this study have evolutionary

implications. So far, our work relates to *Ae. aegypti*, where the *Wolbachia w*AlbB was

artificially introduced recently from a close relative of *Ae. aegypti*, *Aedes albopictus*, in order

to reduce the transmission of arboviral diseases [65]. A *Wolbachia* strain may induce similar

fitness costs in a new host when compared to its native host, such as in the case of life-

shortening caused by *w*MelPop which is expressed in both *Drosophila melanogaster* and *Ae.*

aegypti [11], and remains stably expressed following long-term laboratory culture [66]. We

do not yet know if there are similar reproductive effects of *Wolbachia w*AlbB expressed in its

native host, *Ae. albopictus*. On the other hand, we have also discovered a smaller loss of

female infertility in *w*Mel infected *Ae. aegypti* [20], though the native host of *w*Mel, *D.*

melanogaster, does not enter egg quiescence, although there is a reduced fecundity of *w*Mel-

infected females under dormancy conditions [67]. Although other *Wolbachia* strains remain

to be investigated, it is possible that *Ae. aegypti* has mechanisms underlying ovarian

formation that can be interrupted by endosymbionts. However, it is unclear why the

frequency of infertile females increases when the density of *Wolbachia* decreases as the result

of increased periods of egg storage or larval development time, and why infertile females

have lower *Wolbachia* densities than fertile females in the same treatment. There may be

cumulative effects of *Wolbachia* not evident from density measures at early stages of

development contributing to failure of ovarian formation. One possible hypothesis which
requires further investigation is that *Wolbachia* competes for some essential but sparse
nutrients with its host *Ae. aegypti* at an early life stage.

In conclusion, we further investigated a novel reproductive alteration of *Wolbachia* that we
discovered previously [20]. We confirmed that *Ae. aegypti* females infected with *Wolbachia*
*w*AlbB can become infertile when they are unable to form functional ovaries during
metamorphosis, but these females retain other female characteristics, leading to an increased
biting frequency. Our study provides significant guidance for future *Wolbachia* releases and
has important evolutionary implications for understanding the reproductive alteration of
Wolbachia, especially in novel hosts.

Funding

This research was funded by the National Health and Medical Research Council (1132412,
1118640, www.nhmrc.gov.au). The funders had no role in the study design, data collection
and analysis, decision to publish, or preparation of the manuscript.

Acknowledgment

We thank Véronique Paris for her assistance in weighing mosquitoes.

1.   Zug R., Hammerstein P. 2012 Still a host of hosts for *Wolbachia*: analysis of recent
data suggests that 40% of terrestrial arthropod species are infected. *PLoS One* **7**(6), e38544.
(doi:10.1371/journal.pone.0038544).
2.   Stouthamer R., Breeuwer J.A., Hurst G.D. 1999 *Wolbachia pipientis*: microbial
manipulator of arthropod reproduction. *Annu Rev Microbiol* **53**, 71-102.
(doi:10.1146/annurev.micro.53.1.71).
3.   Werren J.H., Baldo L., Clark M.E. 2008 *Wolbachia*: master manipulators of
invertebrate biology. *Nat Rev Microbiol* **6**(10), 741.
4.   O'Neill S.L., Hoffmann A., Werren J. 1997 *Influential passengers: inherited
microorganisms and arthropod reproduction*, Oxford University Press.
5.   Zug R., Hammerstein P. 2015 Bad guys turned nice? A critical assessment of
Wolbachia mutualisms in arthropod hosts. *Biol Rev Camb Philos Soc* **90**(1), 89-111.
(doi:10.1111/brv.12098).
6.   Hoffmann A.A., Turelli M., Harshman L.G. 1990 Factors affecting the distribution of
cytoplasmic incompatibility in *Drosophila simulans*. *Genetics* **126**(4), 933-948.
7.   Perrot‐Minnot M.J., Cheval B., Migeon A., Navajas M. 2002 Contrasting effects of
Wolbachia on cytoplasmic incompatibility and fecundity in the haplodiploid mite
Tetranychus urticae. *J Evol Biol* **15**(5), 808-817.

8.   Ross P.A., Turelli M., Hoffmann A.A. 2019 Evolutionary ecology of *Wolbachia*
releases for disease control. *Annu Rev Genet* **53**, 93-116. (doi:10.1146/annurev-genet-
112618-043609).
9.   Ryan P.A., Turley A.P., Wilson G., Hurst T.P., Retzki K., Brown-Kenyon J., Hodgson L.,
Kenny N., Cook H., Montgomery B.L. 2019 Establishment of *w*Mel *Wolbachia* in *Aedes*
aegypti mosquitoes and reduction of local dengue transmission in Cairns and surrounding
locations in northern Queensland, Australia. *Gates Open Res* **3**, 1547.
10.   Nazni W.A., Hoffmann A.A., NoorAfizah A., Cheong Y.L., Mancini M.V., Golding N.,
Kamarul G.M., Arif M.A., Thohir H., NurSyamimi H. 2019 Establishment of *Wolbachia* strain
*w*AlbB in Malaysian populations of *Aedes aegypti* for dengue control. *Curr Biol* **29**(24), 4241-
4248.
11.   McMeniman C.J., Lane R.V., Cass B.N., Fong A.W.C., Sidhu M., Wang Y.-F., O'Neill S.L.
2009 Stable introduction of a life-shortening *Wolbachia* infection into the mosquito *Aedes*
aegypti. *Science* **323**(5910), 141-144. (doi:10.1126/science.1165326).
12.   Turley A.P., Moreira L.A., O'Neill S.L., McGraw E.A. 2009 *Wolbachia* infection reduces
blood-feeding success in the dengue fever mosquito, *Aedes aegypti*. *PLoS Negl Trop Dis* **3**(9),
e516.
13.   Yeap H.L., Mee P., Walker T., Weeks A.R., O'Neill S.L., Johnson P., Ritchie S.A.,
Richardson K.M., Doig C., Endersby N.M., et al. 2011 Dynamics of the "Popcorn" *Wolbachia*
infection in outbred *Aedes aegypti* informs prospects for mosquito vector control. *Genetics*
187(2), 583-595. (doi:10.1534/genetics.110.122390).
14.   Ware-Gilmore F., Sgrò C.M., Xi Z., Dutra H.L.C., Jones M.J., Shea K., Hall M.D.,
Thomas M.B., McGraw E.A. 2021 Microbes increase thermal sensitivity in the mosquito
Aedes aegypti, with the potential to change disease distributions. *PLoS Negl Trop Dis* **15**(7),
e0009548. (doi:10.1371/journal.pntd.0009548).
15.   Fenn K., Blaxter M. 2006 *Wolbachia* genomes: revealing the biology of parasitism
and mutualism. *Trends Parasitol* **22**(2), 60-65. (doi:10.1016/j.pt.2005.12.012).
16.   Weeks A.R., Turelli M., Harcombe W.R., Reynolds K.T., Hoffmann A.A. 2007 From
parasite to mutualist: rapid evolution of *Wolbachia* in natural populations of *Drosophila*.
PLoS Biol **5**(5), e114.
17.   Lau M.J., Schmidt T.L., Yang Q., Chung J., Sankey L., Ross P.A., Hoffmann A.A. 2021
Genetic stability of *Aedes aegypti* populations following invasion by *w*Mel *Wolbachia*. *BMC*
Genomics **22**(1), 894. (doi:10.1186/s12864-021-08200-1).
18.   Carrington L.B., Hoffmann A.A., Weeks A.R. 2010 Monitoring long-term evolutionary
changes following *Wolbachia* introduction into a novel host: the *Wolbachia* popcorn
infection in *Drosophila simulans*. *Proc Biol Sci* **277**(1690), 2059-2068.
19.   Ford S.A., Allen S.L., Ohm J.R., Sigle L.T., Sebastian A., Albert I., Chenoweth S.F.,
McGraw E.A. 2019 Selection on *Aedes aegypti* alters *Wolbachia*-mediated dengue virus
blocking and fitness. *Nat Microbiol* **4**(11), 1832-1839. (doi:10.1038/s41564-019-0533-3).
20.   Lau M.J., Ross P.A., Hoffmann A.A. 2021 Infertility and fecundity loss of *Wolbachia*-
infected *Aedes aegypti* hatched from quiescent eggs is expected to alter invasion dynamics.
PLoS Negl Trop Dis **15**(2), e0009179. (doi:10.1371/journal.pntd.0009179).
21.   Ross P.A., Hoffmann A.A. 2022 Local adaptation of *Aedes aegypti* mosquitoes to
Wolbachia-induced fitness costs. *bioRxiv*.
22.   Allman M.J., Fraser J.E., Ritchie S.A., Joubert D.A., Simmons C.P., Flores H.A. 2020
Wolbachia's deleterious impact on *Aedes aegypti* egg development: The potential role of
nutritional parasitism. *Insects* **11**(11), 735. (doi:10.3390/insects11110735).

23. Xi Z.Y., Khoo C.C.H., Dobson S.L. 2005 *Wolbachia* establishment and invasion in an
Aedes aegypti laboratory population. *Science* **310**(5746), 326-328.
(doi:10.1126/science.1117607).
24. Garcia G.d.A., Sylvestre G., Aguiar R., da Costa G.B., Martins A.J., Lima J.B.P.,
Petersen M.T., Lourenço-de-Oliveira R., Shadbolt M.F., Rašić G., et al. 2019 Matching the
genetics of released and local *Aedes aegypti* populations is critical to assure *Wolbachia*
invasion. *PLoS Negl Trop Dis* **13**(1), e0007023. (doi:10.1371/journal.pntd.0007023).
25. Hoffmann A.A., Montgomery B., Popovici J., Iturbe-Ormaetxe I., Johnson P., Muzzi F.,
Greenfield M., Durkan M., Leong Y., Dong Y. 2011 Successful establishment of *Wolbachia* in
Aedes populations to suppress dengue transmission. *Nature* **476**(7361), 454-457.
26. Ross P.A., Axford J.K., Richardson K.M., Endersby-Harshman N.M., Hoffmann A.A.
2017 Maintaining *Aedes aegypti* mosquitoes infected with *Wolbachia*. *J Vis Exp* **14**(126),
e56124.
27. Ross P.A., Gu X., Robinson K.L., Yang Q., Cottingham E., Zhang Y., Yeap H.L., Xu X.,
Endersby-Harshman N.M., Hoffmann A.A. 2021 A *w*AlbB *Wolbachia* Transinfection Displays
Stable Phenotypic Effects across Divergent *Aedes aegypti* Mosquito Backgrounds. *Appl*
Environ Microbiol **87**(20), e0126421. (doi:10.1128/aem.01264-21).
28. Unckless R.L., Boelio L.M., Herren J.K., Jaenike J. 2009 *Wolbachia* as populations
within individual insects: causes and consequences of density variation in natural
populations. *Proc Biol Sci* **276**(1668), 2805-2811. (doi:10.1098/rspb.2009.0287).
29. Lee S.F., White V.L., Weeks A.R., Hoffmann A.A., Endersby N.M. 2012 High-
throughput PCR assays to monitor *Wolbachia* infection in the dengue mosquito (*Aedes*
aegypti) and *Drosophila simulans*. *Appl Environ Microbiol* **78**(13), 4740-4743.
30. Lau M.-J., Hoffmann A.A., Endersby-Harshman N.M. 2021 A diagnostic primer pair to
distinguish between *w*Mel and *w*AlbB *Wolbachia* infections. *PLoS One* **16**(9), e0257781.
(doi:10.1371/journal.pone.0257781).
31. Raikhel A.S., Lea A.O. 1983 Previtellogenic development and vitellogenin synthesis in
the fat body of a mosquito: an ultrastructural and immunocytochemical study. *Tissue Cell*
15(2), 281-299. (doi:10.1016/0040-8166(83)90023-x).
32. Sappington T.W., Raikhel A.S. 1998 Molecular characteristics of insect vitellogenins
and vitellogenin receptors. *Insect Biochem Mol Biol* **28**(5-6), 277-300. (doi:10.1016/s0965-
1748(97)00110-0).
33. Tufail M., Takeda M. 2009 Insect vitellogenin/lipophorin receptors: molecular
structures, role in oogenesis, and regulatory mechanisms. *J Insect Physiol* **55**(2), 87-103.
(doi:10.1016/j.jinsphys.2008.11.007).
34. Hodin J., Riddiford L.M. 1998 The ecdysone receptor and ultraspiracle regulate the
timing and progression of ovarian morphogenesis during *Drosophila metamorphosis*. *Dev*
Genes Evol **208**(6), 304-317. (doi:10.1007/s004270050186).
35. Martín D., Wang S.F., Raikhel A.S. 2001 The vitellogenin gene of the mosquito *Aedes*
aegypti is a direct target of ecdysteroid receptor. *Mol Cell Endocrinol* **173**(1-2), 75-86.
(doi:10.1016/s0303-7207(00)00413-5).
36. Isoe J., Koch L.E., Isoe Y.E., Rascón A.A., Jr., Brown H.E., Massani B.B., Miesfeld R.L.
2019 Identification and characterization of a mosquito-specific eggshell organizing factor in
Aedes aegypti mosquitoes. *PLoS Biol* **17**(1), e3000068. (doi:10.1371/journal.pbio.3000068).
37. Dzaki N., Ramli K.N., Azlan A., Ishak I.H., Azzam G. 2017 Evaluation of reference
genes at different developmental stages for quantitative real-time PCR in *Aedes aegypti*. *Sci*
Rep **7**, 43618. (doi:10.1038/srep43618).

38.   Team R.C. 2019 R: A language and environment for statistical computing. R
Foundation for Statistical Computing, Vienna, Austria.
39.   Fox J., Weisberg S., Adler D., Bates D., Baud-Bovy G., Ellison S., Firth D., Friendly M.,
Gorjanc G., Graves S. 2012 Package 'car'. *Vienna: R Foundation for Statistical Computing*.
40.   Wickham H. 2016 *ggplot2: elegant graphics for data analysis*, Springer.
41.   Livak K.J., Schmittgen T.D. 2001 Analysis of relative gene expression data using real-
time quantitative PCR and the 2(-delta delta C(T)) method. *Methods* **25**(4), 402-408.
(doi:10.1006/meth.2001.1262).
42.   Tukey J.W. 1949 Comparing individual means in the analysis of variance. *Biometrics*
5(2), 99-114.
43.   Hastie T.J. 2017 Generalized additive models. In *Statistical models in S* (ed. Science
S.), pp. 249-307, Routledge.
44.   Lau M.J., Ross P.A., Endersby-Harshman N.M., Hoffmann A.A. 2020 Impacts of low
temperatures on *Wolbachia* (Rickettsiales: Rickettsiaceae)-infected *Aedes aegypti* (Diptera:
Culicidae). *J Med Entomol* **57**(5), 1567-1574. (doi:10.1093/jme/tjaa074).
45.   Werren J.H. 1997 Biology of *Wolbachia*. *Annu Rev Entomol* **42**, 587-609.
46.   Rigaud T., Juchault P., Mocquard J.P. 1997 The evolution of sex determination in
isopod crustaceans. *Bioessays* **19**(5), 409-416.
47.   Bandi C., Dunn A.M., Hurst G.D., Rigaud T. 2001 Inherited microorganisms, sex-
specific virulence and reproductive parasitism. *Trends Parasitol* **17**(2), 88-94.
(doi:10.1016/s1471-4922(00)01812-2).
48.   Koelle M.R., Talbot W.S., Segraves W.A., Bender M.T., Cherbas P., Hogness D.S. 1991
The *Drosophila EcR* gene encodes an ecdysone receptor, a new member of the steroid
receptor superfamily. *Cell* **67**(1), 59-77. (doi:10.1016/0092-8674(91)90572-g).
49.   Belles X., Piulachs M.D. 2015 Ecdysone signalling and ovarian development in insects:
from stem cells to ovarian follicle formation. *Biochim Biophys Acta* **1849**(2), 181-186.
(doi:10.1016/j.bbagrm.2014.05.025).
50.   Dansereau D.A., Lasko P. 2008 The development of germline stem cells in *Drosophila*.
Methods Mol Biol **450**, 3-26. (doi:10.1007/978-1-60327-214-8_1).
51.   Juliano S.A., O'Meara G.F., Morrill J.R., Cutwa M.M. 2002 Desiccation and thermal
tolerance of eggs and the coexistence of competing mosquitoes. *Oecologia* **130**(3), 458-469.
52.   Faull K.J., Williams C.R. 2015 Intraspecific variation in desiccation survival time of
Aedes aegypti (L.) mosquito eggs of Australian origin. *J Vector Ecol* **40**(2), 292-300.
(doi:10.1111/jvec.12167).
53.   Ross P.A., Endersby N.M., Hoffmann A.A. 2016 Costs of three *Wolbachia* infections
on the survival of *Aedes aegypti* larvae under starvation conditions. *PLoS Negl Trop Dis* **10**(1),
e0004320.
54.   Barrera R., Medialdea V. 1996 Development time and resistance to starvation of
mosquito larvae. *J Nat Hist* **30**(3), 447-458.
55.   Barrera R. 1996 Competition and resistance to starvation in larvae of container-
inhabiting *Aedes* mosquitoes. *Ecol Entomol* **21**(2), 117-127.
56.   Koh C., Audsley M.D., Di Giallonardo F., Kerton E.J., Young P.R., Holmes E.C., McGraw
E.A. 2019 Sustained *Wolbachia*-mediated blocking of dengue virus isolates following serial
passage in *Aedes aegypti* cell culture. *Virus Evol* **5**(1), vez012.
57.   Ahmad N.A., Mancini M.V., Ant T.H., Martinez J., Kamarul G.M.R., Nazni W.A.,
Hoffmann A.A., Sinkins S.P. 2021 *Wolbachia* strain *w*AlbB maintains high density and dengue

inhibition following introduction into a field population of *Aedes aegypti*. *Philos Trans R Soc*
Lond B Biol Sci **376**(1818), 20190809. (doi:10.1098/rstb.2019.0809).

58.   Gulia-Nuss M., Elliot A., Brown M.R., Strand M.R. 2015 Multiple factors contribute to
anautogenous reproduction by the mosquito *Aedes aegypti*. *J Insect Physiol* **82**, 8-16.
(doi:10.1016/j.jinsphys.2015.08.001).

59.   Scott T.W., Naksathit A., Day J.F., Kittayapong P., Edman J.D. 1997 A fitness
advantage for *Aedes aegypti* and the viruses it transmits when females feed only on human
blood. *Am J Trop Med Hyg* **57**(2), 235-239. (doi:10.4269/ajtmh.1997.57.235).

60.   Farjana T., Tuno N. 2013 Multiple blood feeding and host-seeking behavior in *Aedes*
aegypti and *Aedes albopictus* (Diptera: Culicidae). *J Med Entomol* **50**(4), 838-846.
(doi:10.1603/me12146).

61.   Almeida L., Duprez M.G., Privat Y., Vauchelet N. 2019 Mosquito population control
strategies for fighting against arboviruses. *Math Biosci Eng* **16**(6), 6274-6297.
(doi:10.3934/mbe.2019313).

62.   Johnson K.N. 2015 The impact of *Wolbachia* on virus infection in mosquitoes. *Viruses*
7(11), 5705-5717. (doi:10.3390/v7112903).

63.   Terradas G., McGraw E.A. 2017 *Wolbachia*-mediated virus blocking in the mosquito
vector *Aedes aegypti*. *Curr Opin Insect Sci* **22**, 37-44. (doi:10.1016/j.cois.2017.05.005).

64.   Moretti R., Lampazzi E., Damiani C., Fabbri G., Lombardi G., Pioli C., Desiderio A.,
Serrao A., Calvitti M. 2022 Increased biting rate and decreased *Wolbachia* density in
irradiated Aedes mosquitoes. *Parasit Vectors* **15**(1), 67. (doi:10.1186/s13071-022-05188-9).

65.   Xi Z., Dean J.L., Khoo C., Dobson S.L. 2005 Generation of a novel *Wolbachia* infection
in *Aedes albopictus* (Asian tiger mosquito) via embryonic microinjection. *Insect Biochem Mol*
Biol **35**(8), 903-910.

66.   Ross P.A., Axford J.K., Callahan A.G., Richardson K.M., Hoffmann A.A. 2020 Persistent
deleterious effects of a deleterious *Wolbachia* infection. *PLoS Negl Trop Dis* **14**(4),
e0008204-e0008204. (doi:10.1371/journal.pntd.0008204).

67.   Kriesner P., Conner W.R., Weeks A.R., Turelli M., Hoffmann A.A. 2016 Persistence of
a *Wolbachia* infection frequency cline in *Drosophila melanogaster* and the possible role of
reproductive dormancy. *Evolution* **70**(5), 979-997. (doi:10.1111/evo.12923).

613