Title: T cell Aryl Hydrocarbon Receptor Activity Tunes the Gut Microenvironment to Sustain Autoimmunity and Neuroinflammation.

Short Title: T cell AHR and neuroinflammation

Authors: Andrea R. Merchak,[1, 2, 3] Hannah J. Cahill,[1] Lucille C. Brown,[1] Ryan M. Brown,[1,2, 3] Courtney Rivet-Noor,[1,2, 3] Rebecca M. Beiter,[1,2, 3] Erica R. Slogar,[1] Deniz G. Olgun,[1] and Alban Gaultier[1, 3]*

Affiliations:
[1]Department of Neuroscience, University of Virginia, Charlottesville, VA, USA

[2]Neuroscience Graduate Program, University of Virginia, Charlottesville, VA, USA

[3]Center for Brain Immunology and Glia, University of Virginia, Charlottesville, VA, USA

*Corresponding author. Email: ag7h@virginia.edu.

Abstract:

Multiple sclerosis (MS) is a T cell driven autoimmune disease that attacks the myelin of the central nervous system and currently has no cure. MS etiology is linked to both the gut flora and external environmental factors but this connection is not well understood. One immune system regulator responsive to non-pathogenic external stimuli is the aryl hydrocarbon receptor (AHR). The AHR, which binds diverse molecules present in the environment in barrier tissues, is a therapeutic target for MS. However, AHR's precise function in T lymphocytes, the orchestrators of MS, has not been described. Here we show that in a mouse model of MS, T cell specific *Ahr* knockout leads to recovery driven by a decrease in T cell fitness. At the mechanistic level, we demonstrate that the absence of AHR changes the gut microenvironment composition to generate metabolites that impact T cell viability, such as bile salts and short chain fatty acids. Our study demonstrates a newly emerging role for AHR in mediating the interdependence between T lymphocytes and the microbiota, while simultaneously identifying new potential molecular targets for the treatment of MS and other autoimmune diseases.

INTRODUCTION

Multiple sclerosis (MS) is a chronic inflammatory demyelinating disease that affects 2.3 million people worldwide. MS etiology and pathology are linked to both genetic and environmental factors[1]. A primary environmental factor impacting disease pathology lies with the gut microbiota[2–5]. Indeed, the connection between the microbiota and the immune system in autoimmune disorders such as MS is well established[6]. The gut flora, composed of bacteria, fungi, and viruses, aids food digestion and maintains pathogen control, but recent studies have also highlighted the integral role of the microbiome in modulation of the homeostatic immune state both locally and systemically[7]. Gut dysbiosis is a hallmark of disease in both patients with MS and in experimental animal models of MS[2,8–11]. The dysbiotic microbiome can cause earlier onset and increase disease severity in mice [10]. Furthermore, germ-free mice lacking a microbiome are resistant to experimental autoimmune encephalomyelitis (EAE), an animal model of MS[12]. These findings spurred the examination of various bacterial supplements in mouse models of MS, and numerous potential candidates have been identified[6,13]. However, the mechanism(s) of action is not well understood. The importance of the microbiome in MS is well described, but a better understanding of the cross-talk between the immune system and the microbiome is required to translate these findings to the clinic. Further, discoveries in the context of MS provide a foundation for understanding environmental factors impacting many other autoimmune disorders.

The aryl hydrocarbon receptor (AHR) is a prime candidate for understanding the microbiome-immune interface. The AHR is a cytoplasmic receptor that, when activated, traffics to the nucleus to execute downstream transcriptional programing[14]. While canonical immune sensors have been primarily characterized based on their response to pathogenic material, the

AHR is a homeostatic regulator that is activated by a variety of nonpathogenic exogenous

ligands present in barrier tissues. These ligands include indoles[15], kynurenines[16], and other

small molecules[17]. Downstream immune effects are dependent on the cell type and ligand. The

AHR has also been directly tied to MS. AHR activation is the suspected mechanism by which

the novel MS therapeutic, Laquinimod, acts[18,19]. It is thought that AHR activation by

Laquinimod in antigen presenting cells reduces the ratio between inflammatory T cells and

regulatory T cells (Treg). Microbiome changes have been identified in AHR null mice and in

mice treated with a potent AHR agonist 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD), further

supporting the connection between AHR, the microbiome and MS [20–24]. While AHR activity

in response to microbiome derived metabolites is well described, a gap of knowledge remains

about the impact of AHR expression on the microbiota composition and function.

Here, we explore the role of AHR activity in CD4+ T cells using EAE. We have

demonstrated that deleting *Ahr* from CD4+ T cells increases recovery in EAE in a microbiome-

dependent manner. This improved recovery was the result of increased T cell apoptosis after

activation in the central nervous system (CNS). While AHR deficiency did not grossly alter the

composition of the microbiome, it significantly impacted the production of microbiome-

mediated metabolites. In particular, AHR deficiency in T cells led to an increase in bile acids

(BA) and a subset of short-chain fatty acids (SCFA) that ultimately impacted T cell viability. Our

study aimed to understand how a microbiota response element can act in the inverse in the

context of autoimmunity. This is the first demonstration, to our knowledge, of a role for AHR in

T lymphocytes as a regulator of the microbiome activity that ultimately influences the outcome

of CNS autoimmunity. Our discovery builds a foundation for further work that could lead to a

microbiome centric approach to dampen the overactive immune system in MS and related

autoimmune disorders.

RESULTS

Separately housed **CD4+ specific *Ahr* knockout mice recover from active EAE:**

Given the emerging role of the microbiome and the well accepted contribution of T cells

to MS pathology, AHR is ideally positioned to influence the pathology of CNS autoimmunity.

Here we explored the role of AHR in CD4+ T cells in EAE. For this study, we used the

previously validated *Cd4^{cre}Ahr$^{fl/fl}$* and littermate control *Ahr$^{fl/fl}$* mouse strains[25]. Cohoused

adult littermate mice were immunized with myelin oligodendrocyte glycoprotein (MOG)$^{35-55}$ to

induce active EAE. The cohoused mice showed no difference in EAE clinical score when *Ahr*

expression was removed from CD4+ cells (**Figure 1A, Supplement 1A**; Females **Supplement**

1B-C; Males).   Further, we observed no difference in myelin content in the spinal cord, as

revealed by Luxol Fast Blue staining at the chronic phase of disease (**Figure 1B-C**). To assess

the role of AHR driven microbiome changes in EAE, we separated mice by genotype at weaning

(3 weeks) and induced EAE between 8 and 16 weeks of age. In this condition, while the time of

onset and the peak clinical scores of EAE were not impacted, mice lacking *Ahr* expression in T

cells presented with a significant recovery when compared to wild type animals (Females;

Figure 1D, Supplement 1D). Furthermore, the change in recovery does not appear to be driven

by sex as it was conserved in both sexes (Males; **Supplement 1E-F**). Supporting the clinical

scores, the demyelination of the spinal cords was not different between the two groups at the

peak of disease (day 16); however, we noted a significant increase in myelin staining at the

chronic phase (day 31) in the *Cd4^{cre}Ahr$^{fl/fl}$* animals when compared to the control (**Figures 1E-**

G). Together, these data indicate that the lack of *Ahr* expression in CD4+ cells promotes EAE

recovery in a microbiome-dependent manner that can be reversed by cohousing.

Fig. 1. **Separately housed *Cd4^{cre}Ahr$^{fl/fl}$* recover from EAE with increased**

myelin staining at chronic phase. (**A**) Clinical score of *Cd4^{cre}Ahr$^{fl/fl}$* and *Ahr$^{fl/fl}$* mice

cohoused (Females; Representative plot includes n= 9 mice/group; total replicates of N=2

experiments). Spinal cords sections of *Cd4^{cre}Ahr$^{fl/fl}$* and *Ahr$^{fl/fl}$* were stained with Luxol

Fast Blue and Hematoxylin/Eosin stain a day 31 post EAE induction (**B**) Representative

images and (C) quantification of myelin stain. Images from four equally spaced spinal

cord levels were averaged for each mouse. (n=5 mice/group; Unpaired T test p=0.0706)

(**D**) Clinical score of *Cd4^{cre}Ahr$^{fl/fl}$* and *Ahr$^{fl/fl}$* littermate controls separated at weaning (3

weeks of age). (Females; Representative plot includes n=8-9 mice/group; total replicates

of N=2 experiments; Mann-Whitney U Test on total scores reported in legend [p=0.0096]

and on single days reported on plot) (**E**) Luxol Fast Blue with Hematoxylin/Eosin stain at

the peak stage of EAE (day 16) and at chronic phase (day 31). (**F**) Quantification of

myelin stain by Luxol Fast Blue alone in *Ahr$^{fl/fl}$* mice and (**G**) *Cd4^{cre}Ahr$^{fl/fl}$* mice. (n=7-8

mice/group; N=2 experiments; Unpaired T tests [p=0.8343, 0.0322]) Scale bars represent

400μm. Error bars represent standard error from the mean.

T cell differentiation and activity are unchanged in *Cd4^{cre}Ahr$^{fl/fl}$* mice:

CD4+ T cells are responsible for orchestrating EAE pathology, and defects in these cells

have been shown to impact disease outcomes[26]. Additionally, AHR agonists can modulate T

cell skewing in a ligand dependent manner *in vivo*[27]. We aimed to explore the role of AHR in

T cells differentiation and cytokines production. First, we examined T cell differentiation and

the production of cytokines in cohoused $Cd4^{cre}Ahr^{fl/fl}$ mice. We found that the deletion of AHR

in T cells did not impact their capacity for *in vitro* differentiation as determined by expression of

lineage transcription factors by flow cytometry (**Figure 2A-E**) or production of the signature

cytokine for each lineage by ELISA (**Figure 2F-I**). This was mirrored in separately housed

animals (**Supplement 2A-G**). Furthermore, there were no differences in the immune cell

composition between the knockout and control animals at baseline (**Figure 2J-M**). We next

measured several cytokines important to EAE induction in the spinal cords of mice at the peak of

disease. We expected to see significant reduction in the pathological cytokines in the separately

housed $Cd4^{cre}Ahr^{fl/fl}$ mice that ultimately recover, but surprisingly we found minimal differences

(**Figure 2N-S**). The only change was a small reduction in TNFα which may warrant further

investigation. These findings suggest that CD4 cell intrinsic AHR activity is not necessary for

normal T cell development and cytokines expression supporting published work indicating that

AHR activation of the antigen presenting cell may be more important to T cell subset generation

in vivo[28–30]. Further, these data reinforce the hypothesis that the T cell-intrinsic reduction of

AHR activity in $Cd4^{cre}Ahr^{fl/fl}$ mice is not responsible for EAE recovery described in separately

housed mice.

Fig. 2. *Ahr* knockout in CD4+ cells does not influence T cell differentiation.

(**A**) Naïve CD4+ T cells were isolated from cohoused $Cd4^{cre}Ahr^{fl/fl}$ and littermate control

animals and treated to promote differentiation. Flow cytometry gating strategy to measure

percent of cells expressing the transcription factors specific to each cell type is shown.

(**B**) Representative histogram and quantification of Tbet+ cells after *in vitro*

differentiation of naïve CD4+ cells to T_H1 cell type, (n=6 mice/group; N=2 experiments;

Unpaired T test [p=0.8715]) (**C**) GATA3 staining of T_H2 differentiated cells, (n=6

mice/group; N=2 experiments; Unpaired T test [p=0.9356]), (**D**) RORγt staining of T_H17

cells (n=8 mice/group; N=3 experiments; Unpaired T test [p=0.3989]) (**E**), and FoxP3

staining for T_{regs}.(n=8 mice/group; N=3 experiments; Unpaired T test [p=0.4078]). (**F**)

ELISA analysis of culture supernatant after 24 hours of anti-CD3 stimulation in $T_H1$cells

(n=9 mice/group; N=3 experiments; Unpaired T test [p=0.3924]) (**G**) T_H2 cells(n=8

mice/group; N=3 experiments; Unpaired T test [p=0.5917]) (**H**) T_H17 cells (n=7-8

mice/group; N=3 experiments; Unpaired T test [p=0.4013]) (**I**) or Treg cells. (n=8

mice/group; N=3 experiments; Unpaired T test [p=0.2278]) (**J**) Gating strategy for (**K**)

flow cytometry of single cell suspensions of Peyer's patches from cohoused $Cd4^{cre}Ahr^{fl/fl}$

and $Ahr^{fl/fl}$ mice. (Multiple T tests with Welch correction [p=0.070748, 0.081520,

0.325104, 0.080798]) (**L**) Analysis of cell compositions from mesenteric lymph nodes

(Multiple T tests with Welch correction [p=0.127038, 0.179526, 0.444670, 0.118472])

(**M**) and the inguinal lymph nodes from cohoused $Cd4^{cre}Ahr^{fl/fl}$ and $Ahr^{fl/fl}$ mice (Multiple

T tests with Welch correction [p=0.381964, 0.224164, 0.664161, 0.465348]) (**N-S**)

Luminex assay on whole spinal cord homogenate of separately housed $Cd4^{cre}Ahr^{fl/fl}$ and

$Ahr^{fl/fl}$ mice at the peak of disease (day 16 after immunization; n=8 mice/group; N=2

experiments; Unpaired T test) (N) [p=0.3087] (O) [p=0.4580] (P) [p=0.8313] (Q)

[p=0.4941] (R) [p=0.0215] (S) [p=0.9333] Error bars represent standard error from the

mean.

$Cd4^{cre}Ahr^{fl/fl}$ mice have fewer T cells in the spinal cord during EAE:

171   To determine whether EAE recovery in $Cd4^{cre}Ahr^{fl/fl}$ was linked to a difference in

172 immune cell number during pathology, we quantified CD45+ immune cells in the spinal cord

173 tissue using immunohistochemistry at the peak and chronic stages of EAE progression (**Figure**

174 **3A-B**). While the quantity of CD45+ cells was equivalent at the peak of the disease, we observed

175 a decrease in the number of CD45+ cells in the $Cd4^{cre}Ahr^{fl/fl}$ spinal cord compared to controls at

176 the chronic phase (**Figure 3B**). We next aimed to determine whether this reduction was a result

177 of changes in the number of infiltrating T cells or macrophages. To accomplish this, we

178 quantified expression of CD3, a pan-T cell marker, by immunohistochemistry. We saw no

179 difference in the CD3+ cells in the spinal cord at the peak of disease (**Figure 3C**); however, at

180 the chronic phase, significantly lower CD3+ T cell coverage was observed in the $Cd4^{cre}Ahr^{fl/fl}$

181 mice compared to the separately housed littermate controls (**Figure 3D**). Meanwhile, monocytes

182 and macrophages, the other primary effector cells in EAE, showed no differences at either peak

183 or chonic phase as shown by CD11b expression (**Figure 3E-F).** We confirmed this outcome

184 using flow cytometry on percoll-isolated immune cells from the spinal cord. At the peak of

185 disease, we again observed no differences in infiltrating T cells (CD4+ or CD8+), T cell subsets

186 (RORgt+, Tbet+ GATA3+ or FoxP3+) or macrophages (CD11b+) (**Figure 3G-H**). At the

187 chronic phase of disease, we found that the loss of T cells was limited to the CD4+ compartment

188 and not the CD8+ compartment. Of the classic CD4 helper cell subtypes, all trended down in

189 $Cd4^{cre}Ahr^{fl/fl}$ mice with no one subtype constituting a significant decrease, indicating a pan-CD4+

190 phenotype, not associated with a specific subtype of CD4+ T cells (**Figure 3F**). Taken together,

191 our data suggest that in separately housed mice, EAE recovery observed in $Cd4^{cre}Ahr^{fl/fl}$ mice is

192 not due to a difference in cytokine expression or T cell differentiation. Instead, recovery

193 correlates with an overall reduction in CD4+ T cell numbers at the chronic phase of disease and

194 suggests an impact for AHR and the microbiome on CD4+ T cell fitness.

196 **Fig. 3. Reduced CD3+ cells in the spinal cord of *Cd4^{cre}Ahr$^{fl/fl}$* mice correlate**

197 **with clinical signs.** (**A**) Spinal cords of separately housed *Cd4^{cre}Ahr$^{fl/fl}$* and *Ahr$^{fl/fl}$*

198 littermate controls were stained for total immune cells (CD45+) by

199 immunohistochemistry at four levels at the peak of disease (day 16 after immunization)

200 (Each dot represents a mouse n=7 mice/group; N=2 experiments; Two-way

201 ANOVA[p=0.5336]) (**B**) or at the chronic phase of disease (day 31 after immunization).

202 (Two-way ANOVA [p=0.0419]; Sidak's multiple comparisons test [p=0.9896, 0.9209,

203 0.7898, 0.1753]) (**C**) Spinal cords of separately housed *Cd4^{cre}Ahr$^{fl/fl}$* and *Ahr$^{fl/fl}$* littermate

204 controls were stained for T cells (CD3+) by immunohistochemistry at four levels at the

205 peak of disease (day 16 after immunization) (n=7 mice/group; N=2 experiments, Two-

206 way ANOVA [p=0.0830]) (**D**) or at chronic phase of disease (day 31 after

207 immunization). (n=7 mice/group; N=2 experiments, Two-way ANOVA [p=0.0054];

208 Sidak's multiple comparisons test [p=0.9578, 0.9029, 0.7095, 0.0070]) (**E**) Spinal cords

209 of separately housed *Cd4^{cre}Ahr$^{fl/fl}$* and *Ahr$^{fl/fl}$* littermate controls were stained for

210 macrophages (CD11b+) by immunohistochemistry at four levels at the peak of disease

211 (day 16 after immunization) (n=4-5 mice/group; N=2 experiments; Two-way ANOVA

212 [p=0.1271]) (**F**) or at chronic phase of disease (day 31 after immunization). (n=5

213 mice/group; N=2 experiments, Two-way ANOVA [p=0.2269]) (**G**) Gating strategy for

214 flow cytometry analysis of immune cells isolated from dissociated spinal cord tissue (**H**)

215 at peak of disease (day 16 after immunization) (n=5-6/group; Multiple Unpaired T Tests

with Benjamini and Yekutieli corrections [p values listed in supplement]). (**I**) or chronic

phase of disease (day 31 after immunization) (n=5 mice/group; Multiple Unpaired T

Tests with Benjamini and Yekutieli corrections [p values listed in supplement]). Scale

bars represent 400μm. Error bars represent standard error from the mean.

Apoptosis of *Ahr* deficient CD4 T cells is dependent on gut-derived small molecules:

We next investigated the mechanism behind the reduced T cell numbers in the spinal

cords of $Cd4^{cre}Ahr^{fl/fl}$ mice during chronic EAE. We performed a gene expression analysis on

whole spinal cord tissues isolated from two $Cd4^{cre}Ahr^{fl/fl}$ and two $Ahr^{fl/fl}$ animals at the peak of

EAE using a qPCR array composed of 384 genes associated with neuroinflammation. We found

that the expression of many genes involved in apoptosis were modulated in the spinal cord tissue

of $Cd4^{cre}Ahr^{fl/fl}$ mice; in particular we noted increased expression of *Fas, GrB, Casp3,* and *Casp9*

and decreased expression of the antiapoptotic transcript of *Xiap,* indicating an increase in

apoptosis in CD4 specific *Ahr*-deficient mice (**Supplement 3A-B**). To test if T cells lacking

AHR were undergoing apoptosis in EAE, we stained spinal cords sections for CD3 expression

by immunofluorescence and TUNEL assay (**Figure 4A**). While we could detect some apoptotic

T cells in the spinal cords of control mice subjected to EAE, CD3+TUNEL+ cell numbers were

significantly increased in the spinal cords of *Ahr* deficient animals (closed triangles; **Figure 4B**).

We confirmed this observation by quantifying cellular death of isolated immune cells from the

spinal cord at the peak of disease using flow cytometry. We observed a significant and specific

increase in the percentage of dead CD4+ T cells without change in the percentage of dead CD8+

and or total TCRB+ cells between genotypes (**Figure 4C**). To further understand the

mechanisms behind these observations, we focused examination on the primary pathogenic

subset of CD4+ cells in EAE, the type 17 T helper cell (T_H17)[31]. To obtain T_H17 cells we

isolated naïve CD4+ cells from the lymph nodes and spleen, then cultured the cells in T_H17

differentiation conditions. Similar to the *in vivo* observations, *Cd4^{cre}Ahr$^{fl/fl}$* T_H17 cells have a

significant increase in apoptosis after mild *in vitro* stimulation by anti-CD3 as measured by

Annexin V staining compared to littermate controls (**Figure 4D**). Increased T_H17 cell death was

only observed in cells isolated from *Cd4^{cre}Ahr$^{fl/fl}$* animals separated at weaning, as T cells

isolated from cohoused *Cd4^{cre}Ahr$^{fl/fl}$* mice did not display elevated apoptosis (**Figure 4E**). Taken

together, these results support the hypothesis that gut microenvironment may be implicated in the

Ahr-dependent T cell fitness phenotype. To directly test this hypothesis, we isolated metabolites

(<3kDa) from cecums of separately housed *Cd4^{cre}Ahr$^{fl/fl}$* and *Ahr$^{fl/fl}$* mice, and added them to *in*

vitro skewed T_H17 cells prepared from C57BL6/J mice during the 24 hour stimulation period.

We next analyzed apoptosis by flow cytometry. T_H17 cells exposed to metabolites derived from

the gut of *Cd4^{cre}Ahr$^{fl/fl}$* mice had elevated apoptosis compared to cells exposed to metabolites

from *Ahr$^{fl/fl}$* microbiome (**Figure 4F**). Collectively, these data demonstrate that exposure to the

microbial metabolites from *Cd4^{cre}Ahr$^{fl/fl}$* mice is sufficient to induce early T cell apoptosis which

likely contributes to recovery in EAE.

Fig 4. T cell apoptosis is increased in animals who recover from EAE. (**A**)

Representative image of spinal cords stained with TUNEL assay and T cell marker at the

peak of disease (CD3). Closed triangles indicate TUNEL+CD3+ cells and arrows

indicate TUNEL+CD3- cells. Scale bar = 400µm. (**B**) Quantification of the number of

TUNEL+ T cells (2-6 lesion sites averaged per animal; n=5-6 mice; N=2 EAE

experiments; Unpaired T test [p=0.0357]). (**C**) T cells isolated from the spinal cord at the

peak of disease stained with viability dye and measured by flow cytometry. Cells gated

on singlets, CD45+, gate shown on x-axis, then dead cells (n=5 mice/group; Two-way

ANOVA [p=0.0192] with Sidak's Multiple Comparison Tests [p=0.0465, 0.2862,

0.6800]). (**D**) T cells skewed to T_H17 *in vitro* from separately housed $Cd4^{cre}Ahr^{fl/fl}$ mice

stained for apoptotic marker Annexin V 24 hours after stimulation with anti-CD3

antibody. (n=6 mice/group; N=2 experiments; Unpaired T test [p=0.0133]). (**E**) T_H17

cells from cohoused $Cd4^{cre}Ahr^{fl/fl}$ and $Ahr^{fl/fl}$ mice show no differences in the number of

Annexin V positive cells 24 hours after stimulation with anti-CD3 antibody. (n=5-6

mice/group; N=2 experiments; Unpaired T test [p=0.5851]). (**F**) *In vitro* differentiated

T_H17 cells from C57BL6/J mice were exposed to the <3kDa fraction of cecal contents

from $Ahr^{fl/fl}$ and $Cd4^{cre}Ahr^{fl/fl}$ mice for 24 hours with anti-CD3 stimulation. (n=6-9

mice/group; N=3 experiments; Mixed-effects analysis, with Geisser-Greenhouse

correction [p=0.0013]; Tukey's post hoc analysis [Ctr v Fl p=0.7729; Ctr v Cre p=0.0333,

Fl v Cre p<0.0001]). Error bars represent standard error from the mean.

Metabolomic differences in the gut microenvironment are evident in separately housed

***Cd4^{cre}Ahr^{fl/fl}* mice:**

To identify the impact of CD4+ cell AHR loss on the gut microenvironment, we began

by performing 16S sequencing of DNA isolated from fecal pellets and cecal contents of $Ahr^{fl/fl}$ or

$Cd4^{cre}Ahr^{fl/fl}$ mice. Neither cohoused nor separately housed animals saw differences in fecal or

cecal microbial signatures (**Supplement 4A-D**). Given the marked effect of the metabolites

prepared from the cecums of $Cd4^{cre}Ahr^{fl/fl}$ and $Ahr^{fl/fl}$ on T cell apoptosis (**Figure 4F**), we next

conducted untargeted metabolomics by LC-MS on these preparations (<3kDa). Two-dimensional

clustering of the annotated products revealed distinct clustering of genotypes, indicating different

chemical profiles (**Figure 5A**). This difference occurred despite there being no significant

difference in the 16S microbial signatures of the cecal contents suggesting that AHR activity in T

cells modify microbial metabolism without altering the microbe composition to a degree that can

be captured by 16S sequencing (**Supplement 4E-F**). Impact analysis of biological pathways

identified primary bile acid biosynthesis as the most significantly upregulated pathway with six

primary (cholic, taurocholic) or secondary (glycocholic, sulfocholic, dehydrocholic, lawsonic)

bile acids significantly increased in the cecal microenvironment of $Cd4^{cre}Ahr^{fl/fl}$ mice (**Figure 5B,**

C). Bile acids are dysregulated systemically in patients with MS and adding back primary bile

acid is sufficient to reduce the severity of EAE[32]. Bile acid biosynthesis is known to be

dependent on the gut microbiota composition[33,34]. Of the changed bile acids, taurocholic acid

was the most highly upregulated with over a 10-fold increase in $Cd4^{cre}Ahr^{fl/fl}$ compared to control

mice (**Figure 5C**). Increased bile acid accumulation in these mice may be due to the

downregulation of bile acid transporters in the small intestine of the $Cd4^{cre}Ahr^{fl/fl}$ mice (**Figure**

5D). In addition to changes in bile acids, alanine, aspartate, and glutamate metabolism were also

upregulated, whereas purine metabolism and tryptophan metabolism were downregulated in

$Cd4^{cre}Ahr^{fl/fl}$ mice. As short chain fatty acids (SCFAs) have been reported to impact the viability

of inflammatory T cell and cannot be captured through untargeted metabolomics, we next

performed targeted SCFA metabolomics [35,36]. Isovaleric acid was the only SCFA with a

significantly higher abundance in the $Cd4^{cre}Ahr^{fl/fl}$ cecums compared to controls (**Figure 5E**).

306   **Fig 5. Elevated bile salts and SCFA detected in the cecum of AHR deficient**

307   **mice.** (**A**) Partial least-squares plot of LC-MS untargeted metabolomics of the <3kDa

fraction of the cecal contents from $Cd4^{cre}Ahr^{fl/fl}$ and $Ahr^{fl/fl}$ mice. (**B**) Metaboanalyst

mouse KEGG pathways significantly changed by genotype. (**C**) MetaMapp network

analysis visualizing the chemically or biologically related significantly changed

molecules. (**D**) qPCR of bile acid transporters (*mrp3, ostb*) and bile acid response

elements (*ezh2, fxr*) in bulk tissue of the duodenum and jejunum of separately housed

animals illustrated as relative quantity (R.Q.) from $Ahr^{fl/fl}$ tissue . (n=6 mice/group; N=2

experiments; Multiple T Tests with Welch correction [duodenum p=0.0104, 0.3591,

0.0071, 0.3828; jejunum p=0.2230, 0.0623, 0.1487, 0.1004]). (**E**) Quantitative targeted

metabolomics for short chain fatty acids show a significant increase in isovaleric acid,

and a trending increase in valeric and hexanoic acid in $Cd4^{cre}Ahr^{fl/fl}$ mice. (n=6

mice/group; N=1 experiment; Unpaired T tests [p=0.0427, 0.1934, 0.1083]) Error bars

represent standard error from the mean.

Taurocholic acid induces T_H17 cell apoptosis and protects from EAE:

The chemical composition of the cecal microenvironment between separately housed

$Cd4^{cre}Ahr^{fl/fl}$ and $Ahr^{fl/fl}$ animals is significantly changed. We next aimed to determine what

functional differences arose from the chemical differences both *in vitro* and *in vivo*. We first

examined the most highly increased bile acid in $Cd4^{cre}Ahr^{fl/fl}$ mice, taurocholic acid. When

administered to C57BL6/J derived T_H17 cells *in vitro,* taurocholic acid was sufficient to drive

apoptosis similarly to cecal content isolates (**Figure 6A-B**). We also examined isovaleric acid,

the short chain fatty acid, elevated in the absence of AHR. Similarly to taurocholic acid,

isovaleric acid could induce apoptosis of *in vitro* generated T_H17 cells (**Figure 6C**). Similar

results were obtained with hexanoic acid, which was also trending higher in $Cd4^{cre}Ahr^{fl/fl}$ mice

(**Figure 6C**). Collectively, our data suggest that the combined effects of the cecal milieu of

Cd4^{cre}Ahr$^{fl/fl}$ mice could induce changes in T cell fitness, ultimately resulting in recovery from

paralysis in EAE. As such, we aimed to determine whether this held true *in vivo* by conducting a

fecal microenvironment transfer to C57BL/6 wildtype mice.   The mice were given a cocktail of

oral antibiotics for two weeks followed by a series of three oral gavages of the flushed contents

of the intestinal tract from separately housed *Cd4^{cre}Ahr$^{fl/fl}$* and *Ahr$^{fl/fl}$* animals. During this time

the animals were also cohoused with an untreated member of the respective genotype to promote

lasting engraftment of the microbiome (**Figure 6D**). When these mice underwent EAE, the

wildtype mice given FMT from *Cd4^{cre}Ahr$^{fl/fl}$* animals had a similar, but less pronounced

reduction in severity as the donor mice (**Figure 6E-F**).   Because the FMT resulted in an

intermediate phenotype, we next aimed to determine whether an individual component of the gut

microenvironment could act as a therapeutic for EAE. As taurocholic acid was the most efficient

driver of apoptosis *in vitro* (**Figure 6B**) we chose this as our prime candidate. We administered

taurocholic acid orally for seven days after EAE induction (**Figure 6G**). Compared to mice given

saline, those receiving taurocholic acid had decreased peak EAE scores (**Figure 6H**) and delayed

onset (**Figure 6I**). These data are in support of recent work showing that bile acid receptor

agonists and bile acid cocktails can reduce T cell activation and suppress EAE [37,38].

347   **Fig 6. Gut Microenvironment and bile salts can drive EAE recovery and T**

348   **cell apoptosis** (**A**) Schematic representation of experiments in B and C. *In vitro* derived

349   T$_H$17 cells from C57/BL6/J mice were treated with a metabolite in addition to mild anti-

350   CD3 stimulation for 24 hours. (**B**) Taurocholic acid (10μM and 100μM) was

351   administered during anti-CD3 stimulation and apoptosis of T$_H$17 cells isolated from

352   C57BL6/J mice was measured using Annexin V staining by flow cytometry. (n=5-6

353   mice/group; N=2 experiments; Mixed effects analysis with Geisser-Greenhouse

correction [p=0.0070]; Dunnett's multiple comparison test [Veh vs 10μM p=0.8178, Veh

vs 100μM p=0.0233]) (**C**) Administration of isovaleric acid and hexanoic acid (short

chain fatty acids that are increased in mice that recover) are sufficient to drive apoptosis

in T_H17 cells isolated from C57BL6/J mice. (n=8 mice/group; N=2 experiments; RM

one-way ANOVA with the Geisser-Greenhouse correction[p=0.0399] Dunnett's multiple

comparisons test [Veh vs 1mM Hexanoic p=0.0764, Veh vs 10mM Hexanoic p=0.0352,

Veh vs 1mM Isovaleric p=0.0026, Veh vs 10mM Isovaleric p=0.0254]). (**D**) Schematic

depicting approach to FMT. (**E**) EAE clinical scores of C57BL6/J mice receiving FMT

from separately housed $Ahr^{fl/fl}$ or $Cd4^{cre}Ahr^{fl/fl}$ mice. (Females; Representative plot

includes n=11 mice/group; total replicates of N=2 experiments; Mann-Whitney U Test on

total scores reported in legend [p=0.0096]. (**F**) Incidence of clinical sign development in

mice treated with FMT. (**G**) Schematic depicting approach to oral supplement of 12.5 mg

taurocholic acid/day in C57BL6/J mice immunized for EAE. (**H**) EAE clinical scores of

C57BL6/J mice receiving 7 days of oral taurocholic acid from separately housed $Ahr^{fl/fl}$ or

$Cd4^{cre}Ahr^{fl/fl}$ mice. (Females; n=12 mice/group; total replicates of N=1 experiment;

Mann-Whitney U Test on single days reported on plot). (**I**) Incidence of clinical sign

development in mice treated with taurocholic acid. Error bars represent standard error

from the mean.

DISCUSSION

Here we demonstrate that the deletion of the aryl hydrocarbon receptor (AHR) in CD4+

cells can promote recovery in chronic EAE while having no impact on the onset nor the initial

magnitude of the disease. We further show that this phenotype is dependent on the microbiome,

as cohousing littermate $Cd4^{cre}Ahr^{fl/fl}$ and $Ahr^{fl/fl}$ mice abrogates recovery in EAE. These data are

in support of previous work in similar models has shown no difference in EAE recovery after T

cell-specific *Ahr* deletion [39]. At the mechanistic level, our data suggest that specific

metabolites elevated in the cecum of $Cd4^{cre}Ahr^{fl/fl}$ mice reduce T cell fitness and viability. In

particular, the cecal environment of $Cd4^{cre}Ahr^{fl/fl}$ animals is characterized by high levels of

isovaleric acid and taurocholic acid, which induce T cell apoptosis *in vitro,* suggesting a potential

mechanism for EAE recovery.

AHR activity can regulate autoimmunity via natural killer (NK) cells[40],

macrophages[41], dendritic cells[42], and T cells[27,30,43]-all contributors to EAE

pathogenesis. AHR activation is a known regulator of T cell differentiation and inflammatory to

regulatory T cell ratios. Importantly, T cell activity is dependent on the specific ligand activation

with some promoting a pro-inflammatory environment, and others having the opposite

effect[27]. It has been postulated that T cell intrinsic AHR activity is necessary for

differentiation but several groups have shown that AHR antagonists do not have a cell intrinsic

effect on T cell differentiation leaving remaining questions in the field[44]. Here we

demonstrate that T cell-specific AHR expression is not necessary for the differentiation of T cells

in vitro supporting the hypothesis that the AHR activity in antigen presenting cell is the

necessary factor influencing T cell differentiation *in vivo*[28–30]. Beyond T cells, AHR

activation in glial cells regulates the pathogenic effects of microglia and astrocytes in EAE via

changed cytokine production [45,46]. AHR's ubiquitous expression in barrier tissues coupled

with its complex ligand binding profile warrants further study of AHR function in autoimmunity.

We have identified that the changes of the luminal microenvironment of the intestine resulting

from *Ahr* knockout in CD4+ cells lead to decreased post-activation lifespan in T cells; however,

the impact of these changes on other cell types and other models of autoimmunity have yet to be

determined. Though we aimed to target T cells using the CD4 cre line, we acknowledge that

other CD4 expressing cells including some gut macrophages may also be contributing to the

observed phenotype.

Bile acids, SCFAs, and other small molecules we identified can move into circulation

and some can gain entry into the central nervous system by crossing the blood brain barrier[47].

There are likely systemic effects that have yet to be described. For example, our data suggest

that significant remyelination is taking place in the chronic phase of EAE in CD4 specific *Ahr*

knockout animals. Given the described impact of the microbiome and SCFA on myelination

[48,49], it would be critical to test if the metabolic signature generated by the lack of *Ahr* in T

cells can influence oligodendrocyte generation.

The data presented here support a bidirectional communication between T cells and the

microbiome via AHR. However, it remains to be understood how T cells can modulate the

metabolic signature of the cecum. We hypothesize that there are many mechanisms that could be

mediating the change. For example, we show that the expression on bile acid transporters are

reduced in the intestines of mice with *Ahr* deletion. This could lead to heightened levels of bile

acids which have an impact on microbial viability, metabolism, and community structure.

Additionally, it is possible that *Ahr* deletion in T cells may lead to modulation of IgA secretion

by B cells[50,51]. IgA is the primary mechanism by which the adaptive immune system can

directly modulate mucosal microbial communities[50]. An alternative hypothesis is that immune

cell activity is modulating antimicrobial peptide release indirectly through communication with

Paneth cells[52,53]. Most work on these systems thus far has focused on models of severe

microbiome dysbiosis. Less is known about the subtler changes in the microbiome subject to

homeostatic regulation. Therefore, the mechanism by which the adaptive immune system can

interface with the microbiota is of great interest for understanding the basis of autoimmune

disorders.

Ahr knockout in T cells results in a change in both primary and conjugated bile acids in

the cecum. Bile acids are produced by the host for digestion and are normally reabsorbed by

transporters (primary) or passively (primary conjugated) in the ileum while secondary bile acids

are not as easily absorbed [54]. There is an increase in conjugated bile acids in *Ahr* knockout

mice, but lower levels of secondary bile acids. This is the inverse of patients with MS indicating

that there may be a causative connection in both patients and mice with EAE[32]. Further studies

are needed to understand which step of this pathway: production, conversion or absorption are

influenced by the gut flora and/or the T cells. Our results support the existing work in synthetic

bile acid receptor agonists showing that T cell fitness and activity can be mediated by bile acid

receptor activation[37,38]. As previous groups have shown that T cells traffic to the gut before

the CNS during EAE, we hypothesize that exposure to these factors during priming in the lymph

node may lead to the early apoptosis after trafficking to the CNS[55]. This evidence that bile

acids can have a direct influence on T cell function warrants further study in the context of

autoimmune diseases.

Microbiome-modulated immune responses become particularly important in patients with

chronic and relapsing autoimmune conditions like MS[56–60]. We have described a novel

function for the environmental sensor AHR in T cells as a regulator of the gut microenvironment.

As we further understand the complexities of harnessing the gut microbiome, we predict tools for

controlling host-derived microbe modulation will be a new frontier for therapeutics. Our work is

an important step in the discovery of a targetable endogenous modulator of the microbiome

opens a potential new therapeutic avenue for MS patients.

MATERIALS AND METHODS

Experimental Design: The objective of these studies was to understand the role of CD4 specific AHR activity in EAE. Immunophenotyping and characterization of the mice occurred in cohoused mice to determine the purely genetic differences. Once no major changes were discovered in the *Ahr* knockout mice, we began to separately house them based on genotype. This allowed us to examine the microbiome-dependent features of autoimmunity and the development of further hypotheses. Sample sizes were determined by power analysis. The final desired n was divided by two and included for each experiment. In this way, we have two independent experiments resulting in appropriate power. Except where otherwise stated, all experiments were conducted twice independently. Data collection for all EAE experiments ended at 30 days- the time point at which we have defined as chronic stage. All experiments were conducted on age- and sex- matched animals. Histological analysis was conducted by a blinded investigator. Clinical scores are reported as the average of two blinded investigators to account for interrater variability. Statistical outliers identified using the ROUT method (Q=1%) are not included in analysis or figures.

Mice: AHR$^{tm3.1Bra}$/J(Ahr^{fx}) (#006203), B6.Cg-Tg(Cd4-cre)1Cwi/BfluJ (CD4Cre) (#022071), and C57BL/6J (#000664) mice were purchased from Jackson Laboratories. Mice were bred in-house with a 12-hour light/dark schedule. Cohoused mice were housed with their littermates of the same sex from birth. Separately housed mice were separated at weaning (3 weeks) based on presence of Cre. Litters born within 4 days of each other were housed together to achieve 2-5 mice per cage. All procedures were approved by the University of Virginia ACUC (protocol #1918).

Experimental autoimmune encephalomyelitis: EAE was induced in both male and

female mice between 8 and 12 weeks of age. MOG$_{35-55}$ peptide (100μg, CSBio, CS0681) was

emulsified in complete Freund's adjuvant containing *Mycobacterium tuberculosis* (1mg/mL,

Sigma, F5881; M. tuberculosis (BD 231141) added to final concentration of 4mg/mL) and was

injected subcutaneously (100 μL volume) at the base of the tail. Pertussis toxin (200 ng, List

Biologicals, 180) was administered i.p. on the day of and 1 day after MOG immunization. Mice

were scored daily by two blinded evaluators using the following scale: 0-no clinical disease, 1-

limp tail, 2-hindlimb incoordination, 3-hindlimb weakness, 4-hindlimb paralysis, 5-moribund. As

is standard practice, animals who did not develop any signs of EAE were excluded from

analysis. Total incidence including excluded animals is reported in the supplement.

Immunohistochemistry and Fluorescent Microscopy: Spinal cords were prepared as

previously described[61]. Slides were deparaffinized using xylenes and an ethanol gradient. IHC

antibodies used were anti-CD3 (Dako; A0452) at 1:200 dilution and anti-CD45 (BD Biosciences;

550539) at 1:80 dilution. Adjacent sections were stained with Luxol Fast Blue and Hemtoxylin

and Eosin for demyelinating plaques. In short, slides were deparaffinized using xylenes and an

ethanol gradient. Slides were stained with Luxol Fast Blue at 56 ℃ overnight. Slides were

rinsed with distilled water and then 0.05% Lithium carbonate until sharp contrast between white

matter and grey matter was evident. Slides were imaged using an EVOS microscope. All

images were analyzed with ImageJ by setting a threshold in the negative control and evaluating

percent positive area within the white matter of the spinal cord. Luxol Fast Blue alone was used

for quantification of myelin coverage while Luxol Fast Blue with H&E images were used in the

figure to highlight lesions. Histological analysis was performed by a blinded evaluator.

For TUNEL staining, cryosections (30μm) were stained with anti-CD3 (eBioscience; 50-

0032-82) or isotype control (Fisher Scientific; 14-0032-82) as previously described[61]. A

positive control slide was treated with DNAse (3000U/mL). After staining, sections were fixed

with paraformaldehyde for 15 minutes at room temperature. Sections were washed two times in

PBS followed by proteinase K (200ug/mL in PBS) treatment for 10 minutes at room

temperature. Sections were washed twice with PBS and incubated with TUNEL reaction

mixture (Roche; 12156792910) for 60 minutes at 37 ºC in a humidified chamber. Sections were

washed three times in PBS and mounted using Prolong Gold Anti-Fade Reagent (Life

Technologies). Slides were imaged with a Leica TCS SP8 confocal microscope. All images

were analyzed with ImageJ. Histological analysis was performed by a blinded evaluator.

RNA Extraction and Quantitative PCR: Whole spinal cords or 0.5 cm sections of gut

tissue were homogenized using a dounce homogenizer and RNA was extracted using the Bioline

Isolate II RNA mini kit as per manufacture's protocol (BIO-52073). RNA was quantified with a

Biotek Epoch Microplate Spectrophotometer. Normalized RNA was reverse transcribed to

cDNA with the Bioline SensiFast cDNA Synthesis Kit (BIO-65054). For the qPCR array, two

spinal cords per group were pooled and analyzed as described by the manufacturer

(Demyelinating Diseases Tier 1-4 M384; 10038953). Primers for targeted qPCR are listed in the

supplemental materials. Results were analyzed with the relative quantity (ΔΔCq) method.

Isolation and Preparation T cells: Mesenteric, inguinal, axial, and brachial lymph

nodes plus the spleen were dissected from the mice after CO_2 euthanasia. They were then

dissociated through sterile 70 μm filters and lysed with ACK lysis buffer (Quality Biological,

118-156-101). Naïve CD4+ T cells were then sorted using an EasySep Mouse CD4+ T cell

Isolation Kit (Stem Cell Technologies, #19852). The cells were incubated at $1x10^6$ cells/mL in

skew media in anti-CD28 (InVivoMab, BE0015-5) and anti-CD3 (InVivoMab, BE0001-1)

coated plates as previously described[62]. T_H1 and T_H2 media contained RPMI (Gibco, 11875-

093) supplemented with 10% FBS (Optima, S12450H), 0.5% Pen/Strep (Gibco, 15240-062),

1mM sodium pyruvate (Gibco, 11360-070), 2mM L-glutamine (Gibco, 25030081), 10mM

HEPES (Gibco, 15603-080), 1:100 NEAA (Gibco, 11140-050), and 50uM B-mercaptoethanol

(Fisher Scientific, O3446I-100). Treg and T_H17 media contained IMDM (Gibco, 12440-053)

using the same supplements. T_H1 skew media contained IL-2 (100U/mL; Tecin, 23-6019), IL-12

(10ng/mL; PeproTech, 210-12), and anti-IL-4 (10ug/mL; InVivoMab, BE0045). T_H2 skew media

contained IL-2 (100U/mL; Tecin, 23-6019), IL-4 (10ng/mL; InVivoMab, BE0045), and anti-

IFNγ (10ug/mL; InVivoMab BE0054).   Treg skew media contained TGF-B (5ng/mL;

BioLegend, 763102) and anti-CD28 (2ug/mL; InVivoMab, BE0015-5). T_H17 skew media

contained IL-6 (20ng/mL; BioLegend, 575704), IL-23 (10ng/mL; Invitrogen, 14-8231), TGF-B

(0.3ng/mL; BioLegend, 763102), anti-IL-4 (10ug/mL; InVivoMab, BE0045), and anti-IFNγ

(10ug/mL; InVivoMab BE0054). T_H1 and T_H2 cell cultures were expanded on day 3 after plating

by transferring cells to larger plates and adding 3x volume of RPMI supplemented media

containing IL-2 (100U/mL). On day 5 skewing was complete and cells were washed and plated

for downstream assays. Treg and T_H17 cell cultures were not expanded but simply washed and

plated for downstream assays on day 4 after plating.

Apoptosis and Cytokines Production Assay: Skewed T cells were plated with

respective treatments at $1x10^6$ cells/mL in 48 well plates coated with anti-CD3. After 24 hours,

the supernatant was collected for ELISA assay or cells were collected for Annexin V staining. In

short, cells were stained with Ghost Live Dead (Tonbo Biosciences; 13-0870), Annexin V

(Fisher Scientific; BDB556420) per manufacturer's instructions (Annexin V, fixed using the

FoxP3 permeabilization, fixation kit (eBioscience; 00-5523-00), and stained with RORgt PE.

Results were gated on singlets, RORgt+, and Annexin V+, L/D negative.

Enzyme-linked immunosorbent Assay: ELISA for IFNγ, IL-4, IL-17, and IL-10 were

performed as previously described[63]. Antibodies used were as follows: anti-IFNγ (BioLegend;

517902), biotin labeled anti-INFγ (BioLegend; 505704), anti-IL-4 (Thermo Fisher; 14-7041-85),

biotin labeled anti-IL-4 (Thermo Fisher; 13-7042-85), anti-IL-17 (eBioscience; 14-7175-85),

biotin labeled anti-IL-17(eBioscience; 13-7177-85), anti-IL-10 (BioLegend; 505002), biotin

labeled anti-IL-10 (BioLegend; 505004).

Isolation of Cecal Contents: For both metabolomics and *in vitro* analysis, the cecal

contents were resuspended in 1mL of IMDM by vortexing. Preparations were then sequentially

processed using first syringe filters to remove insoluble materials (70 μm, 40 μm, .22 μm) and

then protein concentrators (50 and 3 kDa from Thermo Fischer scientific) to exclude

macromolecules with a molecular weight > 3kDa. Preparations were used immediately, or flash

frozen at -80C for up to 3 weeks.

Spinal Cord Dissociation Spinal cords were collected in 2.5 mL ice cold HBSS (Gibco,

14025134). Equal amounts of 4mg/mL collagenase 4 (Worthington; LS002139) with 50U/mL

DNase (Worthington, LS002139) were added and samples were shaken three times at 37ºC for

15 minutes, triturating in between. Spinal cords were strained through a 70mm filter into 10mL

DMEM (Gibco, 11965-092) with 10% FBS (R&D Systems; S12450H). The samples were then

centrifuged at 260 g for 8 minutes. The pellet was resuspended in a 15mL tube in 12 mL of

Percoll (GE Healthcare, 17-0891-01). The sample was then centrifuged at 650g for 25 min

without brake. The myelin layer and supernatant were aspirated and the pellet was resuspended

in 2mL of FACS buffer.

Spinal Cord Luminex Mice were perfused with PBS containing heparin (5U/mL;

Sigma; H-3125) solution. Spinal cords were dissected and flash frozen. Frozen spinal cords

were homogenized in 1mL of PBS with protease inhibitor (MedChem Express; HY-K0010)

using a Dounce homogenizer. Triton X-100 was added to final concentration of 0.2% and

samples were vortexed. Samples were centrifuged at 10,000g for 5 minutes at 4 °C and 200 μL

was acquired from the top layer to be flash frozen for later analysis. Mouse T_H17 Luminex was

run per manufacturer's instructions on a MagPix by the University of Virginia Flow Cytometry

Core , RRID: SCR_017829.

Microbiome Transfer Female C57BL6/J mice were given drinking water containing

1g/L Ampicillin (Sigma-Aldrich; A8351-25G), 1g/L Neomycin (Sigma-Aldrich; N6386-25G),

1g/L Metronidazol (Sigma-Aldrich; M1547-25G), 500mg/mL Vancomycin (Sigma-Aldrich;

V1130-5G), and 8mg/mL Splenda for two weeks. Mice were then orally gavaged with fresh

filtered (40μm) cecal contents from pooled donor animals on day 0, day 2, and day 4 after

removal from antibiotics. On the day of antibiotics withdrawal, a naïve microbiome donor

animal was added to the cage and remained for the duration of the experiment. EAE was induced

2 weeks post reconstitution.

Bile Acid Supplement Female and male C57BL6/J mice were immunized with MOG as

described above. Daily gavage of either 100μL of 125 mg/mL taurocholic acid (in saline) or

100μL of saline started on day 4 and continued through day 10. Mice were then monitored and

scored as described earlier.

Flow Cytometry Single cell suspensions were incubated with Fc Block CD16/32

(Invitrogen, 14-0161-85). The cells were then incubated with a 1:200 dilution of antibody for

surface stains and A Live/Dead Ghost Dye Violet 510 (Tonbo Biosciences; 13-0870). Surface

antibodies: 488-conjugated CD8 (Invitrogen; 53-008182), APCe780-conjugated TCRβ

(Invitrogen; 47-5961-82), e450-conjugated CD4 (Invitrogen; 48-0042-820), 488 conjugated-

CD11b (Invitrogen; 53-0112-82), and APC-conjugated CD45.2 (Biolegend; 109813). For

transcription factor staining, the eBioscience FoxP3/Transcription Factor Staining Kit (00-5523-

00) was used per manufacturer's instructions with intracellular antibodies: PE-conjugated

RORγT (Invitrogen; 12-6981-82), PE-Cy7- conjugated FoxP3 (Invitrogen; 25-5773-82), PE-

conjugated GATA3 (Invitrogen; 12-9966-42), PE-Cy7-conjugated Tbet (Biolegend; 644824).

OneComp eBeads (Thermo Fisher Scientific, 01-111-42) were used for color controls. Flow

cytometry was performed using a Beckman Coulter Gallios flow cytometer and data were

analyzed with FlowJo software v10.7.1.

LC-MS Untargeted Metabolomics was performed by Creative Proteomics. Cecal samples

(<3kDa) were prepared and concentrated as before as described before. Samples were thawed

and lyophilized, 200μL of 80% methanol was added. Samples were vortexed and then sonicated

for 30 minutes at 4 ℃. Samples were kept at -20 for 1 h, vortexed for 30s and kept at 4 ℃ for

15min until they were centrifuged at 12000 rpm at 4 ℃ for 15 min. 100μL of the supernatant

and 2.5μL of DL-o-Chlorophenylalanine were used for LC-MS analysis. Separation is performed

by Ultimate 3000 LC combined with Q Exactive MS (Thermo) and screened with ESI-MS. The

LC system is comprised of ACQUITY UPLC HSS T3 (100×2.1mm×1.8 μm) with Ultimate 3000

LC. The mobile phase is composed of solvent A (0.05% formic acid water) and solvent B

(acetonitrile) with a gradient elution (0-1.0 min, 5%B; 1.0-12.0 min, 5%-95%B; 12.0-13.5 min,

95%B; 13.5-13.6 min, 95%-5%B; 13.6-16.0 min, 5%B). The flow rate of the mobile phase is 0.3

mL·min-1. The column temperature is maintained at 40°C, and the sample manager temperature

is set at 4°C. Mass spectrometry parameters in ESI+ and ESI- mode are listed as follows:

ESI+: Heater Temp 300 °C; Sheath Gas Flow rate, 45arb; Aux Gas Flow Rate, 15arb; Sweep

Gas Flow Rate, 1arb; spray voltage, 3.0KV; Capillary Temp, 350 °C; S-Lens RF Level, 30%.

ESI-: Heater Temp 300 °C, Sheath Gas Flow rate, 45arb; Aux Gas Flow Rate, 15arb; Sweep Gas

Flow Rate, 1arb; spray voltage, 3.2KV; Capillary Temp,350 °C; S-Lens RF Level,60%.

LC-MS Data Processing LC-MS data was processed using R studio. Analysis was

conducted using POMA Shiny version 1.4.0[64]. In short, we removed features with more than

20% missing values, normalized with log scaling and then completed sparse partial least squares

discriminate analysis (sPLS-DA). Top ranked products were utilized in MetaMapp[65,66]

analysis to produce a web of chemically and biologically related compounds. Finally, the top

ranked products were also inputted into Metaboanalyst to complete pathway enrichment and

determine mouse KEGG terms.

16S Sequencing Data Processing 16S Sequencing was conducted at Microbiome

Insights (Cohoused Feces), ZymoBIOMICS (Separately Housed Cecum), and at the University

of Virginia Genomics Core (Separately housed Feces). Initial sequence data was analyzed using

the latest version of Quantitative Insights Into Microbial Ecology 2 (Qiime2 v2021.11)[67].

Demultiplexed paired-end sequence reads were preprocessed using DADA2[68]. The first 20

base pairs were trimmed from forward and reverse reads before they were merged to remove

adaptors. Taxonomy was assigned to Amplicon Sequence Variants (ASVs) using a Naïve Bayes

classifier trained on full length 16S sequences from the latest Greengenes16S database (13_8)

clustered at 99% sequence similarity. Samples were rarified before core diversity analysis.

Rarefied sampling depth was 16000 for cecal contents, 30000 for separately housed fecal

contents, and 10,411 for cohoused fecal contents. Core diversity metrics were analyzed,

including number of ASVs. Nonmetric multidimensional scaling was performed in RStudio

using the phyloseq package[69].

Statistical Analysis Unless otherwise noted, all statistical tests were run on Graph Pad

Prism 9 Version 9.1.0. Statistical tests and p-values for each comparison are listed in the figure

legends or in the supplementary report. In the figure legends, n denotes the total number of

animals displayed in the figure used in analysis. In the figure legends, N denotes the total

number of replicates that were completed for the given experiment. Testing level of $\alpha=0.05$ was

used throughout.

References and Notes

1.   Compston A, Coles A. Multiple sclerosis. Vol. 372, The Lancet. Lancet; 2008. p. 1502–
17.

2.   Miyake S, Kim S, Suda W, Oshima K, Nakamura M, Matsuoka T, et al. Dysbiosis in the
gut microbiota of patients with multiple sclerosis, with a striking depletion of species
belonging to clostridia XIVa and IV clusters. PLoS One. 2015 Sep 14;10(9).

3.   Aaen G, Belman A, Benson L, Casper C, Chitnis T, Gorman M, et al. Associations
between the gut microbiota and host immune markers in pediatric multiple sclerosis and
controls. BMC Neurol. 2016 Sep 21;16(1).

4.   Chen J, Chia N, Kalari KR, Yao JZ, Novotna M, Soldan MMP, et al. Multiple sclerosis
patients have a distinct gut microbiota compared to healthy controls. Sci Rep. 2016 Jun
27;6.

5.   Jangi S, Gandhi R, Cox LM, Li N, Von Glehn F, Yan R, et al. Alterations of the human
gut microbiome in multiple sclerosis. Nat Commun 2016 71. 2016 Jun 28;7(1):1–11.

6.   Boziki MK, Kesidou E, Theotokis P, Mentis AFA, Karafoulidou E, Melnikov M, et al.
Microbiome in multiple sclerosis; where are we, what we know and do not know. Vol. 10,
Brain Sciences. MDPI AG; 2020.

7.   Merchak A, Gaultier A. Microbial metabolites and immune regulation: New targets for
major depressive disorder. Brain, Behav Immun - Heal. 2020 Nov 1;9:100169.

8.   Levinthal DJ, Rahman A, Nusrat S, O'Leary M, Heyman R, Bielefeldt K. Adding to the
Burden: Gastrointestinal Symptoms and Syndromes in Multiple Sclerosis. Mult Scler Int.
2013;2013:1–9.

9.   Cox LM, Maghzi AH, Liu S, Tankou SK, Dhang FH, Willocq V, et al. Gut Microbiome in
Progressive Multiple Sclerosis. Ann Neurol. 2021;89(6):1195–211.

10. Berer K, Gerdes LA, Cekanaviciute E, Jia X, Xiao L, Xia Z, et al. Gut microbiota from
multiple sclerosis patients enables spontaneous autoimmune encephalomyelitis in mice.
Proc Natl Acad Sci U S A. 2017 Oct 3;114(40):10719–24.

11. Jangi S, Gandhi R, Cox LM, Li N, Von Glehn F, Yan R, et al. Alterations of the human
gut microbiome in multiple sclerosis. Nat Commun. 2016 Jun 28;7.

12. Berer K, Mues M, Koutrolos M, AlRasbi Z, Boziki M, Johner C, et al. Commensal
microbiota and myelin autoantigen cooperate to trigger autoimmune demyelination.
Nature. 2011 Nov 24;479(7374):538–41.

13. Johanson DM, Goertz JE, Marin IA, Costello J, Overall CC, Gaultier A. Experimental
autoimmune encephalomyelitis is associated with changes of the microbiota composition
in the gastrointestinal tract. Sci Rep. 2020 Dec 1;10(1).

14. Barroso A, Mahler JV, Fonseca-Castro PH, Quintana FJ. The aryl hydrocarbon receptor
and the gut-brain axis. Cell Mol Immunol. 2021;18:259–68.

15. Hubbard TD, Murray IA, Bisson WH, Lahoti TS, Gowda K, Amin SG, et al. Adaptation
of the human aryl hydrocarbon receptor to sense microbiota-derived indoles. Sci Rep.
2015;5.

16. Jin UH, Lee SO, Sridharan G, Lee K, Davidson LA, Jayaraman A, et al. Microbiome-
Derived Tryptophan Metabolites and Their Aryl Hydrocarbon Receptor-Dependent
Agonist and Antagonist Activities. Mol Pharmacol. 2014;85(5).

17. Denison MS, Faber SC. And now for something completely different: Diversity in ligand-
dependent activation of Ah receptor responses. Vol. 2, Current Opinion in Toxicology.
2017.

18. Ott M, Avendanõ-Guzmán E, Ullrich E, Dreyer C, Strauss J, Harden M, et al.
Laquinimod, a prototypic quinoline-3-carboxamide and aryl hydrocarbon receptor agonist,
utilizes a CD155-mediated natural killer/dendritic cell interaction to suppress CNS
autoimmunity. J Neuroinflammation. 2019 Feb 26;16(1):1–14.

19. Kaye J, Piryatinsky V, Birnberg T, Hingaly T, Raymond E, Kashi R, et al. Laquinimod
arrests experimental autoimmune encephalomyelitis by activating the aryl hydrocarbon
receptor. Proc Natl Acad Sci U S A. 2016 Oct 11;113(41):E6145–52.

20. Korecka A, Dona A, Lahiri S, Tett AJ, Al-Asmakh M, Braniste V, et al. Bidirectional
communication between the Aryl hydrocarbon Receptor (AhR) and the microbiome tunes
host metabolism. npj Biofilms Microbiomes 2016 21. 2016 Aug 24;2(1):1–10.

21. Yang F, DeLuca JAA, Menon R, Garcia-Vilarato E, Callaway E, Landrock KK, et al.
Effect of diet and intestinal AhR expression on fecal microbiome and metabolomic
profiles. Microb Cell Fact. 2020 Dec 1;19(1):1–18.

22. Murray IA, Nichols RG, Zhang L, Patterson AD, Perdew GH. Expression of the aryl
hydrocarbon receptor contributes to the establishment of intestinal microbial community
structure in mice. Sci Rep. 2016;6.

23. Fling RR, Zacharewski TR. Aryl hydrocarbon receptor (Ahr) activation by 2,3,7,8-
tetrachlorodibenzo-p-dioxin (tcdd) dose-dependently shifts the gut microbiome consistent
with the progression of non-alcoholic fatty liver disease. Int J Mol Sci. 2021 Nov

1;22(22).

24.   Abdulla OA, Neamah W, Sultan M, Alghetaa HK, Singh N, Busbee PB, et al. The Ability
of AhR Ligands to Attenuate Delayed Type Hypersensitivity Reaction Is Associated With
Alterations in the Gut Microbiota. Front Immunol. 2021 Jun 29;12:2505.

25.   Rivet-Noor CR, Merchak AR, Li S, Beiter RM, Lee S, Thomas JA, et al. Stress-induced
despair behavior develops independently of the Ahr-RORγt axis in CD4□+□cells. Sci
Reports 2022 121. 2022 May 21;12(1):1–10.

26.   Kurschus FC. T cell mediated pathogenesis in EAE: Molecular mechanisms. Biomed J.
2015 May 1;38(3):183–93.

27.   Ehrlich AK, Pennington JM, Bisson WH, Kolluri SK, Kerkvliet NI. TCDD, FICZ, and
Other High Affinity AhR Ligands Dose-Dependently Determine the Fate of CD4+ T Cell
Differentiation. Toxicol Sci. 2018;161(2):310–20.

28.   Hong CH, Lin SH, Clausen BE, Lee CH. Selective AhR knockout in langerin-expressing
cells abates Langerhans cells and polarizes Th2/Tr1 in epicutaneous protein sensitization.
Proc Natl Acad Sci U S A. 2020 Jun 9;117(23):12980–90.

29.   Duarte JH, Di Meglio P, Hirota K, Ahlfors H, Stockinger B. Differential Influences of the
Aryl Hydrocarbon Receptor on Th17 Mediated Responses in vitro and in vivo. PLoS One.
2013 Nov 14;8(11):e79819.

30.   Quintana FJ, Murugaiyan G, Farez MF, Mitsdoerffer M, Tukpah AM, Burns EJ, et al. An
endogenous aryl hydrocarbon receptor ligand acts on dendritic cells and T cells to
suppress experimental autoimmune encephalomyelitis. Proc Natl Acad Sci U S A. 2010
Nov 30;107(48):20768–73.

31.   Angelou CC, Wells AC, Vijayaraghavan J, Dougan CE, Lawlor R, Iverson E, et al.
Differentiation of Pathogenic Th17 Cells Is Negatively Regulated by Let-7 MicroRNAs in
a Mouse Model of Multiple Sclerosis. Front Immunol. 2020 Jan 17;10:3125.

32.   Bhargava P, Smith MD, Mische L, Harrington E, Fitzgerald KC, Martin K, et al. Bile acid
metabolism is altered in multiple sclerosis and supplementation ameliorates
neuroinflammation. J Clin Invest. 2020 Jul 1;130(7):3467.

33.   Sayin SI, Wahlström A, Felin J, Jäntti S, Marschall HU, Bamberg K, et al. Gut microbiota
regulates bile acid metabolism by reducing the levels of tauro-beta-muricholic acid, a
naturally occurring FXR antagonist. Cell Metab. 2013 Feb 5;17(2):225–35.

34.   Ma C, Han M, Heinrich B, Fu Q, Zhang Q, Sandhu M, et al. Gut microbiome–mediated
bile acid metabolism regulates liver cancer via NKT cells. Science (80-). 2018 May
25;360(6391).

35.   MA Z, N S, PM M, M T, V G, JL W, et al. Butyrate suppresses colonic inflammation
through HDAC1-dependent Fas upregulation and Fas-mediated apoptosis of T cells. Am J
Physiol Gastrointest Liver Physiol. 2012 Jun 15;302(12).

36.   Kurita-Ochiai T, Ochiai K, Fukushima K. Butyric Acid-Induced T-Cell Apoptosis Is
Mediated by Caspase-8 and -9 Activation in a Fas-Independent Manner. Clin Diagn Lab
Immunol. 2001;8(2):325.

37.   Lewis ND, Patnaude LA, Pelletier J, Souza DJ, Lukas SM, King FJ, et al. A GPBAR1
(TGR5) Small Molecule Agonist Shows Specific Inhibitory Effects on Myeloid Cell
Activation In Vitro and Reduces Experimental Autoimmune Encephalitis (EAE) In Vivo.
PLoS One. 2014 Jun 26;9(6):e100883.

38.   Ho PP, Steinman L. Obeticholic acid, a synthetic bile acid agonist of the farnesoid X
receptor, attenuates experimental autoimmune encephalomyelitis. Proc Natl Acad Sci U S
A. 2016 Feb 9;113(6):1600.

39.   Duarte JH, Di Meglio P, Hirota K, Ahlfors H, Stockinger B. Differential Influences of the
Aryl Hydrocarbon Receptor on Th17 Mediated Responses in vitro and in vivo.
2013;8(11):e79819.

40.   Shin JH, Moreno-Nieves UY, Zhang LH, Chen C, Dixon AL, Linde MH, et al. AHR
Regulates NK Cell Migration via ASB2–Mediated Ubiquitination of Filamin A. Front
Immunol. 2021 Feb 24;12:94.

41.   Wang C, Ye Z, Kijlstra A, Zhou Y, Yang P. Activation of the aryl hydrocarbon receptor
affects activation and function of human monocyte-derived dendritic cells. Clin Exp
Immunol. 2014 Aug 1;177(2):521–30.

42.   Quintana FJ, Murugaiyan G, Farez MF, Mitsdoerffer M, Tukpah AM, Burns EJ, et al. An
endogenous aryl hydrocarbon receptor ligand acts on dendritic cells and T cells to
suppress experimental autoimmune encephalomyelitis. Proc Natl Acad Sci U S A.
2010;107(48):20768–73.

43.   Lee Y, Awasthi A, Yosef N, Quintana FJ, Xiao S, Peters A, et al. Induction and molecular
signature of pathogenic T H 17 cells. Nat Immunol. 2012;

44.   McAleer JP, Fan J, Roar B, Primerano DA, Denvir J. Cytokine Regulation in Human CD4
T Cells by the Aryl Hydrocarbon Receptor and Gq-Coupled Receptors. Sci Reports 2018
81. 2018 Jul 19;8(1):1–12.

45.   Rothhammer V, Borucki DM, Tjon EC, Takenaka MC, Chao CC, Ardura-Fabregat A, et
al. Microglial control of astrocytes in response to microbial metabolites. Nature.
2018;557:724–8.

46.   Rothhammer V, Mascanfroni ID, Bunse L, Takenaka MC, Kenison JE, Mayo L, et al.
Type i interferons and microbial metabolites of tryptophan modulate astrocyte activity and
central nervous system inflammation via the aryl hydrocarbon receptor. Nat Med.
2016;22(6).

47.   Parker A, Fonseca S, Carding SR. Gut microbes and metabolites as modulators of blood-
brain barrier integrity and brain health. Gut Microbes. 2020 Mar 3;11(2):135.

48.   Keogh CE, Kim DHJ, Pusceddu MM, Knotts TA, Rabasa G, Sladek JA, et al. Myelin as a
regulator of development of the microbiota-gut-brain axis. Brain Behav Immun. 2021 Jan
1;91:437–50.

49.   Hoban AE, Stilling RM, Ryan FJ, Shanahan F, Dinan TG, Claesson MJ, et al. Regulation
of prefrontal cortex myelination by the microbiota. Transl Psychiatry. 2016 Apr 5;6(4).

50.   Lycke NY, Bemark M. The regulation of gut mucosal IgA B-cell responses: recent
developments. Mucosal Immunol 2017 106. 2017 Jul 26;10(6):1361–74.

51.   Vaidyanathan B, Chaudhry A, Yewdell WT, Angeletti D, Yen WF, Wheatley AK, et al.
The aryl hydrocarbon receptor controls cell-fate decisions in B cells. J Exp Med.
2017;214(1):197.

52.   Araujo A, Safronova A, Burger E, López-Yglesias A, Giri S, Camanzo ET, et al. Ifn-g
mediates paneth cell death via suppression of mtor. Elife. 2021 Oct 1;10.

53.   Kamioka M, Goto Y, Nakamura K, Yokoi Y, Sugimoto R, Ohira S, et al. Intestinal
commensal microbiota and cytokines regulate Fut2+ Paneth cells for gut defense. Proc
Natl Acad Sci U S A. 2022 Jan 18;119(3).

54.   Šarenac TM, Mikov M. Bile acid synthesis: From nature to the chemical modification and
synthesis and their applications as drugs and nutrients. Front Pharmacol. 2018 Sep
25;9(SEP):939.

55.   Duc D, Vigne S, Bernier-Latmani J, Yersin Y, Ruiz F, Gaïa N, et al. Disrupting Myelin-
Specific Th17 Cell Gut Homing Confers Protection in an Adoptive Transfer Experimental
Autoimmune Encephalomyelitis. Cell Rep. 2019 Oct 8;29(2):378-390.e4.

56.   JG B, MV T, NL D, S O, JA S, E G, et al. Regulation of autoimmune myocarditis by host
responses to the microbiome. Exp Mol Pathol. 2017 Oct 1;103(2):141–52.

57.   Bergot AS, Giri R, Thomas R. The microbiome and rheumatoid arthritis. Best Pract Res
Clin Rheumatol. 2019;33(6).

58.   KL G, BP A, EMM Q. The microbiome and inflammatory bowel disease. J Allergy Clin
Immunol. 2020 Jan 1;145(1):16–27.

59.   J OR, TO K, LH K. The Gut Microbiome and Multiple Sclerosis. Cold Spring Harb
Perspect Med. 2018 Jun 1;8(6).

60.   YJ H, HA B. The microbiome in asthma. J Allergy Clin Immunol. 2015 Jan 1;135(1):25–
30.

61.   Chuang TY, Guo Y, Seki SM, Rosen AM, Johanson DM, Mandell JW, et al. LRP1
expression in microglia is protective during CNS autoimmunity. Acta Neuropathol
Commun. 2016 Jul 11;4(1):68.

62.   Seki SM, Stevenson M, Rosen AM, Arandjelovic S, Gemta L, Bullock TNJ, et al.
Lineage-Specific Metabolic Properties and Vulnerabilities of T Cells in the Demyelinating
Central Nervous System. J Immunol. 2017 Jun 15;198(12):4607–17.

63.   Remick DG, Bolgos GR, Siddiqui J, Shin J, Nemzek JA. Six at six: interleukin-6
measured 6 h after the initiation of sepsis predicts mortality over 3 days. Shock.
2002;17(6):463–7.

64.   Castellano-Escuder P, Andrés-Lacueva C, Sánchez-Pla A. POMA: User-friendly
Workflow for Metabolomics and Proteomics Data Analysis. PLoS Comput Biol.
2021;17(7):e1009148.

65.   Hartman AL, Lough DM, Barupal DK, Fiehn O, Fishbein T, Zasloff M, et al. Human gut
microbiome adopts an alternative state following small bowel transplantation. Proc Natl
Acad Sci U S A. 2009 Oct 6;106(40):17187–92.

66.   Barupal DK, Haldiya PK, Wohlgemuth G, Kind T, Kothari SL, Pinkerton KE, et al.

MetaMapp: mapping and visualizing metabolomic data by integrating information from
biochemical pathways and chemical and mass spectral similarity. BMC Bioinformatics.
2012 May 16;13(1):1–15.

67.   Bolyen E, Rideout JR, Dillon MR, Bokulich NA, Abnet CC, Al-Ghalith GA, et al.
Reproducible, interactive, scalable and extensible microbiome data science using QIIME
2. Nat Biotechnol 2019 378. 2019 Jul 24;37(8):852–7.

68.   Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. DADA2:
High-resolution sample inference from Illumina amplicon data. Nat Methods 2016 137.
2016 May 23;13(7):581–3.

69.   McMurdie PJ, Holmes S. phyloseq: An R Package for Reproducible Interactive Analysis
and Graphics of Microbiome Census Data. PLoS One. 2013 Apr 22;8(4):e61217.

Supplementary Materials

Figure S1. Separately housed $Cd4^{cre}Ahr^{fl/fl}$ recover from EAE phase. (A) No genotype differences in EAE incidence from the experiment shown in Figure 1A. (B) No differences in EAE clinical score or (C) incidence in cohoused male mice (Males; n= 7-8; N=1; Mann-Whitney U Test [p=0.5680]) (A) No genotype differences in EAE incidence from the experiment shown in Figure 1D. (B) Separately housed male $Cd4^{cre}Ahr^{fl/fl}$ mice recover from EAE while $Ahr^{fl/fl}$ mice maintain paralysis. (Males; n=7-8/group; N=1; Mann-Whitney U Test on total scores reported in legend [p=0.0010] and on single days reported on plot) (C) No difference in incidence of cohoused male mice.

Figure S2. Lack of AHR in T cells isolated from separately housed mice does not impact differentiation capacity or cytokine production. Naïve T cells were isolated from separately housed at weaning animals and differentiated toward T_H1, T_H2, T_H17 and T_{reg} *in vitro*. Transcription factors expression was quantified by Flow Cytometry. (**A**) Quantification of Tbet+ cells differentiated to T_H1 cell type and stimulated for 24 hours with either anti-CD3 or anti-

CD28 (n=4 mice/group; N=1 experiment; Ordinary one-way ANOVA [p=0.2927]). (**B**)

Quantification of GATA3+ cells differentiated to T_H2 cell type and stimulated for 24 hours with

either anti-CD3 or anti-CD28 (n=4 mice/group; N=1 experiment; Ordinary one-way ANOVA

[p=0.0076] followed by Tukey's Post-Hoc). (**C**) Quantification of RORγt+ cells differentiated to

T_H17 cell type and stimulated for 24 hours with either anti-CD3 or anti-CD28 (n=4 mice/group;

N=1 experiment; Ordinary one-way ANOVA [p=0.0163] followed by Tukey's Post-Hoc). (**D**)

Quantification of FoxP3+ cells differentiated to T_{reg} cell type and stimulated for 24 hours with

either anti-CD3 or anti-CD28 (n=4 mice/group; N=1 experiment; Ordinary one-way ANOVA

[p=0.8801]). (**E**) IFNγ ELISA on supernatant of differentiated T_H1 cells stimulated for 24 hours

with either anti-CD3 or anti-CD28 (n=4 mice/group; N=1 experiment; Ordinary one-way

ANOVA [p=0.8639]). (**F**) IL-4 ELISA on supernatant of differentiated T_H2 cells stimulated for

24 hours with either anti-CD3 or anti-CD28 (n=4 mice/group; N=1 experiment; Ordinary one-

way ANOVA [p=0.0064] followed by Tukey's Post-Hoc). (**G**) IL-17a ELISA on supernatant of

differentiated T_H17 cells stimulated for 24 hours with either anti-CD3 or anti-CD28 (n=4

mice/group; N=1 experiment; Ordinary one-way ANOVA [p=0.4826]).

Figure S3. Higher apoptosis in CD4 T cells lacking AHR. (**A**) Gene expression analysis

(qPCR array) performed on spinal cord RNAs prepared from separately housed $Cd4^{cre}Ahr^{fl/fl}$

mice and $Ahr^{fl/fl}$ littermate controls at the peak of disease (Day 16) shows increased pro-apoptotic

transcripts and decreased anti-apoptotic transcript (cDNA prepared from 2 spinal cords were

pooled for each sample). (**B**) Gating strategy for Annexin V+ apoptotic and necrotic cells shown

in show in Figure 4 and S3C-D. (**C**) Quantification of total dead cells in separately housed T_H17

differentiated cells (n=3 mice/group; N=1 experiment; Unpaired T Test [p=0.5130]). (**D**) T cells

differentiated to T_H1, T_H2, or T_{reg} *in vitro* from separately housed $Cd4^{cre}Ahr^{fl/fl}$ mice have higher

expression of the apoptotic marker Annexin V after 24 hours of CD3 stimulation. (n=3/group;

N=1 experiment; Unpaired T tests [p=0.4276, 0.0078, 0.0442]).

Figure S4. Only subtle microbiota population changes in *Cd4^{cre}Ahr^{fl/fl}* mice (**A**) PCA plot

and (**B**) microbiome composition at the genus level obtained from 16S sequencing performed on

feces collected from cohoused animals. (Males; n=6/group) (**C**) PCA plot and (**D**) microbiome

composition at the genus level obtained from 16S sequencing performed on feces collected from

separately housed animals. (Males and Females; n= 10/group) (**E**) PCA plot and (**F**) microbiome

composition at the genus level obtained from 16S sequencing performed on cecum collected

from separately housed animals. (Males and Females; n=12/group)

Data S1. Statistical report for all data presented in the manuscript.

Data S2. Raw Data files from Negative ion mode LC-MS corresponding to Figure 5 A-C.

Data S3. Raw Data files from Positive ion mode LC-MS corresponding to Figure 5 A-C.

DECLARATION:

Ethics approval and consent to participate:

Not applicable.

Consent for publication:

Not applicable.

Availability of data and materials:

The datasets supporting the conclusions of this article are available in the NCBI Gene Expression

Omnibus repository accession number GSE200440 and in additional files.

https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE200440

Competing interests:

899 Authors declare that they have no competing interests.

900 All authors have read and approved the manuscript.

Funding:

903 This work received funding from the National Institutes of Health (R33 MH108156 to A.G., T32

904 NS115657 to A.R.M., T32 GM008136 to R.M.B.), from the UVA Wagner fellowship (R.M.B.),

905 from the Owens Family Foundation (A.G.) from the UVA Trans University Microbiome

906 Initiative pilot grant (A.G. and A.R.M.) and from the UVA Presidential Fellowship in

907 Neuroscience (C.R.N.).

Author contributions:

All authors have read and approved the manuscript

Conceptualization: ARM, AG

Methodology: ARM, AG

Investigation: ARM, HJC, LCB, RMB, CRN, RMB, ERS

Data Curation: DGO

Formal Analysis: ARM, DGO

Supervision: AG

Visualization: ARM

Writing—original draft: ARM, AG

Writing—review & editing: HJC, LCB, RMB, CRN, RMB, DGO

Funding Acquisition: AG

Acknowledgments:

The authors wish to thank the University of Virginia Biorepository and Tissue Research facility

for their support with immunohistochemistry.

The authors would also like to thank Tula Raghavan, Dr. Sanja Arandjelovic, and Dr. Sharon

Walsh for critical reading and members the University of Virginia Brain Immunology and Glia

Center for scientific input.

Authors information:

Department of Neuroscience, University of Virginia, Charlottesville, VA, USA:

Andrea R. Merchak, Hannah J. Cahill, Lucille C. Brown, Ryan M. Brown, Courtney Rivet-Noor,

Rebecca M. Beiter, Erica R. Slogar, Deniz G. Olgun, and Alban Gaultier

Neuroscience Graduate Program, University of Virginia, Charlottesville, VA, USA:

Andrea R. Merchak, Ryan M. Brown, Courtney Rivet-Noor, Rebecca M. Beiter and Alban

Gaultier

Center for Brain Immunology and Glia, University of Virginia, Charlottesville, VA, USA:

Andrea R. Merchak, Ryan M. Brown, Courtney Rivet-Noor, Rebecca M. Beiter and Alban

Gaultier

A **Cohoused**

Ahr^{fl/fl}
Cd4^{cre} Ahr^{fl/fl}

B   *Ahr^{fl/fl}*   *Cd4^{cre} Ahr^{fl/fl}*

Chronic

400 µm

C

ns

D **Separately Housed**

Ahr^{fl/fl}
Cd4^{cre} Ahr^{fl/fl} **

*

E   *Ahr^{fl/fl}*   *Cd4^{cre} Ahr^{fl/fl}*

Peak

400 µm

Chronic

400 µm

F   *Ahr^{fl/fl}*

Peak   Chronic

G   *Cd4^{cre} Ahr^{fl/fl}*

*

Peak   Chronic

A

CD3   CD3
TUNEL

Ahr^{fl/fl}

Cd4^{cre} *Ahr*^{fl/fl}

B

TUNEL Quantification

#TUNEL+CD3+ Cells / FOV

Ahr^{fl/fl}   *Cd4*^{cre}*Ahr*^{fl/fl}

C

Dead Cells at Chronic Phase

% Dead Cells

CD4+   CD8+   TCRB+
Cell Type

D

Apoptosis in TH17 Cells
Separately Housed

% Annexin V+

Ahr^{fl/fl}   *Cd4*^{cre}*Ahr*^{fl/fl}

E

Apoptosis in TH17 Cells
Cohoused

% Annexin V+

Ahr^{fl/fl}   *Cd4*^{cre}*Ahr*^{fl/fl}

F

Apoptosis by Cecum Contents
C57BL6 TH17 Cells

% Annexin V+

Control   *Ahr*^{fl/fl}Cecum   *Cd4*^{cre}*Ahr*^{fl/fl}Cecum