A robust and transformation-free joint model with matching and regularization for metagenomic trajectory and disease onset

Qian Li[a,*], Kendra Vehik[b], Cai Li[a], Eric W. Triplett[c], Luiz Roesch[c], Yi-Juan Hu[d], Jeffery P. Krischer[b]

Department of Biostatistics, St. Jude Children's Research Hospital, Memphis, TN 38105 [a]; Health Informatics Institute, University of South Florida, Tampa, FL 33620 [b]; Department of Microbiology and Cell Science, University of Florida, Gainesville ,FL 32611 [c]; Department of Biostatistics and Bioinformatics, Emory University, Atlanta, GA 30322 [d]

ABSTRACT To identify operational taxonomy units (OTUs) signaling disease onset in an observational study, a powerful strategy was selecting participants by matched sets and profiling temporal metagenomes, followed by trajectory analysis. Existing trajectory analyses modeled individual OTU or microbial community without adjusting for the within-community correlation and matched-set-specific latent factors. We proposed a joint model with matching and regularization (JMR) to detect OTU-specific compositional trajectory predictive of host disease status, using nested random effects and covariate taxa pre-selected by Bray-Curtis distance and elastic net regression. We demonstrated that JMR effectively controlled the false discovery and pseudo biomarkers in a simulation study that generated temporal high-dimensional metagenomic counts with random intercept or slope. Application of the competing methods in the simulated data and the TEDDY cohort showed that JMR outperformed the other methods and identified important taxa in infants' fecal samples with dynamics preceding host disease status.

IMPORTANCE We proposed a framework to link taxon-specific compositional trajectory and the risk of host developing disease without transformation of the relative abundance. The heterogeneity between matched sets was considered in our model, with improved detection power and well-controlled false positive rate. The inherent negative correlation in microbiota composition due to sum-to-one constraint was adjusted by incorporating the top-correlated taxa as covariate. We designed a simulation pipeline to generate true biomarkers (i.e., causal OTUs and/or those interacting with causal OTUs) for disease onset and the pseudo biomarkers caused by sum-to-one or latent noises. Our method successfully reduced pseudo biomarker rate in simulation results, as well as identified more microbial trajectories signaling autoimmune status of young children enrolled in the TEDDY cohort.

KEYWORDS: metagenomic, trajectory, compositional data, joint model, nested random effect, zero-inflated, pseudo biomarker

INTRODUCTION

Gut microbiota profiled by 16s rRNA gene sequencing or metagenomic (i.e., whole-genome shotgun) sequencing has been frequently used in observational studies of

environmental exposures, immune biomarkers, and disease onset ([1], [2], [3], [4], [5]). One of the challenges in analyzing microbiota in an observational study is to incorporate the matching between participants by certain confounding risk factors (e.g. gender, clinical site, etc.) and/or disease status (case-control), such as the DIABIMMUNE and TEDDY cohorts ([1], [2], [5]). A matching design effectively eliminates the noise effect of sample collection, storage, shipment, sequencing batch, and environmental exposures confounding with disease outcomes, as well as reduces the costs of sample collection and sequencing. Statistical analyses for microbiota in matched sets included, but are not limited to, conditional logistic regression ([1]), non-parametric comparison PERMANOVA ([6]) and LDM ([7]) with extension to compare cases and controls within a matched set ([8]), which aimed to model and analyze microbiome data at independent time points.

Longitudinal profiling is a powerful strategy for the microbiome studies that aim to identify differential microbial trajectories between exposure groups or phenotypes ([9], [10]) or detect the time intervals of differential abundance ([11]). However, most of these studies failed to test if the compositional trajectory of an operational taxonomic unit (OTU) signaled host disease status. To detect microbial trajectories predictive of disease outcome in matched sets, an intuitive method is the generalized linear mixed effect model with or without the zero-inflation component ([9], [10], [12]), in which a taxon's abundance and/or presence is the outcome variable and the disease status is the co-variate of interest. The ZIBR model ([9]) tests the association between OTU and a co-variate factor using a two-part model for the non-zero relative abundance and presence of each OTU, assuming the non-zero relative abundance and presence being independent. A similar framework ([10]) was proposed to analyze the longitudinal zero-inflated counts per OTU using a Negative Binomial distribution, without converting the raw counts to relative abundance. A semi-parametric approach for longitudinal taxon-specific relative abundance is the linear mixed effect model (LMM) with asin-square-root transformation, which has been implemented in MaAsLin 2 ([12]).

One concern about using generalized linear mixed model to test the association between 16S rRNA or metagenomic trajectory and disease onset is that the covari-ates in this model may contribute to disease risk. For example, the HLA haplogeno-types and early use of probiotics may affect infants' gut microbiota and should be included as covariates. These factors were also found associated with islet autoim-munity among children enrolled in TEDDY ([13]). One usually added interaction terms between each covariate and the disease outcome ([3], [12]) to adjust for the association. However, a linear model with many interaction terms may lead to overfitting and re-duce the detection power ([14]). A sensible choice is the joint modeling of longitudinal biomarker and survival outcomes ([15], [16]), but there are limitations in applying this model to microbiome data in observational studies. First, the cost of metagenomic sequencing and the availability of fecal samples in a multi-center study may restrict the metagenome profiling to a subgroup of participants selected by certain criteria ([1], [2], [3]), whose survival outcome may deviate from common statistical assumptions. Second, the classic joint modeling approach aims to address repeated measurements of biomarkers in a time-to-event analysis rather than to test if a biomarker's intercept or slope is predictive of host health condition. Third, in an observational study that selects and matches participants by certain factors, their risk of developing disease is also matched. Thus, a survival submodel may not be capable of characterizing the disease risk between matched participants.

Many of the existing methods for microbiome data are built on the transformed relative abundance, such as centered log-ratio or inter-quartile log-ratio. In our new method, transformation of compositional data is not considered, since transforma-

tion strategy may have profound impact on analysis result and interpretation ([17]). The compositional change in true biomarkers (e.g., causal OTUs contributing to disease onset) always leads to simultaneous change in some other OTUs' composition because of sum-to-one constraint. In an observational study with matching design, it is common to collect and profile microbiota at many time points. The sum-to-one constraint and latent noise effect on microbiota may yield pseudo biomarkers with relative abundance associated with host disease status but not contributing to disease development. Hence, a taxon-level model is built for relative abundance trajectory that adjusts for the within-community correlation at each time point and reduces pseudo biomarker rate. In addition, we illustrate the performance of our method by a simulation pipeline that mimics the negative correlation in microbial community.

The latent technical noise in microbiome was removed by converting raw counts to relative abundance, and Zero-Inflated Beta density ([9]) was adopted to model an OTU's non-zero relative abundance and presence, respectively. We employed a subject-level random effect to link the logistic regression model of disease to a two-part longitudinal submodel. The latent effect of exposures related to matched set indicator was modeled by another random effect nested with the subject-level random effect. The OTU-disease association was assessed by jointly testing the scaling parameters for the subject-level random effect in the two-part submodel. We benchmarked the robustness and power of our method by a comprehensive simulation study and an application in the TEDDY cohort. The results illustrated that our method controlled the rates of false discovery and pseudo biomarkers, as well as improved the efficacy of detecting microbial trajectories signaling disease outcome.

RESULTS

We developed a Joint model with Matching and Regularization (JMR) to detect taxon-specific compositional trajectory associated with disease onset, adjusting for the top-correlated taxa and matched-set-specific latent noises. According to the characteristics of disease risk and infant-age gut microbiota in the TEDDY cohort, we designed a simulation pipeline similar to ([8]), generated the observed counts of temporal microbiota and compared our method to LMM and ZIBR using the simulated data. We also applied these methods to the shotgun metagenomic sequencing data profiled from the 4-9 months stool samples of infants enrolled in TEDDY cohort.

Overview of TEDDY microbiome study TEDDY is an observational prospective study of children at increased genetic risk of type 1 diabetes (T1D) conducted in six clinical centers in the U.S. and Europe (Finland, Germany, and Sweden). A total of 8,676 children were enrolled from birth and followed every 3 months for blood sample collection and islet autoantibody measurement up to 4 years of age, then every 3-6 months based on autoantibody status until the age of 15 years or diabetes onset ([18]). A primary disease endpoint in TEDDY is islet autoimmunity (IA), defined as persistent positive for insulin autoantibodies (IAA), glutamic acid decarboxylase autoantibodies (GADA), or insulinoma-associated-2 autoantibodies (IA-2A) at two consecutive visits confirmed by the two TEDDY laboratories ([18]). The participants' monthly stool samples were collected from 3-month age until the onset of IA or censoring with random missing samples ([1], [2]). Based on the sample availability and metagenomic sequencing cost, the microbiome study in TEDDY selected all the participants (cases) who developed IA by the design cutoff date May 31, 2012 and the controls at 1:1 case-control ratio matched by clinical center, gender, family history of T1D to profile the temporal gut microbiota, resulting in S = 418 matched sets (or pairs ([19])). These matching factors are known risk factors for type 1 diabetes. Some of the matched sets are at higher

risk of IA than the others due to higher risk human leukocyte antigen (HLA) genotypes,
geography or having family history of T1D. Hence, the matched participants have com-
parable risk of IA, but heterogeneity still exists between them according to the case-
control status by the design freeze date. The observed metagenomic counts table
in TEDDY was generated by the standard procedure of DNA extraction, PCR amplifi-
cation, shotgun metagenomic sequencing, assembly, annotation and quantification,
as described in (1). We visualized the top abundant species in the metagenomes of
TEDDY participants who had matched IA endpoint no later than 2 years of age (FIG 1).

FIG 1 **Trajectories of top abundant species in infant-age metagenomes grouped by host islet autoimmunity status at 2 years of age for a subgroup of TEDDY participants.**

Simulation Disease outcome for the matched participants are simulated by the
procedure below. The observed relative abundance per taxon were simulated by dif-
ferent scenarios. We first generated raw counts for a single OTU by Beta-Binomial
distribution to assess the robustness and power of our method JMR without covariate
taxa. We also designed a shifting procedure to mimic inherent negative correlation
in the true composition of microbiota and generated the temporal high-dimensional
raw counts table to evaluate the performance of compared methods.

1. **Generate disease outcome in matched sets** We defined matched sets and
subjects as 'high-risk' and 'low-risk' to generate the temporal OTU counts prior
to disease onset. Subjects are matched at 1:1 ratio. We first generated subject-
level and set-level random effects from a standard Normal distribution $a_{s_j} \sim$
$N(0, 1)$, $b_s \sim N(0, 1)$, $j = 1, \ldots, J, s = 1, \ldots, S$. Each random effect was converted
to a binary variable by the median value. That is $A_{s_j} = \mathbf{I}(a_{s_j} > \text{median}(a_{s_j}))$,
$B_s = \mathbf{I}(b_s > \text{median}(b_s))$, where $A_{s_j} = 1$ (or $B_s = 1$) represents a 'high-risk'
subject (or set). Next, we simulated a genetic risk factor G_{s_j} at subject-level with
effect size estimated from TEDDY data, and the disease status per subject by a
Bernoulli distribution $O_{s_j} \sim B(p_{s_j})$, where $\text{logit}(p_{s_j}) = \alpha_0 + \alpha_1 G_{s_j} + \alpha_2 A_{s_j} + \alpha_3 B_s$. We

set $(\alpha_0, \alpha_1, \alpha_3) = (0.5, -2, 1)$ and $\alpha_2 \in \{0.5, 0.75, 1, 1.25, 1.5\}$ to generate different

datasets, with α_1 estimated by JMR using the real data.

2. **Scenario A: single OTU counts.** We first simulated the observed counts of a

single OTU by Beta-Binomial (14) distribution to compare the univariate trajec-

tory methods without adjusting for covariate taxa. The true relative abundance

of an OTU at the earliest time point $t = 1$ was drawn from a Beta distribution

$\mu_1 \sim Beta(\mu_0, \phi_0)$, where parameters μ_0, ϕ_0 were estimated by applying Beta-

Binomial MLE to the metagenomic raw counts of an OTU selected at a given

relative abundance level in the TEDDY data. The relative abundance at later

time points $t > 1$ was generated by linearly increasing μ_1 to μ_t. The baseline rel-

ative abundance at time t in a matched set s was generated by $\mu_{st} \sim Beta(\mu, \phi_t)$,

and was increased or decreased by $\Delta\mu_{st}$ if the set was labeled as 'high-risk'. The

true relative abundance of this OTU for subject j in set s at time point t was sim-

ulated by $\mu_{s_j t} \sim Beta(\mu_{st}, \phi_{st})$, and was increased or decreased by $\Delta\mu_{s_j t}$ if the

subject was 'high-risk'. The total counts per sample, i.e., library size was drawn

from a Poisson distribution $N_{s_j t} \sim PS(100000)$, and the counts for this OTU is

generated from a Binomial (BN) distribution $Y_{s_j t} \sim BN(N_{s_j t}, \mu_{s_j t})$.

3. **Scenario B1: counts table with random intercept and pseudo biomarkers**

We also generated a high-dimensional counts table with $P = 1030$ OTUs to

demonstrate the performance of each method, so that the covariate taxa can be

used in JMR. First, we simulated the true composition of each microbiome sam-

ple by a shifting procedure combined with Dirichlet distribution, generating the

negative correlation within microbial community. Next, the sample-wise library

size was generated by a Poisson distribution, and the observed raw counts were

sampled from a Multinomial distribution, using the aforementioned true com-

position and sequencing library size. We visualized the simulation procedure of

Scenario B1 in FIG 2 for dimension of $P = 4$.

Step 1: Estimate the mean composition (or frequency) of microbiota ($\bar{\eta}_0$) and

the overdispersion ($\xi_0 = 0.04$) at the starting time point $t = 1$ in TEDDY data by

Dirichlet-Multinomial (DM) maximum likelihood estimate (MLE) of the observed

counts. Generate the mean frequency of microbiota at the first time point by

Dirichlet (DL) distribution: $\bar{\eta}_{01} \sim DL(\bar{\eta}_0, \xi_0)$.

Step 2: The mean frequency $\bar{\eta}_{0t}$ at later time points, $t > 1$, are generated by

the following shifting procedure: increase the composition of some OTUs in $\bar{\eta}_{01}$

(denoted by M^+_{base}) with a sum of Δ_t and simultaneously reduce the frequency

of other OTUs in $\bar{\eta}_{01}$ (denoted by M^-_{base}) by Δ_t. The absolute shift size Δ_t rep-

resented the age effect on microbiota and linearly increased with t. This shift-

ing strategy characterized the inherent correlation between M^+_{base} and M^-_{base}

because of the simultaneous shift-up and -down in true composition. All the

OTUs in $\bar{\eta}_{0t}$ are assigned to either M^+_{base} or M^-_{base} to account for the impact of

various latent exposures across time points.

Step 3: At each time point, the heterogeneity between matched sets is the

overdispersion estimated by DM MLE based on the samples per time point in

TEDDY, denoted by ξ_t. The overdispersion at the first time point is $\xi_1 = 0.05$

and linearly decreases over time, which mimics the time-dependent overdisper-

sion observed in the infant-age metagenome in TEDDY. We generated a mean

frequency for each matched set s at time point t by $\bar{\eta}_{st} \sim DL(\bar{\eta}_{0t}, t)$. If a set

is labeled as 'high-risk', we shifted all the OTUs in $\bar{\eta}_{st}$ using the procedure in

Step 2 with shift size Δ_{st}, which is a proportion of the maximum shift size, i.e.,

$\Delta_{st} = \gamma\Delta^0_{st}$. We set $\gamma \in \{0.6, 0.7, 0.8\}$ to evaluate method performance at different

FIG 2 Shifting procedure in simulation scenario B1.

set-level random effect size.

Step 4: The between-subject heterogeneity within each matched set was the

median DM MLE of overdispersion per matched set based on the real data, that

is $\xi^* = 0.03$. Hence, we generated the true microbiota composition for a sample

collected from a 'low-risk' subject j in set s at time t by $\bar{\eta}_{sjt} \sim DL(\bar{\eta}_{st}, \xi^*)$. For a

subject labeled as 'high-risk', we increased 15% OTUs (denoted by M^+) in $\bar{\eta}_{sjt}$ by

Δ_{sjt}, and reduced another 15% OTUs (denoted by M^-) by $d\Delta_{sjt}$ ($0 < d < 1$). The

subsets M^+ and M^- are the true biomarkers for disease status. We randomly

selected a third subset (denoted by M^0) from the remaining 70% OTUs in $\bar{\eta}_{sjt}$ and

reduced the composition of M^0 by a total of $(1 - d)\Delta_{sjt}$. There may exist OTUs

never selected in M^+, M^-, or M^0, which are the 'null' OTUs. The OTUs selected in

M^0 are the pseudo biomarkers due to random shift in frequency. Similarly, we

set the total shift $\Delta_{sjt} = \lambda\Delta^0_{sjt}$ at distinct effect size $\lambda \in \{0.5, 0.6, 0.7, 0.8\}$, where

Δ^0_{sjt} is the maximum shift restricted by sum-to-one.

Step 5: The library size for each sample is simulated by a Poisson distribution

$N_{sjt} \sim PS(100000)$, truncated by a minimum of 10000. The raw counts per sam-

ple is generated by Multinomial (MN) distribution $C_{sjt} \sim MN(N_{sjt}, \bar{\eta}_{sjt})$.

4. **Scenario B2: counts table with random slope and pseudo biomarkers** Data

generation process for this scenario is similar to Scenario B1, except that the

shift size Δ_{sjt} from 'low-risk' to 'high-risk' subjects varies by time points. It's worth

to note that we cannot distinguish 'false positive' from 'pseudo positive' in sce-

narios B1 and B2. Hence, we use the sum of false positive rate (FPR) and pseudo

positive rate (PPR), i.e., false or pseudo positive rate (FPPR) as a performance

metric for scenarios B1,B2. That is FPPR $= \frac{\text{\# of positives in } M^0}{\text{\# of OTUs in } M^0}$.

5. **Scenario C: counts table without pseudo biomarkers** In this scenario we con-

sidered random intercept signaling the disease onset and fixed half OTUs in $\bar{\eta}_{sjt}$

as 'null' in order to evaluate the FPR and FDR of each method, although this sce-
nario is not applicable to real data. Among the other half OTUs, we selected 10%
OTUs in $\bar{\eta}_{s_jt}$ as M^+ and 40% OTUs as M^- without a subset of pseudo biomarkers
(M^0).

In scenario A, we compared JMR not adjusting for correlated taxa (JMR-NC) with the
following methods : a) a joint model with regularization but without matching indicator
and correlated taxa (JR-NC); b) the ZIBR model with a Wald statistic jointly testing OTU-
specific abundance or presence using either a single random effect (ZIBR-S) or nested
random effects (ZIBR-N); c) LMM with arcsin-square-root transformation using either
a single random effect (LMM-S) or nested random effects (LMM-N). For ZIBR and LMM
methods, we used R package gamlss and set the sample age, genotype, disease status,
and genotype-disease interaction term as the fixed effect covariates.

We randomly selected 6 OTUs with different relative abundance from TEDDY data
and estimated the baseline parameters for each. Then we generated $n = 10000$ repli-
cates for each OTU with $S \in \{50, 100\}$. The type I error rate and power of each method
was calculated at statistical significance level $p < 0.05$, shown in Table 1. The results
showed that JMR-NC persistently controlled the type I error and provided higher detec-
tion power at distinct abundance levels except for the OTUs with relative abundance
between $(10^{-2}, 10^{-3})$ and $(10^{-6}, 10^{-5})$. Type I error of the reduced model JR-NC was
severely inflated in some datasets and its power was lower than JMR. LMM consis-
tently controlled type I error, while the power of LMM was lower than that of JMR-NC
in most of the simulated OTUs. The ZIBR method yielded inflated type I error rate and
low efficacy regardless of sample size.

FIG 3   FPPR of each method for scenario B1 at different size of matched-set-specific random effect.

We generated 10 replicates of each OTU table to assess the performance of com-
peting methods in scenarios B1, B2 and C. The taxa associated with disease onset in
each OTU table are detected by FDR cutoff $q < 0.15$. The FPPR in scenario B1 (FIG 3)
showed that adjusting for the top-correlated taxa in JMR successfully controlled the
rate of pseudo biomarkers across distinct scenarios with power higher than LMM at
larger sample size ($N = 200$), although this strategy resulted in lower detection power
compared to JMR-NC. The results in FIG 4 also demonstrated the outperformance of
JMR-NC in terms of sensitivity and FDR in the taxon-specific trajectory analysis for an
OTU table. The LMM methods were powerful in the test of intercept, but this model oc-

TABLE 1 The type I error and power based on 10000 simulated replicates for a taxon at different levels of mean relative abundance (y).

$-\log_{10}(y)$		(0, 1]	(1, 2]	(2, 3]	(3, 4]	(4, 5]	(5, 6]
				Type I error			
	JMR-NC	0.01	0.003	0.008	0.0004	0	0
	JR-NC	0.061	0.03	0.001	0.002	0.0003	0.0005
$N = 100$	LMM-N	0.045	0.031	0.026	0.033	0.068	0.052
($S = 50$)	LMM-S	0.041	0.036	0.023	0.036	0.066	0.049
	ZIBR-N	0.057	0.066	0.080	0.053	0.164	0.369
	ZIBR-S	0.049	0.079	0.076	0.063	0.163	0.366
	JMR-NC	0.018	0.004	0.022	0.051	0.001	0.0003
	JR-NC	0.162	0.091	0.049	0.066	0.035	0.002
$N = 200$	LMM-N	0.043	0.039	0.068	0.057	0.079	0.058
($S = 100$)	LMM-S	0.046	0.051	0.066	0.060	0.079	0.056
	ZIBR-N	0.056	0.067	0.099	0.078	0.170	0.324
	ZIBR-S	0.059	0.088	0.099	0.088	0.174	0.324
				Power			
	JMR-NC	0.399	0.5	0.372	0.833	0.798	0.136
	JR-NC	0.32	0.65	0.051	0.057	0.548	0.040
$N = 100$	LMM-N	0.040	0.084	0.474	0.107	0.552	0.512
($S = 50$)	LMM-S	0.043	0.090	0.468	0.108	0.347	0.468
	ZIBR-N	0.06	0.088	0.412	0.123	0.743	0.666
	ZIBR-S	0.057	0.095	0.405	0.123	0.686	0.664
	JMR-NC	0.452	0.723	0.619	0.944	0.917	0.388
	JR-NC	0.677	0.699	0.194	0.836	0.737	0.324
$N = 200$	LMM-N	0.06	0.115	0.429	0.478	0.287	0.253
($S = 100$)	LMM-S	0.068	0.134	0.423	0.321	0.273	0.206
	ZIBR-N	0.057	0.332	0.433	0.667	0.280	0.379
	ZIBR-S	0.063	0.344	0.430	0.648	0.265	0.377

casionally produced inflated FPPR as shown in FIG 3 regardless of the set-level random effect size. Furthermore, the power of LMM was unstable in intercept analysis, while its power in slope analysis was nearly zero. A larger set-level random effect size yielded higher FPPR in LMM, but this impact was trivial in the other methods. JMR showed the best performance in slope analysis, with higher sensitivity (FIG 4) and the lowest FPPR. Results of scenario C in FIG 5 showed that JMR and LMM effectively controlled the FPR, while LMM produced higher FPR at larger matched-set-specific random effect (γ). The FDR of JMR at $\gamma = 0.6$ was relatively higher than that of LMM due to lower sensitivity. The ZIBR methods yielded inflated FPPR in the high-dimensional scenarios, although the power of this model was higher than the competing methods regardless of sample size. The severely-inflated FPR and FDR by ZIBR in scenario C (FIG 5) is consistent with the results of scenario B (FIG 3).

Application in TEDDY We applied the competing methods to the longitudinal metagenomes profiled from TEDDY children's monthly stool samples collected at the age of 4-9 months (1). We included the cases developing IA between 9-month and 24-month age and their matched controls who remained IA-negative by the cases' diagnosis age. For each matched pair included in the present analysis, one participant was IA positive and the other one was negative at the age of 24 months. We excluded the participant(s) matched to multiple pairs, yielding $N = 152$ subjects ($S = 76$ pairs) and $n = 672$ metagenome samples. The cases who experienced IA onset after 24 months and their matched controls were not included in this analysis.

FIG 4 **Sensitivity of each method for scenarios B1 and B2 at different sample size (N) and different effect size (λ).**

FIG 5 **FDR and FPR for each method in scenario C at different size of matched-set-specific random effect.**

We first filtered OTUs at genus and species level by relative abundance $> 10^{-6}$ and prevalence $> 5\%$, selecting 125 genera and 365 species in downstream analysis. The sample age and the hosts' breastfeeding status per time point were used as longitudinal covariates, while HLA DR3&4 haplotype was included as time-invariant covariates. For the LMM and ZIBR methods, we used the interaction term between IA status and binarized HLA category (DR3&4 vs. others) as a covariate to adjust for the association. We tested each OTU's association with IA by FDR cutoff $q < 0.05$ or $q < 0.1$, individually. The results in Table 2 showed that JMR identified more OTUs than LMM in both

intercept and slope analysis. The LMM methods only found a small subgroup of taxa associated with IA at either genus or species level. We also visualized the overlap and difference between JMR, JMR-NC, LMM-N in FIG 6, and then compared Akaike Information Criterion (AIC) of JMR and JMR-NC for the 76 species detected by both methods. Adjusting for the correlated taxa in JMR did improve model fitting with lower mean AIC (-2631.847) compared to JMR-NC (-2615.571). LMM-N is not comparable to JMR or JMR-NC in terms of information criteria, since the taxon-specific relative abundance was transformed by asin-square-root.

TABLE 2 The number of genera and species associated with IA detected by each method in a subgroup of TEDDY participants.

		Intercept		Slope	
FDR		$q < 0.05$	$q < 0.1$	$q < 0.05$	$q < 0.1$
Genera	JMR	27	31	27	34
	JMR-NC	44	52	36	44
	LMM-N	11	16	3	4
	LMM-S	49	82	10	19
	ZIBR-N	26	30	31	36
	ZIBR-S	26	32	31	36
Species	JMR	75	94	37	46
	JMR-NC	147	166	112	125
	LMM-N	43	60	13	21
	LMM-S	40	63	7	16
	ZIBR-N	89	106	119	138
	ZIBR-S	83	105	120	140

FIG 6 Venn diagram for the intercept analysis in TEDDY data by JMR, JMR-NC, LMM-N.

The taxa with mean abundance (intercept) associated with IA onset exclusively detected by both JMR and JMR-NC at $q < 0.1$ include *Bifidobacterium.breve*, *Bacteroides.fragilis*, *Lactobacillus.ruminis*, *Veillonella.ratti*. *B.breve*, as one of the three species dominating infant-age gut microbiota in TEDDY, was less abundant in intercept (i.e. at 4- and 9-month) during infancy among IA cases, with density shown in FIG 7. The species *B.fragilis* as part of the normal microbiota in human colon was found more abundant among IA cases compared to their matched controls (FIG 1). This *Bacteroide* species was also found differential between T1D cases and controls at only one time point in a small-size Finnish cohort ([20]).

There are two more abundant species *Faecalibacterium.prausnitzlii* and *Escherichia.coli*

FIG 7 Distribution of relative abundance for *B.breve* and *E.coli* per time point grouped by 2-year IA status.

visualized in FIG 1 associated with IA in slope and exclusively detected by JMR. *F.prausnitzlii*, as one of the most abundant and important commensal bacteria of human gut microbiota that produces butyrate and short-chain fatty acids from the fermentation of dietary fiber, increased faster in IA cases after 6-month of age. This rapid change and abnormally higher level of *F.prausnitzlii* prior to IA seroconversion may be a result of the sudden change of dietary pattern during infancy.

Our method successfully detected the case-control difference in the slope of *E.coli*, which was found as an amyloid-producing bacteria with temperal dynamics heralding IA onset in a subset analysis in DIABIMMUNE cohort (21). The relative abundance of *E.coli* in TEDDY smoothly decreased from 4-month to 9-month for both cases and controls (FIG 1), and it was relatively more abundant in controls between 7- and 9-month with stratified densities shown in FIG 7. The temporal change of *E.coli* prior to IA seroconversion in TEDDY detected by JMR was consistent with the decrease of *E.coli* reported in DIABIMMUNE cohort (21), which was possibly due to prophage activation according to the *E.coli* phage/*E.coli* ratio prior to *E.coli* depletion in that research.

DISCUSSION

We developed a joint model with nested random effects to test the association between taxa and disease risk, and adjusted for the correlated taxa screened by a pre-selection procedure in abundance and prevalence, individually. We implemented our method in an R package mtradeR (metagenomic trajectory analysis with disease endpoint and risk factors), available at https://github.com/qianli10000/mtradeR. The JMR function implemented the framework in equation (1) by parallel computing. We also provided simulation functions StatSim and TaxaSim to generate (binary) disease status and temporal high-dimensional metagenomic counts of matched sets.

The simulation of single OTU demonstrated the performance of each method at

different relative abundance levels, implying that LMM with either single or nested random effect is still a robust method. The simulation of high-dimensional OTU tables also illustrated LMM's overall performance in intercept analysis, but the unstableness of LMM is a concern in the real data analysis. JMR yielded lower false or pseudo positive rate in the simulated datasets and higher detection power in slope analysis by adjusting for the top-correlated taxa in modeling. The pre-selection of top-correlated taxa in JMR was performed in relative abundance and presence, individually, being consistent with the two-part model strategy. According to the simulation study, a disadvantage of JMR is the power being relatively low at small sample size and depends on the tuning of shrinkage parameter. Adding nodes in the GH approximation may improve the power of JMR, but more nodes will also lead to additional computation costs. Hence, future work should focus on improvement of JMR in both detection power and computation efficiency.

Another limitation of our method is the bias of scaling parameter (λ_r, λ_p) estimate, potentially caused by the L_2 penalty. Our current work only focused on the unsigned association between a taxon and host disease status by using a Wald statistic. An improvement in the estimate of scaling parameter and statistical inference should be considered in future work, such as the algorithm in ZINQ ([14]). We did not use quantile regression in current research, since the performance of ZINQ required tuning of grid. But ZINQ provided an alternative approach for modeling zero-inflation in microbiota composition with fewer statistical assumptions.

There are other important topics to be considered in the modeling of longitudinal microbiome data. One potential direction is high dimensional modeling framework, such as tensor singular value decomposition ([22]). A promising extension of the current work in JMR is to exploit functional data analysis for multiple microbial trajectories. By employing a non-parametric joint modeling, we may be able to capture nonlinear trends and heterogeneous patterns of longitudinal biomarkers in microbiota, as well as negative correlations among taxa ([23]).

MATERIALS AND METHODS

Joint model with matching and regularization (JMR) The probability for participant j ($j = 1, ..., J$) in matched set s ($s = 1, ..., S$) developing the disease of interest is $p_{s_j} = P(O_{s_j} = 1)$, where O_{s_j} is the binary disease status. There are J participants in each matched set. Let $y_{s_j t}$ be the relative abundance of an OTU for participant j in matched set s at time point t ($t = 1, ..., T_{s_j}$). We denote the expected non-zero abundance by $\mu_{s_j t} = E(y_{s_j t} | y_{s_j t} > 0)$, and the probability of presence (or zero-inflation) by $\pi_{s_j t} = P(y_{s_j t} > 0)$, similar to ([9]). For a microbiome study matching participants by the disease-associated factors and/or disease status (e.g., DIABIMMUNE, TEDDY), the matched participants are assumed to have comparable but distinct disease risk. Hence, we model the disease status by a logistic mixed effect model with nested random effects. A joint model for the host disease status and microbial trajectory in matched set is

$$
\begin{aligned}
\text{logit}(p_{s_j}) &= \boldsymbol{u}_{s_j}\boldsymbol{\alpha} + a_{s_j} + b_s \\
\text{logit}(\mu_{s_j t}) &= \boldsymbol{x}_{s_j t}^{(1)}\boldsymbol{\beta}_{11} + \boldsymbol{z}_{s_j t}\boldsymbol{\beta}_{12} + \tilde{z}_{s_j t}(\lambda_r a_{s_j} + \gamma_r b_s) \\
\text{logit}(\pi_{s_j t}) &= \boldsymbol{x}_{s_j t}^{(2)}\boldsymbol{\beta}_{21} + \boldsymbol{z}_{s_j t}\boldsymbol{\beta}_{22} + \tilde{z}_{s_j t}(\lambda_p a_{s_j} + \gamma_p b_s)
\end{aligned}
\tag{1}
$$

The host disease status is determined by a vector of fixed effect covariates $\boldsymbol{u}_{s_j}$ and the independent nested random effects a_{s_j}, b_s. The non-zero relative abundance $\mu_{s_j t}$ and presence $\pi_{s_j t}$ per OTU are predicted by the same random effects rescaled by parameters λ_r, λ_p and a vector of clinical or bioinformatics technical covariates $\boldsymbol{z}_{s_j t}$. To

account for the inherent correlation between taxa, this OTU's non-zero abundance
and presence per time point are also determined by a vector of top-correlated taxa's
relative abundance $\boldsymbol{x}_{s_j t}^{(1)}$ and presence-absence $\boldsymbol{x}_{s_j t}^{(2)}$ measured at the same time point,
pre-selected by a procedure described below. The two-part submodel of $y_{s_j t}$ character-
izes how the trajectory is affected by subject- and set-level latent factors contributing
to disease risk, and how the OTU trajectory interacts with correlated taxa over time. If
an OTU is a pseudo biomarker, then its relative abundance ($y_{s_j t}$) should be driven by
the top-correlated taxa per time point instead of the disease-associated random effect
a_{s_j}. On the other hand, the abundance of a true biomarker OTU at each time point is
mainly determined by the latent risk of disease onset (a_{s_j}, b_s) and possibly associated
with the top-correlated taxa.

We set $\tilde{z}_{s_j t} = 1$ in equation (1) to test intercept, and $\tilde{z}_{s_j t} = age$ to test slope. The
nested random effects and parameters λ_r, λ_p provide flexibility in the modeling of
between-subjects and between-sets heterogeneity, as well as model the abundance-
presence correlation in each taxon by shared nested random effects instead of assum-
ing independence between the two processes as in (9).

Parameter estimation and hypothesis testing To account for the sum-to-one
restriction on non-zero relative abundance ($0 < \mu_{s_j t} < 1$) and the binarized measure-
ment $\boldsymbol{I}(y_{s_j t} > 0)$ of an OTU, we intuitively employ the Zero-Inflated Beta (ZIB) density
function (9) to define the match-set-specific marginal likelihood for parameter estima-
tion. That is $L(\theta; \boldsymbol{y}, \boldsymbol{O}) = \prod_{s=1}^{S} L_s$, where

$$L_s = \int_{-\infty}^{\infty} \ldots \int_{-\infty}^{\infty} g_b(b_s) \prod_{j=1}^{J} I_{s_j}^{(1)}(a_{s_j}, b_s) I_{s_j}^{(2)}(a_{s_j}, b_s) g_a(a_{s_j}) \, da_{s_1} \ldots da_{s_J} db_s \tag{2}$$

$$\begin{aligned}
I_{s_j}^{(1)}(a_{s_j}, b_s) &= \prod_{t=1}^{T_{s_j}} [(1 - \pi_{s_j t}) \boldsymbol{I}(y_{s_j t} = 0) + \pi_{s_j t} \boldsymbol{I}(y_{s_j t} > 0) f(y_{s_j t} | y_{s_j t} > 0)] \\
I_{s_j}^{(2)}(a_{s_j}, b_s) &= p_{s_j} \boldsymbol{I}(O_{s_j} = 1) + (1 - p_{s_j}) \boldsymbol{I}(O_{s_j} = 0)
\end{aligned} \tag{3}$$

$g_a(a_{s_j}), g_b(b_s)$ are the Gaussian density functions with mean 0 and variance σ_a^2, σ_b^2, in-
dividually, and $f(y_{s_j t} | y_{s_j t} > 0)$ is the Beta density function with mean $\mu_{s_j t}$ and overdis-
persion ϕ. In the simulation study, we demonstrated that the robustness and perfor-
mance of this model does not require the observed relative abundance being gener-
ated from ZIB distribution.

The estimate of overdispersion $\hat{\phi}$ without regularization is severely inflated and
also leads to bias in the estimate of other parameters. Hence, we use L_2 (ridge) regu-
larization to control the overdispersion and type I error in hypothesis testing. All the
parameters θ are estimated by maximizing a penalized marginal likelihood function
$\hat{\theta} = \arg \max \tilde{L}(\theta; \boldsymbol{y}, \boldsymbol{O})$, where

$$\tilde{L}(\theta; \boldsymbol{y}, \boldsymbol{O}) = \ln L(\theta; \boldsymbol{y}, \boldsymbol{O}) - \rho ||\theta||_2^2 \tag{4}$$

425 and ρ is selected by a cross-validation described below.

426 There is no closed form of the multivariate integral L_s in equation (2) because of
427 the Beta density in $I_{s_j}^{(1)}(a_{s_j}, b_s)$. Hence, L_s can be approximated by Gauss-Hermite (GH)
428 quadrature, with details explained in Appendix. We test the association between OTU
429 trajectory and host disease status with null hypothesis:

$$H_0 : \lambda_r = \lambda_p = 0 \tag{5}$$

and a Wald statistic $W = \frac{\hat{\lambda}_r^2}{SE_{\lambda_r}^2} + \frac{\hat{\lambda}_p^2}{SE_{\lambda_p}^2}$, which follows a Chi-Square distribution $W \sim$
$\chi^2(2)$. The false discovery rate (FDR) for multiple testing is corrected by the Benjamini-
Hochberg (BH) procedure.

Pre-selection of correlated taxa and tuning parameter selection For each OTU
($y_{s_j t}$) in equation (1), using all the other taxa as covariates is computationally inefficient.
Hence, we use a two-step procedure to pre-select $\boldsymbol{x}_{s_j t}^{(1)}$ and $\boldsymbol{x}_{s_j t}^{(2)}$, and then perform a
post-selection hypothesis testing. The first step screens the taxa correlated with $y_{s_j t}$
in abundance and presence, individually, using the Bray–Curtis distance (24) less than
0.1 quantile. This step may still result in many covariate taxa at species level in meta-
geonmic data due to high dimensionality. Thus, we employ elastic net regression to
further select the taxa with relative abundance $\boldsymbol{x}_{s_j t}^{(1)}$ associated with $y_{s_j t}$ or the taxa
with presence $\boldsymbol{x}_{s_j t}^{(2)}$ associated with $\boldsymbol{I}(y_{s_j t} > 0)$, individually. In this pre-selection proce-
dure, we model all the longitudinal metagenomes as independent samples regardless
of time points (or age).

To reduce the computational burden of cross-validation for a high-dimensional
OTU table, we randomly select P_0 OTUs from distinct relative abundance levels to rep-
resent the complexity of microbiota composition. The matched sets are divided into
5 folds, each being a validation fold for the model built on the other four (training)
folds. The penalized log likelihood in equation (4) is the negative objective function in
cross-validation. For each validation fold f and the selected OTU i, the loss function is
$S_{fi} = -\tilde{L}(\hat{\theta}_{-f}^i; \boldsymbol{y}_f^i, \boldsymbol{O}_f)$, where $\hat{\theta}_{-f}^i = \arg\max \tilde{L}(\theta^i; \boldsymbol{y}_{-f}^i, \boldsymbol{O}_{-f})$. The optimal ρ is selected by
the 'elbow point' minimizing $S = \sum_{f=1}^{5} \sum_{i=1}^{P_0} S_{fi}/(5P_0)$.

Availability of data and materials. The TEDDY Microbiome WGS data that sup-
ports the findings of this study have been deposited in NCBI's database of Genotypes
and Phenotypes (dbGaP) with the primary accession code phs001443.v1.p1. The R
package mtradeR that implements JMR and the simulation pipeline is available at https:
//github.com/qianli10000/mtradeR.

ACKNOWLEDGMENTS

The work of QL was in part supported by U24DK097771 from the National Institute of
Diabetes, Digestive and Kidney Diseases via the NIDDK Information Network's (dkNET)
New Investigator Pilot Program in Bioinformatics. QL and CL are in part supported
by P30 Cancer Center Support Grant (CA21765) funded by National Cancer Institute;
and the American Lebanese Syrian Associated Charities (ALSAC). The TEDDY study is
funded by the National Institute of Diabetes and Digestive and Kidney Diseases, Na-
tional Institute of Allergy and Infectious Diseases, National Institute of Child Health
and Human Development, National Institute of Environmental Health Sciences, Cen-
ters for Disease Control and Prevention, and JDRF. We thank the TEDDY study data co-
ordinating center at Health Informatics Institute, University of South Florida for data
processing and sharing. We thank Suraj Sarvode Mothi for data cleaning support.

QL proposed and implemented the model and algorithm, performed the exper-
iments, analyzed the data, and drafted the manuscript. KV conceived the real data
analysis, interpreted results, and contributed to manuscript writing. QL and YH de-
signed the experiments. CL contributed to methodology discussion and manuscript
writing. ET and LR contributed to manuscript writing. JK supervised the study design,
sample collection, data generation and analysis in the TEDDY cohort.

REFERENCES

1. **Stewart CJ, Ajami NJ, O'Brien JL, Hutchinson DS, Smith DP, Wong MC, Ross MC, Lloyd RE, Doddapaneni H, Metcalf GA, et al.**. 2018. Temporal development of the gut microbiome in early childhood from the TEDDY study. Nature 562 (7728):583–588.

2. **Vatanen T, Franzosa EA, Schwager R, Tripathi S, Arthur TD, Ve-hik K, Lernmark Å, Hagopian WA, Rewers MJ, She JX, et al.**. 2018.

The human gut microbiome in early-onset type 1 diabetes from the TEDDY study. Nature 562 (7728):589–594.

3. **Wang DD, Nguyen LH, Li Y, Yan Y, Ma W, Rinott E, Ivey KL, Shai I, Willett WC, Hu FB, et al.**. 2021. The gut microbiome modulates the protective association between a Mediterranean diet and car-diometabolic disease risk. Nat medicine 27 (2):333–343.

4. **Schirmer M, Smeekens SP, Vlamakis H, Jaeger M, Oosting M, Franzosa EA, Ter Horst R, Jansen T, Jacobs L, Bonder MJ, et al.**. 2016. Linking the human gut microbiome to inflammatory cytokine production capacity. Cell 167 (4):1125–1136.

5. **Vatanen T, Kostic AD, d'Hennezel E, Siljander H, Franzosa EA, Yassour M, Kolde R, Vlamakis H, Arthur TD, Hämäläinen AM, et al.**. 2016. Variation in microbiome LPS immunogenicity contributes to autoimmunity in humans. Cell 165 (4):842–853.

6. **Anderson MJ**. 2001. A new method for non-parametric multivariate analysis of variance. Austral ecology 26 (1):32–46.

7. **Hu YJ, Satten GA**. 2020. Testing hypotheses about the microbiome using the linear decomposition model (LDM). Bioinformatics 36 (14):4106–4115.

8. **Zhu Z, Satten GA, Mitchell C, Hu YJ**. 2021. Constraining PERMANOVA and LDM to within-set comparisons by projection improves the efficiency of analyses of matched sets of microbiome data. Microbiome 9 (1):1–19.

9. **Chen EZ, Li H**. 2016. A two-part mixed-effects model for analyzing longitudinal microbiome compositional data. Bioinformatics 32 (17):2611–2617.

10. **Zhang X, Yi N**. 2020. NBZIMM: negative binomial and zero-inflated mixed models, with application to microbiome/metagenomics data analysis. BMC bioinformatics 21 (1):1–19.

11. **Metwally AA, Yang J, Ascoli C, Dai Y, Finn PW, Perkins DL**. 2018. MetaLonDA: a flexible R package for identifying time intervals of differentially abundant features in metagenomic longitudinal studies. Microbiome 6 (1):1–12.

12. **Mallick H, Rahnavard A, McIver LJ, Ma S, Zhang Y, Nguyen LH, Tickle TL, Weingart G, Ren B, Schwager EH, et al.**. 2021. Multivariable association discovery in population-scale meta-omics studies. PLoS computational biology 17 (11):e1009442.

13. **Uusitalo U, Liu X, Yang J, Aronsson CA, Hummel S, Butterworth M, Lernmark Å, Rewers M, Hagopian W, She JX, et al.**. 2016. Association of early exposure of probiotics and islet autoimmunity in the TEDDY study. JAMA pediatrics 170 (1):20–28.

14. **Ling W, Zhao N, Plantinga AM, Launer LJ, Fodor AA, Meyer KA, Wu MC**. 2021. Powerful and robust non-parametric association testing for microbiome data via a zero-inflated quantile approach (ZINQ). Microbiome 9 (1):1–19.

15. **Luna PN, Mansbach JM, Shaw CA**. 2020. A joint modeling approach for longitudinal microbiome data improves ability to detect microbiome associations with disease. PLoS computational biology 16 (12):e1008473.

16. **Hu J, Wang C, Blaser MJ, Li H**. 2021. Joint modeling of zero-inflated longitudinal proportions and time-to-event data with application to a gut microbiome study. Biometrics .

17. **Quinn TP, Erb I, Gloor G, Notredame C, Richardson MF, Crowley TM**. 2019. A field guide for the compositional analysis of any-omics data. GigaScience 8 (9):giz107.

18. **Group TS**. 2008. The environmental determinants of diabetes in the young (TEDDY) study. Ann New York Acad Sci 1150 (1):1–13.

19. **Lee HS, Burkhardt BR, McLeod W, Smith S, Eberhard C, Lynch K, Hadley D, Rewers M, Simell O, She JX, et al.**. 2014. Biomarker discovery study design for type 1 diabetes in The Environmental Determinants of Diabetes in the Young (TEDDY) study. Diabetes/metabolism research reviews 30 (5):424–434.

20. **Giongo A, Gano KA, Crabb DB, Mukherjee N, Novelo LL, Casella G, Drew JC, Ilonen J, Knip M, Hyöty H, et al.**. 2011. Toward defining the autoimmune microbiome for type 1 diabetes. The ISME journal 5 (1):82–91.

21. **Tetz G, Brown SM, Hao Y, Tetz V**. 2019. Type 1 diabetes: an association between autoimmunity, the dynamics of gut amyloid-producing E. coli and their phages. Sci reports 9 (1):1–11.

22. **Han R, Shi P, Zhang AR**. 2021. Guaranteed Functional Tensor Singular Value Decomposition. arXiv preprint arXiv:2108.04201 .

23. **Li C, Xiao L, Luo S**. 2021. Joint model for survival and multivariate sparse functional data with application to a study of Alzheimer's Disease. Biometrics .

24. **Weiss S, Van Treuren W, Lozupone C, Faust K, Friedman J, Deng Y, Xia LC, Xu ZZ, Ursell L, Alm EJ, et al.**. 2016. Correlation detection strategies in microbial data sets vary widely in sensitivity and precision. The ISME journal 10 (7):1669–1681.

25. **Sharma D, Xu W**. 2021. phyLoSTM: a novel deep learning model on disease prediction from longitudinal microbiome data. Bioinformatics 37 (21):3707–3714.

26. **Han Y, Baker C, Vogtmann E, Hua X, Shi J, Liu D**. 2021. Modeling longitudinal microbiome compositional data: a two-part linear mixed model with shared random effects. Stat Biosci 13 (2):243–266.

27. **Benjamini Y, Hochberg Y**. 1995. Controlling the false discovery rate: a practical and powerful approach to multiple testing. J Royal statistical society: series B (Methodological) 57 (1):289–300.

28. **Schiffer C, Lalanne AI, Cassard L, Mancardi DA, Malbec O, Bruhns P, Dif F, Daëron M**. 2011. A strain of Lactobacillus casei inhibits the effector phase of immune inflammation. The J Immunol 187 (5):2646–2655.

29. **Schiffer C, Lalanne AI, Cassard L, Mancardi DA, Malbec O, Bruhns P, Dif F, Daëron M**. 2011. A Strain of Lactobacillus casei Inhibits the Effector Phase of Immune Inflammation. The J Immunol 187 (5):2646–2655. doi:10.4049/jimmunol.1002415.

Mean composition per time point

| | Taxon A | Taxon B | Taxon C | Taxon D |

T1 — 0.33, 0.33, 0.33, 0.01
T2 — 0.43, 0.23, 0.3, 0.04
T3 — 0.5, 0.23, 0.23, 0.04
T4 — 0.53, 0.25, 0.17, 0.05

Mean composition per low-risk set

Mean composition per high-risk set

A low-risk subject in high-risk set

A high-risk subject in high-risk set

Low-risk set T1 — 0.32, 0.34, 0.325, 0.015

High-risk set T1 — 0.51, 0.2, 0.27, 0.02

Low-risk subject in high-risk set T1 — 0.52, 0.195, 0.273, 0.012 — M^0, M^+, M^-

High-risk subject in high-risk set T1 — 0.338, 0.3, 0.25, 0.012