1 **Mitochondrial effects on fertility and longevity in *Tigriopus***

2 ***californicus* contradict predictions of the mother's curse**

3 **hypothesis**

4 ## ABSTRACT

5

6 Strict maternal inheritance of mitochondria favors the evolutionary accumulation of sex-biased

7 fitness effects, as mitochondrial evolution occurs exclusively in female lineages. The "mother's

8 curse" hypothesis proposes that male-harming mutations should accumulate in mitochondrial

9 genomes when they have neutral or beneficial effects on female fitness. Rigorous empirical tests

10 have largely focused on *Drosophila*, where support for the predictions of mother's curse has

11 been mixed. We investigated the impact of mother's curse mutations in *Tigriopus californicus,* a

12 minute crustacean. Using nonrecombinant backcrosses, we introgressed four divergent

13 mitochondrial haplotypes into two nuclear backgrounds and recorded measures of fertility and

14 longevity. We found that the phenotypic effects of mitochondrial mutations were context-

15 dependent, being influenced by the nuclear-background in which they were expressed, as well as

16 the sex of the individual and rearing temperature. Mitochondrial haplotype effects were greater

17 for fertility than longevity, and temperature effects were greater for longevity. However, in

18 opposition to mother's curse expectations, females had higher mitochondrial genetic variance

19 than males for fertility and longevity, little evidence of sexual antagonism favoring females was

20 found, and the impacts of mitonuclear mismatch harmed females but not males. Together, this

21 indicates that selection on mitochondrial variation has not resulted in the accumulation of male

22 mutation load in *Tigriopus californicus*.

Introduction

In eukaryotes, life is powered by mitochondria - cytoplasmic organelles capable of producing

energy in the form of ATP. Originating from an ancient symbiosis over 2 billion years ago,

mitochondria provided the energy required for the evolution and diversification of multicellular

organisms[1–3]. Over evolutionary history, mitochondrial genomes have retained some aspects

of their ancestral genetic codes and a few dozen genes while dramatically reducing their genome

size through deletion and gene transfer to the nucleus. Beyond the generation of ATP by

oxidative phosphorylation (OXPHOS), mitochondria perform additional cellular functions,

including initiation of apoptosis, amino acid and steroid biosynthesis, β-oxidation of fatty acids,

cellular signaling, and nuclear epigenetic regulation [4–8]. Nevertheless, out of 1,500 or more

proteins found in the mitochondrion, only the 13 membrane-bound electron transport chain

complexes are encoded by the mitochondrial genome, with the rest being encoded by nuclear

DNA[9,10]. The coevolution of mitochondrial and nuclear genomes is thus an ancient yet

ongoing process, involving the evolutionary fine-tuning of various molecular interactions (i.e.

protein-protein, protein-DNA, and protein-RNA) required for the accurate replication and

transcription of mtDNA, translation of mtDNA-encoded proteins, and efficient cellular

respiration. Mutation rates are generally higher in mtDNA than in nDNA[11,12] leading to a

continuous introduction of new mitochondrial alleles. With asexual transmission and lack of

recombination in mitochondria, novel alleles are likely to persist even when slightly deleterious

due to the potential masking effect of linkage disequilibrium on mitochondrial genetic variation

and reduced efficacy of natural selection[13], although debate over the efficacy of selection on

mtDNA continues to produce compelling counterarguments (see [14–18]). Nevertheless, mtDNA

likely experiences recurrent adaptive evolution via selective sweeps acting on traits such as

respiratory efficiency and mitonuclear coadaptation[19–23]. In natural populations,

mitochondrial fitness variation is likely to be present at substantial levels and may make a

meaningful contribution to adaptation and diversification[24]. The distribution of fitness effects

among mtDNA mutations is also likely to differ between the sexes, owing to the maternal

inheritance of mitochondria[25–27]. Strict maternal inheritance favors the evolutionary

accumulation of sex-biased fitness effects, as it is expected to preclude any opportunity for

evolution in males, forming a sex-specific selective sieve.

Under maternal inheritance, mitochondrial evolution is thought to be restricted to female

lineages, leading to the accumulation of male-harming mutations when they have neutral or

beneficial effects on female fitness (i.e. "mother's curse"[26,28]). Initial evidence for the

mother's curse hypothesis (MCH) centered on evidence that sperm motility in humans and other

taxa was linked to mitochondrial function[29,30,28] therefore sperm are more exposed to

deleterious effects of mtDNA mutations than ova. The MCH has also been proposed to impact

viability, and even to contribute to the taxonomically widespread pattern of shorter male

lifespans[31]. Theoretical work, however, shows that conditions favoring the spread of mother's

curse mutations can be limited by even modest levels of inbreeding and kin selection, allowing

for an adaptive response to mtDNA alleles impacting male viability and fertility[32–34].

Nevertheless, interest in MCH has increased steadily in recent years leading to formal tests of the

hypothesis, mostly in *Drosophila* but also in plants (reviewed in [35]), utilizing carefully

constructed mitochondrial-nuclear (mitonuclear) hybrids with divergent mtDNA haplotypes

backcrossed into an isogenic nuclear background. The most direct test of MCH focuses on the

effects of mutational variation in mtDNA, which are predicted to confer larger trait variance in

males. A second major prediction focuses on the subset of alleles that are directly sexually

antagonistic, which could lead to a negative intersexual genetic correlation across haplotypes. A

third prediction is that experimental disruption of mitonuclear coadaptation will be more

detrimental to males than females. Studies using the mitonuclear hybrid approach have shown

both support[31,36–40], and rejection[41–44] of the mother's curse hypothesis.

Understanding the pervasiveness of mother's curse is crucial given its fundamental implications

for the evolution of sexual dimorphism, as well as very practical consequences for conservation

breeding programs and medicine (particularly mitochondrial replacement theory and male-biased

mitochondrial disease). While past work has focused on *Drosophila*, the predominance of

maternal inheritance of mitochondria in eukaryotes allows for a taxonomically diverse

investigation of the generality of mother's curse. Here we test predictions of the MCH using the

splash-pool copepod *Tigriopus californicus*, an arthropod that shares a common ancestor with

Drosophila around 409.3 – 536.6 million years ago[45]. We used a genetic panel of mitonuclear

hybrids to assay both fertility and longevity. As fertility genes are likely sex-specific and

viability genes are likely shared between the sexes (see review[46]), we predict that mitonuclear

mismatch will lead to stronger sex bias for fertility than longevity. Additionally, we measured

the impact of high environmental temperature on these traits, as mild mitonuclear mismatch can

be expected to be unmasked under higher metabolic demand[47]. Populations of *T. californicus*

show extreme genetic divergence over short geographical distances[48–50] while remaining

somewhat tolerant of hybridization, producing viable and fertile offspring[51,52]. Nucleotide

divergence in *T. californicus* extends across the mtDNA genome, involving coding and

noncoding genes, and can exceed 20% between distant populations. *T. californicus* also differ

from standard animal models in that they lack heteromorphic sex chromosomes - sex is

determined by the inheritance of multiple, unlinked genes in combination with environmental

effects (polygenic sex determination)[53–55]. While rarely utilized in evolutionary biology,

polygenic sex determination provides an advantage for testing sex-specific mitochondrial effects

as sex differences cannot be attributed to the effects of sex chromosomes. Importantly, despite

the lack of heteromorphic sex chromosomes, there is marked sexual dimorphism in *T.*

californicus and considerable evidence for sex-specific effects in multiple traits (thermal

tolerance[56,57]; salinity tolerance[58,59]; resistance to toxins[59]; intrinsic hybridization

stress[54]) in which males are generally the weaker sex.

Methods

Population sampling

Populations of *T. californicus* are distributed in supralittoral "splash pools" along the Pacific

coast of North America. Copepods were originally collected from supralittoral pools at four

locations (Figure 1A): Friday Harbor, WA (FHL: 48.55 N,123.01 W), La Jolla, CA (LJ: 32.83

N,117.27 W), Bird Rock, CA (BR: 32.82 N,117.27 W), and San Diego, CA (SD: 32.7460

N,117.2551 W). Isofemale lines were initiated from a single female fertilized by a single male,

and maintained at 20°C on a 12h:12h light:dark cycle in standard culture media (three times

filtered seawater from Wrigley Marine Science Center, Santa Catalina Island, CA; 0.1g/l of

Spirulina and 0.1g/l ground Tetramin (Tetra®) Tropical fish food flakes). All lines were

maintained in petri dishes for ≥ 5 years under these conditions and periodically fed additional

Spirulina and ground Tetramin® tropical fish food and rehydrated with DI water whenever

necessary to maintain approximately 35 ppt salinity. Prior to experimentation, we sequenced

samples from each isofemale line at the 12S ribosomal RNA locus (Forward primer:

CCAAGATACTTTAGGGATAACAGC; Reverse primer: CTGTTCTATCAAAACAGCCCT;

fragment size: 687 base pairs) to rule out cross-contamination.

Mitonuclear hybrid panel

In *Tigriopus*, meiosis is chiasmatic in males but achiasmatic in females, yielding recombinant

sperm and non-recombinant ova. This allows us to backcross mitochondria from the population

of the dam into that of the sire while preventing nuclear recombination between populations.

Because sexes in *T. californicus* are not easily distinguished before maturity, we allowed males

to clasp female siblings with their geniculate first antennae in order to ensure the sex of females.

We moved clasped pairs onto filter paper and quickly separated them using entomological pins

under a microscope. We backcrossed FHL, BR, LJ, and SD dams to SD and BR sires for 10

generations (Figure 1B), resulting in an average proportion of $1 - \frac{1}{2}^{(b+1)}$ (where b is equal to the

number of backcross generations) or 99.95% of the sire's nuclear genome, assuming no

selection. We chose the SD and BR lines for our isogenic nuclear backgrounds because SD was

the source for the <u>*Tigriopus californicus*</u> genome assembly, BR is the genetically closest

resequenced population to SD, and it would be experimentally intractable to perform a full

factorial design. For each cross, we established 4 replicates, each maintained in 12-well plates

containing standard culture media. All crosses were fed weekly with additional Spirulina and

ground Tetramin® fish food and rehydrated with DI water whenever necessary to maintain

approximately 35 ppt salinity.

Trait measurement and modeling

For each of our mitonuclear hybrid lines, we measured three fitness components (fertility,

lifespan, and age-specific rate of survival) for males and females reared at two contrasting

temperature regimes (15ºC and 25ºC). In order to achieve an adequate sample size for each

mtDNA haplotype and sex, it was necessary to set up multiple families from each mitonuclear

hybrid line. Each family was composed of up to four clutches of fertilized eggs derived from a

single mitonuclear hybrid or control female (dam). Dams were maintained at standard conditions

and checked daily for the presence of hatched eggsacs. For each dam, we alternated moving up

to four hatched eggsacs to 15ºC and 25ºC (Figure 1B) so that clutch order was orthogonal to

temperature in both fertility and longevity experiments. Individuals were maintained at the same

temperature for the duration of the fertility or longevity experiment.

Longevity

Individual animals were separated into wells in a 24-well plate once they developed to the

juvenile copepodid stage and were checked three times a week. The first measurement we

recorded was the sex-specific age of reproductive maturity. For males, this occurred with the

development of geniculate first antennae (claspers) and for females with the development of

eggs, which can be seen through the translucent body. We continued monitoring longevity

experiments three times a week, recording the age of death of individuals until all had expired.

We performed linear mixed-effects modeling on family sex ratio using sex information from our

longevity experiment, comparing mtDNA x nDNA interactions with the impact of mitonuclear

status on family sex ratio. We also conducted a separate experiment, using different families,

measuring mortality occurring before sexual maturity (Document S1, Figure S1). We were able

to measure the naupliar, copepodid, and total development rate, and perform linear mixed-effects

modeling investigating the influence of genetic (mtDNA, nDNA) and environmental

(temperature) influence on these juvenile life-history parameters.

Sex-specific fertility

For female fertility experiments, all individuals from hatched eggsacs were separated into

individual wells in a 24-well plate once they developed to the juvenile copepodid stage. To each

well, we added a single wild-caught adult male from the appropriate locality (SD or BR),

matching the paternal line of the individual. Males were added to every well because sexes are

difficult to distinguish morphologically in immature *T. californicus* and we wanted to ensure that

each female was paired with an adult male before sexual maturity. Mate clasping typically occurs

between adult males and juvenile females, and rarely occurs with mature females once they

begin extruding non-fertilized eggsacs. We also ran experiments utilizing sib-pairs in parallel to

compare the impact of mating with mitonuclear hybrid males versus wild-type males on female

fertility (Document S1). Plates were maintained at the appropriate temperature (15ºC or 25ºC)

and checked three times a week until females produced an eggsac. As we utilized several

hundred plates in our experiment, it became necessary to maximize incubator space. Therefore,

we placed gravid females from the same family and clutch together in petri dishes until they

reached experimental age. Because individuals develop twice as fast at 25ºC as at 15ºC, we used

temperature-specific ages for our experiment in order to minimize age-related variation. Once

individuals reared at 25☐ reached 35 days of age, and individuals reared at 15☐ reached 70 days

of age, we moved them into separate dishes and maintained them at the correct temperature,

checking them three times a week until eggsacs hatched. Upon hatching, each individual larval

nauplius was counted under a dissecting microscope to measure female hatching number.

For male fertility experiments, we removed males from each of the four clutches in a family and

allowed them to age for the appropriate temperature (35 days old for 25ºC and 70 days old for

15ºC). Once at the appropriate age, each male was pipetted into a petri dish with six wild-caught

juvenile females from the appropriate locality (BR or SD) to match the paternal nuclear line. To

ensure that wild-caught juveniles were female, we selected clasped pairs from culture within 0-2

weeks of collection and separated pairs under a dissecting microscope. Care was taken to provide

males with females of consistent and reasonably large size, to reduce variation across

experiments. Each experimental male and six wild-caught females were allowed 14 days to mate

in a petri dish at the appropriate temperature. After 14 days, females were isolated into a 6-well

plate and scored for the presence or absence of any offspring indicating a successful mating. We

recorded male remating success as the ratio of successful matings to the number of surviving

females.

Parametric and nonparametric trait modeling

For our trait modeling, we performed three types of linear models: sex-effect modeling,

coevolutionary status modeling, and best-fit models. Sex-effect models measure sex effects

impacted by mtDNA and nDNA variation and their interactions. If mother's curse mutations are

present, they may exist in additive (Sex x mtDNA) or epistatic (Sex x mtDNA x nDNA) form.

Sex differences may also be encoded by nuclear genes, described by the (Sex x nDNA) term. In

order to compare fertility experiments across the two sexes, individual measurements of fertility

were normalized to the mean of the appropriate matched control line reared at the same

temperature (e.x. male *i* from FHLxSD at 15□ / mean of all males from SDxSD at 15□). We

performed model testing on the four sex-effect models using the *model.sel* function in the

MuMIn[60] R package, which ranks models based on their AIC_C values. Model contrasts for sex

differences were performed using the *pairs* function from the emmeans[61] R package .

Coevolutionary status models measure the impact of disrupting coevolved mitonuclear

combinations on sex-specific data grouped by mtDNA haplotype. These models included

temperature and coevolutionary status as fixed-effect terms, compared mitonuclear hybrids of

both nDNA backgrounds with both matched controls and included nDNA as a random term.

Mitonuclear hybrids with the SD or BR mtDNA were only compared to one matched control.

Modeling for best-fit utilized an "all-subset" approach, where all possible combinations of

predictors were compared using the *MuMIn* function *dredge*. Models with the highest weight and

lowest AIC_C were designated as best-fit. Both fertility and longevity models included

temperature, nDNA, mtDNA and all possible interactions as fixed-effects terms in initial models

and were performed separately for each sex. For fertility, we used raw sex-specific fertility

(males: remating success, females: hatching number). All linear mixed-effects modeling was

performed using the *lmer* function from the lme4[62] R package and included clutch nested

within family as a random-effect term. We used type III ANOVA to estimate significance for

model main effects and interaction terms using the *Anova* function from the car[63] R package.

We performed parametric regression of survivorship for each nuclear, temperature, and sex

combination against the Gompertz distribution [64], estimating the shape (α) and rate (β)

parameters using the R package flexsurv[65]. Parameters were estimated by maximum

likelihood, with those defined to be positive estimated on a log scale. Confidence intervals were

estimated from the Hessian at maximum, and then transformed back to the original scale (days).

The shape of the resulting distributions, $\ln(\alpha)$, describe the initial mortality rate or frailty of the

samples, while the rate (β) parameter describes the age-independent rate of mortality acceleration

or rate of senescence. We employed Kruskal-Wallis one-way analysis of variance to estimate the

significance of parameter estimates produced by parametric survival models using the

kruskal.test function from the stats[66] R package.

Mitochondrial coefficient of variation

One prediction of the mother's curse hypothesis is that the accumulation of male-affecting

variation should lead to a higher mitochondrial coefficient of variation (CV_{mt}) in males than in

females. Using the *boot*[67,68] package in R, bootstrapped CV_{mt} was computed by resampling

raw trait means for each mitochondrial line using 1000 replicates with replacement and

calculating the coefficient of variation among mitochondrial lines ($CV = \sigma/\mu$) for each nuclear,

sex, and temperature combination. Hypothesis testing was performed to compare male and

female CV_{mt} using a sign test as implemented in the *SIGN.test* function in the BSDA[69] R

package.

Genetic correlation among haplotypes

Mother's curse mutations are likely to reach fixation fastest when they confer a fitness advantage

to females, resulting in sexual conflict when they simultaneously result in low fitness in males.

When enough mtDNA variation is involved in sexual conflict, a negative intersexual correlation

may be observed across haplotypes for a given trait. We measured the correlation between

haplotype means for fertility and median longevity, comparing traits within sexes (intrasexual

correlation) and also comparing the sexes for a given trait (intersexual correlation). We used

bootstrapped estimates of Pearson's coefficient using the boot R package along with the *cor*

function from the stats R package, recording 1000 samples for each comparison. Equi-tailed,

two-sided nonparametric confidence intervals were generated from the first order normal

approximation with a confidence level of 0.95 using the *boot.ci* function from the boot package

in R.

Whole genome sequencing

For each isofemale and mitonuclear hybrid line, we extracted DNA from 10 males and 10

females using Zymo quick-DNA preps and quantified DNA yields using Qubit HS dsDNA

quantification kits. Because female *T. californicus* yield approximately 2.12x more DNA than

males, we pooled 2 ng of DNA from each of the 20 individuals per line for library preparation.

DNA pools were sheared using a Covaris S2 sonicator with microTUBE-50 AFA tubes, and

sequencing libraries were prepared using Qiagen QIAseq Ultralow Input kits following

instructions, using 8 PCR cycles during library amplification. Mitonuclear hybrid libraries were

pooled together in equimolar amounts and sequenced together on a single lane of Illumina

HiSeq4000 (PE-150), and the parental isofemale libraries were pooled and sequenced on a

separate lane. Sequence reads were quality trimmed using trimmomatic[70] v0.36, and paired

reads were filtered for mitochondrial reads using the filter_reads.pl perl script included with the

NOVOPlasty[71] software. We assembled consensus mtDNA genomes from each sequenced

isofemale line, by utilizing the appropriate mtDNA reference from a previous publication which

included the SD, BR, and FHL populations[23]. A reference mtDNA genome did not exist for

the LJ population, so those reads were mapped to the BR mtDNA reference. We annotated our

de novo assembled genomes using the Mitos[72] 2.0 software, each containing all 13 protein

coding genes, and the majority of non-coding RNAs. Following this, we mapped all trimmed and

paired reads to the *T. californicus* reference genome assembly (NCBI accession:

GCA_007210705.1) along with the appropriate mitochondrial genome using stampy[73] v1.0.32

(substitutionrate=0.05). Following the removal of PCR and optical duplicates using picard[74]

MarkDuplicates, we called variants for each sequenced pool using samtools[75] mpileup.

We estimated pairwise nuclear divergence for isofemale lines in a sliding window approach. For

each comparison, we calculated F_{ST} using a sliding window of 10,000 bp, requiring a depth of

40x coverage, and minimum PHRED base quality score of 30 using the fst-sliding.pl perl script

included with popoolation v1.2.2. For parental mitochondrial divergence, we aligned whole

mtDNA genomes using progressiveMauve[76] along with mtDNA genome sequences from an

outgroup population from San Roque, Baja California Sur, Mexico to produce rooted trees, and

measured divergence (d_{XY}) using the calculate-dxy.pl perl script included with popoolation

v1.2.2. We also measured divergence at synonymous and nonsynonymous sites by aligning

protein-coding sequences with MACSE [77] v2, constructing neighbor joining trees with

ClustalW [78] and analyzing these data with codeml from the PAML suite [79] using the null

model (M0), assuming a uniform selective pressure among sites.

Confirming mitonuclear genetic status of hybrids

To confirm the isogenic nuclear status of our lines, we measured allele frequencies at a set of

divergent sites for each paternal line's nuclear genome. A Fisher's exact test for significant allele

frequency differences for nDNA from the maternal and paternal lines was conducted using

popoolation[80] v1.201, requiring a minimum depth of 20x coverage. In mitonuclear hybrids,

sites with no significant differences (alpha > 0.05) to the paternal line were considered

"paternally concordant" and significantly different sites (alpha <= 0.05) were considered

"paternally discordant". Because significance may be determined by variations in the same

alleles between lines, we recorded paternally discordant sites with alleles matching maternal

lines as "maternally concordant" when they were absent in the paternal nDNA. Paternal

concordance rates were recorded by measuring the proportion of sites that were not maternally

concordant to the total number of sites. Lee and Willett[81] recently reported the presence of

paternally inherited mtDNA, likely due to paternal leakage, in F1 hybrid *T. californicus*. This

paternal mtDNA was found in a high frequency of individuals (up to 59%) but a very low

frequency of reads (<0.01% in pooled samples). We therefore tested for paternally inherited

mtDNA in our mitonuclear hybrid lines. This was accomplished by mapping trimmed and paired

sequence reads to the reference genome as well as all mtDNA haplotypes simultaneously. We

measured depth of coverage for each mtDNA haplotype using samtools depth command to

identify any potential sources of contamination.

Results

DNA sequencing

Illumina HiSeq 4000 sequencing yielded an average of 34.7 million reads for mitonuclear hybrid line pools, and 76.1 million reads for inbred line pools. The read sequences were deposited at the National Center for Biotechnology Information (NCBI) Sequence Read Archive (SRA) under the project accession number PRJNA840866. Following quality trimming and mapping of paired reads to the appropriate references, an average of 29 million reads mapped for cybrid pools, and 61 million reads mapped for maternal and paternal line pools. Average nuclear genome coverage across inbred lines was 91.88% at a mean depth of coverage of 26x. Across the mitochondrial genome, the mean depth of coverage was 2670X, and genome coverage was 100%. Nuclear divergence (F_{ST}) between paternal lines ranged from 0.049 to 0.369 in lines with BR nDNA, and from 0.102 to 0.378 in lines with SD nDNA (Figure 1C). Mitochondrial divergence (d_{XY}) between parental lines ranged from 0.7% to 21.1% in BR nDNA lines, and 9.6% to 20.6% in SD nDNA lines. The mitonuclear hybrid panel represented a wide range of mitochondrial variation, with as few as 13 amino acid substitutions between the BR and LJ mitotypes, 170 aa substitutions between the BR and SD nuclear line mitotypes, and 555 aa substitutions between SD and FHL haplotypes. The highly divergent FHL haplotype also differed from the BR & LJ haplotypes by 546 aa substitutions.

For each mitonuclear hybrid line, we measured allele frequencies at an average of 580k divergent nDNA SNPs between parental lines. We found that alleles were mostly concordant with paternal nDNA, with limited evidence of introgression from maternal nDNA, with paternal concordance rates ranging from 99.15% - 100% (Figure 1C). To test for potential mtDNA contamination via

paternal leakage or experimental error, we compared depth of coverage across all mtDNA

haplotypes, since mtDNA positions and alleles are not comparable across haplotypes. We found

that the percent maternally matching depth ranged from 94.09% to 99.81%, while the percent of

paternally matching depth ranged from 0.02% to 0.28% providing limited evidence for

contaminating mtDNA (Figure 1C). While we cannot determine with certainty whether this result

confirms paternal leakage, it is important to note that paternally matching depth measured in our

study may also be confounded with nuclear-encoded mitochondrial DNA fragments (numts)

which also match the paternal mtDNA haplotype. Further, because pooled-DNA sequencing

lacks haplotype information, we are also unable to differentiate such low-frequency variants

from sequencing error (0.1-1%).

Trait measurements and modeling

Despite initially setting up four replicates for each mitonuclear hybrid line, low productivity led

to extinction of several replicates, forcing us to proceed with one replicate per line to keep our

experiment balanced. We measured the hatching number for 182 females coming from 41

families reared at 15ºC and 216 females from 40 families reared at 25ºC. Male remating success

was measured for 491 males, using a total of 2,458 wild-caught females. Since 378 females died

over the course of the experiment, we discarded 31 males that had fewer than four out of six

females remaining in order to reduce variation associated with differential access to females.

Therefore, our male remating experiment is based on 235 males from 68 families reared at 15ºC

and 225 males from 76 families reared at 25ºC. The average male successfully copulated with

2.45 females in the span of 14 days when reared at 15□, declining to 2.04 females at 25□

representing a 17% reduction. In the average female clutch, 30.05 eggs hatched at 15□, declining

to 21.50 eggs hatching at 25☐ representing a 28% reduction. Female hatching number declined

an average of 31.16% in mitonuclear hybrid lines when females were bred with siblings as

opposed to wild-caught males (Document S1, Figure S1). Lifespan, frailty, and rate of

senescence was measured for 3,072 individuals, with 1,531 individuals from 89 families reared

at 15ºC, and 1,541 individuals from 77 families reared at 25ºC.

Experiments on early developmental stages indicate that mortality rate occurring before sexual

maturity was very high, ranging from 14.6% to 90.6% mortality depending on cross and

temperature (Document S1, Figure S2). Naupliar stages had the highest mortality, ranging from

4.2% to 84.4%, while copepodid stages ranged from 1.5% to 27.9% mortality. Linear mixed-

effects modeling showed that measurements describing the development rate varied significantly

with mtDNA x nDNA interactions and rearing temperature, with rates increasing at higher

temperature.

Sex-effect modeling

Model contrasts revealed mostly male-biased values for longevity (longer lifespan), and mixed

bias for fertility with three out of six significant contrasts being female biased (Figure 2A).

Model selection comparing the four sex-effects models (Table S1) revealed that models

including three-way sex x nDNA x mtDNA interaction term fit the trait data best in all

temperature and trait combinations except for relative fertility at 15☐, which was better fit with

sex x nDNA interactions (ΔAIC = 17.35). None of the sex-effects models uncovered significant

sex x mtDNA interactions (Table S2). However, we do see that sex x nDNA and sex x mtDNA x

nDNA interactions are a significant influence on fertility variation (Figure 2B).

Mitonuclear coevolutionary status modeling

Fertility was negatively impacted by mitonuclear status (matched vs. mismatched), but only in

females possessing the FHL mtDNA haplotype (females: $\chi2 = 17.90$, $p_{Bonf.} = 7.2e\text{-}05$; males: $\chi2$

$= 0.13$, $p_{Bonf.} = 1.0$), while no other trait was significant using the Bonferroni-corrected cutoff

(Table S3). The estimated marginal mean for matched female hatching number was 26.9

offspring, while for mismatched FHL females it was 16.5 offspring. Females with FHL mtDNA

also showed a trend of delayed development at the naupliar (females: $\chi2 = 5.20$, $p_{Bonf.} = 0.0675$;

males: $\chi2 = 1.16$, $p_{Bonf} = 1.0$) and copepodid (females: $\chi2 = 4.39$, $p_{Bonf} = 0.1086$; males: $\chi2$

$= 1.52$, $p_{Bonf} = 1.0$) stages, although these values were less than alpha $= 0.05$ and not the

Bonferroni-corrected alpha. The estimated marginal means for naupliar and copepodid

development rate in matched females were 8.68 and 12.1 days respectively, while the means for

mismatched FHL females were 12.08 and 14.9 days. Family sex ratio was significantly

influenced by mitonuclear status ($\chi^2 = 18.51$, $p = 1.7e\text{-}5$), as indicated in linear mixed-effects

modeling (Document S1, Figure S3), as well as nDNA x mitonuclear status interactions ($\chi^2 =$

28.77, $p = 8.1e\text{-}8$). Results using mortality-corrected family sex ratio data were also consistent

with uncorrected family sex ratio. (Table S4).

Best-fit models

Temperature significantly impacted all traits by reducing their average values (Table S5).

Notably, mean longevity was reduced by 102 days in 25ºC, an almost 10 day decrease per degree

reduction. Significant mtDNA effects were only seen in females for both hatching number ($\chi^2 =$

20.36, $p = 1.43e\text{-}4$) and longevity ($\chi^2 = 14.66$, $p = 0.0021$). Variation in nDNA impacted

longevity in both sexes (males: χ^2= 20.36, p = 6.42e-6; females: χ^2= 8.31, p = 0.0040) but

neither measure of fertility (female hatching number: χ^2= 1.37, p = 0.2416; male remating

success: χ^2= 0.15, p = 0.6991). Additionally, effects of nDNA were contingent upon rearing

temperature, with significant interactions between the two terms for male remating success (χ^2=

7.41, p = 0.0065), male longevity (χ^2= 4.19, p = 0.0407) and female longevity (χ^2= 4.05, p =

0.0442). Interactions between mtDNA and nDNA were also significant for longevity in females

(χ^2= 12.03, p = 0.0073) and in a three-way interaction with temperature for male remating

success (χ^2= 16.95, p = 7.22e-4). Mitochondrial effects impacted fertility more than longevity

(Figure 2C, p = 3.2e-4), while nuclear effects impacted both traits equally (p = 0.95) and

temperature affected longevity more than fertility (p = 1.5e-6). Linear mixed-effects modeling of

family sex ratio indicated significant influence of temperature (χ^2 = 10.02, p = 0.0015), mtDNA

haplotype (χ^2 = 28.35, p = 3.1e-6), as well as interactions between mtDNA and nDNA (χ^2 =

31.98, p =1.6e-8) and temperature and nDNA (χ^2 = 29.94, p = 1.4e-6; Document S1, Figure S3).

Parametric survivorship modeling

Survivorship curves indicate a drastic reduction in longevity shifting from 15□ to 25□, with SD

nDNA individuals having higher survivorship than BR nDNA individuals at 25□ (Figure 3A).

Additionally, we see that median longevity is sex-biased in SD nDNA individuals reared at 15□,

with higher male longevity (Figure 3A insert, Wilcoxon rank sum test, p = 0.004). Rearing

temperature affected both frailty (χ^2= 22.55, p = 2.05e-6) and rate of senescence (χ^2= 23.27, p =

1.41e-6) in a similar manner, with the higher temperature leading to increased frailty and

increased rate of senescence (Table S6). Mitochondrial haplotype showed no significant

contribution to Gompertz parameters when reared at either 15ºC or 25ºC (Table S6). We found

significant sexual dimorphism in frailty and senescence in individuals reared at 25ºC, but not

those reared at 15ºC (Figure 3B). When reared at 25ºC, females had lower frailty than males, but

a higher rate of senescence and while this pattern held in both nDNA backgrounds, it was

nonsignificant in BR nDNA lines (Table S6).

Mitochondrial coefficient of variation

Females had significantly higher CV_{mt} than males in seven out of eight experimental groupings

(Figure 4A,Table S7). The median difference between male and female coefficients in these

groupings ranged from 0.0039 to 0.1061. In contrast, fertility in BR nDNA individuals at 25ºC

showed higher male CV_{mt} than females (median difference = -0.0993, CI = (-0.11, -0.09), $p <$

2.2e-16).

Genetic correlation among haplotypes

Intersexual correlations for median longevity ($r = 0.99$; $CI_{95} = 0.20, 2.00$) and fertility ($r = 0.84$;

$CI_{95} = 0.08, 1.72$) were significantly positive in BR nDNA individuals reared at 25ºC, while

correlations for median longevity in SD nDNA individuals were also significantly positive ($r =$

0.85; $CI_{95} = 0.04, 1.75$) in individuals reared at 25ºC (Figure 4B). Intrasexual correlations

between fertility and longevity were significantly positive in BR nDNA individuals of both sexes

reared at 25ºC (males: $r = 0.96$; $CI_{95} = 0.21, 1.95$; females: $r = 0.90$; $CI_{95} = 0.08, 1.86$). No

measures of intersexual or intrasexual correlation were significantly negative.

Discussion

Our study sought to explicitly test three predictions arising from the mother's curse hypothesis: higher mitochondrial genetic variance in males for fitness traits, negative intersexual correlation among mitochondrial haplotypes for fitness traits, and higher impact of experimental disruption of putatively coevolved mitonuclear genotypes on males. Our results did not support these predictions. Females had a higher level of mitochondrial genetic variance in seven experimental groupings and lower variance in one grouping (fertility in individuals with BR nDNA reared at 25ºC). We also found no evidence for sexual antagonism for fertility or longevity, with negative intersexual correlations found in two of eight experimental groupings (fertility in BR nDNA at 15□ and SD nDNA at 25□) having upper confidence intervals that extended well into positive ranges (0.90 and 1.41, respectively). Overall, the intersexual comparisons mimicked the intrasexual comparisons. Lastly, we found a significant negative influence of coevolutionary status on female fertility in hybrids with the FHL mtDNA, but not for male fertility contrary to the predictions of the mother's curse hypothesis. Female hybrids with FHL mtDNA also show a trend of delayed larval development, while males do not. The FHL mtDNA is the most divergent haplotype ($\geq$ 20.6% nucleotide divergence) while the remaining haplotypes are 0.7 - 9.6% divergence).

Detecting sex-bias in mitochondrial coefficient of variation for fitness traits is arguably the most direct test of the hypothesis since it emphasizes the differential accumulation of mutational variation between the sexes[82]. According to the mother's curse hypothesis, mitochondrial mutations causing male infertility that are neutral in females are more likely to spread in a population through random drift since males are a dead-end for natural selection on mtDNA.

However, this argument is best applied to randomly mating populations with no inbreeding. In

populations with even modest levels of inbreeding, such as those with geographical population

structure, females will also suffer fitness insults from mother's curse mutations to a certain

degree since they are more likely to mate with sub-fertile siblings[32,33,83]. The degree to

which male subfertility influences female fertility fitness can also influence these dynamics. In

monandrous species like *Tigriopus*, females mate only once[84,85], resulting in a direct

relationship between male fertility and female offspring production. In this study, we showed

that female hatching number dropped substantially when mated with siblings compared to wild-

caught males, indicating the influence of male subfertility on female fertility. However, in

mating systems where females mate with several males (e.g. *Drosophila*), females can be

safeguarded against male infertility or subfertility[86]. As a result, selection to purge male-

harming mtDNA mutations is expected to be strongest in species with monandry and weakest

with polyandry whenever inbreeding is present[83]. Studies show that *T. californicus* avoid close

inbreeding[87], but live in highly isolated, ephemeral populations making inbreeding all but

inevitable[88,89]. Additionally, while inbreeding can limit mother's curse by decreasing the

conditions under which male-harming alleles can spread, it can also result in the spread of

female-harming alleles that benefit males and lead to female-biased fitness variance [27,32]. The

widespread female-biased fitness variance observed in our fertility and longevity experiments

may therefore indicate the impact of historical inbreeding on the fixation of mitochondrial

divergence sampled in our study.

Another factor that has been proposed to limit the spread of male-harming mtDNA mutations is

paternal leakage of mtDNA, although we do not believe it played a major role in our results. We

found a very low level of mtDNA reads of potential paternal origin (0.02% to 0.28%), and these

could not be cleanly distinguished from numts or sequencing error. Haploid models of

cytoplasmic inheritance suggest that small amounts of paternal leakage are not sufficient to

rescue mother's curse[90], with indirect selection on human males via inbreeding estimated to be

up to 370 times stronger than direct selection on sperm performance due to paternal leakage [32].

Indeed, low levels of paternal mtDNA leakage do not appear to reliably cure mother's curse,

since levels of leakage higher than those reported in *Tigriopus* [80] are common in *Drosophila*

crosses[91,92], where the strongest (albeit inconsistent) support for mother's curse has been

found.

We predicted that sex differences due to mitochondrial haplotype effects are greater for fertility

than longevity, based on the expectation that fertility genes are more sex-specific than viability

genes[93,94]. We also predicted that the consequences of mitonuclear mismatch should increase

with metabolic rate associated with increased temperature. Efficient energy production depends

on mitochondrial function, including mitonuclear coordination. Mild mitonuclear mismatches

may be tolerated under low aerobic demand and yet become lethal when energetic demand

become higher [40,but see 82]. Increased temperature reduced measures of both fertility and longevity

but was more dramatic with the latter. We uncovered significant mitonuclear effects impacting

longevity at 25□ but not 15□. Additionally, we found that mitonuclear x sex effects were

significant for fertility at 25□, but not 15□. Mitochondrial haplotype effects in our experiment

impacted fertility more than longevity, however experiments on juvenile stages indicate a

substantial amount of mortality, influenced by temperature and mitonuclear interactions,

occurring before sexual maturity. Because *T. californicus* sperm are amotile, declines in male

fertility are not attributed to loss of sperm motility – a key marker of male sterility in organisms

with flagellated, motile sperm. Instead, declines in male fertility in our study are likely due to the

number of mature spermatids in a spermatophore, the ability of sperm to maintain homeostasis

while residing in the female reproductive tract, or behavior. High juvenile mortality may also

impact our study as it could produce indirect selection on the adult traits under study, although

any juvenile mortality precludes analysis of sex differences. Mortality occurring during the

juvenile stages could not be assigned to either sex, as it is difficult to distinguish the sex of

Tigriopus californicus juveniles before sexual maturity. However, while family sex ratio

differed among crosses and by temperature, most crosses indicate male-biased sex ratios, which

may indicate either increased mortality of females during the larval stages or the misexpression

of sex determining loci in mitonuclear hybrids.

Survivorship analysis indicates that mtDNA variation had an inverse impact on frailty and

senescence between the sexes, similar to the 'Strehler-Mildvan' (S-M) relationship[96]. When

reared at 25ºC, females had lower frailty than males while also having a higher rate of

senescence, despite having similar median longevity. One explanation for this relationship is that

the preferential survival of the most-frail female *T. californicus* may lead to an increase in the

rate of senescence for females, since frail individuals are more susceptible to aging. Male *T.*

californicus typically have lower tolerance than females to a variety of stressors, including

temperature, and a greater transcriptomic response to oxidative stress[56,57,59,97,98].

Conclusion

Like previous work[99,100], we found that the functional mtDNA effects were highly context-

dependent, being influenced by the nuclear-background in which they were expressed, the

temperature in which individuals developed, and/or the sex of the individual. We explicitly

sought to test the main predictions of the mother's curse hypothesis, finding patterns opposite to

those predictions. Support for the mother's curse hypothesis has been highly mixed to date, both

theoretically and empirically, with many studies rejecting these predictions. While many studies

directly testing predictions of the mother's curse hypothesis have used *Drosophila*, the

predominance of maternal inheritance of mitochondria in eukaryotes allows for a taxonomically

diverse investigation of the generality of mother's curse. Particularly, diversity in mating system

should be considered for future studies, as the fitness of females that mate with only one male

will be strongly determined by the fitness of her mate when there is inbreeding. Furthermore,

sperm diversity (motile/amotile, storage/remating) should be highlighted in order to address the

relevance of various mutational insults to male fitness potentially harbored by mitochondrial

DNA. In light of the evidence presented here and elsewhere, we suggest that accumulation of

male-harming mitochondrial mutations may not be a generalizable phenomenon, despite the

prevalence of maternal mitochondrial transmission across eukaryotes.

Acknowledgments

We are thankful to MacKenzie Partridge, Manuel Morales and Mi Rae Park for their help during

the experiments and population maintenance. We also thank Ning Li and Scott Applebaum for

their constructive criticisms and suggestions, which greatly improved the quality of the

manuscript. Comments provided by anonymous reviewers also aided in improving the

manuscript. The experiments reported here were supported by a grant (#1656048) from the

National Science Foundation, Division of Environmental Biology, awarded to SE.

Declaration of interests

The authors declare no competing interests.

References

1. Yang D, Oyaizu Y, Oyaizu H, Olsen GJ, Woese CR. 1985 Mitochondrial origins. *PNAS* **82**, 4443–
4447. (doi:10.1073/pnas.82.13.4443)

2. Gray MW, Burger G, Lang BF. 1999 Mitochondrial Evolution. *Science* **283**, 1476–1481.
(doi:10.1126/science.283.5407.1476)

3. Hedges SB, Blair JE, Venturi ML, Shoe JL. 2004 A molecular timescale of eukaryote evolution
and the rise of complex multicellular life. *BMC Evol Biol* **4**, 2. (doi:10.1186/1471-2148-4-2)

4. McBride HM, Neuspiel M, Wasiak S. 2006 Mitochondria: More Than Just a Powerhouse. *Curr.*
Biol. **16**, R551–R560. (doi:10.1016/j.cub.2006.06.054)

5. Pagliarini DJ, Rutter J. 2013 Hallmarks of a new era in mitochondrial biochemistry. *Genes*
Dev. **27**, 2615–2627. (doi:10.1101/gad.229724.113)

6. Chandel NS. 2014 Mitochondria as signaling organelles. *BMC Biol* **12**, 34. (doi:10.1186/1741-
7007-12-34)

7. Birsoy K, Wang T, Chen WW, Freinkman E, Abu-Remaileh M, Sabatini DM. 2015 An Essential
Role of the Mitochondrial Electron Transport Chain in Cell Proliferation Is to Enable
Aspartate Synthesis. *Cell* **162**, 540–551. (doi:10.1016/j.cell.2015.07.016)

8. Santos JH. 2021 Mitochondria signaling to the epigenome: A novel role for an old organelle.
Free Radical Biol. Med. **170**, 59–69. (doi:10.1016/j.freeradbiomed.2020.11.016)

9. Bar-Yaacov D, Blumberg A, Mishmar D. 2012 Mitochondrial-nuclear co-evolution and its
effects on OXPHOS activity and regulation. *Biochim. Biophys. Acta* **1819**, 1107–1111.
(doi:10.1016/j.bbagrm.2011.10.008)

10. Lotz C *et al.* 2014 Characterization, design, and function of the mitochondrial proteome:
from organs to organisms. *J. Proteome Res.* **13**, 433–446. (doi:10.1021/pr400539j)

11. Brown WM, George M, Wilson AC. 1979 Rapid evolution of animal mitochondrial DNA.
PNAS **76**, 1967–1971. (doi:10.1073/pnas.76.4.1967)

12. Lynch M. 2010 Rate, molecular spectrum, and consequences of human mutation. *PNAS* **107**,
961–968. (doi:10.1073/pnas.0912629107)

13. Lynch M, Blanchard JL. 1998 Deleterious mutation accumulation in organelle genomes.
Genetics **102/103**, 29–39. (doi:10.1023/A:1017022522486)

14. Cooper BS, Burrus CR, Ji C, Hahn MW, Montooth KL. 2015 Similar Efficacies of Selection
Shape Mitochondrial and Nuclear Genes in Both *Drosophila melanogaster* and *Homo*
sapiens. *G3-Genes Genom Genet* **5**, 2165–2176. (doi:10.1534/g3.114.016493)

15. Christie JR, Beekman M. 2016 Uniparental Inheritance Promotes Adaptive Evolution in
Cytoplasmic Genomes. *Mol. Biol. Evol.* , msw266. (doi:10.1093/molbev/msw266)

16. Christie JR, Beekman M. 2017 Selective sweeps of mitochondrial DNA can drive the
evolution of uniparental inheritance. *Evolution* **71**, 2090–2099. (doi:10.1111/evo.13291)

17. Radzvilavicius AL, Kokko H, Christie JR. 2017 Mitigating Mitochondrial Genome Erosion
Without Recombination. *Genetics* **207**, 1079–1088. (doi:10.1534/genetics.117.300273)

18. Edwards DM, Røyrvik EC, Chustecki JM, Giannakis K, Glastad RC, Radzvilavicius AL, Johnston
IG. 2021 Avoiding organelle mutational meltdown across eukaryotes with or without a
germline bottleneck. *PLOS Biology* **19**, e3001153. (doi:10.1371/journal.pbio.3001153)

19. Schmidt TR, Wu W, Goodman M, Grossman LI. 2001 Evolution of nuclear-and
mitochondrial-encoded subunit interaction in cytochrome c oxidase. *Mol. Biol. Evol.* **18**,
563–569. (doi:10.1093/oxfordjournals.molbev.a003836)

20. Willett CS, Burton RS. 2004 Evolution of Interacting Proteins in the Mitochondrial Electron
Transport System in a Marine Copepod. *Mol Biol Evol* **21**, 443–453.
(doi:10.1093/molbev/msh031)

21. Bazin E, Glémin S, Galtier N. 2006 Population Size Does Not Influence Mitochondrial Genetic
Diversity in Animals. *Science* **312**, 570–572. (doi:10.1126/science.1122033)

22. James JE, Piganeau G, Eyre-Walker A. 2016 The rate of adaptive evolution in animal
mitochondria. *Mol. Ecol.* **25**, 67–78. (doi:10.1111/mec.13475)

23. Barreto FS, Watson ET, Lima TG, Willett CS, Edmands S, Li W, Burton RS. 2018 Genomic
signatures of mitonuclear coevolution across populations of *Tigriopus californicus*. *Nat.*
Ecol. Evol. **2**, 1250–1257. (doi:10.1038/s41559-018-0588-1)

24. Hood WR *et al.* 2018 The Mitochondrial Contribution to Animal Performance, Adaptation,
and Life-History Variation. *Integr. Comp. Biol.* **58**, 480–485. (doi:10.1093/icb/icy089)

25. Wolstenholme DR. 1992 Animal Mitochondrial DNA: Structure and Evolution. In
International Review of Cytology (eds DR Wolstenholme, KW Jeon), pp. 173–216. Academic
Press. (doi:10.1016/S0074-7696(08)62066-5)

26. Frank SA, Hurst LD. 1996 Mitochondria and male disease. *Nature* **383**, 224–224.
(doi:10.1038/383224a0)

27. Smith SRT, Connallon T. 2017 The contribution of the mitochondrial genome to sex-specific
fitness variance. *Evolution* **71**, 1417–1424. (doi:10.1111/evo.13238)

28. Gemmell NJ, Metcalf VJ, Allendorf FW. 2004 Mother's curse: the effect of mtDNA on
individual fitness and population viability. *TREE* **19**, 238–244.
(doi:10.1016/j.tree.2004.02.002)

29. Cardullo RA, Baltz JM. 1991 Metabolic regulation in mammalian sperm: Mitochondrial
volume determines sperm length and flagellar beat frequency. *Cell Motility* **19**, 180–188.
(doi:10.1002/cm.970190306)

30. Holyoake AJ, McHugh P, Wu M, O'Carroll S, Benny P, Sin IL, Sin FYT. 2001 High incidence of
single nucleotide substitutions in the mitochondrial genome is associated with poor semen
parameters in men. *Int. J. Androl.* **24**, 175–182. (doi:10.1046/j.1365-2605.2001.00292.x)

31. Camus MF, Clancy DJ, Dowling DK. 2012 Mitochondria, Maternal Inheritance, and Male
Aging. *Curr. Biol.* **22**, 1717–1721. (doi:10.1016/j.cub.2012.07.018)

32. Wade MJ, Brandvain Y. 2009 Reversing Mother's Curse: Selection on Male Mitochondrial
Fitness Effects. *Evolution* **63**, 1084–1089. (doi:10.1111/j.1558-5646.2009.00614.x)

33. Unckless RL, Herren JK. 2009 Population genetics of sexually antagonistic mitochondrial
mutants under inbreeding. *J. Theor. Biol.* **260**, 132–136. (doi:10.1016/j.jtbi.2009.06.004)

34. Hedrick PW. 2012 Reversing Mother's Curse Revisited. *Evolution* **66**, 612–616.
(doi:10.1111/j.1558-5646.2011.01465.x)

35. Chase CD. 2007 Cytoplasmic male sterility: a window to the world of plant mitochondrial–
nuclear interactions. *TRIG* **23**, 81–90. (doi:10.1016/j.tig.2006.12.004)

36. Innocenti P, Morrow EH, Dowling DK. 2011 Experimental Evidence Supports a Sex-Specific
Selective Sieve in Mitochondrial Genome Evolution. *Science* **332**, 845–848.
(doi:10.1126/science.1201157)

37. Camus MF, Wolf JBW, Morrow EH, Dowling DK. 2015 Single Nucleotides in the mtDNA
Sequence Modify Mitochondrial Molecular Function and Are Associated with Sex-Specific
Effects on Fertility and Aging. *Curr Biol* **25**, 2717–2722. (doi:10.1016/j.cub.2015.09.012)

38. Camus MF, Dowling DK. 2018 Mitochondrial genetic effects on reproductive success:
signatures of positive intrasexual, but negative intersexual pleiotropy. *Proc. R. Soc. B* **285**,
10. (doi:10.1098/rspb.2018.0187)

39. Wolff JN, Pichaud N, Camus MF, Côté G, Blier PU, Dowling DK. 2016 Evolutionary
implications of mitochondrial genetic variation: mitochondrial genetic effects on OXPHOS
respiration and mitochondrial quantity change with age and sex in fruit flies. *J. Evol. Biol.*
29, 736–747. (doi:10.1111/jeb.12822)

40. Carnegie L, Reuter M, Fowler K, Lane N, Camus MF. 2021 Mother's curse is pervasive across
a large mitonuclear *Drosophila* panel. *Evol. Lett.* **5**, 230–239. (doi:10.1002/evl3.221)

41. Mossman JA, Tross JG, Li N, Wu Z, Rand DM. 2016 Mitochondrial-Nuclear Interactions
Mediate Sex-Specific Transcriptional Profiles in *Drosophila*. *Genetics* **204**, 613–630.
(doi:10.1534/genetics.116.192328)

42. Mossman JA, Biancani LM, Rand DM. 2016 Mitonuclear Epistasis for Development Time and
its Modification by Diet in *Drosophila*. *Genetics* **203**, 463–484.
(doi:10.1534/genetics.116.187286)

43. Immonen E, Collet M, Goenaga J, Arnqvist G. 2016 Direct and indirect genetic effects of sex-
specific mitonuclear epistasis on reproductive ageing. *Heredity* **116**, 338–339.
(doi:10.1038/hdy.2015.112)

44. Đorđević M, Stojković B, Savković U, Immonen E, Tucić N, Lazarević J, Arnqvist G. 2017 Sex-
specific mitonuclear epistasis and the evolution of mitochondrial bioenergetics, ageing, and
life history in seed beetles: SEX-SPECIFIC MITONUCLEAR EPISTASIS. *Evolution* **71**, 274–288.
(doi:10.1111/evo.13109)

45. Kumar S, Suleski M, Craig JM, Kasprowicz AE, Sanderford M, Li M, Stecher G, Hedges SB.
2022 TimeTree 5: An Expanded Resource for Species Divergence Times. *Mol. Biol. Evol.* **39**,
msac174. (doi:10.1093/molbev/msac174)

46. Ashburner M, Golic KG, Hawley RS, others. 2004 *Drosophila: a laboratory handbook.* Cold
spring harbor laboratory press.

47. Lane N. 2011 The Costs of Breathing. *Science* **334**, 184–185. (doi:10.1126/science.1214012)

48. Burton RS, Feldman MW, Curtsinger JW. 1979 Population genetics of *Tigriopus californicus*
(Copepoda: Harpacticoida): I. Population structure along the central California coast.
Marine Ecology Progress Series , 29–39. (doi:10.3354/meps001029)

49. Burton RS. 1998 Intraspecific Phylogeography Across the Point Conception Biogeographic
Boundary. *Evolution* **52**, 734–745. (doi:10.1111/j.1558-5646.1998.tb03698.x)

50. Edmands S. 2001 Phylogeography of the intertidal copepod *Tigriopus californicus* reveals
substantially reduced population differentiation at northern latitudes. *Mol. Ecol.* **10**, 1743–
1750. (doi:10.1046/j.0962-1083.2001.01306.x)

51. Edmands S. 1999 Heterosis and Outbreeding Depression in Interpopulation Crosses
Spanning a Wide Range of Divergence. *Evolution* **53**, 1757–1768. (doi:10.1111/j.1558-
5646.1999.tb04560.x)

52. Pritchard VL, Edmands S. 2013 The Genomic Trajectory of Hybrid Swarms: Outcomes of
Repeated Crosses Between Populations of *Tigriopus californicus*. *Evolution* **67**, 774–791.
(doi:10.1111/j.1558-5646.2012.01814.x)

53. Voordouw MJ, Anholt BR. 2002 Heritability of sex tendency in a harpacticoid copepod,
Tigriopus californicus. *Evolution* **56**, 1754–1763. (doi:10.1111/j.0014-3820.2002.tb00189.x)

54. Foley BR, Rose CG, Rundle DE, Leong W, Edmands S. 2013 Postzygotic isolation involves
strong mitochondrial and sex-specific effects in *Tigriopus californicus*, a species lacking
heteromorphic sex chromosomes. *Heredity* **111**, 391–401. (doi:10.1038/hdy.2013.61)

55. Alexander HJ, Richardson JML, Edmands S, Anholt BR. 2015 Sex without sex chromosomes:
genetic architecture of multiple loci independently segregating to determine sex ratios in
the copepod *Tigriopus californicus*. *J. Evol. Biol.* **28**, 2196–2207. (doi:10.1111/jeb.12743)

56. Willett CS. 2010 Potential Fitness Trade-Offs for Thermal Tolerance in the Intertidal
Copepod *Tigriopus Californicus*. *Evolution* **64**, 2521–2534. (doi:10.1111/j.1558-
5646.2010.01008.x)

57. Kelly MW, Sanford E, Grosberg RK. 2012 Limited potential for adaptation to climate change
in a broadly distributed marine crustacean. *Proc Biol Sci* **279**, 349–356.
(doi:10.1098/rspb.2011.0542)

58. Leong W, Sun PY, Edmands S. 2018 Latitudinal Clines in Temperature and Salinity Tolerance
in Tidepool Copepods. *J. Hered.* **109**, 71–77. (doi:10.1093/jhered/esx061)

59. Foley HB, Sun PY, Ramirez R, So BK, Venkataraman YR, Nixon EN, Davies KJA, Edmands S.
2019 Sex-specific stress tolerance, proteolysis, and lifespan in the invertebrate *Tigriopus*
californicus. *Exp. Gerontol.* **119**, 146–156. (doi:10.1016/j.exger.2019.02.006)

60. Bartoń K. 2020 *MuMIn: Multi-Model Inference*. See https://CRAN.R-
project.org/package=MuMIn.

61. Lenth RV. 2020 *emmeans: Estimated Marginal Means, aka Least-Squares Means*. See
https://CRAN.R-project.org/package=emmeans.

62. Bates D, Mächler M, Bolker B, Walker S. 2015 Fitting Linear Mixed-Effects Models Using
lme4. *J. Stat. Softw.* **67**, 1–48. (doi:10.18637/jss.v067.i01)

63. Fox J, Weisberg S. 2019 *An R Companion to Applied Regression*. Third. Thousand Oaks CA:
Sage. See https://socialsciences.mcmaster.ca/jfox/Books/Companion/.

64. Gompertz B. 1833 On the Nature of the Function Expressive of the Law of Human Mortality,
and on a New Mode of Determining the Value of Life Contingencies. *Proc. R. Soc. Lond.* **2**.
(doi:10.1098/rspl.1815.0271)

65. Jackson C. 2016 flexsurv: A Platform for Parametric Survival Modeling in R. *J. Stat. Softw.* **70**,
1–33. (doi:10.18637/jss.v070.i08)

66. R Core Team. 2020 *R: A Language and Environment for Statistical Computing*. Vienna,
Austria: R Foundation for Statistical Computing. See https://www.R-project.org/.

67. Canty A, Ripley BD. 2020 *boot: Bootstrap R (S-Plus) Functions*.

68. Davison AC, Hinkley DV. 1997 *Bootstrap Methods and Their Applications*. Cambridge:
Cambridge University Press. See http://statwww.epfl.ch/davison/BMA/.

69. Arnholt AT, Evans B. 2017 *BSDA: Basic Statistics and Data Analysis*. See https://CRAN.R-
project.org/package=BSDA.

70. Bolger AM, Lohse M, Usadel B. 2014 Trimmomatic: a flexible trimmer for Illumina sequence
data. *Bioinformatics* **30**, 2114–2120. (doi:10.1093/bioinformatics/btu170)

71. Dierckxsens N, Mardulyn P, Smits G. 2017 NOVOPlasty: de novo assembly of organelle
genomes from whole genome data. *Nucleic Acids Res.* **45**, e18. (doi:10.1093/nar/gkw955)

72. Bernt M, Donath A, Jühling F, Externbrink F, Florentz C, Fritzsch G, Pütz J, Middendorf M,
Stadler PF. 2013 MITOS: Improved de novo metazoan mitochondrial genome annotation.
Mol. Phylogenet. Evol. **69**, 313–319. (doi:10.1016/j.ympev.2012.08.023)

73. Lunter G, Goodson M. 2011 Stampy: A statistical algorithm for sensitive and fast mapping of
Illumina sequence reads. *Genome Res.* **21**, 936–939. (doi:10.1101/gr.111120.110)

74. In press. Picard Tools - By Broad Institute. See https://broadinstitute.github.io/picard/
(accessed on 24 October 2021).

75. Li H. 2011 A statistical framework for SNP calling, mutation discovery, association mapping
and population genetical parameter estimation from sequencing data. *Bioinformatics* **27**,
2987–2993. (doi:10.1093/bioinformatics/btr509)

76. Darling AE, Mau B, Perna NT. 2010 progressiveMauve: Multiple Genome Alignment with
Gene Gain, Loss and Rearrangement. *PLoS ONE* **5**, e11147.
(doi:10.1371/journal.pone.0011147)

77. Ranwez V, Douzery EJP, Cambon C, Chantret N, Delsuc F. 2018 MACSE v2: Toolkit for the
Alignment of Coding Sequences Accounting for Frameshifts and Stop Codons. *Mol Biol Evol*
35, 2582–2584. (doi:10.1093/molbev/msy159)

78. Larkin MA *et al.* 2007 Clustal W and Clustal X version 2.0. *Bioinformatics* **23**, 2947–2948.
(doi:10.1093/bioinformatics/btm404)

79. Yang Z. 2007 PAML 4: Phylogenetic Analysis by Maximum Likelihood. *Mol Biol Evol* **24**,
1586–1591. (doi:10.1093/molbev/msm088)

80. Kofler R, Orozco-terWengel P, De Maio N, Pandey RV, Nolte V, Futschik A, Kosiol C,
Schlötterer C. 2011 PoPoolation: A Toolbox for Population Genetic Analysis of Next
Generation Sequencing Data from Pooled Individuals. *PLoS ONE* **6**.
(doi:10.1371/journal.pone.0015925)

81. Lee J, Willett CS. 2022 Frequent Paternal Mitochondrial Inheritance and Rapid Haplotype
Frequency Shifts in Copepod Hybrids. *J. Hered.* **113**, 171–183. (doi:10.1093/jhered/esab068)

82. Dowling DK, Adrian RE. 2019 Challenges and Prospects for Testing the Mother's Curse
Hypothesis. *Integr. Comp. Biol.* **59**, 875–889. (doi:10.1093/icb/icz110)

83. Zhang H, Guillaume F, Engelstädter J. 2012 The Dynamics of Mitochondrial Mutations
Causing Male Infertility in Spatially Structured Populations. *Evolution* **66**, 3179–3188.
(doi:10.1111/j.1558-5646.2012.01675.x)

84. Vittor BA. 1971 Effects of the environment of fitness-related life history characters in
Tigriopus californicus. Doctoral dissertation, University of Oregon.

85. Burton RS. 1985 Mating system of the intertidal copepod *Tigriopus californicus*. *Marine*
Biology **86**, 247–252. (doi:10.1007/BF00397511)

86. Singh SR, Singh BN, Hoenigsberg HF. 2002 Female remating, sperm competition and sexual
selection in *Drosophila*. *Genet Mol Res* **1**, 178–215.

87. Palmer CA, Edmands S. 2000 Mate choice in the face of both inbreeding and outbreeding
depression in the intertidal copepod *Tigriopus californicus*. *Marine Biology* **136**, 693–698.
(doi:10.1007/s002270050729)

88. Burton RS. 1997 Genetic Evidence for Long Term Persistence of Marine Invertebrate
Populations in an Ephemeral Environment. *Evolution* **51**, 993. (doi:10.2307/2411174)

89. Dybdahl MF. 1994 Extinction, recolonization, and the genetic structure of tidepool copepod
populations. *Evol Ecol* **8**, 113–124. (doi:10.1007/BF01238245)

90. Wade MJ. 2014 Paradox of Mother's Curse and the Maternally Provisioned Offspring
Microbiome. *Cold Spring Harbor Perspectives in Biology* **6**, a017541.

91. Sherengul W, Kondo R, Matsuura ET. 2006 Analysis of paternal transmission of
mitochondrial DNA in *Drosophila*. *Genes Genet Syst* **81**, 399–404. (doi:10.1266/ggs.81.399)

92. Wolff JN, Nafisinia M, Sutovsky P, Ballard JWO. 2013 Paternal transmission of mitochondrial
DNA as an integral part of mitochondrial inheritance in metapopulations of *Drosophila*
simulans. *Heredity* **110**, 57–62. (doi:10.1038/hdy.2012.60)

93. Orr HA. 1989 Genetics of sterility in hybrids between two subspecies of *Drosophila*.
Evolution **43**, 180–189. (doi:10.1111/j.1558-5646.1989.tb04216.x)

94. Turelli M, Orr HA. 2000 Dominance, Epistasis, and the Genetics of Postzygotic Isolation.
Genetics **154**, 1663–1679. (doi:10.1093/genetics/154.4.1663)

95. Pichaud N, Ballard JWO, Tanguay RM, Blier PU. 2013 Mitochondrial haplotype divergences
affect specific temperature sensitivity of mitochondrial respiration. *J Bioenerg Biomembr*
45, 25–35. (doi:10.1007/s10863-012-9473-9)

96. Strehler BL, Mildvan AS. 1960 General theory of mortality and aging: A stochastic model
relates observations on aging, physiologic decline, mortality, and radiation. *Science* **132**, 14–
21. (doi:10.1126/science.132.3418.14)

97. Li N, Arief N, Edmands S. 2019 Effects of oxidative stress on sex-specific gene expression in
the copepod *Tigriopus californicus* revealed by single individual RNA-seq. *Comp. Biochem.*
Physiol. Part D Genomics **31**, 100608. (doi:10.1016/j.cbd.2019.100608)

98. Li N, Flanagan BA, Partridge M, Huang EJ, Edmands S. 2020 Sex differences in early
transcriptomic responses to oxidative stress in the copepod *Tigriopus californicus*. *BMC*
Genomics **21**, 759. (doi:10.1186/s12864-020-07179-5)

99. Montooth KL, Dhawanjewar AS, Meiklejohn CD. 2019 Temperature-Sensitive Reproduction
and the Physiological and Evolutionary Potential for Mother's Curse. *Integr. Comp. Biol.* **59**,
890–899. (doi:10.1093/icb/icz091)

100.   Rand DM, Mossman JA. 2020 Mitonuclear conflict and cooperation govern the
integration of genotypes, phenotypes and environments. *Phil. Trans. R. Soc. B* **375**,
20190188. (doi:10.1098/rstb.2019.0188)

SUPPLEMENTAL INFORMATION

Supplemental Information includes one document, three figures, and seven tables and can be

found online.

Backcross	Dam line	Sire line	Nuclear F_{ST}	mtDNA D_{XY}	mtDNA d_N	mtDNA d_S	% Paternal nDNA	% Maternal mtDNA	% Paternal mtDNA
SDxSD	SD	SD	0	0	0	0	100.00	94.09	94.09
BRxSD	BR	SD	0.102	0.096	0.023	0.532	99.90	97.22	0.28
LJxSD	LJ	SD	0.108	0.096	0.023	0.523	99.97	97.63	0.15
FHLxSD	FHL	SD	0.378	0.206	0.085	2.675	99.70	99.65	0.13
BRxBR	BR	BR	0	0	0	0	100.00	96.91	96.91
LJxBR	LJ	BR	0.049	0.007	0.002	0.046	99.93	95.96	0.08
SDxBR	SD	BR	0.102	0.096	0.023	0.532	99.72	95.32	0.02
FHLxBR	FHL	BR	0.369	0.211	0.085	4.347	99.15	99.81	0.02

Figure 1. Construction of mitonuclear hybrid panel

(A) Geographic location of populations used in this study – SD (San Diego, CA), BR (Bird Rock, CA), LJ (La Jolla, CA), FHL (Friday Harbor, WA). (B) Experimental design. Dams from the central panel were backcrossed to sires from BR and SD for 10 generations. Grey arrows indicate experimental control lines. Individuals from each line were reared at two contrasting temperatures, 15℃ and 25℃. (C) DNA sequencing results from pool-seq. Measures of divergence (F_{ST}, D_{XY}, d_N, and d_S) were calculated between dam and sire line. Percent paternal nDNA and percent maternal/paternal mtDNA indicate minimal contaminating nDNA remaining in backcrossed individuals and the absence of paternal leakage of mtDNA.

A

Model contrast (female − male), dotted line = 0

B

Trait	Temp.	Rank	Model	df	AICc	ΔAIC	weight
Fertility	15℃	1	**Sex x nDNA** + mtDNA	9	638.6	0.00	0.934
		2	**Sex** + mtDNA + nDNA	8	644.0	5.46	0.061
		3	**Sex x mtDNA** + nDNA	11	649.1	10.53	0.005
		4	**Sex x mtDNA x nDNA**	18	655.9	17.35	0.000
	25℃	1	**Sex x mtDNA x nDNA**	18	1042.0	0.00	0.991
		2	**Sex x nDNA** + mtDNA	9	1052.3	10.30	0.006
		3	**Sex** + mtDNA + nDNA	8	1053.4	11.37	0.003
		4	**Sex x mtDNA** + nDNA	11	1061.6	19.64	0.000
Longevity	15℃	1	**Sex x mtDNA x nDNA**	18	15907.2	0.00	1.000
		2	**Sex x mtDNA** + nDNA	11	15945.0	37.84	0.000
		3	**Sex x nDNA** + mtDNA	9	15952.6	45.41	0.000
		4	**Sex** + mtDNA + nDNA	8	15956.1	48.93	0.000
	25℃	1	**Sex x mtDNA x nDNA**	18	12923.6	0.00	1.000
		2	**Sex x nDNA** + mtDNA	9	12959.8	36.20	0.000
		3	**Sex x mtDNA** + nDNA	11	12960.1	36.49	0.000
		4	**Sex** + mtDNA + nDNA	8	12965.7	42.05	0.000

C

Figure 2. Trait modeling results.

(**A**) Model contrasts between the sexes for the four sex-effect models. Models were performed separately for each temperature. Model contrasts for the "Sex" model were summed over all mtDNA and nDNA sources, "Sex x nDNA" contrasts were summed over mtDNA, and "Sex x mtDNA" contrasts were summed over nDNA. Circles represent sex contrasts for fertility and triangles represent longevity. The vertical dotted line represents contrasts (female – male) = 0.0, indicating identical estimated marginal means. Contrasts with 95% confidence intervals that fall on the left side are negative and thus significantly male-biased and coloured blue, while those falling on the right side are positive, female-biased, and coloured red. (**B**) Sex-effect model type III ANOVA table. Each row represents a separate model, with columns representing modeled fixed effects. Cells are marked with a dash when a variable is not included in the model. Abbreviations: NS – non significant (p <0.05), ✱ – p ≤ 0.05, ✱✱ – p ≤ 0.01, ✱✱✱– p ≤ 0.001. (**C**)Genetic and temperature effect size comparisons between fitness traits. We estimated Cohen's *d* for all pairwise contrasts among temperature, mtDNA, and nDNA groupings. Contrasts were made on best fit linear mixed effects models for each sex and trait grouping (sex-specific fertility, longevity). Significance was determined using two-sample unpaired Wilcoxon rank sum tests with a two-sided alternate hypothesis and 95% confidence intervals. Sample sizes for contrasts were mtDNA, n = 96; nDNA, n = 32; and Temperature, n = 32.

Figure 3. Males outlive females and have lower rates of senescence and higher frailty.

(A) Sex-specific survivorship curves based on nDNA and temperature groupings. Median longevity shown in insert with p-values indicating results of two-sample unpaired Wilcoxon rank sum tests with a two-sided alternate hypothesis and 95% confidence intervals.

(B) Sex contrasts for Gompertz parameters. We performed Wilcoxon tests as above to identify parametric differences in the rate of aging between the sexes.

Figure 4. Testing the mother's curse hypothesis with mitonuclear hybrid _T. californicus._

(A) Sex differences in mitochondrial coefficient of variation (CV_{mt}) for fertility and longevity. We estimated CV_{mt} for both fertility and median longevity by temperature and nuclear line and performed a two-sided sign test on bootstrapped estimates (1000 samples). The sign test compares the difference between female and male median values with negative values indicating higher male CV_{mt}, while positive values indicate that females have higher $CV_{mt.}$. The

majority of groupings indicate higher mitochondrial genetic variation in females in contradiction to the mother's curse hypothesis. Only fertility in BR nDNA individuals reared at 15°C had significantly higher mitochondrial genetic variation in males.

(B) Intra- and intersexual genetic correlation for female and male fertility and longevity across mitochondrial haplotypes. Bootstrapped Pearson's correlation coefficients (95% confidence intervals) were plotted for both rearing temperatures (15℃: solid lines, 25℃: dotted lines). Abbreviations: F (females), M (males), fertility (females: hatching number, males: remating success), longevity (mean longevity). All significant intersexual correlations indicate positive values, suggesting the absence of sexually antagonistic pleiotropy predicted by the mother's curse hypothesis.