Mitochondrial effects on fertility and longevity in *Tigriopus californicus*

contradict predictions of the mother's curse hypothesis

SUMMARY

Strict maternal inheritance of mitochondria favors the evolutionary accumulation of sex-biased

fitness effects, as mitochondrial evolution occurs exclusively in female lineages. The "mother's

curse" hypothesis proposes that male-harming mutations should accumulate in mitochondrial

genomes when they have neutral or beneficial effects on female fitness. Rigorous empirical tests

have largely focused on *Drosophila*, where support for the predictions of mother's curse has

been mixed. We investigated the impact of mother's curse mutations in *Tigriopus californicus,* a

minute crustacean. Using nonrecombinant backcrosses, we introgressed four divergent

mitochondrial haplotypes into two nuclear backgrounds and recorded measures of fertility and

longevity. We found that the phenotypic effects of mitochondrial mutations were context-

dependent, being influenced by the nuclear-background in which they were expressed, as well as

the sex of the individual and rearing temperature. Mitochondrial haplotype effects were greater

for fertility than longevity, and temperature effects were greater for longevity. However, in

opposition to mother's curse expectations, females had a higher level of mitochondrial genetic

variance than males for both fertility and longevity, and there was no evidence for mitochondrial

mutations favoring female performance at the cost of male performance. Together, this indicates

that selection on mitochondrial variation has not resulted in the accumulation of male mutation

load in *Tigriopus californicus*.

Introduction

In eukaryotes, life is powered by mitochondria - cytoplasmic organelles capable of producing
energy in the form of ATP. Originating from an ancient symbiosis over 2 billion years ago,
mitochondria provided the energy required for the evolution and diversification of multicellular
organisms[1–3]. Over evolutionary history, mitochondrial genomes have retained their ancestral
genetic codes and a few dozen genes while dramatically reducing their genome size through
deletion and gene transfer to the nucleus. Beyond the generation of ATP by oxidative
phosphorylation (OXPHOS), mitochondria perform additional cellular functions, including
initiation of apoptosis, amino acid and steroid biosynthesis, β-oxidation of fatty acids, cellular
signaling, and nuclear epigenetic regulation [4–8]. Nevertheless, out of 1,500 or more proteins
found in the mitochondrion, only the 13 membrane-bound ETC complexes are encoded by the
mitochondrial genome, with the rest being encoded by nuclear DNA[9,10]. The coevolution of
mitochondrial and nuclear genomes is thus an ancient yet ongoing process, involving the
evolutionary fine-tuning of various molecular interactions (i.e. protein-protein, protein-DNA,
and protein-RNA) required for the accurate replication and transcription of mtDNA, translation
of mtDNA-encoded proteins, and efficient cellular respiration. Mutation rates are generally
higher in mtDNA than in nDNA[11,12] leading to a continuous introduction of new mitochondrial
alleles. With asexual transmission and lack of recombination in mitochondria, novel alleles are
likely to persist, even when slightly deleterious, due to the potential masking effect of linkage
disequilibrium on mitochondrial genetic variation and reduced efficacy of natural selection[13].
Nevertheless, mtDNA likely experiences recurrent adaptive evolution via selective sweeps acting
on traits such as respiratory efficiency and mitonuclear coadaptation[14–18]. In natural populations,

mitochondrial fitness variation is likely to be present at substantial levels and may make a

meaningful contribution to adaptation and diversification[19]. The distribution of fitness effects

among mtDNA mutations is also likely to differ between the sexes, owing to the maternal

inheritance of mitochondria[20–22]. Strict maternal inheritance favors the evolutionary

accumulation of sex-biased fitness effects, as it is expected to preclude any opportunity for

evolution in males, forming a sex-specific selective sieve.

Under maternal inheritance, mitochondrial evolution is thought to be restricted to female

lineages, leading to the accumulation of male-harming mutations when they have neutral or

beneficial effects on female fitness (i.e. "mother's curse"[21,23]). Initial evidence for the mother's

curse hypothesis (MCH) centered on evidence that sperm motility in humans and other taxa was

linked to mitochondrial function[24,25,23] therefore sperm are more exposed to deleterious effects

of mtDNA mutations than ova. The MCH has also been proposed to impact viability, and even to

contribute to the taxonomically widespread pattern of shorter male lifespans[26]. Theoretical work,

however, shows that conditions favoring the spread of mother's curse mutations can be limited

by even modest levels of inbreeding and kin selection, allowing for an adaptive response to

mtDNA alleles impacting male viability and fertility[27–29]. Nevertheless, interest in MCH has

increased steadily in recent years leading to formal tests of the hypothesis, mostly in *Drosophila*,

utilizing carefully constructed mitochondrial-nuclear (mitonuclear) hybrids with divergent

mtDNA haplotypes backcrossed into an isogenic nuclear background. The most direct test of

MCH focuses on the effects of mutational variation in mtDNA, which are predicted to confer

larger trait variance in males. A second major prediction focuses on the subset of alleles that are

directly sexually antagonistic, which could lead to a negative intersexual genetic correlation

across haplotypes. A third prediction is that experimental disruption of mitonuclear coadaptation

will be more detrimental to males than females. Studies using the mitonuclear hybrid approach

have shown both support[26,30–34], and rejection[35–38] of the mother's curse hypothesis.

Understanding the pervasiveness of mother's curse is crucial given its fundamental

implications for the evolution of sexual dimorphism, as well as very practical consequences for

conservation breeding programs and medicine (particularly mitochondrial replacement theory

and male-biased mitochondrial disease). While past work has focused on *Drosophila*, the

predominance of maternal inheritance of mitochondria in eukaryotes allows for a taxonomically

diverse investigation of the generality of mother's curse. Here we test predictions of the MCH

using the splash-pool copepod *Tigriopus californicus*, an arthropod that shares a common

ancestor with *Drosophila* at least 450 million years ago. We used a genetic panel of mitonuclear

hybrids to assay both fertility and longevity. As fertility genes are likely sex-specific and

viability genes are likely shared between the sexes (see review[39]), we predict that mitonuclear

mismatch will lead to stronger sex bias for fertility than longevity. Additionally, we measured

the impact of high environmental temperature on these traits, as mild mitonuclear mismatch can

be expected to be unmasked under higher metabolic demand[40]. Populations of *T. californicus*

show extreme genetic divergence over short geographical distances[41–43] while remaining

somewhat tolerant of hybridization, producing viable and fertile offspring[44,45]. Nucleotide

divergence in *T. californicus* extends across the mtDNA genome, involving coding and

noncoding genes, and can exceed 20% between distant populations. *T. californicus* also differ

from most model organisms in that they lack heteromorphic chromosomes - sex is determined by

the inheritance of multiple, unlinked genes in combination with environmental effects (polygenic

sex determination)[46–48]. While rarely utilized in evolutionary biology, polygenic sex

determination provides an advantage for testing sex-specific mitochondrial effects as sex

differences cannot be attributed to the effects of sex chromosomes. Importantly, despite the lack

of heteromorphic sex chromosomes, there is marked sexual dimorphism in *T. californicus* and

considerable evidence for sex-specific effects in multiple traits (thermal tolerance[49,50]; salinity

tolerance[51,52]; resistance to toxins[52]; intrinsic hybridization stress[47]) in which males are generally

the weaker sex.

Methods

Population sampling

Populations of *T. californicus* are distributed in supralittoral "splash pools" along the Pacific

coast of North America. Copepods were originally collected from supralittoral pools at four

locations (Figure 1A): Friday Harbor, WA (FHL: 48.55 N,123.01 W), La Jolla, CA (LJ: 32.83

N,117.27 W), Bird Rock, CA (BR: 32.82 N,117.27 W), and San Diego, CA (SD: 32.7460

N,117.2551 W). Isofemale lines were initiated from a single female fertilized by a single male,

and maintained at 20°C on a 12h:12h light:dark cycle in standard culture media (three times

filtered seawater from Wrigley Marine Science Center, Santa Catalina Island, CA; 0.1g/l of

Spirulina (BRAND) and 0.1g/l ground Tetramin (Tetra®) Tropical fish food flakes). All lines

were maintained in petri dishes for ≥ 5 years under these conditions and periodically fed

additional Spirulina and ground Tetramin® tropical fish food and rehydrated with DI water

whenever necessary to maintain approximately 35 ppt salinity. Prior to experimentation, we

sequenced samples from each isofemale line at the 12S ribosomal RNA locus (Forward primer:

CCAAGATACTTTAGGGATAACAGC; Reverse primer: CTGTTCTATCAAAACAGCCCT;

fragment size: 687 base pairs) to rule out cross-contamination.

Mitonuclear hybrid panel

In *Tigriopus*, meiosis is chiasmatic in males but achiasmatic in females, yielding recombinant
sperm and non-recombinant ova. This allows us to backcross mitochondria from the population
of the dam into that of the sire while preventing nuclear recombination between populations.
Because sexes in *T. californicus* are not easily distinguished before maturity, we allowed males
to clasp female siblings with their geniculate first antennae in order to ensure the sex of females.
We moved clasped pairs onto filter paper and quickly separated them using entomological pins
under a microscope. We backcrossed FHL, BR, LJ, and SD dams to SD and BR sires for 10
generations (Figure 1B), resulting in an average proportion of $1 - \frac{1}{2}^{(b+1)}$ (where b is equal to the
number of backcross generations) or 99.95% of the sire's nuclear genome, assuming no
selection. For each cross, we established 4 replicates, each maintained in 12-well plates
containing standard culture media. All crosses were fed weekly with additional Spirulina and
ground Tetramin® fish food and rehydrated with DI water whenever necessary to maintain
approximately 35 ppt salinity.

Trait measurement and modeling

For each of our mitonuclear hybrid lines, we measured three fitness components (fertility,
lifespan, and age-specific rate of survival) for males and females reared at two contrasting
temperature regimes (15ºC and 25ºC). In order to achieve an adequate sample size for each
mtDNA haplotype and sex, it was necessary to set up multiple families from each mitonuclear
hybrid line. Each family was composed of up to four clutches of fertilized eggs derived from a
single mitonuclear hybrid or control female (dam). Dams were maintained at standard conditions
and checked daily for the presence of hatched eggsacs. For each dam, we alternated moving up

to four hatched eggsacs to 15ºC and 25ºC (Figure 1B) so that clutch order was orthogonal to

temperature in both fertility and longevity experiments. Individuals were maintained at the same

temperature for the duration of the fertility or longevity experiment.

Longevity

Individual animals were separated into wells in a 24-well plate once they developed to the

juvenile copepodid stage and were checked three times a week. The first measurement we

recorded was the sex-specific age of reproductive maturity. For males, this occurred with the

development of geniculate first antennae (claspers) and for females with the development of

eggs, which can be seen through the translucent body. We continued monitoring longevity

experiments three times a week, recording the age of death of individuals until all had expired.

We performed linear mixed-effects modeling on family sex ratio using sex information

from our longevity experiment, comparing mtDNA x nDNA interactions with the impact of

mitonuclear status on family sex ratio. We also conducted a separate experiment, using different

families, measuring mortality occurring before sexual maturity (Document S1, Figure S1). We

were able to measure the age at the naupliar-copepodid transition, duration of the copepodid

stage, and the sex-specific age of reproductive maturity, and perform linear mixed-effects

modeling investigating the influence of genetic (mtDNA, nDNA) and environmental

(temperature) influence on these larval life-history parameters.

Sex-specific fertility

For female fertility experiments, all individuals from hatched eggsacs were separated into

individual wells in a 24-well plate once they developed to the juvenile copepodid stage. To each

well, we added a single wild-caught adult male from the appropriate locality (SD or BR),

matching the paternal line of the individual. Males were added to every well because sexes are

difficult to distinguish morphologically in immature *T. californicus* and we wanted to ensure that

each female was paired with an adult male before sexual maturity. Mate clasping typically occurs

between adult males and juvenile females, and rarely occurs with mature females once they

begin extruding non-fertilized eggsacs. We also ran experiments utilizing sib-pairs in parallel to

compare the impact of mating with mitonuclear hybrid males versus wild-type males on female

fertility (Document S1). Plates were maintained at the appropriate temperature (15ºC or 25ºC)

and checked three times a week until females produced an eggsac. As we utilized several

hundred plates in our experiment, it became necessary to maximize incubator space. Therefore,

we placed gravid females from the same family and clutch together in petri dishes until they

reached experimental age. Because individuals develop twice as fast at 25ºC as at 15ºC, we used

temperature-specific ages for our experiment in order to minimize age-related variation. Once

individuals reared at 25°C reached 35 days of age, and individuals reared at 15°C reached 70

days of age, we moved them into separate dishes and maintained them at the correct temperature,

checking them three times a week until eggsacs hatched. Upon hatching, each individual larval

nauplius was counted under a dissecting microscope to measure female hatching number.

For male fertility experiments, we removed males from each of the four clutches in a

family and allowed them to age for the appropriate temperature (35 days old for 25ºC and 70

days old for 15ºC). Once at the appropriate age, each male was pipetted into a petri dish with six

wild-caught juvenile females from the appropriate locality (BR or SD) to match the paternal

nuclear line. To ensure that wild-caught juveniles were female, we selected clasped pairs from

culture within 0-2 weeks of collection and separated pairs under a dissecting microscope. Care

was taken to provide males with females of consistent and reasonably large size, to reduce

variation across experiments. Each experimental male and six wild-caught females were allowed

14 days to mate in a petri dish at the appropriate temperature. After 14 days, females were

isolated into a 6-well plate and scored for the presence or absence of any offspring indicating a

successful mating. We recorded male remating success as the ratio of successful matings to the

number of surviving females.

Parametric and nonparametric trait modeling

For our trait modeling, we performed three types of linear models: sex-effect modeling,

coevolutionary status modeling, and best-fit models. Sex-effect models measure sex effects

impacted by mtDNA and nDNA variation and their interactions. If mother's curse mutations are

present, they may exist in additive (Sex x mtDNA) or epistatic (Sex x mtDNA x nDNA) form.

Sex differences may also be encoded by nuclear genes, described by the (Sex x nDNA) term. In

order to compare fertility experiments across the two sexes, individual measurements of fertility

were normalized to the mean of the appropriate matched control line reared at the same

temperature (e.x. male *i* from FHLxSD at 15°C / mean of all males from SDxSD at 15°C). We

performed model testing on the four sex-effect models using the *model.sel* function in the

MuMIn[53] R package, which ranks models based on their AIC_C values. Model contrasts for sex

differences were performed using the *pairs* function from the emmeans[54] R package .

Coevolutionary status models measure the impact of disrupting coevolved mitonuclear

combinations on sex-specific data grouped by temperature and nDNA. These models included

coevolutionary status as a fixed-effect term. Modeling for best-fit utilized an "all-subset"

approach, where all possible combinations of predictors were compared using the *MuMIn*

function *dredge*. Models with the highest weight and lowest AIC_C were designated as best-fit.

Both fertility and longevity models included temperature, nDNA, mtDNA and all possible

interactions as fixed-effects terms in initial models and were performed separately for each sex.

For fertility, we used raw sex-specific fertility (males: remating success, females: hatching

number). All linear mixed-effects modeling was performed using the *lmer* function from the

lme4[55] R package and included clutch nested within family as a random-effect term. We used

type III ANOVA to estimate significance for model main effects and interaction terms using the

Anova function from the car[56] R package.

We performed parametric regression of survivorship for each nuclear, temperature, and

sex combination against the Gompertz distribution [57], estimating the shape (α) and rate (β)

parameters using the R package flexsurv[58]. Parameters were estimated by maximum likelihood,

with those defined to be positive estimated on a log scale. Confidence intervals were estimated

from the Hessian at maximum, and then transformed back to the original scale (days). The shape

of the resulting distributions, $\ln(\alpha)$, describe the initial mortality rate or frailty of the samples,

while the rate (β) parameter describes the age-independent rate of mortality acceleration or rate

of senescence. We employed Kruskal-Wallis one-way analysis of variance to estimate the

significance of parameter estimates produced by parametric survival models using the

kruskal.test function from the stats[59] R package.

Mitochondrial coefficient of variation

One prediction of the mother's curse hypothesis is that the accumulation of male-affecting

variation should lead to a higher mitochondrial coefficient of variation (CV_{mt}) in males than in

females. Using the *boot*[60,61] package in R, bootstrapped CV_{mt} was computed by resampling raw

trait means for each mitochondrial line using 1000 replicates with replacement and calculating

the coefficient of variation among mitochondrial lines ($CV = \sigma/\mu$) for each nuclear, sex, and

temperature combination. Hypothesis testing was performed to compare male and female CV_{mt}

using a sign test as implemented in the *SIGN.test* function in the BSDA[62] R package.

Genetic correlation among haplotypes

Mother's curse mutations are likely to reach fixation fastest when they confer a fitness advantage

to females, resulting in sexual conflict when they simultaneously result in low fitness in males.

When enough mtDNA variation is involved in sexual conflict, a negative intersexual correlation

may be observed across haplotypes for a given trait. We measured the correlation between

haplotype means for fertility and median longevity, comparing traits within sexes (intrasexual

correlation) and also comparing the sexes for a given trait (intersexual correlation). We used

bootstrapped estimates of Pearson's coefficient using the boot R package along with the *cor*

function from the stats R package, recording 1000 samples for each comparison. Equi-tailed,

two-sided nonparametric confidence intervals were generated from the first order normal

approximation with a confidence level of 0.95 using the *boot.ci* function from the boot package

in R.

Whole genome sequencing

For each isofemale and mitonuclear hybrid line, we extracted DNA from 10 males and 10 females using Zymo quick-DNA preps and quantified DNA yields using Qubit HS dsDNA quantification kits. Because female *T. californicus* yield approximately 2.12x more DNA than males, we pooled 2 ng of DNA from each of the 20 individuals per line for library preparation. DNA pools were sheared using a Covaris S2 sonicator with microTUBE-50 AFA tubes, and sequencing libraries were prepared using Qiagen QIAseq Ultralow Input kits following instructions. Mitonuclear hybrid libraries were pooled together in equimolar amounts and sequenced together on a single lane of Illumina HiSeq4000 (PE-150), and the parental isofemale libraries were pooled and sequenced on a separate lane. Sequence reads were quality trimmed using trimmomatic[63] v0.36, and paired reads were filtered for mitochondrial reads using the filter_reads.pl perl script included with the NOVOPlasty[64] software. We assembled consensus mtDNA genomes from each sequenced isofemale line, by utilizing the appropriate mtDNA reference from a previous publication which included the SD, BR, and FHL populations[18]. A reference mtDNA genome did not exist for the LJ population, so those reads were mapped to the BR mtDNA reference. We annotated our *de novo* assembled genomes using the Mitos[65] 2.0 software, each containing all 13 protein coding genes, and the majority of non-coding RNAs. Following this, we mapped all trimmed and paired reads to the *T. californicus* reference genome assembly (NCBI accession: GCA_007210705.1) along with the appropriate mitochondrial genome using stampy[66] v1.0.32 (substitutionrate=0.05). Following the removal of PCR and optical duplicates using picard[67] MarkDuplicates, we called variants for each sequenced pool using samtools[68] mpileup.

We estimated pairwise nuclear divergence for isofemale lines in a sliding window

approach. For each comparison, we calculated F_{ST} using a sliding window of 10,000 bp,

requiring a depth of 40x coverage, and minimum PHRED base quality score of 30 using the fst-

sliding.pl perl script included with popoolation v1.2.2. For parental mitochondrial divergence,

we aligned whole mtDNA genomes using progressiveMauve[69] along with mtDNA genome

sequences from an outgroup population from San Roque, Baja California Sur, Mexico to produce

rooted trees, and measured divergence (d_{XY}) using the calculate-dxy.pl perl script included with

popoolation v1.2.2. We also measured divergence at synonymous and nonsynonymous sites by

aligning protein-coding sequences with MACSE [70] v2, constructing neighbor joining trees with

ClustalW [71] and analyzing these data with codeml from the PAML suite [72] using the null model

(M0), assuming a uniform selective pressure among sites.

Confirming mitonuclear genetic status of hybrids

To test for potential selection or accidental introgression, we measured allele frequencies at a set

of divergent sites for each paternal line's nuclear genome. A Fisher's exact test for significant

allele frequency differences for nDNA from the maternal and paternal lines was conducted using

popoolation[73] v1.201, requiring a minimum depth of 20x coverage. In mitonuclear hybrids, sites

with no significant differences (alpha > 0.05) to the paternal line were considered "paternally

concordant" and significantly different sites (alpha <= 0.05) were considered "paternally

discordant". Because significance may be determined by variations in the same alleles between

lines, we recorded paternally discordant sites with alleles matching maternal lines as "maternally

concordant" when they were absent in the paternal nDNA. Paternal concordance rates were

recorded by measuring the proportion of sites that were not maternally concordant to the total

number of sites. We also examined mitonuclear hybrid lines for the presence of paternally

inherited mtDNA. This was accomplished by mapping trimmed and paired sequence reads to the

reference genome as well as all mtDNA haplotypes simultaneously. We measured depth of

coverage using samtools depth command to identify any potential sources of contamination.

Results

DNA sequencing

Illumina HiSeq 4000 sequencing yielded an average of 34.7 million reads for mitonuclear hybrid

line pools, and 76.1 million reads for inbred line pools. Following quality trimming and mapping

of paired reads to the appropriate references, an average of 29 million reads mapped for cybrid

pools, and 61 million reads mapped for maternal and paternal line pools. Average nuclear

genome coverage across inbred lines was 91.88% at a mean depth of coverage of 26x. Across

the mitochondrial genome, the mean depth of coverage was 2670X, and genome coverage was

100%. Nuclear divergence (F_{ST}) between paternal lines ranged from 0.049 to 0.369 in lines with

BR nDNA, and from 0.102 to 0.378 in lines with SD nDNA (Figure 1C). Mitochondrial

divergence (d_{XY}) between parental lines ranged from 0.7% to 21.1% in BR nDNA lines, and

9.6% to 20.6% in SD nDNA lines. The mitonuclear hybrid panel represented a wide range of

mitochondrial variation, with as few as 13 amino acid substitutions between the BR and LJ

mitotypes, 170 aa substitutions between the BR and SD nuclear line mitotypes, and 555 aa

substitutions between SD and FHL haplotypes. The highly divergent FHL haplotype also

differed from the BR & LJ haplotypes by 546 aa substitutions.

For each mitonuclear hybrid line, we measured allele frequencies at an average of 580k

divergent nDNA SNPs between parental lines. We found that alleles were mostly concordant

with paternal nDNA, with limited evidence of introgression from maternal nDNA, with paternal

concordance rates ranging from 99.15% - 100% (Figure 1C). To test for potential mtDNA

contamination via paternal leakage or experimental error, we compared depth of coverage across

all mtDNA haplotypes, since mtDNA positions and alleles are not comparable across haplotypes.

We found that the percent maternally matching depth ranged from 94.09% to 99.81%, while the

percent of paternally matching depth ranged from 0.02% to 0.28% providing limited evidence

for contaminating mtDNA (Figure 1C).

325   Trait measurements and modeling

Despite initially setting up four replicates for each mitonuclear hybrid line, low productivity led

to extinction of several replicates, forcing us to proceed with one replicate per line to keep our

experiment balanced. We measured the hatching number for 182 females coming from 41

families reared at 15ºC and 216 females from 40 families reared at 25ºC. Male remating success

was measured for 491 males, using a total of 2,458 wild-caught females. Since 378 females died

over the course of the experiment, we discarded 31 males that had fewer than four out of six

females remaining in order to reduce variation associated with differential access to females.

Therefore, our male remating experiment is based on 235 males from 68 families reared at 15ºC

and 225 males from 76 families reared at 25ºC. The average male successfully copulated with

2.45 females in the span of 14 days when reared at 15°C, declining to 2.04 females at 25°C

representing a 17% reduction. In the average female clutch, 30.05 eggs hatched at 15°C,

declining to 21.50 eggs hatching at 25°C representing a 28% reduction. Female hatching number

declined an average of 31.16% in mitonuclear hybrid lines when females were bred with siblings

as opposed to wild-caught males (Document S1, Figure S2. Lifespan, frailty, and rate of

senescence was measured for 3,072 individuals, with 1,531 individuals from 89 families reared

at 15ºC, and 1,541 individuals from 77 families reared at 25ºC.

Experiments on early developmental stages indicate that mortality rate occurring before sexual

maturity was very high, ranging from 14.6% to 90.6% mortality depending on cross and

temperature (Document S1, Figure S1. Naupliar stages had the highest mortality, ranging from

4.2% to 84.4%, while copepodid stages ranged from 1.5% to 27.9% mortality. Linear mixed-

effects modeling showed that measurements describing the development rate varied significantly

with mtDNA x nDNA interactions and rearing temperature, with rates increasing at higher

temperature.

Sex-effect modeling

Model contrasts revealed mostly male-biased values for longevity (longer lifespan), and mixed

bias for fertility with three out of six significant contrasts being female biased (Figure 2A).

Model selection comparing the four sex-effects models (Table S1) revealed that models

including three-way sex x nDNA x mtDNA interaction term fit the trait data best in all

temperature and trait combinations except for relative fertility at 15°C, which was better fit with

sex x nDNA interactions (ΔAIC = 17.35). None of the sex-effects models uncovered significant

sex x mtDNA interactions (Table S2). However, we do see that sex x nDNA and sex x mtDNA x

nDNA interactions are a significant influence on fertility variation (Figure 2B).

Mitonuclear coevolutionary status modeling

We found that neither fertility nor longevity were influenced by mitonuclear status (matched vs.

mismatched), as indicated from type III ANOVA on fitted parameters (Table S3). Family sex

ratio was significantly influenced by mitonuclear status (χ^2= 18.51, p = 1.7e-5), as indicated in

linear mixed-effects modeling (Document S1, Figure S3), as well as nDNA x mitonuclear status

interactions (χ^2= 28.77, p = 8.1e-8). Results using mortality-corrected family sex ratio data were

also consistent with uncorrected family sex ratio. (Table S5).

Best-fit models

Temperature significantly impacted all traits by reducing their average values (Table S4).

Notably, mean longevity was reduced by 102 days in 25ºC, an almost 10 day decrease per degree

reduction. Significant mtDNA effects were only seen in females for both hatching number (χ^2=

20.36, p = 1.43e-4) and longevity (χ^2= 14.66, p = 0.0021). Variation in nDNA impacted

longevity in both sexes (males: χ^2= 20.36, p = 6.42e-6; females: χ^2= 8.31, p = 0.0040) but

neither measure of fertility (female hatching number: χ^2= 1.37, p = 0.2416; male remating

success: χ^2= 0.15, p = 0.6991). Additionally, effects of nDNA were contingent upon rearing

temperature, with significant interactions between the two terms for male remating success (χ^2=

7.41, p = 0.0065), male longevity (χ^2= 4.19, p = 0.0407) and female longevity (χ^2= 4.05, p =

0.0442). Interactions between mtDNA and nDNA were also significant for longevity in females

(χ^2= 12.03, p = 0.0073) and in a three-way interaction with temperature for male remating

success (χ^2= 16.95, p = 7.22e-4). Mitochondrial effects impacted fertility more than longevity

(Figure 2C, p = 3.2e-4), while nuclear effects impacted both traits equally (p = 0.95) and

temperature affected longevity more than fertility (p = 1.5e-6). Linear mixed-effects modeling of

family sex ratio indicated significant influence of temperature ($\chi^2 = 10.02$, $p = 0.0015$), mtDNA

haplotype ($\chi^2 = 28.35$, $p = 3.1$e-6), as well as interactions between mtDNA and nDNA ($\chi^2 =$

31.98, $p = 1.6$e-8) and temperature and nDNA ($\chi^2 = 29.94$, $p = 1.4$e-6; Document S1, Figure S3).

Parametric survivorship modeling

Survivorship curves indicate a drastic reduction in longevity shifting from 15°C to 25°C, with

SD nDNA individuals having higher survivorship than BR nDNA individuals at 25°C (Figure

3A). Additionally, we see that median longevity is sex-biased in SD nDNA individuals reared at

15°C, with higher male longevity (Figure 3A insert, Wilcoxon rank sum test, p = 0.004). Rearing

temperature affected both frailty ($\chi^2 = 22.55$, $p = 2.05$e-6) and rate of senescence ($\chi^2 = 23.27$, $p =$

1.41e-6) in a similar manner, with the higher temperature leading to increased frailty and

increased rate of senescence (Table S6). Mitochondrial haplotype showed no significant

contribution to Gompertz parameters when reared at either 15ºC or 25ºC (Table S6). We found

significant sexual dimorphism in frailty and senescence in individuals reared at 25ºC, but not

those reared at 15ºC (Figure 3B). When reared at 25ºC, females had lower frailty than males, but

a higher rate of senescence and while this pattern held in both nDNA backgrounds, it was

nonsignificant in BR nDNA lines (Table S6).

Mitochondrial coefficient of variation

Females had significantly higher CV_{mt} than males in seven out of eight experimental groupings

(Figure 4A, Table S3). The median difference between male and female coefficients in these

groupings ranged from 0.0039 to 0.1061. In contrast, fertility in BR nDNA individuals at 25ºC

showed higher male CV_{mt} than females (median difference = -0.0993, CI = (-0.11, -0.09), $p <$

2.2e-16).

Genetic correlation among haplotypes

Intersexual correlations for median longevity ($r = 0.99$; $CI_{95} = 0.20, 2.00$) and fertility ($r = 0.84$;

$CI_{95} = 0.08, 1.72$) were significantly positive in BR nDNA individuals reared at 25ºC, while

correlations for median longevity in SD nDNA individuals were also significantly positive ($r =$

0.85; $CI_{95} = 0.04, 1.75$) in individuals reared at 25ºC (Figure 4B). Intrasexual correlations

between fertility and longevity were significantly positive in BR nDNA individuals of both sexes

reared at 25ºC (males: $r = 0.96$; $CI_{95} = 0.21, 1.95$; females: $r = 0.90$; $CI_{95} = 0.08, 1.86$). No

measures of intersexual or intrasexual correlation were significantly negative.

Discussion

Our study sought to explicitly test three predictions arising from the mother's curse

hypothesis: higher mitochondrial genetic variance in males for fitness traits, negative intersexual

correlation among mitochondrial haplotypes for fitness traits, and higher impact of experimental

disruption of putatively coevolved mitonuclear genotypes on males. Our results did not support

these predictions. Females had a higher level of mitochondrial genetic variance in seven

experimental groupings and lower variance in one grouping (fertility in individuals with BR

nDNA reared at 25ºC). We also found no evidence for sexual antagonism for fertility or

longevity, with negative intersexual correlations found in two of eight experimental groupings

(fertility in BR nDNA at 15°C and SD nDNA at 25°C) having upper confidence intervals that

extended well into positive ranges (0.90 and 1.41, respectively). Overall, the intersexual

comparisons mimicked the intrasexual comparisons. Lastly, we did not find the significant

influence of coevolutionary status on fertility or longevity traits required to test the third

prediction.

Detecting sex-bias in mitochondrial coefficient of variation for fitness traits is arguably

the most direct test of the hypothesis since it emphasizes the differential accumulation of

mutational variation between the sexes[74]. According to the mother's curse hypothesis,

mitochondrial mutations causing male infertility that are neutral in females are more likely to

spread in a population through random drift since males are a dead-end for natural selection on

mtDNA. However, this argument is best applied to randomly mating populations with no

inbreeding. In populations with even modest levels of inbreeding, such as those with

geographical population structure, females will also suffer fitness insults from mother's curse

mutations to a certain degree since they are more likely to mate with sub-fertile siblings[27,28,75].

The degree to which male subfertility influences female fertility fitness can also influence these

dynamics. In monandrous species like *Tigriopus*, females mate only once[76,77], resulting in a

direct relationship between male fertility and female offspring production. In this study, we

showed that female hatching number dropped substantially when mated with siblings compared

to wild-caught males, indicating the influence of male subfertility on female fertility. However,

in mating systems where females mate with several males (e.g. *Drosophila*), females can be

safeguarded against male infertility or subfertility[78]. As a result, selection to purge male-harming

mtDNA mutations is expected to be strongest in species with monandry and weakest with

polyandry whenever inbreeding is present[75]. Studies show that *T. californicus* avoid close

inbreeding[79], but live in highly isolated, ephemeral populations making inbreeding all but

452 inevitable[80,81]. Additionally, while inbreeding can limit mother's curse by decreasing the

453 conditions under which male-harming alleles can spread, it can also result in the spread of

454 female-harming alleles that benefit males and lead to female-biased fitness variance [22,27]. The

455 widespread female-biased fitness variance observed in our fertility and longevity experiments

456 may therefore indicate the impact of historical inbreeding on the fixation of mitochondrial

457 divergence sampled in our study.

459   We predicted that sex differences due to mitochondrial haplotype effects are greater for

460 fertility than longevity, based on the expectation that fertility genes are more sex-specific than

461 viability genes[82,83]. We also predicted that the consequences of mitonuclear mismatch should

462 increase with metabolic rate associated with increased temperature. Efficient energy production

463 depends on mitochondrial function, including mitonuclear coordination. Mild mitonuclear

464 mismatches may be tolerated under low aerobic demand and yet become lethal when energetic

465 demand become higher [40,but see 82]. Increased temperature reduced measures of both fertility and

466 longevity but was more dramatic with the latter. We uncovered significant mitonuclear effects

467 impacting longevity at 25°C but not 15°C. Additionally, we found that mitonuclear x sex effects

468 were significant for fertility at 25°C, but not 15°C. Mitochondrial haplotype effects in our

469 experiment impacted fertility more than longevity, however experiments on the larval stages

470 indicate a substantial amount of mortality, influenced by temperature and mitonuclear

471 interactions, occurring before sexual maturity. Because *T. californicus* sperm are amotile,

472 declines in male fertility are not attributed to loss of sperm motility – a key marker of male

473 sterility in organisms with flagellated, motile sperm. Instead, declines in male fertility in our

study are likely due to the number of mature spermatids in a spermatophore, the ability of sperm

to maintain homeostasis while residing in the female reproductive tract, or behavior.

Survivorship analysis indicates that mtDNA variation had an inverse impact on frailty

and senescence between the sexes, similar to the 'Strehler-Mildvan (S-M) relationship[85]. When

reared at 25ºC, females had lower frailty than males while also having a higher rate of

senescence, despite having similar median longevity. One explanation for this relationship is that

the preferential survival of the most-frail female *T. californicus* may lead to an increase in the

rate of senescence for females, since frail individuals are more susceptible to aging. Male *T.*

californicus typically have lower tolerance than females to a variety of stressors, including

temperature, and a greater transcriptomic response to oxidative stress[49,50,52,86,87].

Conclusion

We found that the functional mtDNA effects were highly context-dependent, being

influenced by the nuclear-background in which they were expressed, the temperature in which

individuals developed, and/or the sex of the individual. We explicitly sought to test the main

predictions of the mother's curse hypothesis, finding patterns opposite to those predictions.

Support for the mother's curse hypothesis has been highly mixed to date, both theoretically and

empirically, with many studies rejecting these predictions. While most studies directly testing

predictions of the mother's curse hypothesis have used *Drosophila*, the predominance of

maternal inheritance of mitochondria in eukaryotes allows for a taxonomically diverse

investigation of the generality of mother's curse. Particularly, diversity in mating system should

be considered for future studies, as the fitness of females that mate with only one male will be

strongly determined by the fitness of her mate when there is inbreeding. Furthermore, sperm

diversity (motile/amotile, storage/remating) should be highlighted in order to address the

relevance of various mutational insults to male fitness potentially harbored by mitochondrial

DNA. In light of the evidence presented here and elsewhere, we suggest that accumulation of

male-harming mitochondrial mutations may not be a generalizable phenomenon, despite the

prevalence of maternal mitochondrial transmission across eukaryotes.

502
503

References

1. Yang, D., Oyaizu, Y., Oyaizu, H., Olsen, G. J. & Woese, C. R. Mitochondrial origins. *PNAS* **82**,

4443–4447 (1985).

2. Gray, M. W., Burger, G. & Lang, B. F. Mitochondrial Evolution. *Science* **283**, 1476–1481

(1999).

3. Hedges, S. B., Blair, J. E., Venturi, M. L. & Shoe, J. L. A molecular timescale of eukaryote

evolution and the rise of complex multicellular life. *BMC Evol Biol* **4**, 2 (2004).

4. McBride, H. M., Neuspiel, M. & Wasiak, S. Mitochondria: More Than Just a Powerhouse.

Current Biology **16**, R551–R560 (2006).

5. Pagliarini, D. J. & Rutter, J. Hallmarks of a new era in mitochondrial biochemistry. *Genes Dev.*

27, 2615–2627 (2013).

6. Chandel, N. S. Mitochondria as signaling organelles. *BMC biology* **12**, 1 (2014).

7. Birsoy, K. *et al.* An Essential Role of the Mitochondrial Electron Transport Chain in Cell

Proliferation Is to Enable Aspartate Synthesis. *Cell* **162**, 540–551 (2015).

8. Santos, J. H. Mitochondria signaling to the epigenome: A novel role for an old organelle. *Free*

Radical Biology and Medicine **170**, 59–69 (2021).

9. Bar-Yaacov, D., Blumberg, A. & Mishmar, D. Mitochondrial-nuclear co-evolution and its

effects on OXPHOS activity and regulation. *Biochimica et Biophysica Acta (BBA) - Gene*

Regulatory Mechanisms **1819**, 1107–1111 (2012).

10.   Lotz, C. *et al.* Characterization, design, and function of the mitochondrial proteome:

from organs to organisms. *J. Proteome Res.* **13**, 433–446 (2014).

11.   Brown, W. M., George, M. & Wilson, A. C. Rapid evolution of animal mitochondrial DNA.

Proc. Natl. Acad. Sci. USA 5 (1979).

12.   Lynch, M. Rate, molecular spectrum, and consequences of human mutation.

Proceedings of the National Academy of Sciences **107**, 961–968 (2010).

13.   Lynch, M. & Blanchard, J. L. Deleterious mutation accumulation in organelle genomes.

Genetics **102/103**, 29–39 (1998).

14.   Schmidt, T. R., Wu, W., Goodman, M. & Grossman, L. I. Evolution of nuclear-and

mitochondrial-encoded subunit interaction in cytochrome c oxidase. *Molecular Biology and*

Evolution **18**, 563–569 (2001).

15.   Willett, C. S. & Burton, R. S. Evolution of Interacting Proteins in the Mitochondrial

Electron Transport System in a Marine Copepod. *Mol Biol Evol* **21**, 443–453 (2004).

16.   Bazin, E., Glémin, S. & Galtier, N. Population Size Does Not Influence Mitochondrial

Genetic Diversity in Animals. *Science* **312**, 570–572 (2006).

17.   James, J. E., Piganeau, G. & Eyre-Walker, A. The rate of adaptive evolution in animal

mitochondria. *Molecular Ecology* **25**, 67–78 (2016).

18.   Barreto, F. S. *et al.* Genomic signatures of mitonuclear coevolution across populations of

Tigriopus californicus. *Nature Ecology & Evolution* **2**, 1250–1257 (2018).

19.   Hood, W. R. *et al.* The Mitochondrial Contribution to Animal Performance, Adaptation,

and Life-History Variation. *Integrative and Comparative Biology* **58**, 480–485 (2018).

20.   Wolstenholme, D. R. Animal Mitochondrial DNA: Structure and Evolution. in

International Review of Cytology (eds. Wolstenholme, D. R. & Jeon, K. W.) vol. 141 173–216

(Academic Press, 1992).

21.   Frank, S. A. & Hurst, L. D. Mitochondria and male disease. *Nature* **383**, 224–224 (1996).

22.   Smith, S. R. T. & Connallon, T. The contribution of the mitochondrial genome to sex-

specific fitness variance. *Evolution* **71**, 1417–1424 (2017).

23.   Gemmell, N. J., Metcalf, V. J. & Allendorf, F. W. Mother's curse: the effect of mtDNA on

individual fitness and population viability. *Trends in Ecology & Evolution* **19**, 238–244 (2004).

24.   Cardullo, R. A. & Baltz, J. M. Metabolic regulation in mammalian sperm: Mitochondrial

volume determines sperm length and flagellar beat frequency. *Cell Motility* **19**, 180–188

(1991).

25. Holyoake, A. J. *et al.* High incidence of single nucleotide substitutions in the

mitochondrial genome is associated with poor semen parameters in men. *International*

Journal of Andrology **24**, 175–182 (2001).

26. M. Florencia Camus, David J. Clancy, & Damian K. Dowling. Mitochondria, Maternal

Inheritance, and Male Aging. *Current Biology* **22**, 1717–1721 (2012).

27. Wade, M. J. & Brandvain, Y. REVERSING MOTHER'S CURSE: SELECTION ON MALE

MITOCHONDRIAL FITNESS EFFECTS. *Evolution* **63**, 1084–1089 (2009).

28. Unckless, R. L. & Herren, J. K. Population genetics of sexually antagonistic mitochondrial

mutants under inbreeding. *Journal of Theoretical Biology* **260**, 132–136 (2009).

29. Hedrick, P. W. REVERSING MOTHER'S CURSE REVISITED. *Evolution* **66**, 612–616 (2012).

30. Innocenti, P., Morrow, E. H. & Dowling, D. K. Experimental Evidence Supports a Sex-

Specific Selective Sieve in Mitochondrial Genome Evolution. *Science* **332**, (2011).

31. Camus, M. F., Wolf, J. B. W., Morrow, E. H. & Dowling, D. K. Single Nucleotides in the

mtDNA Sequence Modify Mitochondrial Molecular Function and Are Associated with Sex-

Specific Effects on Fertility and Aging. *Curr Biol* **25**, 2717–2722 (2015).

32. Camus, M. F. & Dowling, D. K. Mitochondrial genetic effects on reproductive success:

signatures of positive intrasexual, but negative intersexual pleiotropy. 10.

33. Wolff, J. N. *et al.* Evolutionary implications of mitochondrial genetic variation:

mitochondrial genetic effects on OXPHOS respiration and mitochondrial quantity change

with age and sex in fruit flies. *Journal of Evolutionary Biology* **29**, 736–747 (2016).

34. Carnegie, L., Reuter, M., Fowler, K., Lane, N. & Camus, M. F. Mother's curse is pervasive

across a large mitonuclear Drosophila panel. *Evolution Letters* **5**, 230–239 (2021).

35.   Mossman, J. A., Tross, J. G., Li, N., Wu, Z. & Rand, D. M. Mitochondrial-Nuclear

Interactions Mediate Sex-Specific Transcriptional Profiles in Drosophila. *Genetics* **204**, 613–

630 (2016).

36.   Mossman, J. A., Biancani, L. M. & Rand, D. M. Mitonuclear Epistasis for Development

Time and its Modification by Diet in Drosophila. *Genetics* (2016)

doi:10.1534/genetics.116.187286.

37.   Immonen, E., Collet, M., Goenaga, J. & Arnqvist, G. Direct and indirect genetic effects of

sex-specific mitonuclear epistasis on reproductive ageing. *Heredity* **116**, 338–339 (2016).

38.   Đorđević, M. *et al.* Sex-specific mitonuclear epistasis and the evolution of mitochondrial

bioenergetics, ageing, and life history in seed beetles: SEX-SPECIFIC MITONUCLEAR

EPISTASIS. *Evolution* **71**, 274–288 (2017).

39.   Ashburner, M., Golic, K. G., Hawley, R. S., & others. *Drosophila: a laboratory handbook.*

(Cold spring harbor laboratory press, 2004).

40.   Lane, N. The Costs of Breathing. *Science* **334**, 184–185 (2011).

41.   Burton, R. S., Feldman, M. W. & Curtsinger, J. W. Population genetics of Tigriopus

californicus (Copepoda: Harpacticoida): I. Population structure along the central California

coast. *Marine Ecology Progress Series* 29–39 (1979).

42.   Burton, R. S. Intraspecific Phylogeography Across the Point Conception Biogeographic

Boundary. *Evolution* **52**, 734–745 (1998).

43.   Edmands, S. Phylogeography of the intertidal copepod Tigriopus californicus reveals

substantially reduced population differentiation at northern latitudes. *Molecular Ecology* **10**,

1743–1750 (2001).

44.   Edmands, S. Heterosis and Outbreeding Depression in Interpopulation Crosses Spanning

a Wide Range of Divergence. *Evolution* **53**, 1757–1768 (1999).

45.   Pritchard, V. L. & Edmands, S. THE GENOMIC TRAJECTORY OF HYBRID SWARMS:

OUTCOMES OF REPEATED CROSSES BETWEEN POPULATIONS OF *TIGRIOPUS CALIFORNICUS*:

TIGRIOPUS CALIFORNICUS HYBRID SWARMS. *Evolution* **67**, 774–791 (2013).

46.   Voordouw, M. J. & Anholt, B. R. Heritability of sex tendency in a harpacticoid copepod,

Tigriopus californicus. *Evolution* **56**, 1754–1763 (2002).

47.   Foley, B. R., Rose, C. G., Rundle, D. E., Leong, W. & Edmands, S. Postzygotic isolation

involves strong mitochondrial and sex-specific effects in Tigriopus californicus, a species

lacking heteromorphic sex chromosomes. *Heredity* **111**, 391–401 (2013).

48.   Alexander, H. J., Richardson, J. M. L., Edmands, S. & Anholt, B. R. Sex without sex

chromosomes: genetic architecture of multiple loci independently segregating to determine

sex ratios in the copepod *Tigriopus californicus*. *Journal of Evolutionary Biology* **28**, 2196–

2207 (2015).

49.   Willett, C. S. Potential Fitness Trade-Offs for Thermal Tolerance in the Intertidal

Copepod Tigriopus Californicus. *Evolution* **64**, 2521–2534 (2010).

50.   Kelly, M. W., Sanford, E. & Grosberg, R. K. Limited potential for adaptation to climate

change in a broadly distributed marine crustacean. *Proc Biol Sci* **279**, 349–356 (2012).

51.   Leong, W., Sun, P. Y. & Edmands, S. Latitudinal Clines in Temperature and Salinity

Tolerance in Tidepool Copepods. *Journal of Heredity* **109**, 71–77 (2018).

52.   Foley, H. B. *et al.* Sex-specific stress tolerance, proteolysis, and lifespan in the

invertebrate Tigriopus californicus. *Experimental Gerontology* **119**, 146–156 (2019).

53.   Bartoń, K. *MuMIn: Multi-Model Inference*. (2020).

54.   Lenth, R. V. *emmeans: Estimated Marginal Means, aka Least-Squares Means*. (2020).

55.   Bates, D., Mächler, M., Bolker, B. & Walker, S. Fitting Linear Mixed-Effects Models Using

630   lme4. *Journal of Statistical Software* **67**, 1–48 (2015).

631   56.   Fox, J. & Weisberg, S. *An R Companion to Applied Regression*. (Sage, 2019).

632   57.   Gompertz, B. On the Nature of the Function Expressive of the Law of Human Mortality,

633   and on a New Mode of Determining the Value of Life Contingencies. (1815)

634   doi:10.1098/rspl.1815.0271.

635   58.   Jackson, C. flexsurv: A Platform for Parametric Survival Modeling in R. *Journal of*

636   *Statistical Software* **70**, 1–33 (2016).

637   59.   R Core Team. *R: A Language and Environment for Statistical Computing*. (R Foundation

638   for Statistical Computing, 2020).

639   60.   Canty, A. & Ripley, B. D. *boot: Bootstrap R (S-Plus) Functions*. (2020).

640   61.   Davison, A. C. & Hinkley, D. V. *Bootstrap Methods and Their Applications*. (Cambridge

641   University Press, 1997).

642   62.   Arnholt, A. T. & Evans, B. *BSDA: Basic Statistics and Data Analysis*. (2017).

643   63.   Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina

644   sequence data. *Bioinformatics* **30**, 2114–2120 (2014).

645   64.   Dierckxsens, N., Mardulyn, P. & Smits, G. NOVOPlasty: de novo assembly of organelle

646   genomes from whole genome data. *Nucleic Acids Research* **45**, e18 (2017).

647   65.   Bernt, M. *et al.* MITOS: Improved de novo metazoan mitochondrial genome annotation.

648   *Molecular Phylogenetics and Evolution* **69**, 313–319 (2013).

66.   Lunter, G. & Goodson, M. Stampy: A statistical algorithm for sensitive and fast mapping

of Illumina sequence reads. *Genome Res.* **21**, 936–939 (2011).

67.   Picard Tools - By Broad Institute. https://broadinstitute.github.io/picard/.

68.   Li, H. A statistical framework for SNP calling, mutation discovery, association mapping

and population genetical parameter estimation from sequencing data. *Bioinformatics* **27**,

2987–2993 (2011).

69.   Darling, A. E., Mau, B. & Perna, N. T. progressiveMauve: Multiple Genome Alignment

with Gene Gain, Loss and Rearrangement. *PLoS ONE* **5**, e11147 (2010).

70.   Ranwez, V., Douzery, E. J. P., Cambon, C., Chantret, N. & Delsuc, F. MACSE v2: Toolkit for

the Alignment of Coding Sequences Accounting for Frameshifts and Stop Codons. *Mol Biol*

Evol **35**, 2582–2584 (2018).

71.   Larkin, M. A. *et al.* Clustal W and Clustal X version 2.0. *Bioinformatics* **23**, 2947–2948

(2007).

72.   Yang, Z. PAML 4: Phylogenetic Analysis by Maximum Likelihood. *Mol Biol Evol* **24**, 1586–

1591 (2007).

73.   Kofler, R. *et al.* PoPoolation: A Toolbox for Population Genetic Analysis of Next

Generation Sequencing Data from Pooled Individuals. *PLoS ONE* **6**, (2011).

74.   Dowling, D. K. & Adrian, R. E. Challenges and Prospects for Testing the Mother's Curse

Hypothesis. *Integrative and Comparative Biology* **59**, 875–889 (2019).

75.   Zhang, H., Guillaume, F. & Engelstädter, J. THE DYNAMICS OF MITOCHONDRIAL

MUTATIONS CAUSING MALE INFERTILITY IN SPATIALLY STRUCTURED POPULATIONS:

MITOCHONDRIA, MALE INFERTILITY & SPATIAL STRUCTURE. *Evolution* **66**, 3179–3188

(2012).

76.   Vittor, B. A. Effects of the environment of fitness-related life history characters in

Tigriopus californicus. (University of Oregon, 1971).

77.   Burton, R. S. Mating system of the intertidal copepod Tigriopus californicus. *Marine*

Biology **86**, 247–252 (1985).

78.   Singh, S. R., Singh, B. N. & Hoenigsberg, H. F. Female remating, sperm competition and

sexual selection in Drosophila. *Genet Mol Res* **1**, 178–215 (2002).

79.   Palmer, C. A. & Edmands, S. Mate choice in the face of both inbreeding and outbreeding

depression in the intertidal copepod Tigriopus californicus. *Marine Biology* **136**, 693–698

(2000).

80.   Burton, R. S. Genetic Evidence for Long Term Persistence of Marine Invertebrate

Populations in an Ephemeral Environment. *Evolution* **51**, 993 (1997).

81.   Dybdahl, M. F. Extinction, recolonization, and the genetic structure of tidepool copepod

populations. *Evol Ecol* **8**, 113–124 (1994).

82.   Orr, H. A. Genetics of sterility in hybrids between two subspecies of Drosophila.

Evolution **43**, 180–189 (1989).

83.   Turelli, M. & Orr, H. A. Dominance, Epistasis, and the Genetics of Postzygotic Isolation.

Genetics **154**, 1663–1679 (2000).

84.   Pichaud, N., Ballard, J. W. O., Tanguay, R. M. & Blier, P. U. Mitochondrial haplotype

divergences affect specific temperature sensitivity of mitochondrial respiration. *J Bioenerg*

Biomembr **45**, 25–35 (2013).

85.   Strehler, B. L. & Mildvan, A. S. General theory of mortality and aging. *Science* **132**, 14–21

(1960).

86.   Li, N., Arief, N. & Edmands, S. Effects of oxidative stress on sex-specific gene expression

in the copepod Tigriopus californicus revealed by single individual RNA-seq. *Comparative*

Biochemistry and Physiology Part D: Genomics and Proteomics **31**, 100608 (2019).

87.   Li, N., Flanagan, B. A., Partridge, M., Huang, E. J. & Edmands, S. Sex differences in early

transcriptomic responses to oxidative stress in the copepod Tigriopus californicus. *BMC*

Genomics **21**, 759 (2020).

SUPPLEMENTAL INFORMATION

703   Supplemental Information includes one document, three figures, and six tables and can be

704   found with this article online at xxx.

Backcross	Dam line	Sire line	Nuclear F_{ST}	mtDNA D_{XY}	mtDNA d_N	mtDNA d_S	% Paternal nDNA	% Maternal mtDNA	% Paternal mtDNA
SDxSD	SD	SD	0	0	0	0	100.00	94.09	94.09
BRxSD	BR	SD	0.102	0.096	0.023	0.532	99.90	97.22	0.28
LJxSD	LJ	SD	0.108	0.096	0.023	0.523	99.97	97.63	0.15
FHLxSD	FHL	SD	0.378	0.206	0.085	2.675	99.70	99.65	0.13
BRxBR	BR	BR	0	0	0	0	100.00	96.91	96.91
LJxBR	LJ	BR	0.049	0.007	0.002	0.046	99.93	95.96	0.08
SDxBR	SD	BR	0.102	0.096	0.023	0.532	99.72	95.32	0.02
FHLxBR	FHL	BR	0.369	0.211	0.085	4.347	99.15	99.81	0.02

Figure 1. Construction of mitonuclear hybrid panel

(A) Geographic location of populations used in this study – SD (San Diego, CA), BR (Bird Rock,

CA), LJ (La Jolla, CA), FHL (Friday Harbor, WA). (B) Experimental design. Dams from the central

panel were backcrossed to sires from BR and SD for 10 generations. Grey arrows indicate

experimental control lines. Individuals from each line were reared at two contrasting

temperatures, 15℃ and 25℃. (C) DNA sequencing results from pool-seq. Measures of

divergence (F_{ST}, D_{XY}, d_N, and d_S) were calculated between dam and sire line. Percent

paternal nDNA and percent maternal/paternal mtDNA indicate minimal contaminating

nDNA remaining in backcrossed individuals and the absence of paternal leakage of mtDNA.

B

Trait	Temp.	nDNA	mtDNA	Sex	mtDNA x nDNA	Sex x mtDNA	Sex x nDNA	Sex x mtDNA x nDNA
Fertility	15℃	***	*	**	-	-	**	-
Fertility	25℃	NS	NS	NS	NS	NS	NS	*
Longevity	15℃	NS	NS	NS	NS	NS	NS	NS
Longevity	25℃	*	NS	NS	*	NS	NS	NS

Figure 2. Trait modeling results.

(A) Model contrasts between the sexes were plotted for each sex-effect model (Sex, Sex x mtDNA, Sex x nDNA, and Sex x mtDNA x nDNA). Fertility contrasts are represented as circles and longevity contrasts as triangles. Points to the left of the dotted zero-line

indicate a higher mean in males and those to the right of the line indicate higher mean in females. Significant contrasts were colored to indicate the sex with the higher mean - red (higher female) or blue (higher male).

(B) Sex-effect model type III ANOVA table. Each row represents a separate model, with columns representing modeled fixed effects. Cells are marked with a dash when a variable is not included in the model. Abbreviations: NS – non significant ($p < 0.05$), ✱ – $p \leq$ 0.05, ✱✱ – $p \leq 0.01$, ✱✱✱ – $p \leq 0.001$.

(C) Genetic and temperature effect size comparisons between fitness traits. We estimated Cohen's *d* for all pairwise contrasts among temperature, mtDNA, and nDNA groupings. Contrasts were made on best fit linear mixed effects models for each sex and trait grouping (sex-specific fertility, longevity).   Significance was determined using two-sample unpaired Wilcoxon rank sum tests with a two-sided alternate hypothesis and 95% confidence intervals. Sample sizes for contrasts were mtDNA, n = 96; nDNA, n = 32; and Temperature, n = 32.

Figure 3. Males outlive females and have lower rates of senescence and higher frailty.

(A) Sex-specific survivorship curves based on nDNA and temperature groupings. Median

longevity shown in insert with p-values indicating results of two-sample unpaired Wilcoxon rank

sum tests with a two-sided alternate hypothesis and 95% confidence intervals.

(B) Sex contrasts for Gompertz parameters. We performed Wilcoxon tests as above to identify

parametric differences in the rate of aging between the sexes.

Figure 4. Testing the mother's curse hypothesis with mitonuclear hybrid _T. californicus._

(A) Sex differences in mitochondrial coefficient of variation (CV_{mt}) for fertility and longevity. We

estimated CV_{mt} for both fertility and median longevity by temperature and nuclear line and

performed a two-sided sign test on bootstrapped estimates (1000 samples). The sign test

compares the difference between female and male median values with negative values

indicating higher male CV_{mt}, while positive values indicate that females have higher $CV_{mt.}$. The

majority of groupings indicate higher mitochondrial genetic variation in females in contradiction to the mother's curse hypothesis. Only fertility in BR nDNA individuals reared at 15°C had significantly higher mitochondrial genetic variation in males.

(B) Intra- and intersexual genetic correlation for female and male fertility and longevity across mitochondrial haplotypes. Bootstrapped Pearson's correlation coefficients (95% confidence intervals) were plotted for both rearing temperatures (15℃: solid lines, 25℃: dotted lines). Abbreviations: F (females), M (males), fertility (females: hatching number, males: remating success), longevity (mean longevity). All significant intersexual correlations indicate positive values, suggesting the absence of sexually antagonistic pleiotropy predicted by the mother's curse hypothesis.