1   **Whole genome sequencing suggests that "non-pathogenicity on banana (NPB)" is**
2   **the ancestral state of the *Ralstonia solanacearum* IIB-4 lineage.**
3

4   Authors

5   Jonathan Beutler, Samuel Holden, Stratton Georgoulis, Darrielle Williams, David J. Norman,
6   Tiffany M. Lowe-Power*
7

8   Affiliations:

Name	Affiliation (current)	Affiliation (time of work)	ORCid
Jonathan Beutler	Faculty of Land and Food Systems, The University of British Columbia, Vancouver, BC, Canada	Department of Plant Pathology, University of California Davis, Davis, CA 95616	0000-0002-1641-537X
Samuel Holden		Land and Food Systems, University of British Columbia, Vancouver, BC, Canada	
Stratton J. Georgoulis		Department of Plant Pathology, University of California Davis, Davis, CA 95616	0000-0002-2733-5463
Darrielle Williams		Department of Plant Pathology, University of California Davis, Davis, CA 95616	
David J. Norman		UF/IFAS Extension, Mid-Florida Research and Education Center, Apopka, FL 32703	
Tiffany M. Lowe-Power		Department of Plant Pathology, University of California Davis, Davis, CA 95616	0000-0003-2681-3563

9
10   *Corresponding author: tlowepower@ucdavis.edu
11

Abstract

The bacterial wilt pathogens in the *Ralstonia solanacearum* species complex (RSSC) have broad but finite host ranges. Population genetic surveys of RSSC pathogens show that many sequevars (subspecies groups) are predominantly recovered from wilting solanaceous plants. In contrast, strains in the IIB-4 sequevar have been isolated from plants in over a dozen families. Certain IIB-4 lineages have been classified as banana-virulent or "not pathogenic to banana (NPB)". Prior analysis suggested that the NPB lineage has diverged from the banana-virulent IIB-4 strains. To test this model, we analyzed the phenotypes and phylogeny of a diverse collection of 19 IIB-4 isolates. We used Illumina sequencing to assemble draft genomes of 12 new strains. Based on whole genome phylogenetic analysis, these IIB-4 strains clustered into five subclades. We quantified virulence of each strain on tomato, banana, melon, and impatiens plants. Overall, the virulence patterns correlated with phylogeny. Banana virulence was restricted to the 4/4 IIB-4D subclade (N=4/4 strains) and IIB-4E subclade (N=1/2 strains). Subclades IIB-4D and IIB-4E are sister subclades and their closest relative, the IIB-4A-C subclade, lacked virulence on banana. Our data support a revised model in which banana virulence is an innovation within the IIB4D/E subclades.

Data Summary

Illumina sequencing and genome assembly data are available as NCBI BioProject PRJNA826884, and Table S1 lists the accession numbers for assemblies in GenBank and raw sequencing data in SRA. To enable future meta-analyses that identify genetic factors that drive host-range, the raw virulence data is included as Table S2.

Introduction

The *Ralstonia solanacearum* species complex is a group of related tropical and temperate pathogens that cause plant wilt diseases after they invade and colonize the xylem (Prior et al. 2016; Lowe-Power et al. 2018; Ingel et al. 2022). Collectively, the species complex has a remarkably broad host range, capable of infecting hundreds of host species in dozens of taxonomic families (Hayward 1991; Lowe-Power et al. 2020). However, the breadth in host range is not uniformly distributed across the species complex. Strains in the *R. solanacearum* phylotype IIB sequevar 4 clade (IIB-4) exhibit a particularly broad host range (Ailloud et al. 2015; Wicker et al. 2007) (Lowe-Power et al. 2020). *R. solanacearum* IIB-4 strains have been isolated from 33 plant species in 19 botanical families (Wicker et al. 2007; Norman et al. 2009; Cellier et al. 2012; Hong et al. 2012; Cellier et al. 2015; Gutarra et al. 2017; Ramírez et al. 2020; Zulperi et al. 2014; Lowe-Power et al. 2020; Cellier and Prior 2010). They are distributed throughout Central and South America, the ancestral range of phylotype II *Ralstonia*. This lineage is also globally invasive as IIB-4 strains have become established in Malaysia (Zulperi et al. 2014) and have been imported into Florida (Norman et al. 2009).

R. solanacearum IIB-4 have caused several well-studied wilt epidemics. In the 1960s, *R. solanacearum* IIB-4 caused a high-impact Moko disease epidemic of plantains and bananas in Colombia and Peru (French and Sequeira 1970; Buddenhagen and Elsasser 1962). These Moko isolates were also virulent on solanaceous crops: tomato, pepper, and potato (French and Sequeira 1970). In the 2000s, a subgroup of *R. solanacearum* IIB-4 from Martinique were recognized as a novel "not pathogenic to banana (NPB) ecotype" with high virulence on cucurbits but low virulence to banana (Wicker et al. 2007). In 2020, a third ecotype of IIB-4 from Colombia was described (Ramírez et al. 2020); these novel Colombian strains are the only IIB-4 strains that are known to be avirulent on tomato. Overall, the well-documented natural history of IIB-4 strains and the diversity of their hosts position *R. solanacearum* IIB-4 as a model clade

for investigating the factors that limit and enable the breadth of host range in bacterial wilt

pathogens.

Our prior comparative genomic analysis of 14 phylotype II strains, including 5 IIB-4 strains, was

consistent with a hypothesis that the NPB lineage diverged from the banana-virulent IIB-4

strains. To test this hypothesis, we collected IIB-4 isolates from additional sources, inferred

their phylogeny by whole-genome sequencing and assembly, and analyzed their virulence

phenotypes on host species from four botanical families: banana, tomato, melon, and

impatiens.

Results

We quantified the natural variation in virulence of 19 *Ralstonia solanacearum* IIB-4 strains

against four natural host species from four botanical families. The strains were isolated from

diverse plants and locations (see Table S1). Seven strains were isolated in Peru and Colombia

from 1960-1963. Of these, six were from plantain during an epidemic of Moko disease, and one

from *Heliconia* (Buddenhagen and Elsasser 1962; French and Sequeira 1970). Three strains were

isolated on *Anthurium* and *Heliconia* during the Martinique epidemic (Wicker et al. 2007).

Seven strains were isolated from ornamentals imported into Florida. Of these, six were isolated

between 1996-2012 from *Pothos*, and one from *Anthurium* in 1997 (Norman et al. 2009). One

strain was isolated in 1999 from cucumber in Brazil (Ailloud et al. 2015), and one in 2017 from

Pothos in the Dominican Republic.

To determine the diversity of these strains, we used Illumina sequencing to assemble draft

genomes of 12 strains. An additional four unpublished genomes were shared by Caitilyn Allen

and Boris Vinatzer. The draft genome assembly of P488 is lower quality than others. The P488

assembly is fragmented into 200 contigs with an N_{50} of ~56 kb while the other assemblies have

40-115 contigs with an N_{50} of 111-414 kb. Full genome assembly statistics are in Table S1. We

collected genomes of five additional IIB-4 strains, the closely related IIB-51 strain CFBP7014,

and two distantly related strains (phylotype I GMI1000 and phylotype IIC K60 (Sharma et al.

2022)) from NCBI.

We constructed a phylogenetic tree using 4,317 genes comparing 369,907 intragenic SNPs (Fig

1). Three of the IIB-4 genomes had longer branch lengths than expected (P488, CFBP6783, and

UW179), consistent with the lower-quality, more fragmented assemblies for these strains. The

tree separated the IIB-4 strains into five branches, which we refer to as subclades A to E. The

Heliconia isolate from 1961 (UW170) was the sole genome on subclade A. Seven *Pothos* and

Anthurium isolates from Florida, and one from the Dominican Republic all clustered in subclade

B. The Brazilian cucumber isolate IBSBF1503 clustered into subclade C with the three strains

from the Martinique epidemic. The four Peruvian strains from the 1960s Moko epidemic

clustered into subclade D. Subclade E contained three Colombian strains isolated from plantain

in the 1960s, three Colombian strains recently isolated from plantain, and one Mexican strain

isolated from potato.

We quantified virulence of 19 IIB-4 strains on four diverse plant species that are documented

natural hosts for *Ralstonia* IIB-4: tomato (*Solanum lycopersicum* cv. Moneymaker), melon

(*Cucumis melo* cv. Sweet Granite), impatiens (*Impatiens walleriana* cv. Beacon Orange), and

banana (AAA *Musa acuminata* cv. Dwarf Cavendish). Tomato, melon, and impatiens were

grown from seed while banana plants were purchased as tissue cultured plantlets. Plants were

grown in growth chambers at 28^{o}C (tomato, melon, and impatiens) or 32^{o}C (banana) with a 12

hr light cycle and 40% humidity. Tomato and melon seeds were sown into Sunshine Mix #1 to a

depth of 1 cm, and impatiens were sown onto the surface of Sunshine Mix and topped with 1

mm vermiculite. Seedlings were transplanted into 3.5" pots with approximately 60 g Sunshine

Mix at 7 days (melon), 14 days (tomato), 21 days post sowing (impatiens), or at the 3-leaf stage

(banana).

We inoculated plants at 21 days post sowing (tomato and melon), 35 days post sowing

(impatiens), or at the 4-leaf stage (banana). We inoculated tomato and impatiens via a standard

cut-petiole approach where the lowest petiole was excised, and a 2 µl droplet of bacterial

suspension (containing approximately 1000 CFU) was placed on the cut surface (Khokhani et al.

2018). Due to rapid wound responses of cucurbits (Zimmermann et al. 2013), we inoculated

melons by applying a 2 µl droplet of 1000 CFU onto a razor blade and excising a petiole with the

infested blade. Banana plants were inoculated by drenching the soil with a bacterial suspension

(50 mL with $5x10^7$ to $1x10^8$ CFU/g soil) after the roots were lightly wounded with a 2 cm-wide

metal spatula (Ailloud et al. 2016). We rated symptoms daily for 14 days (Fig. 2). We rated

tomato symptoms on a standard 0-4 disease index (DI) where 0 = no wilt, 1 = 0.1-25% of leaflets

wilted, 2 = 25.1-50% of leaflets wilted, 3 = 50.1-75% of leaflets wilted, and 4 = 75.1-100% of

leaflets wilted (Khokhani et al. 2018). We measured symptoms on melon, impatiens, and

banana as the percentage of leaves with wilt. Symptom measurements, area under the disease

progress curve (AUDPC), and incidence are displayed in Table S2.

Although all strains were pathogenic to tomato, three strains had reduced virulence: UW170

(subclade A), UW162 (subclade D), and UW175 (subclade E). Because these strains have been

stored as "waterstocks" (in distilled water at room temperature) for decades, we suspect that

these strains acquired mutations in culture storage that reduced their virulence. The subclade A

strain UW170 was moderately virulent on both tomato and impatiens but caused minor wilt on

only a few banana or melon plants. The subclade D strain UW162 had moderate virulence on

tomato and mild virulence on impatiens, melon, and banana. The subclade E strain UW175 had

moderate virulence on tomato, mild virulence on impatiens and melon, and no virulence on

banana.

Virulence of strains generally correlated with their phylogenetic relationships. The subclade B

and C strains lacked virulence on banana but highly virulent on tomato, impatiens, and melon.

The subclade C strains displayed slightly higher virulence on melon than the subclade B strains.

Three other subclade D strains were highly virulent on tomato and had low-to-moderate

virulence on impatiens, melon, and banana. The subclade E strains included UW70 which was

highly virulent on tomato, melon, and banana. UW70 was more virulent on impatiens than

subclade D strains, but less virulent than the subclade B and subclade C strains.

Discussion

Pathogen populations are collections of genotypically and phenotypically diverse individuals. By

inferring phylogenetic relationships of strains, we can set the groundwork for genome-wide

analyses that associate genotypic differences with phenotypes. For plant pathogenic *Ralstonia*,

the sequevar system has been widely used to classify strains to sub-species groups (Fegan and

Prior 2005; Lowe-Power et al. 2020). Although the sequevar system is merely based on

sequence alignments of a single 750 bp DNA marker, we recently validated that sequevars for

Ralstonia phylotype II (i.e. *Ralstonia solanacearum*) accurately reflect the strains' core genome

phylogenic trees (Sharma et al. 2022). However, the sequevar system broke down for the

phylotype I branch of *Ralstonia pseudosolanacearum* (Sharma et al. 2022), which is known to

be highly recombinogenic (Wicker et al. 2012; Peeters et al. 2013). Moving forward, bacterial

taxonomy will continue to be influenced by whole-genome comparisons (Parks et al. 2020).

Taxonomic systems like the Life Identification Number (LIN) system (Tian et al. 2020; Vinatzer et

al. 2017) are likely to supersede the sequevar system. LINs are a systematic taxonomy based on

multiple levels of whole genome average nucleotide identity (ANI). Strains in the IIB-4 sequevar

have pairwise ANI of over 99.5% on their shared genes, and they are circumscribed in the LIN

group $14_A1_B0_C0_D0_E3_F0_G0_H1_I0_J0_K0_L0_M$ (Sharma et al. 2022).

Although many *Ralstonia* isolates only wilt plants efficiently at tropical temperatures, some

Ralstonia isolates can wilt plants at temperate temperatures. Cool-virulence is famously a trait

of the global pandemic lineage of *Ralstonia* (IIB-1 AKA the U.S. Select Agent-regulated "Race 3

Biovar 2" group)(Champoiseau et al. 2009), but recent studies have shown that multiple

lineages of *Ralstonia* can wilt tomato plants at cool temperatures: IIB-27, III-48, and the focal

clade of this study, IIB-4 (Bocsanczy et al. 2014, 2017). Although we carried out this study's

virulence assays at warm, tropical temperatures, several of our isolates, P673 (subclade B),

CFBP6783 (subclade C), and UW163 (subclade D) can wilt tomatoes at cool temperatures (18°C)

(Bocsanczy et al. 2014, 2017).

Early studies suggested that IIB-4 strains have binary host range phenotype with either a

"Moko" phenotype with virulence on banana/ tomato/ plantain or a "NPB" phenotype with

virulence on tomato/ plantain/ anthurium/ cucurbits (Ailloud et al. 2015; Wicker et al. 2007;

Ailloud et al. 2016). However, our study shows that the virulence patterns are more complex.

First, we show that banana virulence is rare among the strains tested. The NPB phenotype was

shared by the subclade C strains previously described as "not pathogenic to banana" (Ailloud et

al. 2015) and strains from subclade A and B. Our results suggest that virulence on banana

might be an acquired trait within the subclade D/E lineage. Consistently, virulence on banana is

a polyphyletic trait shared by distantly related lineages in phylotypes IIA, IIB, and IV (Ray et al.

2021; Albuquerque et al. 2014; Obrador-Sánchez et al. 2017). Our study did not include any

genomes from the recently identified IIB-4 strains from Colombia that lacks virulence on

tomato (Ramírez et al. 2020). If those strains branch independently from the five subclades of

the 24 IIB-4 strains in this study, we propose to name the newly described strains as "subclade

F".

One subclade E strain (UW70 also known as "CIP417") was highly virulent on all host species

tested. UW70 was isolated from plantain in Colombia in the 1960s and is closely related to

strains isolated from plantain in Colombia in 2012 (UA-1591 and UA-1617)(Ramírez et al. 2020).

Interestingly, UW70 was highly virulent on melons, but UA-1591 is reported to be avirulent on

cucumber (Ramírez et al. 2020). Cucurbit virulence has also been documented in phylotype I

strains isolated in Taiwan, Thailand, China, and Japan (He et al. 2021; Yahiaoui et al. 2017;

Horita et al. 2014; Lin et al. 2014).

Traits of several of this study's isolates were studied: subclade C strain CFBP6783, subclade D

strain UW163 and subclade E strain UW70. Lebeau et al. (2011) measured virulence of

CFBP6783 and 11 other diverse *Ralstonia* strains against 30 accessions of tomato, eggplant, or

pepper accessions in the "Core-TEP" panel. In this study, CFBP6783 exhibited a unique pattern

of high virulence on 10/10 tomato, 10/10 pepper, and 6/10 eggplant accessions. CFBP6783 is

virulent on plantain (AAB *Musa* sp. cv. Dominico-Harton)(Wicker et al. 2007) and potato (cv.

Bintje Munterschen × Franschen)(Cellier and Prior 2010), but is not virulent on Cavendish

banana (Wicker et al. 2007 and this study). UW70 and UW163 were highly virulent on *Musa*

balbisiana Colla, tomato (*Solanum lycopersicum* L. 'Bonny Best'), pepper (*Capsicum annuum* L.

'Sweet Yellow'), and potato (*Solanum tuberosum* L. 'Russet Burbank')(French and Sequeira

1970). In contrast, they exhibited low virulence on eggplant (*Solanum melongena* L. 'Black

Beauty') and tobacco (*Nicotiana tobacum* L. 'Bottom Special'). In a separate study, UW163 (D

subclade) and UW179 (E subclade) lacked virulence on Anthurium while subclade C strains

CFBP6783 and IBSBF1503 were virulent on Anthurium (Ailloud et al. 2015).

Our findings reinforce the long-recognized phenotypic plasticity within the *Ralstonia*

solanacearum species complex (Buddenhagen 1985). In addition to host range variation within

sequevar IIB-4, banana-virulent strains in IIA-24 and IIA-41 clades vary in their virulence on

tomato (Albuquerque et al. 2014). A study of phylotype I *Ralstonia* from Guanxi, China showed

that closely related strains shared virulence on tomato, eggplant, tobacco, and potato but

varied in their virulence on pepper, mulberry, ginger, and cucurbits (He et al. 2021).

In attempts to classify strains by their host range, scientists have attempted to sort *Ralstonia*

into "Races", "Pathoprofiles", "Pathotypes", and "Ecotypes" (Lebeau et al. 2011; Cellier et al.

2012; Buddenhagen 1985; Peeters et al. 2013). Having clear phenotypic classifications would be

both intrinsically satisfying and immensely helpful for clarifying regulations. In the future, this

may be possible through sophisticated statistical analysis. However, these data science

approaches will require a concerted and sustained effort to generate robust data on the

genomic and phenotypic diversity of the global *Ralstonia* population. In parallel, a systematic

database should be developed to consolidate and communicate knowledge on host range of

different *Ralstonia* lineages. Continued study is needed to gain deeper insight into the genetic

and evolutionary drivers of host range in the *Ralstonia solanacearum* species complex

Acknowledgements

We thank Caitilyn Allen for sharing strains used in this study and Caitilyn Allen, Boris Vinatzer,

and Parul Sharma for sharing unpublished genomes. Work was funded by a Henry A. Jastro

Graduate Research Award to J. Beutler; the UC Davis College of Agricultural and Environmental

Sciences and USDA CA-D-PPA-2610-H to T. Lowe-Power; the Plant Agricultural Biology Graduate

Admissions Pathways Program to D. Williams; the USDA Floriculture and Nursery Research

Initiative and the University of Florida Institute of Food and Agricultural Sciences to D. Norman.

Conflicts of Interest

The authors declare that there are no conflicts of interest.

References

Ailloud, F., Lowe, T., Cellier, G., Roche, D., Allen, C., and Prior, P. 2015. Comparative genomic
analysis of *Ralstonia solanacearum* reveals candidate genes for host specificity. BMC Genomics.
16:270.
Ailloud, F., Lowe, T. M., Robène, I., Cruveiller, S., Allen, C., and Prior, P. 2016. *In planta*
comparative transcriptomics of host-adapted strains of *Ralstonia solanacearum*. PeerJ.
4:e1549.
Albuquerque, G. M. R., Santos, L. A., Felix, K. C. S., Rollemberg, C. L., Silva, A. M. F., Souza, E. B.,
et al. 2014. Moko disease-causing strains of *Ralstonia solanacearum* from Brazil extend known
diversity in paraphyletic Phylotype II. Phytopathology. 104:1175–1182.
Bayliss, S. C., Thorpe, H. A., Coyle, N. M., Sheppard, S. K., and Feil, E. J. 2019. PIRATE: A fast and
scalable pangenomics toolbox for clustering diverged orthologues in bacteria. Gigascience. 8
Available at: https://academic.oup.com/gigascience/article/8/10/giz119/5584409 [Accessed
May 5, 2021].
Bocsanczy, A. M., Achenbach, U. C. M., Mangravita-Novo, A., Chow, M., and Norman, D. J. 2014.
Proteomic comparison of *Ralstonia solanacearum* strains reveals temperature dependent
virulence factors. BMC Genomics. 15:280.
Bocsanczy, A. M., Huguet-Tapia, J. C., and Norman, D. J. 2017. Comparative Genomics of
Ralstonia solanacearum Identifies Candidate Genes Associated with Cool Virulence. Front. Plant
Sci. 8:1565.
Buddenhagen. 1985. Bacterial wilt revisited. Bacterial wilt disease in Asia and the. Available at:
https://ageconsearch.umn.edu/record/134643/files/PR013.pdf#page=124.
Buddenhagen, I. W., and Elsasser, T. A. 1962. An insect-spread bacterial wilt epiphytotic of
bluggoe banana. Nature. 194:164–165.
Cellier, G., Moreau, A., Chabirand, A., Hostachy, B., Ailloud, F., and Prior, P. 2015. A duplex PCR
assay for the detection of *Ralstonia solanacearum* phylotype II strains in *Musa* spp. PLoS One.
10:e0122182.
Cellier, G., and Prior, P. 2010. Deciphering Phenotypic Diversity of *Ralstonia solanacearum*
Strains Pathogenic to Potato. Phytopathology®. 100:1250–1261.
Cellier, G., Remenant, B., Chiroleu, F., Lefeuvre, P., and Prior, P. 2012. Phylogeny and
population structure of brown rot- and Moko disease-causing strains of *Ralstonia solanacearum*
phylotype II. Appl. Environ. Microbiol. 78:2367–2375.
Champoiseau, P. G., Jones, J. B., and Allen, C. 2009. *Ralstonia solanacearum* Race 3 Biovar 2
causes tropical losses and temperate anxieties. Plant Health Prog. Available at:
https://apsjournals.apsnet.org/doi/abs/10.1094/PHP-2009-0313-01-RV.
Fegan, M., and Prior, P. 2005. How complex is the *Ralstonia solanacearum* species complex. In
Bacterial wilt disease and the Ralstonia solanacearum species complex, APS press St. Paul, p.
449–461.

French, E. R., and Sequeira, L. 1970. Strains of *Pseudomonas solanacearum* from Central and
South America: a comparative study. Phytopathology. 60:506–512.
Gutarra, L., Herrera, J., Fernandez, E., Kreuze, J., and Lindqvist-Kreuze, H. 2017. Diversity,
pathogenicity, and current occurrence of bacterial wilt bacterium *Ralstonia solanacearum* in
Peru. Front. Plant Sci. 8:1221.
Hayward, A. C. 1991. Biology and epidemiology of bacterial wilt caused by *Pseudomonas*
solanacearum. Annu. Rev. Phytopathol. 29:65–87.
He, Y., Chen, Y., Zhang, Y., Qin, X., Wei, X., Zheng, D., et al. 2021. Genetic diversity of *Ralstonia*
solanacearum species complex strains obtained from Guangxi, China and their pathogenicity on
plants in the Cucurbitaceae family and other botanical families. Plant Pathol. Available at:
https://onlinelibrary.wiley.com/doi/10.1111/ppa.13389.
Hong, J. C., Norman, D. J., Reed, D. L., Momol, M. T., and Jones, J. B. 2012. Diversity among
Ralstonia solanacearum strains isolated from the Southeastern United States. Phytopathology.
102:924–936.
Horita, M., Tsuchiya, K., Suga, Y., Yano, K., Waki, T., Kurose, D., et al. 2014. Current classification
of *Ralstonia solanacearum* and genetic diversity of the strains in Japan. J. Gen. Plant Pathol.
80:455–465.
Ingel, B., Caldwell, D., Duong, F., Parkinson, D. Y., McCulloh, K. A., Iyer-Pascuzzi, A. S., et al.
2022. Revisiting the source of wilt symptoms: X-ray microcomputed tomography provides
direct evidence that *Ralstonia* biomass clogs xylem vessels. Phytofrontiers. Available at:
http://dx.doi.org/10.1094/PHYTOFR-06-21-0041-R [Accessed April 9, 2021].
Khokhani, D., Tran, T. M., Lowe-Power, T. M., and Allen, C. 2018. Plant assays for quantifying
Ralstonia solanacearum virulence. Bio-protocol. Available at: https://en.bio-
protocol.org/CN/e3028.
Lebeau, A., Daunay, M.-C., Frary, A., Palloix, A., Wang, J.-F., Dintinger, J., et al. 2011. Bacterial
wilt resistance in tomato, pepper, and eggplant: genetic resources respond to diverse strains in
the *Ralstonia solanacearum* species complex. Phytopathology. 101:154–165.
Letunic, I., and Bork, P. 2021. Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic
tree display and annotation. Nucleic Acids Res. 49:W293–W296.
Lin, C.-H., Tsai, K.-C., Prior, P., and Wang, J.-F. 2014. Phylogenetic relationships and population
structure of *Ralstonia solanacearum* isolated from diverse origins in Taiwan. Plant Pathol.
63:1395–1403.
Lowe-Power, T., Avalos, J., Charco Munoz, M., and Chipman, K. 2020. A meta-analysis of the
known global distribution and host range of the *Ralstonia* species complex. bioRxiv. Available
at: https://www.biorxiv.org/content/10.1101/2020.07.13.189936v1.abstract.
Lowe-Power, T. M., Khokhani, D., and Allen, C. 2018. How *Ralstonia solanacearum* exploits and
thrives in the flowing plant xylem environment. Trends Microbiol. 26:929–942.
Norman, D. J., Zapata, M., Gabriel, D. W., Duan, Y. P., Yuen, J. M. F., Mangravita-Novo, A., et al.
2009. Genetic diversity and host range variation of *Ralstonia solanacearum* strains entering
North America. Phytopathology. 99:1070–1077.
Obrador-Sánchez, J. A., Tzec-Simá, M., Higuera-Ciapara, I., and Canto-Canché, B. 2017. Genetic
diversity of *Ralstonia solanacearum* strains from Mexico associated with Moko disease. Eur. J.
Plant Pathol. 149:817–830.

Parks, D. H., Chuvochina, M., Chaumeil, P.-A., Rinke, C., Mussig, A. J., and Hugenholtz, P. 2020. A
complete domain-to-species taxonomy for Bacteria and Archaea. Nat. Biotechnol. 38:1079–
1086.

Peeters, N., Guidot, A., Vailleau, F., and Valls, M. 2013. Ralstonia solanacearum, a widespread
bacterial plant pathogen in the post-genomic era. Mol. Plant Pathol. 14:651–662.

Prior, P., Ailloud, F., Dalsing, B. L., Remenant, B., Sanchez, B., and Allen, C. 2016. Genomic and
proteomic evidence supporting the division of the plant pathogen *Ralstonia solanacearum* into
three species. BMC Genomics. 17:90.

Ramírez, M., Moncada, R. N., Villegas-Escobar, V., Jackson, R. W., and Ramírez, C. A. 2020.
Phylogenetic and pathogenic variability of strains of *Ralstonia solanacearum* causing moko
disease in Colombia. Plant Pathol. 69:360–369.

Ray, J. D., Subandiyah, S., Rincon-Florez, V. A., Prakoso, A. B., Mudita, I. W., Carvalhais, L. C., et
al. 2021. Geographic Expansion of Banana Blood Disease in Southeast Asia. Plant Dis. 105:2792–
2800.

Sharma, P., Johnson, M. A., Mazloom, R., Allen, C., Heath, L. S., Lowe-Power, T. M., et al. 2022.
Meta-analysis of the *Ralstonia solanacearum* species complex (RSSC) based on comparative
evolutionary genomics and reverse ecology. Microb Genom. 8 Available at:
http://dx.doi.org/10.1099/mgen.0.000791.

Stamatakis, A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large
phylogenies. Bioinformatics. 30:1312–1313.

Tian, L., Huang, C., Mazloom, R., Heath, L. S., and Vinatzer, B. A. 2020. LINbase: a web server for
genome-based identification of prokaryotes as members of crowdsourced taxa. Nucleic Acids
Res. 48:W529–W537.

Vinatzer, B. A., Weisberg, A. J., Monteil, C. L., Elmarakeby, H. A., Sheppard, S. K., and Heath, L. S.
2017. A Proposal for a Genome Similarity-Based Taxonomy for Plant-Pathogenic Bacteria that Is
Sufficiently Precise to Reflect Phylogeny, Host Range, and Outbreak Affiliation Applied to
Pseudomonas syringae sensu lato as a Proof of Concept. Phytopathology. 107:18–28.

Wicker, E., Grassart, L., Coranson-Beaudu, R., Mian, D., Guilbaud, C., Fegan, M., et al. 2007.
Ralstonia solanacearum strains from Martinique (French West Indies) exhibiting a new
pathogenic potential. Appl. Environ. Microbiol. 73:6790–6801.

Wicker, E., Lefeuvre, P., de Cambiaire, J.-C., Lemaire, C., Poussier, S., and Prior, P. 2012.
Contrasting recombination patterns and demographic histories of the plant pathogen Ralstonia
solanacearum inferred from MLSA. ISME J. 6:961–974.

Yahiaoui, N., Chéron, J.-J., Ravelomanantsoa, S., Hamza, A. A., Petrousse, B., Jeetah, R., et al.
2017. Genetic diversity of the *Ralstonia solanacearum* species complex in the Southwest Indian
Ocean Islands. Front. Plant Sci. 8:2139.

Zimmermann, M. R., Hafke, J. B., van Bel, A. J. E., and Furch, A. C. U. 2013. Interaction of xylem
and phloem during exudation and wound occlusion in *Cucurbita maxima*. Plant Cell Environ.
36:237–247.

Zulperi, D., Sijam, K., Ahmad, Z. A. M., Awang, Y., and Mohd Hata, E. 2014. Phylotype
classification of *Ralstonia solanacearum* biovar 1 strains isolated from banana (*Musa* spp.) in
Malaysia. Archives of Phytopathology and Plant Protection. 47:2352–2364.

Figures

Fig 1. Maximum likelihood tree describing 27 strains of the *Ralstonia solanacearum* species complex, based on 369,907 intragenic SNPs. The tree includes 24 IIB-4 strains and three reference strains from phylotype IIB-51 (CFBP7014), IIC-7 (K60) (Sharma et al. 2022), and I (GMI1000). Strains are described as belonging to one of five IIB-4 subclades (A-E). 95% bootstrap support is indicated by a white circle. Some branch lengths were condensed for clarity. SNPs were identified by grouping orthologous genes across all isolates using PIRATE v1.0.4 with default parameters and nucleotide-based clustering (Bayliss et al. 2019). Orthologous gene groups identified by PIRATE were concatenated and reduced to positions present in at least 20 accessions, with at least one variable nucleotide, by a python script Phylip_reducer.py (github.com/Erysiphe-graminis/Utilities). These positions were used to generate the tree using RAxML v8.2.12 with the GTRGAMMA model, Lewis ascertainment bias correction, and 1000 bootstraps (Stamatakis 2014). GMI1000 was the outgroup. The tree was

380   visualized using ITOL v6.5.2 (Letunic and Bork 2021), and the figure prepared using Adobe

381   Illustrator. To evaluate the impact of excluding infrequently occurring genes, two additional

382   trees were generated using the same parameters except that SNP positions could be present in

383   (1) at least 10 accessions, or (2) in any number of accessions. These trees were topologically

384   identical with subtle variations in lengths of some branches.

Fig 2. Disease progress of 19 *R. solanacearum* IIB-4 strains on four host species: tomato cv.

Moneymaker (n≥44 plants), melon cv. Sweet Granite (n≥43), impatiens cv. Beacon Orange

(n≥44), and Banana cv. Dwarf Cavendish (n≥19). Each graph represents one strain. Virulence on

tomato was quantified on 0-4 disease index (left Y-axes), and virulence on other hosts was

quantified as a percentage of leaves with wilt (right Y-axes). Symbols represent the mean of 2-3

392   trials and bars show standard error of the mean (SEM). Strains are grouped based on the

393   phylogeny from Fig 1. Raw virulence data is available in Table S2.