HYBRIDIZATION IN THE LATE PLEISTOCENE: Merging Morphological and Genetic Evidence

Harvati K. [1,2] (*), Ackermann R.R.[3,4,2] (*)

1 Paleoanthropology section, Senckenberg Centre for Human Evolution and Palaeoenvironment,
Institute for Archaeological Sciences, Eberhard Karls Universität Tübingen
2 DFG Centre for Advanced Studies 'Words, Bones, Genes, Tools', Eberhard Karls Universität
Tübingen
3 Human Evolution Research Institute, University of Cape Town, South Africa
4 Department of Archaeology, University of Cape Town, South Africa
* Addresses for correspondence
Prof. Katerina Harvati
Paleoanthropology Section
Rümelinstrasse 23
Tübingen 72070
Germany
21
Prof. Rebecca R. Ackermann
Department of Archaeology
University of Cape Town
Private Bag X1
Rondebosch 7701
South Africa
28

This is an exciting time for our understanding of the origin of our species. Previous scientific consensus saw human evolution as defined by adaptive differences (behavioural and/or biological) and the emergence of *Homo sapiens* as the ultimate replacement of non-modern groups by a modern, adaptively more competitive one. However, recent research has shown that the process underlying our origins was considerably more complex. While archaeological and fossil evidence suggests that behavioural complexity may not be confined to the modern human lineage, recent paleogenomic work shows that gene flow between distinct lineages (e.g. Neanderthals, Denisovans, early *H. sapiens*) occurred repeatedly in the Late Pleistocene, likely contributing elements to our genetic make-up that might have been crucial to our success as a diverse, adaptable species. Following these advances, the prevailing human origins model has shifted from one of near-complete replacement to a more nuanced view of partial replacement with considerable reticulation. Here we provide a brief introduction to the current genetic evidence for hybridization among hominins, its prevalence in, and effects on, comparative mammal groups, and especially how it manifests in the skull. We then explore the degree to which cranial variation seen in the fossil record of Late Pleistocene hominins from Western Eurasia corresponds with our current genetic and comparative data. We are especially interested in understanding the degree to which skeletal data can reflect admixture. Our findings indicate some correspondence between these different lines of evidence, flag individual fossils as possibly admixed, and suggest that different cranial regions may preserve hybridisation signals differentially. We urge further studies of the phenotype in order to expand our ability to detect the ways in which migration, interaction and genetic exchange have shaped the human past, beyond what is currently visible with the lens of ancient DNA.

Current genetic evidence for gene exchange and its effects in hominins

Natural hybridization promotes evolutionary innovation, creating novel and diverse outcomes in subsequent generations, thereby providing a rich substrate on which selection can further act to shape evolutionary trajectories [1,2]. Since 2010, methodological advances allowing unprecedented, high-resolution insights into ancient genomes have provided increasing evidence for hybridization and resultant gene flow among Late Pleistocene humans. Currently, indications for gene exchange include movement of genes from Neanderthals into early *Homo sapiens* (conventionally called "early modern humans")[3-8], resulting in approximately 2-3% Neanderthal ancestry of non-African living modern humans [7]; as well as evidence that *Homo sapiens* contributed to the Neanderthal gene pool as early as 150 to >200ka[9,10]. Gene flow from Denisovans into the ancestors of modern Asian populations [11,12], from Neanderthals into Denisovans [13,14], and from some unknown hominin into Denisovans [13] has also been reported, and the genome of a first generation descendent of a Neanderthal mother and a Denisovan father living ca. 90ka was recently discovered [15]. Finally, genetic exchanges between ancient and recent lineages may have also occurred within Africa [9,16-21]. Taken together, these studies indicate that gene flow has been multidirectional, was much more common than previously appreciated by most (but see e.g. [22]), and may have been instrumental in structuring genetic diversity across our ancestral lineage over the last half a million years. Given the speed at which new discoveries and methodological breakthroughs are occurring, such as the retrieval of hominin DNA from cave sediments [14], our expectation is that such evidence will likely continue to accumulate in the future.

Gene flow among hominins has had variable effects, best documented over the last 100K
years. These include genetic evidence for some level of introgression affecting phenotypes in a
beneficial manner, including those involved in immunity, spermatogenesis, adaptation to low-
oxygen contexts, response to ultraviolet radiation, and other traits [23-31] (but see [32]). For example,
Neanderthal genes affecting skin and hair phenotypes are retained in humans living today, (e.g. [27,31]),
suggesting that these genes might have been important in the dispersal and adaptation of people
emerging from Africa and migrating into environments inhabited by Neanderthals. In other cases,
gene exchange may have been detrimental. For example, the existence of chromosomal regions in
living humans devoid of Neanderthal-derived alleles, such as the X-chromosome and genes related
to testes and therefore reproduction [27,31], suggests that selection may have acted to purge these
genes from descendants. Neanderthal alleles present in living people have also been associated
with a range of phenotypes considered detrimental in modern (but not necessarily ancient) contexts,
including depression, neurodevelopmental disorders, hypercoagulation, altered carbohydrate
metabolism, and addiction [27,29,33] (but see [32]). A few recent studies suggest that Neanderthal-derived
genetic variation also influences brain phenotypes [29,34,35] and susceptibility to infectious diseases
[36,37].

Taken together, the genetic evidence to date indicates that gene flow played an important
role in shaping the evolutionary fate of our lineage [38], although its exact effects appear to vary
considerably across time, population, and environmental/geographical context.

Hybridization in extant primates and other mammals and its relevance to hominins

Although the genetic evidence for hybridization in hominins has shifted the prevailing
narrative about human origins over the last decade, there was already a growing realisation prior to
this (e.g. [39,40]) that its role may have been underappreciated, based on an increasing understanding
of its prevalence across other mammals, including primates. We now know that approximately ten
percent of animal species produce hybrids, and occasionally "phylogenetic hotspots" occur in which
hybridization rates in animals exceed those seen in plants [41,42]. Within mammals, hybridization
occurs across a wide range of lineages, including (but not limited to) a number of large-bodied
terrestrial groups such as bovids [43-45]; bears [46,47]; cats [48,49]; canids [50-54] and primates (see below).
These studies have provided compelling evidence that gene flow impacts the evolutionary trajectory
of large-bodied mammals, acting as a particularly strong force for accelerating evolution in novel or
changing environmental contexts [1,2], a scenario that resonates with the narrative of human origins.

Non-human primates are arguably the most relevant models for human evolution, and there
is considerable evidence for hybridization in the wild within all the major lineages at both specific
and intraspecific levels including strepsirrhines (e.g. [55,56]), American monkeys (e.g. [57-61]) and Afro-
Eurasian monkeys (e.g. [62-66]). Among these, perhaps the best studied are baboons (genus *Papio)*,
which have also repeatedly been put forth as models for human evolution [40,67]. The six recognised
baboon species (or 'allotaxa'; see [40]) have parapatric ranges, with natural hybridization recorded
between the species that are most phylogenetically distant (*Papio ursinus* vs. *P. cynocephalus*),
morphologically distinct (*P. ursinus* vs. Kinda baboons), and behaviourally different (*P. hamadryas* vs.
P. anubis)[68]. Like our own genus *Homo*, *Papio* is the evolutionary product of a radiation that began in

non-forested regions of tropical Africa around two million years ago; both genera have inhabited
similar regions in Africa and been subject to comparable climatic fluctuations.

Hybridization has also occurred among our closest primate relatives, the apes (Superfamily
Hominoidea). It is well-documented among the small-bodied apes (e.g. [69,70]), and there are also
genetic signatures of gene flow both among subspecies (e.g. [71]) and between species [72] of great
apes. One percent of the central chimpanzee genome has been shown to derive from the bonobo [72],
indicating two ancient hybridization events comparable to the admixture seen between modern
humans and Neanderthals.

Hybridization can have a wide range of effects on anatomy, behaviour and speciation (see
[73]), but it is in its interplay with adaptation that its impact may be most powerful. However, the
impact of adaptive introgression can differ even among closely related taxa. For example, in
chimpanzees the regions of adaptive introgression are subspecies-specific (e.g. regions involving
male reproduction versus immune system) [74]. As a species-specific example, the region around the
FOXP2 locus is devoid of introgression in humans[75,76], but not in either chimpanzees [77] or bonobos [78].

How does gene exchange manifest itself in skeletal morphology?

Genetic evidence for the effect of gene flow on the hominin skeleton remains limited,
despite its importance in linking the genetic and fossil record, as well as potentially understanding
the functional implications of skeletal variation. Recent studies suggest that Neanderthal-derived
genetic variation influences shape variation in the crania and brains of Europeans living today [34,35]. In
particular, Neanderthal ancestry was found to be associated with a more Neanderthal-like,
elongated, cranial and endocranial shape in these Europeans, including morphology in the occipital
and parietal regions, as well as differences in brain morphology [34,35].

Genetic evidence aside, some researchers have proposed hybrid individuals in the human
fossil record on the basis of their morphology. Such proposed hybrids include Lagar Velho 1 [79],
Mladeč 5 and 6 [80], Cioclovina 1 [81], Peştera cu Oase 1 [82,83] and 2 [82,84], Skhul IV and V [84], Vindija [85],
Klasies River Mouth [86], Jebel Irhoud and Mughâret el 'Aliya in North Africa [86,87] and possibly others [88-
90]. However, these hypotheses have generally not been possible to test, and the hybrid status of
these specimens has been disputed [91-93] or considered inconclusive [94,95]. This was mainly due to the
lack of clear expectations about hybrid morphology that could be empirically applied to the fossil
record (but see [84,96-99]), but also because of the problem of equifinality, as phenotypes consistent
with hybridization, especially "intermediate" morphology, may also be produced through other
processes, most importantly by the retention of primitive features. In the face of these
shortcomings, admixed status has almost exclusively been recognized on the basis of genetic
evidence (as discussed above). However, such evidence is limited in many respects. For example, the
application of ancient DNA is constrained due to preservation issues, which can vary from site to site
and specimen to specimen, but become particularly severe as we move further back in time or into
warmer climates. Additionally, knowledge derived from comparisons among extant genomes can
only provide partial insight into the past, given the extinction of many ancient lineages. Therefore,
evidence for hybridization present in the skeletal phenotype remains essential to the interpretation

of the fossil record, as it can help us to locate such potential events in time and place, and
particularly within lineages for which we do not have a genetic record.

The taxon for which most empirical evidence for the effects of hybridization on the skeleton
is available is baboons. Studies of baboons have revealed visible perturbations in dental and sutural
formation at high frequencies in early generation inter-specific hybrids, as well as atypical expression
of some dental traits [1,96,97], suggesting that hybridization breaks down the coordination of early
development, although this does not appear to meaningfully affect fitness [84,97,100]. These results are
consistent with what is seen in the skeletal anatomy of hybrids in other mammalian lineages,
including ungulates [43], rodents [101], and most recently canids [102], although they manifest somewhat
differently in each taxon. Hybrid baboons also have, on average, crania that are larger than an
intermediate value between their parents [96,97,103], with some measurements that are extreme
relative to both parents. The production of extreme hybrid phenotypes, or transgressive
phenotypes, outside of the range of both parental taxa (in a negative or positive direction) is called
transgressive segregation [104], and in the case of the mammals mentioned above could include
atypical traits as well as extreme size/shape.

Large cranial size in hybrids (relative to either a parental midpoint value, or the mean of the
largest parent) has also been identified in mice [98,105-110] (as well as tamarins [111]). Inter-subspecific
mouse hybrids (F1s, F2s, and backcrossed individuals) are typically as large as or larger than the
larger parent taxon, with associated size-related shape changes [98,110]. They also tend to have cranial
and mandibular shape variation that is somewhat intermediate to the parents, but more closely
resembling the smaller parent (Figure 1), with high levels of heterosis in certain features such as
molar length [98,110]. Later generations (F2, B2) are more variable than first generation hybrids, with
backcrosses expectedly moving towards the shape of the parent taxon with which they are
hybridizing. These patterns hold for crosses of taxa that hybridize in the wild but have low levels of
gene flow and low hybrid fertility; for taxa that hybridize in the wild and produce successful
offspring; and for taxa that are geographically separated in nature but nevertheless hybridize under
laboratory conditions [98,110], making them robust across different scenarios of contact and hybrid
fitness. Importantly, the fact that these models show similar patterns for hybridization across both
species and subspecies provides a robust model for assessing its impact in taxa where the specific
status is debated, such as Neanderthals and *H. sapiens* (See text box "Are Neanderthals Distinct
Species and Does It Matter?").

The studies above have focused on skulls. Unfortunately, considerably less information
exists on the effects of hybridization on the postcranial skeleton, outside of the observation from a
number of previous studies in mice and primates that hybrids generally exhibit both longer limbs
and increased body size relative to parents (e.g. [84,98,105,106,109,112-114]), though a new study of macaques
suggests that effects of admixture on the pelvis may be relatively small, possibly due to functional or
developmental constraints, or relatively minor divergence of the parent taxa (in this case at or
possibly below the subspecies level) [115].

Figure 1: Principal components analysis, mice crania (top) and mandibles (bottom). Red and Green represent parent taxa; Blue represents a pooled sample of first (F1), second (F2), and unidirectionally (with the green parent) backcrossed (B1) inter-subspecific hybrids. Data from [110].

Do

Do Late Pleistocene Western Eurasian humans fit the morphological predictions of a hybrid sample?

Although current genetic evidence indicates that hybridization occurred repeatedly among
Pleistocene hominins, there have been few efforts to link this genetic evidence to morphological
evidence from the fossil record itself, despite such a link being key for ascertaining the status and
relevance of the bulk of the fossil record (for which genetic data are not available). This is further
exacerbated by the fact that our ability to extrapolate from genotype to skeletal phenotype is
currently very limited. While it is true that some individuals that show genetic evidence of admixture
have limited morphology (e.g.[15]), making establishing these links difficult, this is not the case for
other specimens. This lack of discourse between morphology and genetics is detrimental to
understanding the dynamics of human evolution in the Late Pleistocene.

Here we explore how insights derived from genetics and model organisms might be applied
to the interpretation of the human fossil record. In particular, we examine the patterns of variation
in cranial shape across the late Middle to Late Pleistocene, interpreted in conjunction with published
genetic and non-metric phenotypic evidence for hybridisation, as discussed further below. The latter
evidence consists of sutural and dental developmental anomalies comparable to what has been
observed in comparative studies on hybridization and its effects on the phenotype. Even though
admixture between hominin lineages has been demonstrated outside of Western Eurasian contexts,
we focus specifically on Neanderthals and early *Homo sapiens*, i.e. hominins from Western Eurasia
and Africa, in order to limit the scope of this inquiry. In keeping with current consensus, we consider
these taxa to represent distinct lineages evolving in large part independently (see, e.g. [116]). However,
in recognizing the important limitations of species concepts and their application to the fossil record,

as well as long standing disagreements on Neanderthal alpha taxonomy, we avoid the term *Homo*
neanderthalensis, while using *Homo sapiens* to refer to extant humans and their ancestors in the late
Middle and Late Pleistocene, following recent literature (e.g. [116-118]). We consider the following
questions: Do Late Pleistocene Eurasian *H. sapiens* as a sample match the anatomical expectations,
based on the mammal comparative data, for a Neanderthal-early *H. sapiens* hybrid population
spanning multiple generations? Among individuals for whom genetic data are available, does a
higher level of Neanderthal ancestry co-occur with Neanderthal-like morphology or with
developmental abnormalities? How does hybrid status manifest itself in different aspects of cranial
shape and size, and are these skeletal indicators useful predictors of admixture in samples where no
genetic evidence is available?

Our sample comprises late Middle and Late Pleistocene (roughly corresponding to MIS 7-2)
fossil human specimens from Europe, Africa and the Middle East assigned to Neanderthals and *H.*
sapiens (Table 1; Figure 2), including, but not limited to, individuals that are genetically known and
morphologically proposed hybrids. We chose an upper age limit of MIS 7 because the suites of
diagnostic morphological features of both Neanderthals and modern humans were largely
established by this time (see e.g. [116,119]). In order to frame our study in a manner that is consistent
with studies from model organisms (e.g. see Figure 2), the Neanderthal portion of this sample is
considered as representative of one of the "parental" (unhybridized) populations. We cannot rule
out *H. sapiens* ancestry in individual Neanderthals, though to date evidence for introgression of *H.*
sapiens genes into Neanderthals is more limited than the reverse. For the second parental
population, because of the poor representation of penecontemporaneous African early *H. sapiens* in
our data set and in the fossil record generally, we consider a pooled sample of ancient and recent
sub-Saharan Africans, expected to have no or minimal Neanderthal ancestry (see [9]), as a proxy for
early *H. sapiens* anatomy. A few individuals from this time period whose attribution to early *H.*
sapiens is uncertain (i.e. Omo 2, Eliye Springs) have also been included in some of our analyses, but
they are labelled and discussed separately. The recent sub-Saharan African portion of the pooled *H.*
sapiens sample is represented by three datasets of individuals from eastern and southern Africa
(face: N=15; hemimandible: N=14; posterior cranial profile: N=15). We will refer to this sample as
African *H. sapiens*. We recognise that the inclusion of this small sample of extant sub-Saharan
Africans is not ideal given the potential effects of recent and ancient demographic processes, as well
as the possibility of admixture in deeper time (e.g. [9,16-21]). We hope to mitigate such effects to the
extent possible by combining the few available ancient individuals with our recent African sample
and limiting the extant sample to sub-Saharan Africa, thereby reducing the likelihood of admixture
from Neanderthals. We cannot rule out the possibility that introgression from other non-
Neanderthal "ghost lineages" might be present in the African sample, or the late survival of archaic
lineages not directly ancestral to *H. sapiens* in our ancient African samples. Specimens explicitly
proposed as such possible hybrids (e.g. the Iwo Eleru calvaria) have been excluded from our
analyses. The results presented below, which largely position the ancient African samples within the
same shape space as the modern ones (and distinct from Neanderthals), support these choices. All
other fossils are considered together as Pleistocene (non-Neanderthal) western Eurasians, which we
will refer to as Eurasian *H. sapiens*, and as such are potentially admixed, and the primary focus of
this study. The Apidima 1 specimen from Greece, whose attribution to early *H. sapiens* is uncertain
[117], is included as a possible Eurasian early *H. sapiens* but labelled separately.

We expect that the morphological datasets investigated will differentiate between
Neanderthals and African *H. sapiens*, reflecting their commonly accepted status as distinct lineages.
Further, we make a series of predictions aimed at determining whether the Eurasian *H. sapiens*
sample shows patterns of variation consistent with hybridization between African *H. sapiens* and
Neanderthals. As discussed briefly above, empirical research on hybridization in primates, mice, and
a handful of other mammals predicts that an admixed sample should contain: (1) individuals with
"mixed" or intermediate morphologies somewhere between their parental (un-admixed) taxa, (2)
individuals with developmentally atypical traits not seen (or seen at extremely low frequency) in the
parental taxa (especially dental or sutural anomalies), and/or (3) individuals that are transgressive in
shape or size relative to the parental taxa [1]; these characteristics will frame our morphological
expectations. Taken together, they are expected to result in hybrid populations that are more
diverse than parental groups. The consistency of these findings across taxa and generations within
taxa [1,97] support the use of this general pattern for determining hybrid status in the fossil record.

In assessing the question of hybridisation between early *H. sapiens* and Neanderthals, some
additional dynamics also need to be taken into account. Introgression from Neanderthals into *H.*
sapiens occurred at a low level, likely mediated by differences in population size (with *H. sapiens*
considerably larger; e.g. [8,120]) as well as directionality of backcrossing and possible reduced hybrid
fitness (e.g. [121-123]). Moreover, any sample is certain to be comprised of multi-generational
recombinants, rather than first generation hybrids. As a result, not all specimens in our Eurasian *H.*
sapiens sample are expected to be admixed, and those that are will likely be represent individuals
with substantially more African than Neanderthal ancestry components. Indeed, all Upper Paleolithic
Eurasian specimens for which genetic information is available show evidence of Neanderthal
admixture at least as great as that observed in modern non-Africans, but specimens with recent
Neanderthal ancestry are rare [6,124]. This differs from the studies of model organisms which focus
primarily on early generation hybrids. Furthermore, our analyses focus on specific aspects of skull
anatomy (mandibular, facial, neurocranial) in order to maximize samples (see below), and therefore
our datasets do not replicate exactly the model organism studies that generally examine size/shape
of overall cranial morphology (in addition to key non-metric traits). As a result of these factors, we
expect to find substantial overlap between the African *H. sapiens* 'parental' population and a
Eurasian admixed sample, with some individuals plotting as expected for hybrids, i.e. intermediate,
atypical, or transgressive. Alternatively, if no admixture occurred, or if such admixture does not
manifest on the aspects of cranial morphology investigated here, the Eurasian *H. sapiens* sample
would be expected to largely conform to the patterns shown by the African *H. sapiens* 'parental'
population. However, it must be stressed again that an important complicating factor in these
assessments is the problem of equifinality, i.e. that similar morphologies can result from different
processes. Some of the predictions outlined above for admixture may also apply to other
evolutionary processes, such as, for example, the retention of primitive features, or selection for
specific phenotypes under particular environmental conditions leading to convergence. The results
presented here, therefore, must be interpreted with caution.

Figure 2: Localities for Pleistocene fossil hominin specimens used in the analyses.

🔴 Neanderthal sites	◆ African HS sites	▲ Eurasian HS sites	
1 Amud 1	10 La Quina 5	17 Aduma 3	23 Abri Pataud	32 Mladeč 1,5
2 Apidima 2	11 Régourdou	18 Hofmeyr	24 Apidima 1	33 Muierii 1
3 Biache-Saint-Vaast	12 Saccopastore 1	19 Omo 1,2	25 Brno 2	34 Oase 1,2
4 La Chapelle-aux-Saints	13 Shanidar 1,5	20 Wadi Kubbaniya	26 Chancelade	35 Oberkassel 1,2
5 Feldhofer	14 Spy 1,2	21 **Eliye Springs**	27 Cioclovina	36 Pavlov 1
6 La Ferrassie 1	15 Tabun C1	22 LH18	28 Cro Magnon 1,2,3	37 Predmost 3,4
7 Gibraltar 1	16 Zafarraya		29 Dolní Věstonice 3,13,14,15,16	**38 Qafzeh 6,9**
8 Guattari 1			30 Grimaldi	**39 Skhul 5**
9 Krapina J			31 Isturitz III	

Due to the fragmentary nature of the fossil record, individuals are generally not fully preserved, and different individuals are often represented by different parts of the skeleton. In order to include as many fossils as possible, we evaluated three anatomical regions: the face, the hemimandible, and the posterior cranial profile (midsagittal profile). Our data were all collected previously by one of us (KH), and consist of three-dimensional landmarks and semilandmarks, processed with Procrustes superimposition and semilandmark sliding (in the case of the posterior cranial profile), and analysed using Principal Components Analysis (PCA). The datasets were specifically designed to capture salient morphological features that are widely considered Neanderthal or *H. sapiens* derived traits in the respective anatomical regions and are routinely used for taxonomic identification (see e.g. [125-127]). However, they may be affected differentially by different evolutionary processes. For example, facial and mandibular traits may be influenced by selection resulting from environmental factors, such as climate or diet (e.g. [128-130]), with facial morphology also possibly affected by stabilizing selection due to its importance in species recognition [131]. In contrast, neurocranial shape is proposed to track neutral evolutionary changes

and population history more closely [128], and has been linked to Neanderthal genetic ancestry in modern Europeans [35]. Our mandibular and facial datasets, therefore, may be expected to reflect a hybridization signal less clearly than our midsagittal profile dataset. Data indicating the presence of non-metric skeletal abnormalities, and genetic information on % Neanderthal ancestry, where available, were compiled from the literature and integrated in our figures and discussion. For each dataset, we also developed a shape index by calculating an axis between the mean Neanderthal and mean African *H. sapiens* shapes and projecting all Eurasian *H. sapiens* onto it [35,117]. Results are presented in Figures 3-5.

Figure 3: Hemimandible analysis. Top left: Principal Components Analysis, landmark dataset shown on modern human mandible model. Top right: Violin plot, centroid size by group, minimum to maximum values, superimposed box plot showing median and 25-75 % quartiles. Bottom: Shape index. Red: Neanderthals; Green: African recent *H. sapiens* (dots), Pleistocene specimen Wadi Kubbaniya (square); Blue: Eurasian Late Pleistocene *H. sapiens* (dots), early *H. sapiens* (square). Individuals with genetic evidence for hybridisation are marked with a DNA symbol in the PCA plot, with % Neanderthal ancestry given in the shape index. Individuals with atypical dental or sutural variation as reviewed in [132] are marked with a TOOTH symbol in the PCA plot and with an (*) in the shape index.

In the hemimandible analysis, the Eurasian *H. sapiens* sample broadly conforms to our
expectations for a hybridised sample, occupying shape space somewhat intermediate to the
Neanderthals and the African *H. sapiens* in the PCA, with transgressive individuals (Figure 3). This is a
similar pattern to that observed in mouse hybrids relative to parental lineages. Although
intermediate, the Eurasian *H. sapiens* sample is largely outside of the range of variation seen in
either Neanderthals or African *H. sapiens*, overlapping only minimally with African *H. sapiens* in the
PCA. Moreover, all the individuals with genetic or morphological signatures of hybridization sit
outside of the range of variation in the "parental" samples (neither African *H. sapiens*, nor
Neanderthal), with many of them transgressive on PC1. The Eurasian *H. sapiens* sample is also partly
intermediate, though closer to the African *H. sapiens*, in centroid size and in the shape index.
However, neither the percentage of Neanderthal genetic ancestry, where known, nor the incidence
of developmental abnormalities, appear to follow a clear relationship with Neanderthal-like
morphology or with the shape index values. A case in point is the Oase 1 mandible. This individual is
known to have approximately 10% Neanderthal ancestry - equivalent to a Neanderthal ancestor four
to six generations previously [4,133] – and is currently the earliest generation Neanderthal-modern
human hybrid known. Oase 1 shows very large overall size (one of the two largest *H. sapiens*
mandibles in centroid size in our sample) and megadont lower third molars [83,132], consistent with its
hybrid status. Yet its mandibular shape index value is less Neanderthal-like than other specimens
with known smaller Neanderthal genetic components (Figure 3). Indeed, Muerii 1 (although there is
no genome data available on Muierii 1, it may represent the same individual as Muierii 2 with 5.2%
Neanderthal ancestry[4]), Oberkassel 2 and Dolní Vestoniče 16 fall closest to Neanderthals in the
mandibular shape index.

A similar PCA pattern is shown by the posterior cranial profile analysis (Figure 4), although
the separation between the African *H. sapiens* and Neanderthal convex hulls is smaller, a
phenomenon largely driven by the position of Omo 1. Although the Eurasian *H. sapiens* sample is
again intermediate between Neanderthals and African *H. sapiens*, it shows much more overlap with
both 'parental' ranges, and especially with the African sample, indicating that a large proportion of
these Eurasian specimens display *H. sapiens*-like shape, while some are more Neanderthal-like (and
some intermediate / transgressive). This dataset essentially investigates a single – albeit very
important – feature, the outline of the posterior part of the cranium in lateral view. A rounded
cranium is considered a derived feature for modern humans, and recent work has linked a relatively
reduced globularity of the parietal and occipital bones in modern Europeans to Neanderthal genetic
ancestry and even to the presence of specific Neanderthal alleles [35]. Our shape index of the posterior
cranial profile, encompassing the midsagittal outline of the parietal region and the upper occipital,
might reasonably be considered as a proxy for an important aspect of the 'globularisation' index
calculated by [35]. The overall observed pattern of separation between our Neanderthal and African *H.*
sapiens samples is consistent with that described by [35], with the exception of Omo 2 – and, to a
lesser extent, Omo 1. These specimens differ from all other African ones in that they plot within the
Neanderthal convex hull (Omo 2) or relatively close to it (Omo 1). Their position on the plot may
indicate high levels of variation and population structure in early *H. sapiens*, as has been argued
previously (e.g. [134,135]). Alternatively, Omo 2, the only African specimen overlapping with
Neanderthals and showing a Neanderthal-like shape index, may not represent an early *H. sapiens*
(see e.g.[116]). The remaining early African *H. sapiens* or possible *H. sapiens,* including LH18, Eliye

Springs and Aduma 3, plot with the African sample, with all but Aduma 3 overlapping with the
Eurasian *H. sapiens* range.

392   In contrast to the African samples, multiple Eurasian *H. sapiens* specimens overlap with
(Qafzeh 6, Pavlov 1, Mladeč 5, Cro Magnon 3), or plot close to (Cro Magnon 1, Mladeč 1, Abri Pataud
1, Predmost 3) the Neanderthal convex hull. Many also show Neanderthal-like shape indices. Some
of these individuals have previously been described as possessing occipital 'hemibuns', posterior
projections of the occipital bone reminiscent of those shown by Neanderthals, possibly due to
Neanderthal ancestry. Unfortunately, no genomic evidence is available for them. Elongated cranial
profiles in fossil *H. sapiens* might also result from the retention of ancestral morphology represented
here by Omo 1 and possibly Omo 2. However, the Omo specimens greatly predate both the
Levantine and the European Upper Paleolithic samples – by ca. 60-90 ky and >160 ky, respectively –
making recent admixture a more likely explanation for the observed variation in the Upper
Paleolithic, and perhaps also the Near Eastern sample, a possibility that requires further
investigation. On the other hand, Oase 2, which exhibits upper third molar megadontia [83,132] and has
also recently been found to have relatively elevated Neanderthal admixture (6.06 %; [124]), plots in the
centre of the African convex hull in the PCA and shows a modern human-like shape index. So does
Dolní Vestoniče 15, which shows 4.3 % Neanderthal ancestry [4] as well as a conical mandibular
supernumerary tooth in the region of the left canine root and rotation of the left mandibular
premolar [132]; the supernumerary tooth in particular might be interpreted as possibly resulting from
admixture (though being a more common form of supernumerary tooth it is not strong evidence).
Furthermore, several additional specimens with known, relatively low, Neanderthal genetic
components (Figure 4) have shape index values within the range of African *H. sapiens*. Finally, the
proposed early *H. sapiens* Apidima 1 specimen plots with African *H. sapiens* both in the PCA as well
as in the shape index, but is characterised by a smaller centroid size, consistent with retention of
ancestral morphology as well as with possible admixture.

Midsagittal Shape PCA

Midsagittal Centroid Size

Midsagittal Shape Index

Figure 4: Posterior cranial (midsagittal) profile analysis. Top left: Principal Components Analysis,
landmark / semilandmark dataset on modern human cranium model. Top right: Violin plot centroid
size by group, minimum to maximum values, superimposed box plot showing median and 25-75 %
quartiles. Bottom: Shape index. Colors and symbols as in Figure 3. Individuals with genetic evidence
for hybridisation are marked with a DNA symbol in the PCA plot, with % Neanderthal ancestry given
in the shape index. Individuals with atypical dental or sutural variation as reviewed in [132] are marked
with a TOOTH symbol in the PCA plot and with an (*) in the shape index.

Although the hemimandible and posterior cranial profile analyses are broadly consistent
with expectations based on animal hybrid models, the facial dataset shows a different pattern
(Figure 5). Here the African *H. sapiens* sample (with the exception of Late Pleistocene specimen
Hofmeyr) falls within the more dispersed shape space of the Eurasian *H. sapiens* group; both plot
away from the tightly clustering Neanderthal sample. The two *H. sapiens* samples also show roughly
equivalent centroid sizes. The Eurasian sample, however, is considerably more variable in shape, as
reflected in their more widely diverging PC scores, with most specimens, including all individuals
with known Neanderthal genetic components, falling outside of the African *H. sapiens* convex hull

(i.e. transgressive relative to African *H. sapiens*). Such increased variability is consistent with an
admixed sample, but could also result from sampling bias, a greater temporal variability in our
Eurasian *H. sapiens* sample, or from within-species geographic variation – though similar temporal
and geographic variation did not lead to this pattern in the other analyses. Similarly, the facial shape
index values of the African *H. sapiens* specimens fall within the Eurasian *H. sapiens* range and away
from that of Neanderthals. Again, there is no relationship between the facial shape index and the
percentage of Neanderthal ancestry in the specimens for which the latter is known (Figure 5). The
early modern humans from the Near East, Qafzeh 9 and Qafzeh 6, plot in more intermediate
positions in the PCA, and have intermediate facial shape indices, although still clearly away from the
Neanderthal range.

446
Figure 5: Face analysis. Top left: Principal Components Analysis, landmark dataset shown on modern
human cranium model. Top right: Violin plot centroid size by group, minimum to maximum values,
superimposed box plot showing median and 25-75 % quartiles. Bottom: Shape index. Colors and
symbols as in Figure 3. Individuals with genetic evidence for hybridisation are marked with a DNA
symbol in the PCA plot, with % Neanderthal ancestry given in the shape index. Individuals with

atypical dental or sutural variation as reviewed in [132] are marked with a TOOTH symbol in the PCA
plot and with an (*) in the shape index.

Discussion and Future Considerations

We did not approach this study by asking whether hybridization was common in Late
Pleistocene Europe – though current evidence suggests that it may have been. Instead we wanted to
evaluate how admixture manifests in the skeleton and whether different lines of evidence,
morphological as well as genetic, can help reveal the presence of admixture, making it possible to
identify hominin hybrids on the basis of either (or both). That being said, our evaluation of the
morphology of Late Pleistocene Eurasian *H. sapiens* against predictions based on model organisms is
based on small and imperfect samples (i.e. poor representation of African early *H. sapiens*; few
individuals with both genomic and morphological data available; representation of primarily later
rather than early generation hybrids). Recent African individuals are also imperfect models for early
H. sapiens given that they have gone through their own process of evolution relative to the
population for which we are using them as a proxy. Furthermore, the interpretation of the observed
patterns is complicated by equifinality, as phenotypic variation consistent with admixture may also
result from other processes, especially retention of ancestral features, and by sampling limitations
which may underestimate the true variability of the groups included in our analyses. We therefore
can only provide tentative and preliminary answers to the questions posed. These answers,
nevertheless, can form the basis for future work exploring hybridization in the human fossil record.

To summarize, we explored whether our Late Pleistocene Eurasian *H. sapiens* sample fits our
predictions for a Neanderthal-early *H. sapiens* population with a history of hybridisation. For our
mandibular and posterior cranial datasets, we found that it was intermediate in shape and size
between Neanderthals and African *H. sapiens* and transgressive in aspects of shape, patterns
consistent with hybridization across the sample as a whole. Facial shape, on the other hand, did not
provide a clear signal, although the large variation in the Eurasian *H. sapiens* sample and high
proportion of transgressive individuals is also consistent with hybridisation in that dataset. It is
unclear why different anatomical regions would differentially preserve the signal of hybridisation, if
indeed that is the signal being detected here. Facial and mandibular shape has been argued to be
affected differentially by selection and by adaptive or plastic responses to external, environmental
factors (e.g. [128-130]). Facial morphology is also widely recognised as important in species recognition
and social interactions among primates [131], and may therefore be under selective pressure to
conform more closely to the backcrossing population. Finally, in all of our analyses Late Pleistocene
Eurasian *H. sapiens* as a sample were closer in shape to African *H. sapiens* than to Neanderthals, as
expected under conditions of asymmetric gene flow (hypothesized for large differences in parental
population sizes, as postulated for early modern Europeans relative to late Neanderthals, e.g. [120]),
or, more importantly, for a sample comprising multiple, later generation (i.e. more backcrossed into
modern humans) hybrids.

In terms of individual specimens, no direct relationship was found between estimated levels
of Neanderthal ancestry based on genomic evidence, where known, and anatomical shape/size, nor

between this genomic evidence and expression of developmentally abnormal dental or sutural
features as reported in the literature. This was the case also in the early generation Neanderthal-
modern human hybrid, the mandible Oase 1, whose only obvious phenotypic signals of hybridization
are its very large overall size and megadontia. This result is perhaps not surprising, as estimated
admixture percentages may vary across most specimens due to noise or sequencing depth.
Furthermore, the critical factor for the expression of Neanderthal-like features is most likely the
presence of particular alleles relevant for the expression of specific phenotypes, rather than overall
percentages of Neanderthal ancestry (as has recently been argued for cranial globularity by [35]).
Assuming that cranio-mandibular morphology is at least in part under genetic control, the
comparatively moderately elevated Neanderthal genetic component shown by, e.g., Dolní Vestoniče
16, may comprise alleles influencing development of the masticatory region and neurocranium,
which resulted in shape similarities to Neanderthals reflected by this specimen's mandibular and
midsagittal profile shape indices and PC scores (Figures 3, 4) and in the known misalignment of the
maxillae along the intermaxillary suture [136] but not in facial similarities (Figure 5).

Are these skeletal morphologies useful predictors of admixture in samples where no genetic
evidence is available? At the moment the patterns observed when considering a larger
sample/population are the most informative. As regards individual specimens, the signals are often
mixed, even across anatomical regions for the same individuals, likely reflecting differential
preservation of the hybridisation signal according to anatomical region (see above). The state of
preservation and degree of completeness of a fossil, therefore, may influence whether an admixture
signal can be detected. This signal will likely further be influenced by the differential expression of
Neanderthal-like or developmentally abnormal features according to the presence of particular
Neanderthal alleles or the degree and/or recency of ancestry. Nevertheless, some observations can
be made. The individuals Cro Magnon 3, Pavlov 1, and Mladeč 5 are the only ones across all our
analyses that plot well within the Neanderthal convex hull in the PCA and show Neanderthal-like
shape index values in their posterior midsagittal cranial outline, the only dataset in which they could
be included. On this basis we may hypothesize that they have a Neanderthal genetic component
comprising alleles important for cranial shape. A similar argument could be made for the Qafzeh 6
individual, who, in contrast to Qafzeh 9 from the same site, also overlaps with Neanderthals in the
posterior cranial analysis. The Qafzeh specimens are also the only ones that show a somewhat
intermediate position in the facial analysis. These results, together with the high levels of variation in
one site, and the geographic origin in the Levant, a postulated contact area between Neanderthals
and modern humans [5], raise the possibility that the Qafzeh individuals may also have some
Neanderthal ancestry [84]. Even though such indications are intriguing, they cannot be considered
conclusive and must be treated as hypotheses, especially since, as mentioned above, similar
phenotypes might be consistent with different underlying causes. Nevertheless, it is possible to
evaluate the likelihood of such alternative explanations on a case-by-case basis. For example,
because a rounded cranium is a derived *H. sapiens* feature, an alternative hypothesis for a relatively
elongated cranial phenotype could be that it results from retention of the ancestral, elongated
condition. An ancestral retention, however, is more convincing for Qafzeh, which represents an early
H. sapiens population dating to ca. 100-130 ka (Table 1), than for the Late Pleistocene European
specimens, which greatly postdate the establishment of the derived condition [117,137].

Finally, recent suggestions that skeletal anomalies in some Upper Paleolithic and
Neanderthal samples result from inbreeding [132,138] may further complicate the interpretation of

developmental abnormalities as indicators of admixture. Indeed both processes are expected to have taken place in the highly dynamic conditions of cyclical environmental change of Pleistocene Eurasia, which likely resulted in repeated isolation of populations in refugia areas, sometimes leading to local extinctions, but also to population expansion and dispersals [139]. Under these conditions, paleodemes have been proposed to resemble 'tidal islands', often isolated but occasionally flooded with expanding / dispersing populations and their genetic material [139]. However, although empirical evidence from primates for the skeletal expression of inbreeding is limited, the evidence that does exist suggests that it is associated with abnormalities (e.g. reduced size, anencephaly, polydactly, syndactly, limb malformations [140-144]) that are different from those shown to occur in hybrids (e.g. increased size, extremely rare dental and sutural traits with no other associated diseases or syndromes [1,84,96,97]). This indicates that it should be possible to distinguish between these causal phenomena, and their relative contributions to the morphology we see in the fossil record, going forward.

This study compared genomic and morphological data sets, in order to interrogate the fossil evidence for Late Pleistocene hybridisation between Neanderthals and early *H. sapiens*, for which we currently have substantial evidence. We urge further studies of the phenotype to expand our ability to detect the ways in which migration, interaction and genetic exchange have shaped the human past, beyond what is currently visible with the lens of ancient DNA. It is particularly important to examine such data sets together to understand the effects of hybridization on the morphology of later generation hybrids, and whether these effects vary by anatomical region. The results provided here should form the basis for developing hypotheses to be tested against the human fossil record in the future.

Acknowledgements

This research was supported by the European Research Council (ERC CoG no. 724703 CROSSROADS), the German Research Foundation (DFG FOR 2237 'Words, Bones, Genes, Tools), and the National Research Foundation of South Africa (Grant# 117670). We thank all curators and institutions that allowed us access to the fossil specimens used in our analyses, K. Warren for collecting and analysing the data used in Figure 1, A. M. Bosman and C. Röding for help with processing the datasets used in Figures 3-4, H. Rathmann and J. Beier for help with the figures, and C. Posth for important feedback. We are grateful to three anonymous reviewers whose comments and suggestions greatly improved our manuscript.

TEXT BOX:

Are Neanderthals Distinct Species and Does It Matter?

There is a longstanding debate around the taxonomy of Neanderthals and especially whether they
represent distinct species from *Homo sapiens*. Even the two authors of this manuscript differ in
opinion on this matter. Applying species concepts that rely on complete reproductive isolation (e.g.
[145]), where species are defined in part according to their ability to interbreed and produce fertile
offspring, it is now clear that they would be considered the same species. However, such species
concepts are of little value in the context of hybridising lineages. Instead, most evolutionary
biologists would agree that species are defined by retained morphological, behavioural and genetic
differences even in the face of gene flow [146], a definition that more clearly argues for species-
distinctiveness for Neanderthals (and possibly others; e.g. [67,147]). However, as pointed out elsewhere
[148], this definition also implies an ability to coexist geographically without the fusion of lineages [149],
which does not appear to be the case between Neanderthals and *H. sapiens*. Although these
lineages appear to have evolved their differences initially in geographic isolation, post-contact gene
flow resulted in the subsequent disappearance, at least morphologically, of all but one lineage, *H.
sapiens*. This may suggest that Neanderthals and *H. sapiens* may not have been able to coexist
without lineage fusion [148]. However, the disappearance of Neanderthals and their relatively low
genetic contribution to later humans may also point to competitive exclusion, in the context of
reported reduced hybrid fitness (e.g. [122]), environmental instability and cultural / demographic
differences (e.g. [150,151]). Given these differences of opinion, whether these taxa represent
genomically coherent, monophyletic species remains an open question, one which may be resolved
in the future as more and more genomes are sequenced across the full distribution of the taxa. More
importantly, the question of specific status is irrelevant for the investigation undertaken here.
Regardless of whether Neanderthals, *H. sapiens* and others represent distinct species, they are
clearly genetically differentiated lineages. How different lineages need to be for gene exchange to be
categorised as hybridization is subjective. Typically, however, it is at or above the rank of subspecies
[152], which is almost certainly the case here. This is also the case for the extant taxa used as models
for considering the morphological effects of admixture. Baboons are particularly interesting in this
respect due to their extensively documented phenotypic, ecological and behavioral distinctiveness
combined with their ability to interbreed extensively (see [40]). The history of disagreement over
specific versus subspecific levels of distinctiveness among baboon 'allotaxa' broadly parallels the
discussion surrounding Neanderthals and *H. sapiens*, pointing to the shortcomings of species
concepts as applied not only to the fossil record, but also to extant mammalian taxa.

	Face	Mandible	Neurocranium	Apx geological age (ka)	Reference
Neanderthal					
Amud 1		x	x	55-60	[153]
Apidima 2	x			ca. 170	[117]
Biache St Vaast			x	ca. 250	[154]
La Chapelle-aux-Saints	x		x	47 / 56	[155]
Feldhofer			x	40	[156]
Ferrassie 1	x	x	x	43-45	[157]
Gibraltar 1	x			ca. 50	[158]
Guattari 1	x		x	50-60	[159,160]
Krapina J		x		140-120	[161]
La Quina 5		x	x	>48 (OIS 3-4)	[162]
Regourdou		x		MIS 4-5	[163]
Saccopastore 1			x	295-220	[164]
Shanidar 1	x	x		46-50 Uncalibrated	[165]
Shanidar 5	x			46-50 Uncalibrated	[165]
Spy 1			x	40.6-44.2 cal BP	[166]
Spy 2			x	40.6-44.2 cal BP	[166]
Tabun C1		x		130-100	[167,168]
Zafarraya		x		ca 30-46, >46	[169,170]
***Homo sapiens* Late Middle – Late Pleistocene Africa**					
Aduma 3			x	79-105	[171]
Hofmeyr	x			36	[172]
Eliye Springs			x	Middle/Late Pleistocene	[173,174]
LH 18			x	120+/-30	[175,176]
Omo 1			x	233+/-22	[177,178]
Omo 2			x	195+/-5	[178]
Wadi Kubbaniya	x	x		ca. 20	[179]
***Homo sapiens* Late Middle – Late Pleistocene Eurasia**					
Abri Pataud 1	x	x	x	28-26 (22 uncalibrated)	[180]
Apidima 1			x	ca. 210	[117]
Brno 2			x	23.7 uncal (ca. 28.5 cal BP)	[181]
Chancelade	x		x	18	[182]
Cioclovina			x	ca. 33	[183]
Cro Magnon 1	x		x	ca 30	[184]
Cro Magnon 2	x		x	ca 30	[184]
Cro Magnon 3			x	ca 30	[184]
Dolní Vestoniče 13	x	x	x	ca. 31	[185]
Dolní Vestoniče 14	x	x		ca. 31	[185]
Dolní Vestoniče 15	x	x	x	ca. 31	[185]
Dolní Vestoniče 16	x	x	x	ca. 30	[185]
Dolní Vestoniče 3	x	x	x	undated	[185]
Grimaldi	x	x		25 uncal (ca. 29.5 cal BP)	[186]

Isturitz III		x		Upper Paleolithic	[187]
Mladec 1	x		x	35-36.5	[188]
Mladec 5			x	35-36.5	[188]
Muierii 1	x	x	x	ca. 35	[81]
Oase 1		x		ca. 40.5	[189]
Oase 2	x		x	ca. 40.5	[82]
Oberkassel 1		x		12 uncalb (ca. 14.2 cal BP)	[190]
Oberkassel 2		x		12 uncalb (ca. 14.2 cal BP)	[190]
Pavlov 1			x	ca. 30	[185]
Predmost 3	x		x	27-29	[191]
Predmost 4	x		x	27-29	[191]
Qafzeh6	x		x	100-130	[168]
Qafzeh9	x	x	x	100-130	[168]
Skhul 5		x	x	100-130	[168]

607 **Table 1**: Hominin samples used in analyses. Dates are reported in calendar years unless otherwise
608 stated

1   Ackermann, R. *et al.* Hybridization in human evolution: insights from other organisms. .
     Evolutionary Anthropology: Issues, News, and Reviews **28**, 189-209,
     doi:https://doi.org/10.1002/evan.21787 (2019).
2   Taylor, S. A. & Larson, E. L. Insights from genomes into the evolutionary importance and
     prevalence of hybridization in nature. *Nature ecology & evolution* **3**, 170-177,
     doi:10.1038/s41559-018-0777-y (2019).
3   Fu, Q. *et al.* Genome sequence of a 45,000-year-old modern human from western Siberia.
     Nature **514**, 445-449, doi:10.1038/nature13810

     http://www.nature.com/nature/journal/v514/n7523/abs/nature13810.html#supplementary-
     information (2014).
4   Fu, Q. *et al.* The genetic history of Ice Age Europe. *Nature* **534**, 200-205,
     doi:10.1038/nature17993 (2016).
5   Green, R. E. *et al.* A Draft Sequence of the Neandertal Genome. *Science* **328**, 710-722 (2010).
6   Hajdinjak, M. *et al.* Initial Upper Palaeolithic humans in Europe had recent Neanderthal
     ancestry. *Nature* **592**, 253-257, doi:10.1038/s41586-021-03335-3 (2021).
7   Prüfer, K. *et al.* A genome sequence from a modern human skull over 45,000 years old from
     Zlatý kůň in Czechia. *Nature ecology & evolution* **5**, 820-825, doi:10.1038/s41559-021-01443-
     x (2021).
8   Villanea, F. A. & Schraiber, J. G. Multiple episodes of interbreeding between Neanderthal
     and modern humans. *Nature ecology & evolution* **3**, 39-44, doi:10.1038/s41559-018-0735-8
     (2019).
9   Chen, L., Wolf, A. B., Fu, W., Li, L. & Akey, J. M. Identifying and Interpreting Apparent
     Neanderthal Ancestry in African Individuals. *Cell* **180**, 677-687.e616,
     doi:10.1016/j.cell.2020.01.012 (2020).
10   Posth, C. *et al.* Deeply divergent archaic mitochondrial genome provides lower time
     boundary for African gene flow into Neanderthals. *Nature Communications* **8**, 16046,
     doi:10.1038/ncomms16046

     https://www.nature.com/articles/ncomms16046#supplementary-information (2017).
11   Meyer, M. *et al.* A high-coverage genome sequence from an archaic Denisovan individual.
     Science **338**, 222-226 (2012).
12   Reich, D. *et al.* Genetic history of an archaic hominin group from Denisova Cave in Siberia.
     Nature **468**, 1053-1060, doi:10.1038/nature09710 (2010).
13   Prufer, K. *et al.* The complete genome sequence of a Neanderthal from the Altai Mountains.
     Nature **505**, 43-49, doi:10.1038/nature12886

     http://www.nature.com/nature/journal/v505/n7481/abs/nature12886.html#supplementary-
     information (2014).
14   Slon, V. *et al.* Neandertal and Denisovan DNA from Pleistocene sediments. *Science*,
     doi:10.1126/science.aam9695 (2017).
15   Slon, V. *et al.* The genome of the offspring of a Neanderthal mother and a Denisovan father.
     Nature **561**, 113-116, doi:10.1038/s41586-018-0455-x (2018).
16   Durvasula, A. & Sankararaman, S. Recovering signals of ghost archaic introgression in African
     populations. *Science Advances* **6**, eaax5097, doi:10.1126/sciadv.aax5097 (2020).
17   Hammer, M. F., Woerner, A. E., Mendez, F. L., Watkins, J. C. & Wall, J. D. Genetic evidence
     for archaic admixture in Africa. *Proceedings of the National Academy of Sciences* **108**, 15123-
     15128, doi:10.1073/pnas.1109300108 (2011).

18 Harvati, K. *et al.* The Later Stone Age Calvaria from Iwo Eleru, Nigeria: Morphology and
Chronology. *PLOS ONE* **6**, e24024, doi:10.1371/journal.pone.0024024 (2011).
19 Lachance, J. *et al.* Evolutionary History and Adaptation from High-Coverage Whole-Genome
Sequences of Diverse African Hunter-Gatherers. *Cell* **150**, 457-469,
doi:10.1016/j.cell.2012.07.009 (2012).
20 Lipson, M. *et al.* Ancient West African foragers in the context of African population history.
Nature **577**, 665-670, doi:10.1038/s41586-020-1929-1 (2020).
21 Wang, K., Mathieson, I., O'Connell, J. & Schiffels, S. Tracking human population structure
through time from whole genome sequences. *PLoS Genet* **16**, e1008552,
doi:10.1371/journal.pgen.1008552 (2020).
22 Smith, F. H., Ahern, J. C. M., Janković, I. & Karavanić, I. The Assimilation Model of modern
human origins in light of current genetic and genomic knowledge. *Quaternary International*
450, 126-136, doi:10.1016/j.quaint.2016.06.008 (2017).
23 Dannemann, M., Andrés, A. M. & Kelso, J. *Adaptive variation in human toll-like receptors is*
contributed by introgression from both Neandertals and Denisovans. (2015).
24 Hu, Y., Ding, Q., He, Y., Xu, S. & Jin, L. Reintroduction of a Homocysteine Level-Associated
Allele into East Asians by Neanderthal Introgression. *Molecular biology and evolution* **32**,
3108-3113, doi:10.1093/molbev/msv176 (2015).
25 Huerta-Sánchez, E. *et al.* Altitude adaptation in Tibetans caused by introgression of
Denisovan-like DNA. *Nature* **512**, 194-197 (2014).
26 Lin, Y. L., Pavlidis, P., Karakoc, E., Ajay, J. & Gokcumen, O. The evolution and functional
impact of human deletion variants shared with archaic hominin genomes. *Molecular biology*
and evolution **32**, 1008-1019, doi:10.1093/molbev/msu405 (2015).
27 Sankararaman, S. *et al.* The genomic landscape of Neanderthal ancestry in present-day
humans. *Nature* **507**, 354-357, doi:10.1038/nature12961

http://www.nature.com/nature/journal/v507/n7492/abs/nature12961.html#supplementary-
information (2014).
28 Schlebusch, C. M. *et al.* Genomic variation in seven Khoe-San groups reveals adaptation and
complex African history. *Science* **338**, 374-379, doi:10.1126/science.1227721 (2012).
29 Simonti, C. N. *et al.* The phenotypic legacy of admixture between modern humans and
Neandertals. *Science* **351**, 737-741, doi:10.1126/science.aad2149 (2016).
30 Sudmant, P. H. *et al.* Global diversity, population stratification, and selection of human copy-
number variation. *Science* **349**, aab3761, doi:10.1126/science.aab3761 (2015).
31 Vernot, B. & Akey, J. M. Resurrecting Surviving Neandertal Lineages from Modern Human
Genomes. *Science* **343**, 1017-1021, doi:10.1126/science.1245938 (2014).
32 Skov, L. *et al.* The nature of Neanderthal introgression revealed by 27,566 Icelandic
genomes. *Nature*, doi:10.1038/s41586-020-2225-9 (2020).
33 Mozzi, A. *et al.* Distinct selective forces and Neanderthal introgression shaped genetic
diversity at genes involved in neurodevelopmental disorders. *Scientific Reports* **7**, 6116,
doi:10.1038/s41598-017-06440-4 (2017).
34 Gregory, M. D. *et al.* Neanderthal-Derived Genetic Variation Shapes Modern Human
Cranium and Brain. *Scientific Reports* **7**, 6308, doi:10.1038/s41598-017-06587-0 (2017).
35 Gunz, P. *et al.* Neandertal Introgression Sheds Light on Modern Human Endocranial
Globularity. *Current Biology* **29**, 120-127.e125,
doi:https://doi.org/10.1016/j.cub.2018.10.065 (2019).
36 Zeberg, H. & Pääbo, S. The major genetic risk factor for severe COVID-19 is inherited from
Neanderthals. *Nature* **587**, 610-612, doi:10.1038/s41586-020-2818-3 (2020).
37 Zeberg, H. & Pääbo, S. A genomic region associated with protection against severe COVID-19
is inherited from Neandertals. *Proceedings of the National Academy of Sciences* **118**,
e2026309118, doi:10.1073/pnas.2026309118 (2021).

38 Gokcumen, O. Archaic hominin introgression into modern human genomes. *American*
journal of physical anthropology **n/a**, doi:10.1002/ajpa.23951.
39 Arnold, M. L. *Evolution Through Genetic Exchange.* (Oxford University Press, 2006).
40 Jolly, C. J. A proper study for mankind: Analogies from the Papionin monkeys and their
implications for human evolution. *American journal of physical anthropology* **116**, 177-204,
doi:doi:10.1002/ajpa.10021 (2001).
41 Mallet, J. Hybridization as an invasion of the genome. *Trends Ecol. Evol.* **20**, 229-237 (2005).
42 Stelkens, R. & Seehausen, O. Genetic distance between species predicts novel trait
expression in their hybrids. *Evolution; international journal of organic evolution* **63**, 884-897,
doi:doi:10.1111/j.1558-5646.2008.00599.x (2009).
43 Ackermann, R., Brink, J., Vrahimis, S. & de Klerk, B. Hybrid wildebeest (Artiodactyla: Bovidae)
provide further evidence for shared signatures of admixture in mammalian crania. *South*
African Journal of Science **106**, 1-4 (2010).
44 Baranov, A. S., Zakharov, V.M. Developmental stability in hybrids of European bison, *Bison*
bonasus, and domestic cattle. *Acta Theriologica* **42**, 87-90 (1997).
45 Brink, J. S. *The evolution of the black wildebeest, Connochaetes gnou, and modern large*
mammal faunas in central southern Africa. PhD thesis, University of Stellenbosch., (2005).
46 Cahill, J. A. *et al.* Genomic evidence of geographically widespread effect of gene flow from
polar bears into brown bears. *Molecular ecology* **24**, 1205-1217, doi:10.1111/mec.13038
(2015).
47 Kumar, V. *et al.* The evolutionary history of bears is characterized by gene flow across
species. *Scientific Reports* **7**, 46487, doi:10.1038/srep46487 (2017).
48 Figueiró, H. V. *et al.* Genome-wide signatures of complex introgression and adaptive
evolution in the big cats. *Science Advances* **3**, doi:10.1126/sciadv.1700299 (2017).
49 Li, G., Davis, B. W., Eizirik, E. & Murphy, W. J. Phylogenomic evidence for ancient
hybridization in the genomes of living cats (Felidae). *Genome research* **26**, 1-11,
doi:10.1101/gr.186668.114 (2016).
50 Benson, J. F., Patterson, B. R. & Wheeldon, T. J. Spatial genetic and morphologic structure of
wolves and coyotes in relation to environmental heterogeneity in a Canis hybrid zone.
Molecular ecology **21**, 5934-5954, doi:10.1111/mec.12045 (2012).
51 Khosravi, R., Rezaei, H. R. & Kaboli, M. Detecting Hybridization between Iranian Wild Wolf
(Canis Lupus Pallipes) and Free-Ranging Domestic Dog (Canis Familiaris) by Analysis of
Microsatellite Markers. *Zoological Science* **30**, 27-34, doi:10.2108/zsj.30.27 (2013).
52 Mahan, B. R., Gipson, P. S. & Case, R. M. Characteristics and Distribution of Coyote X Dog
Hybrids Collected in Nebraska. *The American Midland Naturalist* **100**, 408-415,
doi:10.2307/2424840 (1978).
53 Monzon, J., Kays, R. & Dykhuizen, D. E. Assessment of coyote-wolf-dog admixture using
ancestry-informative diagnostic SNPs. *Molecular ecology* **23**, 182-197,
doi:10.1111/mec.12570 (2014).
54 Vila, C. *et al.* Combined use of maternal, paternal and bi-parental genetic markers for the
identification of wolf-dog hybrids. *Heredity* **90**, 17-24, doi:10.1038/sj.hdy.6800175 (2003).
55 Pastorini, J., Zaramody, A., Curtis, D. J., Nievergelt, C. M. & Mundy, N. I. Genetic analysis of
hybridization and introgression between wild mongoose and brown lemurs. *BMC*
evolutionary biology **9**, 32, doi:10.1186/1471-2148-9-32 (2009).
56 Wyner, Y. M., Johnson, S. E., Stumpf, R. M. & Desalle, R. Genetic assessment of a white-
collared×red-fronted lemur hybrid zone at Andringitra, Madagascar. *American Journal of*
Primatology **57**, 51-66, doi:doi:10.1002/ajp.10033 (2002).
57 Aguiar, L. M. *et al.* Sympatry between Alouatta caraya and Alouatta clamitans and the
rediscovery of free-ranging potential hybrids in Southern Brazil. *Primates* **48**, 245-248,
doi:10.1007/s10329-007-0039-0 (2007).

58   Cortés-Ortiz, L., Agostini, I., Aguiar, L. M., Kelaita, M., Silva, F. E., & Bicca-Marques, J. C. . in
Howler monkeys: Behaviour, Ecology and Conservation   (eds M.M. Kowalewski *et al.*) 107-
131 (Springer-Verlag, 2015).
59   Malukiewicz, J. *et al.* Hybridization effects and genetic diversity of the common and black-
tufted marmoset (Callithrix jacchus and Callithrix penicillata) mitochondrial control region.
American journal of physical anthropology **155**, 522-536, doi:10.1002/ajpa.22605 (2014).
60   Peres, C. A., Patton, J. L. & Nazareth F. da Silva, M. Riverine Barriers and Gene Flow in
Amazonian Saddle-Back Tamarins. *Folia Primatologica* **67**, 113-124, doi:10.1159/000157213
(1996).
61   Rossan, R. N. & Baerg, D. C. Laboratory and feral hybridization ofAteles geoffroyi panamensis
Kellogg and Goldman 1944 andA. fusciceps robustus Allen 1914 in Panama. *Primates* **18**,
235-237, doi:10.1007/BF02382962 (1977).
62   Detwiler, K. M., Burrell, A. S. & Jolly, C. J. Conservation Implications of Hybridization in
African Cercopithecine Monkeys. *International Journal of Primatology* **26**, 661-684,
doi:10.1007/s10764-005-4372-0 (2005).
63   Fooden, J. Rhesus and crab-eating macaques: Intergradation in Thailand. *Science* **143**, 363-
364 (1964).
64   Schillaci, M. A., Froehlich, J. W., Supriatna, J. & Jones-Engel, L. The effects of hybridization on
growth allometry and craniofacial form in Sulawesi macaques. *Journal of human evolution*
49, 335-369, doi:https://doi.org/10.1016/j.jhevol.2005.04.010 (2005).
65   Wildman, D. E. *et al.* Mitochondrial evidence for the origin of hamadryas baboons. *Molecular*
Phylogenetics and Evolution **32**, 287-296, doi:https://doi.org/10.1016/j.ympev.2003.12.014
(2004).
66   Zinner, D., Groeneveld, L. F., Keller, C. & Roos, C. Mitochondrial phylogeography of baboons
(Papio spp.): indication for introgressive hybridization? *BMC evolutionary biology* **9**, 83,
doi:10.1186/1471-2148-9-83 (2009).
67   Harvati, K., Frost, S. R. & McNulty, K. P. Neanderthal taxonomy reconsidered: Implications of
3D primate models of intra- and interspecific differences. *Proceedings of the National*
Academy of Sciences of the United States of America **101**, 1147-1152,
doi:10.1073/pnas.0308085100 (2004).
68   Jolly, C. J. in *Species, Species Concepts and Primate Evolution*   (eds William H. Kimbel &
Lawrence B. Martin) 67-107 (Springer US, 1993).
69   Brockelman, W. Y. Maintenance and evolution of social structure in gibbons. *The Lesser Apes*
: Evolutionary and Behavioural Biology (1984).
70   Marshall, J. & Sugardjito, J. in *Comparative Primate Biology, 1. Systematics, Evolution and*
Anatomy   (eds D.R. Swindler & J. Erwin) 137-185 (Liss, 1986).
71   Prado-Martinez, J. *et al.* Great ape genetic diversity and population history. *Nature* **499**, 471,
doi:10.1038/nature12228

https://www.nature.com/articles/nature12228#supplementary-information (2013).
72   de Manuel, M. *et al.* Chimpanzee genomic diversity reveals ancient admixture with bonobos.
Science (New York, N.Y.) **354**, 477-481, doi:10.1126/science.aag2602 (2016).
73   Arnold, M. L. & Kunte, K. Adaptive Genetic Exchange: A Tangled History of Admixture and
Evolutionary Innovation. *Trends in Ecology & Evolution* **32**, 601-611,
doi:https://doi.org/10.1016/j.tree.2017.05.007 (2017).
74   Nye, J. *et al.* Selection in the Introgressed Regions of the Chimpanzee Genome. *Genome*
Biology and Evolution **10**, 1132-1138, doi:10.1093/gbe/evy077 (2018).
75   Sankararaman, S., Mallick, S., Patterson, N. & Reich, D. The Combined Landscape of
Denisovan and Neanderthal Ancestry in Present-Day Humans. *Current Biology* **26**, 1241-
1247, doi:10.1016/j.cub.2016.03.037 (2016).
76   Vernot, B. *et al.* Excavating Neandertal and Denisovan DNA from the genomes of Melanesian
individuals. *Science* **352**, 235-239, doi:10.1126/science.aad9416 (2016).

77   Kuhlwilm, M. The evolution of FOXP2 in the light of admixture. *Current Opinion in Behavioral*
Sciences **21**, 120-126, doi:https://doi.org/10.1016/j.cobeha.2018.04.006 (2018).
78   Kuhlwilm, M., Han, S., Sousa, V. C., Excoffier, L. & Marques-Bonet, T. Ancient admixture from
an extinct ape lineage into bonobos. *Nature ecology & evolution* **3**, 957-965,
doi:10.1038/s41559-019-0881-7 (2019).
79   Duarte, C. *et al.* The early Upper Paleolithic human skeleton from the Abrigo do Lagar Velho
(Portugal) and modern human emergence in Iberia. *Proc Nat Acad Sci USA* **96**, 7604-7609
(1999).
80   Wolpoff, M., Hawks, J., Frayer, D. & Hunley, K. Modern human ancestry at the peripheries: a
test of the replacement theory. *Science* **291**, 293-297 (2001).
81   Soficaru, A., Petrea, C., Dobos, A. & Trinkaus, E. Early modern humans from the Pestera
Muierii, Baia de Fier, Romania. *Proc Natl Acad Sci U S A* **103**, 17196-17201 (2006).
82   Rougier, H. *et al.* Peştera cu Oase 2 and the cranial morphology of early modern Europeans.
Proceedings of the National Academy of Sciences **104**, 1165-1170,
doi:10.1073/pnas.0610538104 (2007).
83   Trinkaus, E., Constantin, S. & Zilhão, J. *Life and Death at the Pestera cu Oase. A Setting for*
Modern Human Emergence in Europe. (Oxford University Press, 2013).
84   Ackermann, R. R. Phenotypic traits of primate hybrids: recognizing admixture in the fossil
record. *Evolutionary Anthropology* **19**, 258-270 (2010).
85   Smith, F., Lacy, K. & Caldwell, S. Morphological Evidence for Modern Human Influences in
Late Central European Neandertals. *Anthropologie (Brno)* **53**, 61-76 (2015).
86   Smith, F. H., Hutchinson, V. T. & Janković, I. in *African Genesis: Perspectives on Hominin*
Evolution (eds S.C. Reynolds & A. Gallagher) 365-393 (Cambridge University Press, 2012).
87   Smith, F. H., Falsetti, A.B., & Simmons, T. . in *Man and Environment in the Paleolithic* (ed H.
Ullrich) 167-179 (ERAUL, 1995).
88   Ahern, J. C., Janković, I., Voisin, J. & Smith, F. H. in *Origins of Modern Humans: Biology*
Reconsidered (eds Smith. F.H. & J.C. Ahern) 151-222 (Wiley-Blackwell, 2013).
89   Cartmill, M. & Smith, F. H. *The human lineage*. (John Wiley & Sons, 2009).
90   Condemi, S. *et al.* Possible interbreeding in late Italian Neanderthals? New data from the
Mezzena Jaw (Monti Lessini, Verona, Italy). *PLoS One* **8**, 1-9 (2013).
91   Harvati, K., Gunz, P. & Grigorescu, D. Cioclovina (Romania): affinities of an early modern
European. *Journal of human evolution* **53**, 732-746, doi:10.1016/j.jhevol.2007.09.009 (2007).
92   Stringer, C. What makes a modern human. *Nature* **485**, 33, doi:10.1038/485033a (2012).
93   Tattersall, I. & Schwartz, J. H. Hominids and hybrids: The place of Neanderthals in human
evolution. *Proceedings of the National Academy of Sciences* **96**, 7117-7119,
doi:10.1073/pnas.96.13.7117 (1999).
94   Klein, R. *The Human Career. Third Edition*. (University of Chicago Press., 2009).
95   Klein, R. G. Paleoanthropology. Whither the Neanderthals? *Science* **299**, 1525-1527,
doi:10.1126/science.1082025 (2003).
96   Ackermann, R. R., Rogers, J. & Cheverud, J. Identifying the morphological signatures of
hybridization in primate and human evolution. *Journal of human evolution* **51**, 632-645
(2006).
97   Ackermann, R. R., Schroeder, L., Rogers, J. & Cheverud, J. Further evidence for phenotypic
signatures of hybridization in descendant baboon populations. *Journal of human evolution*
74 (2014).
98   Warren, K. A. *et al.* Craniomandibular form and body size variation of first generation mouse
hybrids: A model for hominin hybridization. *Journal of human evolution* **116**, 57-74,
doi:10.1016/j.jhevol.2017.12.002 (2018).
99   Harvati, K. & Roksandic, M. in *Paleoanthropology of the Balkans and Anatolia: Human*
Evolution and its Context (eds Katerina Harvati & Mirjana Roksandic) 51-68 (Springer
Netherlands, 2016).

100　Ackermann, R. R. in *Tinkering: the Micro-evolution of Development* Vol. 284 (ed Novartis
Foundation Symposium) 262-279 (Wiley, 2007).

101　Goodwin, T. *Supernumerary Teeth in Pleistocene, Recent, and Hybrid Individuals of the*
Spermophilus richardsonii Complex (Sciuridae). Vol. 79 (1998).

102　Zdjelar, N., Nagendran, L., Kendall, C., Ackermann, R. R. & Schroeder, L. The hybrid skull of
the eastern coyote (Canis latrans var.): Nonmetric traits and craniomandibular shape.
Journal of morphology **282**, 1745-1764, doi:10.1002/jmor.21417 (2021).

103　Eichel, K. & Ackermann, R. R. Variation in the nasal cavity of baboon hybrids with
implications for late Pleistocene hominins. *Journal of human evolution* **94**, 134-145 (2016).

104　Rieseberg, L., Archer, M. & Wayne, R. Transgressive segregation, adaptation and speciation.
Heredity **83**, 363-372 (1999).

105　Leamy, L. Morphometric studies in inbred and hybrid house miceI. Patterns in the mean
values. *Journal of Heredity* **73**, 171-176, doi:10.1093/oxfordjournals.jhered.a109612 (1982).

106　Leamy, L. *Morphometric studies in inbred and hybrid house mice. VII. Heterosis in fluctuating*
asymmetry at different ages. Vol. 191 (1992).

107　Leamy, L. & Thorpe, R. *Morphometric studies in inbred and hybrid house mice. Heterosis,*
homeostasis and heritability of size and shape. Vol. 22 (1984).

108　Percival, C. J. *et al.* Genetics of murine craniofacial morphology: diallel analysis of the eight
founders of the Collaborative Cross. *Journal of anatomy* **228**, 96-112, doi:10.1111/joa.12382
(2016).

109　Thorpe, R. & L.s, L. *Morphometric studies in inbred and hybrid House mice (Mus sp.):*
Multivariate analysis of size and shape. Vol. 199 (1983).

110　Warren, K. A. *Using the craniomandibular morphology of hybrid mice to better understand*
hybrid morphologies in the hominin fossil record PhD thesis, University of Cape Town, (2017).

111　Cheverud, J. M., Jacobs, S. C. & Moore, A. J. Genetic differences among subspecies of the
saddle-back tamarin (Saguinus fuscicollis):evidence from hybrids. *American Journal of*
Primatology **31**, 23-39, doi:doi:10.1002/ajp.1350310104 (1993).

112　Carmon, J. L. Heterosis, combining ability, and maternal effects in mice. *Journal of Genetics*
58, 225-231, doi:10.1007/BF02986141 (1963).

113　Kohn, L. A. P., Langton, L. B. & Cheverud, J. M. Subspecific genetic differences in the saddle-
back tamarin (Saguinus fuscicollis) postcranial skeleton. *American Journal of Primatology* **54**,
41-56, doi:doi:10.1002/ajp.1011 (2001).

114　Kurnianto, E., Shinjo, A., Suga, D. & Uema, N. Diallel Cross Analysis of Body Weight in
Subspecies of Mice. *Experimental Animals* **48**, 277-283, doi:10.1538/expanim.48.277 (1999).

115　Buck, L. T. *et al.* Effects of admixture on pelvic morphology: a macaque model. *Journal of*
human evolution **159**, 1030-1049 (2021).

116　Bergström, A., Stringer, C., Hajdinjak, M., Scerri, E. M. L. & Skoglund, P. Origins of modern
human ancestry. *Nature* **590**, 229-237, doi:10.1038/s41586-021-03244-5 (2021).

117　Harvati, K. *et al.* Apidima Cave fossils provide earliest evidence of Homo sapiens in Eurasia.
Nature **571**, 500-504, doi:10.1038/s41586-019-1376-z (2019).

118　Tryon, C. A. *et al.* Late Pleistocene age and archaeological context for the hominin calvaria
from GvJm-22 (Lukenya Hill, Kenya). *Proceedings of the National Academy of Sciences* **112**,
2682-2687, doi:10.1073/pnas.1417909112 (2015).

119　Hublin, J. J. The origin of Neandertals. *Proceedings of the National Academy of Sciences* **106**,
16022-16027, doi:10.1073/pnas.0904119106 (2009).

120　Mellars, P. & French, J. C. Tenfold Population Increase in Western Europe at the Neandertal–
to–Modern Human Transition. *Science* **333**, 623-627, doi:10.1126/science.1206930 (2011).

121　Currat, M. & Excoffier, L. Strong reproductive isolation between humans and Neanderthals
inferred from observed patterns of introgression. *Proc Natl Acad Sci U S A* **108**, 15129-
15134, doi:10.1073/pnas.1107450108 (2011).

122 Juric, I., Aeschbacher, S. & Coop, G. The Strength of Selection against Neanderthal
Introgression. *PLOS Genetics* **12**, e1006340, doi:10.1371/journal.pgen.1006340 (2016).
123 McCoy, R. C., Wakefield, J. & Akey, J. M. Impacts of Neanderthal-Introgressed Sequences on
the Landscape of Human Gene Expression. *Cell* **168**, 916-927.e912,
doi:10.1016/j.cell.2017.01.038 (2017).
124 Siska, V. *Human population history and its interplay with natural selection* PhD thesis,
Cambridge, (2018).
125 Gunz, P. & Harvati, K. The Neanderthal "chignon": variation, integration, and homology.
Journal of human evolution **52**, 262-274, doi:10.1016/j.jhevol.2006.08.010 (2007).
126 Harvati, K., Hublin, J.-J. & Gunz, P. Evolution of middle-late Pleistocene human cranio-facial
form: A 3-D approach. *Journal of human evolution* **59**, 445-464,
doi:https://doi.org/10.1016/j.jhevol.2010.06.005 (2010).
127 Nicholson, E. & Harvati, K. Quantitative analysis of human mandibular shape using three-
dimensional geometric morphometrics. *American journal of physical anthropology* **131**, 368-
383, doi:10.1002/ajpa.20425 (2006).
128 Harvati, K. & Weaver, T. D. Human cranial anatomy and the differential preservation of
population history and climate signatures. *The anatomical record. Part A, Discoveries in*
molecular, cellular, and evolutionary biology **288**, 1225-1233, doi:10.1002/ar.a.20395 (2006).
129 Hubbe, M., Hanihara, T. & Harvati, K. Climate Signatures in the Morphological Differentiation
of Worldwide Modern Human Populations. *The Anatomical Record* **292**, 1720-1733,
doi:doi:10.1002/ar.20976 (2009).
130 Noback, M. L. & Harvati, K. The contribution of subsistence to global human cranial
variation. *Journal of human evolution* **80**, 34-50, doi:10.1016/j.jhevol.2014.11.005 (2015).
131 Schmidt, K. L. & Cohn, J. F. Human facial expressions as adaptations: Evolutionary questions
in facial expression research. *American journal of physical anthropology* **116**, 3-24,
doi:doi:10.1002/ajpa.20001 (2001).
132 Trinkaus, E. An abundance of developmental anomalies and abnormalities in Pleistocene
people. *Proceedings of the National Academy of Sciences* **115**, 11941-11946,
doi:10.1073/pnas.1814989115 (2018).
133 Fu, Q. *et al.* An early modern human from Romania with a recent Neanderthal ancestor.
Nature **524**, 216-219, doi:10.1038/nature14558 (2015).
134 Gunz, P. *et al.* Early modern human diversity suggests subdivided population structure and a
complex out-of-Africa scenario. *Proceedings of the National Academy of Sciences* **106**, 6094-
6098, doi:10.1073/pnas.0808160106 (2009).
135 Scerri, E. M. L. *et al.* Did Our Species Evolve in Subdivided Populations across Africa, and Why
Does It Matter? *Trends in Ecology & Evolution* **33**, 582-594, doi:10.1016/j.tree.2018.05.005
(2018).
136 Franciscus, R. & Vlček, E. in *Early Modern Human Evolution in Central Europe: The*
People of Dolní Věstonice and Pavlov, (eds E Trinkaus & JA Svoboda) 63-152 (Oxford
University Press, 2006).
137 Galway-Witham, J. & Stringer, C. How did Homo sapiens evolve? *Science* **360**,
1296-1298, doi:10.1126/science.aat6659 (2018).
138 Ríos, L. *et al.* Skeletal Anomalies in The Neandertal Family of El Sidrón (Spain) Support A Role
of Inbreeding in Neandertal Extinction. *Scientific Reports* **9**, 1697, doi:10.1038/s41598-019-
38571-1 (2019).
139 Dennell, R. W., Martinón-Torres, M. & Bermúdez de Castro, J. M. Hominin variability,
climatic instability and population demography in Middle Pleistocene Europe. *Quaternary*
Science Reviews **30**, 1511-1524, doi:https://doi.org/10.1016/j.quascirev.2009.11.027 (2011).
140 Charpentier, M. J. E., Widdig, A. & Alberts, S. C. Inbreeding depression in non-human
primates: a historical review of methods used and empirical data. *American Journal of*
Primatology **69**, 1370-1386, doi:doi:10.1002/ajp.20445 (2007).

141 G. Rawlins, R. & J. Kessler, M. *Congenital and hereditary anomalies in the rhesus monkeys*
(Macaca mulatta) of Cayo Santiago. Vol. 28 (1983).
142 Nakamichi, M., Nobuhara, H., Nobuhara, T., Nakahashi, M. & Nigi, H. *Birth rate and mortality*
rate of infants with congenital malformations of the limbs in the Awajishima free-ranging
group of Japanese monkeys (Macaca fuscata). Vol. 42 (1997).
143 V Chalifoux, L. & W Elliott, M. *Congenital anomalies in two neonatal tamarins (Saguinus*
oedipus and Saguinus fuscicollis). Vol. 15 (1986).
144 van der Valk, T., Díez-del-Molino, D., Marques-Bonet, T., Guschanski, K. & Dalén, L. Historical
Genomes Reveal the Genomic Consequences of Recent Population Decline in Eastern
Gorillas. *Current Biology* **29**, 165-170.e166, doi:10.1016/j.cub.2018.11.055 (2019).
145 Mayr, E. *Systematics and the Origin of Species*. (Columbia University Press, 1942).
146 Coyne, J. A. & Orr, H. A. *Speciation*. (Sinauer Associates, 2004).
147 Harvati, K. The Neanderthal taxonomic position: models of intra- and inter-specific
craniofacial variation. *Journal of human evolution* **44**, 107-132 (2003).
148 Ackermann, R. R., Mackay, A. & Arnold, M. L. The Hybrid Origin of "Modern" Humans.
Evolutionary Biology **43**, 1-11, doi:10.1007/s11692-015-9348-1 (2016).
149 Mallet, J. Hybridization, ecological races and the nature of species: empirical evidence for
the ease of speciation. *Philosophical Transactions of the Royal Society B* **363**, 2971–2986
(2008).
150 Stringer, C. *et al.* in *Neanderthals and Modern Humans in the European Landscape During*
the Last Glaciation: Archaeological Results of the Stage 3 Project   (eds TH. Van Andel & W.
Davies) Ch. 13, 233-240 (McDonald Institute for Archaeological Research, 2004).
151 Timmermann, A. Quantifying the potential causes of Neanderthal extinction: Abrupt climate
change versus competition and interbreeding. *Quaternary Science Reviews* **238**, 106331,
doi:https://doi.org/10.1016/j.quascirev.2020.106331 (2020).
152 Holliday, T. Species concepts, reticulation, and human evolution. *Current Anthropology* **44**,
653-673 (2003).
153 Valladas, H. *et al.* TL Dates for the Neanderthal Site of the Amud Cave, Israel. *Journal of*
Archaeological Science **26**, 259-268, doi:https://doi.org/10.1006/jasc.1998.0334 (1999).
154 Bahain, J. J., Sarcia, M. N., Falguères, C. & Yokoyama, Y. Attempt at ESR dating of tooth
enamel of French middle Pleistocene sites. *Applied Radiation and Isotopes* **44**, 267-272,
doi:https://doi.org/10.1016/0969-8043(93)90230-8 (1993).
155 Grün, R. & Stringer, C. *ESR dating and the evolution of modern humans*. Vol. 33 (2007).
156 Schmitz, R. W. *et al.* The Neandertal type site revisited: Interdisciplinary investigations of
skeletal remains from the Neander Valley, Germany. *Proceedings of the National Academy of*
Sciences **99**, 13342-13347, doi:10.1073/pnas.192464099 (2002).
157 Guérin, G. *et al. A Multi-Method Luminescence Dating of the Palaeolithic Sequence of La*
Ferrassie Based on New Excavations Adjacent to the La Ferrassie 1 and 2 Skeletons. Vol. 58
(2015).
158 Oakley, K., Campbell, B. & Molleson, T.   (Mus. Nat. Hist., London, 1971).
159 Grün, R. & Stringer, C. B. Electron spin resonance dating and the evolution of modern
humans. *Archaeometry* **33**, 153-199, doi:10.1111/j.1475-4754.1991.tb00696.x (1991).
160 Schwarcz, H. P. *et al.* On the Reexamination of Grotta Guattari: Uranium-Series and Electron-
Spin-Resonance Dates. *Current Anthropology* **32**, 313-316, doi:10.1086/203959 (1991).
161 Rink, W. J., P Schwarcz, H., H Smith, F. & Radovĉić. *ESR ages for Krapina hominids [5]*. Vol.
378 (1995).
162 Debénath, A. & Jelinek, A. J. *Nouvelles fouilles à La Quina (Charente). Résultats préliminaires*.
Vol. 40 (1998).
163 Vandermeersch, B. & Trinkaus, E. The postcranial remains of the Régourdou 1 Neandertal:
the shoulder and arm remains. *Journal of human evolution* **28**, 439-476,
doi:https://doi.org/10.1006/jhev.1995.1034 (1995).

1010	164	Marra, F. *et al.* The Aggradational Successions of the Aniene River Valley in Rome: Age
1011		Constraints to Early Neanderthal Presence in Europe. *PLOS ONE* **12**, e0170434,
1012		doi:10.1371/journal.pone.0170434 (2017).
1013	165	Solecki, R. S. *Shanidar, the first flower people*. (Knopf, 1971).
1014	166	Devièse, T. *et al.* Reevaluating the timing of Neanderthal disappearance in Northwest
1015		Europe. *Proceedings of the National Academy of Sciences* **118**, e2022466118,
1016		doi:doi:10.1073/pnas.2022466118 (2021).
1017	167	Grün, R. & Stringer, C. *Tabun revisited: Revised ESR chronology and new ESR and U-series*
1018		*analyses of dental material from Tabun C1*. Vol. 39 (2001).
1019	168	Grün, R. *et al.* U-series and ESR analyses of bones and teeth relating to the human burials
1020		from Skhul. *Journal of human evolution* **49**, 316-334,
1021		doi:https://doi.org/10.1016/j.jhevol.2005.04.006 (2005).
1022	169	Michel, V., Delanghe-Sabatier, D., Bard, E. & Barroso Ruiz, C. U-series, ESR and 14C studies of
1023		the fossil remains from the Mousterian levels of Zafarraya Cave (Spain): A revised
1024		chronology of Neandertal presence. *Quaternary Geochronology* **15**, 20-33,
1025		doi:https://doi.org/10.1016/j.quageo.2012.12.008 (2013).
1026	170	Wood, R. E. *et al.* Radiocarbon dating casts doubt on the late chronology of the Middle to
1027		Upper Palaeolithic transition in southern Iberia. *Proceedings of the National Academy of*
1028		*Sciences* **110**, 2781-2786, doi:10.1073/pnas.1207656110 (2013).
1029	171	Haile-Selassie, Y., Asfaw, B. & White, T. D. Hominid cranial remains from upper pleistocene
1030		deposits at Aduma, Middle Awash, Ethiopia. *American journal of physical anthropology* **123**,
1031		1-10, doi:10.1002/ajpa.10330 (2004).
1032	172	Grine, F. *et al. Late Pleistocene human skull from Hofmeyr*. Vol. 315 (2007).
1033	173	Schwartz, J. H. & Tattersall, I. *The Human Fossil Record, Craniodental Morphology of Genus*
1034		*Homo (Africa and Asia) (Volume 2)*. (Wiley-Liss, 2003).
1035	174	Wood, B. *Wiley-Blackwell Encyclopedia of Human Evolution 1st Edition*. (Wiley-Blackwell,
1036		2011).
1037	175	Day, M. H., Leakey, M. D. & Magori, C. A new hominid fossil skull (L.H. 18) from the Ngaloba
1038		Beds, Laetoli, northern Tanzania. *Nature* **284**, 55-56, doi:10.1038/284055a0 (1980).
1039	176	Leakey, M. D. & Harris, J. M. *Laetoli: A Pliocene Site in Northern Tanzania*. 05/01 edn, Vol.
1040		125 (Oxford University Press, 1987).
1041	177	Vidal, C. M. *et al.* Age of the oldest known Homo sapiens from eastern Africa. *Nature* **601**,
1042		579-583, doi:10.1038/s41586-021-04275-8 (2022).
1043	178	McDougall, I., Brown, F. H. & Fleagle, J. G. Stratigraphic placement and age of modern
1044		humans from Kibish, Ethiopia. *Nature* **433**, 733-736, doi:10.1038/nature03258 (2005).
1045	179	Schild, R. & Wendorf, F. Palaeolithic Living Sites in Upper and Middle Egypt: A Review Article.
1046		*Journal of Field Archaeology* **29**, 447-461, doi:10.2307/3250903 (2002).
1047	180	Mellars, P. A., Bricker, H. M., Gowlett, J. A. J. & Hedges, R. E. M. Radiocarbon Accelerator
1048		Dating of French Upper Palaeolithic Sites. *Current Anthropology* **28**, 128-133 (1987).
1049	181	Holt, B. M. & Formicola, V. Hunters of the Ice Age: The biology of Upper Paleolithic people.
1050		*American journal of physical anthropology* **137**, 70-99, doi:10.1002/ajpa.20950 (2008).
1051	182	Barshay-Szmidt, C. *et al.* New extensive focused AMS 14C dating of the Middle and Upper
1052		Magdalenian of the western Aquitaine/Pyrenean region of France (ca. 19–14 ka cal BP):
1053		Proposing a new model for its chronological phases and for the timing of occupation.
1054		*Quaternary International* **414**, 62-91, doi:https://doi.org/10.1016/j.quaint.2015.12.073
1055		(2016).
1056	183	Soficaru, A., Petrea, C., Doboş, A. & Trinkaus, E. The Human Cranium from the Peştera
1057		Cioclovina Uscată, Romania: Context, Age, Taphonomy, Morphology, and Paleopathology.
1058		*Current Anthropology* **48**, 611-619, doi:10.1086/519915 (2007).
1059	184	Henry-Gambier, D. *Les fossiles de Cro-Magnon (Les Eyzies-de-Tayac, Dordogne): Nouvelles*
1060		*données sur leur position chronologique et leur attribution culturelle*. Vol. 14 (2002).

1061   185   Trinkaus, E. & Svoboda, J. *Early modern human evolution in central Europe: the people of*
1062   *Dolní Věstonice and Pavlov*. (2006).
1063   186   Formicola, V., Pettitt, P. B. & Del Lucchese, A. A Direct AMS Radiocarbon Date on the Barma
1064   Grande 6 Upper Paleolithic Skeleton. *Current Anthropology* **45**, 114-118,
1065   doi:10.1086/381008 (2004).
1066   187   Schwartz, J. H. & Tattersall, I. *The Human Fossil Record, Terminology and Craniodental*
1067   *Morphology of Genus I Homo/I (Europe) (Volume 1)*. (Wiley-Liss, 2002).
1068   188   Wild, E. M. *et al.* Direct dating of Early Upper Palaeolithic human remains from Mladeč.
1069   *Nature* **435**, 332-335, doi:10.1038/nature03585 (2005).
1070   189   Trinkaus, E. *et al.* An early modern human from the Peştera cu Oase, Romania. *Proceedings*
1071   *of the National Academy of Sciences* **100**, 11231-11236, doi:10.1073/pnas.2035108100
1072   (2003).
1073   190   Street, M., Terberger, T. & Orschiedt, J. A critical review of the German Paleolithic hominin
1074   record. *Journal of human evolution* **51**, 551-579,
1075   doi:https://doi.org/10.1016/j.jhevol.2006.04.014 (2006).
1076   191   Svoboda, J. A. The Upper Paleolithic burial area at Předmostí: ritual and taphonomy. *Journal*
1077   *of human evolution* **54**, 15-33, doi:https://doi.org/10.1016/j.jhevol.2007.05.016 (2008).

1078