Speciation without gene-flow in hybridising deer

Camille Kessler[1], Eric Wootton[2], Aaron B.A. Shafer[1]

[1]Environmental and Life Sciences Graduate Program, Trent University, 7K9J 7B8 Peterborough,

Ontario, Canada

[2]Biochemistry & Molecular Biology, Trent University, K9J 7B8 Peterborough, Ontario, Canada

Corresponding authors:

Camille Kessler: camillekessler@trentu.ca

Aaron Shafer: aaronshafer@trentu.ca

Abstract

Under the ecological speciation model, divergent selection acts on ecological differences

between populations, gradually creating barriers to gene flow and ultimately leading to

reproductive isolation. Hybridisation is part of this continuum and can both promote and

inhibit the speciation process. Here, we used white-tailed (*Odocoileus virginianus*) and mule

deer (*O. hemionus*) to investigate patterns of speciation in hybridising sister species. We

quantified genome-wide historical introgression and performed genome scans to look for

signatures of four different selection scenarios. Despite ample modern evidence of

hybridisation, we found negligible patterns of ancestral introgression and no signatures of

divergence with gene flow, rather localised patterns of allopatric and balancing selection were

detected across the genome. Genes under balancing selection were related to immunity, MHC

and sensory perception of smell, the latter of which is consistent with deer biology. The

deficiency of historical gene-flow suggests that white-tailed and mule deer were spatially

separated during the glaciation cycles of the Pleistocene and genome wide differentiation

accrued via genetic drift. Dobzhansky-Muller incompatibilities and selection against hybrids

are hypothesised to be acting, and diversity correlations to recombination rates suggests these

sister species are far along the speciation continuum.

29
30 Keywords: Speciation genomics, hybridisation, allopatric speciation, secondary contact,

31 *Odocoileus*.

Introduction

Hybridisation is a widespread phenomenon that occurs at variable rates (Adavoudi & Pilot, 2021; Iacolina et al., 2019; Mallet, 2005; Ragavan et al., 2017; Taylor & Larson, 2019). The prevalence of hybridisation suggests speciation follows a continuum as opposed to the more discrete and allopatric view introduced by Mayr (1942) (Mallet, 2005; Stankowski & Ravinet, 2021). However, both ideas are not incompatible as the speciation continuum ultimately leads to reproductive isolation with hybridisation being a natural outcome (Stankowski & Ravinet, 2021). Under the ecological speciation model (sensu Darwin, 1859), hybrids are common and can even facilitate speciation (Nosil, 2012). Ecological speciation works when ecological differences lead to divergent selection between populations; the underlying genetic mechanisms (i.e. causative loci) gradually create barriers to gene flow that accumulate which leads to reproductive isolation (Nosil, 2012; Rundle & Nosil, 2005; Shafer & Wolf, 2013).

The genetic mechanisms involved in ecological speciation can take the form of barrier loci which act as a restraint to gene flow in different ways: for example, such loci can be under divergent selection, involved in mate choice, contribute to assortative mating, or reduce hybrid fitness (Abbott et al., 2013; Ravinet et al., 2017). Divergent selection should produce genomic islands in this context, also referred to as genomic islands of speciation, that are regions where the differentiation (e.g. F_{ST}) is higher than the neutral genomic background (Campbell et al., 2018; Ravinet et al., 2017). Such loci and islands have been observed in several species underpinning a wide array of speciation and divergence processes (Lavretsky et al., 2019; Marques et al., 2019; Momigliano et al., 2017; Poelstra et al., 2014). Scanning the genome for just F_{ST} peaks when speciation with gene flow is ongoing, however, is problematic as F_{ST} is dependent on genetic diversity (π), and other mechanisms can create similar F_{ST} profiles such

as genetic drift, global adaptation, or simply reduced genetic diversity due to background

selection (Booker et al., 2021; Burri, 2017; Cruickshank & Hahn, 2014; Ravinet et al., 2017). Joint

genome scans including F_{ST}, d_{xy} and π facilitate a better depiction of the processes of selection

and ecological speciation, particularly the use of d_{xy} which is not influenced by current levels

of diversity (Campbell et al., 2018; Cruickshank & Hahn, 2014; Ravinet et al., 2017; Shang et al.,

2021). Shang et al. (2021) recently conceptualised the expected patterns F_{ST}, d_{xy} and π under

four main modes of selection: divergence with gene flow, allopatric selection (i.e., positive

selection post species split), balancing selection and background selection. Using pairs of

Populus species along the speciation continuum, they investigated the genomic landscape to

test clear predictions on the behaviour of F_{ST}, d_{xy} and π. Including correlations of π and F_{ST} to

recombination rate (ρ) can help place a species on the continuum, with for example the

correlation between F_{ST} and ρ is expected to become stronger with increasing divergence

(Burri, 2017; Shang et al., 2021). This approach ultimately allows for disentangling selection

patterns from a neutral background based on those statistics, and helps decipher the

evolutionary history of species.

White-tailed (*Odocoileus virginianus*; WTD) and mule deer (*O. hemionus*; MD) are

abundant in North America with similar morphology, activity patterns and life-history traits,

but they differ in several ecological aspects (Douzery & Randi, 1997; Gilbert et al., 2006; Pitra et

al., 2004; Brunjes et al., 2006; Berry et al., 2019). Mule deer favour food availability whereas WTD

prioritise security and thermoregulation (Whittaker & Lindzey, 2004); consequently, WTD prefer

habitats at lower altitude and with denser visual cover than MD that favour open areas at higher

elevation (Anthony & Smith, 1977; Brunjes et al., 2006). White-tailed deer and MD also differ in

their sociality and associated predator response. Mule deer live in large cohesive groups

including both sexes, and, as a group, adopt an aggressive behaviour in presence of predators,

whereas WTD live in smaller female-biased groups and flee in response to predators (Lingle,

2001, 2003). Both species represent a high economic value in North America as hunting-related

activities generate billions of dollars annually (Cambronne, 2013), and both species are an

important cultural component of Indigenous communities (Adams & Hamilton, 2011; Peres &

Altman, 2018).

Despite a species divergence date estimated at ~3.13 mya (Wright et al., 2022), WTD and

MD hybridise in areas of sympatry with estimated hybridisation rates ranging from 1 to 19%

depending on the region (Carr & Hughes, 1993; Combe et al., 2021; Hornbeck & Mahoney, 2000;

Russell et al., 2021). Based on divergent species distributions (Fig. 1A) and likely separate

refugia during glacial events (Greenslade, 1998), both species likely spent considerable time in

allopatry. Recent hybrid zone analyses mainly found backcrossed individuals rather than F1

hybrids (Combe et al., 2021; Russell et al., 2021), and an interspecies F_{ST} up to 0.4 (Combe et al.,

2021). It has also been shown that gene flow appears restricted but bidirectional, and there are

signs of introgression at both mitochondrial and nuclear levels (Bradley et al., 2003; Carr et al.,

1986; Carr & Hughes, 1993; Cathey et al., 1998; Cronin, 1991; Derr, 1991; Russell et al., 2019),

such that some MD acquired WTD mitochondrial DNA around ~1.32 Mya (Wright et al., 2022).

This pattern would suggest ancestral hybridisation and gene flow has taken place. Given their

clear behavioural differences, and their documented hybridisation, white-tailed and mule deer

do not present the reproductive isolation required by Mayr's biological species concept (Mayr,

1942). Here, we hypothesised that WTD and MD have evolved via ecological speciation, and

more specifically a speciation with gene flow scenario. To test this hypothesis, we quantified

genome-wide introgression and past admixture events and performed genome scans to look

for signatures of four different divergence scenarios. We expected to find genetic signs of

divergent selection, such as speciation islands, consistent with patterns of divergence with

gene flow (DwGF), and a stronger F_{ST} - ρ correlation between species compared to within. We

also predicted higher rates of admixture and historical introgression in areas of sympatry.

Materials and Methods

Sampling & Sequencing

We obtained tissue samples from harvested deer collected across the range of WTD & MD

including areas of sympatry and allopatry (Fig. 1A, Table 1). These areas were determined from

NatureServe records and adjusted with IUCN range data (IUCN, 2015a, 2015b; NatureServe,

2021a, 2021b). Specifically, the state of Washington and the province of British Columbia, for

which the coasts are largely MD allopatric whereas the eastern parts are sympatric, were

divided in two by the 120[th] meridian West. We extracted DNA from tissue using the Qiagen

DNeasy Blood and Tissue Kit following manufacturer's instructions and checked the samples

concentration using Invitrogen Qubit assays. WGS libraries were generated at The Centre for

Applied Genomics in Toronto, Canada and sequenced to an average of 4x coverage on an

Illumina HiSeqX.

Data processing

Raw reads quality was examined using FastQC (v0.11.9; Andrews S., 2010). We trimmed the

reads using Trimmomatic (v0.36; Bolger et al., 2014), and aligned them to the WTD genome

(GCA_014726795.1) with bwa-mem (v0.7.17; Li & Durbin, 2009). We sorted the reads using

SAMtools sort (v1.10; Li et al., 2009), identified duplicates reads with Picard MarkDuplicates

(v2.23.2; Broad Institute, GitHub Repository., 2019) and removed them using Sambamba view

(v0.7.0; Tarasov et al., 2015). We performed a local re-alignment using GATK

RealignerTargetCreator and IndelRealigner (v4.1.7.0; McKenna et al., 2010). For further quality

checks, we used Sambamba flagstat, mosdepth (v0.3.1; Pedersen and Quinlan 2018) and

MultiQC (v1.10; Ewels et al., 2016). For the MSMC analysis, some samples were sequenced to a

higher depth on multiple lanes. For those, before the duplicate removal step, the read groups

were specified and both sequencing files merged using Picard AddOrReplaceReadGroups and

MergeSamFiles. The rest of the pipeline was performed as described above with the addition

of a final step: to be able to carry out other analyses with a coverage consistent across samples,

we reduced the final coverage to the average coverage of all other samples (4x) using Picard

DownsampleSam.

We created three datasets for our different analyses. The first contained the variants

among all deer individuals (hereafter called DeerSNP dataset). We produced it using ANGSD

(v0.918; Korneliussen et al., 2014) and estimated genotype likelihoods following the GATK

model (-gl 2) and called genotypes (-doGeno 4) with the following filtering: SNPs with a

minimum p-value of $1e^{-6}$, a minimum base quality of 20 and Minimum mapQ quality of 20 (-

SNP_pval 1e-6, -minMapQ 20 & -minQ 20). The second dataset additionally included invariant

sites (referred to as DeerALL), it was produced following the same method as DeerSNP except

for the filtering on the SNP p-value (-SNP_pval 1e-6) which was removed. Our third dataset was

designed for an ABBA-BABA analysis which requires samples from an additional species as

outgroup. It comprises the 28 deer individuals and one caribou sample, all mapped to the

caribou genome (GCA_014898785.1; afterwards called DeerCAR dataset). For this dataset, we

first sorted the deer bam files by read name and converted them to fastq using SAMtools sort

and fastq respectively. We mapped these fastq to the caribou reference genome using bwa-

mem. We ran ANGSD using the same procedure as for DeerSNP dataset, with the addition of a

caribou genome sequence in bam format, obtained for this analysis from Dedato et al. (2022).

Population structure analyses and cross-species coalescent rate

To infer admixture proportions and compute a PCA on allele frequencies based on genotype

likelihood data we used PCAngsd (Meisner & Albrechtsen, 2018) and NGSadmix (Skotte et al.,

2013) on the DeerSNP dataset. NGSadmix was run with K = 1 up to K =7 and the best K value

was determined with CLUMPAK (Kopelman et al., 2015) following the Evanno method (Evanno

et al., 2005) with 7 replicates for each K value. We investigated the correlation between the

distribution on each PC and each individual's range in R using the package "stats" with a linear

model including the species and either latitude or longitude, and a Pearson's correlation (e.g.

lm(PC ~ Species + Latitude)).

For two samples with high coverage (Ov_ON6 at 18x - and Oh_WA1 at 22x) we

implemented the multiple sequentially Markovian coalescent (MSMC) to infer the cross-

coalescent rate. To do this we first generated a 35-mer mappability mask file using the SNPable

pipeline as is required for MSMC (Schiffels & Wang, 2020). We restricted analyses to scaffolds

>500 Kbp (Gower et al., 2018) and phased samples using whatsHap 1.3 (Martin et al., 2016) prior

to generating the MSMC input files. We ran MSMC2 using the time segment pattern

1×2+25×1+1×2+1×3 and estimated the cross-coalescence rate (CCR) across populations by

comparing the first haplotype from each sample in a pairwise fashion using the -P flag and

skipping any sites with ambiguous phasing. To quantify the migration rate over time m(t) and

the cumulative migration probability M(t) – which is related to the CCR (Schiffels & Durbin,

2014) -– we implemented the IM model to the MSMC output (Wang et al., 2020). We assumed

a generation time of 2 years (Demarais et al., 2000; Deyoung et al., 2003) and mutation rate of

1.33×10^{-8} mutations/site/generation.

Genome-wide ancestral introgression

To evaluate the extent of historical introgression between sympatric MD and WTD, we used the

ABBA-BABA test, which allows for the detection of introgression between three populations P1,

P2 & P3 and an outgroup (Durand et al., 2011; Green et al., 2010). We ran the ABBA-BABA

analysis implemented in ANGSD on the DeerCAR dataset between individuals (-doAbbababa

1) and between populations (-doAbbababa2 1; Soraggi et al. 2018), using the caribou as

outgroup. The D-statistic, standard error and Z-score were computed with ANGSD's

accompanying R scripts: jackKnife.R and estAvgError.R. To further our understanding of the

admixture events and build a maximum likelihood tree of our system, we used treemix (v1.13;

Pickrell & Pritchard, 2012) on both DeerCAR and DeerSNP datasets. We constructed the

maximum likelihood trees with migration events ranging from 0 to 5, either WTD_allopatry or

Caribou as root, and accounted for linkage using 1000 SNPs per block. Tree and residuals

visualisations were performed with associated R script plotting_funcs.R and the variance

explained by each migration event was computed with the get_f() function.

In-windows historical introgression

The D statistic is sensitive to genomic variation (π), and should not be used to determine

introgression on a small scale (Martin et al., 2015). The f_d statistic, proposed by Martin et al.

(2015), is less dependent on diversity than D and therefore allows for the detection of

potentially introgressed regions of the genome. We used the python script

ABBABABAwindows.py (Martin, 2021) to estimate in D and f_d in 50Kbp windows to detect

potentially introgressed loci in our DeerCAR dataset on two comparisons: 1) P1 = WTD

allopatry, P2 = WTD sympatry, P3 = MD sympatry and 2) P1 = MD allopatry, P2 = MD sympatry,

P3 = WTD sympatry, both with the caribou as outgroup. We then identified potentially

introgressed windows as those having a f_d value higher than the 97.5% quantile. Their position

was used in BEDTools intersect (v2.30.0; Quinlan and Hall 2010) with the caribou annotation

file to identify the genes present in putatively introgressed windows.

Genome scans

To detect islands of divergence between WTD & MD, we used the python script

popgenWindows.py (Martin, 2021) to estimate individual heterozygosity, F_{ST}, D_{xy} and π in 50Kbp

windows (note we filtered out scaffolds with less than 2 windows). Since the accurate

computation of D_{xy} and π require a dataset including the invariant sites, we performed this

analysis on the DeerALL dataset. To detect the outlier loci, we based our approach on the four

selection scenarios developed in (Shang et al., 2021): I) Divergence with gene flow that

presents as high F_{ST} and D_{xy} but low π; II) allopatric selection that shows high F_{ST}, low π and

stable D_{xy}; III) background selection that presents as high F_{ST} but low π and D_{xy}; and IV)

balancing selection that displays as low F_{ST} but high π and D_{xy}. Thresholds were set as the upper

or lower 5% for a high or low criteria, and between 45 and 55% for a stable condition. Outliers

were flagged as belonging to one of the four scenarios when they met the specific criteria

shown in Table 2.

We identified genes present in outlier regions by comparing the coordinates of the

outlier windows and the WTD annotation file in BEDTools intersect. We extracted the gene

names from those regions as well as those from their associated scaffolds. The gene list from

outlier windows was uploaded to ShinyGO for enrichment analysis with the scaffold's gene list

as background information and no species specification. We downloaded the enrichment

analysis results for the molecular function, biological process, and cellular component GOs

and analysed the results in R. We also uploaded the gene list to UniProt's Retrieve/ID mapping

tool (Pundir et al., 2016) and downloaded the information for visual inspection.

To estimate the position of WTD & MD along the speciation continuum through the

correlation between recombination rate (ρ), F_{ST}, and π, we computed ρ in windows using

FastEPRR (v.2.0; Gao et al., 2016). Here, we used our four categories as a proxy for a speciation

continuum with six comparisons: WTDa - WTDs and MDa - MDs representing no speciation,

WTDs - MDs, WTDs - MDa and WTDa - MDs as intermediate stages and WTDa - MDa as full

differentiation. We first phased the vcf using SHAPEIT (Delaneau et al., 2012) and split the

resulting file into scaffolds. We then ran FastEPRR separately on our four populations to obtain

the ρ estimation in 50kbp windows (winLength = 50000) on the 15 longest scaffolds of the WTD

genome, with 100 repetitions. To be able to correlate ρ with F_{ST} and π, we re-ran the genome

scan analysis on the population level. FastEPRR outputs ρ in $4N_e r$, to convert this value to ρ in

cM/Mb, we computed N_e per population using the relationship $\pi = 4N_e\mu$. We used $\mu= 1.33\times10^{-8}$

8 and the genome scans result to average π across windows in each population. After filtering

to keep only windows that were present across all comparisons (12,500 windows), we

generated Pearson's correlations between I) ρ in both populations (ρ_1 vs ρ_2), II) average ρ

between populations and F_{ST} (ρ_{1-2} vs F_{ST}), and III) average ρ and average π between populations

(ρ_{1-2} vs π_{1-2}) following Shang et al. (2021). The resulting correlation coefficients were compared

between all comparisons to infer trends over divergence time. All results were analysed in R

version 4.1.0 "Camp Pontanezen" (R Core Team, 2021) .

Results

We sequenced 28 individuals to an average coverage of approximately 4x. We called

103,970,889 SNPs in the DeerSNP dataset, which represented 4% of the genome. In the PCA

based on allele frequencies, PC1 accounted for 59% of the variation and separated both

species (Fig. 1B). PC2 explained an additional 8% of the variation and showed a spread of MD

individuals that is consistent with their longitudinal distribution (Pearson correlation = 0.8; *p-*

value = 0.00057). Latitude was not important in shaping PC1 or PC2 variation (Pearson

correlation *p-value* > 0.05). Admixture analysis showed a well-defined genetic clustering

between species (Evanno's method: best K = 2, Fig. 1C, Fig. S1). The WTD individual Ov_MX2

was partially admixed in both analyses, with NGSAdmix assigning it 92.5% WTD and 7.44% MD

ancestry (Fig. 1C). This individuals' admixed pattern disappeared in the admixture analysis with

higher K values (K = 3 to 7, Fig. S1), though additional within species clusters are observed.

The MSMC2-IM model showed no support for migration between species once they split, with

the CCR suggesting the speciation event for *Odocoileus* took place just under 1 Mya (Fig. 2).

To detect historical gene flow, we performed an ABBA-BABA test based on genotype

likelihoods between all combinations of our 28 individuals in a first analysis and between all

four populations in a second, with the caribou as outgroup (DeerCAR dataset, 72,438,766

SNPs). We computed 9828 ABBA-BABA individual topologies in ANGSD, of which 30 presented

signatures of introgression. We found six potential inter-species introgression events with a D-

statistic ranging from 0.011 to -0.013, two of which are significant (Z-score > |3| (Barlow et al.,

2021; Kirch et al., 2021), Fig. 3A). D-statistic in intra-species comparisons range from 0.25 to

0.916 in ABBA and from -0.046 to -0.914 in BABA, all intra-species comparison show a significant

Z-score (Fig. 3A). In the comparisons between the four populations, three ABBA-BABA

topologies presented signatures of introgression on a total of 12. The single inter-species

comparison presents a D-statistic of -0.07 (Z-score = -7.66) for an introgression between

sympatric MD and WTD. The other two comparisons show a reciprocal introgression between

MD in allopatry and sympatry of 0.9 and 0.89 (Z-score = 1342.83 & 1376.69; Fig. S3). The treemix

analyses showed a topology with no migration that explained over 99% of the variation (Fig.

3B). We further analysed historical introgression through the measure of f_d across 50kb non-

overlapping windows in two different introgression scenarios: 1) WTDa, WTDs, MDs, Caribou

and 2) MDa, MDs, WTDs, Caribou (Fig. S4). We found 1243 outlier windows with a f_d higher than

the 97.5% percentile in the first scenario and 1248 in the second, each representing ~2.45% of

the genome with 68 windows overlapping between comparisons (Fig. S4).

268   Measures of relative genomic divergence between WTD and MD were elevated (F_{ST} =

0.26) and absolute divergence d_{xy} was 0.011; genetic diversity (π) was higher in WTD than in MD

(WTD = 0.008, MD = 0.004, Mann-Whitney test: p < 2.2e-16). We identified a total of 1183 windows

presenting patterns consistent with one of the four selection scenarios (Fig. 4, Fig. S5). Of those,

1016 suggest a pattern of balancing selection, they were distributed across 236 scaffolds and

represent 1.99% of the genome. Those windows contained 121 genes, some of which showed

an enrichment in ontologies associated with the sense of smell and chemical stimuli, including

three categories presenting an enrichment above 15-fold (Fig. S6). We also detected GOs

related to the MHC and immunity, these include three categories with an enrichment above

24-fold (Fig. S6). We identified patterns of background selection in 165 windows across 58

scaffolds, representing 0.32% of the genome. These windows harboured 208 genes identified

with a UniProt ID and for which we found enrichment in GOs related to epigenetic factors such

as "Unmethylated CpG binding" (25-fold) or "Histone demethylation" (15-fold) (Fig. S7). These

windows were either isolated or clustered together into putative islands of divergence (Fig. S5).

We found 2 windows under allopatric selection, each window containing one gene: ACAP2, a

GTPase activating protein, and PCDHB4 potentially involved in cell-binding (UniProt, 2022a,

2022b). When we applied a more liberal cut-offs for F_{ST}, d_{xy}, and π we still observed no evidence

for divergence with gene flow (Table S1). The correlations between π, F_{ST} and ρ are highly

significant in all six comparisons (Fig. S8). As predicted correlations coefficients for F_{ST} - ρ

became stronger with divergent comparisons (Fig. 5). Likewise, the π - ρ correlation coefficients

were constant over divergence time, while the recombination landscape (ρ - ρ) differed

between species (Fig. 5).

Discussion

We investigated the speciation history of the *Odocoileus* genera through introgression analyses

and genome scans. Our results suggest speciation between WTD & MD took place with

negligible historical gene flow, despite contemporary and historical evidence of interbreeding

(Cathey et al., 1998; Combe et al., 2021; Cronin et al., 1988; Cronin, 1991; Derr, 1991; Hornbeck

& Mahoney, 2000; Hughes & Carr, 1993; Russell et al., 2021; Stubblefield et al., 1986). Consistent

with other estimates of divergence (Combe et al., 2021; Douzery & Randi, 1997), WTD and MD

appear to have split ~1 Mya (Fig. 2). The main pattern of selection, however, appears to be that

of balancing selection, with patterns of divergence with gene flow notably absent. Moreover,

the scarcity of allopatric selection (i.e., positive selection post divergence) suggests isolation

and drift primarily underlies the species differentiation, with WTD and MD likely only recently

coming into secondary contact. The genome-wide scans combined with the absence of

introgression suggests that WTD & MD are far along the speciation continuum (i.e. Feder 2012),

despite contemporary hybridisation. Some signals of selection might have been lost by

recombination, elevated F_{ST}, and time, but the paucity of DwGF and AS signals even at liberal

cut-offs suggest speciation was driven primarily by drift.

Speciation with negligible historical introgression

Contemporary hybridisation in wild WTD & MD appears highly variable: rates between 1 and

19% have been observed, which can be explained by the region and degree of overlap, but

also by the methodologies used that vary in resolution (Cathey et al., 1998; Combe et al., 2021;

Cronin et al., 1988; Cronin, 1991; Derr, 1991; Hornbeck & Mahoney, 2000; Hughes & Carr, 1993;

Russell et al., 2021; Stubblefield et al., 1986). Moreover, a proposed historical mtDNA

introgression would suggest ancestral hybridisation occurred (Wright et al., 2022). We therefore

expected to find signs of introgression in our samples (e.g. Combe et al. 2021) assuming that

historic matings would have left some detectable signal as is seen in humans (e.g. Green et al.,

2010) and other species (Liu et al., 2021; Paijmans et al., 2021; Poelstra et al., 2014). Surprisingly,

we observed neglible levels of ancestral gene flow and introgression. While Combe et al. (2021)

did find evidence of introgression between *Odocoileus* species; we note, this inference was

based off a small number of topologies and included recent hybrids, both of which could skew

D values. We did find signs for some contemporary admixture in one sample (Fig. 1C), but

supplementary analysis with an additional WTD from Mexico showed that the admixture

pattern likely reflects population structure rather than hybridisation since it is detected in both

individuals (Fig. S2).

Our sampling surely is limited for the detection of recent hybridisation, as previous

studies suggest gene flow is rather restricted (Cronin et al., 1988; Cronin, 1991; Derr, 1991;

Hughes & Carr, 1993; Russell et al., 2021). For example, Russell et al., (2021) estimated a 1%

hybridisation rate in Alberta by sampling 987 individuals in a range overlap of approximately

230,000 km². While signatures of fd identified potentially introgressed windows in ~1% of the

genome, this metric can be a sign of incomplete lineage sorting (ILS) rather than introgression,

particularly with no signs of genome-wide introgression as evidenced in the D-statistic (Durand

et al., 2011). The independent treemix and MSMC analyses do not support historical gene flow,

thus collectively, we consider the fd signal to be reflective of ILS which would be expected given

the high genome-wide diversity and a relatively recent species-split (Cutter, 2013; Combe et al.

2021).

The absence of signatures of ancestral gene flow suggest that WTD & MD evolved in

allopatry and have recently come in secondary contact. The dynamic history of the North

American continent, notably the glacial cycles of the Quaternary, shaped the evolution and

distribution of many taxa (Avise et al., 1998; G. Hewitt, 2000; Shafer et al., 2010). The use of

different refugia for prolonged periods of time during the glaciation events has increased the

differentiation between populations in several species, including deer (Colella et al., 2021;

Dussex et al., 2020; Ito et al., 2021; Kinoshita et al., 2020; Latch et al., 2014). Previous studies

show that MD persisted in several refugia during the glacial cycles of the Pleistocene, increasing

the intraspecific divergence (Latch et al., 2009, 2014; Wright et al., 2022). This divergence is

highlighted here in the PCA which shows two distinct MD clusters, potentially segregating MD

and its black-tailed deer subspecies (Fig. 1B). Environmental shifts following the LGM also seem

to have impacted the divergence between subspecies of WTD & MD in Florida and Mexico for

example (Alminas et al., 2021; Ellsworth et al., 1994). More recently, a form of allopatry was

mediated through overharvest and habitat destruction of WTD where populations greatly

decreased, even extirpated, in some regions (Budd et al., 2018; Chafin et al., 2021; Deyoung et

al., 2003; McDonald et al., 2004). Here, the core of the current sympatric zone was greatly

affected, notably Colorado, Montana, Idaho, Nebraska and Wyoming where WTD was almost

extirpated before restocking efforts took place starting in the mid 40's (McDonald et al., 2004).

Historical introgression could further be hindered by selection against hybrids, either through

sexual selection, intrinsic or ecological incompatibilities (Nosil, 2012; Rundle & Nosil, 2005,

2005). Given the genome-wide level of differentiation, we hypothesise that Dobzhansky-Muller

incompatibles (DMIs) are in play, noting that they often occur early and rapidly (Schumer et al.,

2015)

357   Little is known about hybrid fitness in the *Odocoileus* genus. Assessment of a single F1

male suggested it was fertile with no overt genetic defects (Derr et al., 1991). In three hybrids

the spermatozoid phenotype showed a gradient of fertility with one sterile individual, one

subfertile or infertile, and one potentially fertile but did not reproduce (Wishart et al., 1988);

sperm phenotypic variation has been implicated in speciation (Albrechtová et al., 2012) and

DMIs (Bhattacharyya et al., 2013). Sexual selection can also be a powerful driver of selection

against hybrids (Servedio, 2004). In WTD, while females might favour males with larger antlers

(Morina et al., 2018), sexual selection is poorly understood and most studies focus on male

breeding success (DeYoung et al. 2006; DeYoung et al. 2009; Jones et al. 2011; Newbolt et al.

2017). Morphologically, however, WTD & MD are similar enough where the identification of their

hybrids is often challenging with the metatarsal gland and their escape gaits being the only

characters consistently distinguishing hybrids (e.g Lingle, 1992; Wishart, 1980). Thus, we

propose a testable hypothesis that ecological incompatibilities combined with DMI maintain

the species boundary. More research is also needed on WTD/MD hybrids to quantify their

fitness and genomic regions underlying any selected hybrid phenotypes.

Mixed signatures of selection underly speciation in North American deer

Given ongoing hybridisation between WTD & MD and a history of recurrent glacial cycles, we

expected to find deer genomes showing patterns of divergence with gene flow. The absence of

introgression and the nonexistence of DwGF in our genome scans, even with more liberal

thresholds, is surprising (Table S1). The further absence of allopatric selection suggests a

diminished role of selection in deer speciation, lending support to drift-induced model more

consistent with Mayr's (1942) view. While the number of windows in allopatric selection

increased with more liberal thresholds (Table S1), they are never the dominant pattern. We

suggest this absence could be a result of species being far along on the speciation continuum

such that patterns of AS (specifically high F_{ST}) are no longer detectable or meaningful because

they were reduced with time and recombination, which is consistent with late stages of

ecological speciation (Feder, 2012). Recombination rates, demography and selection can still

influence the patterns investigated here. We inferred ρ estimates that are consistent with those

in other mammals (Jensen-Seaman et al., 2004), their correlations with divergence would

suggest *Odocoileus* are well defined species and reflects a shift in recombination landscape

between WTD & MD (Fig. 5; (Burri, 2017; Shang et al., 2021).

Signatures of balancing selection (BLS) were widespread with windows containing

genes consistent with deer biology (olfactory receptors) and previous literature (i.e., MHC

genes). The major histocompatibility complex (MHC) is involved in pathogen recognition in

vertebrate species (Bernatchez & Landry, 2003; Piertney & Oliver, 2006). Polymorphism in this

complex is directly linked to disease susceptibility, the most diverse are the most resistant, and

there is evidence that MHC diversity is maintained by BLS (Bernatchez and Landry 2003;

Piertney and Oliver 2006; Aguilar et al. 2004; Pierini and Lenz 2018; Zhang et al. 2018). The same

is true for the results on sensory perception and particularly the olfactory receptors genes, both

of which are under BLS (Liu et al., 2021). Each allele in the olfactory receptors genes is

expressed by a single sensory neuron (Degl'Innocenti & D'Errico, 2017), it is therefore plausible

that diversity in these genes would be maintained by BLS. Olfactory receptors in WTD and MD,

and more broadly deer, is critical to both predator detection and rutting behaviour (Ditchkoff,

2011). Anatomically, WTD has an architecture optimised for the delivery of sensory stimuli to

the receptors in the nose (Ranslow et al., 2014). Chamaillé-Jammes et al. (2014) showed that

MD inspect and respond differently to different predators' olfactory cues, and both human and

mammalian predators utilise wind direction and other mechanism to minimise scent

detection by deer (Cherry & Barton, 2017; Zagata & Haugen, 1974). In other ungulates, olfactory

reception is associated with sexual activity (Cann et al., 2019), maternal behaviour (Blank &

Yang, 2017; Keller & Lévy, 2012), territory choice (Deutsch & Nefdt, 1992), predator response

(Kuijper et al., 2014; Wikenros et al., 2015) and foraging (Hirata & Kusatake, 2021). Low

differentiation (F_{ST}) but high diversity (π) of olfactory receptors in deer is consistent with their

biology and behaviour and would be expected to be under strong selection pressures.

Conclusion

Our results suggests that WTD & MD do not conform to a speciation with gene flow scenario

despite evidence of contemporary hybridisation. We propose they were spatially separated

during the Quaternary glaciation cycles where genome-wide differentiation accrued via drift.

This is evidenced by the majority of the genome (> 97%) not matching a selection scenario.

Increased sampling and model-based demographic assessment should help clarify the role of

glaciers and secondary contact in North American deer. The near absence of patterns of

allopatry and ancestral gene flow suggest that white-tailed and mule deer are far along the

speciation continuum, the absence of introgression signs could suggest DMIs and selection

against hybrids which would contribute to the reinforcement of reproductive isolation. Future

studies should focus on assaying hybrid phenotypes and vigour.

Acknowledgements

Camille Kessler was supported by an International Graduate Scholarship. Eric Wootton was

supported by three Natural Sciences and Engineering Research Council (NSERC)

Undergraduate Student Research Awards. This work was supported by NSERC Discovery Grant

(Grant Number: RGPIN-2017- 03934); ComputeCanada Resources for Research Groups (Grant

Number: RRG gme-665-ab); Canadian Foundation for Innovation: John R. Evans Leaders Fund

and the Ontario Early Researcher Award (Grant Number: #36905). We thank Catherine

Cullingham, Anh Dao, Orrin Duvuvuei, Brad Fulk, Steve Griffin, Levi Jaster, Lee Jeffers, Emily

Latch, the NRDPFC, Charles Ruth, David Walter, Geoff Williams, Kevin White, Mark Wong, Kiana

Young and Liana Zanette, for providing samples. We are also grateful to Andrew Foote, Jose

Alberto Lòpez Alemàn and Emily Latch for their comments on the manuscript. Trent University

is located on the traditional territory of the Michi Saagiig Anishnaabeg. Our sampling also

covers most of North America, a stolen land which remains a home to many First Peoples. We

are grateful to have had the opportunity to work on this land, would like to show our respect

to the First Peoples, and thank them for their care, stewardship, and teachings. Miigwetch.

References

Abbott, R., Albach, D., Ansell, S., Arntzen, J. W., Baird, S. J. E., Bierne, N., Boughman, J.,
Brelsford, A., Buerkle, C. A., Buggs, R., Butlin, R. K., Dieckmann, U., Eroukhmanoff, F.,
Grill, A., Cahan, S. H., Hermansen, J. S., Hewitt, G., Hudson, A. G., Jiggins, C., … Zinner,
D. (2013). Hybridization and Speciation. *Journal of Evolutionary Biology*, *26*(2), 229–246.
https://doi.org/10.1111/j.1420-9101.2012.02599.x

Adams, K. P., & Hamilton, R. J. (2011). Management History. In D. G. Hewitt (Ed.), *Biology and*
management of white-tailed deer. CRC Press. http://site.ebrary.com/id/11007093

Adavoudi, R., & Pilot, M. (2021). Consequences of Hybridization in Mammals: A Systematic
Review. *Genes*, *13*(1), 50. https://doi.org/10.3390/genes13010050

Aguilar, A., Roemer, G., Debenham, S., Binns, M., Garcelon, D., & Wayne, R. K. (2004). High MHC
Diversity Maintained by Balancing Selection in an otherwise Genetically Monomorphic
Mammal. *Proceedings of the National Academy of Sciences*, *101*(10), 3490–3494.
https://doi.org/10.1073/pnas.0306582101

Albrechtová, J., Albrecht, T., Baird, S. J. E., Macholán, M., Rudolfsen, G., Munclinger, P., Tucker,
P. K., & Piálek, J. (2012). Sperm-related phenotypes implicated in both maintenance
and breakdown of a natural species barrier in the house mouse. *Proceedings of the*
Royal Society B: Biological Sciences, *279*(1748), 4803–4810.
https://doi.org/10.1098/rspb.2012.1802

Alminas, O. S. V., Heffelfinger, J. R., Statham, M. J., & Latch, E. K. (2021). Phylogeography of
Cedros and Tiburón Island Mule Deer in North America's Desert Southwest. *Journal of*
Heredity, *112*(3), 260–275. https://doi.org/10.1093/jhered/esab013

Andrews S. (2010). *FastQC: a Quality Control Tool for High Throughput Sequence Data. Available*

460   *online at: Http://www.bioinformatics.babraham.ac.uk/projects/fastqc.*

461   Anthony, R. G., & Smith, N. S. (1977). Ecological Relationships between Mule Deer and White-

462   Tailed Deer in Southeastern Arizona. *Ecological Monographs*, *47*(3), 255–277.

463   https://doi.org/10.2307/1942517

464   Avise, J. C., Walker, D., & Johns, G. C. (1998). Speciation Durations and Pleistocene Effects on

465   Vertebrate Phylogeography. *Proceedings of the Royal Society of London. Series B:*

466   *Biological Sciences*, *265*(1407), 1707–1712. https://doi.org/10.1098/rspb.1998.0492

467   Barlow, A., Paijmans, J. L. A., Alberti, F., Gasparyan, B., Bar-Oz, G., Pinhasi, R., Foronova, I.,

468   Puzachenko, A. Y., Pacher, M., Dalén, L., Baryshnikov, G., & Hofreiter, M. (2021). Middle

469   Pleistocene genome calibrates a revised evolutionary history of extinct cave bears.

470   *Current Biology*, *31*(8), 1771-1779.e7. https://doi.org/10.1016/j.cub.2021.01.073

471   Bernatchez, L., & Landry, C. (2003). MHC Studies in Nonmodel Vertebrates: What Have we

472   Learned about Natural Selection in 15 Years? *Journal of Evolutionary Biology*, *16*(3),

473   363–377. https://doi.org/10.1046/j.1420-9101.2003.00531.x

474   Berry, S. L., Shipley, L. A., Long, R. A., & Loggers, C. (2019). Differences in Dietary Niche and

475   Foraging Behavior of Sympatric Mule and White-Tailed Deer. *Ecosphere*, *10*(7).

476   https://doi.org/10.1002/ecs2.2815

477   Bhattacharyya, T., Gregorova, S., Mihola, O., Anger, M., Sebestova, J., Denny, P., Simecek, P., &

478   Forejt, J. (2013). Mechanistic basis of infertility of mouse intersubspecific hybrids.

479   *Proceedings of the National Academy of Sciences*, *110*(6).

480   https://doi.org/10.1073/pnas.1219126110

Blank, D. A., & Yang, W. (2017). Mother-young Recognition in Goitered Gazelle during Hiding

Period. *Behavioural Processes*, *142*, 21–28.

https://doi.org/10.1016/j.beproc.2017.05.016

Bolger, A. M., Lohse, M., & Usadel, B. (2014). Trimmomatic: A flexible trimmer for Illumina

sequence data. *Bioinformatics*, *30*(15), 2114–2120.

https://doi.org/10.1093/bioinformatics/btu170

Booker, T. R., Yeaman, S., & Whitlock, M. C. (2021). Global Adaptation Complicates the

Interpretation of Genome Scans for Local Adaptation. *Evolution Letters*, *5*(1), 4–15.

https://doi.org/10.1002/evl3.208

Bradley, R. D., Bryant, F. C., Bradley, L. C., Haynie, M. L., & Baker, R. J. (2003). IMPLICATIONS OF

HYBRIDIZATION BETWEEN WHITE-TAILED DEER AND MULE DEER. *The Southwestern*

Naturalist, *48*(4), 7.

Broad Institute, GitHub Repository. (2019). *"Picard Toolkit."*

http://broadinstitute.github.io/picard/

Brunjes, K. J., Ballard, W. B., Humphrey, M. H., Harwell, F., Mcintyre, N. E., Krausman, P. R., &

Wallace, M. C. (2006). Habitat Use by Sympatric Mule and White-Tailed Deer in Texas.

Journal of Wildlife Management, *70*(5), 1351–1359. https://doi.org/10.2193/0022-

541X(2006)70[1351:HUBSMA]2.0.CO;2

Budd, K., Berkman, L. K., Anderson, M., Koppelman, J., & Eggert, L. S. (2018). Genetic Structure

and Recovery of White-tailed Deer in Missouri. *The Journal of Wildlife Management*,

82(8), 1598–1607. https://doi.org/10.1002/jwmg.21546

Burri, R. (2017). Interpreting Differentiation Landscapes in the Light of Long-term Linked

Selection. *Evolution Letters*, *1*(3), 118–131. https://doi.org/10.1002/evl3.14

Cambronne, A. (2013). *Deerland: America's Hunt for Ecological Balance and the Essence of*

Wildness.

http://www.freading.com/ebooks/details/r:download/MDAwMDA5LTEwOTA1NDYz

Campbell, C. R., Poelstra, J. W., & Yoder, A. D. (2018). What is Speciation Genomics? The Roles

of Ecology, Gene Flow, and Genomic Architecture in the Formation of Species.

Biological Journal of the Linnean Society, 124(4), 561–583.

https://doi.org/10.1093/biolinnean/bly063

Cann, P., Chabi, M., Delsart, A., Le Danvic, C., Saliou, J.-M., Chasles, M., Keller, M., & Nagnan-Le

Meillour, P. (2019). The Olfactory Secretome Varies According to Season in Female

Sheep and Goat. *BMC Genomics, 20*(1), 794. https://doi.org/10.1186/s12864-019-6194-z

Carr, S. M., Ballinger, S. W., Derr, J. N., Blankenship, L. H., & Bickham, J. W. (1986). Mitochondrial

DNA Analysis of Hybridization between Sympatric White-tailed Deer and Mule Deer in

West Texas. *Proceedings of the National Academy of Sciences, 83*(24), 9576–9580.

https://doi.org/10.1073/pnas.83.24.9576

Carr, S. M., & Hughes, G. A. (1993). Direction of Introgressive Hybridization between Species of

North American Deer (Odocoileus) as Inferred from Mitochondrial-Cytochrome-b

Sequences. *Journal of Mammalogy, 74*(2), 331–342. https://doi.org/10.2307/1382388

Cathey, J. C., Bickham, J. W., & Patton, J. C. (1998). Introgressive Hybridization and

Nonconcordant Evolutionary History of Maternal and Paternal Lineages in North

American Deer. *Evolution, 52*(4), 1224–1229. https://doi.org/10.1111/j.1558-

5646.1998.tb01850.x

Chafin, T. K., Zbinden, Z. D., Douglas, M. R., Martin, B. T., Middaugh, C. R., Gray, M. C., Ballard, J.

R., & Douglas, M. E. (2021). Spatial Population Genetics in Heavily Managed Species:

Separating Patterns of Historical Translocation from Contemporary Gene Flow in

White-tailed Deer. *Evolutionary Applications*, *14*(6), 1673–1689.

https://doi.org/10.1111/eva.13233

Chamaillé-Jammes, S., Malcuit, H., Le Saout, S., & Martin, J.-L. (2014). Innate Threat-sensitive

Foraging: Black-tailed Deer Remain More Fearful of Wolf than of the less Dangerous

Black Bear even after 100 years of Wolf Absence. *Oecologia*, *174*(4), 1151–1158.

https://doi.org/10.1007/s00442-013-2843-0

Cherry, M. J., & Barton, B. T. (2017). Effects of Wind on Predator-Prey Interactions. *Food Webs*,

13, 92–97. https://doi.org/10.1016/j.fooweb.2017.02.005

Colella, J. P., Lan, T., Talbot, S. L., Lindqvist, C., & Cook, J. A. (2021). Whole-genome

Resequencing Reveals Persistence of Forest-Associated Mammals in Late Pleistocene

Refugia along North America's North Pacific Coast. *Journal of Biogeography*, *48*(5),

1153–1169. https://doi.org/10.1111/jbi.14068

Combe, F. J., Jaster, L., Ricketts, A., Haukos, D., & Hope, A. G. (2021). Population Genomics of

Free-ranging Great Plains White-tailed and Mule deer Reflects a Long History of Inter-

specific Hybridization. *Evolutionary Applications*, eva.13330.

https://doi.org/10.1111/eva.13330

Cronin, M. A. (1991). Mitochondrial and Nuclear Genetic Relationships of Deer (Odocoileus

spp.) in Western North America. *Canadian Journal of Zoology*, *69*(5), 1270–1279.

https://doi.org/10.1139/z91-179

Cronin, M. A., Vyse, E. R., & Cameron, D. G. (1988). Genetic Relationships between Mule Deer and

White-Tailed Deer in Montana. *The Journal of Wildlife Management*, *52*(2), 320.

https://doi.org/10.2307/3801243

Cruickshank, T. E., & Hahn, M. W. (2014). Reanalysis Suggests that Genomic Islands of Speciation are Due to Reduced Diversity, not Reduced Gene Flow. *Molecular Ecology*, *23*(13), 3133–3157. https://doi.org/10.1111/mec.12796

Cutter, A. D. (2013). Integrating Phylogenetics, Phylogeography and Population Genetics through Genomes and Evolutionary Theory. *Molecular Phylogenetics and Evolution*, *69*(3), 1172–1185. https://doi.org/10.1016/j.ympev.2013.06.006

Darwin, C., 1809-1882. (1859). *On the Origin of Species by Means of Natural Selection, or Preservation of Favoured Races in the Struggle for Life*. London : John Murray, 1859. https://search.library.wisc.edu/catalog/9934839413602122

Dedato, M., Robert, C., Taillon, J., Shafer, A., & Cote, S. (2022). *Demographic History and Conservation Genomics of Caribou (Rangifer tarandus) in Québec* [Preprint]. Preprints. https://doi.org/10.22541/au.164165840.02956352/v1

Degl'Innocenti, A., & D'Errico, A. (2017). Regulatory Features for Odorant Receptor Genes in the Mouse Genome. *Frontiers in Genetics*, *8*, 8.

Delaneau, O., Marchini, J., & Zagury, J.-F. (2012). A linear complexity phasing method for thousands of genomes. *Nature Methods*, *9*(2), 179–181. https://doi.org/10.1038/nmeth.1785

Demarais, S., Miller, K. V., & Jacobson, H. A. (2000). White-tailed deer. In S. Demarais & P. R. Krausman (Eds.), *Ecology and management of large mammals in North America*. Prentice Hall.

Derr, J. N. (1991). Genetic Interactions between White-Tailed and Mule Deer in the Southwestern United States. *The Journal of Wildlife Management*, *55*(2), 228. https://doi.org/10.2307/3809144

Derr, J. N., Hale, D. W., Ellsworth, D. L., & Bickham, J. W. (1991). *Fertility in an F1 Male Hybrid of*

White-tailed Deer (Odocoileus virginianus) x Mule deer (O. hemionus). 7.

Deutsch, J. C., & Nefdt, R. J. C. (1992). Olfactory Cues Influence Female Choice in two Lek-

breeding Antelopes. *Nature*, *356*(6370), 596–598. https://doi.org/10.1038/356596a0

Deyoung, R. W., Demarais, S., Honeycutt, R. L., Rooney, A. P., Gonzales, R. A., & Gee, K. L. (2003).

Genetic Consequences of White-tailed Deer (Odocoileus virginianus) Restoration in

Mississippi. *Molecular Ecology*, *12*(12), 3237–3252. https://doi.org/10.1046/j.1365-

294X.2003.01996.x

Ditchkoff, S. S. (2011). Anatomy and Physiology. In D. G. Hewitt (Ed.), *Biology and management*

of white-tailed deer. CRC Press. http://site.ebrary.com/id/11007093

Douzery, E., & Randi, E. (1997). The Mitochondrial Control Region of Cervidae: Evolutionary

Patterns and Phylogenetic Content. *Molecular Biology and Evolution*, *14*(11), 1154–

1166. https://doi.org/10.1093/oxfordjournals.molbev.a025725

Durand, E. Y., Patterson, N., Reich, D., & Slatkin, M. (2011). Testing for Ancient Admixture

between Closely Related Populations. *Molecular Biology and Evolution*, *28*(8), 2239–

2252. https://doi.org/10.1093/molbev/msr048

Dussex, N., Alberti, F., Heino, M. T., Olsen, R.-A., van der Valk, T., Ryman, N., Laikre, L., Ahlgren,

H., Askeyev, I. V., Askeyev, O. V., Shaymuratova, D. N., Askeyev, A. O., Döppes, D.,

Friedrich, R., Lindauer, S., Rosendahl, W., Aspi, J., Hofreiter, M., Lidén, K., … Díez-del-

Molino, D. (2020). Moose Genomes Reveal Past Glacial Demography and the Origin of

Modern Lineages. *BMC Genomics*, *21*(1), 854. https://doi.org/10.1186/s12864-020-

07208-3

Ellsworth, D. L., Honeycutt, R. L., Silvy, N. J., Bickham, J. W., & Klimstra, W. D. (1994). Historical

Biogeography and Contemporary Patterns Of Mitochondrial DNA Variation In White-

Tailed Deer From The Southeastern United States. *Evolution*, *48*(1), 122–136.

https://doi.org/10.1111/j.1558-5646.1994.tb01299.x

Evanno, G., Regnaut, S., & Goudet, J. (2005). Detecting the Number of Clusters of Individuals

Using the Software Structure: A Simulation Study. *Molecular Ecology*, *14*(8), 2611–2620.

https://doi.org/10.1111/j.1365-294X.2005.02553.x

Ewels, P., Magnusson, M., Lundin, S., & Käller, M. (2016). MultiQC: Summarize Analysis Results

for Multiple Tools and Samples in a Single Report. *Bioinformatics*, *32*(19), 3047–3048.

https://doi.org/10.1093/bioinformatics/btw354

Feder, J. L. (2012). *The Genomics of Speciation-With-Gene-Flow*. *28*(7), 9.

Gao, F., Ming, C., Hu, W., & Li, H. (2016). New Software for the Fast Estimation of Population

Recombination Rates (FastEPRR) in the Genomic Era. *G3 Genes|Genomes|Genetics*, *6*(6),

1563–1571. https://doi.org/10.1534/g3.116.028233

Gilbert, C., Ropiquet, A., & Hassanin, A. (2006). Mitochondrial and Nuclear Phylogenies of

Cervidae (Mammalia, Ruminantia): Systematics, Morphology, and Biogeography.

Molecular Phylogenetics and Evolution, *40*(1), 101–117.

https://doi.org/10.1016/j.ympev.2006.02.017

Gower, G., Tuke, J., Rohrlach, A., Soubrier, J., Llamas, B., Bean, N., & Cooper, A. (2018).

Population Size History from Short Genomic Scaffolds: How Short is Too Short? [Preprint].

Genetics. https://doi.org/10.1101/382036

Green, R. E., Krause, J., Briggs, A. W., Maricic, T., Stenzel, U., Kircher, M., Patterson, N., Li, H., Zhai,

W., Fritz, M. H. Y., Hansen, N. F., Durand, E. Y., Malaspinas, A. S., Jensen, J. D., Marques-

Bonet, T., Alkan, C., Prufer, K., Meyer, M., Burbano, H. A., … Paabo, S. (2010). A Draft

Sequence of the Neandertal Genome. *Science*, *328*(5979), 710–722.

https://doi.org/10.1126/science.1188021

Greenslade, A. D. (1998). *Inter- and intraspecific phylogeography of North American Odocoileus*

Deer based on mitochondrial DNA sequences. Memorial University of Newfoundland.

https://research.library.mun.ca/1190/

Hewitt, G. (2000). The Genetic Legacy of the Quaternary Ice Ages. *Nature*, *405*(6789), 907–913.

https://doi.org/10.1038/35016000

Hirata, M., & Kusatake, N. (2021). Relative Importance of Senses in Forage Discrimination by

Cattle Depends on the Sensory Contrast between the Discrimination Targets: A

Preliminary Study. *Animal Cognition*, *24*(1), 99–106. https://doi.org/10.1007/s10071-

020-01422-y

Hornbeck, G. E., & Mahoney, J. M. (2000). *Introgressive Hybridization of Mule Deer and White-*

Tailed Deer in Southwestern Alberta. 5.

Hughes, G. A., & Carr, S. M. (1993). Reciprocal Hybridization between White-tailed Deer

(Odocoileus virginianus) and Mule Deer (O.hemionus) in Western Canada: Evidence

from Serum Albumin and mtDNA Sequences. *Canadian Journal of Zoology*, *71*(3), 524–

530. https://doi.org/10.1139/z93-073

Iacolina, L., Corlatti, L., Buzan, E., Safner, T., & Šprem, N. (2019). Hybridisation in European

Ungulates: An Overview of the Current Status, Causes, and Consequences. *Mammal*

Review, *49*(1), 45–59. https://doi.org/10.1111/mam.12140

Ito, T., Hayakawa, T., Suzuki–Hashido, N., Hamada, Y., Kurihara, Y., Hanya, G., Kaneko, A.,

Natsume, T., Aisu, S., Honda, T., Yachimori, S., Anezaki, T., Omi, T., Hayama, S., Tanaka,

M., Wakamori, H., Imai, H., & Kawamoto, Y. (2021). Phylogeographic History of Japanese

Macaques. *Journal of Biogeography*, jbi.14087. https://doi.org/10.1111/jbi.14087

IUCN. (2015a). *Odocoileus virginianus: Gallina, S. and Lopez Arevalo, H.: The IUCN Red List of*

Threatened Species 2016: e.T42394A22162580 [Data set]. International Union for

Conservation   of   Nature.   https://doi.org/10.2305/IUCN.UK.2016-

2.RLTS.T42394A22162580.en

IUCN. (2015b). *Odocoileus hemionus: Sanchez Rojas, G. and Gallina Tessaro, S.: The IUCN Red*

List of Threatened Species 2016: e.T42393A22162113 [Data set]. International Union for

Conservation   of   Nature.   https://doi.org/10.2305/IUCN.UK.2016-

1.RLTS.T42393A22162113.en

Jensen-Seaman, M. I., Furey, T. S., Payseur, B. A., Lu, Y., Roskin, K. M., Chen, C.-F., Thomas, M. A.,

Haussler, D., & Jacob, H. J. (2004). Comparative Recombination Rates in the Rat, Mouse,

and   Human   Genomes.   *Genome   Research*,   *14*(4),   528–538.

https://doi.org/10.1101/gr.1970304

Keller, M., & Lévy, F. (2012). The Main but not the Accessory Olfactory System is Involved in the

Processing of Socially Relevant Chemosignals in Ungulates. *Frontiers in Neuroanatomy*,

6. https://doi.org/10.3389/fnana.2012.00039

Kinoshita, E., Kosintsev, P. A., Abramov, A. V., Solovyev, V. A., Saveljev, A. P., Nishita, Y., &

Masuda, R. (2020). Holocene Changes in the Distributions of Asian and European

Badgers (Carnivora: Mustelidae: Meles) Inferred from Ancient DNA Analysis. *Biological*

Journal   of   the   Linnean   Society,   *129*(3),   594–602.

https://doi.org/10.1093/biolinnean/blaa007

Kirch, M., Romundset, A., Gilbert, M. T. P., Jones, F. C., & Foote, A. D. (2021). Ancient and modern

stickleback genomes reveal the demographic constraints on adaptation. *Current*

Biology, S0960982221002682. https://doi.org/10.1016/j.cub.2021.02.027

Kopelman, N. M., Mayzel, J., Jakobsson, M., Rosenberg, N. A., & Mayrose, I. (2015). Clumpak: A

Program for Identifying Clustering Modes and Packaging Population Structure

Inferences Across K. *Molecular Ecology Resources*, *15*(5), 1179–1191.

https://doi.org/10.1111/1755-0998.12387

Korneliussen, T. S., Albrechtsen, A., & Nielsen, R. (2014). ANGSD: Analysis of Next Generation

Sequencing Data. *BMC Bioinformatics*, *15*(1), 356. https://doi.org/10.1186/s12859-014-

0356-4

Kuijper, D. P. J., Verwijmeren, M., Churski, M., Zbyryt, A., Schmidt, K., Jędrzejewska, B., & Smit,

C. (2014). What Cues Do Ungulates Use to Assess Predation Risk in Dense Temperate

Forests? *PLoS ONE*, *9*(1), e84607. https://doi.org/10.1371/journal.pone.0084607

Latch, E. K., Heffelfinger, J. R., Fike, J. A., & RHODES Jr, O. E. (2009). Species-wide

Phylogeography of North American Mule Deer (Odocoileus hemionus): Cryptic Glacial

Refugia and Postglacial Recolonization. *Molecular Ecology*, *18*(8), 1730–1745.

https://doi.org/10.1111/j.1365-294X.2009.04153.x

Latch, E. K., Reding, D. M., Heffelfinger, J. R., Alcalá-Galván, C. H., & Rhodes, O. E. (2014). Range-

wide Analysis of Genetic Structure in a Widespread, Highly Mobile Species (Odocoileus

hemionus) Reveals the Importance of Historical Biogeography. *Molecular Ecology*,

23(13), 3171–3190. https://doi.org/10.1111/mec.12803

Lavretsky, P., DaCosta, J. M., Sorenson, M. D., McCracken, K. G., & Peters, J. L. (2019). DdRAD-

seq Data Reveal Significant Genome-Wide Population Structure and divergent

Genomic Regions that Distinguish the Mallard and Close Relatives in North America.

Molecular Ecology, *28*(10), 2594–2609. https://doi.org/10.1111/mec.15091

Li, H., & Durbin, R. (2009). Fast and Accurate Short Read Alignment with Burrows-Wheeler

Transform. *Bioinformatics*, *25*(14), 1754–1760.

https://doi.org/10.1093/bioinformatics/btp324

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G., Durbin,

R., & 1000 Genome Project Data Processing Subgroup. (2009). The Sequence

Alignment/Map format and SAMtools. *Bioinformatics*, *25*(16), 2078–2079.

https://doi.org/10.1093/bioinformatics/btp352

Lingle, S. (1992). Escape Gaits of White-Tailed Deer, Mule Deer and Their Hybrids: Gaits

Observed and Patterns of Limb Coordination. *Behaviour*, *122*(3–4), 153–181.

https://doi.org/10.1163/156853992X00499

Lingle, S. (2001). Anti-Predator Strategies and Grouping Patterns in White-Tailed Deer and Mule

Deer. *Ethology*, *107*(4), 295–314. https://doi.org/10.1046/j.1439-0310.2001.00664.x

Lingle, S. (2003). Group Composition and Cohesion in Sympatric White-tailed Deer and Mule

deer. *Canadian Journal of Zoology*, *81*(7), 1119–1130. https://doi.org/10.1139/z03-097

Liu, Q., Mishra, M., Saxena, A. S., Wu, H., Qiu, Y., Zhang, X., You, X., Ding, S., & Miyamoto, M. M.

(2021). Balancing Selection Maintains Ancient Polymorphisms at Conserved Enhancers

for the Olfactory Receptor Genes of a Chinese Marine Fish. *Molecular Ecology*, *30*(16),

4023–4038. https://doi.org/10.1111/mec.16016

Liu, S., Westbury, M. V., Dussex, N., Mitchell, K. J., Sinding, M.-H. S., Heintzman, P. D., Duchêne,

D. A., Kapp, J. D., von Seth, J., Heiniger, H., Sánchez-Barreiro, F., Margaryan, A., André-

Olsen, R., De Cahsan, B., Meng, G., Yang, C., Chen, L., van der Valk, T., Moodley, Y., …

709 Gilbert, M. T. P. (2021). Ancient and modern genomes unravel the evolutionary history

710 of the rhinoceros family. *Cell*, *184*(19), 4874-4885.e16.

711 https://doi.org/10.1016/j.cell.2021.07.032

712 Mallet, J. (2005). Hybridization as an Invasion of the Genome. *Trends in Ecology & Evolution*,

713 *20*(5), 229–237. https://doi.org/10.1016/j.tree.2005.02.010

714 Marques, D. A., Lucek, K., Sousa, V. C., Excoffier, L., & Seehausen, O. (2019). Admixture Between

715 Old Lineages Facilitated Contemporary Ecological Speciation in Lake Constance

716 Stickleback. *Nature Communications*, *10*(1), 4240. https://doi.org/10.1038/s41467-019-

717 12182-w

718 Martin, M., Patterson, M., Garg, S., O Fischer, S., Pisanti, N., Klau, G. W., Schöenhuth, A., &

719 Marschall, T. (2016). *WhatsHap: Fast and Accurate Read-based Phasing* [Preprint].

720 Bioinformatics. https://doi.org/10.1101/085050

721 Martin, S. (2021). *Genomics_general*. GitHub.

722 https://github.com/simonhmartin/genomics_general

723 Martin, S., Davey, J. W., & Jiggins, C. D. (2015). Evaluating the Use of ABBA–BABA Statistics to

724 Locate Introgressed Loci. *Molecular Biology and Evolution*, *32*(1), 244–257.

725 https://doi.org/10.1093/molbev/msu269

726 Mayr, E. (1942). *Systematics and the Origin of Species*.

727 McDonald, J. S., Miller, K. V., Association, Q. D. M., & Resources, D. B. W. S. of F. (2004). *A History*

728 *of White-tailed Deer Restocking in the United States, 1878 to 2004*. Quality Deer

729 Management Assoc. https://books.google.ca/books?id=fv_oGwAACAAJ

730 McKenna, A., Hanna, M., Banks, E., Sivachenko, A., Cibulskis, K., Kernytsky, A., Garimella, K.,

731 Altshuler, D., Gabriel, S., Daly, M., & DePristo, M. A. (2010). The Genome Analysis Toolkit:

A MapReduce framework for analyzing next-generation DNA sequencing data. *Genome*

Research, *20*(9), 1297–1303. https://doi.org/10.1101/gr.107524.110

Meisner, J., & Albrechtsen, A. (2018). Inferring Population Structure and Admixture Proportions

in Low-Depth NGS Data. *Genetics*, *210*(2), 719–731.

https://doi.org/10.1534/genetics.118.301336

Momigliano, P., Jokinen, H., Fraimout, A., Florin, A.-B., Norkko, A., & Merilä, J. (2017).

Extraordinarily Rapid Speciation in a Marine Fish. 6.

Morina, D. L., Demarais, S., Strickland, B. K., & Larson, J. E. (2018). While Males Fight, Females

Choose: Male Phenotypic Quality Informs Female Mate Choice in Mammals. *Animal*

Behaviour, *138*, 69–74. https://doi.org/10.1016/j.anbehav.2018.02.004

NatureServe. (2021a). *NatureServe Explorer* [Web application]. NatureServe Mule Deer.

https://explorer.natureserve.org/Taxon/ELEMENT_GLOBAL.2.101365/Odocoileus_he

mionus

NatureServe. (2021b). *NatureServe Explorer [web application] WTD*. NatureServe, Arlington,

Virginia.

https://explorer.natureserve.org/Taxon/ELEMENT_GLOBAL.2.104777/Odocoileus_virg

inianus

Nosil, P. (2012). *Ecological Speciation*. Oxford University Press.

Paijmans, J. L. A., Barlow, A., Becker, M. S., Cahill, J. A., Fickel, J., Förster, D. W. G., Gries, K.,

Hartmann, S., Havmøller, R. W., Henneberger, K., Kern, C., Kitchener, A. C., Lorenzen, E.

D., Mayer, F., OBrien, S. J., von Seth, J., Sinding, M.-H. S., Spong, G., Uphyrkina, O., …

Hofreiter, M. (2021). African and Asian leopards are highly differentiated at the genomic

level. *Current Biology*, S0960982221004577. https://doi.org/10.1016/j.cub.2021.03.084

Pedersen, B. S., & Quinlan, A. R. (2018). Mosdepth: Quick Coverage Calculation for Genomes

and Exomes. *Bioinformatics*, *34*(5), 867–868.

https://doi.org/10.1093/bioinformatics/btx699

Peres, T. M., & Altman, H. (2018). The Magic of Improbable Appendages: Deer Antler Objects in

the Archaeological Record of the American South. *Journal of Archaeological Science:*

Reports, *20*, 888–895. https://doi.org/10.1016/j.jasrep.2017.10.028

Pickrell, J. K., & Pritchard, J. K. (2012). Inference of Population Splits and Mixtures from

Genome-Wide Allele Frequency Data. *PLoS Genetics*, *8*(11), e1002967.

https://doi.org/10.1371/journal.pgen.1002967

Pierini, F., & Lenz, T. L. (2018). Divergent Allele Advantage at Human MHC Genes: Signatures of

Past and Ongoing Selection. *Molecular Biology and Evolution*, *35*(9), 2145–2158.

https://doi.org/10.1093/molbev/msy116

Piertney, S. B., & Oliver, M. K. (2006). The Evolutionary Ecology of the Major Histocompatibility

Complex. *Heredity*, *96*(1), 7–21. https://doi.org/10.1038/sj.hdy.6800724

Pitra, C., Fickel, J., Meijaard, E., & Groves, C. (2004). Evolution and Phylogeny of Old World Deer.

Molecular Phylogenetics and Evolution, *33*(3), 880–895.

https://doi.org/10.1016/j.ympev.2004.07.013

Poelstra, J. W., Vijay, N., Bossu, C. M., Lantz, H., Ryll, B., Muller, I., Baglione, V., Unneberg, P.,

Wikelski, M., Grabherr, M. G., & Wolf, J. B. W. (2014). The Genomic Landscape Underlying

Phenotypic Integrity in the Face of Gene Flow in Crows. *Science*, *344*(6190), 1410–1414.

https://doi.org/10.1126/science.1253226

Pundir, S., Martin, M. J., O'Donovan, C., & The UniProt Consortium. (2016). UniProt Tools.

Current Protocols in Bioinformatics, *53*(1).

https://doi.org/10.1002/0471250953.bi0129s53

Quinlan, A. R., & Hall, I. M. (2010). BEDTools: A Flexible Suite of Utilities for Comparing Genomic

Features. *Bioinformatics*, *26*(6), 841–842.

https://doi.org/10.1093/bioinformatics/btq033

R Core Team. (2021). *R: A Language and Environment for Statistical Computing*. R Foundation

for Statistical Computing. https://www.R-project.org/

Ragavan, P., Zhou, R., Ng, W. L., Rana, T. S., Mageswaran, T., Mohan, P. M., & Saxena, A. (2017).

Natural Hybridization in Mangroves – an Overview. *Botanical Journal of the Linnean*

Society, *185*(2), 208–224. https://doi.org/10.1093/botlinnean/box053

Ranslow, A. N., Richter, J. P., Neuberger, T., Van Valkenburgh, B., Rumple, C. R., Quigley, A. P.,

Pang, B., Krane, M. H., & Craven, B. A. (2014). Reconstruction and Morphometric Analysis

of the Nasal Airway of the White-Tailed Deer (Odocoileus virginianus) and Implications

Regarding Respiratory and Olfactory Airflow: White-Tailed Deer Nasal Anatomy and

Function. *The Anatomical Record*, *297*(11), 2138–2147. https://doi.org/10.1002/ar.23037

Ravinet, M., Faria, R., Butlin, R. K., Galindo, J., Bierne, N., Rafajlović, M., Noor, M. A. F., Mehlig, B.,

& Westram, A. M. (2017). Interpreting the Genomic Landscape of Speciation: A Road Map

for Finding Barriers to Gene Flow. *Journal of Evolutionary Biology*, *30*(8), 1450–1477.

https://doi.org/10.1111/jeb.13047

Rundle, H. D., & Nosil, P. (2005). Ecological Speciation. *Ecology Letters*, *8*(3), 336–352.

https://doi.org/10.1111/j.1461-0248.2004.00715.x

Russell, T., Cullingham, C., Ball, M., Pybus, M., & Coltman, D. (2021). Extent and Direction of

Introgressive Hybridization of Mule and White-tailed Deer in Western Canada.

Evolutionary Applications, eva.13250. https://doi.org/10.1111/eva.13250

Russell, T., Cullingham, C., Kommadath, A., Stothard, P., Herbst, A., & Coltman, D. (2019).

Development of a Novel Mule Deer Genomic Assembly and Species-Diagnostic SNP

Panel for Assessing Introgression in Mule Deer, White-Tailed Deer, and Their

Interspecific Hybrids. *G3: Genes|Genomes|Genetics*, g3.200838.2018.

https://doi.org/10.1534/g3.118.200838

Schiffels, S., & Durbin, R. (2014). Inferring Human Population Size and Separation History from

Multiple Genome Sequences. *Nature Genetics*, *46*(8), 919–925.

https://doi.org/10.1038/ng.3015

Schiffels, S., & Wang, K. (2020). MSMC and MSMC2: The Multiple Sequentially Markovian

Coalescent. In J. Y. Dutheil (Ed.), *Statistical Population Genomics* (Vol. 2090, pp. 147–

166). Springer US. https://doi.org/10.1007/978-1-0716-0199-0_7

Schumer, M., Cui, R., Rosenthal, G. G., & Andolfatto, P. (2015). Reproductive Isolation of Hybrid

Populations Driven by Genetic Incompatibilities. *PLOS Genetics*, *11*(3), e1005041.

https://doi.org/10.1371/journal.pgen.1005041

Servedio, M. R. (2004). The Evolution of Premating Isolation: Local Adaptation and Natural and

Sexual Selection Against Hybrids. *Evolution*, *58*(5), 913–924.

https://doi.org/10.1111/j.0014-3820.2004.tb00425.x

Shafer, A. B. A., Cullingham, C. I., Côté, S. D., & Coltman, D. W. (2010). Of Glaciers and Refugia: A

Decade of Study Sheds New Light on the Phylogeography of Northwestern North

America. *Molecular Ecology*, *19*(21), 4589–4621. https://doi.org/10.1111/j.1365-

294X.2010.04828.x

Shafer, A. B. A., & Wolf, J. B. W. (2013). Widespread Evidence for Incipient Ecological Speciation:

A Meta-analysis of Isolation-by-ecology. *Ecology Letters*, *16*(7), 940–950.

https://doi.org/10.1111/ele.12120

Shang, H., Rendón-Anaya, M., Paun, O., Field, D. L., Hess, J., Vogl, C., Liu, J., Ingvarsson, P. K.,

Lexer, C., & Leroy, T. (2021). *Conserved Genomic Landscapes of Differentiation Across*

Populus Speciation Continuum [Preprint]. Evolutionary Biology.

https://doi.org/10.1101/2021.08.26.457771

Skotte, L., Korneliussen, T. S., & Albrechtsen, A. (2013). Estimating Individual Admixture

Proportions from Next Generation Sequencing Data. *Genetics*, *195*(3), 693–702.

https://doi.org/10.1534/genetics.113.154138

Soraggi, S., Wiuf, C., & Albrechtsen, A. (2018). Powerful Inference with the D-statistic on Low-

coverage Whole-genome Data. *G3: Genes, Genomes, Genetics*, *8*(2).

https://doi.org/10.1534/g3.117.300192

Stankowski, S., & Ravinet, M. (2021). Defining the Speciation Continuum. *Evolution*, *75*(6), 1256–

1273. https://doi.org/10.1111/evo.14215

Stubblefield, S. S., Warren, R. J., & Murphy, B. R. (1986). Hybridization of Free-Ranging White-

Tailed and Mule Deer in Texas. *The Journal of Wildlife Management*, *50*(4), 688.

https://doi.org/10.2307/3800982

Tarasov, A., Vilella, A. J., Cuppen, E., Nijman, I. J., & Prins, P. (2015). Sambamba: Fast processing

of NGS alignment formats. *Bioinformatics*, *31*(12), 2032–2034.

https://doi.org/10.1093/bioinformatics/btv098

Taylor, S. A., & Larson, E. L. (2019). Insights from Genomes into the Evolutionary Importance
and Prevalence of Hybridization in Nature. *Nature Ecology & Evolution*, *3*(2), 170–177.
https://doi.org/10.1038/s41559-018-0777-y

UniProt.   (2022a).   *Q9Y5E5   ·   PCDB4_HUMAN*.   UniProt   Beta.
https://beta.uniprot.org/uniprotkb/Q9Y5E5/entry

UniProt.   (2022b).   *Q15057   ·   ACAP2_HUMAN*.   UniProt   Beta.
https://beta.uniprot.org/uniprotkb/Q15057/entry

Wang, K., Mathieson, I., O'Connell, J., & Schiffels, S. (2020). Tracking Human Population
Structure through Time from Whole Genome Sequences. *PLOS Genetics*, *16*(3),
e1008552. https://doi.org/10.1371/journal.pgen.1008552

Whittaker, D. G., & Lindzey, F. G. (2004). Habitat Use Patterns of Sympatric Deer Species on
Rocky Mountain Arsenal, Colorado. *Wildlife Society Bulletin*, *32*(4), 1114–1123.
https://doi.org/10.2193/0091-7648(2004)032[1114:HUPOSD]2.0.CO;2

Wikenros, C., Kuijper, D. P. J., Behnke, R., & Schmidt, K. (2015). Behavioural Responses of
Ungulates to Indirect Cues of an Ambush Predator. *Behaviour*, *152*(7–8), 1019–1040.
https://doi.org/10.1163/1568539X-00003266

Wishart, W. D. (1980). Hybrids of White-Tailed and Mule Deer in Alberta. *Journal of Mammalogy,*
61(4), 716–719. https://doi.org/10.2307/1380318

Wishart, W. D., Hrudka, F., Schmutz, S. M., & Flood, P. F. (1988). Observations on
Spermatogenesis, Sperm Phenotype, and Fertility in White-tailed × Mule Deer Hybrids
and a Yak × Cow Hybrid. *Canadian Journal of Zoology*, *66*(7), 1664–1671.
https://doi.org/10.1139/z88-240

Wright, E. A., Roberts, E. K., Platt, R. N., Bayouth, J. V., Conway, W. C., & Bradley, R. D. (2022).

Mitochondrial capture and subsequent genetic divergence generates a novel

haplogroup: Evidence from ancient and ongoing hybridization in mule and white-tailed

deer. *Journal of Mammalogy*. https://doi.org/10.1093/jmammal/gyab156

Zagata, M. D., & Haugen, A. O. (1974). Influence of Light and Weather on Observability of Iowa

Deer. *The Journal of Wildlife Management*, *38*(2), 220. https://doi.org/10.2307/3800727

Zhang, P., Huang, K., Zhang, B., Dunn, D. W., Chen, D., Li, F., Qi, X., Guo, S., & Li, B. (2018). High

Polymorphism in MHC-DRB Genes in Golden Snub-nosed Monkeys Reveals Balancing

Selection in Small, Isolated Populations. *BMC Evolutionary Biology*, *18*(1), 29.

https://doi.org/10.1186/s12862-018-1148-7

875
876
877 *Data Accessibility and Benefit-Sharing*

Data Accessibility

Raw reads for the 28 deer individuals were deposited on the NCBI Sequence Read Archive

(Accession number PRJNA830519). All scripts are available on GitLab

(https://gitlab.com/WiDGeT_TrentU/graduate_theses/-/tree/master/Kessler/CH_01).

Benefit-Sharing

The contributions of tissue samples to the research are described in the

ACKNOWLEDGEMENTS, benefits from this research results in data accessible on public

databases as described above.

886 *Author contributions*

CK and ABAS conceived the study, CK performed bioinformatic analyses with contribution

from ABAS, CK and EW performed the molecular laboratory work, CK and ABAS wrote the

manuscript. All authors commented on the text.

Speciation without gene-flow in hybridising deer

Camille Kessler[1], Eric Wootton[2], Aaron B.A. Shafer[1]

[1]Environmental and Life Sciences Graduate Program, Trent University, 7K9J 7B8 Peterborough,

Ontario, Canada

[2]Biochemistry & Molecular Biology, Trent University, K9J 7B8 Peterborough, Ontario, Canada

Corresponding authors:

Camille Kessler: camillekessler@trentu.ca

Aaron Shafer: aaronshafer@trentu.ca

Figure 1: Deer individuals in the study. (A) Sampling locations, colours represent the different groups: WTD allopatry (WTDa, blue), WTD sympatry (WTDs, violet), MD sympatry (MDs, green), MD allopatry (MDa, orange). Areas of sympatry are coloured in dark grey and areas of allopatry in light grey, from the IUCN range data for both species (IUCN, 2015a, 2015b). (B) Principal component analysis performed in PCAngsd. (C) Admixture proportions for K = 2 calculated in NGSAdmix.

Figure 2. MSMC-IM analysis of white-tailed deer (Ov) and mule deer (Oh) from Ontario (ON) and Washington (WA). (A) Migration rates over time where dashed line indicates the time point where 50% of ancestry has merged. Blue shading indicates 99% percentile of the cumulative migration probability. (B) Cumulative migration probabilities $M(t)$. Dashed lines indicate the relative cross coalescence rate obtained from MSMC2.

909
910

Figure 3: Introgression analyses. (A) ABBA-BABA analysis between individuals, individuals
involved in gene flow are coloured depending on the species (WTD in blue, MD in orange), the
colour gradient represents Z score, excess of ABBA depicted as points, excess of BABA shown
as triangles, significant tests highlighted with an *(Z-score > |3|). (B) Maximum likelihood tree
inferred by Treemix for 0 migration (right) and the variance explained by each migration event
(left).

Scenarios: ● Allopatric selection ● Background selection ● Balancing selection

Figure 4: Distribution of windows presenting pattern of selection across the genome for d_{xy} (top), F_{ST} (middle) and π (bottom). Windows following a pattern of selection are coloured according to its corresponding evolutionary scenario, grey windows do not exhibit pattern of selection. The continuous and dashed line represent the mean and median respectively.

Figure 5: Correlation coefficients over divergence time for π - ρ (dark blue) F_{ST} - ρ (dark green) and ρ - ρ (yellow), trend lines represent a linear model. The speciation continuum is represented by six comparisons of our four groups).

933
934
935
936

Table 1: Deer sample information

Genome_ID	Species	Location	Sex	Group
Ov_ON3	*O_virginianus*	Ontario	F	WTD_allopatry
Ov_ON5	*O_virginianus*	Ontario	M	WTD_allopatry
Ov_ON6	*O_virginianus*	Ontario	M	WTD_allopatry
Ov_PA3	*O_virginianus*	Pennsylvania	F	WTD_allopatry
Ov_VA3	*O_virginianus*	Virginia	F	WTD_allopatry
Ov_AL3	*O_virginianus*	Alabama	Unk	WTD_allopatry
Ov_SC1	*O_virginianus*	South Carolina	F	WTD_allopatry
Ov_AB1	*O_virginianus*	Alberta	F	WTD_sympatry
Ov_SK2	*O_virginianus*	Saskatchewan	M	WTD_sympatry
Ov_SK3	*O_virginianus*	Saskatchewan	M	WTD_sympatry
Ov_SD1	*O_virginianus*	South Dakota	M	WTD_sympatry
Ov_KS1	*O_virginianus*	Kansas	F	WTD_sympatry
Ov_MT1	*O_virginianus*	Montana	M	WTD_sympatry
Ov_MX2	*O_virginianus*	Mexico	M	WTD_sympatry
Oh_NV2	*O_hemionus*	Nevada	M	MD_allopatry
Oh_CA3	*O_hemionus*	California	Unk	MD_allopatry
Oh_BC6	*O_hemionus*	British Columbia	M	MD_allopatry
Oh_BC8	*O_hemionus*	British Columbia	M	MD_allopatry
Oh_BC1	*O_hemionus*	British Columbia	F	MD_allopatry
Oh_AK1	*O_hemionus*	Alaska	Unk	MD_allopatry
Oh_WA1	*O_hemionus*	Washington	M	MD_allopatry
Oh_AB1	*O_hemionus*	Alberta	F	MD_sympatry
Oh_UT2	*O_hemionus*	Utah	M	MD_sympatry
Oh_CO1	*O_hemionus*	Colorado	F	MD_sympatry
Oh_SK5	*O_hemionus*	Saskatchewan	Unk	MD_sympatry
Oh_OR2	*O_hemionus*	Oregon	Unk	MD_sympatry
Oh_SD2	*O_hemionus*	South Dakota	F	MD_sympatry
Oh_NM2	*O_hemionus*	New Mexico	M	MD_sympatry

941   **Table 2:** Four different evolutionary scenario and their thresholds for F_{st}, π and D_{xy} percentiles
942   applied in the genome scans analysis.

Scenario	F_{st}	π	D_{xy}
Divergence with gene flow (DwGF)	> 0.95	< 0.05	> 0.95
Allopatric selection (AS)	> 0.95	< 0.05	Between 0.45 & 0.55
Background selection (BGS)	> 0.95	< 0.05	< 0.05
Balancing selection (BLS)	< 0.05	> 0.95	> 0.95