Evolutionary investigations of the biosynthetic diversity in the skin microbiome using *lsa*BGC

Authors

Rauf Salamzade[1], J.Z. Alex Cheong[1], Shelby Sandstrom[1], Mary Hannah Swaney[1], Reed M. Stubbendieck[2], Nicole Lane Starr[1], Cameron R. Currie[2], Anne Marie Singh[3], and Lindsay R. Kalan[1,4,*]

Affiliations

[1]Department of Medical Microbiology and Immunology, School of Medicine and Public Health, University of Wisconsin, Madison, WI, USA

[2]Department of Bacteriology, University of Wisconsin—Madison, Madison, Wisconsin, USA

[3]Department of Pediatrics, School of Medicine and Public Health, University of Wisconsin, Madison, WI, USA

[4]Department of Medicine, School of Medicine and Public Health, University of Wisconsin, Madison, WI, USA

[*]Corresponding author. Email: lkalan@wisc.edu

Abstract

We developed *lsa*BGC, a bioinformatics suite that introduces several new methods to expand on the available infrastructure for genomic and metagenomic-based comparative and evolutionary investigation of biosynthetic gene clusters (BGCs). Through application of the suite to four genera commonly found in skin microbiomes, we uncover multiple novel findings on the evolution and diversity of their BGCs. We show that the virulence associated carotenoid staphyloxanthin in *Staphylococcus aureus* is ubiquitous across the *Staphylococcus* genus but has largely been lost in the skin-commensal species *Staphylococcus epidermidis*. We further identify thousands of novel single nucleotide variants (SNVs) within BGCs from the *Corynebacterium tuberculostearicum* sp. complex, which we describe here to be a narrow, multi-species clade that features the most prevalent *Corynebacterium* in healthy skin microbiomes. Although novel SNVs were approximately ten times as likely to correspond to synonymous changes when located in the top five percentile of conserved sites, *lsa*BGC identified SNVs which defied this trend and are predicted to underlie amino acid changes within functionally key enzymatic domains. Ultimately, beyond supporting evolutionary investigations, *lsa*BGC provides important functionalities to aid efforts for the discovery or synthesis of natural products.

Introduction

The secondary metabolome of bacteria has served as a valuable reservoir of natural products with great societal benefit[1]. Historically, many commercialized secondary metabolites, such as antibiotics, have been identified from particular taxonomic groups, such as the genus of *Streptomyces* within Actinomycetota[1–3]. Several studies have explored the secondary metabolome of these metabolically rich microbial taxa to develop a comprehensive understanding of their chemical diversity and unique traits[4–7]. Most of these studies begin by first identifying biosynthetic gene clusters within genomes using the popular antiSMASH software[8] and then group BGCs into gene cluster families (GCFs) using BiG-SCAPE[9]. This approach, similar to other bioinformatics software for genomics annotation and analysis of BGCs[10–12], uses a set of key protein domains as their base to group BGCs predicted to encode similar metabolites. However, when working in the context of a single genus or species, it becomes more appropriate to use full protein sequences which enables the use of advanced algorithms for the identification of orthologs and paralogs[13] to reliably determine evolutionary relationships between BGCs.

Recent studies which explored the vertical descent and conservation of BGCs across the genus of *Salinospora* and the fungal species *Aspergillus flavus* have shown that intra-genus or intra-species evolution can result in chemical and regulatory diversification of the encoded metabolites[6,14]. Thus, improved understanding of the evolutionary emergence of BGCs could be informative towards elucidating the function of individual genes and encoded metabolites[9,15–20].

Due to difficulties in the cultivation of many bacterial taxa, several recent studies have begun to explore metagenomic datasets to unearth novel secondary metabolites[21–23]. While most of these endeavors have relied on performing metagenomic assembly, recently, two read-based approaches were described to reliably and sensitively search for key BGC domains in metagenomes[22,24]. This highlights a larger trend in which these approaches primarily aim to identify highly novel secondary metabolites and thus prioritize investigation of poorly studied or newly discovered taxa[25]. However, provided that most metabolites have been mined from select genera, mostly within the *Actinomycetia* class[2], there could be tremendous reward for efforts to advance our understanding of cataloged but uncharacterized BGCs from well-studied taxa. This is especially important because many species within the metabolically rich *Streptomyces* only have limited representative genomes, thus methods to leverage metagenomic datasets and identify homologous instances of rare BGCs can lead to critical insight into their function and ecological distribution[4].

To aid the advancement of such taxa-specific analyses of BGCs, we introduce the comprehensive software suite *Lineage Specific Analysis* of BGCs (*lsa*BGC; https://github.com/Kalan-Lab/lsaBGC), which consists of several programs for comparative genetics of BGCs as well as metagenomic mining. We showcase the utility of *lsa*BGC in this study through application to four major genera which are commonly found in healthy human skin microbiomes. Several secondary metabolites which function as antibiotics[26–28], virulence factors[29], or in microbe-host interactions[30,31] have been identified in bacteria from skin microbiomes, primarily from staphylococci. Human skin is also easily accessible to sampling and most species can be cultivated for follow up studies. Using *lsa*BGC, we reveal new ecological and evolutionary insights for both well characterized and unknown predicted metabolites, including signatures of inter-species transfer for the virulence-associated staphylococcal carotenoid staphyloxanthin and a highly conserved predicted siderophore found

81 encoded by divergent *Corynebacterium* species. In addition, through mining skin metagenomic
82 datasets, we find novel coding variants within the functionally important ketide synthase (KS)
83 domain of the polyketide synthase (PKS) responsible for mycolic acid biosynthesis, a defining
84 cell wall component of *Corynebacterium* species implicated in immune recognition.
85
86

New Approaches

Comprehensive studies of the BGCs for specific taxa have become commonplace in the last five
years owing to advances and standardization in methods to cluster BGCs into GCFs[9,12]. Deeper
investigations into individual GCFs, however, remains a largely manual process. Here we
introduce *lsa*BGC which simplifies such investigations with easy-to-use workflows that require
minimal knowledge of the command-line. *lsa*BGC features a comprehensive set of
functionalities to cluster BGCs into GCFs, compute evolutionary statistics, and perform
metagenomic mining. Finalized reports include a spreadsheet which lists all genes found in
GCFs together with annotation information, conservation, and evolutionary statistics to allow
quick assessment of a taxa's biosynthetic potential. Additionally, visualizations are automatically
generated including a view of GCFs across a species phylogeny, which is commonly reported in
recent studies[6,7,14] but for which no software currently exists to automate the process. Finally,
through utilization of the metagenomic mining functionality in *lsa*BGC, users can obtain a report
on whether metagenomes contain novel alleles of known BGCs found in available assemblies.
101
102

Results

*lsa*BGC allows for sensitive and systematic identification of BGC homolog groups

The *lsa*BGC suite consists of eight core programs, four workflows, and several additional
scripts for analyses (Fig. 1, S1; Supplementary Text). These programs allow for clustering BGCs
into GCFs, refining boundaries of BGCs, and sensitive, yet specific, rapid detection of GCF
instances in draft-quality assemblies. The suite additionally includes dedicated tools for GCF
visualization, evolutionary and population genetic analysis of individual genes within GCFs, and
base-resolution identification of novel SNVs in complex metagenomes.

To demonstrate the capabilities of the *lsa*BGC suite we applied it to four genera which
are stable and integral members of the human skin microbiome[32–36]. These included three genera
in the Actinomycetota (formerly Actinobacteria) phylum: *Corynebacterium*, *Cutibacterium*, and
Micrococcus as well as the Bacillota (formerly Firmicutes) genus *Staphylococcus*. *lsa*BGC was
also independently applied to the species or species-complex from each genus most commonly
represented in skin microbiomes. These were the *C. tuberculostearicum* species-complex,
Cutibacterium acnes, *Micrococcus luteus*, and *S. epidermidis* (Table S1).

We used *lsa*BGC-AutoProcess to run gene-calling, BGC annotation, and *de novo*
homolog group delineation with complete or chromosomally complete genomes for each genus
or species. We then applied *lsa*BGC-Cluster to group homologous BGCs into GCFs and
systematically searched for instances of GCFs using *lsa*BGC-AutoExpansion (Fig. 1b; Table S2,
S3; Supplementary Text). To assess the validity and sensitivity of our approach, two
benchmarking tests using BGCs identified in *M. luteus* were performed (Fig. 1c, S3;

Supplementary Text) showing that the *lsa*BGC-AutoExpansion framework leads to greater
sensitivity compared to running *de novo* antiSMASH on fragmented draft-quality assemblies.
This increased sensitivity was due to more robust, yet reliable, detection of auxiliary genes
located on the edges of scaffolds not containing core domains used by antiSMASH for BGC
detection. In addition, *lsa*BGC-AutoExpansion is highly efficient and allowed us to search for 63
GCFs identified from high-quality *Staphylococcus* genomes in 14,978 draft-quality genomes in
approximately 14 hours using 40 cores and less than 150 Gb of memory. We found a positive
association between detection of complete BGC instances and assembly quality (Fig. 1d;
Supplementary Text).

lsaBGC enables reliable comparative genomics to explore evolutionary trends of BGCs

The increased sensitivity of *lsa*BGC to uniformly detect auxiliary components of BGCs
in each genome enabled us to perform reliable comparative genetic analyses across GCF
instances. We focused on two of the four genera with the greatest breadth of diversity,
Corynebacterium and *Staphylococcus*, and selected a subset of 456 and 229 dereplicated,
representative genomes from each genera, respectively (Table S1). To better understand which
species in these genera were most relevant to skin health, we first profiled how often distinct
species were found among healthy skin microbiomes using a large, recently generated
metagenomics dataset[37,38] (Table S4). Skin-associated species (found in more than 10 of 270
metagenomes) were phylogenetically clustered for both genera, within the *S. epidermidis/aureus*
clade (Fig. S4a) and within the *C. tuberculostearicum* species complex (Fig. S4b; Supplementary
Text).

Association of individual homolog groups from these GCFs revealed several were
enriched within skin-associated staphylococci, but only two were enriched for skin-associated
corynebacteria, both encoding hypothetical proteins (Table S5). Nearly half of the homolog
groups enriched for skin-associated staphylococci (5 of 12; 41.7%) were related to GCF-10,
predicted to encode for an NRP synthetase corresponding to pyrazinone biosynthesis, molecules
reported to regulate virulence in *S. aureus*[29,39]. Staphylococcal pyrazinones were previously
identified as being a self-contained BGC consisting of two genes, *pznA* and *pznB*, specific to
species commonly associated with human skin, including *S. aureus*, *S. epidermidis*, and
Staphylococcus capitis[29,39]. Using *lsa*BGC, we discover novel variants of pyrazinones, which
encapsulate seven distinct GCFs (Fig. 2a), that belong to species not commonly found on human
skin, such as *Staphylococcus felis* (GCF-56), the dominant species of *Staphylococcus* on cat skin.
While speciation had resulted in orthologous copies of *pznA* and *pznB* being highly diverged in
sequence, it is surprising to find that the surrounding genomic context of the two genes is highly
variable for all seven GCFs (Fig. 2b), suggesting their ancestral mobilization or relocation within
or between genomes.

Application of *lsaBGC*-Divergence followed by Bayesian shrinkage analysis was then
used to systematically assess whether pyrazinones or other *Staphylococcus* and *Corynebacterium*
GCFs exhibited indications of HGT between species, as suggested by high Beta-RD values (Fig.
S4cd; Table S6; Supplementary Text). Despite the small size of pyrazinones, we do not find
support for the frequent mobilization of any of the seven GCFs encoding them across
staphylococcal species. However, high Beta-RD posterior ranges were observed for two non-
ribosomal peptide synthetase (NRPS) associated GCFs from *Corynebacterium*. One of these

GCFs, predicted to encode for an NRPS-dependent siderophore[40], was found in several species
commonly isolated from skin[37], including *C. kefirresidentii*, from the *C. tuberculostearicum* sp.
complex. Using results from *lsa*BGC-PopGene for this GCF, which includes determination of a
consensus gene order across BGC instances, we find that the NRPS is highly conserved in
sequence across species and found flanked by mobile genetic elements (MGEs), as previously
reported in *Corynebacterium jeikeium*[40] (Fig. 2cd). The taxonomic breadth and high sequence
conservation of this GCF suggest that it has undergone recent HGT between distantly related
species.

We next used *lsa*BGC-PopGene to infer selective pressures acting on homolog groups
within GCFs through calculating inter- and intra-species Tajima's D using select representative
genomes, obtained through genomic dereplication, for each genus, including *Cutibacterium* and
Micrococcus (Table S1, S7; Supplementary Text). While *lsa*BGC-PopGene is highly scalable, it
is important to perform uniform dereplication of genomes to accurately infer species-wide
signatures of selection without being biased by highly represented strains in genomic datasets.
Further, only homolog groups which were found as a single copy within the GCF context and
observed in four of more genomes were considered. We found that the median intra-species
Tajima's D was centered at a value of roughly 0, in accordance with expectations of the null
model (Fig. S5a). Also in alignment with expectations, we observed high values of inter-species
Tajima's D for homolog groups in GCFs found in multiple species (Fig. 5b). A focused intra-
species analysis within *S. aureus* revealed that a hybrid terpene/T3PKS GCF (GCF-3) had the
lowest Tajima's D values and seven homolog groups with values below -2.0, suggestive they are
either highly conserved or, potentially, under sweeping selection (Fig. 5c). The terpene
component of this hybrid GCF encodes for staphyloxanthin carotenoid production[41] and manual
inspection determined that the T3PKS prediction actually corresponds to hydroxymethylglutaryl-
CoA (HMG-CoA) synthase, which is involved in the mevalonate pathway for isoprenoid
synthesis and is homologous to KS domains of polyketide synthases[42]. Further, the regions in
between and flanking the staphyloxanthin encoding operon, *crt*, and HMG-CoA synthase include
several genes important to staphylococcal virulence, some of which were found in multiple
copies in certain species (Fig. S6ab). Because this GCF was predicted to be one of the four GCFs
ancestrally acquired by the *S. epidermidis/aureus* clade, the five gene *crt* operon encoded by it
was found in most species belonging to the clade. Surprisingly however, *crt* genes were missing
in more than 95% of the *S. epidermidis* genomes, the most prevalent staphylococcal species on
skin[34,43], likely due to gene loss (Fig. 4a, S6b).

The virulence-associated carotenoid staphyloxanthin is ubiquitous across the
202
Staphylococcus genus
203

Although staphyloxanthin is well characterized as an *S. aureus* virulence factor[44], we
were unable to identify a report of its distribution across the *Staphylococcus* genus, with only a
single study, to our knowledge, suggesting its production in another species[45]. *lsa*BGC analysis
revealed a total of five staphylococcal GCFs encoding for staphyloxanthin. Following GCF-3,
the second most prevalent GCF encoding staphyloxanthin, GCF-6, had a high Beta-RD
distribution (Fig. S4c) and was widely distributed across divergent species in the genus (Fig. 3a),
as well as *Mammaliicoccus,* the closest phylogenetic neighbors of staphylococci[46]. Constructing
a phylogeny based on a concatenated alignment of *crtM* and *crtN* revealed the appropriate
delineation of GCF-3 and GCF-6 despite their co-occurrence in the genomes of particular

species, such as *Staphylococcus warneri* (Fig. 3b, S7a). Additional analyses showed that GCF-6
has mobilized across species and can be found on plasmids (Fig. 3bcd).

To assess whether carriage of the *crt* operon translates to staphyloxanthin production in
non-aureus species, we identified staphylococcal isolates cultivated from human skin with
whole-genome sequences generated by our lab that encode a staphyloxanthin related GCF. We
identified a rare instance of an *S. epidermidis* isolate with GCF-3 (only 2-3% of *S. epidermidis*
genomes encode GCF-3), strain LK1136, and an *S. warneri* isolate with both GCF-3 and GCF-6,
strain LK413. While we observe GCF-3 in all *S. warneri* genomes, the GCF is degraded
encoding only the dehydrosqualene synthase gene, *crtM*, and the dehydrosqualene desaturase
gene, *crtN* (Fig. 3a; Table S8). However, roughly half of the available *S. warneri* genomes
(47.5%) encode GCF-6 in full and thus have two copies of *crtM* and *crtN*. We used long-read
sequencing to generate near-complete genomes for *S. epidermidis* LK1136 and *S. warneri*
LK413 and found that both strains encode GCF-3 within their chromosomes while GCF-6 is
encoded on a plasmid within *S. warneri*. (Fig. S7b; Supplementary Text).

To test for phenotypic effects associated with strains carrying different staphyloxanthin
GCFs, we assessed pigmentation levels in strains of *S. aureus*, *S. epidermidis*, and *S. warneri*
with and without full *crt* operons (Table S9). Cells of *S. epidermidis* encoding GCF-3 are
pigmented compared to strains without the cluster (Fig. 3e-f). Cells of *S. warneri* LK413
containing the complete *crt* operon (GCF-6) and additional copies of *crtMN* displayed the
strongest golden pigmentation, confirmed as staphyloxanthin by spectrophotometry[41,47] (Fig. 3f).

lsaBGC provides a framework for mining for base-resolution novelty in BGCs

We have demonstrated how the *lsa*BGC suite can be used to identify homologous BGC
instances into GCFs and subsequently uncover their evolutionary trends and phylogenetic
distribution. With increasing metagenomic datasets becoming available, often from complex
microbiomes, we developed *lsa*BGC-DiscoVary to directly profile BGCs and rapidly identify
novel variation within GCF related genes from metagenomes (Fig. S2; Supplementary Text). The
*lsa*BGC framework thus allows users to first identify all instances of GCFs in genomic
assemblies to build a comprehensive database of known alleles for GCF genes and subsequently
apply *lsa*BGC-DiscoVary to find novel SNVs within genes which are absent in the available
assemblies. Further, because *lsa*BGC-DiscoVary extracts the subset of reads supporting the
presence of novel SNVs, exhaustive methods to assess the validity of metagenomic SNV calling,
which would be computationally intensive and impractical to perform for the full set of reads
from a metagenome, are made accessible for further validation of novel SNVs.

We first assessed the sensitivity and specificity of *lsa*BGC-DiscoVary and found high
concordance (>96% overlap) between novel SNVs reported by it when compared to an
assembly-based approach using whole-genome sequencing readsets for 132 *M. luteus* isolates
(Fig. S8; Table S10, S11; Supplementary Text). Next, to demonstrate *lsa*BGC-DiscoVary's
application in metagenomic datasets, its primary intended usage, we investigated the
microdiversity of BGCs from *C. acnes*, the most abundant species at sebaceous sites, within
healthy skin metagenomes generated by our lab[37]. We found that the BGC encoding for
cutimycin, an anti-Staphylococcal thiopeptide, exhibited genomic signatures of horizontal, yet
likely ancestral, acquisition by *C. acnes* and is now largely restricted to isolates belonging to
either clades IB or III(Fig. 4ab, S9, S10a; Supplementary Text). Extensive metagenomic analysis

demonstrated the reliability of *lsa*BGC-DiscoVary to type clade-specific alleles of genes within
the cutimycin BGC in metagenomic datasets which correlated with strain presence inferences
(Fig. 4cd, S10bc; Table S12). We used *lsa*BGC to further investigate novel SNVs detected
within cutimycin genes and found that many were from the metagenomes of a single individual
and belonged to clade III *C. acnes* (Fig. 4efgh). Through observation that the majority of novel
SNVs from this individual also corresponded to cytosine to thymine changes, we discovered that
clade III *C. acnes* have elevated rates of cytosine deamination, which coincides with the clade's
carriage of a truncated uracil DNA glycosylase, the enzyme responsible for the repair of such
deaminations (Fig. 4ijk).

lsaBGC identifies novel SNVs within polyketide synthase catalytic domains

Phylogenetic and average nucleotide identity (ANI)[48] based investigations revealed that
C. tuberculostearicum exhibits high genomic similarity (ANI >88%) to four other species in
GTDB (Fig. S4b, S11a), which we thus refer to, in aggregate, as the *C. tuberculostearicum*
species complex. While metagenomic profiling found that this complex features the most
prevalent *Corynebacterium* species within healthy skin microbiomes (Fig. S4b, Table S4), only
22 genomic assemblies were available for the complex at the time of this analysis. This clade
thus represents an ideal taxonomic group to examine novel BGC microdiversity using *lsa*BGC-
DiscoVary. We ran *lsa*BGC-DiscoVary systematically on six GCFs identified in the species
complex, largely encoding for unknown metabolites, and found 40,019 putatively novel SNV
instances across 105 homolog groups and 109 metagenomic samples. Of these, 5,474 instances
were filtered because reads supporting their presence aligned with a higher score to other regions
in a concatenated database of all *Corynebacterium* genomes from GTDB as compared to the
reference database used for *lsa*BGC-DiscoVary. While ubiquitous, *C. tuberculostearicum*
species complex members are present at very low abundances in some body-sites where
metagenomic assembly would struggle to construct their genomes (Fig. S11b). We performed
sample-specific metagenomic assemblies[49] and found that at least 11,659 novel SNV instances
detected by *lsa*BGC-DiscoVary would be missed by assembly-based investigation (Fig. 5a;
Supplementary Text).

To further investigate the novel SNVs detected, we focused on 68 homolog groups
overlapping or in close proximity to protocore regions of BGCs and further filtered based on
standardized coverage metrics, retaining 5,802 instances of 2,343 unique novel SNVs. Novel
SNVs were more likely to correspond to synonymous changes if they were found in multiple
metagenomic samples (Fig. S11c). Further, metagenomic samples from the same subject at
different body sites and from the same body sites across different subjects were more likely to
share novel SNVs when compared to metagenomic samples from different body sites and
different subjects (*p*<1E-8; two-sided Wilcoxon rank sum test) (Fig. S11d).

Finally, we expected and observed that the ratio of predicted synonymous to non-
synonymous novel SNVs increases at deeply conserved sites along homolog groups (Fig. 5b;
Table S13; Supplementary Text). The homolog group with the greatest number of novel SNVs,
and novel, non-synonymous SNVs within highly conserved regions, was the PKS gene
responsible for synthesis of mycolic acid, a major component of the cell wall in most
Corynebacterium species recently shown to have immunomodulatory effects on the skin[30,50,51].
We identified and manually validated three non-synonymous SNVs along conserved sites of the

PKS gene, including two adjacent SNVs predicted to affect one codon within the ketoacyl
synthase domain, which catalyzes condensation[52], and could alter reaction kinetics[53] or scaffold
structure[54] (Fig. 5c, S12; Table S14).
304
305

306 **Discussion**
307

The *lsa*BGC suite packages several fundamental, as well as novel, functionalities for
high-throughput comparative and evolutionary analysis of BGCs from any taxa in a structured
framework, such as the ability to detect GCFs within draft-quality genomic assemblies. Of
practical relevance, it produces a comprehensive spreadsheet that includes annotation,
conservation, and evolutionary statistics for homolog groups found in GCFs to enable efficient
and high-throughput assessment by users. By incorporating GToTree[55], *lsa*BGC will also
automatically generate visualizations showing the distribution of GCFs across a species
phylogeny.

One of the current limitations is that OrthoFinder based homology inference can be
coarse and similar proteins encoded by recently duplicated genes can be collapsed into the same
homolog group[13]. To mitigate this limitation, we clearly mark which homolog groups do not
correspond to single-copy-orthologs within GCF contexts in resulting reports.

The metagenomic mining functionalities of *lsa*BGC further provide insight into the
presence and sequence diversity of a specific taxa's BGCs across diverse microbiomes. This is a
particularly important feature for rare BGCs and taxa with a limited numbers of genomic
assemblies available, such as taxa that are difficult to cultivate from complex metagenomes. In
the future, the use of pan-genome alignment approaches to potentially increase the efficiency of
*lsa*BGC for metagenomic identification of novel variants within BGC genes[56] may be explored.

Here, we apply *lsa*BGC's functionalities to major taxa commonly found in skin
microbiomes as a proof of concept to demonstrate the potential to advance understanding of the
diversity, evolutionary trends, and prevalence of both well studied and uncharacterized
secondary metabolites. For example, we uncovered the ubiquity of the carotenoid
staphyloxanthin across the genus of *Staphylococcus,* a molecule conferring the signature golden
pigmentation in *S. aureus*[57,58]. These findings raise the hypothesis that staphyloxanthin, which
has been identified primarily as a virulence factor conferring oxidative stress resistance in the
pathogen *S. aureus*[44,59], could similarly contribute to virulence in other staphylococcal species or
serve additional unknown functions. Supporting this hypothesis, we observe that one
staphyloxanthin encoding GCF appears ancestral to the skin-associated, multi-species clade
which features both *S. aureus* and *S. epidermidis*. The observation that multiple copies of *crt*
genes can be found within a single genome further suggests staphyloxanthin is of functional
importance since evolutionary retention of paralogous genes has previously been shown to
underlie ecological advantages[60,61]. Additionally, while this GCF is present in most skin-
associated species, it is absent in greater than 95% of *S. epidermidis* genomes, suggesting large-
scale loss within the species. It will thus be interesting to further explore how the loss of
staphyloxanthin, if it contributes to oxidative stress resistance within *S. epidermidis*, coincides
with the species' reputation as a beneficial symbiont of host skin[34,62].

Using the novel metagenomic mining functionality of *lsa*BGC, we identified novel SNVs distinguishing uncultured *C. acnes* and *Corynebacterium* strains from isolates with genomes available in public databases. With regards to *Corynebacterium*, we determined that some of the most common skin residing species[33,37] belong to the *C. tuberculostearicum* species complex, which were widely distributed across skin metagenomes and highly abundant at moist body sites. Because this species complex is vastly underrepresented in public genomic databases and additional species and strains belonging to it likely remain to be discovered, application of *lsa*BGC-DiscoVary led to the identification of thousands of novel SNVs within the BGCs of the clade. Critically, some of these variants could have functional implications as they lie within highly conserved regions of important domains for natural product biosynthesis.

The comprehensive and open-source packaging of *lsa*BGC allows analyses to be applied to both public and proprietary microbial genomic databases. Besides accelerating fundamental evolutionary research of BGCs from diverse taxa, we envision *lsa*BGC will be particularly useful for natural product discovery and guiding mechanism of action studies. Evolutionary insight gathered from *lsa*BGC can be used as prior information to train algorithms in an emerging field using artificial intelligence to identify new antibiotics[63,64]. Further, an improved understanding of the relationship between intra-taxa genomic diversity and BGC content novelty at the base-resolution can also highlight which sub-lineages or sub-clades bear the largest reservoir of untapped secondary metabolic potential. Finally, we posit that identifying evolutionary trends of BGCs detected by highly-reliable rule-based approaches, such as antiSMASH[8], can be used to assess the validity of BGC predictions from newly developed machine learning approaches, such as DeepBGC[11] or GECCO[65], which we now support usage of in *lsa*BGC.

Acknowledgments

The authors gratefully acknowledge additional members of the Kalan lab, Dr. John-Demian Sauer, Dr. Jason Kwan, Dr. Karthik Anantharaman, Dr. Caroline Grunenwald, Dr. Chase Clark, and Dr. Milton Drott for helpful discussions.

Funding: This work was supported by grants from the National Institutes of Health (NIAID U19AI142720 and NIGMS R35GM137828 [L.R.K]). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

Author contributions: Conceptualization: RS, LRK. Computational Analysis and Software Development: RS. Investigation: RS, JC, LRK. Isolate Culturing and Sequencing: SS, MS, NLS, AS, LRK. Visualization: RS, JC. Supervision: LRK. Writing—original draft: RS, LRK. Writing—review & editing: RS, JC, SS, MS, RMS, NLS, CRC, AS, LRK.

Competing interests: The authors declare that they have no competing interests.

Materials and Methods

Software availability

The *lsa*BGC software suite was developed in Python3 and R and is available on Github at: https://github.com/Kalan-Lab/lsaBGC. Algorithmic descriptions of programs can be found on

389 the Github wiki and in the Supplementary Text document. A small test case of *Cutibacterium*
390 genomes is included in the Github repository. Version 1.0 of the software was used for the
391 analyses described in this study.
392
An overview of the *lsa*BGC software suite

The first core program is *lsa*BGC-Cluster which clusters homologous and non-
fragmented instances of BGCs, identified by antiSMASH from chromosomally complete
genomic assemblies, into GCFs, similar to BiG-SCAPE[9].Users can then generate a report to
select for the most appropriate parameters for downstream analyses of interest. Because
antiSMASH groups neighboring BGCs into hybrid gene clusters if they are located close to one
another on the genome[7], *lsa*BGC-Refiner has the option to manually separate hybrid clusters
based on user-defined boundary homolog groups. Homologous instances of GCFs across all
available genomic assemblies for the taxa of interest can then be identified through *lsa*BGC-
Expansion (Figure S1b). Similar to ClusterFinder[10], *lsa*BGC-Expansion uses profile Hidden
Markov models (HMMs) to identify homologs of GCF associated genes in assemblies followed
by a classical HMM approach to define BGC regions and their boundaries. Unlike ClusterFinder,
*lsa*BGC-Expansion: (i) uses all coding genes associated with a GCF instead of being restricted to
core protein domains, (ii) is specifically designed to account for assembly fragmentation through
conditional assessment of predicted regions, and (iii) features a "quick mode" option for fast
detection of BGCs using DIAMOND[66].

The remaining four core programs are analytical and designed to be run on a single GCF
for ease of parallelization: *lsa*BGC-See, *lsa*BGC-PopGene, *lsa*BGC-Divergence, and *lsa*BGC-
DiscoVary. *lsa*BGC-See permits visualization of BGCs across a user-provided species tree or a
BGC based phylogeny and can handle multiple fragmented BGC instances from the same
genome, depicting them as neighboring leaves on phylogenies. *lsa*BGC-PopGene can then infer
evolutionary and population genetics statistics for each homolog group associated with a GCF. It
generates codon based alignments for each homolog group which can then be used as input for
*lsa*BGC-Divergence and *lsa*BGC-DiscoVary. *lsa*BGC-PopGene also determines the consensus
order and direction of homolog groups for a GCF across all BGC instances, which provides
intuitive assessment of its reports. *lsa*BGC-Divergence calculates a comprehensive statistic
between pairs of homologous BGCs from the same GCF but different genomes, which we refer
to as Beta-RD, that captures how similar the BGC sequences are relative to what would be
expected based on genome-wide similarity. Finally, *lsa*BGC-DiscoVary is a multi-functional
program which can mine raw sequencing readsets from metagenomes or single genomes, for
GCF instances and then reliably identify whether they possess novel single nucleotide variants
(SNVs) yet to be observed in available genomic assemblies for a taxa (Figure S2).

Automated workflows for running *lsa*BGC

To automate the application of these programs, we had initially developed three
workflow programs called lsaBGC-AutoProcess.py, lsaBGC-AutoExpansion.py, and lsaBGC-
AutoAnalyze.py. lsaBGC-AutoProcess.py is a preliminary workflow which does not execute any
of the aforementioned core programs of the suite. It functions to take in a listing of genomic
assemblies and perform gene-calling and basic annotation using Prokka[67], annotate BGCs using
antiSMASH[8], and finally run OrthoFinder2[13] for *de novo* delineation of homolog groups. It runs
antiSMASH and OrthoFinder2 using the same Prokka output to ensure locus tag identifiers are
able to be matched between the output of the two former programs (Figure S1a). lsaBGC-

AutoExpansion.py is a wrapper of lsaBGC-Expansion.py which performs additional GCF
instance identification in draft assemblies across all GCFs identified for a taxa. It is
recommended that users run lsaBGC-AutoExpansion.py instead of lsaBGC-Expansion.py
individually because the workflow additionally features a critical consolidation step in which
BGC instances identified as potentially belonging to multiple GCFs are re-assessed and assigned
to only the single best fitting GCF (Figure S1c). Finally, lsaBGC-AutoAnalyze.py is a major
workflow which runs the analytical core programs across each GCF and creates a few
consolidated reports after completion. This workflow starts by computing pairwise whole-
genome similarity metrics using either CompareM[68] or FastANI[48] and then runs lsaBGC-See.py,
lsaBGC-PopGene.py, lsaBGC-RelativeDivergence.py and, optionally, lsaBGC-DiscoVary.py for
each GCF. At the end of the workflow, it generates consolidated report tables and visualizations
from GCF-specific results for lsaBGC-PopGene.py and lsaBGC-RelativeDivergence.py. These
workflows, similar to the core programs, are documented on the Github wiki.

Recent additions to the *lsa*BGC suite

More recently, we have introduced two new core and workflow programs into the suite
(release 1.2). The first is the core program lsaBGC-Ready.py, which serves to format existing
BGC predictions in GenBank format and, optionally, BiG-SCAPE[9] results, together with
corresponding genomes, as input for downstream core programs in *lsa*BGC. Briefly, lsaBGC-
Ready.py performs gene-calling on genomes using prodigal[69], consolidates gene calling with
those present in BGC GenBanks, performs annotation using KOfam and PGAP profile
HMMs[70,71], and can automatically run lsaBGC-Cluster.py and lsaBGC-AutoExpansion.py. We
now recommend lsaBGC-Ready.py in place of the workflow lsaBGC-AutoProcess.py due to its
simplified usage.

The second is an automated workflow for running *lsa*BGC on any bacterial species or
genus called lsaBGC-Easy.py. Briefly, this workflow assesses GTDB[72] for high-quality genomic
assemblies belonging to a specified genus or species, downloads the genomes using ncbi-
genome-download (https://github.com/kblin/ncbi-genome-download), performs gene-calling
using prodigal[69], constructs an alignment of single copy genes and phylogeny using GToTree[55],
dereplicates genomes based on the constructed alignment, and then runs lsaBGC-Ready.py,
lsaBGC-Cluster.py, and lsaBGC-AutoAnalyze.py. The workflow also allows for users to include
their own genomes belonging to the taxa, which might not yet be available on NCBI.

Analytical changes to the core and workflow programs since the initial release of *lsa*BGC
(version 1.0) in relation to the most recent release of the software (version 1.3) have been
minimal (Supplementary Text).

Usage of genomic and metagenomic datasets

To overcome issues with misclassification and contamination in NCBI, we used GTDB
(release 202) [72], to query for reliable genomic assemblies and respective species designations for
the genera of *Corynebacterium*, *Cutibacterium*, *Micrococcus*, and *Staphylococcus* (Table S1).
For *Micrococcus*, we further included 132 additional genomes of the type species *M. luteus*
which our lab had sequenced (BioProject ID PRJNA803478) but were not yet uploaded to NCBI
to be included in the GTDB release used. These isolates were classified as *M. luteus* by GTDB-
tk[73] and exhibited > 95% ANI to the NCBI representative genome for the species *M. luteus*
NCTC 2665. A similar approach was used for gathering genome sets for species-level analyses
performed (Table S1). For the *C. tuberculostearicum* species complex, an additional five

genomes from NCBI were gathered, classified using GTDB-tk[73], and incorporated into analysis
if they belonged to one of the five species designated as belonging to the complex.

Because a single representative per species as provided by GTDB might not adequately
capture the diversity of BGCs found across a genus, we instead employed a more granular
approach to select for representative genomes for a genus. We performed dereplication using
MASH[74] with an estimated ANI cutoff of 0.99 to pair similar genomes and dynamically select
representative genomes from each cluster based on assembly N50 (Table S1). This dereplication
was performed using the script popSizeAndSampleSelector.py provided in the *lsa*BGC suite.
Complete or chromosome level assemblies among the representative genomes were run through
lsaBGC-AutoProcess.py and lsaBGC-Cluster.py to identify the core set of GCFs per genus.
GCFs determined from the core set of genomes were comprehensively identified in the
remaining genomes using lsaBGC-AutoExpansion.py, including those which were not deemed as
representatives from the MASH based analysis to showcase scalability (e.g. all 15,055
Staphylococcus and not just the 229 representative genomes of which 77 corresponded to
complete/chromosome level assemblies). To expedite population genetics analysis however, we
only ran the downstream functional analytics of *lsa*BGC using the representative genomes for
each genus. For species-level analyses, all genomes available were used for such downstream
analyses and to comprehensively profile known alleles for genes in BGCs, which was critical to
enable metagenomic mining for base-resolution novelty previously unseen in assemblies
available for a species. For population level analyses within species, strain delineations were
performed using the popSizeAndSampleSelector.py script in *lsa*BGC with strain groups
determined from single-linkage clustering of pairs of genomes found to exhibit ≥98% ANI and
≥80% genomic content similarity by FastANI[48].

For determining skin-associated species amongst the genera investigated in our study and
to demonstrate mining functionalities with *lsa*BGC-DiscoVary, we used a paired-end
metagenomics dataset sequenced on the Illumina NovaSeq where individual samples
corresponded to microbiome surveys of one of eight body sites for one of 34 participants,
collected in the state of Wisconsin (BioProject ID PRJNA763232)[37]. As previously described[37],
adapter removal, quality filtering, human sequence decontamination, and tandem repeat removal
were performed using fastp (v0.21.0)[75] and KneadData (v0.8.0)
(https://huttenhower.sph.harvard.edu/kneaddata/).

Genomic assessment for staphyloxanthin carriage in the lab isolate collection

We had previously constructed draft genomic assemblies for 44 *Staphylococcus* isolates
from skin (BioProject ID PRJNA803478; BioProject ID PRJNA830888). Illumina sequencing
was performed at the Microbial Genome Sequencing Center (MiGS). Briefly, assemblies were
generated using Unicycler[76] with default settings after quality and adapter trimming using
fastp[75]. The set of 44 *Staphylococcus* genomes were searched for carriage of staphyloxanthin
encoding GCFs identified in our study using *lsa*BGC-AutoExpansion (Table S9).

Library preparation, sequencing and genomic assembly

Reconstruction of Illumina draft assemblies for 132 M. luteus: We assessed the
performance of multiple programs in the *lsa*BGC suite using a set of genomic assemblies for 132

M. luteus isolates which were cultured from skin. For these isolates, assemblies were
reconstructed from raw-reads (BioProject ID PRJNA803478) and were uploaded to NCBI's
Genbank database (Table S1). Briefly, Illumina sequencing reads were processed using fastp (v.
0.20.1)[75] to trim for Poly-G tail artifacts and TrimGalore (v. 0.6.5)[77] was additionally used to
detect and remove adapters. Afterwards, assembly was performed using Unicycler (v. 0.4.6)[76]
with default settings.

Construction of Illumina+Nanopore hybrid assemblies for staphyloxanthin producing S.
warneri *LK413 and* S. epidermidis *LK1136*: To learn the genomic location of staphyloxanthin
encoding GCFs in *S. epidermidis* LK1136 and *S. warneri* LK413, we performed additional long-
read sequencing using Oxford Nanopore Technologies (ONT). Library construction, using a
PCR-free ligation method, and sequencing were performed at the Microbial Genome Sequencing
Center (MiGS). Hybrid assemblies were constructed using a modified version of the Hybrid
Assembly workflow in seQuoia (https://github.com/broadinstitute/seQuoia), Illumina reads, used
for construction of the initial draft genomic assemblies for the isolates, were first reprocessed
using fastp (v. 0.20.1)[75] to trim for Poly-G tail artifacts and TrimGalore (v. 0.6.5)[77] was
subsequently used to detect and remove adapters. Similarly, ONT FASTQ reads, provided by
MIGS after basecalling, were filtered for potential adapters using PoreChop (v. 0.2.3)
(https://github.com/rrwick/Porechop) and filtered to retain only reads greater than 3 kb in length
as well as subsampled to 300 mb (while being inclusive of all reads greater than 20 kb) using the
script fastqfilter.py included in seQuoia. Afterwards, hybrid-assemblies were constructed using
Unicycler (v. 0.4.6)[76] with default settings and additional Pilon (v. 1.23)[78] polishing was
iteratively performed by aligning ONT reads[79] as well as the Illumina reads[80] to assemblies and
refining them, allowing for a maximum of 10 iterations. Quality assessment of the two near-
complete assemblies was performed using GAEMR (https://github.com/broadinstitute/GAEMR)
(Supplementary Text).

Experimental validation of staphyloxanthin production in non-aureus *Staphylococcus*

Isolates were grown overnight at 37°C on tryptone soy agar plates and inoculated into
tryptone soy broth at a concentration of 1×10^6 CFU/mL. Liquid cultures were grown in shaking
conditions at 37°C for 24 hours. 10 mL of the liquid culture was then pelleted and washed twice
with phosphate-buffered saline before pigment extraction in 1 mL of methanol at 55°C for 15
mins. Spectral scans from 300 nm to 600 nm were taken using a plate reader (EPOCH2, Biotek,
Winooski, VT). Absorbance readings were blanked with methanol and normalized to liquid
culture cell density based on OD600. Spectra shown are representative of at least two biological
replicates.

MASH distance neighbor-joining tree construction

Average nucleotide identity estimates between pairs of genomes were computed with
MASH[74] and used to create a distance matrix. The distance matrix was then used to construct a
neighbor-joining tree with the ape library in R[81].

Ribosomal and BGC phylogeny construction

Ribosomal phylogeny construction: Ribosomal phylogenies for each species were
constructed based on a concatenated alignment of 16 ribosomal proteins[82]. HMMER (v3) was
used to search for profile HMMs of each ribosomal protein in the predicted proteomes of
samples of interest belonging to a particular taxonomic group. Afterwards, the best matching hits
for each sample based on alignment E-value were identified and used to construct individual
alignments for each ribosomal protein with the L-INS-I method in MAFFT[83]. Protein alignments
were then converted to codon alignments using PAL2NAL[84] and subsequently concatenated
together. Samples which featured ≥ 25% gaps in the concatenated alignment were filtered after
which core SNV sites were extracted and used for constructing a maximum likelihood phylogeny
with RAxML[85], specifying a GTRCAT model and performing 100 bootstraps. Where needed,
ribosomal phylogenies were subset to showcase relationships between select samples using
PareTree (http://emmahodcroft.com/PareTree.html ; v1.0.2).

Staphyloxanthin phylogeny construction: A maximum likelihood phylogeny was
constructed from a concatenated protein alignment of CrtM and CrtN amongst representative
Staphylococcus genomes to understand the relationship between different GCFs encoding for
staphyloxanthin. Proteins designated as HG0001876 (CrtM) and HG0001904 (CrtN) were
gathered for each staphyloxanthin encoding GCF found in each sample and aligned using the L-
INS-I method in MAFFT[83]. To account for assembly fragmentation resulting in multiple proteins
from a sample being assigned to the same homolog group for a specific GCF, a consensus
sequence was determined and multi-allele sites were replaced as gaps. A concatenated alignment
of such consensus sequences was subsequently constructed and used for phylogeny construction
with RAxML using a PROTCAT model with JM selected as the best amino-acid replacement
matrix through the automatic maximum likelihood model selector[85]. The phylogeny of CrtMN
sequences exhibited high correspondence to GCF delineations with the exception of a single *S.*
lugdunensis BGC which is likely misclassified as GCF-17 and should be GCF-3. Upon further
examination, it was identified that the reason for the misclassification was likely due to the
absence of mevalonate pathway genes, classified as a type III polyketide synthase (T3PKS),
within *S. lugdunensis* which are found downstream of the *crt* operon in all other species with
GCF-3 (Figure S7b).

Pyrazinone phylogeny construction: A maximum likelihood phylogeny was constructed
from a concatenated protein alignment of PznA and PznB amongst representative
Staphylococcus genomes using the same methods as described for phylogeny construction from
CrtMN proteins.

Designation of species belonging to *Mammaliicoccus*

The *Staphylococcus_A* genus designation in GTDB R202 was largely concordant with
the recent reclassification of five species from *Staphylococcus* to *Mammaliicoccus*[46]. In the latest
GTDB R207 release, this genus has now been reclassified to *Mammaliicoccus*. The recent update
to taxonomic names also includes the reclassification of species *S. pasteuri_A* to *M. fleurettii*.
Notably, we do not regard *S. schleiferi* as part of *Mammaliicoccus* because it is categorized as
Staphylococcus by GTDB (both R202 and R207) and groups within the *Staphylococcus* genus in
our ribosomal protein based maximum-likelihood phylogeny. Similar to our assessment, the
reclassification of this species to *Mammaliicoccus* was recently questioned[86].

Ubiquity assessment of skin residing species and strains of *Staphylococcus* and *Corynebacterium*

We searched for the presence of representative *Staphylococcus* and *Corynebacterium* strains in 270 skin metagenomes from 34 participants across 8 body sites using StrainGST[38]. To construct the database of distinct representative genomes for each genus, the protocol described in the StrainGE documentation was followed. *S. aureus* genomes 16405 and 278 were dropped from the analysis because at the time of downloading them, the plasmid and chromosome were switched. Briefly, genomes were downloaded from NCBI using ncbi-genome-download (https://github.com/kblin/ncbi-genome-download) and dereplication was performed using built-in StrainGE functionalities. As some representative genomes were not part of the GTDB release R202, GTDB-tk[73] was used to annotate such genomes and perform species classification. We considered a species as skin-associated if it was found in at least 10 of the 270 skin metagenomes, regardless of the abundance (Table S4). Certain skin-associated species from each genus were also identified in the 19 negative control metagenomic samples, which included air-swab, bench, desk, and door-handle samplings (Table S4). Because we expect heavy overlap in species content between these environmental samples and skin microbiome samples, we still considered such species as skin-associated if they were found in 10 or more skin metagenomes.

We found the majority of skin-associated *Staphylococcus* belonged to the *S. epidermidis/aureus* clade, as has previously been noted[42]. Similarly, most of the skin-associated *Corynebacterium* belonged to *C. tuberculostearicum* and it's close neighboring species (including *C. kefirresidentii*, *C. aurimucosum* type E, *C.* sp. 900539985, *C. tuberculostearicum*, *C. tuberculostearicum* type C), which we denote in this study as the *C. tuberculostearicum* species complex. The minimal ANI separating genomes belonging to this species complex was determined to be 88.59% using FastANI[48]. To assess whether any of the 24 new *Corynebacterium* species reported by Saheb Kashaf et al. 2022[36] belonged to this species complex, FastANI was used to look at their ANI similarity to genomes from the clade and regard them as additional members if they exhibited 88.59% ANI or greater. Of note, many additional *Corynebacterium* species have been routinely isolated from the skin[37] which we did not regard as skin-associated based on our criteria in this study. In particular, the species of *C. amycolatum* (detected in 4 metagenomes), *C. kroppenstedtii* (detected in 4 metagenomes)*,* and *C. simulans* (detected in 5 metagenomes) are commonly found on skin.

To estimate the relative abundance of *Cutibacterium* (see Supplementary Text section "Identification of *Cutibacterium* Strains within Metagenomes and Determination of Near-Fixed Sites Differentiating Cutimycin Biosynthesis Alleles between Clade I and Clade III *C. acnes*") and *Corynebacterium* strains within metagenomic samples we normalized the estimated relative abundances for strains (*rapct*) by the estimated relative abundance of the genus (*pan%*) as reported by StrainGST[38].

Identification of homolog groups associated with skin residing species of *Staphylococcus* and *Corynebacterium*

We developed a script called crawlingFisher.py, provided in the *lsa*BGC suite, which tests for enrichment or depletion of homolog group presence amongst genomes in focal clades using Fisher's exact test. Provided with a phylogenetic tree and matrix specifying the presence of homolog groups across genomes, the program automatically performs such tests for each

homolog group at each inner-node of the phylogeny. Multiple testings correction was performed
comprehensively at the end using Benjamini-Hochberg false discovery rate and cases were
considered statistically significant if the adjusted p-value was less than 0.05. Results were further
filtered to retain only homolog groups which were present in at least 80% of genomes under the
focal node and at most 20% of other genomes or, alternatively, at most 20% of genomes under
the focal node and at least 80% of other genomes. Finally, only results pertaining to focal nodes
where at least 80% of genomes were classified as skin-associated (see section "Ubiquity
Assessment of Skin Residing Species and Strains of *Staphylococcus* and *Corynebacterium*")
were reported (Table S5).

Assessing the divergence of BGCs relative to their background genomic contexts

BGC divergence does not always track with background genomic divergence and
speciation[14]. The program *lsa*BGC-Divergence allows for easy assessment of how BGC
divergence compares to whole-genome divergence, primarily through computing a statistic we
refer to as Beta-Relative-Divergence (Beta-RD) between pairs of genomes for a single GCF. The
Beta-RD statistic is simply the average amino or nucleotide identity between homologous
instances of proteins for a GCF observed in two genomes normalized by the average amino or
nucleotide identity observed between the full genomes. *lsa*BGC allows users to specify whether
amino or nucleotide divergence should be used and automated workflows (see section
Automated Workflows for Running *lsa*BGC) can be used to compute pairwise AAI or ANI
between genomes using CompareM (https://github.com/dparks1134/CompareM) or FastANI[48],
respectively.

To better understand how distributions of Beta-RD differ across GCFs, in this study, we
applied a Bayesian hierarchical model to perform shrinking analysis and appropriately account
for differences in the number of genomes GCFs are found (Figure S4cd; Table S6). For
computational efficiency, a maximum of 500 Beta-RD observations were used for each GCF.
Posterior predictions for Beta-RD were made using rstan (Stan Development Team 2021) with
the following hierarchical model:

$$\bar{\mu} \sim Norm(\mu_0, \sigma_0^2)$$

$$\tau \sim half - t_7(1)$$

$$\mu_1 \dots \mu_J \sim Norm(\bar{\mu}, \tau)$$

$$\sigma \sim half - t_7$$

$$y_j \sim Norm(\mu_j, \sigma^2)$$

where for each GCF j, the posterior Beta-RD distribution is sampled as y_j using Markov chain
Monte Carlo with four chains, each featuring 2,000 iterations, including a burnin of 1,000
iterations. A hierarchical prior was specified for μ_j, which represents a GCF-specific baseline.
The values of μ_0 and σ_0 were set to the respective mean and standard deviation of raw Beta-RD
values across all GCFs. R and STAN code used for this analysis are available within the scripts
subdirectory of the *lsa*BGC software package.

Scrutinization of putative novel variants for *lsa*BGC-DiscoVary application to cutimycin in *C. acnes* and the comprehensive set of GCFs for the *C. tuberculostearicum* species complex

By default, lsaBGC-DiscoVary.py only considers reads aligning at 95% identity to reference gene sequences and avoids reporting novel variants on genes if either annotation or coverage (relative to other genes in the BGC context) suggests they might be MGEs. The program also automatically extracts the subset of reads supporting the existence of novel variants, which can then be further screened with high stringency filters using otherwise computationally expensive methods. This can involve mapping reads supporting putative novel SNVs to a representative database of whole-genomes to observe for higher quality alignments or checking whether reads are classified as a particular taxonomy by Kraken2 (v2.0.8-beta)[87]. As described previously[37], we used a customized database for Kraken2 analysis constructed from chromosomally complete bacterial, viral, archaeal, fungal, protozoan genomes as well as the human genome available on RefSeq, where plasmid sequences were distinguished to alleviate potential misclassifications of reads to a particular taxonomy.

For reads supporting putative novel variants in the cutimycin BGC of *C. acnes*, we required them to be classified as *Cutibacterium* or *Priopionibacterium* by Kraken2[87] and to not map with a higher alignment score to a comprehensive database of all *Cutibacterium* genomes than was observed for the alignment to the reference gene sequence in lsaBGC-DiscoVary.py (Figure S2c). We generally do not recommend screening variant-supporting reads for an appropriate taxonomic match with Kraken2 as it can generally lead to false negatives (on average only 92.8% of read-pairs per sample were taxonomically classified and misclassification can also occur; Figure S12) but performed this here to provide further assurance that novel variants discovered are reliable. Standardized coverage metrics within lsaBGC-DiscoVary.py's novel SNV report were further used to conservatively filter sites and variants if they exhibited site or allele-specific coverage values greater than two median absolute deviations from the median coverage for the entire BGC.

For reads supporting putative novel variants within GCFs of the *C. tuberculostearicum* species complex, we required them to not align with a higher alignment score to a comprehensive database of all *Corynebacterium* genomes. We did not require reads to be classified by Kraken2 as *Corynebacterium* because genomes from the species complex are likely to be poorly represented in the Kraken2 database. We did however further validate that reads supporting the existence of variants at conserved sites in the mycolic acid biosynthesis gene mapped best to the *C. tuberculostearicum* species and confirmed our suspicion that they could be misclassified by Kraken2 given the representation in the underlying database (Figure S12). This *ad hoc* analysis was performed using paired-end information when mapping in Bowtie2 and accounting for concordant read pair alignments. While many reads supporting the existence of novel SNVs in the biosynthesis gene were classified as *C. acnes* by Kraken2, we found that none of the reads concordantly aligned to the comprehensive *Cutibacterium* genomes Bowtie 2 database whereas 85.07% concordantly aligned to the comprehensive *Corynebacterium* genomes Bowtie 2 database.

Note, the reasoning behind mapping SNV supporting reads to comprehensive genomic databases to purge false positive SNV calls, is that if a read truly aligns best to the BGC-associated homolog group where it was used to call a SNV, then it should at most generate a mapping to this comprehensive database of equal mapping quality. We align paired-end reads individually as was performed in lsaBGC-DiscoVary.py to have mapping scores be comparable

with those from the original alignment. Furthermore, because our initial database within lsaBGC-DiscoVary.py consisted of allelic representatives of homolog groups, we realized that reads mapping to the start and end of genes could result in a lower mapping score than if a larger genomic context was provided. To ensure we do not remove such reads mapping to gene edges, scrutunizeNovelSNVSupport.py, the program we developed to perform Kraken 2 and genomic database alignment based filtering of novel SNV reports from lsaBGC-DiscoVary.py, also compares the reference sequences when it finds a better Bowtie 2 mapping for SNV-supporting reads in genomic databases. If the reference sequence in the genomic database encompasses the reference sequence in the original alignment to the lsaBGC-DiscoVary.py homolog group database and the read was previously noted to map to a gene edge, then reads are still retained and SNV read support is not decremented. Five retained and supportive reads are required for an SNV to be featured in the filtered report produced by the program.

lsaBGC-DiscoVary based analyses of BGCs from the *C. tuberculostearicum* species complex:

For the following analyses, we used 5,802 instances of 2,343 unique novel SNVs found in 68 homolog groups which were core or adjacent to core biosynthesis machinery of BGCs from the *C. tuberculostearicum* species complex. These novel SNVs had site and allele-specific coverage values less than two median absolute deviations from the median coverage for the entire BGC.

<u>*Comparing SNVs shared across body sites and subjects*</u>: To comprehensively assess whether novel SNVs were shared more frequently for microbiomes of the same body site or from the same individual, we performed a pairwise, multi-iteration resampling analysis. For each pair of metagenomes, where each sample had at least 30 novel SNVs, we calculated the average Jaccard similarity across 1,000 simulations in which we randomly drew sets of 30 SNVs from each metagenomic sample to control for sequencing depth. Pairs of metagenomes were classified as one of three categories: "different body site, different participant", "different body site, same participant", and "same body site, different participant" (Figure S11c). A two-sided Wilcoxon rank sum test for differences between the three distributions of average Jaccard similarities revealed that there was a statistically significant difference between "different body site, same participant' and "same body site, different participant" (p=6.01E-3). Of greater relevance, there were statistically strong significant differences between the "different body site, different participant" distribution and the two other distributions: "different body site, same participant" and "same body site, different participant" (p=3.67E-09 and p=1.39E-116, respectively). Thus, pairs of metagenomes were more likely to share novel SNVs if they were either from the same body site or same participant compared to if they were from different participants and different body sites. Two metagenomic samples for participant S002 were not used because they corresponded to a resampling of two of their body sites and we did not have enough of such samples to pursue a temporal analysis. We assessed this testing using different considerations/cutoffs for: (i) the number of novel SNVs needed for a metagenome to be considered in the analysis and (ii) whether or not to account for singleton SNVs. In all cases, statistical testing for differences between distributions of pairwise metagenome categories yielded similar conclusions.

Assessing trends for novel SNV predicted synonymous to non-synonymous rate with
metagenome ubiquity and site conservation: Using reports from lsaBGC-DiscoVary.py, we
performed a systematic prediction of whether novel SNVs identified on the 68 homolog groups
nearby or overlapping with protocore regions of BGCs in *C. tuberculostearicum* corresponded to
synonymous or non-synonymous substitutions. As described in other sections, SNV alleles were
compared to corresponding alleles observed at the site on the reference gene sequences which
SNVs were called upon. We aimed to check whether the rate of synonymous to non-synonymous
novel SNVs showed a relationship to: (i) how common SNVs were across metagenomic samples
and (ii) conservation levels across protein sequences.

Novel SNV predicted synonymous to non-synonymous rate increases with SNV
metagenome ubiquity amongst metagenomic samples: The occurrence of novel SNVs, a novel
allele observed at a specific site in the multiple sequence codon alignment of a particular
homolog group, was tabulated across metagenomic samples. While novel SNVs found in a single
metagenome were only twice as likely to correspond to a synonymous change as opposed to a
non-synonymous change, more prevalent novel SNVs, found in ten or more samples, were eight
times more likely to correspond to a synonymous change (Figure S11b).

Novel SNV predicted synonymous to non-synonymous rate increases at conserved sites in
protein sequences: We searched RefSeq's bacterial NR database for remote homologs of the 68
homolog groups within or adjacent to protocore regions of BGCs. The top 20 homologs which
belonged to classified bacterial species outside of the *Corynebacterium* genus or
Corynebacteriales family were identified using HMMER3 with profile-HMMs gathered from
*lsa*BGC-Expansion for each homolog group and an E-value threshold of 1E-20 (Table S13). The
sequences of these representative homologs were added to protein alignments for homolog
groups in *lsa*BGC-PopGene results using the '--add' option in MAFFT[83]. Afterwards, we used
the alignments for scoring protein sequence conservation with Jensen-Shannon divergence[88].
Conservation scores were used to determine conservation percentiles of sites along protein
alignments for individual homolog groups. Rates of predicted synonymous to non-synonymous
novel SNVs were calculated for each conservation percentile across all homolog groups. For the
three SNVs predicted to encode for non-synonymous differences and also in the top five
percentile of conserved sites along the mycolic acid PKS, we inspected their codon contexts to
check that there were no additional variants which, in aggregate, might result in a synonymous
change (Table S14).

References
1.   Katz, L. & Baltz, R. H. Natural product discovery: past, present, and future. *J. Ind.*

Microbiol. Biotechnol. **43**, 155–176 (2016).

2.   Watve, M. G., Tickoo, R., Jog, M. M. & Bhole, B. D. How many antibiotics are produced by

the genus Streptomyces? *Arch. Microbiol.* **176**, 386–390 (2001).

3.   Hopwood, D. A. *Streptomyces in nature and medicine: The antibiotic makers*. (Oxford

University Press, 2007).

4. Chevrette, M. G. *et al.* The antimicrobial potential of Streptomyces from insect microbiomes.

Nat. Commun. **10**, 516 (2019).

5. Waterworth, S. C. *et al.* Horizontal Gene Transfer to a Defensive Symbiont with a Reduced

Genome in a Multipartite Beetle Microbiome. *MBio* **11**, (2020).

6. Drott, M. T. *et al.* Microevolution in the pansecondary metabolome of Aspergillus flavus and

its potential macroevolutionary implications for filamentous fungi. *Proc. Natl. Acad. Sci. U.*

S. A. **118**, (2021).

7. Steinke, K., Mohite, O. S., Weber, T. & Kovács, Á. T. Phylogenetic Distribution of

Secondary Metabolites in the Bacillus subtilis Species Complex. *mSystems* **6**, (2021).

8. Blin, K. *et al.* antiSMASH 6.0: improving cluster detection and comparison capabilities.

Nucleic Acids Res. **49**, W29–W35 (2021).

9. Navarro-Muñoz, J. C. *et al.* A computational framework to explore large-scale biosynthetic

diversity. *Nat. Chem. Biol.* **16**, 60–68 (2020).

10. Cimermancic, P. *et al.* Insights into secondary metabolism from a global analysis of

prokaryotic biosynthetic gene clusters. *Cell* **158**, 412–421 (2014).

11. Hannigan, G. D. *et al.* A deep learning genome-mining strategy for biosynthetic gene cluster

prediction. *Nucleic Acids Res.* **47**, e110 (2019).

12. Kautsar, S. A., van der Hooft, J. J. J., de Ridder, D. & Medema, M. H. BiG-SLiCE: A highly

scalable tool maps the diversity of 1.2 million biosynthetic gene clusters. *Gigascience* **10**,

(2021).

13. Emms, D. M. & Kelly, S. OrthoFinder: phylogenetic orthology inference for comparative

genomics. *Genome Biol.* **20**, 238 (2019).

14. Chase, A. B., Sweeney, D., Muskat, M. N., Guillén-Matus, D. G. & Jensen, P. R. Vertical

Inheritance Facilitates Interspecies Diversification in Biosynthetic Gene Clusters and

Specialized Metabolites. *MBio* **12**, e0270021 (2021).

15. Chevrette, M. G. *et al.* Evolutionary dynamics of natural product biosynthesis in bacteria.

Nat. Prod. Rep. **37**, 566–599 (2020).

16. Sélem-Mojica, N., Aguilar, C., Gutiérrez-García, K., Martínez-Guerrero, C. E. & Barona-

Gómez, F. EvoMining reveals the origin and fate of natural product biosynthetic enzymes.

Microb. Genom. **5**, (2019).

17. Felnagle, E. A., Rondon, M. R., Berti, A. D., Crosby, H. A. & Thomas, M. G. Identification

of the biosynthetic gene cluster and an additional gene for resistance to the antituberculosis

drug capreomycin. *Appl. Environ. Microbiol.* **73**, 4162–4170 (2007).

18. Del Carratore, F. *et al.* Computational identification of co-evolving multi-gene modules in

microbial biosynthetic gene clusters. *Commun Biol* **2**, 83 (2019).

19. Crits-Christoph, A., Bhattacharya, N., Olm, M. R., Song, Y. S. & Banfield, J. F. Transporter

genes in biosynthetic gene clusters predict metabolite characteristics and siderophore

activity. *Genome Res.* (2020) doi:10.1101/gr.268169.120.

20. Mungan, M. D. *et al.* ARTS 2.0: feature updates and expansion of the Antibiotic Resistant

Target Seeker for comparative genome mining. *Nucleic Acids Res.* **48**, W546–W552 (2020).

21. Crits-Christoph, A., Diamond, S., Butterfield, C. N., Thomas, B. C. & Banfield, J. F. Novel

soil bacteria possess diverse genes for secondary metabolite biosynthesis. *Nature* **558**, 440–

444 (2018).

22. Sugimoto, Y. *et al.* A metagenomic strategy for harnessing the chemical repertoire of the

human microbiome. *Science* **366**, (2019).

23. Andreu, V. P. *et al.* A systematic analysis of metabolic pathways in the human gut

microbiota. *bioRxiv* 2021.02.25.432841 (2021) doi:10.1101/2021.02.25.432841.

24. Pereira-Flores, E. *et al.* Mining metagenomes for natural product biosynthetic gene clusters:

unlocking new potential with ultrafast techniques. *bioRxiv* 2021.01.20.427441 (2021)

doi:10.1101/2021.01.20.427441.

25. Chevrette, M. G. & Handelsman, J. Needles in haystacks: reevaluating old paradigms for the

discovery of bacterial secondary metabolites. *Nat. Prod. Rep.* **38**, 2083–2099 (2021).

26. Zipperer, A. *et al.* Human commensals producing a novel antibiotic impair pathogen

colonization. *Nature* **535**, 511–516 (2016).

27. Claesen, J. *et al.* A Cutibacterium acnes antibiotic modulates human skin microbiota

composition in hair follicles. *Sci. Transl. Med.* **12**, (2020).

28. Nakatsuji, T. *et al.* Antimicrobials from human skin commensal bacteria protect against

Staphylococcus aureus and are deficient in atopic dermatitis. *Sci. Transl. Med.* **9**, (2017).

29. Wyatt, M. A. *et al.* Staphylococcus aureus nonribosomal peptide secondary metabolites

regulate virulence. *Science* **329**, 294–296 (2010).

30. Ridaura, V. K. *et al.* Contextual control of skin immunity and inflammation by

Corynebacterium. *J. Exp. Med.* **215**, 785–799 (2018).

31. Swaney, M. H. & Kalan, L. R. Living in Your Skin: Microbes, Molecules, and Mechanisms.

Infect. Immun. **89**, (2021).

32. Grice, E. A. *et al.* Topographical and temporal diversity of the human skin microbiome.

Science **324**, 1190–1192 (2009).

33. Oh, J. *et al.* Temporal Stability of the Human Skin Microbiome. *Cell* **165**, 854–866 (2016).

34. Byrd, A. L., Belkaid, Y. & Segre, J. A. The human skin microbiome. *Nat. Rev. Microbiol.*
16, 143–155 (2018).

35. Acosta, E. M. *et al.* Bacterial DNA on the skin surface overrepresents the viable skin
microbiome. *bioRxiv* 2021.08.16.455933 (2021) doi:10.1101/2021.08.16.455933.

36. Saheb Kashaf, S. *et al.* Integrating cultivation and metagenomics for a multi-kingdom view
of skin microbiome diversity and functions. *Nat Microbiol* **7**, 169–179 (2022).

37. Swaney, M. H., Sandstrom, S. & Kalan, L. R. Cobamide sharing drives skin microbiome
dynamics. *bioRxiv* 2020.12.02.407395 (2021) doi:10.1101/2020.12.02.407395.

38. van Dijk, L. R. *et al.* StrainGE: a toolkit to track and characterize low-abundance strains in
complex microbial communities. *Genome Biol.* **23**, 74 (2022).

39. Zimmermann, M. & Fischbach, M. A. A family of pyrazinone natural products from a
conserved nonribosomal peptide synthetase in Staphylococcus aureus. *Chem. Biol.* **17**, 925–
930 (2010).

40. Tauch, A. *et al.* Complete genome sequence and analysis of the multiresistant nosocomial
pathogen Corynebacterium jeikeium K411, a lipid-requiring bacterium of the human skin
flora. *J. Bacteriol.* **187**, 4671–4682 (2005).

41. Pelz, A. *et al.* Structure and biosynthesis of staphyloxanthin from Staphylococcus aureus. *J.*
Biol. Chem. **280**, 32493–32498 (2005).

42. Balibar, C. J., Shen, X. & Tao, J. The mevalonate pathway of Staphylococcus aureus. *J.*
Bacteriol. **191**, 851–861 (2009).

43. Grice, E. A. *et al.* A diversity profile of the human skin microbiota. *Genome Res.* **18**, 1043–
1050 (2008).

44. Clauditz, A., Resch, A., Wieland, K.-P., Peschel, A. & Götz, F. Staphyloxanthin plays a role

in the fitness of Staphylococcus aureus and its ability to cope with oxidative stress. *Infect.*

Immun. **74**, 4950–4953 (2006).

45. Seel, W. *et al.* Carotenoids are used as regulators for membrane fluidity by Staphylococcus

xylosus. *Sci. Rep.* **10**, 330 (2020).

46. Madhaiyan, M., Wirth, J. S. & Saravanan, V. S. Phylogenomic analyses of the

Staphylococcaceae family suggest the reclassification of five species within the genus

Staphylococcus as heterotypic synonyms, the promotion of five subspecies to novel species,

the taxonomic reassignment of five Staphylococcus species to Mammaliicoccus gen. nov.,

and the formal assignment of Nosocomiicoccus to the family Staphylococcaceae. *Int. J. Syst.*

Evol. Microbiol. **70**, 5926–5936 (2020).

47. Dong, P.-T. *et al.* Photolysis of Staphyloxanthin in Methicillin-Resistant Staphylococcus

aureus Potentiates Killing by Reactive Oxygen Species. *Adv. Sci.* **6**, 1900030 (2019).

48. Jain, C., Rodriguez-R, L. M., Phillippy, A. M., Konstantinidis, K. T. & Aluru, S. High

throughput ANI analysis of 90K prokaryotic genomes reveals clear species boundaries. *Nat.*

Commun. **9**, 1–8 (2018).

49. Li, D., Liu, C.-M., Luo, R., Sadakane, K. & Lam, T.-W. MEGAHIT: an ultra-fast single-

node solution for large and complex metagenomics assembly via succinct de Bruijn graph.

Bioinformatics **31**, 1674–1676 (2015).

50. Gande, R. *et al.* Acyl-CoA carboxylases (accD2 and accD3), together with a unique

polyketide synthase (Cg-pks), are key to mycolic acid biosynthesis in Corynebacterianeae

such as Corynebacterium glutamicum and Mycobacterium tuberculosis. *J. Biol. Chem.* **279**,

44847–44857 (2004).

51. Portevin, D. *et al.* A polyketide synthase catalyzes the last condensation step of mycolic acid

biosynthesis in mycobacteria and related organisms. *Proc. Natl. Acad. Sci. U. S. A.* **101**, 314–

319 (2004).

52. Chen, Y. *et al.* Structural classification and properties of ketoacyl synthases. *Protein Sci.* **20**,

1659–1667 (2011).

53. Klaus, M., Buyachuihan, L. & Grininger, M. Ketosynthase Domain Constrains the Design of

Polyketide Synthases. *ACS Chem. Biol.* **15**, 2422–2432 (2020).

54. Medema, M. H., Cimermancic, P., Sali, A., Takano, E. & Fischbach, M. A. A systematic

computational analysis of biosynthetic gene cluster evolution: lessons for engineering

biosynthesis. *PLoS Comput. Biol.* **10**, e1004016 (2014).

55. Lee, M. D. GToTree: a user-friendly workflow for phylogenomics. *Bioinformatics* **35**, 4162–

4164 (2019).

56. Garrison, E. *et al.* Variation graph toolkit improves read mapping by representing genetic

variation in the reference. *Nat. Biotechnol.* **36**, 875–879 (2018).

57. Kloos, W. E. & Schleifer, K. H. Isolation and Characterization of Staphylococci from Human

Skin II. Descriptions of Four New Species: Staphylococcus warneri, Staphylococcus capitis,

Staphylococcus hominis, and Staphylococcus simulans. *Int. J. Syst. Bacteriol.* **25**, 62–79

(1975).

58. Becker, K., Heilmann, C. & Peters, G. Coagulase-negative staphylococci. *Clin. Microbiol.*

Rev. **27**, 870–926 (2014).

59. Liu, C.-I. *et al.* A cholesterol biosynthesis inhibitor blocks Staphylococcus aureus virulence.

Science **319**, 1391–1394 (2008).

60. Blount, Z. D., Barrick, J. E., Davidson, C. J. & Lenski, R. E. Genomic analysis of a key

innovation in an experimental Escherichia coli population. *Nature* **489**, 513–518 (2012).

61. Ghosh, S. & O'Connor, T. J. Beyond Paralogs: The Multiple Layers of Redundancy in

Bacterial Pathogenesis. *Front. Cell. Infect. Microbiol.* **7**, 467 (2017).

62. Otto, M. Staphylococcus epidermidis--the "accidental" pathogen. *Nat. Rev. Microbiol.* **7**,

555–567 (2009).

63. Stokes, J. M. *et al.* A Deep Learning Approach to Antibiotic Discovery. *Cell* **180**, 688-

702.e13 (2020).

64. Melo, M. C. R., Maasch, J. R. M. A. & de la Fuente-Nunez, C. Accelerating antibiotic

discovery through artificial intelligence. *Commun Biol* **4**, 1050 (2021).

65. Carroll, L. M. *et al.* Accurate de novo identification of biosynthetic gene clusters with

GECCO. *bioRxiv* 2021.05.03.442509 (2021) doi:10.1101/2021.05.03.442509.

66. Buchfink, B., Xie, C. & Huson, D. H. Fast and sensitive protein alignment using

DIAMOND. *Nat. Methods* **12**, 59–60 (2014).

67. Seemann, T. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* **30**, 2068–2069

(2014).

68. Parks, D. *CompareM: A toolbox for comparative genomics*. (Github).

69. Hyatt, D. *et al.* Prodigal: prokaryotic gene recognition and translation initiation site

identification. *BMC Bioinformatics* **11**, 119 (2010).

70. Aramaki, T. *et al.* KofamKOALA: KEGG Ortholog assignment based on profile HMM and

adaptive score threshold. *Bioinformatics* **36**, 2251–2252 (2019).

71. Tatusova, T. *et al.* NCBI prokaryotic genome annotation pipeline. *Nucleic Acids Res.* **44**,

6614–6624 (2016).

72. Parks, D. H. *et al.* GTDB: an ongoing census of bacterial and archaeal diversity through a

phylogenetically consistent, rank normalized and complete genome-based taxonomy. *Nucleic*

Acids Res. (2021) doi:10.1093/nar/gkab776.

73. Chaumeil, P.-A., Mussig, A. J., Hugenholtz, P. & Parks, D. H. GTDB-Tk: a toolkit to

classify genomes with the Genome Taxonomy Database. *Bioinformatics* **36**, 1925–1927

(2019).

74. Ondov, B. D. *et al.* Mash: fast genome and metagenome distance estimation using MinHash.

Genome Biol. **17**, 132 (2016).

75. Chen, S., Zhou, Y., Chen, Y. & Gu, J. fastp: an ultra-fast all-in-one FASTQ preprocessor.

Bioinformatics **34**, i884–i890 (2018).

76. Wick, R. R., Judd, L. M., Gorrie, C. L. & Holt, K. E. Unicycler: Resolving bacterial genome

assemblies from short and long sequencing reads. *PLoS Comput. Biol.* **13**, e1005595 (2017).

77. Krueger, F., James, F., Ewels, P., Afyounian, E. & Schuster-Boeckler, B.

FelixKrueger/TrimGalore: v0.6.7 - DOI via Zenodo. (2021). doi:10.5281/zenodo.5127899.

78. Walker, B. J. *et al.* Pilon: an integrated tool for comprehensive microbial variant detection

and genome assembly improvement. *PLoS One* **9**, e112963 (2014).

79. Li, H. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* **34**, 3094–

3100 (2018).

80. Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows–Wheeler

transform. *Bioinformatics* **25**, 1754–1760 (2009).

81. Paradis, E., Claude, J. & Strimmer, K. APE: Analyses of Phylogenetics and Evolution in R

language. *Bioinformatics* **20**, 289–290 (2004).

82. Hug, L. A. *et al.* A new view of the tree of life. *Nature Microbiology* **1**, 1–6 (2016).

83. Katoh, K. & Standley, D. M. MAFFT multiple sequence alignment software version 7:

improvements in performance and usability. *Mol. Biol. Evol.* **30**, 772–780 (2013).

84. Suyama, M., Torrents, D. & Bork, P. PAL2NAL: robust conversion of protein sequence

alignments into the corresponding codon alignments. *Nucleic Acids Res.* **34**, W609-12

(2006).

85. Stamatakis, A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large

phylogenies. *Bioinformatics* **30**, 1312–1313 (2014).

86. Kania, S. A. Reclassification of Staphylococcus schleiferi by Madhaiyan et al. lacks key

supporting data. *Int. J. Syst. Evol. Microbiol.* **72**, (2022).

87. Wood, D. E., Lu, J. & Langmead, B. Improved metagenomic analysis with Kraken 2.

Genome Biol. **20**, 1–13 (2019).

88. Capra, J. A. & Singh, M. Predicting functionally important residues from sequence

conservation. *Bioinformatics* **23**, 1875–1882 (2007).

Figure 1: *Isa*BGC offers an efficient and sensitive means for GCF homology detection. a) A schematic overview of the full *Isa*BGC suite. **b**) Neighbor-joining trees were constructed for each of the four genera based on pairwise MASH ANI estimates between genomes and heatmaps showcase the presence of GCFs found in ≥ 5% of genomes. Red dots on neighbor-joining trees signify complete or chromosomal genomes used for initial identification of BGCs. Heatmap cell color indicates the BGC class as predicted by antiSMASH. **c**) Results from benchmarking *Isa*BGC-AutoExpansion sensitivity on 14 completed genomes for the species of *M. luteus*. AntiSMASH and *Isa*BGC-AutoExpansion were run on artificially fragmented genomes and resulting BGC predictions were compared to initial BGCs identified on the unfragmented, complete genomes. **d**) The relationship between the assembly quality measured by N50 and the total number of homolog groups detected across 14,978 *Staphylococcus* draft genomes is shown. The coloring corresponds to the percentage of BGC homolog groups co-located in a segment with a homolog group commonly found in protocore regions of BGC predictions by antiSMASH run on completed genomes.

Figure 2: *lsa*BGC reveals new insight about characterized and unstudied BGCs from skin-associated species. a) A maximum-likelihood phylogeny was constructed from PznA and PznB showing the relationship between GCFs encoding for pyrazinones. Branch color corresponds to different GCFs and the outer circular color denotes the species classification for the genome from which the sequences were gathered. **b**) Homolog groups across the seven pyrazinone encoding GCFs are depicted as a network where edges indicate that two homolog groups co-occur in the same GCF. Homolog group color of nodes depict distinct GCFs. **c**) Phylogeny constructed from ribosomal proteins showing the conservation of *Corynebacterium* GCF-50 across the diverse species encoding the NRPS using results from *lsa*BGC-PopGene analysis. The gene schematics show the consensus gene order for the GCF. Gene lengths correspond to the median length across all BGC instances and only genes found in two or more genomes are shown. For each species, the gene is colored based on the median similarity of gene sequences from the species to the consensus sequence across all species. The barplot displays the total number of GCF instances identified per species across the full set of *Corynebacterium* genomes. Broad annotations for homolog groups are shown as a color strip. Below it are bar plots corresponding to the conservation of homolog groups across samples and their median Beta-RD. **d**) A histogram of the amino acid identities between GCF-50 NRP synthase instances which are complete (excluding partial instances due to assembly fragmentation included in panel **c**) is compared to the average amino acid identities (AAI) between genomes they were gathered from.

Figure 3: Staphyloxanthin is ubiquitous across *Staphylococcus* and found on plasmids. a) A maximum-likelihood phylogeny of the full *Staphylococcus* genus was constructed from ribosomal protein encoding genes. The clade in green represents the newly recognized genus of *Mammaliicoccus*. Heatmaps showcase the presence of the five *crt* genes for each species for five separate GCFs containing the *crt* operon. The shading indicates the proportion of a species found to carry homolog groups within a particular GCF context in log₁₀ scale. **b)** A maximum-likelihood phylogeny was built from CrtM and CrtN protein alignments. Branch colors represent the GCF classification of the CrtMN sequences and bootstrap values are shown for key nodes where GCFs partition. The species the CrtMN sequence was extracted from is shown as a color strip followed by a bar plot depicting the comparative codon usage cosine distance (CCU-cd). CCU-cd was only calculated for the full five gene *crt* operon and represents the codon frequency dissimilarity with the codon frequency of the background genome. The heatmap showcases the genome-wide average amino acid identity (AAI) between pairs of genomes from which CrtMN sequences were gathered. **c)** AAI and shared homolog group content were calculated between pairs of GCF instances from different genomes for GCF-3 and GCF-6 individually. The coloring represents genome-wide AAI similarity. **d)** A schematic of the completed genome for *S. equorum* C2014, which was predicted to feature GCF-6 in three distinct copies. Bottom panel shows the BGC alignment between the three instances of GCF-6 found within this genome. **e)** Representative isolates with and without staphyloxanthin encoding GCFs were grown on 20% BHI agar to visualize pigmentation production. **f)** Methanol extractions and wavelength absorption analysis were performed to identify signature peaks in the 400 to 500 nm range associated with the presence of staphyloxanthin in *S. aureus*[39,44]. Spectra are normalized to cell density.

Figure 4: Mining for microdiversity within the cutimycin BGC leads to genetic insights for clade III *C. acnes*. a) Codon usage dissimilarity between BGC regions and background genomes shown for six common GCFs in *C. acnes*. **b**) The cutimycin encoding GCF is found primarily in clade I (subclade IB; orange) and clade III (tan) *C. acnes*. **c**) A schematic of the cutimycin encoding GCF. Lines in black correspond to sites of nearly-fixed differences between clades of *C. acnes*. **d**) *Isa*BGC-DiscoVary was used to identify the presence of the cutimycin GCF in 270 skin metagenomes. The proportion of alleles representative of clade IB or clade III *C. acnes* at sites of nearly fixed-differences was quantified. High-resolution strain quantification within metagenomes was similarly performed with StrainGE and used to overlay clade-specific typing results from *Isa*BGC-DiscoVary analysis. **e**) The distribution of novel SNVs identified by *Isa*BGC-DiscoVary within metagenomes, post filtering for removal of false positives, is shown across homolog groups and subjects. **f**) The location of novel SNVs is shown along the lantibiotic dehydratase N-terminal encoding gene (*HG2417*) for each subject. Coloring of the novel SNVs corresponds to which body-site(s) they were identified within. **g**) The clade distribution of reference genes on which novel SNVs were identified across the cutimycin GCF for subject S012 are shown individually for the predicted transitions or transversions. **i**) The proportion of differences which corresponded to cytosine to thymine or thymine to cytosine transitions along a core-genome alignment was calculated between pairs of *C. acnes* and categorized by clade comparisons. **j**) Comparative analysis of presumed directional deaminations from *C. namnetense*, treated as ancestral, to *C. acnes* clade IB and clade III were estimated using a multi-iteration sub-sampling based approach for sites along the core-genome alignment, whereby a greater number of suspected C to T transitions were estimated for clade III *C. acnes*. **k**) The presence of a complete uracil DNA glycosylase (*ung*), from clade I *C. acnes*, is shown across available *Cutibacterium* genomes.

Figure 5: Thousands of novel SNVs identified within BGCs from the *C. tuberculostearicum* species complex. a) Assessment of novel SNVs identified by *Isa*BGC-DiscoVary, post-filtering, for each BGC from the *C. tuberculostearicum* species complex for presence within metagenomic assemblies. **b**) The number of novel SNVs is shown for ranges of conservation percentile across all homolog groups and metagenomes. The coloring corresponds to the ratio of synonymous to non-synonymous novel SNVs within each conservation range. **c**) Novel SNVs are shown for the polyketide synthase involved in mycolic acid biosynthesis. The top panel showcases the non-synonymous (up-facing) and synonymous (down-facing) novel SNVs found along the T1PKS gene, with heights corresponding to the number of metagenomes with the novel SNV. The middle panel shows the conservation scores colored by percentile ranges (lightest = least conserved sites; darkest = most conserved sites). Key domains common to T1PKS are shown underneath along with the location of predicted non-synonymous mutations within the top 5% of conserved sites for the gene. Two adjacent SNVs affecting the coding of the residue at position 343 were identified and these could result in a combined effect of encoding for alanine.