*lsa*BGC provides a comprehensive framework for evolutionary analysis of biosynthetic gene clusters within focal taxa

Rauf Salamzade[1], J.Z. Alex Cheong[1], Shelby Sandstrom[1], Mary Hannah Swaney[1], Nicole Lane Starr[1], Anne Marie Singh[2], and Lindsay R. Kalan[1,3]

[1]Department of Medical Microbiology and Immunology, School of Medicine and Public Health, University of Wisconsin, Madison, WI, USA
[2]Department of Pediatrics, School of Medicine and Public Health, University of Wisconsin, Madison, WI, USA
[3]Department of Medicine, School of Medicine and Public Health, University of Wisconsin, Madison, WI, USA

Address for Correspondence: Lindsay Kalan, 1550 Linden Dr, 6155 Microbial Sciences Building, Madison, WI, USA, 53706. lkalan@wisc.edu

Abstract:

Bacterial secondary metabolites, encoded by biosynthetic gene clusters (BGCs), can underlie microbiome homeostasis and be developed into high-value commercialized products, which have historically been mined from a select group of taxa. Despite this, dedicated bioinformatics tools for comparative analysis of BGCs are limited. We have thus developed *lsa*BGC to aid exploration of microdiversity and evolutionary trends across BGCs in any bacterial taxa of interest through calculation of population genetic statistics and metagenomic exploration. As a proof of concept, we applied *lsa*BGC to four genera commonly found in skin microbiomes to reveal novel insights on both well and less studied taxa and their secondary metabolites. We find evidence of horizontal gene transfer (HGT) within multiple genera. The BGC encoding for the carotenoid staphyloxanthin, a virulence factor in *S. aureus*, is ubiquitous across the genus of *Staphylococcus* and exhibits signatures of inter-species HGT, including mobilization onto plasmids. We also identified a non-ribosomal peptide synthetase (NRPS) BGC flanked by transposable elements, encoding for a predicted siderophore, which is highly conserved across several divergent *Corynebacterium* species commonly isolated from skin such as *C. kefirresidentii*. From strain-resolution metagenomics, we further determined that the *C. kefirresidentii* species complex is the most prevalent clade of *Corynebacterium* in skin metagenomes and is found in the skin microbiome of every participant surveyed. While ubiquitous across skin, the number of publicly-available genomes for this species complex is currently limited. Using *lsa*BGC's framework for metagenomic mining, we identified 34,545 novel single nucleotide variants (SNVs) in predicted BGCs of the species complex, including several novel, non-synonymous SNVs within highly conserved sites of enzymes. Ultimately, we envision that application of *lsa*BGC can highlight important, yet poorly studied, secondary metabolites as well as aid in the discovery and design of more efficacious commercialized metabolites.

Introduction:

The secondary metabolome of bacteria has served as a valuable reservoir of natural products with great societal benefit[1]. Historically, many commercialized secondary metabolites, such as antibiotics, have been identified from particular taxonomic groups, such as the genus of *Streptomyces* within Actinomycetota[2,3]. Several studies have explored the secondary metabolome of these metabolically rich microbial taxa to develop a comprehensive understanding of their chemical diversity and unique traits[4–7]. Most of these studies begin by first identifying biosynthetic gene clusters within genomes using the popular antiSMASH software[8] and then group BGCs into gene cluster families (GCFs) using BiG-SCAPE[9]. Critically, this approach, in addition to other bioinformatics software for genomics annotation and analysis of BGCs[10–12], uses a set of key protein domains as their base to group BGCs predicted to encode similar metabolites. However, when working in the context of a single genus or species, it becomes more appropriate to use full protein sequences which enables the use of advanced algorithms for the identification of orthologs and paralogs[13] to reliably determine evolutionary relationships between BGCs.

Recent studies which explored the vertical descent and conservation of BGCs across the genus of *Salinospora* and the fungal species *Aspergillus flavus* have shown that intra-genus or intra-species evolution can result in chemical and regulatory diversification of the encoded metabolites[6,14]. Thus, improved understanding of the evolutionary emergence of BGCs could be informative towards elucidating the function of individual genes and encoded metabolites[9,15–17].

Due to difficulties in the cultivation of many bacterial taxa, several recent studies have begun to explore metagenomic datasets to unearth novel secondary metabolites[18–21]. While most of these endeavors have relied on performing metagenomic assembly followed by BGC identification using software such as antiSMASH, recently, two read-based approaches were described to reliably and sensitively search for key BGC domains in metagenomes[20,22]. This highlights a larger trend in which these approaches primarily aim to identify highly novel secondary metabolites and thus prioritize investigation of poorly studied or newly discovered taxa[23]. However, provided that most metabolites have been mined from select genera, mostly within the *Actinomycetia* class[2], there could be tremendous reward for efforts to advance our understanding of cataloged but uncharacterized BGCs from well studied taxa. This is especially important because many species within the metabolically rich *Streptomyces* only have limited representative genomes, thus methods to leverage metagenomic datasets and identify homologous instances of rare BGCs can lead to critical insight into their function and ecological distribution[4].

To aid the advancement of such taxa-specific analyses of BGCs, we introduce the comprehensive software suite *Lineage Specific Analysis* of BGCs (*lsa*BGC; https://github.com/Kalan-Lab/lsaBGC), which consists of several programs for comparative genetics of BGCs as well as metagenomic mining. We showcase the utility of *lsa*BGC in this study through application to four major genera which are commonly found in healthy human skin microbiomes. Several secondary metabolites which function as antibiotics[24–26], virulence

factors[27], or in microbe-host interactions[28,29] have been identified in bacteria from skin
microbiomes, primarily from staphylococci. Human skin is also easily accessible to sampling
and most species can be cultivated for follow up studies. Using *Isa*BGC, we reveal several new
ecological and evolutionary insights for both well characterized and unknown predicted
metabolites, including signatures of inter-species transfer for the virulence-associated
staphylococcal carotenoid staphyloxanthin and a predicted siderophore found to be highly
conserved among divergent *Corynebacterium* species. Additionally, through mining skin
metagenomic datasets, we find novel coding variants within the functionally important ketide
synthase (KS) domain of the polyketide synthase (PKS) responsible for mycolic acid
biosynthesis, a defining cell wall component of *Corynebacterium* species implicated in immune
recognition.
82
Results:

***Isa*BGC allows for sensitive and systematic identification of BGC homolog groups**

The *Isa*BGC suite consists of seven core programs, three workflow programs, and
several additional scripts for analyses (Figure 1, S1; Supplementary Methods and Results).
These programs allow for clustering BGCs into GCFs, refining boundaries of BGCs, and
sensitive, yet specific, rapid detection of GCF instances in draft-quality assemblies. The suite
additionally includes dedicated tools for GCF visualization, evolutionary and population genetic
analysis of individual genes within GCFs, and base-resolution identification of novel SNVs in
complex metagenomes.

To demonstrate the capabilities of the *Isa*BGC suite we applied it to four genera which
are stable and integral members of the human skin microbiome[30–34]. These included three
genera in the Actinomycetota (formerly Actinobacteria) phylum: *Corynebacterium*,
Cutibacterium, and *Micrococcus* as well as the Bacillota (formerly Firmicutes) genus
Staphylococcus. *Isa*BGC was also independently applied to the species or species-complex
from each genus most commonly represented in skin microbiomes. These were the *C.*
kefirresidentii species-complex, *C. acnes*, *M. luteus*, and *S. epidermidis* (Table S1).

We used *Isa*BGC-AutoProcess to run gene-calling, BGC annotation, and *de novo*
homolog group delineation with complete or chromosomally complete genomes for each genus
or species. We then applied *Isa*BGC-Cluster to group homologous BGCs into GCFs and
systematically searched for instances of GCFs using *Isa*BGC-AutoExpansion (Figure 1b; Table
S2, S3; Supplementary Methods and Results). To assess the validity and sensitivity of our
approach, two benchmarking tests using BGCs identified in *M. luteus* were performed (Figure
1c, S3; Supplementary Methods and Results) showing that the *Isa*BGC-AutoExpansion
framework leads to greater sensitivity compared to running *de novo* antiSMASH on fragmented
draft-quality assemblies. This increased sensitivity was due to more robust, yet reliable,
detection of auxiliary genes located on the edges of scaffolds not containing core domains used
by antiSMASH for BGC detection. In addition, *Isa*BGC-AutoExpansion is highly efficient and
allowed us to search for 63 GCFs identified from high-quality *Staphylococcus* genomes in

14,978 draft-quality genomes in approximately 14 hours using 40 cores and less than 150 Gb of
memory. We found a positive association between detection of complete BGC instances and
assembly quality (Figure 1d; Supplementary Methods and Results).
118
119 **_Isa_BGC enables reliable comparative genomics to explore evolutionary trends of BGCs**
120
The increased sensitivity of _Isa_BGC to uniformly detect auxiliary components of BGCs in
each genome enabled us to perform reliable comparative genetic analyses on them. We
focused on two of the four genera with the greatest breadth of diversity, _Corynebacterium_ and
Staphylococcus, and selected a subset of 456 and 229 representative genomes from each
genera, respectively (Table S1). To better understand which species in these genera were most
relevant to skin health, we first profiled how often distinct species were found among healthy
skin microbiomes using a large, recently generated metagenomics dataset[35,36] (Table S4). Skin-
associated species (found in more than 10 of 270 metagenomes) were phylogenetically
clustered for both genera, within the _S. epidermidis_ clade (Figure S4a) and within the _C._
kefirresidentii species complex (Figure S4b; Supplementary Methods and Results).
131
Association of individual homolog groups from these GCFs revealed several were
enriched within skin-associated staphylococci, but only two were enriched for skin-associated
corynebacteria, both encoding hypothetical proteins (Table S5). Nearly half of the homolog
groups enriched for skin-associated staphylococci (5 of 12; 41.7%) were related to GCF-10,
predicted to encode for an NRP synthetase corresponding to pyrazinone biosynthesis,
molecules reported to regulate virulence in _S. aureus_[27,37]. Staphylococcal pyrazinones were
previously identified as being a self-contained BGC consisting of two genes, _pznA_ and _pznB_,
specific to species commonly associated with human skin, including _S. aureus_, _S. epidermidis_,
and _S. capitis_[27,37]. Using _Isa_BGC, we discover novel variants of pyrazinones, which encapsulate
seven distinct GCFs (Figure 2a), that belong to species not commonly found on human skin,
such as _S. felis_ (GCF-56), the dominant species of _Staphylococcus_ on cat skin. While speciation
had resulted in orthologous copies of _pznA_ and _pznB_ being highly diverged in sequence, it is
surprising to find that the surrounding genomic context of the two genes is highly variable for all
seven GCFs (Figure 2b), suggesting their ancestral mobilization or relocation within or between
genomes.
147
Application of _Isa_BGC-Divergence followed by Bayesian shrinkage analysis was then
used to systematically assess whether pyrazinones or other _Staphylococcus_ and
Corynebacterium GCFs exhibited indications of HGT between species, as suggested by high
Beta-RD values (Figure S4cd; Table S6; Supplementary Methods and Results). Despite the
small size of pyrazinones, we do not find support for the frequent mobilization of any of the
seven GCFs encoding them across staphylococcal species. However, high Beta-RD posterior
ranges were observed for two NRPS associated GCFs from _Corynebacterium_. One of these
GCFs, predicted to encode for an NRPS-dependent siderophore[38], was found in several species
commonly isolated from skin[35], including _C. kefirresidentii_. Using results from _Isa_BGC-PopGene
for this GCF, which includes determination of a consensus gene order across BGC instances,
we find that the NRPS is highly conserved in sequence across species and found flanked by

mobile genetic elements, as previously reported in *C. jeikeium*[38] (Figure 2cd). The taxonomic
breadth and high sequence conservation of this GCF suggest that it has undergone recent HGT
between distantly related species.
162
We next used *lsa*BGC-PopGene to identify genes under sweeping or balancing selection
by comprehensively calculating Tajima's D for each homolog group within a GCF across all four
genera, including *Micrococcus* and *Cutibacterium* (Table S7). We found that a hybrid
terpene/T3PKS GCF, in *Staphylococcus* exhibits the greatest signal for intra-species sweeping
selection and is one of four GCFs observed to be ancestral to the *S. epidermidis* clade (Figure
S5; Supplementary Methods and Results). The terpene component of this hybrid GCF encodes
for staphyloxanthin carotenoid production[39] and manual inspection determined that the T3PKS
prediction actually corresponds to hydroxymethylglutaryl-CoA (HMG-CoA) synthase, which is
involved in the mevalonate pathway for isoprenoid synthesis and is homologous to KS domains
of polyketide synthases[40]. In addition to these two BGC predictions, the regions in between and
flanking the staphyloxanthin encoding BGC and HMG-CoA synthase include several genes
important to staphylococcal virulence, some of which were found in multiple copies in certain
species (Figure S6ab). Further, while the five *crt* operon genes encoding for staphyloxanthin
were found in multiple species within the *S. epidermidis/aureus* clade, they are particularly
conserved in *S. aureus* as indicated by low Tajima's D values (Figure S6b).
178
179 **The virulence-associated carotenoid staphyloxanthin is ubiquitous across the**
180 ***Staphylococcus* genus**
181
Although staphyloxanthin is well characterized as an *S. aureus* virulence factor[41], we
were unable to identify a report of its distribution across the *Staphylococcus* genus, with only a
single study, to our knowledge, suggesting its production in another species[42]. *lsa*BGC analysis
revealed a total of five staphylococcal GCFs encoding for staphyloxanthin. Following GCF-3, the
second most prevalent GCF encoding staphyloxanthin, GCF-6, had a high Beta-RD distribution
(Figure S4c) and was widely distributed across divergent species in the genus (Figure 3a), as
well as *Mammaliicoccus,* the closest phylogenetic neighbors of staphylococci[43]. Constructing a
phylogeny based on a concatenated alignment of *crtM* and *crtN* revealed the appropriate
delineation of GCF-3 and GCF-6 despite their co-occurrence in the genomes of particular
species, such as *S. warneri* (Figure 3b, S7a). Additional analyses showed that GCF-6 has
mobilized across species and can be found on plasmids (Figure 3bcd).
193
To assess whether carriage of the *crt* operon translates to staphyloxanthin production in
non-aureus species, we identified staphylococcal isolates cultivated from human skin with
whole-genome sequences generated by our lab that encode a staphyloxanthin related GCF. We
identified a rare instance of an *S. epidermidis* isolate with GCF-3 (only 1-3% of *S. epidermidis*
genomes encode GCF-3), strain LK1136, and an *S. warneri* isolate with both GCF-3 and GCF-
6, strain LK413. While we observe GCF-3 in all *S. warneri* genomes, the GCF is degraded
encoding only the dehydrosqualene synthase gene, *crtM*, and the dehydrosqualene desaturase
gene, *crtN* (Figure 3a; Table S8). However, roughly half of the available *S. warneri* genomes
(47.5%) encode GCF-6 in full and thus have two copies of *crtM* and *crtN*. We used long-read

sequencing to generate near-complete genomes for *S. epidermidis* LK1136 and *S. warneri* LK413 and found that both strains encode GCF-3 within their chromosomes while GCF-6 is encoded on a plasmid within *S. warneri*. (Figure S7b; Supplementary Methods and Results).

To check for phenotypic effects in strains carrying different staphyloxanthin GCFs, we assessed pigmentation levels in strains of *S. aureus*, *S. epidermidis*, and *S. warneri* with and without full *crt* operons (Table S9). Cells of *S. epidermidis* encoding GCF-3 are pigmented compared to strains without the cluster (Figure 3e-f). Cells of *S. warneri* LK413 containing the complete *crt* operon (GCF-6) and additional copies of *crtMN* displayed the strongest golden pigmentation, confirmed as staphyloxanthin by spectrophotometry[39,44] (Figure 3f).

_Isa_BGC provides a framework for mining for base-resolution novelty in BGCs

We have demonstrated how the *Isa*BGC suite can be used to identify homologous BGC instances into GCFs and subsequently uncover their evolutionary trends and phylogenetic distribution. With increasing metagenomic datasets becoming available, often from complex microbiomes, we developed *Isa*BGC-DiscoVary to directly profile BGCs and rapidly identify novel variation within GCF related genes from metagenomes (Figure S2; Supplementary Methods and Results). The *Isa*BGC framework thus allows users to first identify all instances of GCFs in genomic assemblies to build a comprehensive database of known alleles for GCF genes and subsequently apply *Isa*BGC-DiscoVary to find novel SNVs within genes which are absent in the available assemblies. Further, because *Isa*BGC-DiscoVary extracts the subset of reads supporting the presence of novel SNVs, exhaustive methods to assess the validity of metagenomic SNV calling, which would be computationally intensive and impractical to perform for the full set of reads from a metagenome, are made accessible for further validation of novel SNVs.

We first assessed and found high concordance (>96% overlap) between novel SNVs reported by *Isa*BGC-DiscoVary compared to an assembly-based approach using whole-genome sequencing readsets for 132 *M. luteus* isolates to benchmark it's sensitivity and specificity (Figure S8; Table S10, S11; Supplementary Methods and Results). Next, to demonstrate *Isa*BGC-DiscoVary's application in metagenomic datasets, its primary intended usage, we investigated the microdiversity of BGCs from *C. acnes*, the most abundant species at sebaceous sites, within healthy skin metagenomes generated by our lab[35]. We found that cutimycin, an anti-Staphylococcal thiopeptide, exhibited genomic signatures of recent acquisition in *C. acnes* and likely has since been lost in most isolates besides those belonging to clade IB and clade III (Figure 4ab, S9, S10a; Supplementary Methods and Results). Extensive metagenomic analysis demonstrated the reliability of *Isa*BGC-DiscoVary to type clade-specific alleles of genes within the cutimycin BGC in metagenomic datasets which correlated with strain presence inferences (Figure 4cd, S10bc; Table S12). We used *Isa*BGC to further investigate novel SNVs detected within cutimycin genes and found that many were from the metagenomes of a single individual and belonged to clade III *C. acnes* (Figure 4efgh). Through observation that the majority of novel SNVs from this individual also corresponded to cytosine to thymine changes, we discovered that clade III *C. acnes* have elevated rates of

246 cytosine deamination, which coincides with the clade's carriage of a truncated uracil DNA
247 glycosylase, the enzyme responsible for the repair of such deaminations (Figure 4ijk).
248
249 **_lsa_BGC identifies novel SNVs within polyketide synthase catalytic domains**
250
We determined the _C. kefirresidentii_ species complex is the most broadly distributed
Corynebacterium across our skin metagenomes (Figure S4b); however, only 22 genomic
assemblies, representing five species, were available for the complex at the time of this
analysis. This clade thus represents an ideal taxonomic group to examine novel BGC
microdiversity using _lsa_BGC-DiscoVary. We ran _lsa_BGC-DiscoVary systematically on six GCFs
identified in the species complex, largely encoding for unknown metabolites, and found 40,019
putatively novel SNV instances across 105 homolog groups and 109 metagenomic samples. Of
these, 5,474 instances were filtered because reads supporting their presence aligned with a
higher score to other regions in a concatenated database of all _Corynebacterium_ genomes from
GTDB as compared to the reference database used for _lsa_BGC-DiscoVary. While ubiquitous, _C._
kefirresidentii species complex members are present at very low abundances in some body-
sites where metagenomic assembly would struggle to construct their genomes (Figure S11a).
We performed sample-specific metagenomic assemblies[45] and found that at least 11,659 novel
SNV instances detected by _lsa_BGC-DiscoVary would be missed by assembly-based
investigation (Figure 5a; Supplementary Methods and Results).
266
To further investigate the novel SNVs detected, we focused on 68 homolog groups
overlapping or in close proximity to protocore regions of BGCs and further filtered based on
standardized coverage metrics, retaining 5,802 instances of 2,343 unique novel SNVs. Novel
SNVs were more likely to correspond to synonymous changes if they were found in multiple
metagenomic samples (Figure S11b). Further, metagenomic samples from the same subject at
different body sites and from the same body sites across different subjects were more likely to
share novel SNVs when compared to metagenomic samples from different body sites and
different subjects (p<1E-8; two-sided Wilcoxon rank sum test) (Figure S11c). We additionally
found that homolog groups for which _lsa_BGC-PopGene estimated lower Tajima's D values,
suggesting the presence of sweeping selection, tended to have fewer novel SNV counts (Figure
5b).
278
Finally, we expected and observed that the ratio of synonymous to non-synonymous
novel SNVs increases at deeply conserved sites along homolog groups (Figure 5c; Table S13).
The homolog group with the greatest number of novel SNVs, and novel, non-synonymous SNVs
within highly conserved regions, was the PKS gene responsible for synthesis of mycolic acid, a
major component of the cell wall in most _Corynebacterium_ species recently shown to have
immunomodulatory effects on the skin[28,46,47]. We identified five non-synonymous SNVs along
conserved sites of the PKS gene, including three within the ketoacyl synthase domain, which
catalyzes condensation[48], and could alter reaction kinetics[49] or scaffold structure[50] (Figure 5d,
S12).
288
Discussion:

The *Isa*BGC suite packages several important, as well as novel, functionalities for large-
scale evolutionary analysis of BGCs from any taxa in a structured framework. In particular we
developed several algorithms to overcome challenges when working with draft-quality genomic
assemblies. The suite's metagenomic mining functionalities further provide valuable insight into
the presence and sequence diversity of a specific taxa's BGCs across diverse microbiomes.
This is a particularly important feature for rare BGCs and taxa with a limited numbers of
genomic assemblies available, such as taxa that are difficult to cultivate from complex
metagenomes. A current limitation of the *Isa*BGC framework includes its basis for using
chromosomally complete genomes to perform initial identification of BGCs and GCFs, which
can miss rare BGCs only observed in draft genomes. To alleviate this limitation, future
development of the suite will aim to allow direct incorporation of cataloged BGC and GCF
instances from publicly available databases or software[12,51–53]. We may also consider the use of
pan-genome alignment approaches to potentially increase the efficiency of *Isa*BGC for
metagenomic identification of novel variants within BGC genes[54,55].

Here, we apply *Isa*BGC's functionalities to major taxa commonly found in skin
microbiomes as a proof of concept to demonstrate its ability to uncover novel insights into the
diversity and prevalence of both well-studied and uncharacterized secondary metabolites. For
example, we uncovered the ubiquity of the carotenoid staphyloxanthin across the genus of
Staphylococcus, a molecule conferring the signature golden pigmentation in *S. aureus*[56,57].
These findings raise the hypothesis that staphyloxanthin, which has been identified primarily as
a virulence factor conferring oxidative stress resistance in the pathogen *S. aureus*[41,58], could
similarly contribute to virulence in other staphylococcal species or serve additional unknown
functions. Supporting this hypothesis, we observe that one staphyloxanthin encoding GCF
appears ancestral to the skin-associated, multi-species clade which features both *S. aureus* and
S. epidermidis. The observation that multiple copies of *crt* genes can be found within a single
genome further suggests staphyloxanthin is of functional importance since evolutionary
retention of paralogous genes has previously been shown to underlie ecological
advantages[59,60]. Additionally, while this GCF is present in most skin-associated species, it is
absent in approximately 95% of *S. epidermidis* genomes, suggesting large-scale loss within the
species. It will thus be interesting to further explore how the loss of staphyloxanthin, if it
contributes to oxidative stress resistance within *S. epidermidis*, coincides with the species'
reputation as a beneficial symbiont of host skin[32,61,62].

Using the novel metagenomic mining functionality of *Isa*BGC, we identified novel SNVs
distinguishing uncultured *C. acnes* and *Corynebacterium* strains from isolates with genomes
available in public databases. With regards to *Corynebacterium*, we determined that some of
the most common skin residing species[31,35] belong to the *C. kefirresidentii* species complex,
which were widely distributed across skin metagenomes and highly abundant at moist body
sites. Because this species complex is vastly underrepresented in public genomic databases
and additional species and strains belonging to it likely remain to be discovered, application of
*Isa*BGC-DiscoVary led to the identification of thousands of novel SNVs within the BGCs of the

clade. Critically, some of these variants could have functional implications as they lie within
highly conserved regions of important domains for natural product biosynthesis.
335
The comprehensive nature of *lsa*BGC allows analyses to be applied to both public and
proprietary microbial genomic databases. Besides accelerating fundamental evolutionary
research of BGCs from diverse taxa, we envision *lsa*BGC will be particularly useful for
secondary metabolite discovery and mechanism of action studies. Specifically, we believe the
evolutionary insight gathered from *lsa*BGC can be used as prior information to guide algorithms
in the emerging field of using artificial intelligence to identify new antibiotics[63,64]. Finally, an
improved understanding of the relationship between intra-taxa genomic diversity and BGC
content novelty at the base-resolution using *lsa*BGC highlights which sub-lineages or sub-
clades bear the largest reservoir of untapped secondary metabolic potential.
345

Acknowledgments:

347
This work was supported by grants from the National Institutes of Health (NIAID U19AI142720
and NIGMS R35GM137828 [L.R.K]). The content is solely the responsibility of the authors and
does not necessarily represent the official views of the National Institutes of Health. The authors
gratefully acknowledge additional members of the Kalan lab, Dr. Cameron Currie, Dr. Reed
Stubbendieck, Dr. John-Demian Sauer, Dr. Jason Kwan, Dr. Karthik Anantharaman, Dr.
Caroline Grunenwald, Dr. Chase Clark, and Dr. Milton Drott for helpful discussions.

Materials and Methods:

356
Code availability
358
359 The *lsa*BGC software suite was developed in Python3 and R and is available on Github at:
360 https://github.com/Kalan-Lab/lsaBGC. Algorithmic descriptions of programs can be found on the
361 Github wiki and in the Supplementary Methods and Results document. A test dataset pertaining
362 to the *C. kefirresidentii* species complex for running the seven core programs is provided on the
363 Github repository, along with expected results.
364
An overview of the *lsa*BGC software suite
366
The first core program is *lsa*BGC-Cluster which clusters homologous and non-
fragmented instances of BGCs, identified by antiSMASH from chromosomally complete
genomic assemblies, into gene cluster families (GCFs), similar to BiG-SCAPE[9].Users can then
generate a report to select for the most appropriate parameters for downstream analyses of
interest. Because antiSMASH groups neighboring BGCs into hybrid gene clusters if they are
located close to one another on the genome[7], *lsa*BGC-Refiner has the option to manually
separate hybrid clusters based on user-defined boundary homolog groups. Homologous
instances of GCFs across all available genomic assemblies for the taxa of interest can then be
identified through *lsa*BGC-Expansion (Figure S1b). Similar to ClusterFinder[10], *lsa*BGC-
Expansion uses profile Hidden Markov models (HMMs) to identify homologs of GCF associated

genes in assemblies followed by a classical HMM approach to define BGC regions and their
boundaries. Unlike ClusterFinder, *lsa*BGC-Expansion: (i) uses all coding genes associated with
a GCF instead of being restricted to core protein domains, (ii) is specifically designed to account
for assembly fragmentation through conditional assessment of predicted regions, and (iii)
features a "quick mode" option for fast detection of BGCs using Diamond[65].
382
The remaining four core programs are analytical and designed to be run on a single
GCF for ease of parallelization: *lsa*BGC-See, *lsa*BGC-PopGene, *lsa*BGC-Divergence, and
*lsa*BGC-DiscoVary. *lsa*BGC-See permits visualization of BGCs across a user-provided species
tree or a BGC based phylogeny and can handle multiple fragmented BGC instances from the
same genome, depicting them as neighboring leaves on phylogenies[66]. *lsa*BGC-PopGene can
then infer evolutionary and population genetics statistics for each homolog group associated
with a GCF. It generates codon based alignments for each homolog group which can then be
used as input for *lsa*BGC-Divergence and *lsa*BGC-DiscoVary. *lsa*BGC-PopGene also
determines the consensus order and direction of homolog groups for a GCF across all BGC
instances, which provides intuitive assessment of its reports. *lsa*BGC-Divergence calculates a
comprehensive statistic between pairs of homologous BGCs from the same GCF but different
genomes, which we refer to as Beta-RD, that captures how similar the BGC sequences are
relative to what would be expected based on genome-wide similarity. Finally, *lsa*BGC-DiscoVary
is a multi-functional program which can mine raw sequencing readsets from metagenomes or
single genomes, for GCF instances and then reliably identify whether they possess novel single
nucleotide variants (SNVs) yet to be observed in available genomic assemblies for a taxa
(Figure S2).
400
401 **Usage of genomic and metagenomic datasets**
402
To overcome issues with misclassification and contamination in NCBI, we used GTDB
(release 202) [67], to query for reliable genomic assemblies and respective species designations
for the genera of *Corynebacterium*, *Cutibacterium*, *Micrococcus*, and *Staphylococcus* (Table
S1). For *Micrococcus*, we further included 132 additional genomes of the type species *M. luteus*
which our lab had sequenced (BioProject ID PRJNA803478) but were not yet uploaded to NCBI
to be included in the GTDB release used. These isolates were classified as *M. luteus* by GTDB-
tk[68] and exhibited > 95% ANI to the NCBI representative genome for the species *M. luteus*
NCTC 2665. A similar approach was used for gathering genome sets for species-level analyses
performed (Table S1). For the *C. kefirresidentii* species complex, an additional five genomes
from NCBI were gathered, classified using GTDB-tk[68], and incorporated into analysis if they
belonged to one of the five species designated as belonging to the complex.
414
Because a single representative per species as provided by GTDB might not adequately
capture the diversity of BGCs found across a genus, we instead employed a more granular
approach to select for representative genomes for a genus. We performed dereplication using
MASH[69] with an estimated ANI cutoff of 0.99 to pair similar genomes and then performed single-
linkage clustering and selected representative genomes from each cluster based on assembly
N50 (Table S1). This dereplication was performed using the script

popSizeAndSampleSelector.py provided in the *Isa*BGC suite. Complete or chromosome level
assemblies among the representative genomes were run through IsaBGC-AutoProcess.py and
IsaBGC-Cluster.py to identify the core set of GCFs per genus. GCFs determined from the core
set of genomes were comprehensively identified in the remaining genomes using IsaBGC-
AutoExpansion.py, including those which were not deemed as representatives from the MASH
based analysis to showcase scalability (e.g. all 15,055 *Staphylococcus* and not just the 229
representative genomes of which 77 corresponded to complete/chromosome level assemblies).
To expedite population genetics analysis however, we only ran the downstream functional
analytics of *Isa*BGC using the representative genomes for each genus. For species-level
analyses, all genomes available were used for such downstream analyses and to
comprehensively profile known alleles for genes in BGCs, which was critical to enable
metagenomic mining for base-resolution novelty previously unseen in assemblies available for a
species. For population level analyses within species, strain delineations were performed using
the popSizeAndSampleSelector.py script in *Isa*BGC with strain groups determined from single-
linkage clustering of pairs of genomes found to exhibit $\geq$98% ANI and $\geq$80% genomic content
similarity by FastANI[70].
437
For determining skin-associated species amongst the genera investigated in our study
and to demonstrate mining functionalities with *Isa*BGC-DiscoVary, we used a paired-end
metagenomics dataset sequenced on the Illumina NovaSeq where individual samples
corresponded to microbiome surveys of one of eight body sites for one of 34 participants,
collected in the state of Wisconsin (BioProject ID PRJNA763232)[35]. As previously described[35],
adapter removal, quality filtering, human sequence decontamination, and tandem repeat
removal were performed using fastp (v0.21.0)[71] and KneadData (v0.8.0)
(https://huttenhower.sph.harvard.edu/kneaddata/).
446
Genomic assessment for staphyloxanthin carriage in the lab isolate collection
448
We had previously constructed draft genomic assemblies for 44 *Staphylococcus* isolates
from skin (BioProject ID PRJNA803478; BioProject ID XXXX). Illumina sequencing was
performed at the Microbial Genome Sequencing Center (MiGS). Briefly, assemblies were
generated using Unicycler[72] with default settings after quality and adapter trimming using
fastp[71]. The set of 44 *Staphylococcus* genomes were searched for carriage of staphyloxanthin
encoding GCFs identified in our study using *Isa*BGC-AutoExpansion (Table S9).
455
Library preparation, sequencing and genomic assembly
457
Reconstruction of Illumina draft assemblies for 132 M. luteus: We assessed the performance of
multiple programs in the *Isa*BGC suite using a set of genomic assemblies for 132 *M. luteus*
isolates which were cultured from skin. For these isolates, assemblies were reconstructed from
raw-reads (BioProject ID PRJNA803478) and were uploaded to NCBI's Genbank database
(Table S1). Briefly, Illumina sequencing reads were processed using fastp (v. 0.20.1)[71] to trim
for Poly-G tail artifacts and TrimGalore (v. 0.6.5)[73] was additionally used to detect and remove
adapters. Afterwards, assembly was performed using Unicycler (v. 0.4.6)[72] with default settings.

Construction of Illumina+Nanopore hybrid assemblies for staphyloxanthin producing S. warneri
LK413 and S. epidermidis *LK1136*: To learn the genomic location of staphyloxanthin encoding
GCFs in *S. epidermidis* LK1136 and *S. warneri* LK413, we performed additional long-read
sequencing using Oxford Nanopore Technologies (ONT). Library construction, using a PCR-free
ligation method, and sequencing were performed at the Microbial Genome Sequencing Center
(MiGS). Hybrid assemblies were constructed using a modified version of the Hybrid Assembly
workflow in seQuoia[74]. Illumina reads, used for construction of the initial draft genomic
assemblies for the isolates, were first reprocessed using fastp (v. 0.20.1)[71] to trim for Poly-G tail
artifacts and TrimGalore (v. 0.6.5)[75] was subsequently used to detect and remove adapters.
Similarly, ONT FASTQ reads, provided by MIGS after basecalling, were filtered for potential
adapters using PoreChop (v. 0.2.3) (https://github.com/rrwick/Porechop) and filtered to retain
only reads greater than 3 kb in length as well as subsampled to 300 mb (while being inclusive of
all reads greater than 20 kb) using the script fastqfilter.py included in seQuoia. Afterwards,
hybrid-assemblies were constructed using Unicycler (v. 0.4.6)[72] with default settings and
additional Pilon (v. 1.23)[76] polishing was iteratively performed by aligning ONT reads[77] as well
as the Illumina reads[78] to assemblies and refining them, allowing for a maximum of 10 iterations.
Quality assessment of the two near-complete assemblies was performed using GAEMR
(https://github.com/broadinstitute/GAEMR) (Supplementary Methods and Results).

Experimental validation of staphyloxanthin production in non-aureus *Staphylococcus*

Isolates were grown overnight at 37°C on tryptone soy agar plates and inoculated into tryptone
soy broth at a concentration of $1x10^6$ CFU/mL. Liquid cultures were grown in shaking conditions
at 37°C for 24 hours. 10 mL of the liquid culture was then pelleted and washed twice with
phosphate-buffered saline before pigment extraction in 1 mL of methanol at 55°C for 15 mins.
Spectral scans from 300 nm to 600 nm were taken using a plate reader (EPOCH2, Biotek,
Winooski, VT). Absorbance readings were blanked with methanol and normalized to liquid
culture cell density based on OD600. Spectra shown are representative of at least two biological
replicates.

MASH distance neighbor-joining tree construction

Average nucleotide identity estimates between pairs of genomes were computed with MASH[69]
and used to create a distance matrix. The distance matrix was then used to construct a
neighbor-joining tree with the ape library in R[79,80].

Ribosomal and BGC phylogeny construction

Ribosomal phylogeny construction: Ribosomal phylogenies for each species were constructed
based on a concatenated alignment of 16 ribosomal proteins[81,82]. HMMER (v3) was used to
search for profile HMMs of each ribosomal protein in the predicted proteomes of samples of
interest belonging to a particular taxonomic group. Afterwards, the best matching hits for each
sample based on alignment E-value were identified and used to construct individual alignments

for each ribosomal protein with the L-INS-I method in MAFFT[83]. Protein alignments were then
converted to codon alignments using PAL2NAL[84] and subsequently concatenated together.
Samples which featured ≥ 25% gaps in the concatenated alignment were filtered after which
core SNV sites were extracted and used for constructing a maximum likelihood phylogeny with
RAxML[85], specifying a GTRCAT model and performing 100 bootstraps. Where needed,
ribosomal phylogenies were subset to showcase relationships between select samples using
PareTree (v1.0.2)[86].
516
Staphyloxanthin phylogeny construction: A maximum likelihood phylogeny was constructed from
a concatenated protein alignment of CrtM and CrtN amongst representative *Staphylococcus*
genomes to understand the relationship between different GCFs encoding for staphyloxanthin.
Proteins designated as HG0001876 (CrtM) and HG0001904 (CrtN) were gathered for each
staphyloxanthin encoding GCF found in each sample and aligned using the L-INS-I method in
MAFFT[83]. To account for assembly fragmentation resulting in multiple proteins from a sample
being assigned to the same homolog group for a specific GCF, a consensus sequence was
determined and multi-allele sites were replaced as gaps. A concatenated alignment of such
consensus sequences was subsequently constructed and used for phylogeny construction with
RAxML using a PROTCAT model with JM selected as the best amino-acid replacement matrix
through the automatic maximum likelihood model selector[85]. The phylogeny of CrtMN
sequences exhibited high correspondence to GCF delineations with the exception of a single *S.*
lugdunensis BGC which is likely misclassified as GCF-17 and should be GCF-3. Upon further
examination, it was identified that the reason for the misclassification was likely due to the
absence of mevalonate pathway genes, classified as a type III polyketide synthase (T3PKS),
within *S. lugdunensis* which are found downstream of the *crt* operon in all other species with
GCF-3 (Figure S7b).
534
535   *Pyrazinone phylogeny construction:* A maximum likelihood phylogeny was constructed from a
536   concatenated protein alignment of PznA and PznB amongst representative *Staphylococcus*
537   genomes using the same methods as described for phylogeny construction from CrtMN
538   proteins.
539
540   **Designation of species belonging to *Mammaliicoccus***
541
The *Staphylococcus_A* genus designation in GTDB R202 was largely concordant with the
recent reclassification of five species from *Staphylococcus* to *Mammaliicoccus*[43]. In the latest
GTDB R207 release, this genus has now been reclassified to *Mammaliicoccus*. The recent
update to taxonomic names also includes the reclassification of species *S. pasteuri_A* to *M.*
fleurettii. Notably, we do not regard *S. schleiferi* as part of *Mammaliicoccus* because it is
categorized as *Staphylococcus* by GTDB (both R202 and R207) and groups within the
Staphylococcus genus in our ribosomal protein based maximum-likelihood phylogeny. Similar to
our assessment, the reclassification of this species to *Mammaliicoccus* was recently
questioned[87].
551

Ubiquity assessment of skin residing species and strains of *Staphylococcus* and *Corynebacterium*

We searched for the presence of representative *Staphylococcus* and *Corynebacterium* strains in 270 skin metagenomes from 34 participants across 8 body sites using StrainGST[36]. To construct the database of distinct representative genomes for each genus, the protocol described in the StrainGE documentation was followed. *S. aureus* genomes 16405 and 278 were dropped from the analysis because at the time of downloading them, the plasmid and chromosome were switched. Briefly, genomes were downloaded from NCBI using ncbi-genome-download[88] and dereplication was performed using built-in StrainGE functionalities. As some representative genomes were not part of the GTDB release R202, GTDB-tk[68] was used to annotate such genomes and perform species classification. We considered a species as skin-associated if it was found in at least 10 of the 270 skin metagenomes, regardless of the abundance (Table S4). Certain skin-associated species from each genus were also identified in the 19 negative control metagenomic samples, which included air-swab, bench, desk, and door-handle samplings (Table S4). Because we expect heavy overlap in species content between these environmental samples and skin microbiome samples, we still considered such species as skin-associated if they were found in 10 or more skin metagenomes.

We found the majority of skin-associated *Staphylococcus* belonged to the *S. epidermidis/aureus* clade, as has previously been noted[40]. Similarly, most of the skin-associated *Corynebacterium* belonged to *C. kefirresidentii* and it's close neighboring species (including *C. kefirresidentii*, *C. aurimucosum* type E, *C.* sp. 900539985, *C. tuberculostearicum*, *C. tuberculostearicum* type C), which we denote in this study as the *C. kefirresidentii* species complex. The minimal ANI separating genomes belonging to this species complex was determined to be 88.59% using FastANI[70]. To assess whether any of the 24 new *Corynebacterium* species reported by Saheb Kashaf et al. 2022[34] belonged to this species complex, FastANI was used to look at their ANI similarity to genomes from the clade and regard them as additional members if they exhibited 88.59% ANI or greater. Of note, many additional *Corynebacterium* species have been routinely isolated from the skin[35] which we did not regard as skin-associated based on our criteria in this study. In particular, the species of *C. amycolatum* (detected in 4 metagenomes), *C. kroppenstedtii* (detected in 4 metagenomes), and *C. simulans* (detected in 5 metagenomes) are commonly found on skin.

To estimate the relative abundance of *Cutibacterium* (see section "Identification of *Cutibacterium* Strains within Metagenomes and Determination of Near-Fixed Sites Differentiating Cutimycin Biosynthesis Alleles between Clade I and Clade III *C. acnes*") and *Corynebacterium* strains within metagenomic samples we normalized the estimated relative abundances for strains (*rapct*) by the estimated relative abundance of the genus (*pan%*) as reported by StrainGST[36].

Identification of homolog groups associated with skin residing species of *Staphylococcus* and *Corynebacterium*

We developed a script called crawlingFisher.py, provided in the *lsa*BGC suite, which tests for enrichment or depletion of homolog group presence amongst genomes in focal clades

using Fisher's exact test. Provided with a phylogenetic tree and matrix specifying the presence
of homolog groups across genomes, the program automatically performs such tests for each
homolog group at each inner-node of the phylogeny. Multiple testings correction was performed
comprehensively at the end using Benjamini-Hochberg false discovery rate and cases were
considered statistically significant if the adjusted p-value was less than 0.05. Results were
further filtered to retain only homolog groups which were present in at least 80% of genomes
under the focal node and at most 20% of other genomes or, alternatively, at most 20% of
genomes under the focal node and at least 80% of other genomes. Finally, only results
pertaining to focal nodes where at least 80% of genomes were classified as skin-associated
(see section "Ubiquity Assessment of Skin Residing Species and Strains of *Staphylococcus* and
Corynebacterium") were reported (Table S5).
607
608　**Assessing the divergence of BGCs relative to their background genomic contexts**
609
BGC divergence does not always track with background genomic divergence and
speciation[14]. The program *Isa*BGC-Divergence allows for easy assessment of how BGC
divergence compares to whole-genome divergence, primarily through computing a statistic we
refer to as Beta-Relative-Divergence (Beta-RD) between pairs of genomes for a single GCF.
The Beta-RD statistic is simply the average amino or nucleotide identity between homologous
instances of proteins for a GCF observed in two genomes normalized by the average amino or
nucleotide identity observed between the full genomes. *Isa*BGC allows users to specify whether
amino or nucleotide divergence should be used and automated workflows (see section
Automated Workflows for Running *Isa*BGC) can be used to compute pairwise AAI or ANI
between genomes using CompareM[89] or FastANI[70], respectively.
To better understand how distributions of Beta-RD differ across GCFs, in this study, we
applied a Bayesian hierarchical model to perform shrinking analysis and appropriately account
for differences in the number of genomes GCFs are found (Figure S4cd; Table S6). For
computational efficiency, a maximum of 500 Beta-RD observations were used for each GCF.
Posterior predictions for Beta-RD were made using rstan (Stan Development Team 2021) with
the following hierarchical model:

$$\overline{\mu} \sim Norm(\mu_0, \sigma_0^2)$$
$$\tau \sim half - t_7(1)$$
$$\mu_1...\mu_J \sim Norm(\overline{\mu}, \tau)$$
$$\sigma \sim half - t_7(1)$$
$$y_j \sim Norm(\mu_j, \sigma^2)$$

626
where for each GCF j, the posterior Beta-RD distribution is sampled as y_j using Markov chain
Monte Carlo with four chains, each featuring 2,000 iterations, including a burnin of 1,000
iterations. A hierarchical prior was specified for　$_j$, which represents a GCF-specific baseline.
The values of　$_0$ and σ_0 were set to the respective mean and standard deviation of raw Beta-RD
values across all GCFs. R and STAN code used for this analysis are available within the scripts
subdirectory of the *Isa*BGC software package.

Scrutinization of putative novel variants for *Isa*BGC-DiscoVary application to cutimycin in *C. acnes* and the comprehensive set of GCFs for the *C. kefirresidentii* species complex

By default, lsaBGC-DiscoVary.py only considers reads aligning at 95% identity to
reference gene sequences and avoids reporting novel variants on genes if either annotation or
coverage (relative to other genes in the BGC context) suggests they might be MGEs. The
program also automatically extracts the subset of reads supporting the existence of novel
variants, which can then be further screened with high stringency filters using otherwise
computationally expensive methods. This can involve mapping reads supporting putative novel
SNVs to a representative database of whole-genomes to observe for higher quality alignments
or checking whether reads are classified as a particular taxonomy by Kraken2 (v2.0.8-beta)[90].
As described previously[35], we used a customized database for Kraken2 analysis constructed
from chromosomally complete bacterial, viral, archaeal, fungal, protozoan genomes as well as
the human genome available on RefSeq, where plasmid sequences were distinguished to
alleviate potential misclassifications of reads to a particular taxonomy.

For reads supporting putative novel variants in the cutimycin BGC of *C. acnes*, we
required them to be classified as *Cutibacterium* or *Priopionibacterium* by Kraken2[90] and to not
map with a higher alignment score to a comprehensive database of all *Cutibacterium* genomes
than was observed for the alignment to the reference gene sequence in lsaBGC-DiscoVary.py
(Figure S2c). We generally do not recommend screening variant-supporting reads for an
appropriate taxonomic match with Kraken2 as it can generally lead to false negatives (on
average only 92.8% of read-pairs per sample were taxonomically classified and
misclassification can also occur; Figure S12), but performed this here to provide further
assurance that novel variants discovered are reliable. Standardized coverage metrics within
lsaBGC-DiscoVary.py's novel SNV report were further used to conservatively filter sites and
variants if they exhibited site or allele-specific coverage values greater than two median
absolute deviations from the median coverage for the entire BGC.

For reads supporting putative novel variants within GCFs of the *C. kefirresidentii* species
complex, we required them to not align with a higher alignment score to a comprehensive
database of all *Corynebacterium* genomes. We did not require reads to be classified by Kraken2
as *Corynebacterium* because genomes from the species complex are likely to be poorly
represented in the Kraken2 database. We did however further validate that reads supporting the
existence of variants at conserved sites in the mycolic acid biosynthesis gene mapped best to
the *C. kefirresidentii* species and confirmed our suspicion that they could be misclassified by
Kraken2 given the representation in the underlying database (Figure S12). This *ad hoc* analysis
was performed using paired-end information when mapping in Bowtie2 and accounting for
concordant read pair alignments. While many reads supporting the existence of novel SNVs in
the biosynthesis gene were classified as *C. acnes* by Kraken2, we found that none of the reads
concordantly aligned to the comprehensive *Cutibacterium* genomes Bowtie 2 database whereas
88% concordantly aligned to the comprehensive *Corynebacterium* genomes Bowtie 2 database.

Note, the reasoning behind mapping SNV supporting reads to comprehensive genomic
databases to purge false positive SNV calls, is that if a read truly aligns best to the BGC-
associated homolog group where it was used to call a SNV, then it should at most generate a
mapping to this comprehensive database of equal mapping quality. We align paired-end reads
individually as was performed in lsaBGC-DiscoVary.py to have mapping scores be comparable
with those from the original alignment. Furthermore, because our initial database within lsaBGC-
DiscoVary.py consisted of allelic representatives of homolog groups, we realized that reads
mapping to the start and end of genes could result in a lower mapping score than if a larger
genomic context was provided. To ensure we do not remove such reads mapping to gene
edges, scrutunizeNovelSNVSupport.py, the program we developed to perform Kraken 2 and
genomic database alignment based filtering of novel SNV reports from lsaBGC-DiscoVary.py,
also compares the reference sequences when it finds a better Bowtie 2 mapping for SNV-
supporting reads in genomic databases. If the reference sequence in the genomic database
encompasses the reference sequence in the original alignment to the lsaBGC-DiscoVary.py
homolog group database and the read was previously noted to map to a gene edge, then reads
are still retained and SNV read support is not decremented. Five retained and supportive reads
are required for an SNV to be featured in the filtered report produced by the program.

_lsa_BGC-DiscoVary based analyses of BGCs fromo the _C. kefirresidentii_ species complex:

For the following analyses, we used 5,802 instances of 2,343 unique novel SNVs found
in 68 homolog groups which were core or adjacent to core biosynthesis machinery of BGCs
from the _C. kefirresidentii_ species complex. These novel SNVs had site and allele-specific
coverage values less than two median absolute deviations from the median coverage for the
entire BGC.

Comparing SNVs shared across body sites and subjects: To comprehensively assess whether
novel SNVs were shared more frequently for microbiomes of the same body site or from the
same individual, we performed a pairwise, multi-iteration resampling analysis. For each pair of
metagenomes, where each sample had at least 30 novel SNVs, we calculated the average
Jaccard similarity across 1,000 simulations in which we randomly drew sets of 30 SNVs from
each metagenomic sample to control for sequencing depth. Pairs of metagenomes were
classified as one of three categories: "different body site, different participant", "different body
site, same participant", and "same body site, different participant" (Figure S11c). A two-sided
Wilcoxon rank sum test for differences between the three distributions of average Jaccard
similarities revealed that there was a statistically significant difference between "different body
site, same participant' and "same body site, different participant" (p=6.01E-3). Of greater
relevance, there were statistically strong significant differences between the "different body site,
different participant" distribution and the two other distributions: "different body site, same
participant" and "same body site, different participant" (p=3.67E-09 and p=1.39E-116,
respectively). Thus, pairs of metagenomes were more likely to share novel SNVs if they were
either from the same body site or same participant compared to if they were from different
participants and different body sites. Two metagenomic samples for participant S002 were not
used because they corresponded to a resampling of two of their body sites and we did not have

enough of such samples to pursue a temporal analysis. We assessed this testing using different
considerations/cutoffs for: (i) the number of novel SNVs needed for a metagenome to be
considered in the analysis and (ii) whether or not to account for singleton SNVs. In all cases,
statistical testing for differences between distributions of pairwise metagenome categories
yielded similar conclusions.
725
Assessing a relationship between normalized SNV rates and Tajima's D for homolog groups:
We assessed how the normalized count of novel SNVs observed across all metagenomic
samples for the 68 homolog groups compared with respective Tajima's D estimate computed
from representative alleles found in genomic assemblies. For this analysis, we only considered
homolog groups with at least 1 novel SNV detected in the metagenomes and at least four
sequences available from genomic assemblies, the minimum number needed to compute the
Tajima's D statistic. To normalize novel SNV counts per homolog group, we multiplied the raw
number of novel SNVs observed for a homolog group by the number of representative
sequences present in genomic assemblies and divided by the number of metagenomic samples
deemed to possess the homolog group. The rationale behind this normalization was that more
novel SNVs would be expected for homolog groups that occur in more metagenomes and
similarly fewer novel SNVs would be expected if a greater number of sequences are already
known from available genomic assemblies. We found that homolog groups with Tajima's D < 0
generally had lower normalized novel SNV rates ($\leq$10), while homolog groups with Tajima's D >
0 generally had higher novel SNV rates (>10) (p=1.63E-3, Fisher's exact test). We observed
similar trends when using non-normalized novel SNV counts. Notably, the homolog groups with
the lowest Tajima's D and novel SNV counts were from GCF-5 and GCF-6, which are specific to
a single species, *C. kefirresidentii*.
744
Assessing trends for novel SNV synonymous to non-synonymous rate with ubiquity and site
conservation: Using reports from lsaBGC-DiscoVary.py, we performed a comprehensive
assessment of whether novel SNVs identified on the 68 homolog groups nearby or overlapping
with protocore regions of BGCs in *C. kefirresidentii* corresponded to synonymous or non-
synonymous substitutions. As described in other sections, SNV alleles were compared to
corresponding alleles observed at the site on the reference gene sequences which SNVs were
called upon. We aimed to check whether the rate of synonymous to non-synonymous novel
SNVs showed a relationship to: (i) how common SNVs were across metagenomic samples and
(ii) conservation levels across protein sequences.
754
Novel SNV synonymous to non-synonymous rate increases with SNV ubiquity amongst
metagenomic samples: The occurrence of novel SNVs, a novel allele observed at a specific site
in the multiple sequence codon alignment of a particular homolog group, was tabulated across
metagenomic samples. While novel SNVs found in a single metagenome were only twice as
likely to correspond to a synonymous change as opposed to a non-synonymous change, more
prevalent novel SNVs, found in ten or more samples, were eight times more likely to correspond
to a synonymous change (Figure S11b).
762

Novel SNV synonymous to non-synonymous rate increases at conserved sites in protein
sequences: We searched RefSeq's bacterial NR database for remote homologs of the 68
homolog groups of interest. The top 20 homologs which belonged to classified bacterial species
outside of the *Corynebacterium* genus or *Corynebacteriales* family were identified using
HMMER3 with profile-HMMs gathered from *lsa*BGC-Expansion for each homolog group and an
E-value threshold of 1E-20 (Table S13). The sequences of these representative homologs were
added to protein alignments for homolog groups in *lsa*BGC-PopGene results using the '--add'
option in MAFFT[83]. Afterwards, we used the alignments for scoring protein sequence
conservation with Jensen-Shannon divergence[91]. Conservation scores were used to determine
conservation percentiles of sites along protein alignments for individual homolog groups. Novel
SNVs found in the top five to ten percentile of conserved sites for a homolog group were
approximately five to seven times more likely to represent a synonymous change as opposed to
a non-synonymous change (Figure 5c). In contrast, the bottom 20% of conserved sites (the
least conserved sites), were almost equally as likely to represent non-synonymous substitutions
as synonymous substitutions. Further, the number of novel SNVs, either synonymous or non-
synonymous observed at conserved sites, amongst the top 20% of conserved sites was 649,
and lower than the number of novel SNVs found in the 20-40 (n=739), 40-60 (n=855), 60-80
(n=806), and 80-100 (n=697) percentile ranges of conserved sites. We suspect that the number
of novel SNVs identified decreases past the top 40-60 percentile range because alignments to
such regions become more challenging.
783

References:

1. Pham, J. V. *et al.* A Review of the Microbial Production of Bioactive Natural Products and

Biologics. *Front. Microbiol.* **10**, 1404 (2019).

2. Watve, M. G., Tickoo, R., Jog, M. M. & Bhole, B. D. How many antibiotics are produced by

the genus Streptomyces? *Arch. Microbiol.* **176**, 386–390 (2001).

3. de Lima Procópio, R. E., da Silva, I. R., Martins, M. K., de Azevedo, J. L. & de Araújo, J. M.

Antibiotics produced by Streptomyces. *Braz. J. Infect. Dis.* **16**, 466–471 (2012).

4. Chevrette, M. G. *et al.* The antimicrobial potential of Streptomyces from insect

microbiomes. *Nat. Commun.* **10**, 516 (2019).

5. Waterworth, S. C. *et al.* Horizontal Gene Transfer to a Defensive Symbiont with a Reduced

Genome in a Multipartite Beetle Microbiome. *MBio* **11**, (2020).

6. Drott, M. T. *et al.* Microevolution in the pansecondary metabolome of Aspergillus flavus and

its potential macroevolutionary implications for filamentous fungi. *Proc. Natl. Acad. Sci. U.*

S. A. **118**, (2021).

7.   Steinke, K., Mohite, O. S., Weber, T. & Kovács, Á. T. Phylogenetic Distribution of

Secondary Metabolites in the Bacillus subtilis Species Complex. *mSystems* **6**, (2021).

8.   Blin, K. *et al.* antiSMASH 6.0: improving cluster detection and comparison capabilities.

Nucleic Acids Res. **49**, W29–W35 (2021).

9.   Navarro-Muñoz, J. C. *et al.* A computational framework to explore large-scale biosynthetic

diversity. *Nat. Chem. Biol.* **16**, 60–68 (2020).

10.   Cimermancic, P. *et al.* Insights into secondary metabolism from a global analysis of

prokaryotic biosynthetic gene clusters. *Cell* **158**, 412–421 (2014).

11.   Hannigan, G. D. *et al.* A deep learning genome-mining strategy for biosynthetic gene

cluster prediction. *Nucleic Acids Res.* **47**, e110 (2019).

12.   Kautsar, S. A., van der Hooft, J. J. J., de Ridder, D. & Medema, M. H. BiG-SLiCE: A highly

scalable tool maps the diversity of 1.2 million biosynthetic gene clusters. *Gigascience* **10**,

(2021).

13.   Emms, D. M. & Kelly, S. OrthoFinder: phylogenetic orthology inference for comparative

genomics. *Genome Biol.* **20**, 238 (2019).

14.   Chase, A. B., Sweeney, D., Muskat, M. N., Guillén-Matus, D. G. & Jensen, P. R. Vertical

Inheritance Facilitates Interspecies Diversification in Biosynthetic Gene Clusters and

Specialized Metabolites. *MBio* **12**, e0270021 (2021).

15.   Felnagle, E. A., Rondon, M. R., Berti, A. D., Crosby, H. A. & Thomas, M. G. Identification of

the biosynthetic gene cluster and an additional gene for resistance to the antituberculosis

drug capreomycin. *Appl. Environ. Microbiol.* **73**, 4162–4170 (2007).

16.   Del Carratore, F. *et al.* Computational identification of co-evolving multi-gene modules in

microbial biosynthetic gene clusters. *Commun Biol* **2**, 83 (2019).

17.   Crits-Christoph, A., Bhattacharya, N., Olm, M. R., Song, Y. S. & Banfield, J. F. Transporter

genes in biosynthetic gene clusters predict metabolite characteristics and siderophore

activity. *Genome Res.* (2020) doi:10.1101/gr.268169.120.

18. Crits-Christoph, A., Diamond, S., Butterfield, C. N., Thomas, B. C. & Banfield, J. F. Novel

soil bacteria possess diverse genes for secondary metabolite biosynthesis. *Nature* **558**,

440–444 (2018).

19. Nurk, S., Meleshko, D., Korobeynikov, A. & Pevzner, P. A. metaSPAdes: a new versatile

metagenomic assembler. *Genome Res.* **27**, 824–834 (2017).

20. Pereira-Flores, E. *et al.* Mining metagenomes for natural product biosynthetic gene

clusters: unlocking new potential with ultrafast techniques. *bioRxiv* 2021.01.20.427441

(2021) doi:10.1101/2021.01.20.427441.

21. Andreu, V. P. *et al.* A systematic analysis of metabolic pathways in the human gut

microbiota. *bioRxiv* 2021.02.25.432841 (2021) doi:10.1101/2021.02.25.432841.

22. Sugimoto, Y. *et al.* A metagenomic strategy for harnessing the chemical repertoire of the

human microbiome. *Science* **366**, (2019).

23. Chevrette, M. G. & Handelsman, J. Needles in haystacks: reevaluating old paradigms for

the discovery of bacterial secondary metabolites. *Nat. Prod. Rep.* **38**, 2083–2099 (2021).

24. Zipperer, A. *et al.* Human commensals producing a novel antibiotic impair pathogen

colonization. *Nature* **535**, 511–516 (2016).

25. Claesen, J. *et al.* A Cutibacterium acnes antibiotic modulates human skin microbiota

composition in hair follicles. *Sci. Transl. Med.* **12**, (2020).

26. Nakatsuji, T. *et al.* Antimicrobials from human skin commensal bacteria protect against

Staphylococcus aureus and are deficient in atopic dermatitis. *Sci. Transl. Med.* **9**, (2017).

27. Wyatt, M. A. *et al.* Staphylococcus aureus nonribosomal peptide secondary metabolites

regulate virulence. *Science* **329**, 294–296 (2010).

28. Ridaura, V. K. *et al.* Contextual control of skin immunity and inflammation by

Corynebacterium. *J. Exp. Med.* **215**, 785–799 (2018).

29. Swaney, M. H. & Kalan, L. R. Living in Your Skin: Microbes, Molecules, and Mechanisms.

Infect. Immun. **89**, (2021).

30. Grice, E. A. *et al.* Topographical and temporal diversity of the human skin microbiome.

Science **324**, 1190–1192 (2009).

31. Oh, J. *et al.* Temporal Stability of the Human Skin Microbiome. *Cell* **165**, 854–866 (2016).

32. Byrd, A. L., Belkaid, Y. & Segre, J. A. The human skin microbiome. *Nat. Rev. Microbiol.* **16**,

143–155 (2018).

33. Acosta, E. M. *et al.* Bacterial DNA on the skin surface overrepresents the viable skin

microbiome. *bioRxiv* 2021.08.16.455933 (2021) doi:10.1101/2021.08.16.455933.

34. Saheb Kashaf, S. *et al.* Integrating cultivation and metagenomics for a multi-kingdom view

of skin microbiome diversity and functions. *Nat Microbiol* **7**, 169–179 (2022).

35. Swaney, M. H., Sandstrom, S. & Kalan, L. R. Cobamide sharing drives skin microbiome

dynamics. *bioRxiv* 2020.12.02.407395 (2021) doi:10.1101/2020.12.02.407395.

36. van Dijk, L. R. *et al.* StrainGE: a toolkit to track and characterize low-abundance strains in

complex microbial communities. *Genome Biol.* **23**, 74 (2022).

37. Zimmermann, M. & Fischbach, M. A. A family of pyrazinone natural products from a

conserved nonribosomal peptide synthetase in Staphylococcus aureus. *Chem. Biol.* **17**,

925–930 (2010).

38. Tauch, A. *et al.* Complete genome sequence and analysis of the multiresistant nosocomial

pathogen Corynebacterium jeikeium K411, a lipid-requiring bacterium of the human skin

flora. *J. Bacteriol.* **187**, 4671–4682 (2005).

39. Pelz, A. *et al.* Structure and biosynthesis of staphyloxanthin from Staphylococcus aureus. *J.*

Biol. Chem. **280**, 32493–32498 (2005).

40. Balibar, C. J., Shen, X. & Tao, J. The mevalonate pathway of Staphylococcus aureus. *J.*

Bacteriol. **191**, 851–861 (2009).

41. Clauditz, A., Resch, A., Wieland, K.-P., Peschel, A. & Götz, F. Staphyloxanthin plays a role

in the fitness of Staphylococcus aureus and its ability to cope with oxidative stress. *Infect.*

Immun. **74**, 4950–4953 (2006).

42. Seel, W. *et al.* Carotenoids are used as regulators for membrane fluidity by Staphylococcus

xylosus. *Sci. Rep.* **10**, 330 (2020).

43. Madhaiyan, M., Wirth, J. S. & Saravanan, V. S. Phylogenomic analyses of the

Staphylococcaceae family suggest the reclassification of five species within the genus

Staphylococcus as heterotypic synonyms, the promotion of five subspecies to novel

species, the taxonomic reassignment of five Staphylococcus species to Mammaliicoccus

gen. nov., and the formal assignment of Nosocomiicoccus to the family Staphylococcaceae.

Int. J. Syst. Evol. Microbiol. **70**, 5926–5936 (2020).

44. Dong, P.-T. *et al.* Photolysis of Staphyloxanthin in Methicillin-Resistant Staphylococcus

aureus Potentiates Killing by Reactive Oxygen Species. *Adv. Sci.* **6**, 1900030 (2019).

45. Li, D., Liu, C.-M., Luo, R., Sadakane, K. & Lam, T.-W. MEGAHIT: an ultra-fast single-node

solution for large and complex metagenomics assembly via succinct de Bruijn graph.

Bioinformatics **31**, 1674–1676 (2015).

46. Gande, R. *et al.* Acyl-CoA carboxylases (accD2 and accD3), together with a unique

polyketide synthase (Cg-pks), are key to mycolic acid biosynthesis in Corynebacterianeae

such as Corynebacterium glutamicum and Mycobacterium tuberculosis. *J. Biol. Chem.* **279**,

44847–44857 (2004).

47. Portevin, D. *et al.* A polyketide synthase catalyzes the last condensation step of mycolic

acid biosynthesis in mycobacteria and related organisms. *Proc. Natl. Acad. Sci. U. S. A.*

101, 314–319 (2004).

48. Chen, Y. *et al.* Structural classification and properties of ketoacyl synthases. *Protein Sci.*

20, 1659–1667 (2011).

49. Klaus, M., Buyachuihan, L. & Grininger, M. Ketosynthase Domain Constrains the Design of

Polyketide Synthases. *ACS Chem. Biol.* **15**, 2422–2432 (2020).

50. Medema, M. H., Cimermancic, P., Sali, A., Takano, E. & Fischbach, M. A. A systematic

computational analysis of biosynthetic gene cluster evolution: lessons for engineering

biosynthesis. *PLoS Comput. Biol.* **10**, e1004016 (2014).

51. Palaniappan, K. *et al.* IMG-ABC v.5.0: an update to the IMG/Atlas of Biosynthetic Gene

Clusters Knowledgebase. *Nucleic Acids Res.* **48**, D422–D430 (2019).

52. Kautsar, S. A. *et al.* MIBiG 2.0: a repository for biosynthetic gene clusters of known

function. *Nucleic Acids Res.* **48**, D454–D458 (2020).

53. Blin, K., Shaw, S., Kautsar, S. A., Medema, M. H. & Weber, T. The antiSMASH database

version 3: increased taxonomic coverage and new query features for modular enzymes.

Nucleic Acids Res. **49**, D639–D643 (2020).

54. Garrison, E. *et al.* Variation graph toolkit improves read mapping by representing genetic

variation in the reference. *Nat. Biotechnol.* **36**, 875–879 (2018).

55. Garrison, E. & Guarracino, A. Unbiased pangenome graphs. *bioRxiv* 2022.02.14.480413

(2022) doi:10.1101/2022.02.14.480413.

56. Kloos, W. E. & Schleifer, K. H. Isolation and Characterization of Staphylococci from Human

Skin II. Descriptions of Four New Species: Staphylococcus warneri, Staphylococcus capitis,

Staphylococcus hominis, and Staphylococcus simulans. *Int. J. Syst. Bacteriol.* **25**, 62–79

(1975).

57. Becker, K., Heilmann, C. & Peters, G. Coagulase-negative staphylococci. *Clin. Microbiol.*

Rev. **27**, 870–926 (2014).

58. Liu, C.-I. *et al.* A cholesterol biosynthesis inhibitor blocks Staphylococcus aureus virulence.

Science **319**, 1391–1394 (2008).

59. Blount, Z. D., Barrick, J. E., Davidson, C. J. & Lenski, R. E. Genomic analysis of a key

innovation in an experimental Escherichia coli population. *Nature* **489**, 513–518 (2012).

60. Ghosh, S. & O'Connor, T. J. Beyond Paralogs: The Multiple Layers of Redundancy in

Bacterial Pathogenesis. *Front. Cell. Infect. Microbiol.* **7**, 467 (2017).

61. Otto, M. Staphylococcus epidermidis--the 'accidental' pathogen. *Nat. Rev. Microbiol.* **7**,

555–567 (2009).

62. Brown, M. M. & Horswill, A. R. Staphylococcus epidermidis-Skin friend or foe? *PLoS*

Pathog. **16**, e1009026 (2020).

63. Stokes, J. M. *et al.* A Deep Learning Approach to Antibiotic Discovery. *Cell* **180**, 688–

702.e13 (2020).

64. Melo, M. C. R., Maasch, J. R. M. A. & de la Fuente-Nunez, C. Accelerating antibiotic

discovery through artificial intelligence. *Commun Biol* **4**, 1050 (2021).

65. Buchfink, B., Xie, C. & Huson, D. H. Fast and sensitive protein alignment using DIAMOND.

Nat. Methods **12**, 59–60 (2014).

66. Huerta-Cepas, J., Serra, F. & Bork, P. ETE 3: Reconstruction, Analysis, and Visualization

of Phylogenomic Data. *Mol. Biol. Evol.* **33**, 1635–1638 (2016).

67. Parks, D. H. *et al.* GTDB: an ongoing census of bacterial and archaeal diversity through a

phylogenetically consistent, rank normalized and complete genome-based taxonomy.

Nucleic Acids Res. (2021) doi:10.1093/nar/gkab776.

68. Chaumeil, P.-A., Mussig, A. J., Hugenholtz, P. & Parks, D. H. GTDB-Tk: a toolkit to classify

genomes with the Genome Taxonomy Database. *Bioinformatics* **36**, 1925–1927 (2019).

69. Ondov, B. D. *et al.* Mash: fast genome and metagenome distance estimation using

MinHash. *Genome Biol.* **17**, 132 (2016).

70. Jain, C., Rodriguez-R, L. M., Phillippy, A. M., Konstantinidis, K. T. & Aluru, S. High

throughput ANI analysis of 90K prokaryotic genomes reveals clear species boundaries.

Nat. Commun. **9**, 1–8 (2018).

71. Chen, S., Zhou, Y., Chen, Y. & Gu, J. fastp: an ultra-fast all-in-one FASTQ preprocessor.

Bioinformatics **34**, i884–i890 (2018).

72. Wick, R. R., Judd, L. M., Gorrie, C. L. & Holt, K. E. Unicycler: Resolving bacterial genome

assemblies from short and long sequencing reads. *PLoS Comput. Biol.* **13**, e1005595

(2017).

953 73. Krueger, F., James, F., Ewels, P., Afyounian, E. & Schuster-Boeckler, B.

954   *FelixKrueger/TrimGalore: v0.6.7 - DOI via Zenodo.* (2021). doi:10.5281/zenodo.5127899.

955 74. *seQuoia: A framework for diverse omics workflows, primarily focused on QC, processing*

956   *and analysis of bacterial genomic sequencing sets.* (Github).

957 75. Krueger, F. *TrimGalore: A wrapper around Cutadapt and FastQC to consistently apply*

958   *adapter and quality trimming to FastQ files, with extra functionality for RRBS data.* (Github).

959 76. Walker, B. J. *et al.* Pilon: an integrated tool for comprehensive microbial variant detection

960   and genome assembly improvement. *PLoS One* **9**, e112963 (2014).

961 77. Li, H. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* **34**, 3094–

962   3100 (2018).

963 78. Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows–Wheeler

964   transform. *Bioinformatics* **25**, 1754–1760 (2009).

965 79. Saitou, N. & Nei, M. The neighbor-joining method: a new method for reconstructing

966   phylogenetic trees. *Mol. Biol. Evol.* **4**, 406–425 (1987).

967 80. Paradis, E., Claude, J. & Strimmer, K. APE: Analyses of Phylogenetics and Evolution in R

968   language. *Bioinformatics* **20**, 289–290 (2004).

969 81. Hug, L. A. *et al.* A new view of the tree of life. *Nature Microbiology* **1**, 1–6 (2016).

970 82. McDaniel, E. A., Anantharaman, K. & McMahon, K. D. metabolisHMM: Phylogenomic

971   analysis for exploration of microbial phylogenies and metabolic pathways. *bioRxiv*

972   2019.12.20.884627 (2020) doi:10.1101/2019.12.20.884627.

973 83. Katoh, K. & Standley, D. M. MAFFT multiple sequence alignment software version 7:

974   improvements in performance and usability. *Mol. Biol. Evol.* **30**, 772–780 (2013).

975 84. Suyama, M., Torrents, D. & Bork, P. PAL2NAL: robust conversion of protein sequence

976   alignments into the corresponding codon alignments. *Nucleic Acids Res.* **34**, W609–12

977   (2006).

978 85. Stamatakis, A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large

phylogenies. *Bioinformatics* **30**, 1312–1313 (2014).

86. PareTree 1.0: Remove Sequences, Bootstraps, and Branch Lengths From Your Trees!

http://emmahodcroft.com/PareTree.html.

87. Kania, S. A. Reclassification of Staphylococcus schleiferi by Madhaiyan et al. lacks key

supporting data. *Int. J. Syst. Evol. Microbiol.* **72**, (2022).

88. Blin, K. *ncbi-genome-download: Scripts to download genomes from the NCBI FTP servers*.

(Github).

89. Parks, D. *CompareM: A toolbox for comparative genomics*. (Github).

90. Wood, D. E., Lu, J. & Langmead, B. Improved metagenomic analysis with Kraken 2.

Genome Biol. **20**, 1–13 (2019).

91. Capra, J. A. & Singh, M. Predicting functionally important residues from sequence

conservation. *Bioinformatics* **23**, 1875–1882 (2007).

Figure 1: *Isa*BGC offers an efficient and sensitive means for GCF homology detection. a) A schematic overview of the full *Isa*BGC suite. **b**) Neighbor-joining trees were constructed for each of the four genera based on pairwise MASH ANI estimates between genomes and heatmaps showcase the presence of GCFs found in ≥ 5% of genomes. Red dots on neighbor-joining trees signify complete or chromosomal genomes used for initial identification of BGCs. Heatmap cell color indicates the BGC class as predicted by antiSMASH. **c**) Results from benchmarking *Isa*BGC-AutoExpansion sensitivity on 14 completed genomes for the species of *M. luteus*. AntiSMASH and *Isa*BGC-AutoExpansion were run on artificially fragmented genomes and resulting BGC predictions were compared to initial BGCs identified on the unfragmented, complete genomes. **d**) The relationship between the assembly quality measured by N50 and the total number of homolog groups detected across 14,978 *Staphylococcus* draft genomes is shown. The coloring corresponds to the percentage of BGC homolog groups co-located in a segment with a homolog group commonly found in protocore regions of BGC predictions by antiSMASH run on completed genomes.

Figure 2: *Isa*BGC reveals new insight about characterized and unstudied BGCs from skin-associated species. **a**) A maximum-likelihood phylogeny was constructed from PznA and PznB showing the relationship between GCFs encoding for pyrazinones. Branch color corresponds to different GCFs and the outer circular color denotes the species classification for the genome from which the sequences were gathered. **b**) Homolog groups across the seven pyrazinone encoding GCFs are depicted as a network where edges indicate that two homolog groups co-occur in the same GCF. Homolog group color of nodes depict distinct GCFs. **c**) Phylogeny constructed from ribosomal proteins showing the conservation of *Corynebacterium* GCF-50 across the diverse species encoding the NRPS using results from *Isa*BGC-PopGene analysis. The gene schematics show the consensus gene order for the GCF. Gene lengths correspond to the median length across all BGC instances and only genes found in two or more genomes are shown. For each species, the gene is colored based on the median similarity of gene sequences from the species to the consensus sequence across all species. The barplot displays the total number of GCF instances identified per species across the full set of *Corynebacterium* genomes. Broad annotations for homolog groups are shown as a color strip. Below it are bar plots corresponding to the conservation of homolog groups across samples and their median Beta-RD. **d**) A histogram of the amino acid identities between NRP synthase instances is compared to the average amino acid identities (AAI) between genomes they were gathered from.

Figure 3: **Staphyloxanthin is ubiquitous across *Staphylococcus* and found on plasmids. a)** A maximum-likelihood phylogeny of the full *Staphylococcus* genus was constructed from ribosomal protein encoding genes. The clade in green represents the newly recognized genus of *Mammaliicoccus*. Heatmaps showcase the presence of the five *crt* genes for each species for five separate GCFs containing the *crt* operon. The shading indicates the proportion of a species found to carry homolog groups within a particular GCF context in $\log_{10}$ scale. **b)** A maximum-likelihood phylogeny was built from CrtM and CrtN protein alignments. Branch colors represent the GCF classification of the CrtMN sequences and bootstrap values are shown for key nodes where GCFs partition. The species the CrtMN sequence was extracted from is shown as a color strip followed by a bar plot depicting the comparative codon usage cosine distance (CCU-cd). CCU-cd was only calculated for the full five gene *crt* operon and represents the codon frequency dissimilarity with the codon frequency of the background genome. The heatmap showcases the genome-wide average amino acid identity (AAI) between pairs of genomes from which CrtMN sequences were gathered. **c)** AAI and shared homolog group content were calculated between pairs of GCF instances from different genomes for GCF-3 and GCF-6 individually. The coloring represents genome-wide AAI similarity. **d)** A schematic of the completed genome for *S. equorum* C2014, which was predicted to feature GCF-6 in three distinct copies. Bottom panel shows the BGC alignment between the three instances of GCF-6 found within this genome. **e)** Representative isolates with and without staphyloxanthin encoding GCFs were grown on 20% BHI agar to visualize pigmentation production. **f)** Methanol extractions and wavelength absorption analysis were performed to identify signature peaks in the 400 to 500 nm range associated with the presence of staphyloxanthin in *S. aureus*[39,44]. Spectra are normalized to cell density.

Figure 4: Mining for microdiversity within the cutimycin BGC leads to genetic insights for clade III *C. acnes*. a) Codon usage dissimilarity between BGC regions and background genomes shown for six common GCFs in *C. acnes*. **b**) The cutimycin encoding GCF is found primarily in clade I (subclade IB; orange) and clade III (tan) *C. acnes*. **c**) A schematic of the cutimycin encoding GCF. Lines in black correspond to sites of nearly-fixed differences between clades of *C. acnes*. **d**) *Isa*BGC-DiscoVary was used to identify the presence of the cutimycin GCF in 270 skin metagenomes. The proportion of alleles representative of clade IB or clade III *C. acnes* at sites of nearly fixed-differences was quantified. High-resolution strain quantification within metagenomes was similarly performed with StrainGE and used to overlay clade-specific typing results from *Isa*BGC-DiscoVary analysis. **e**) The distribution of novel SNVs identified by *Isa*BGC-DiscoVary within metagenomes, post filtering for removal of false positives, is shown across homolog groups and subjects. **f**) The location of novel SNVs is shown along the lantibiotic dehydratase N-terminal encoding gene (*HG2417*) for each subject. Coloring of the novel SNVs corresponds to which body-site(s) they were identified within. **g**) The clade distribution of reference genes on which novel SNVs were identified across the cutimycin GCF for subject S012 are shown individually for the predicted transitions or transversions. **i**) The proportion of differences which corresponded to cytosine to thymine or thymine to cytosine transitions along a core-genome alignment was calculated between pairs of *C. acnes* and categorized by clade comparisons. **j**) Comparative analysis of presumed directional deaminations from *C. namnetense*, treated as ancestral, to *C. acnes* clade IB and clade III were estimated using a multi-iteration sub-sampling based approach for sites along the core-genome alignment, whereby a greater number of suspected C to T transitions were estimated for clade III *C. acnes*. **k**) The presence of a complete uracil DNA glycosylase (*ung*), from clade I *C. acnes*, is shown across available *Cutibacterium* genomes.

Figure 5: Thousands of novel SNVs identified within BGCs from the *C. kefirresidentii* species complex. a) Assessment of novel SNVs identified by *lsa*BGC-DiscoVary, post-filtering, for each BGC from the *C. kefirresidentii* species complex for presence within metagenomic assemblies. **b**) Tajima's D statistic, calculated for homolog groups using sequences in available genomic assemblies, is compared to the normalized number of novel SNVs found per homolog groups within metagenomes. **c**) The number of novel SNVs is shown for ranges of conservation percentile across all homolog groups and metagenomes. The coloring corresponds to the ratio of synonymous to non-synonymous novel SNVs within each conservation range. **d**) Novel SNVs are shown for the polyketide synthase involved in mycolic acid biosynthesis. The top panel showcases the non-synonymous (up-facing) and synonymous (down-facing) novel SNVs found along the T1PKS gene, with heights corresponding to the number of metagenomes with the novel SNV. The middle panel shows the conservation scores colored by percentile ranges (lightest = least conserved sites; darkest = most conserved sites). Key domains common to T1PKS are shown underneath along with the location of predicted non-synonymous mutations within the top 5% of conserved sites for the gene. Two adjacent SNVs affecting the coding of the residue at position 342 were identified and these could result in a combined effect of encoding for alanine.