*lsa*BGC provides a comprehensive framework for evolutionary analysis of biosynthetic gene clusters within focal taxa

Rauf Salamzade[1], J.Z. Alex Cheong[1], Shelby Sandstrom[1], Mary Hannah Swaney[1], Reed M. Stubbendieck[2], Nicole Lane Starr[1], Cameron R. Currie[2], Anne Marie Singh[3], and Lindsay R. Kalan[1,4]

[1]Department of Medical Microbiology and Immunology, School of Medicine and Public Health, University of Wisconsin, Madison, WI, USA
[2]Department of Bacteriology, University of Wisconsin—Madison, Madison, Wisconsin, USA
[3]Department of Pediatrics, School of Medicine and Public Health, University of Wisconsin, Madison, WI, USA
[4]Department of Medicine, School of Medicine and Public Health, University of Wisconsin, Madison, WI, USA

Address for Correspondence: Lindsay Kalan, 1550 Linden Dr, 6155 Microbial Sciences Building, Madison, WI, USA, 53706. lkalan@wisc.edu

Abstract:

Bacterial secondary metabolites, encoded by biosynthetic gene clusters (BGCs), can underlie microbiome homeostasis and be developed into pharmaceutical drugs or agricultural chemicals. While several software and databases have been developed for genomic exploration of BGCs, recent advances have largely focused on clustering orthologous instances of BGCs from different genomes into gene cluster families (GCFs). Here we present *lsa*BGC, a software suite which allows for detailed genomic investigation of individual GCFs within specific taxonomic contexts. The suite provides several novel functionalities, including: (i) direct, sensitive, and precise identification of GCFs in genomes, (ii) base-resolution mining of novel variants within BGCs from metagenomes, and (iii) generating comprehensive overviews of the conservation, function, syntenic organization, and evolutionary trends of genes found in GCFs. To showcase the suite, we applied *lsa*BGC to four genera commonly found in skin microbiomes to reveal novel insights on both well and less studied taxa and their secondary metabolites. We find evidence of inter-species horizontal gene transfer (HGT) within multiple genera, including the *crt* BGC encoding for staphyloxanthin, a virulence factor in *Staphylococcus aureus*. We show the *crt* BGC is ubiquitous across the *Staphylococcus* genus and exhibits varying evolutionary trajectories in different species contexts, being highly conserved amongst *S. aureus* while being lost in approximately 95% of *Staphylococcus epidermidis*, the most abundant staphylococcal species on healthy skin. *lsa*BGC's framework was also applied to healthy-skin metagenomes to uncover thousands of novel single nucleotide variants (SNVs) within BGCs from the *Corynebacterium kefirresidentii* sp. complex, which we determine here to be the most prevalent group of *Corynebacterium* in skin microbiomes. Although novel SNVs were approximately ten times as likely to correspond to synonymous changes when located in the top five percentile of conserved sites, *lsa*BGC identified SNVs which defied this trend and are predicted to underlie amino acid changes within functionally key enzymatic domains. Ultimately, *lsa*BGC provides a new repertoire of tools to ease evolutionary and comparative analysis of BGCs across ever expanding genomic datasets to gain novel insights into uncharacterized secondary metabolites and their genomic diversity. Such tools may expedite efforts in the discovery of new natural products with high commercial value.

Introduction:

The secondary metabolome of bacteria has served as a valuable reservoir of natural products with great societal benefit[1]. Historically, many commercialized secondary metabolites, such as antibiotics, have been identified from particular taxonomic groups, such as the genus of *Streptomyces* within Actinomycetota[2]. Several studies have explored the secondary metabolome of these metabolically rich microbial taxa to develop a comprehensive understanding of their chemical diversity and unique traits[3–6]. Most of these studies begin by first identifying biosynthetic gene clusters within genomes using the popular antiSMASH software[7] and then group BGCs into GCFs using BiG-SCAPE[8]. Critically, this approach, in addition to other bioinformatics software for genomics annotation and analysis of BGCs[9–11], uses a set of key protein domains as their base to group BGCs predicted to encode similar metabolites.

44　However, when working in the context of a single genus or species, it becomes more
45　appropriate to use full protein sequences which enables the use of advanced algorithms for the
46　identification of orthologs and paralogs[12] to reliably determine evolutionary relationships
47　between BGCs.
48
49　　　Recent studies which explored the vertical descent and conservation of BGCs across
50　the genus of *Salinospora* and the fungal species *Aspergillus flavus* have shown that intra-genus
51　or intra-species evolution can result in chemical and regulatory diversification of the encoded
52　metabolites[5,13]. Thus, improved understanding of the evolutionary emergence of BGCs could be
53　informative towards elucidating the function of individual genes and encoded metabolites[8,14–16].
54
Due to difficulties in the cultivation of many bacterial taxa, several recent studies have
begun to explore metagenomic datasets to unearth novel secondary metabolites[17–19]. While
most of these endeavors have relied on performing metagenomic assembly, recently, two read-
based approaches were described to reliably and sensitively search for key BGC domains in
metagenomes[18,20]. This highlights a larger trend in which these approaches primarily aim to
identify highly novel secondary metabolites and thus prioritize investigation of poorly studied or
newly discovered taxa[21]. However, provided that most metabolites have been mined from select
genera, mostly within the *Actinomycetia* class[2], there could be tremendous reward for efforts to
advance our understanding of cataloged but uncharacterized BGCs from well studied taxa. This
is especially important because many species within the metabolically rich *Streptomyces* only
have limited representative genomes, thus methods to leverage metagenomic datasets and
identify homologous instances of rare BGCs can lead to critical insight into their function and
ecological distribution[3].
68
To aid the advancement of such taxa-specific analyses of BGCs, we introduce the
comprehensive software suite *Lineage Specific Analysis* of BGCs (*lsa*BGC;
https://github.com/Kalan-Lab/lsaBGC), which consists of several programs for comparative
genetics of BGCs as well as metagenomic mining. We showcase the utility of *lsa*BGC in this
study through application to four major genera which are commonly found in healthy human skin
microbiomes. Several secondary metabolites which function as antibiotics[22–24], virulence
factors[25], or in microbe-host interactions[26,27] have been identified in bacteria from skin
microbiomes, primarily from staphylococci. Human skin is also easily accessible to sampling
and most species can be cultivated for follow up studies. Using *lsa*BGC, we reveal new
ecological and evolutionary insights for both well characterized and unknown predicted
metabolites, including signatures of inter-species transfer for the virulence-associated
staphylococcal carotenoid staphyloxanthin and a highly conserved predicted siderophore found
encoded by divergent *Corynebacterium* species. In addition, through mining skin metagenomic
datasets, we find novel coding variants within the functionally important ketide synthase (KS)
domain of the polyketide synthase (PKS) responsible for mycolic acid biosynthesis, a defining
cell wall component of *Corynebacterium* species implicated in immune recognition.
85
86　# Results:
87

IsaBGC allows for sensitive and systematic identification of BGC homolog groups

The *Isa*BGC suite consists of seven core programs, three workflow programs, and several additional scripts for analyses (Figure 1, S1; Supplementary Methods and Results). These programs allow for clustering BGCs into GCFs, refining boundaries of BGCs, and sensitive, yet specific, rapid detection of GCF instances in draft-quality assemblies. The suite additionally includes dedicated tools for GCF visualization, evolutionary and population genetic analysis of individual genes within GCFs, and base-resolution identification of novel SNVs in complex metagenomes.

To demonstrate the capabilities of the *Isa*BGC suite we applied it to four genera which are stable and integral members of the human skin microbiome[28–32]. These included three genera in the Actinomycetota (formerly Actinobacteria) phylum: *Corynebacterium*, *Cutibacterium*, and *Micrococcus* as well as the Bacillota (formerly Firmicutes) genus *Staphylococcus*. *Isa*BGC was also independently applied to the species or species-complex from each genus most commonly represented in skin microbiomes. These were the *C. kefirresidentii* species-complex, *Cutibacterium acnes*, *Micrococcus luteus*, and *S. epidermidis* (Table S1).

We used *Isa*BGC-AutoProcess to run gene-calling, BGC annotation, and *de novo* homolog group delineation with complete or chromosomally complete genomes for each genus or species. We then applied *Isa*BGC-Cluster to group homologous BGCs into GCFs and systematically searched for instances of GCFs using *Isa*BGC-AutoExpansion (Figure 1b; Table S2, S3; Supplementary Methods and Results). To assess the validity and sensitivity of our approach, two benchmarking tests using BGCs identified in *M. luteus* were performed (Figure 1c, S3; Supplementary Methods and Results) showing that the *Isa*BGC-AutoExpansion framework leads to greater sensitivity compared to running *de novo* antiSMASH on fragmented draft-quality assemblies. This increased sensitivity was due to more robust, yet reliable, detection of auxiliary genes located on the edges of scaffolds not containing core domains used by antiSMASH for BGC detection. In addition, *Isa*BGC-AutoExpansion is highly efficient and allowed us to search for 63 GCFs identified from high-quality *Staphylococcus* genomes in 14,978 draft-quality genomes in approximately 14 hours using 40 cores and less than 150 Gb of memory. We found a positive association between detection of complete BGC instances and assembly quality (Figure 1d; Supplementary Methods and Results).

IsaBGC enables reliable comparative genomics to explore evolutionary trends of BGCs

The increased sensitivity of *Isa*BGC to uniformly detect auxiliary components of BGCs in each genome enabled us to perform reliable comparative genetic analyses across GCF instances. We focused on two of the four genera with the greatest breadth of diversity, *Corynebacterium* and *Staphylococcus*, and selected a subset of 456 and 229 representative genomes from each genera, respectively (Table S1). To better understand which species in these genera were most relevant to skin health, we first profiled how often distinct species were found among healthy skin microbiomes using a large, recently generated metagenomics

132　dataset[33,34] (Table S4). Skin-associated species (found in more than 10 of 270 metagenomes)
133　were phylogenetically clustered for both genera, within the *S. epidermidis* clade (Figure S4a)
134　and within the *C. kefirresidentii* species complex (Figure S4b; Supplementary Methods and
135　Results).
136
Association of individual homolog groups from these GCFs revealed several were
enriched within skin-associated staphylococci, but only two were enriched for skin-associated
corynebacteria, both encoding hypothetical proteins (Table S5). Nearly half of the homolog
groups enriched for skin-associated staphylococci (5 of 12; 41.7%) were related to GCF-10,
predicted to encode for an NRP synthetase corresponding to pyrazinone biosynthesis,
molecules reported to regulate virulence in *S. aureus*[25,35]. Staphylococcal pyrazinones were
previously identified as being a self-contained BGC consisting of two genes, *pznA* and *pznB*,
specific to species commonly associated with human skin, including *S. aureus, S. epidermidis*,
and *Staphylococcus capitis*[25,35]. Using *lsa*BGC, we discover novel variants of pyrazinones,
which encapsulate seven distinct GCFs (Figure 2a), that belong to species not commonly found
on human skin, such as *Staphylococcus felis* (GCF-56), the dominant species of
Staphylococcus on cat skin. While speciation had resulted in orthologous copies of *pznA* and
pznB being highly diverged in sequence, it is surprising to find that the surrounding genomic
context of the two genes is highly variable for all seven GCFs (Figure 2b), suggesting their
ancestral mobilization or relocation within or between genomes.
152
Application of *lsa*BGC-Divergence followed by Bayesian shrinkage analysis was then
used to systematically assess whether pyrazinones or other *Staphylococcus* and
Corynebacterium GCFs exhibited indications of HGT between species, as suggested by high
Beta-RD values (Figure S4cd; Table S6; Supplementary Methods and Results). Despite the
small size of pyrazinones, we do not find support for the frequent mobilization of any of the
seven GCFs encoding them across staphylococcal species. However, high Beta-RD posterior
ranges were observed for two NRPS associated GCFs from *Corynebacterium*. One of these
GCFs, predicted to encode for an NRPS-dependent siderophore[36], was found in several species
commonly isolated from skin[33], including *C. kefirresidentii*. Using results from *lsa*BGC-PopGene
for this GCF, which includes determination of a consensus gene order across BGC instances,
we find that the NRPS is highly conserved in sequence across species and found flanked by
mobile genetic elements, as previously reported in *Corynebacterium jeikeium*[36] (Figure 2cd).
The taxonomic breadth and high sequence conservation of this GCF suggest that it has
undergone recent HGT between distantly related species.
167
We next used *lsa*BGC-PopGene to identify genes under sweeping or balancing selection
by comprehensively calculating Tajima's D for each homolog group within a GCF across all four
genera, including *Micrococcus* and *Cutibacterium* (Table S7). We found that a hybrid
terpene/T3PKS GCF, in *Staphylococcus* exhibits the greatest signal for intra-species sweeping
selection and is one of four GCFs observed to be ancestral to the *S. epidermidis* clade (Figure
S5; Supplementary Methods and Results). The terpene component of this hybrid GCF encodes
for staphyloxanthin carotenoid production[37] and manual inspection determined that the T3PKS
prediction actually corresponds to hydroxymethylglutaryl-CoA (HMG-CoA) synthase, which is

involved in the mevalonate pathway for isoprenoid synthesis and is homologous to KS domains
of polyketide synthases[38]. In addition to these two BGC predictions, the regions in between and
flanking the staphyloxanthin encoding BGC and HMG-CoA synthase include several genes
important to staphylococcal virulence, some of which were found in multiple copies in certain
species (Figure S6ab). Further, while the five *crt* operon genes encoding for staphyloxanthin
were found in multiple species within the *S. epidermidis/aureus* clade, they are particularly
conserved in *S. aureus* as indicated by low Tajima's D values (Figure S6b).
183
184   **The virulence-associated carotenoid staphyloxanthin is ubiquitous across the**
185   ***Staphylococcus* genus**
186
Although staphyloxanthin is well characterized as an *S. aureus* virulence factor[39], we
were unable to identify a report of its distribution across the *Staphylococcus* genus, with only a
single study, to our knowledge, suggesting its production in another species[40]. *Isa*BGC analysis
revealed a total of five staphylococcal GCFs encoding for staphyloxanthin. Following GCF-3, the
second most prevalent GCF encoding staphyloxanthin, GCF-6, had a high Beta-RD distribution
(Figure S4c) and was widely distributed across divergent species in the genus (Figure 3a), as
well as *Mammaliicoccus,* the closest phylogenetic neighbors of staphylococci[41]. Constructing a
phylogeny based on a concatenated alignment of *crtM* and *crtN* revealed the appropriate
delineation of GCF-3 and GCF-6 despite their co-occurrence in the genomes of particular
species, such as *Staphylococcus warneri* (Figure 3b, S7a). Additional analyses showed that
GCF-6 has mobilized across species and can be found on plasmids (Figure 3bcd).
198
To assess whether carriage of the *crt* operon translates to staphyloxanthin production in
non-aureus species, we identified staphylococcal isolates cultivated from human skin with
whole-genome sequences generated by our lab that encode a staphyloxanthin related GCF. We
identified a rare instance of an *S. epidermidis* isolate with GCF-3 (only 1-3% of *S. epidermidis*
genomes encode GCF-3), strain LK1136, and an *S. warneri* isolate with both GCF-3 and GCF-
6, strain LK413. While we observe GCF-3 in all *S. warneri* genomes, the GCF is degraded
encoding only the dehydrosqualene synthase gene, *crtM*, and the dehydrosqualene desaturase
gene, *crtN* (Figure 3a; Table S8). However, roughly half of the available *S. warneri* genomes
(47.5%) encode GCF-6 in full and thus have two copies of *crtM* and *crtN*. We used long-read
sequencing to generate near-complete genomes for *S. epidermidis* LK1136 and *S. warneri*
LK413 and found that both strains encode GCF-3 within their chromosomes while GCF-6 is
encoded on a plasmid within *S. warneri*. (Figure S7b; Supplementary Methods and Results).
211
To test for phenotypic effects associated with strains carrying different staphyloxanthin
GCFs, we assessed pigmentation levels in strains of *S. aureus*, *S. epidermidis*, and *S. warneri*
with and without full *crt* operons (Table S9). Cells of *S. epidermidis* encoding GCF-3 are
pigmented compared to strains without the cluster (Figure 3e-f). Cells of *S. warneri* LK413
containing the complete *crt* operon (GCF-6) and additional copies of *crtMN* displayed the
strongest golden pigmentation, confirmed as staphyloxanthin by spectrophotometry[37,42] (Figure
3f).
219

_Isa_BGC provides a framework for mining for base-resolution novelty in BGCs

We have demonstrated how the _Isa_BGC suite can be used to identify homologous BGC instances into GCFs and subsequently uncover their evolutionary trends and phylogenetic distribution. With increasing metagenomic datasets becoming available, often from complex microbiomes, we developed _Isa_BGC-DiscoVary to directly profile BGCs and rapidly identify novel variation within GCF related genes from metagenomes (Figure S2; Supplementary Methods and Results). The _Isa_BGC framework thus allows users to first identify all instances of GCFs in genomic assemblies to build a comprehensive database of known alleles for GCF genes and subsequently apply _Isa_BGC-DiscoVary to find novel SNVs within genes which are absent in the available assemblies. Further, because _Isa_BGC-DiscoVary extracts the subset of reads supporting the presence of novel SNVs, exhaustive methods to assess the validity of metagenomic SNV calling, which would be computationally intensive and impractical to perform for the full set of reads from a metagenome, are made accessible for further validation of novel SNVs.

We first assessed the sensitivity and specificity of _Isa_BGC-DiscoVary and found high concordance (>96% overlap) between novel SNVs reported by it when compared to an assembly-based approach using whole-genome sequencing readsets for 132 _M. luteus_ isolates (Figure S8; Table S10, S11; Supplementary Methods and Results). Next, to demonstrate _Isa_BGC-DiscoVary's application in metagenomic datasets, its primary intended usage, we investigated the microdiversity of BGCs from _C. acnes_, the most abundant species at sebaceous sites, within healthy skin metagenomes generated by our lab[33]. We found that the BGC encoding for cutimycin, an anti-Staphylococcal thiopeptide, exhibited genomic signatures of horizontal, yet likely ancestral, acquisition by _C. acnes_ and is now largely restricted to isolates belonging to either clades IB or III(Figure 4ab, S9, S10a; Supplementary Methods and Results). Extensive metagenomic analysis demonstrated the reliability of _Isa_BGC-DiscoVary to type clade-specific alleles of genes within the cutimycin BGC in metagenomic datasets which correlated with strain presence inferences (Figure 4cd, S10bc; Table S12). We used _Isa_BGC to further investigate novel SNVs detected within cutimycin genes and found that many were from the metagenomes of a single individual and belonged to clade III _C. acnes_ (Figure 4efgh). Through observation that the majority of novel SNVs from this individual also corresponded to cytosine to thymine changes, we discovered that clade III _C. acnes_ have elevated rates of cytosine deamination, which coincides with the clade's carriage of a truncated uracil DNA glycosylase, the enzyme responsible for the repair of such deaminations (Figure 4ijk).

_Isa_BGC identifies novel SNVs within polyketide synthase catalytic domains

We determined the _C. kefirresidentii_ species complex is the most broadly distributed _Corynebacterium_ across our skin metagenomes (Figure S4b); however, only 22 genomic assemblies, representing five species, were available for the complex at the time of this analysis. This clade thus represents an ideal taxonomic group to examine novel BGC microdiversity using _Isa_BGC-DiscoVary. We ran _Isa_BGC-DiscoVary systematically on six GCFs identified in the species complex, largely encoding for unknown metabolites, and found 40,019 putatively novel SNV instances across 105 homolog groups and 109 metagenomic samples. Of

these, 5,474 instances were filtered because reads supporting their presence aligned with a
higher score to other regions in a concatenated database of all *Corynebacterium* genomes from
GTDB as compared to the reference database used for *lsa*BGC-DiscoVary. While ubiquitous, *C.*
kefirresidentii species complex members are present at very low abundances in some body-
sites where metagenomic assembly would struggle to construct their genomes (Figure S11a).
We performed sample-specific metagenomic assemblies[43] and found that at least 11,659 novel
SNV instances detected by *lsa*BGC-DiscoVary would be missed by assembly-based
investigation (Figure 5a; Supplementary Methods and Results).
272
To further investigate the novel SNVs detected, we focused on 68 homolog groups
overlapping or in close proximity to protocore regions of BGCs and further filtered based on
standardized coverage metrics, retaining 5,802 instances of 2,343 unique novel SNVs. Novel
SNVs were more likely to correspond to synonymous changes if they were found in multiple
metagenomic samples (Figure S11b). Further, metagenomic samples from the same subject at
different body sites and from the same body sites across different subjects were more likely to
share novel SNVs when compared to metagenomic samples from different body sites and
different subjects (p<1E-8; two-sided Wilcoxon rank sum test) (Figure S11c).
281
Finally, we expected and observed that the ratio of predicted synonymous to non-
synonymous novel SNVs increases at deeply conserved sites along homolog groups (Figure 5b;
Table S13; Supplemental Methods and Results). The homolog group with the greatest number
of novel SNVs, and novel, non-synonymous SNVs within highly conserved regions, was the
PKS gene responsible for synthesis of mycolic acid, a major component of the cell wall in most
Corynebacterium species recently shown to have immunomodulatory effects on the skin[26,44,45].
We identified and manually validated three non-synonymous SNVs along conserved sites of the
PKS gene, including two adjacent SNVs predicted to affect one codon within the ketoacyl
synthase domain, which catalyzes condensation[46], and could alter reaction kinetics[47] or scaffold
structure[48] (Figure 5c, S12; Table S14).
292

Discussion:

294
The *lsa*BGC suite packages several foundational, as well as novel, functionalities for
high-throughput evolutionary analysis of BGCs from any taxa in a structured framework. In
particular it includes several algorithms to overcome challenges when working with draft-quality
genomic assemblies. The suite's metagenomic mining functionalities further provide insight into
the presence and sequence diversity of a specific taxa's BGCs across diverse microbiomes.
This is a particularly important feature for rare BGCs and taxa with a limited numbers of
genomic assemblies available, such as taxa that are difficult to cultivate from complex
metagenomes. A current limitation of the *lsa*BGC framework includes its basis for using
chromosomally complete genomes to perform initial identification of BGCs and GCFs, which
can miss rare BGCs only observed in draft genomes. To alleviate this limitation, future
development of the suite will aim to allow direct incorporation of cataloged BGC and GCF
instances from publicly available databases or software[11,49,50]. We may also consider the use of

pan-genome alignment approaches to potentially increase the efficiency of *lsa*BGC for
metagenomic identification of novel variants within BGC genes[51].
309
Here, we apply *lsa*BGC's functionalities to major taxa commonly found in skin
microbiomes as a proof of concept to demonstrate the potential to advance understanding of the
diversity, evolutionary trends, and prevalence of both well studied and uncharacterized
secondary metabolites. For example, we uncovered the ubiquity of the carotenoid
staphyloxanthin across the genus of *Staphylococcus,* a molecule conferring the signature
golden pigmentation in *S. aureus*[52,53]. These findings raise the hypothesis that staphyloxanthin,
which has been identified primarily as a virulence factor conferring oxidative stress resistance in
the pathogen *S. aureus*[39,54], could similarly contribute to virulence in other staphylococcal
species or serve additional unknown functions. Supporting this hypothesis, we observe that one
staphyloxanthin encoding GCF appears ancestral to the skin-associated, multi-species clade
which features both *S. aureus* and *S. epidermidis*. The observation that multiple copies of *crt*
genes can be found within a single genome further suggests staphyloxanthin is of functional
importance since evolutionary retention of paralogous genes has previously been shown to
underlie ecological advantages[55,56]. Additionally, while this GCF is present in most skin-
associated species, it is absent in approximately 95% of *S. epidermidis* genomes, suggesting
large-scale loss within the species. It will thus be interesting to further explore how the loss of
staphyloxanthin, if it contributes to oxidative stress resistance within *S. epidermidis*, coincides
with the species' reputation as a beneficial symbiont of host skin[30,57].
328
Using the novel metagenomic mining functionality of *lsa*BGC, we identified novel SNVs
distinguishing uncultured *C. acnes* and *Corynebacterium* strains from isolates with genomes
available in public databases. With regards to *Corynebacterium*, we determined that some of
the most common skin residing species[29,33] belong to the *C. kefirresidentii* species complex,
which were widely distributed across skin metagenomes and highly abundant at moist body
sites. Because this species complex is vastly underrepresented in public genomic databases
and additional species and strains belonging to it likely remain to be discovered, application of
*lsa*BGC-DiscoVary led to the identification of thousands of novel SNVs within the BGCs of the
clade. Critically, some of these variants could have functional implications as they lie within
highly conserved regions of important domains for natural product biosynthesis.
339
The comprehensive nature of *lsa*BGC allows analyses to be applied to both public and
proprietary microbial genomic databases. Besides accelerating fundamental evolutionary
research of BGCs from diverse taxa, we envision *lsa*BGC will be particularly useful for
secondary metabolite discovery and mechanism of action studies. Specifically, we believe the
evolutionary insight gathered from *lsa*BGC can be used as prior information to guide algorithms
in the emerging field of using artificial intelligence to identify new antibiotics[58,59]. Finally, an
improved understanding of the relationship between intra-taxa genomic diversity and BGC
content novelty at the base-resolution using *lsa*BGC highlights which sub-lineages or sub-
clades bear the largest reservoir of untapped secondary metabolic potential.
349

350 ## Acknowledgments:

351
This work was supported by grants from the National Institutes of Health (NIAID U19AI142720
and NIGMS R35GM137828 [L.R.K]). The content is solely the responsibility of the authors and
does not necessarily represent the official views of the National Institutes of Health. The authors
gratefully acknowledge additional members of the Kalan lab, Dr. John-Demian Sauer, Dr. Jason
Kwan, Dr. Karthik Anantharaman, Dr. Caroline Grunenwald, Dr. Chase Clark, and Dr. Milton
Drott for helpful discussions.
358

Materials and Methods:

Code availability

The *Isa*BGC software suite (version 1.0) was developed in Python3 and R and is available on
Github at: https://github.com/Kalan-Lab/lsaBGC. Algorithmic descriptions of programs can be
found on the Github wiki and in the Supplementary Methods and Results document. A test
dataset pertaining to the *C. kefirresidentii* species complex for running the seven core programs
is provided on the Github repository, along with expected results.

An overview of the *Isa*BGC software suite

The first core program is *Isa*BGC-Cluster which clusters homologous and non-
fragmented instances of BGCs, identified by antiSMASH from chromosomally complete
genomic assemblies, into GCFs, similar to BiG-SCAPE[8].Users can then generate a report to
select for the most appropriate parameters for downstream analyses of interest. Because
antiSMASH groups neighboring BGCs into hybrid gene clusters if they are located close to one
another on the genome[6], *Isa*BGC-Refiner has the option to manually separate hybrid clusters
based on user-defined boundary homolog groups. Homologous instances of GCFs across all
available genomic assemblies for the taxa of interest can then be identified through *Isa*BGC-
Expansion (Figure S1b). Similar to ClusterFinder[9], *Isa*BGC-Expansion uses profile Hidden
Markov models (HMMs) to identify homologs of GCF associated genes in assemblies followed
by a classical HMM approach to define BGC regions and their boundaries. Unlike ClusterFinder,
*Isa*BGC-Expansion: (i) uses all coding genes associated with a GCF instead of being restricted
to core protein domains, (ii) is specifically designed to account for assembly fragmentation
through conditional assessment of predicted regions, and (iii) features a "quick mode" option for
fast detection of BGCs using Diamond[60].

The remaining four core programs are analytical and designed to be run on a single
GCF for ease of parallelization: *Isa*BGC-See, *Isa*BGC-PopGene, *Isa*BGC-Divergence, and
*Isa*BGC-DiscoVary. *Isa*BGC-See permits visualization of BGCs across a user-provided species
tree or a BGC based phylogeny and can handle multiple fragmented BGC instances from the
same genome, depicting them as neighboring leaves on phylogenies. *Isa*BGC-PopGene can
then infer evolutionary and population genetics statistics for each homolog group associated
with a GCF. It generates codon based alignments for each homolog group which can then be
used as input for *Isa*BGC-Divergence and *Isa*BGC-DiscoVary. *Isa*BGC-PopGene also
determines the consensus order and direction of homolog groups for a GCF across all BGC

instances, which provides intuitive assessment of its reports. *Isa*BGC-Divergence calculates a
comprehensive statistic between pairs of homologous BGCs from the same GCF but different
genomes, which we refer to as Beta-RD, that captures how similar the BGC sequences are
relative to what would be expected based on genome-wide similarity. Finally, *Isa*BGC-DiscoVary
is a multi-functional program which can mine raw sequencing readsets from metagenomes or
single genomes, for GCF instances and then reliably identify whether they possess novel single
nucleotide variants (SNVs) yet to be observed in available genomic assemblies for a taxa
(Figure S2).
404
405   **Usage of genomic and metagenomic datasets**
406
To overcome issues with misclassification and contamination in NCBI, we used GTDB
(release 202) [61], to query for reliable genomic assemblies and respective species designations
for the genera of *Corynebacterium*, *Cutibacterium*, *Micrococcus*, and *Staphylococcus* (Table
S1). For *Micrococcus*, we further included 132 additional genomes of the type species *M. luteus*
which our lab had sequenced (BioProject ID PRJNA803478) but were not yet uploaded to NCBI
to be included in the GTDB release used. These isolates were classified as *M. luteus* by GTDB-
tk[62] and exhibited > 95% ANI to the NCBI representative genome for the species *M. luteus*
NCTC 2665. A similar approach was used for gathering genome sets for species-level analyses
performed (Table S1). For the *C. kefirresidentii* species complex, an additional five genomes
from NCBI were gathered, classified using GTDB-tk[62], and incorporated into analysis if they
belonged to one of the five species designated as belonging to the complex.
418
Because a single representative per species as provided by GTDB might not adequately
capture the diversity of BGCs found across a genus, we instead employed a more granular
approach to select for representative genomes for a genus. We performed dereplication using
MASH[63] with an estimated ANI cutoff of 0.99 to pair similar genomes and dynamically select
representative genomes from each cluster based on assembly N50 (Table S1). This
dereplication was performed using the script popSizeAndSampleSelector.py provided in the
*Isa*BGC suite. Complete or chromosome level assemblies among the representative genomes
were run through IsaBGC-AutoProcess.py and IsaBGC-Cluster.py to identify the core set of
GCFs per genus. GCFs determined from the core set of genomes were comprehensively
identified in the remaining genomes using IsaBGC-AutoExpansion.py, including those which
were not deemed as representatives from the MASH based analysis to showcase scalability
(e.g. all 15,055 *Staphylococcus* and not just the 229 representative genomes of which 77
corresponded to complete/chromosome level assemblies). To expedite population genetics
analysis however, we only ran the downstream functional analytics of *Isa*BGC using the
representative genomes for each genus. For species-level analyses, all genomes available
were used for such downstream analyses and to comprehensively profile known alleles for
genes in BGCs, which was critical to enable metagenomic mining for base-resolution novelty
previously unseen in assemblies available for a species. For population level analyses within
species, strain delineations were performed using the popSizeAndSampleSelector.py script in
*Isa*BGC with strain groups determined from single-linkage clustering of pairs of genomes found
to exhibit ≥98% ANI and ≥80% genomic content similarity by FastANI[64].

For determining skin-associated species amongst the genera investigated in our study
and to demonstrate mining functionalities with *lsa*BGC-DiscoVary, we used a paired-end
metagenomics dataset sequenced on the Illumina NovaSeq where individual samples
corresponded to microbiome surveys of one of eight body sites for one of 34 participants,
collected in the state of Wisconsin (BioProject ID PRJNA763232)[33]. As previously described[33],
adapter removal, quality filtering, human sequence decontamination, and tandem repeat
removal were performed using fastp (v0.21.0)[65] and KneadData (v0.8.0)
(https://huttenhower.sph.harvard.edu/kneaddata/).

Genomic assessment for staphyloxanthin carriage in the lab isolate collection

We had previously constructed draft genomic assemblies for 44 *Staphylococcus* isolates
from skin (BioProject ID PRJNA803478; BioProject ID PRJNA830888). Illumina sequencing was
performed at the Microbial Genome Sequencing Center (MiGS). Briefly, assemblies were
generated using Unicycler[66] with default settings after quality and adapter trimming using
fastp[65]. The set of 44 *Staphylococcus* genomes were searched for carriage of staphyloxanthin
encoding GCFs identified in our study using *lsa*BGC-AutoExpansion (Table S9).

Library preparation, sequencing and genomic assembly

Reconstruction of Illumina draft assemblies for 132 M. luteus: We assessed the performance of
multiple programs in the *lsa*BGC suite using a set of genomic assemblies for 132 *M. luteus*
isolates which were cultured from skin. For these isolates, assemblies were reconstructed from
raw-reads (BioProject ID PRJNA803478) and were uploaded to NCBI's Genbank database
(Table S1). Briefly, Illumina sequencing reads were processed using fastp (v. 0.20.1)[65] to trim
for Poly-G tail artifacts and TrimGalore (v. 0.6.5)[67] was additionally used to detect and remove
adapters. Afterwards, assembly was performed using Unicycler (v. 0.4.6)[66] with default settings.

Construction of Illumina+Nanopore hybrid assemblies for staphyloxanthin producing S. warneri
LK413 and S. epidermidis *LK1136*: To learn the genomic location of staphyloxanthin encoding
GCFs in *S. epidermidis* LK1136 and *S. warneri* LK413, we performed additional long-read
sequencing using Oxford Nanopore Technologies (ONT). Library construction, using a PCR-free
ligation method, and sequencing were performed at the Microbial Genome Sequencing Center
(MiGS). Hybrid assemblies were constructed using a modified version of the Hybrid Assembly
workflow in seQuoia (https://github.com/broadinstitute/seQuoia), Illumina reads, used for
construction of the initial draft genomic assemblies for the isolates, were first reprocessed using
fastp (v. 0.20.1)[65] to trim for Poly-G tail artifacts and TrimGalore (v. 0.6.5)[67] was subsequently
used to detect and remove adapters. Similarly, ONT FASTQ reads, provided by MIGS after
basecalling, were filtered for potential adapters using PoreChop (v. 0.2.3)
(https://github.com/rrwick/Porechop) and filtered to retain only reads greater than 3 kb in length
as well as subsampled to 300 mb (while being inclusive of all reads greater than 20 kb) using
the script fastqfilter.py included in seQuoia. Afterwards, hybrid-assemblies were constructed
using Unicycler (v. 0.4.6)[66] with default settings and additional Pilon (v. 1.23)[68] polishing was

iteratively performed by aligning ONT reads[69] as well as the Illumina reads[70] to assemblies and
refining them, allowing for a maximum of 10 iterations. Quality assessment of the two near-
complete assemblies was performed using GAEMR (https://github.com/broadinstitute/GAEMR)
(Supplementary Methods and Results).

Experimental validation of staphyloxanthin production in non-aureus *Staphylococcus*

Isolates were grown overnight at 37°C on tryptone soy agar plates and inoculated into tryptone
soy broth at a concentration of 1×10^6 CFU/mL. Liquid cultures were grown in shaking conditions
at 37°C for 24 hours. 10 mL of the liquid culture was then pelleted and washed twice with
phosphate-buffered saline before pigment extraction in 1 mL of methanol at 55°C for 15 mins.
Spectral scans from 300 nm to 600 nm were taken using a plate reader (EPOCH2, Biotek,
Winooski, VT). Absorbance readings were blanked with methanol and normalized to liquid
culture cell density based on OD600. Spectra shown are representative of at least two biological
replicates.

MASH distance neighbor-joining tree construction

502 Average nucleotide identity estimates between pairs of genomes were computed with MASH[63]
503 and used to create a distance matrix. The distance matrix was then used to construct a
504 neighbor-joining tree with the ape library in R[71].

Ribosomal and BGC phylogeny construction

Ribosomal phylogeny construction: Ribosomal phylogenies for each species were constructed
based on a concatenated alignment of 16 ribosomal proteins[72]. HMMER (v3) was used to
search for profile HMMs of each ribosomal protein in the predicted proteomes of samples of
interest belonging to a particular taxonomic group. Afterwards, the best matching hits for each
sample based on alignment E-value were identified and used to construct individual alignments
for each ribosomal protein with the L-INS-I method in MAFFT[73]. Protein alignments were then
converted to codon alignments using PAL2NAL[74] and subsequently concatenated together.
Samples which featured $\geq$ 25% gaps in the concatenated alignment were filtered after which
core SNV sites were extracted and used for constructing a maximum likelihood phylogeny with
RAxML[75], specifying a GTRCAT model and performing 100 bootstraps. Where needed,
ribosomal phylogenies were subset to showcase relationships between select samples using
PareTree (http://emmahodcroft.com/PareTree.html ; v1.0.2).

Staphyloxanthin phylogeny construction: A maximum likelihood phylogeny was constructed from
a concatenated protein alignment of CrtM and CrtN amongst representative *Staphylococcus*
genomes to understand the relationship between different GCFs encoding for staphyloxanthin.
Proteins designated as HG0001876 (CrtM) and HG0001904 (CrtN) were gathered for each
staphyloxanthin encoding GCF found in each sample and aligned using the L-INS-I method in
MAFFT[73]. To account for assembly fragmentation resulting in multiple proteins from a sample
being assigned to the same homolog group for a specific GCF, a consensus sequence was

determined and multi-allele sites were replaced as gaps. A concatenated alignment of such
consensus sequences was subsequently constructed and used for phylogeny construction with
RAxML using a PROTCAT model with JM selected as the best amino-acid replacement matrix
through the automatic maximum likelihood model selector[75]. The phylogeny of CrtMN
sequences exhibited high correspondence to GCF delineations with the exception of a single *S.
lugdunensis* BGC which is likely misclassified as GCF-17 and should be GCF-3. Upon further
examination, it was identified that the reason for the misclassification was likely due to the
absence of mevalonate pathway genes, classified as a type III polyketide synthase (T3PKS),
within *S. lugdunensis* which are found downstream of the *crt* operon in all other species with
GCF-3 (Figure S7b).
538
539 *Pyrazinone phylogeny construction:* A maximum likelihood phylogeny was constructed from a
540 concatenated protein alignment of PznA and PznB amongst representative *Staphylococcus*
541 genomes using the same methods as described for phylogeny construction from CrtMN
542 proteins.
543
544 **Designation of species belonging to *Mammaliicoccus***
545
The *Staphylococcus_A* genus designation in GTDB R202 was largely concordant with the
recent reclassification of five species from *Staphylococcus* to *Mammaliicoccus*[41]. In the latest
GTDB R207 release, this genus has now been reclassified to *Mammaliicoccus*. The recent
update to taxonomic names also includes the reclassification of species *S. pasteuri_A* to *M.
fleurettii*. Notably, we do not regard *S. schleiferi* as part of *Mammaliicoccus* because it is
categorized as *Staphylococcus* by GTDB (both R202 and R207) and groups within the
Staphylococcus genus in our ribosomal protein based maximum-likelihood phylogeny. Similar to
our assessment, the reclassification of this species to *Mammaliicoccus* was recently
questioned[76].
555
556 **Ubiquity assessment of skin residing species and strains of *Staphylococcus* and**
557 ***Corynebacterium***
558
We searched for the presence of representative *Staphylococcus* and *Corynebacterium*
strains in 270 skin metagenomes from 34 participants across 8 body sites using StrainGST[34].
To construct the database of distinct representative genomes for each genus, the protocol
described in the StrainGE documentation was followed. *S. aureus* genomes 16405 and 278
were dropped from the analysis because at the time of downloading them, the plasmid and
chromosome were switched. Briefly, genomes were downloaded from NCBI using ncbi-genome-
download (https://github.com/kblin/ncbi-genome-download) and dereplication was performed
using built-in StrainGE functionalities. As some representative genomes were not part of the
GTDB release R202, GTDB-tk[62] was used to annotate such genomes and perform species
classification. We considered a species as skin-associated if it was found in at least 10 of the
270 skin metagenomes, regardless of the abundance (Table S4). Certain skin-associated
species from each genus were also identified in the 19 negative control metagenomic samples,
which included air-swab, bench, desk, and door-handle samplings (Table S4). Because we

expect heavy overlap in species content between these environmental samples and skin
microbiome samples, we still considered such species as skin-associated if they were found in
10 or more skin metagenomes.
We found the majority of skin-associated *Staphylococcus* belonged to the *S.*
epidermidis/aureus clade, as has previously been noted[38]. Similarly, most of the skin-associated
Corynebacterium belonged to *C. kefirresidentii* and it's close neighboring species (including *C.*
kefirresidentii, *C. aurimucosum* type E, *C.* sp. 900539985, *C. tuberculostearicum*, *C.*
tuberculostearicum type C), which we denote in this study as the *C. kefirresidentii* species
complex. The minimal ANI separating genomes belonging to this species complex was
determined to be 88.59% using FastANI[64]. To assess whether any of the 24 new
Corynebacterium species reported by Saheb Kashaf et al. 2022[32] belonged to this species
complex, FastANI was used to look at their ANI similarity to genomes from the clade and regard
them as additional members if they exhibited 88.59% ANI or greater. Of note, many additional
Corynebacterium species have been routinely isolated from the skin[33] which we did not regard
as skin-associated based on our criteria in this study. In particular, the species of *C.*
amycolatum (detected in 4 metagenomes), *C. kroppenstedtii* (detected in 4 metagenomes)*,* and
C. simulans (detected in 5 metagenomes) are commonly found on skin.
To estimate the relative abundance of *Cutibacterium* (see section "Identification of
Cutibacterium Strains within Metagenomes and Determination of Near-Fixed Sites
Differentiating Cutimycin Biosynthesis Alleles between Clade I and Clade III *C. acnes*") and
Corynebacterium strains within metagenomic samples we normalized the estimated relative
abundances for strains (*rapct*) by the estimated relative abundance of the genus (*pan%*) as
reported by StrainGST[34].
595
596 **Identification of homolog groups associated with skin residing species of**
597 ***Staphylococcus* and *Corynebacterium***
598
We developed a script called crawlingFisher.py, provided in the *lsa*BGC suite, which
tests for enrichment or depletion of homolog group presence amongst genomes in focal clades
using Fisher's exact test. Provided with a phylogenetic tree and matrix specifying the presence
of homolog groups across genomes, the program automatically performs such tests for each
homolog group at each inner-node of the phylogeny. Multiple testings correction was performed
comprehensively at the end using Benjamini-Hochberg false discovery rate and cases were
considered statistically significant if the adjusted p-value was less than 0.05. Results were
further filtered to retain only homolog groups which were present in at least 80% of genomes
under the focal node and at most 20% of other genomes or, alternatively, at most 20% of
genomes under the focal node and at least 80% of other genomes. Finally, only results
pertaining to focal nodes where at least 80% of genomes were classified as skin-associated
(see section "Ubiquity Assessment of Skin Residing Species and Strains of *Staphylococcus* and
Corynebacterium") were reported (Table S5).
612
613 **Assessing the divergence of BGCs relative to their background genomic contexts**
614

BGC divergence does not always track with background genomic divergence and
speciation[13]. The program *Isa*BGC-Divergence allows for easy assessment of how BGC
divergence compares to whole-genome divergence, primarily through computing a statistic we
refer to as Beta-Relative-Divergence (Beta-RD) between pairs of genomes for a single GCF.
The Beta-RD statistic is simply the average amino or nucleotide identity between homologous
instances of proteins for a GCF observed in two genomes normalized by the average amino or
nucleotide identity observed between the full genomes. *Isa*BGC allows users to specify whether
amino or nucleotide divergence should be used and automated workflows (see section
Automated Workflows for Running *Isa*BGC) can be used to compute pairwise AAI or ANI
between genomes using CompareM (https://github.com/dparks1134/CompareM) or FastANI[64],
respectively.
To better understand how distributions of Beta-RD differ across GCFs, in this study, we
applied a Bayesian hierarchical model to perform shrinking analysis and appropriately account
for differences in the number of genomes GCFs are found (Figure S4cd; Table S6). For
computational efficiency, a maximum of 500 Beta-RD observations were used for each GCF.
Posterior predictions for Beta-RD were made using rstan (Stan Development Team 2021) with
the following hierarchical model:

$$\overline{\mu} \sim Norm(\mu_0, \sigma_0^2)$$
$$\tau \sim half - t_7(1)$$
$$\mu_1...\mu_J \sim Norm(\overline{\mu}, \tau)$$
$$\sigma \sim half - t_7(1)$$
$$y_j \sim Norm(\mu_j, \sigma^2)$$

632
where for each GCF j, the posterior Beta-RD distribution is sampled as y_j using Markov chain
Monte Carlo with four chains, each featuring 2,000 iterations, including a burnin of 1,000
iterations. A hierarchical prior was specified for $_j$, which represents a GCF-specific baseline.
The values of $_0$ and σ_0 were set to the respective mean and standard deviation of raw Beta-RD
values across all GCFs. R and STAN code used for this analysis are available within the scripts
subdirectory of the *Isa*BGC software package.
639
640   **Scrutinization of putative novel variants for *Isa*BGC-DiscoVary application to cutimycin**
641   **in *C. acnes* and the comprehensive set of GCFs for the *C. kefirresidentii* species complex**
642
By default, IsaBGC-DiscoVary.py only considers reads aligning at 95% identity to
reference gene sequences and avoids reporting novel variants on genes if either annotation or
coverage (relative to other genes in the BGC context) suggests they might be MGEs. The
program also automatically extracts the subset of reads supporting the existence of novel
variants, which can then be further screened with high stringency filters using otherwise
computationally expensive methods. This can involve mapping reads supporting putative novel
SNVs to a representative database of whole-genomes to observe for higher quality alignments
or checking whether reads are classified as a particular taxonomy by Kraken2 (v2.0.8-beta)[77].
As described previously[33], we used a customized database for Kraken2 analysis constructed

from chromosomally complete bacterial, viral, archaeal, fungal, protozoan genomes as well as
the human genome available on RefSeq, where plasmid sequences were distinguished to
alleviate potential misclassifications of reads to a particular taxonomy.
655
For reads supporting putative novel variants in the cutimycin BGC of *C. acnes*, we
required them to be classified as *Cutibacterium* or *Priopionibacterium* by Kraken2[77] and to not
map with a higher alignment score to a comprehensive database of all *Cutibacterium* genomes
than was observed for the alignment to the reference gene sequence in lsaBGC-DiscoVary.py
(Figure S2c). We generally do not recommend screening variant-supporting reads for an
appropriate taxonomic match with Kraken2 as it can generally lead to false negatives (on
average only 92.8% of read-pairs per sample were taxonomically classified and
misclassification can also occur; Figure S12), but performed this here to provide further
assurance that novel variants discovered are reliable. Standardized coverage metrics within
lsaBGC-DiscoVary.py's novel SNV report were further used to conservatively filter sites and
variants if they exhibited site or allele-specific coverage values greater than two median
absolute deviations from the median coverage for the entire BGC.
668
For reads supporting putative novel variants within GCFs of the *C. kefirresidentii* species
complex, we required them to not align with a higher alignment score to a comprehensive
database of all *Corynebacterium* genomes. We did not require reads to be classified by Kraken2
as *Corynebacterium* because genomes from the species complex are likely to be poorly
represented in the Kraken2 database. We did however further validate that reads supporting the
existence of variants at conserved sites in the mycolic acid biosynthesis gene mapped best to
the *C. kefirresidentii* species and confirmed our suspicion that they could be misclassified by
Kraken2 given the representation in the underlying database (Figure S12). This *ad hoc* analysis
was performed using paired-end information when mapping in Bowtie2 and accounting for
concordant read pair alignments. While many reads supporting the existence of novel SNVs in
the biosynthesis gene were classified as *C. acnes* by Kraken2, we found that none of the reads
concordantly aligned to the comprehensive *Cutibacterium* genomes Bowtie 2 database whereas
85.07% concordantly aligned to the comprehensive *Corynebacterium* genomes Bowtie 2
database.
683
Note, the reasoning behind mapping SNV supporting reads to comprehensive genomic
databases to purge false positive SNV calls, is that if a read truly aligns best to the BGC-
associated homolog group where it was used to call a SNV, then it should at most generate a
mapping to this comprehensive database of equal mapping quality. We align paired-end reads
individually as was performed in lsaBGC-DiscoVary.py to have mapping scores be comparable
with those from the original alignment. Furthermore, because our initial database within lsaBGC-
DiscoVary.py consisted of allelic representatives of homolog groups, we realized that reads
mapping to the start and end of genes could result in a lower mapping score than if a larger
genomic context was provided. To ensure we do not remove such reads mapping to gene
edges, scrutinizeNovelSNVSupport.py, the program we developed to perform Kraken 2 and
genomic database alignment based filtering of novel SNV reports from lsaBGC-DiscoVary.py,
also compares the reference sequences when it finds a better Bowtie 2 mapping for SNV-

supporting reads in genomic databases. If the reference sequence in the genomic database
encompasses the reference sequence in the original alignment to the lsaBGC-DiscoVary.py
homolog group database and the read was previously noted to map to a gene edge, then reads
are still retained and SNV read support is not decremented. Five retained and supportive reads
are required for an SNV to be featured in the filtered report produced by the program.
701
702 ***lsa*BGC-DiscoVary based analyses of BGCs from the *C. kefirresidentii* species complex:**
703
704   For the following analyses, we used 5,802 instances of 2,343 unique novel SNVs found
705 in 68 homolog groups which were core or adjacent to core biosynthesis machinery of BGCs
706 from the *C. kefirresidentii* species complex. These novel SNVs had site and allele-specific
707 coverage values less than two median absolute deviations from the median coverage for the
708 entire BGC.
709
Comparing SNVs shared across body sites and subjects: To comprehensively assess whether
novel SNVs were shared more frequently for microbiomes of the same body site or from the
same individual, we performed a pairwise, multi-iteration resampling analysis. For each pair of
metagenomes, where each sample had at least 30 novel SNVs, we calculated the average
Jaccard similarity across 1,000 simulations in which we randomly drew sets of 30 SNVs from
each metagenomic sample to control for sequencing depth. Pairs of metagenomes were
classified as one of three categories: "different body site, different participant", "different body
site, same participant", and "same body site, different participant" (Figure S11c). A two-sided
Wilcoxon rank sum test for differences between the three distributions of average Jaccard
similarities revealed that there was a statistically significant difference between "different body
site, same participant' and "same body site, different participant" (p=6.01E-3). Of greater
relevance, there were statistically strong significant differences between the "different body site,
different participant" distribution and the two other distributions: "different body site, same
participant" and "same body site, different participant" (p=3.67E-09 and p=1.39E-116,
respectively). Thus, pairs of metagenomes were more likely to share novel SNVs if they were
either from the same body site or same participant compared to if they were from different
participants and different body sites. Two metagenomic samples for participant S002 were not
used because they corresponded to a resampling of two of their body sites and we did not have
enough of such samples to pursue a temporal analysis. We assessed this testing using different
considerations/cutoffs for: (i) the number of novel SNVs needed for a metagenome to be
considered in the analysis and (ii) whether or not to account for singleton SNVs. In all cases,
statistical testing for differences between distributions of pairwise metagenome categories
yielded similar conclusions.
733
Assessing trends for novel SNV predicted synonymous to non-synonymous rate with
metagenome ubiquity and site conservation: Using reports from lsaBGC-DiscoVary.py, we
performed a systematic prediction of whether novel SNVs identified on the 68 homolog groups
nearby or overlapping with protocore regions of BGCs in *C. kefirresidentii* corresponded to
synonymous or non-synonymous substitutions. As described in other sections, SNV alleles
were compared to corresponding alleles observed at the site on the reference gene sequences

which SNVs were called upon. We aimed to check whether the rate of synonymous to non-
synonymous novel SNVs showed a relationship to: (i) how common SNVs were across
metagenomic samples and (ii) conservation levels across protein sequences.
743
Novel SNV predicted synonymous to non-synonymous rate increases with SNV metagenome
ubiquity amongst metagenomic samples: The occurrence of novel SNVs, a novel allele
observed at a specific site in the multiple sequence codon alignment of a particular homolog
group, was tabulated across metagenomic samples. While novel SNVs found in a single
metagenome were only twice as likely to correspond to a synonymous change as opposed to a
non-synonymous change, more prevalent novel SNVs, found in ten or more samples, were
eight times more likely to correspond to a synonymous change (Figure S11b).
751
Novel SNV predicted synonymous to non-synonymous rate increases at conserved sites in
protein sequences: We searched RefSeq's bacterial NR database for remote homologs of the
68 homolog groups within or adjacent to protocore regions of BGCs. The top 20 homologs
which belonged to classified bacterial species outside of the *Corynebacterium* genus or
Corynebacteriales family were identified using HMMER3 with profile-HMMs gathered from
*lsa*BGC-Expansion for each homolog group and an E-value threshold of 1E-20 (Table S13). The
sequences of these representative homologs were added to protein alignments for homolog
groups in *lsa*BGC-PopGene results using the '--add' option in MAFFT[73]. Afterwards, we used
the alignments for scoring protein sequence conservation with Jensen-Shannon divergence[78].
Conservation scores were used to determine conservation percentiles of sites along protein
alignments for individual homolog groups. Rates of predicted synonymous to non-synonymous
novel SNVs were calculated for each conservation percentile across all homolog groups. For
the three SNVs predicted to encode for non-synonymous differences and also in the top five
percentile of conserved sites along the mycolic acid PKS, we inspected their codon contexts to
check that there were no additional variants which, in aggregate, might result in a synonymous
change (Table S14).
768

769 References:

1. Pham, J. V. *et al.* A Review of the Microbial Production of Bioactive Natural Products and

Biologics. *Front. Microbiol.* **10**, 1404 (2019).

2. Watve, M. G., Tickoo, R., Jog, M. M. & Bhole, B. D. How many antibiotics are produced by

the genus Streptomyces? *Arch. Microbiol.* **176**, 386–390 (2001).

3. Chevrette, M. G. *et al.* The antimicrobial potential of Streptomyces from insect

microbiomes. *Nat. Commun.* **10**, 516 (2019).

4. Waterworth, S. C. *et al.* Horizontal Gene Transfer to a Defensive Symbiont with a Reduced

Genome in a Multipartite Beetle Microbiome. *MBio* **11**, (2020).

5.   Drott, M. T. *et al.* Microevolution in the pansecondary metabolome of Aspergillus flavus and

its potential macroevolutionary implications for filamentous fungi. *Proc. Natl. Acad. Sci. U.*

S. A. **118**, (2021).

6.   Steinke, K., Mohite, O. S., Weber, T. & Kovács, Á. T. Phylogenetic Distribution of

Secondary Metabolites in the Bacillus subtilis Species Complex. *mSystems* **6**, (2021).

7.   Blin, K. *et al.* antiSMASH 6.0: improving cluster detection and comparison capabilities.

Nucleic Acids Res. **49**, W29–W35 (2021).

8.   Navarro-Muñoz, J. C. *et al.* A computational framework to explore large-scale biosynthetic

diversity. *Nat. Chem. Biol.* **16**, 60–68 (2020).

9.   Cimermancic, P. *et al.* Insights into secondary metabolism from a global analysis of

prokaryotic biosynthetic gene clusters. *Cell* **158**, 412–421 (2014).

10. Hannigan, G. D. *et al.* A deep learning genome-mining strategy for biosynthetic gene

cluster prediction. *Nucleic Acids Res.* **47**, e110 (2019).

11. Kautsar, S. A., van der Hooft, J. J. J., de Ridder, D. & Medema, M. H. BiG-SLiCE: A highly

scalable tool maps the diversity of 1.2 million biosynthetic gene clusters. *Gigascience* **10**,

(2021).

12. Emms, D. M. & Kelly, S. OrthoFinder: phylogenetic orthology inference for comparative

genomics. *Genome Biol.* **20**, 238 (2019).

13. Chase, A. B., Sweeney, D., Muskat, M. N., Guillén-Matus, D. G. & Jensen, P. R. Vertical

Inheritance Facilitates Interspecies Diversification in Biosynthetic Gene Clusters and

Specialized Metabolites. *MBio* **12**, e0270021 (2021).

14. Felnagle, E. A., Rondon, M. R., Berti, A. D., Crosby, H. A. & Thomas, M. G. Identification of

the biosynthetic gene cluster and an additional gene for resistance to the antituberculosis

drug capreomycin. *Appl. Environ. Microbiol.* **73**, 4162–4170 (2007).

15. Del Carratore, F. *et al.* Computational identification of co-evolving multi-gene modules in

microbial biosynthetic gene clusters. *Commun Biol* **2**, 83 (2019).

16. Crits-Christoph, A., Bhattacharya, N., Olm, M. R., Song, Y. S. & Banfield, J. F. Transporter

genes in biosynthetic gene clusters predict metabolite characteristics and siderophore

activity. *Genome Res.* (2020) doi:10.1101/gr.268169.120.

17. Crits-Christoph, A., Diamond, S., Butterfield, C. N., Thomas, B. C. & Banfield, J. F. Novel

soil bacteria possess diverse genes for secondary metabolite biosynthesis. *Nature* **558**,

440–444 (2018).

18. Sugimoto, Y. *et al.* A metagenomic strategy for harnessing the chemical repertoire of the

human microbiome. *Science* **366**, (2019).

19. Andreu, V. P. *et al.* A systematic analysis of metabolic pathways in the human gut

microbiota. *bioRxiv* 2021.02.25.432841 (2021) doi:10.1101/2021.02.25.432841.

20. Pereira-Flores, E. *et al.* Mining metagenomes for natural product biosynthetic gene

clusters: unlocking new potential with ultrafast techniques. *bioRxiv* 2021.01.20.427441

(2021) doi:10.1101/2021.01.20.427441.

21. Chevrette, M. G. & Handelsman, J. Needles in haystacks: reevaluating old paradigms for

the discovery of bacterial secondary metabolites. *Nat. Prod. Rep.* **38**, 2083–2099 (2021).

22. Zipperer, A. *et al.* Human commensals producing a novel antibiotic impair pathogen

colonization. *Nature* **535**, 511–516 (2016).

23. Claesen, J. *et al.* A Cutibacterium acnes antibiotic modulates human skin microbiota

composition in hair follicles. *Sci. Transl. Med.* **12**, (2020).

24. Nakatsuji, T. *et al.* Antimicrobials from human skin commensal bacteria protect against

Staphylococcus aureus and are deficient in atopic dermatitis. *Sci. Transl. Med.* **9**, (2017).

25. Wyatt, M. A. *et al.* Staphylococcus aureus nonribosomal peptide secondary metabolites

regulate virulence. *Science* **329**, 294–296 (2010).

26. Ridaura, V. K. *et al.* Contextual control of skin immunity and inflammation by

Corynebacterium. *J. Exp. Med.* **215**, 785–799 (2018).

27. Swaney, M. H. & Kalan, L. R. Living in Your Skin: Microbes, Molecules, and Mechanisms.

Infect. Immun. **89**, (2021).

28. Grice, E. A. *et al.* Topographical and temporal diversity of the human skin microbiome.

Science **324**, 1190–1192 (2009).

29. Oh, J. *et al.* Temporal Stability of the Human Skin Microbiome. *Cell* **165**, 854–866 (2016).

30. Byrd, A. L., Belkaid, Y. & Segre, J. A. The human skin microbiome. *Nat. Rev. Microbiol.* **16**,

143–155 (2018).

31. Acosta, E. M. *et al.* Bacterial DNA on the skin surface overrepresents the viable skin

microbiome. *bioRxiv* 2021.08.16.455933 (2021) doi:10.1101/2021.08.16.455933.

32. Saheb Kashaf, S. *et al.* Integrating cultivation and metagenomics for a multi-kingdom view

of skin microbiome diversity and functions. *Nat Microbiol* **7**, 169–179 (2022).

33. Swaney, M. H., Sandstrom, S. & Kalan, L. R. Cobamide sharing drives skin microbiome

dynamics. *bioRxiv* 2020.12.02.407395 (2021) doi:10.1101/2020.12.02.407395.

34. van Dijk, L. R. *et al.* StrainGE: a toolkit to track and characterize low-abundance strains in

complex microbial communities. *Genome Biol.* **23**, 74 (2022).

35. Zimmermann, M. & Fischbach, M. A. A family of pyrazinone natural products from a

conserved nonribosomal peptide synthetase in Staphylococcus aureus. *Chem. Biol.* **17**,

925–930 (2010).

36. Tauch, A. *et al.* Complete genome sequence and analysis of the multiresistant nosocomial

pathogen Corynebacterium jeikeium K411, a lipid-requiring bacterium of the human skin

flora. *J. Bacteriol.* **187**, 4671–4682 (2005).

37. Pelz, A. *et al.* Structure and biosynthesis of staphyloxanthin from Staphylococcus aureus. *J.*

Biol. Chem. **280**, 32493–32498 (2005).

38. Balibar, C. J., Shen, X. & Tao, J. The mevalonate pathway of Staphylococcus aureus. *J.*

Bacteriol. **191**, 851–861 (2009).

39. Clauditz, A., Resch, A., Wieland, K.-P., Peschel, A. & Götz, F. Staphyloxanthin plays a role

in the fitness of Staphylococcus aureus and its ability to cope with oxidative stress. *Infect.*

Immun. **74**, 4950–4953 (2006).

40. Seel, W. *et al.* Carotenoids are used as regulators for membrane fluidity by Staphylococcus

xylosus. *Sci. Rep.* **10**, 330 (2020).

41. Madhaiyan, M., Wirth, J. S. & Saravanan, V. S. Phylogenomic analyses of the

Staphylococcaceae family suggest the reclassification of five species within the genus

Staphylococcus as heterotypic synonyms, the promotion of five subspecies to novel

species, the taxonomic reassignment of five Staphylococcus species to Mammaliicoccus

gen. nov., and the formal assignment of Nosocomiicoccus to the family Staphylococcaceae.

Int. J. Syst. Evol. Microbiol. **70**, 5926–5936 (2020).

42. Dong, P.-T. *et al.* Photolysis of Staphyloxanthin in Methicillin-Resistant Staphylococcus

aureus Potentiates Killing by Reactive Oxygen Species. *Adv. Sci.* **6**, 1900030 (2019).

43. Li, D., Liu, C.-M., Luo, R., Sadakane, K. & Lam, T.-W. MEGAHIT: an ultra-fast single-node

solution for large and complex metagenomics assembly via succinct de Bruijn graph.

Bioinformatics **31**, 1674–1676 (2015).

44. Gande, R. *et al.* Acyl-CoA carboxylases (accD2 and accD3), together with a unique

polyketide synthase (Cg-pks), are key to mycolic acid biosynthesis in Corynebacterianeae

such as Corynebacterium glutamicum and Mycobacterium tuberculosis. *J. Biol. Chem.* **279**,

44847–44857 (2004).

45. Portevin, D. *et al.* A polyketide synthase catalyzes the last condensation step of mycolic

acid biosynthesis in mycobacteria and related organisms. *Proc. Natl. Acad. Sci. U. S. A.*

101, 314–319 (2004).

46. Chen, Y. *et al.* Structural classification and properties of ketoacyl synthases. *Protein Sci.*

20, 1659–1667 (2011).

47. Klaus, M., Buyachuihan, L. & Grininger, M. Ketosynthase Domain Constrains the Design of

Polyketide Synthases. *ACS Chem. Biol.* **15**, 2422–2432 (2020).

48. Medema, M. H., Cimermancic, P., Sali, A., Takano, E. & Fischbach, M. A. A systematic

computational analysis of biosynthetic gene cluster evolution: lessons for engineering

biosynthesis. *PLoS Comput. Biol.* **10**, e1004016 (2014).

49. Palaniappan, K. *et al.* IMG-ABC v.5.0: an update to the IMG/Atlas of Biosynthetic Gene

Clusters Knowledgebase. *Nucleic Acids Res.* **48**, D422–D430 (2019).

50. Kautsar, S. A. *et al.* MIBiG 2.0: a repository for biosynthetic gene clusters of known

function. *Nucleic Acids Res.* **48**, D454–D458 (2020).

51. Garrison, E. *et al.* Variation graph toolkit improves read mapping by representing genetic

variation in the reference. *Nat. Biotechnol.* **36**, 875–879 (2018).

52. Kloos, W. E. & Schleifer, K. H. Isolation and Characterization of Staphylococci from Human

Skin II. Descriptions of Four New Species: Staphylococcus warneri, Staphylococcus capitis,

Staphylococcus hominis, and Staphylococcus simulans. *Int. J. Syst. Bacteriol.* **25**, 62–79

(1975).

53. Becker, K., Heilmann, C. & Peters, G. Coagulase-negative staphylococci. *Clin. Microbiol.*

Rev. **27**, 870–926 (2014).

54. Liu, C.-I. *et al.* A cholesterol biosynthesis inhibitor blocks Staphylococcus aureus virulence.

Science **319**, 1391–1394 (2008).

55. Blount, Z. D., Barrick, J. E., Davidson, C. J. & Lenski, R. E. Genomic analysis of a key

innovation in an experimental Escherichia coli population. *Nature* **489**, 513–518 (2012).

56. Ghosh, S. & O'Connor, T. J. Beyond Paralogs: The Multiple Layers of Redundancy in

Bacterial Pathogenesis. *Front. Cell. Infect. Microbiol.* **7**, 467 (2017).

57. Otto, M. Staphylococcus epidermidis--the 'accidental' pathogen. *Nat. Rev. Microbiol.* **7**,

555–567 (2009).

58. Stokes, J. M. *et al.* A Deep Learning Approach to Antibiotic Discovery. *Cell* **180**, 688–

702.e13 (2020).

59. Melo, M. C. R., Maasch, J. R. M. A. & de la Fuente-Nunez, C. Accelerating antibiotic

discovery through artificial intelligence. *Commun Biol* **4**, 1050 (2021).

60.   Buchfink, B., Xie, C. & Huson, D. H. Fast and sensitive protein alignment using DIAMOND.

Nat. Methods **12**, 59–60 (2014).

61.   Parks, D. H. *et al.* GTDB: an ongoing census of bacterial and archaeal diversity through a

phylogenetically consistent, rank normalized and complete genome-based taxonomy.

Nucleic Acids Res. (2021) doi:10.1093/nar/gkab776.

62.   Chaumeil, P.-A., Mussig, A. J., Hugenholtz, P. & Parks, D. H. GTDB-Tk: a toolkit to classify

genomes with the Genome Taxonomy Database. *Bioinformatics* **36**, 1925–1927 (2019).

63.   Ondov, B. D. *et al.* Mash: fast genome and metagenome distance estimation using

MinHash. *Genome Biol.* **17**, 132 (2016).

64.   Jain, C., Rodriguez-R, L. M., Phillippy, A. M., Konstantinidis, K. T. & Aluru, S. High

throughput ANI analysis of 90K prokaryotic genomes reveals clear species boundaries.

Nat. Commun. **9**, 1–8 (2018).

65.   Chen, S., Zhou, Y., Chen, Y. & Gu, J. fastp: an ultra-fast all-in-one FASTQ preprocessor.

Bioinformatics **34**, i884–i890 (2018).

66.   Wick, R. R., Judd, L. M., Gorrie, C. L. & Holt, K. E. Unicycler: Resolving bacterial genome

assemblies from short and long sequencing reads. *PLoS Comput. Biol.* **13**, e1005595

(2017).

67.   Krueger, F., James, F., Ewels, P., Afyounian, E. & Schuster-Boeckler, B.

FelixKrueger/TrimGalore: v0.6.7 - DOI via Zenodo. (2021). doi:10.5281/zenodo.5127899.

68.   Walker, B. J. *et al.* Pilon: an integrated tool for comprehensive microbial variant detection

and genome assembly improvement. *PLoS One* **9**, e112963 (2014).

69.   Li, H. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* **34**, 3094–

3100 (2018).

70.   Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows–Wheeler

transform. *Bioinformatics* **25**, 1754–1760 (2009).

71.   Paradis, E., Claude, J. & Strimmer, K. APE: Analyses of Phylogenetics and Evolution in R

language. *Bioinformatics* **20**, 289–290 (2004).

72. McDaniel, E. A., Anantharaman, K. & McMahon, K. D. metabolisHMM: Phylogenomic

analysis for exploration of microbial phylogenies and metabolic pathways. *bioRxiv*

2019.12.20.884627 (2020) doi:10.1101/2019.12.20.884627.

73. Katoh, K. & Standley, D. M. MAFFT multiple sequence alignment software version 7:

improvements in performance and usability. *Mol. Biol. Evol.* **30**, 772–780 (2013).

74. Suyama, M., Torrents, D. & Bork, P. PAL2NAL: robust conversion of protein sequence

alignments into the corresponding codon alignments. *Nucleic Acids Res.* **34**, W609–12

(2006).

75. Stamatakis, A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large

phylogenies. *Bioinformatics* **30**, 1312–1313 (2014).

76. Kania, S. A. Reclassification of Staphylococcus schleiferi by Madhaiyan et al. lacks key

supporting data. *Int. J. Syst. Evol. Microbiol.* **72**, (2022).

77. Wood, D. E., Lu, J. & Langmead, B. Improved metagenomic analysis with Kraken 2.

Genome Biol. **20**, 1–13 (2019).

78. Capra, J. A. & Singh, M. Predicting functionally important residues from sequence

conservation. *Bioinformatics* **23**, 1875–1882 (2007).

Figure 1: *Isa*BGC offers an efficient and sensitive means for GCF homology detection. **a)** A schematic overview of the full *Isa*BGC suite. **b)** Neighbor-joining trees were constructed for each of the four genera based on pairwise MASH ANI estimates between genomes and heatmaps showcase the presence of GCFs found in ≥ 5% of genomes. Red dots on neighbor-joining trees signify complete or chromosomal genomes used for initial identification of BGCs. Heatmap cell color indicates the BGC class as predicted by antiSMASH. **c)** Results from benchmarking *Isa*BGC-AutoExpansion sensitivity on 14 completed genomes for the species of *M. luteus*. AntiSMASH and *Isa*BGC-AutoExpansion were run on artificially fragmented genomes and resulting BGC predictions were compared to initial BGCs identified on the unfragmented, complete genomes. **d)** The relationship between the assembly quality measured by N50 and the total number of homolog groups detected across 14,978 *Staphylococcus* draft genomes is shown. The coloring corresponds to the percentage of BGC homolog groups co-located in a segment with a homolog group commonly found in protocore regions of BGC predictions by antiSMASH run on completed genomes.

Figure 2: *Isa*BGC reveals new insight about characterized and unstudied BGCs from skin-associated species. a) A maximum-likelihood phylogeny was constructed from PznA and PznB showing the relationship between GCFs encoding for pyrazinones. Branch color corresponds to different GCFs and the outer circular color denotes the species classification for the genome from which the sequences were gathered. **b)** Homolog groups across the seven pyrazinone encoding GCFs are depicted as a network where edges indicate that two homolog groups co-occur in the same GCF. Homolog group color of nodes depict distinct GCFs. **c)** Phylogeny constructed from ribosomal proteins showing the conservation of *Corynebacterium* GCF-50 across the diverse species encoding the NRPS using results from *Isa*BGC-PopGene analysis. The gene schematics show the consensus gene order for the GCF. Gene lengths correspond to the median length across all BGC instances and only genes found in two or more genomes are shown. For each species, the gene is colored based on the median similarity of gene sequences from the species to the consensus sequence across all species. The barplot displays the total number of GCF instances identified per species across the full set of *Corynebacterium* genomes. Broad annotations for homolog groups are shown as a color strip. Below it are bar plots corresponding to the conservation of homolog groups across samples and their median Beta-RD. **d)** A histogram of the amino acid identities between NRP synthase instances is compared to the average amino acid identities (AAI) between genomes they were gathered from.

Figure 3: Staphyloxanthin is ubiquitous across *Staphylococcus* and found on plasmids. a) A maximum-likelihood phylogeny of the full *Staphylococcus* genus was constructed from ribosomal protein encoding genes. The clade in green represents the newly recognized genus of *Mammaliicoccus*. Heatmaps showcase the presence of the five *crt* genes for each species for five separate GCFs containing the *crt* operon. The shading indicates the proportion of a species found to carry homolog groups within a particular GCF context in log₁₀ scale. **b)** A maximum-likelihood phylogeny was built from CrtM and CrtN protein alignments. Branch colors represent the GCF classification of the CrtMN sequences and bootstrap values are shown for key nodes where GCFs partition. The species the CrtMN sequence was extracted from is shown as a color strip followed by a bar plot depicting the comparative codon usage cosine distance (CCU-cd). CCU-cd was only calculated for the full five gene *crt* operon and represents the codon frequency dissimilarity with the codon frequency of the background genome. The heatmap showcases the genome-wide average amino acid identity (AAI) between pairs of genomes from which CrtMN sequences were gathered. **c)** AAI and shared homolog group content were calculated between pairs of GCF instances from different genomes for GCF-3 and GCF-6 individually. The coloring represents genome-wide AAI similarity. **d)** A schematic of the completed genome for *S. equorum* C2014, which was predicted to feature GCF-6 in three distinct copies. Bottom panel shows the BGC alignment between the three instances of GCF-6 found within this genome. **e)** Representative isolates with and without staphyloxanthin encoding GCFs were grown on 20% BHI agar to visualize pigmentation production. **f)** Methanol extractions and wavelength absorption analysis were performed to identify signature peaks in the 400 to 500 nm range associated with the presence of staphyloxanthin in *S. aureus*[39,44]. Spectra are normalized to cell density.

Figure 4: Mining for microdiversity within the cutimycin BGC leads to genetic insights for clade III *C. acnes*. a) Codon usage dissimilarity between BGC regions and background genomes shown for six common GCFs in *C. acnes*. **b**) The cutimycin encoding GCF is found primarily in clade I (subclade IB; orange) and clade III (tan) *C. acnes*. **c**) A schematic of the cutimycin encoding GCF. Lines in black correspond to sites of nearly-fixed differences between clades of *C. acnes*. **d**) *Isa*BGC-DiscoVary was used to identify the presence of the cutimycin GCF in 270 skin metagenomes. The proportion of alleles representative of clade IB or clade III *C. acnes* at sites of nearly fixed-differences was quantified. High-resolution strain quantification within metagenomes was similarly performed with StrainGE and used to overlay clade-specific typing results from *Isa*BGC-DiscoVary analysis. **e**) The distribution of novel SNVs identified by *Isa*BGC-DiscoVary within metagenomes, post filtering for removal of false positives, is shown across homolog groups and subjects. **f**) The location of novel SNVs is shown along the lantibiotic dehydratase N-terminal encoding gene (*HG2417*) for each subject. Coloring of the novel SNVs corresponds to which body-site(s) they were identified within. **g**) The clade distribution of reference genes on which novel SNVs were identified across the cutimycin GCF for subject S012 are shown individually for the predicted transitions or transversions. **i**) The proportion of differences which corresponded to cytosine to thymine or thymine to cytosine transitions along a core-genome alignment was calculated between pairs of *C. acnes* and categorized by clade comparisons. **j**) Comparative analysis of presumed directional deaminations from *C. namnetense*, treated as ancestral, to *C. acnes* clade IB and clade III were estimated using a multi-iteration sub-sampling based approach for sites along the core-genome alignment, whereby a greater number of suspected C to T transitions were estimated for clade III *C. acnes*. **k**) The presence of a complete uracil DNA glycosylase (*ung*), from clade I *C. acnes*, is shown across available *Cutibacterium* genomes.

Figure 5: Thousands of novel SNVs identified within BGCs from the *C. kefirresidentii* species complex. a) Assessment of novel SNVs identified by *Isa*BGC-DiscoVary, post-filtering, for each BGC from the *C. kefirresidentii* species complex for presence within metagenomic assemblies. **b**) The number of novel SNVs is shown for ranges of conservation percentile across all homolog groups and metagenomes. The coloring corresponds to the ratio of synonymous to non-synonymous novel SNVs within each conservation range. **c**) Novel SNVs are shown for the polyketide synthase involved in mycolic acid biosynthesis. The top panel showcases the non-synonymous (up-facing) and synonymous (down-facing) novel SNVs found along the T1PKS gene, with heights corresponding to the number of metagenomes with the novel SNV. The middle panel shows the conservation scores colored by percentile ranges (lightest = least conserved sites; darkest = most conserved sites). Key domains common to T1PKS are shown underneath along with the location of predicted non-synonymous mutations within the top 5% of conserved sites for the gene. Two adjacent SNVs affecting the coding of the residue at position 343 were identified and these could result in a combined effect of encoding for alanine.