Intrinsic furin-mediated cleavability of the spike S1/S2 site from SARS-CoV-2 variant B.1.529 (Omicron)

Bailey Lubinski [1,2], **Javier A. Jaimes** * [2] **and Gary R. Whittaker** *+ [2,3]

[1] Graduate Field of Biological & Biomedical Sciences, Cornell University, Ithaca NY, 14853, USA.

[2] Department of Microbiology & Immunology, College of Veterinary Medicine, Cornell University, Ithaca NY,

14853, USA.

[3] Department of Public and Ecosystem Health, College of Veterinary Medicine, Cornell University, Ithaca NY,

14853, USA.

* Corresponding authors
+ Lead corresponding
618 Tower Rd., Ithaca NY 14853, USA
jaj246@cornell.edu; grw7@cornell.edu

Abstract

The ability of SARS-CoV-2 to be primed for viral entry by the host cell protease furin has become one of the most investigated of the numerous transmission and pathogenicity features of the virus. Here, we analyzed the S1/S2 cleavage site (also called "furin cleavage site") of the spike protein of SARS-CoV-2 B.1.529 (Omicron variant) *in vitro*, to assess the role of two key mutations (spike gene, N679K and P681H) compared to the ancestral B.1 virus. We observed significantly increased intrinsic cleavability with furin compared to an original B lineage virus (Wuhan-Hu-1) and two variants, B.1.1.7 (Alpha) and B.1.617 (Delta), that subsequently had wide circulation. Increased furin-mediated cleavage was attributed to the N679K mutation, which lies outside the conventional furin binding pocket. Our findings suggest that B.1.529 (Omicron variant) has gained genetic features linked to intrinsic furin cleavability, in line with its evolution within the population as the COVID-19 pandemic has proceeded.

Introduction

The so-called furin cleavage site has become one of the most established transmission and pathogenicity factors of the SARS-CoV-2 spike protein (1), recently reviewed in (2). A structural loop containing a $_{682}$-RRAR$_{685}$ amino acid motif was found to be present in the original SARS-CoV-2 isolates (3), with this "furin cleavage site" (FCS) located between the spike receptor-binding (S1) and fusion (S2) domains (S1/S2). An "FCS" has been subsequently found in all circulating variants but has evolved during the course of the pandemic, with the upstream proline (P$_{681}$) residue being replaced by histidine (H) and arginine (R) in several variants of concern (notably B.1.1.7 – Alpha and B.1.167 – Delta, respectively). This loop/FCS is missing from all other SARS-like viruses (Sarbecoviruses, betacoronavirus lineage B), and is actively de-selected upon adaptation of SARS-CoV-2 to common cell culture systems (e.g., Vero cells); see ref (4) as an example.

In late November 2021, a distinct SARS-CoV-2 variant emerged (B.1.529) that had an unusual cluster of mutations, especially in the spike gene (5, 6). There was an unprecedented and rapidly disseminated series

of studies on this variant, which was subsequently named "Omicron" by the World Health Organization. These studies described the quite different infection properties of B.1.529 compared to all other variants that had emerged by that time. In summary, B.1.529 has generally been shown to be less fusogenic and with lowered S1/S2 cleavage than the other variants (7-14), especially in comparison to B.1.617.2 (Delta) the predominant variant at the time of B.1.529 emergence. Conversely, it has also been reported that the furin motif in the recombinant Omicron spike ectodomain is more efficient than that in the prototypic spike ectodomain (15), and many details of the cleavage-activation process of SARS-CoV-2 B.1.529 (Omicron) remain to be resolved.

The furin cleavage site for proteins was initially described as a four amino acid pattern $_{P4}$R-X-[K/R]-R$_{P1}$ (16), with the cleavage occurring between the P1 and P1' residues, as defined for protease substrates (17). This motif had become established as a prototype FCS, although the minimal residues for furin cleavage are generally defined as $_{P4}$R-X-X-R$_{P1}$ (18). Notably, the sequence context of the four amino acid "FCS" readily modulates cleavability, and subsequent analysis has characterized the furin cleavage motif as a 20 amino acid sequence running from position P14 to P6'—having a "core" region (P6-P2'), along with two flanking solvent accessible regions, P14-P7 and P3'-P6' (19, 20).

The emergence of Omicron was reminiscent of that of the Alpha variant approximately one year earlier; both were notable for a suddenly increased accumulation of mutations, and both have been linked to long-term infection of individuals with persistent SARS-CoV-2 infection (21). Both Alpha and Omicron contain the same P681H mutation in the FCS (P5 residue), but Omicron contains an additional mutation N679K, which is the P7 residue for furin-mediated cleavage. While within the 20 amino acid sequence comprising a furin cleavage motif, P7 is outside of the core region comprising the established furin binding pocket (19). N679K introduces an additional basic residue into the FCS, which may result in a more "polybasic" sequence and increase furin-mediated cleavage.

Here, we analyze the intrinsic furin-mediated cleavage of the B.1.529 (Omicron) spike "FCS" in comparison to other SARS-CoV-2 variants, and discuss the impact of our observed increased cleavage on the COVID-19 pandemic and virus evolution.

Methods

Furin prediction calculations

Prop: CoV sequences were analyzed using the ProP 1.0 Server hosted at: https://services.healthtech.dtu.dk/service.php?ProP-1.0. PiTou: CoV sequences were analyzed using the PiTou V3 software hosted at: http://www.nuolan.net/reference.html.

Fluorogenic peptide assays

Fluorogenic peptide cleavage assays were performed as described previously (22), with minor modifications as described below. Reactions were carried out in a 96 well plate with a 100 µl volume consisting of cleavage buffer, protease, and fluorogenic peptide. Peptides were labeled with an MCA/DNP fluorescence resonance energy transfer (FRET) pair. Peptide sequences are listed in Table 1. Trypsin catalyzed reactions acted as a positive control, where 0.8 nM/well TPCK trypsin was diluted in PBS buffer. For furin catalyzed reactions, 1 U/well recombinant furin was diluted in a buffer consisting of 20 mM HEPES, 0.2 mM CaCl2, and 0.2 mM β-mercaptoethanol, at pH 7.5. A negative control consisted of peptide suspended in buffer for furin cleavage with no protease added (not shown). Fluorescence emission was measured once per minute for 60 continuous minutes using a SpectraMax fluorometer (Molecular Devices) at 30°C with an excitation wavelength of 330 nm and an emission wavelength of 390 nm. V_{max} was calculated by fitting the linear rise in fluorescence to the equation of the line.

Table 1.

Peptide Name	Peptide Sequence
Wuhan Hu 1	MCA-TQTNSPRRARSVA-Lys(DNP)
Alpha	MCA-TQTNSHRRARSVA-Lys(DNP)
Delta	MCA-TQTNSRRARSVA-Lys(DNP)
Omicron	MCA-TQTKSHRRARSVA-Lys(DNP)
N679K P681R	MCA-TQTKSRRRARSVA-Lys(DNP)
N679K P681	MCA-TQTKSPRRARSVA-Lys(DNP)

Results

As an initial bioinformatic approach to assess biochemical function, we utilized the PiTou (19) and ProP (23) protein cleavage prediction tools, comparing the spike proteins from B.1.529 to B.1.1.7, B.1.617 and the prototype SARS-CoV-2 sequence from the A.1 and B.1 lineages (e.g., Wuhan-Hu-1), as well as to MERS-CoV and selected other human respiratory betacoronaviruses that have identifiable furin cleavage sites (i.e., HCoV-HKU1 and HCoV-OC43). PiTou is expected to be a more accurate prediction tool as it utilizes biological knowledge-based cumulative probability score functions, combined with a hidden Markov model, and is reported to have a sensitivity of 96.9% and a specificity of 97.3% (19). ProP is a widely-used, but simpler, algorithm based on a pure machine learning neural network, with a reported sensitivity of 94.7% and a specificity of only 83.7 % (19)

For SARS-CoV-2 spike, only the ProP algorithm predicted an increase in the furin cleavage for the B.1.529 lineage, compared to other lineages (Figure 1). Pitou calculated B.1.529 as having the same score as B.1.1.7. As expected, both algorithms predicted MERS-CoV to have a relatively low furin cleavage score, with HCoV-HKU1 and HCoV-OC43 showing relatively high furin cleavage scores using both approaches.

To directly examine the activity of furin on the SARS-CoV-2 A.23.1 S1/S2 site, we used a biochemical peptide cleavage assay to directly measure furin cleavage activity *in vitro* (24). The specific peptide sequences used here are described in Table 1.

Virus	S1/S2 Sequence	Furin Score PiTou	Furin Score ProP
SARS-CoV-2 (A.1/B.1)	672-ASYQTQTNSP**RRAR**|SVASQS-691	+9.196	0.626
SARS-CoV-2 (B.1.1.7 – Alpha)	669-ASYQTQTNS**HRRA**R|SVASQS-688	+9.907	0.706
SARS-CoV-2 (B.1.617.2 – Delta)	669-ASYQTQTNS**RRRA**R|SVASQS-688	+12.209	0.704
SARS-CoV-2 (B.1.1.529 – Omicron)	**669-ASYQTQTKSHRRAR|SVASQS-688**	**+9.907**	**0.725**
MERS-CoV	738-LPDTPSTLTP**R**SV**R**|SVPGEM-757	+5.155	0.556
HCoV-HKU1	747-YNSPSSSSS**RRKRR**|SISASY-766	+14.634	0.868
HCoV-OC43 (clinical)	750-GYCVDYFKN**RR**S**RR**|AITTGY-769	+10.10	0.736

Figure 1: Predicted furin cleavage scores of the S1/S2 region of selected coronavirus spike proteins.

We found a noticeable increase in cleavability for Omicron compared to B.1 and all other variants tested, including Delta. With the peptides tested cleavage of Alpha (P681H) was actually decreased compared to B.1, indicating that the P7 K residue has a major impact on furin-mediated cleavage, despite it being outside the proposed "core" furin binding pocket. Trypsin was used as a control (see Appendix).

Figure 2: Fluorogenic cleavage assays of synthetic peptides labeled with an MCA/DNP FRET pair. Peptides were designed and synthesized to reproduce the S1/S2 sites of the spike protein of selected SARS-CoV-2 variants. Peptides were incubated with furin, fluorescence was monitored for 1 h and the V_{max} calculated. Error bars denote the standard error of the mean (SEM) (n=9). Two-tailed unpaired T tests were performed comparing Wuhan-Hu-1 to each variant, with significance marked by asterisks. (****, p < 0.0001)

To explore this increased cleavage in more detail, we tested peptides with a P7 lysine residue (K) on the background of either B.1 and Delta FCS sequences (i.e., containing P681 and P681R). In both cases, the P7 K had a notable impact on cleavability, as with Omicron. The peptide with a P7 lysine residue on a B.1 background (P681) had a significantly higher cleavage rate when compared with Omicron, but how this increase in peptide cleavability translates to cleavability in the context of a full spike remains to be seen.

Figure 3: Fluorogenic cleavage assays of synthetic peptides labeled with an MCA/DNP FRET pair. Peptides were designed and synthesized to reproduce the S1/S2 site of the SARS-CoV-2 spike protein, examining the effect of introducing a lysine (K) residue at P7 (position 679) along with different established amino acids present at P5 (position 681). Peptides were incubated with furin, fluorescence was monitored for 1 h and the V_{max} calculated. Error bars denote the standard error of the mean (SEM) (n=9). Two-tailed unpaired T tests were performed comparing Omicron to each peptide, with significance marked by asterisks. (NS, p = 0.0308; **, p = 0.0085; ****, p < 0.0001)

Discussion

Our data show that SARS-CoV-2 B.1.529 (Omicron) has intrinsically high ability to be cleaved by furin at its spike S1/S2 position, despite early studies indicating the contrary, and despite the lowered score predicted by the PiTou FCS cleavage algorithm. In our study, *in vitro* peptide cleavage data are in line with the ProP FCS prediction tool, and with the data on recombinant S from Ye *et al.* (15). Our data further suggest that N679K is a key mutation impacting SARS-CoV-2 spike protein function, especially when combined with additional changes at position 681 (R/H). While our data suggest that these changes are linked to furin processivity, other investigators have recently attributed altered functionality to other proteases, notably cathepsin G (25).

In addition to N679K/P681H is also notable B.1.529 (Omicron) has addition spike mutations such as H655Y and S375F that have been shown to allow altered spike cleavage-activation (26-28), with sub-variant BA.1 (but not BA.2) having ΔH69/V70 deletion in the N-terminal domain. All of these mutations, along with D614G (29), are likely to have an impact on cleavage efficiency and fusogencity for the intact spike protein. Our view is that SARS-CoV-2 B.1.529 (Omicron) has increased intrinsic S1/S2 spike cleavability, which is balanced by epistatic effects of other mutations to drive viral infection and cell tropism in a highly context-dependent manner.

It is noteworthy that mutations upstream of the S1/S2 cleavage motif, including Q675H/R, Q677 H/P as well as N679K have also occurred independently in many SARS-CoV-2 variants (30) and may confer more of a direct impact on furin processing, as they occur in the P9, P11 and P13 positions within the 20 amino acid furin cleavage motif. Despite overall mixed conclusions on Omicron spike cleavage, fusion and infection, our biochemical data support the hypothesis that SARS-CoV-2 is evolving toward increased inherent furin cleavability, with reside 679 (N-K) being a major site of selection.

Acknowledgements

This work was funded in part by the National Institute of Health research grant R01AI35270 (to GW). We thank the global SARS-CoV-2 sequencing groups for their open and rapid sharing of sequence data and GISAID for providing an effective platform to make these data available, and all members of the Whittaker lab for helpful discussions.

Appendix

As a control, we also tested the ability of trypsin to cleave selected S1/S2 peptides. All variants were cleaved more effectively than for the prototype B.1 virus (Wuhan Hu 1), with B.1.529 (Omicron) showing the most cleavability.

References

1. Whittaker GR. 2021. SARS-CoV-2 spike and its adaptable furin cleavage site. Lancet Microbe 2:e488-e489.PMC8346238

2. Chan YA, Zhan SH. 2021. The emergence of the spike furin cleavage site in SARS-CoV-2. Molecular Biology and Evolution doi:10.1093/molbev/msab327

3. Jaimes JA, Andre NM, Chappie JS, Millet JK, Whittaker GR. 2020. Phylogenetic Analysis and Structural Modeling of SARS-CoV-2 Spike Protein Reveals an Evolutionary Distinct and Proteolytically Sensitive Activation Loop. J Mol Biol 432:3309-3325.PMC7166309

4. Johnson BA, Xie X, Bailey AL, Kalveram B, Lokugamage KG, Muruato A, Zou J, Zhang X, Juelich T, Smith JK, Zhang L, Bopp N, Schindewolf C, Vu M, Vanderheiden A, Winkler ES, Swetnam D, Plante JA, Aguilar P, Plante KS, Popov V, Lee B, Weaver SC, Suthar MS, Routh AL, Ren P, Ku Z, An Z, Debbink K, Diamond MS, Shi P-Y, Freiberg AN, Menachery VD. 2021. Loss of furin cleavage site attenuates SARS-CoV-2 pathogenesis. Nature 591:293-299

5. Martin DP, Lytras S, Lucaci AG, Maier W, Grüning B, Shank SD, Weaver S, MacLean OA, Orton RJ, Lemey P, Boni MF, Tegally H, Harkins GW, Scheepers C, Bhiman JN, Everatt J, Amoako DG, San JE, Giandhari J, Sigal A, NGS-SA, Williamson C, Hsiao N-y, von Gottberg A, De Klerk A, Shafer RW, Robertson DL, Wilkinson RJ, Sewell BT, Lessells R, Nekrutenko A, Greaney AJ, Starr TN, Bloom JD, Murrell B, Wilkinson E, Gupta RK, de Oliveira T, Kosakovsky Pond SL. 2022. Selection analysis identifies clusters of unusual mutational changes in Omicron lineage BA.1 that likely impact Spike function. Molecular Biology and Evolution doi:10.1093/molbev/msac061

6. Jung C, Kmiec D, Koepke L, Zech F, Jacob T, Sparrer KMJ, Kirchhoff F. 2022. Omicron: What Makes the Latest SARS-CoV-2 Variant of Concern So Concerning? Journal of Virology 96:e02077-21

7. Du X, Tang H, Gao L, Wu Z, Meng F, Yan R, Qiao S, An J, Wang C, Qin FX-F. 2022. Omicron adopts a different strategy from Delta and other variants to adapt to host. Signal Transduction and Targeted Therapy 7:45

8. Zhao H, Lu L, Peng Z, Chen L-L, Meng X, Zhang C, Ip JD, Chan W-M, Chu AW-H, Chan K-H, Jin D-Y, Chen H, Yuen K-Y, To KK-W. 2022. SARS-CoV-2 Omicron variant shows less efficient replication and fusion activity when compared with Delta variant in TMPRSS2-expressed cells. Emerging Microbes & Infections 11:277-283

9. Xia S, Chan JF-W, Wang L, Jiao F, Chik KK-H, Chu H, Lan Q, Xu W, Wang Q, Wang C, Yuen K-Y, Lu L, Jiang S. 2022. Peptide-based pan-CoV fusion inhibitors maintain high potency against SARS-CoV-2 Omicron variant. Cell Research 32:404-406

10. Meng B, Abdullahi A, Ferreira IATM, Goonawardane N, Saito A, Kimura I, Yamasoba D, Gerber PP, Fatihi S, Rathore S, Zepeda SK, Papa G, Kemp SA, Ikeda T, Toyoda M, Tan TS, Kuramochi J, Mitsunaga S, Ueno T, Shirakawa K, Takaori-Kondo A, Brevini T, Mallery DL, Charles OJ, Baker S, Dougan G, Hess C, Kingston N, Lehner PJ, Lyons PA, Matheson NJ, Ouwehand WH, Saunders C, Summers C, Thaventhiran JED, Toshner M, Weekes MP, Maxwell P, Shaw A, Bucke A, Calder J, Canna L, Domingo J, Elmer A, Fuller S, Harris J, Hewitt S, Kennet J, Jose S, Kourampa J, et al. 2022. Altered TMPRSS2 usage by SARS-CoV-2 Omicron impacts infectivity and fusogenicity. Nature 603:706-714

11. Peacock TP, Brown JC, Zhou J, Thakur N, Newman J, Kugathasan R, Sukhova K, Kaforou M, Bailey D, Barclay WS. 2022. The SARS-CoV-2 variant, Omicron, shows rapid replication in human

primary nasal epithelial cultures and efficiently uses the endosomal route of entry. bioRxiv doi:10.1101/2021.12.31.474653:2021.12.31.474653

12. Hui KPY, Ho JCW, Cheung M-c, Ng K-c, Ching RHH, Lai K-l, Kam TT, Gu H, Sit K-Y, Hsin MKY, Au TWK, Poon LLM, Peiris M, Nicholls JM, Chan MCW. 2022. SARS-CoV-2 Omicron variant replication in human bronchus and lung ex vivo. Nature 603:715-720

13. Willett BJ, Grove J, MacLean OA, Wilkie C, Logan N, Lorenzo GD, Furnon W, Scott S, Manali M, Szemiel A, Ashraf S, Vink E, Harvey W, Davis C, Orton R, Hughes J, Holland P, Silva V, Pascall D, Puxty K, da Silva Filipe A, Yebra G, Shaaban S, Holden MTG, Pinto RM, Gunson R, Templeton K, Murcia P, Patel AH, investigators TC-DVCS, Consortium TC-GU, Consortium TGP-UNV, investigators TEoVADC-V, Haughney J, Robertson DL, Palmarini M, Ray S, Thomson EC. 2022. The hyper-transmissible SARS-CoV-2 Omicron variant exhibits significant antigenic change, vaccine escape and a switch in cell entry mechanism. medRxiv doi:10.1101/2022.01.03.21268111:2022.01.03.21268111

14. Wang Q, Anang S, Iketani S, Guo Y, Liu L, Ho DD, Sodroski JG. 2021. Functional properties of the spike glycoprotein of the emerging SARS-CoV-2 variant B.1.1.529. bioRxiv doi:10.1101/2021.12.27.474288:2021.12.27.474288

15. Ye G, Liu B, Li F. 2022. Cryo-EM structure of a SARS-CoV-2 omicron spike protein ectodomain. Nature Communications 13:1214

16. NAKAYAMA K. 1997. Furin: a mammalian subtilisin/Kex2p-like endoprotease involved in processing of a wide variety of precursor proteins. Biochemical Journal 327:625-635

17. Polgár L. 1989. General Aspects of Proteases, p 43-86, Mechanisms of Protease Action. CRC press, Boca Raton, FL.

18. Seidah NG, Prat A. 2012. The biology and therapeutic targeting of the proprotein convertases. Nat Rev Drug Discov 11:367-83

19. Tian S, Huajun W, Wu J. 2012. Computational prediction of furin cleavage sites by a hybrid method and understanding mechanism underlying diseases. Sci Rep 2:261.PMC3281273

20. Tian S. 2009. A 20 Residues Motif Delineates the Furin Cleavage Site and its Physical Properties May Influence Viral Fusion. Biochemistry Insights 2:BCI.S2049

21. Lemieux JE, Luban J. 2021. Consulting the Oracle of SARS-CoV-2 Infection. The Journal of Infectious Diseases 225:1115-1117

22. Jaimes JA, Millet JK, Goldstein ME, Whittaker GR, Straus MR. 2019. A Fluorogenic Peptide Cleavage Assay to Screen for Proteolytic Activity: Applications for coronavirus spike protein activation. JoVE doi:doi:10.3791/58892:e58892

23. Duckert P, Brunak S, Blom N. 2004. Prediction of proprotein convertase cleavage sites. Protein Eng Des Sel 17:107-12

24. Jaimes JA, Millet JK, Whittaker GR. 2020. Proteolytic Cleavage of the SARS-CoV-2 Spike Protein and the Role of the Novel S1/S2 Site. iScience 23:101212 https://doi.org/10.1016/j.isci.2020.101212.

25. Mustafa Z, Kalbacher H, Burster T. 2022. Occurrence of a novel cleavage site for cathepsin G adjacent to the polybasic sequence within the proteolytically sensitive activation loop of the SARS-CoV-2 Omicron variant: The amino acid substitution N679K and P681H of the spike protein. PLOS ONE 17:e0264723

26. Kimura I, Yamasoba D, Nasser H, Zahradnik J, Kosugi Y, Wu J, Nagata K, Uriu K, Tanaka YL, Ito J, Shimizu R, Tan TS, Butlertanaka EP, Asakura H, Sadamasu K, Yoshimura K, Ueno T, Takaori-Kondo A, Schreiber G, Consortium TGtPJ, Toyoda M, Shirakawa K, Irie T, Saito A, Nakagawa S, Ikeda T, Sato K. 2022. SARS-CoV-2 spike S375F mutation characterizes the Omicron BA.1 variant. bioRxiv doi:10.1101/2022.04.03.486864:2022.04.03.486864

27. Yamamoto M, Tomita K, Hirayama Y, Inoue J-i, Kawaguchi Y, Gohda J. 2022. SARS-CoV-2 Omicron spike H655Y mutation is responsible for enhancement of the endosomal entry pathway and reduction of cell surface entry pathways. bioRxiv doi:10.1101/2022.03.21.485084:2022.03.21.485084

28. Escalera A, Gonzalez-Reiche AS, Aslam S, Mena I, Laporte M, Pearl RL, Fossati A, Rathnasinghe R, Alshammary H, van de Guchte A, Farrugia K, Qin Y, Bouhaddou M, Kehrer T, Zuliani-Alvarez L, Meekins DA, Balaraman V, McDowell C, Richt JA, Bajic G, Sordillo EM, Dejosez M, Zwaka TP, Krogan NJ, Simon V, Albrecht RA, van Bakel H, García-Sastre A, Aydillo T. 2022. Mutations in SARS-CoV-2 variants of concern link to increased spike cleavage and virus transmission. Cell Host & Microbe 30:373-387.e7

29. Yurkovetskiy L, Wang X, Pascal KE, Tomkins-Tinch C, Nyalile TP, Wang Y, Baum A, Diehl WE, Dauphin A, Carbone C, Veinotte K, Egri SB, Schaffner SF, Lemieux JE, Munro JB, Rafique A, Barve A, Sabeti PC, Kyratsous CA, Dudkina NV, Shen K, Luban J. 2020. Structural and Functional Analysis of the D614G SARS-CoV-2 Spike Protein Variant. Cell 183:739-751.e8.PMC7492024

30. Tao K, Tzou PL, Nouhin J, Gupta RK, de Oliveira T, Kosakovsky Pond SL, Fera D, Shafer RW. 2021. The biological and clinical significance of emerging SARS-CoV-2 variants. Nature Reviews Genetics 22:757-773