Candidatus Nealsonbacteria (OD1) are biomass recycling ectosymbionts of methanogenic archaea in a stable benzene-degrading enrichment culture

Xu Chen[1], Olivia Molenda[1], Christopher T. Brown[2,3†], Courtney R. A. Toth[1], Shen Guo[1], Fei Luo[1††], Jane Howe[4], Camilla L. Nesbø[1], Christine He[5], Elizabeth A. Montabana[2,6], Jamie H. D. Cate[2,6,7], Jillian F. Banfield[5,6,8,9], and Elizabeth A. Edwards[1,*]

[1]Department of Chemical Engineering and Applied Chemistry, University of Toronto, Toronto, ON, M5S 3E5, Canada.
[2]Department of Molecular and Cell Biology, University of California, Berkeley, CA, USA
[3]California Institute for Quantitative Biosciences, University of California, Berkeley, CA, USA
[4]Department of Materials Science and Engineering, University of Toronto, Toronto, ON, M5S 3E4, Canada
[5]Innovative Genomics Institute, University of California, Berkeley, CA, USA
[6]Molecular Biophysics and Integrated Bioimaging, Lawrence Berkeley National Laboratory, Berkeley, CA, USA
[7]Department of Chemistry, University of California, Berkeley, CA, USA
[8]Department of Earth and Planetary Science, University of California, Berkeley, CA, USA;
[9]Environmental Science, Policy and Management, University of California, Berkeley, CA, USA;
†Current Address: Metagenomi, Inc., 1545 Park Avenue, Emeryville, CA, USA
††Current Address: Liven Proteins Corporation, 200 College Street, Toronto, ON, M5S 3E5, Canada.

Keywords: Candidate Phyla Radiation, Nealsonbacteria, genomics, genome-resolved metagenomics, fluorescent in situ hybridization, CryoEM, Benzene, Anaerobic, Methanothrix, methanogenic benzene biodegradation, biomass recycling

*Corresponding author mailing address: University of Toronto, Toronto, Ontario, M5S 3E5, Canada. Phone: (416) 946-0996. Fax: (416) 978-8605.
E-mail: elizabeth.edwards@utoronto.ca;

1   The authors declare no competing interests.

Summary

The Candidate Phyla Radiation (CPR) is a very large group of bacteria with no pure culture

representatives, first discovered by metagenomic analyses. Within the CPR, candidate phylum

Parcubacteria (previously referred to as OD1) within the candidate superphylum Patescibacteria

is prevalent in anoxic sediments and groundwater. Previously, we had identified a specific

member of the Parcubacteria (referred to as DGGOD1a) as an important member of a

methanogenic benzene-degrading consortium. Phylogenetic analyses herein place DGGOD1a

within the *Candidate* clade Nealsonbacteria. Because of its persistence over many years, we

hypothesized that *Ca.* Nealsonbacteria DGGOD1a must serve an important role in sustaining

anaerobic benzene metabolism in the consortium. To try to identify its growth substrate, we

amended the culture with a variety of defined compounds (pyruvate, acetate, hydrogen, DNA,

phospholipid), as well as crude culture lysate and three subfractions thereof. We observed the

greatest (10 fold) increase in the absolute abundance of *Ca.* Nealsonbacteria DGGOD1a only

when the consortium was amended with crude cell lysate. These results implicate *Ca.*

Nealsonbacteria in biomass recycling. Fluorescent in situ hybridization and cryogenic

transmission electron microscope images revealed that *Ca.* Nealsonbacteria DGGOD1a cells

were attached to larger archaeal *Methanothrix* cells. This apparent epibiont lifestyle was

supported by metabolic predictions from a manually curated complete genome. This is one of

the first examples of bacterial-archaeal episymbiosis and may be a feature of other *Ca.*

Nealsonbacteria found in anoxic environments.

Introduction

The candidate phylum Parcubacteria belongs to a superphylum referred to as the

Candidate Phyla Radiation (CPR) which was first defined by culture-independent metagenomic

analysis (1). Members of the CPR are phylogenetically diverse and ubiquitous, but are especially

abundant in lakes, sediment, and groundwater (1, 2). Although their roles are still not well

characterized, clues are emerging from the rapidly growing number of metagenome sequencing

projects and genome reconstructions. To date, most members of the Parcubacteria remain

uncultured and are only known from metagenome assembled genomes (MAGs). In 2020, Chen

et al. (3) summarized 38 closed, complete MAGs or cMAGs belonging to the CPR. As of 2022,

19 cMAGs from the Parcubacteria within the CPR can be retrieved from NCBI, but none is for a

member of the Nealsonbacteria subgroup. The genomes of Parcubacteria are extremely small (~1

Mb) and include many hypothetical proteins which lack homologs in well-studied microbes (4).

The reduced genomes of Parcubacteria encode few metabolic capacities, lacking the ability to

synthesize lipids and often missing complete pathways for the biosynthesis of several vitamins,

amino acids, and nucleotides (5, 6). Coupled with a lack of an electron transport system, the

small cells typical of Parcubacteria and other members of the CPR are thought to obtain nutrients

and energy through close contact with an obligate host or partner cell (7). Several recent

publications support this hypothesis. For example, Huddy et al. (8) inferred an association

between CPR member Saccharibacteria (TM7) and Actinobacteria based on abundance patterns.

He at el. (9) reveled episymbiotic interaction between ultrasmall cells and hosts in groundwater

by Cryo-TEM. Scanning transmission x-ray microscopy (STXM) data obtained by Alvarado et

al. (10) showed an association of ultra-small cells with filamentous bacteria which encapsulated

with elemental sulfur spherical granules. The most definitive study was a study of

Saccharibacteria (TM7) and its bacterial host, *Actinomyces odontolyticus* XH001, in cultivated

co-cultures (11). Saccharibacteria genomes share similarities with the Parcubacteria in that they

are small and lack complete biosynthetic pathways. Metagenomic analyses suggest that

Parcubacteria may also be closely associated with a host microbe, perhaps as an episymbiont (5,

9, 12, 13) but experimental evidence is scant because of the difficulty in cultivating and

enriching these tiny microorganisms from complex ecosystems.

Interestingly, a stable methanogenic benzene-degrading enrichment culture maintained in

our laboratory, referred to as the OR consortium (for Oil Refinery), provides an opportunity to

study a member of the Parcubacteria in vivo. The OR consortium was originally derived from

oil-contaminated sediments in 1995 and has since been maintained in anaerobic minimal

medium and provided with benzene as the only added electron donor and carbon source (14).

Multiple subcultures have been established over the years, including a scaled-up lineage

currently being used for bioaugmentation field trials (15-17). The dominant microbes in these

subcultures include specialized benzene-fermenting *Desulfobacterota* (ORM2), several

methanogenic archaea, and - most interestingly for this study - a member of the Parcubacteria

originally referred to as OD1 but now further classified as *Candidatus* Nealsonbacteria (Fig. S1)

in this study. The consortium repeatedly degrades benzene to methane and CO_2 at rates from 0.2

to 10 mg benzene/L/day, however the biochemical mechanism for anaerobic benzene

biodegradation is still poorly understood. Toth et al. (17) recently described the current state of

knowledge of the OR consortium, highlighting the coordinated metabolic interactions of ORM2

and acetoclastic and hydrogenotrophic methanogens for complete mineralization of benzene. *Ca.*

Nealsonbacteria might also play role in the OR consortium because they have consistently been

detected in every culture survey (18) although a clear correlation between the abundance of *Ca.*

Nealsonbacteria and benzene degradation rate has not been found. OR consortium cultures are

maintained in fed-batch mode with infrequent (every 1-2 months) shallow transfers into fresh

medium resulting in long cell residence times. Despite years of maintenance solely on benzene

the total microbial concentration remains below 10^9 cells/mL (19). Accumulated dead cells are a

likely an alternative carbon and energy source for microbial growth. More generally, *Ca.*

Nealsonbacteria are also found in anoxic hydrocarbon- and organohalide-contaminated sites (20,

21), high pH spring water (22), and other nutrient-limited groundwater environments (9), where

biomass turnover rates are sometimes high, perhaps in response to stress, therefore we

hypothesized that these microbes may play a role in biomass recycling.

We sought to identify the substrates of *Ca.* Nealsonbacteria and any possible host associations

within the benzene-degrading OR consortium using enrichment experiments on various

substrates, quantitative PCR (qPCR), sequencing and genome reconstruction, and various types

of microscopy. The most growth of *Ca.* Nealsonbacteria was seen in cultures amended with

crude cell lysate (prepared from the consortium itself or from *Escherichia coli*). Fluorescence *in*

situ hybridization (FISH) and electron microscopy (EM) revealed *Ca.* Nealsonbacteria as an

epibiont on the surface of *Methanothrix*, a methanogenic archaeon essential for completing the

biotransformation of benzene to methane. Hybrid assemblies of short read (Illumina) and long

read (PacBio) metagenomic sequences were used to successfully reconstruct the complete closed

genome of the uncultured Parcubacteria in OR consortium, designated DGGOD1a. This 1.16

Mbp genome displays biosynthetic deficiencies and other episymbiotic traits as observed in most

other Parcubacteria genomes. A maximum likelihood concatenated ribosomal protein tree

placed the DGGOD1a genome within the candidate group Nealsonbacteria (Figure S1) as

defined in Anantharaman et al. (23). These observations begin to shed light on uncharacterized

members of the CPR and bring into focus the phenomenon of bacterial episymbiosis on archaeal

hosts.

Results and Discussion

qPCR reveals abundance of *Ca.* Nealsonbacteria in methanogenic cultures

A qPCR survey of 15 different methanogenic benzene-degrading subculture bottles was

conducted to measure the abundance of benzene-fermenting *Desulfobacterota* ORM2 and *Ca.*

Nealsonbacteria (referred to as OD1 in previous studies), as well as the abundance of total

bacteria and total archaea. One subculture bottle named "OR-1bBig" was selected for use in two

substrate "Donor Trial" experiments because it contained a relatively high abundance of *Ca.*

Nealsonbacteria (~5% based on 16S rRNA amplicon sequencing results) and sufficient culture

volume was available (1.5 L). Other than benzene, the products of cell decay could feasibly

serve as electron donors in these bottles. This led us to hypothesize that components of lysed

cells might serve as growth substrates for *Ca.* Nealsonbacteria.

Ca. Nealsonbacteria is most enriched in cultures amended with lysed cells.

In Donor Trial #1, we tested three common substrates (pyruvate, acetate, and hydrogen), as well

as French-Press lysed benzene culture, and two specific components of cells (salmon sperm DNA,

and the phospholipid L-α-phosphatidylethanolamine) to see if any of these putative electron donors

could support the growth of *Ca*. Nealsonbacteria, directly or indirectly (Table S1). We found that

Ca. Nealsonbacteria concentrations increased in vials amended with lysed culture (Figure 1a). In

these vials, *Ca.* Nealsonbacteria relative abundance increased from 5% to 30% and absolute

abundance by an order of magnitude to 5.1×10^6 copies (cells)/mL over 4 months (Figure 1; data

in Table S2a). None of the other electron donors tested resulted in discernable enrichment of *Ca.*

Nealsonbacteria relative to benzene-amended controls. In bottles amended with pyruvate, *Ca.*

Nealsonbacteria absolute abundance increased ~5 times (up to 9.8×10^5 copies/mL), but relative

abundance decreased to 1% because other microbes grew much more (Figure 1b; Table S3). *Ca.*

Nealsonbacteria growth in pyruvate and acetate amended bottles may have resulted from the direct

substrate utilization or as an indirect consequence of cell growth and corresponding increase in

dead cells. Either way, neither pyruvate nor acetate were selective for *Ca.* Nealsonbacteria. As

expected, the abundance of benzene-degrading Desulfobacterota ORM2 dropped significantly in

the absence of benzene (Figure 1b). In the vials amended with cell lysate (second bar; Figure 1b),

in addition to enrichment of *Ca.* Nealsonbacteria to 30% of the population, we also observed

enrichment of *Syntrophaceae* while the absolute abundance of methanogenic archaea was

relatively stable (Table S2a). The ASV corresponding to *Syntrophaceae* was found to share 100%

16S RNA gene sequence identity to an uncultured *Syntrophaceae* (MH665869) from a

methanogenic pyrite forming culture fed with ferrous sulfide (FeS), H_2S, and CO_2 (24). A similar

metabolism may be occurring as our medium is reduced with FeS and buffered with bicarbonate

(25).

Nonmetric dimensional scaling (NMDS) was applied to further analyze Donor Trial #1 ASV data

(Figure S2a). The unfed "starved control" culture community clustered with communities from

culture vials amended with cell lysate, consistent with products of cell decay supporting the

microbial community in both cases. Methane production data from each treatment indicated that

the added substrate equivalents (chemical oxygen demand [COD]) were consumed by 4 months

(Table S4). Therefore, data collected at 7 months were from cultures that were starved for more

than 3 months and marked with "+" in the NMDS analysis. Interestingly, we found that after 7

months, the communities did not converge. We expected to see enrichment of *Ca.* Nealsonbacteria

in every vial after prolonged starvation. However, such enrichment was only clearly observed in

acetate and pyruvate-amended vials (Figure 2). Perhaps there was insufficient biomass growth,

therefore insufficient dead biomass in the other vials to detect enrichment of *Ca.* Nealsonbacteria.

Indeed, total cell concentrations were highest in acetate and sodium pyruvate amended vials.

Interestingly, a second ASV belonging to *Ca.* Nealsonbacteria (designated ASVb) was enriched

in the acetate- and pyruvate-amended cultures (Figure 2) and became more abundant than *Ca.*

Nealsonbacteria ASVa – the strain that is typically found in the benzene-degrading consortium

(18). These two *Ca.* Nealsonbacteria ASVs share only 91.2% 16S rRNA sequence identity. This

intriguing result suggests that different species may exhibit distinct preferences for the type of

lysed cell material they can metabolize, because the acetate- and pyruvate-amended cultures

selected for very different communities.

Building on the first trial, a second trial (Donor Trial #2) compared the growth of *Ca.*

Nealsonbacteria on lysates from the benzene culture as well as *E. coli* lysate. Further, we tested

three sub-fractions of the benzene culture lysate, which were separated by centrifugation and

tangential flow filtration with a 50kDa cut off as depicted in Figure S3. The pyruvate treatment

from Donor Trial #1 was also repeated but this time with added antibiotics (1 mM each kanamycin

and vancomycin) to try to increase selectivity for *Ca.* Nealsonbacteria. The results of Donor Trial

#2 (raw data in Table S2b) were similar to those of Donor Trial #1, with whole cell lysate resulting

in significant (>10×) growth (Figure S4) regardless of source (*E. coli* or benzene culture). At first

glance, the *E. coli* crude lysate appeared to be most stimulatory, but we later realized it was

provided at 10× the concentration of the lysate from the benzene culture based on measured COD

(Table S1c). The fact that *E. coli* lysate also supported the growth of *Ca.* Nealsonbacteria ruled

out involvement of culture specific substrates (e.g., archaeal lipids). The sub-fractions of benzene

culture lysate also resulted in enrichment although we observed considerable variability between

duplicates (Figure S4). Of the three cell lysate fractions, the >50 kDa fraction (supernatant

retentate) was most effective at enriching *Ca.* Nealsonbacteria after 4 months of incubation which

may suggest that larger molecules may be important for growth. An NMDS analysis was also

conducted on amplicon sequencing data from Donor Trial #2 (Figure S2b). These results show

that the microbial communities enriched on each fraction of benzene culture lysate were slightly

different after the first two months (triangles), but that they converged after four months (squares).

Perhaps after four months, only larger, more slowly degraded biomolecules remained in all

fractions and results in similar communities.

To look more closely at the microbial composition in each treatment, a heatmap of absolute

abundance was generated showing the top 18 ASVs in Donor Trial #2 (Figure 3); raw data are

reported in Table S5. *Ca.* Nealsonbacteria ASVa was the most abundant sequence variant detected.

Only two other microbes (*Methanothrix* and *Spirochaetaceae)* were consistently enriched with *Ca.*

Nealsonbacteria in vials amended with cell lysate, regardless of source or sub-fraction tested. Vials

amended with lysed *E. coli* also enriched for ASVs belonging to *Clostridium* and *Bacteroides* that

were not enriched in other treatments (Figure 3). Comparing the culture amended with only

pyruvate in Donor Trial #1, the vials amended with antibiotics kanamycin and vancomycin

successfully inhibited the growth of most microbes including *Ca.* Nealsonbacteria and

Methanothrix. This indicates that *Ca.* Nealsonbacteria is antibiotic-sensitive or indirectly affected

by its antibiotic-sensitive metabolic partners. It is interesting to note that the culture's benzene

fermenter (ORM2) seems to have persisted in those treatments. The growth of

Methanomethylovorans and *Methanobacterium* in the presence of antibiotics confirms their

resistance and together these data point to a possible means of enriching for benzene-degrading

Desulfobacterota ORM2.

The two donor trials confirmed that compound(s) present in bacterial cell lysate can enrich *Ca.*

Nealsonbacteria. The exact identity of the compound(s) remains elusive, but the data point to

higher molecular weight and/or more recalcitrant proteins or extracellular polymeric substances,

as these would present in all of the fractions tested. Two very distinct strains of *Ca.*

Nealsonbacteria grew in the culture depending on treatment revealing subtle ecological

preferences yet to be discovered.

Fluorescence in situ microscopy (FISH) reveals *Ca.* Nealsonbacteria DGGOD1a as an

epibiont of *Methanothrix*.

To visualize the enriched *Ca.* Nealsonbacteria in the methanogenic benzene degrading culture by

microscopy, we attempted to develop specific mono-fluorescent *in situ* hybridization (FISH)

probes targeting various single stranded regions in the 16S rRNA gene. However, none of the

probes worked, possibly because of low ribosome counts in these cells (26) or non-ideal targeting

in the secondary structure. Very recently, Kuroda et al. (27) successfully designed and validated a

mono-FISH probe for a *Ca.* Nealsonbacteria with a nearly identical 16S RNA gene sequence by

targeting a different location on the gene; clearly this new FISH probe will be tested in the future.

As an alternative to a *Ca.* Nealsonbacteria-specific FISH probe, we used a combination of a

universal Archaea FISH probe (ARCH915, Table S6) and DAPI staining to distinguish

methanogens from bacteria. Given that *Ca.* Nealsonbacteria DGGOD1a was the most abundant

bacterial phylotype in several of our lysate-amended enrichment cultures (Figure 3), we compared

microscopic cell counts to qPCR abundance estimates of the same sample to confirm that the ultra-

small cells were *Ca.* Nealsonbacteria. We focused primarily on one specific cell lysate enrichment

culture (Lysate replicate #2 at 2 months) because of its high *Ca.* Nealsonbacteria abundance ($>10^6$

cells/mL in this culture compared to all other bacterial phylotypes (Figure 3). In this *culture*, we

observed many DAPI-stained ultra-small (~0.2 µm) cells attached to a single filamentous structure

that stained with the archaeal FISH probe (Figure 4a). The ultra-small cell size matches the typical

size of CPR members (5, 9) and is presumed to be *Ca.* Nealsonbacteria. The shape of the

filamentous archaea is distinctive of *methanothrix* cells. We counted the number of small cells

attached to long archaeal filaments as well as the number of filamentous archaea. We also counted

the number of free-living microbes with similar size and shape to that of the attached small cells.

Twenty-one fields of view were imaged and counted for each of two independent samples from a

Ca. Nealsonbacteria enrichment culture. These small cells were by far the most numerous bacteria

in all views, and thus must correspond to *Ca.* Nealsonbacteria, as it was most numerous via qPCR

in this culture. The count of small *Ca.* Nealsonbacteria cells was $3.2\text{-}6.6\times10^6$ cells /ml with

attached cells ~2.5 times more abundant than free-living cells. (Table S7). Gene copies by qPCR

in samples of DNA extracted from the same culture using primers specific for *Ca.* Nealsonbacteria

gave a comparable result of 8.3×10^6 copies/ml. Thus, we are confident that the small cells attached

to the archaeal filaments are *Ca.* Nealsonbacteria. *Methanothrix* cell counts were also estimated at

$1.9\text{-}3.4 \times 10^6$ cells/mL, which also matches reasonably well to the value of 0.9×10^6 cells/mL from

qPCR results (Table S7). Interestingly, a similar association of *Ca.* Patescibacteria and

Methanothrix was very recently reported in an artificial mixture of a culture of *Methanothrix*

soehngenii GP6 with wastewater biosludge (27).

We had previously unsuccessfully attempted to enrich *Ca.* Nealsonbacteria by filtration through a

0.22-μm filter. Analysis of the filtrate by qPCR showed no enrichment: both bacterial and the *Ca.*

Nealsonbacteria concentrations decreased 100-fold (data not shown). Other studies have reported

success enriching ultra-small microbes this way (1, 26, 28). Perhaps the strong attachment of *Ca.*

Nealsonbacteria to its host caused this approach to fail. Interestingly, we also noticed that the

archaeal FISH probe seemed to bind more strongly to methanogens that had many *Ca.*

Nealsonbacteria cells attached, compared to naked methanogens (Figure 4a, bottom right; Figure

S5). Naked methanogens stained brightly with DAPI (DNA) but only faintly with the archaeal

FISH probe (rRNA). Given that FISH signal intensity is a measure of ribosome count and thus

metabolic activity (29), perhaps methanogens with *Ca.* Nealsonbacteria attached are more active.

This observation is in contrast to the recent study by Kuroda et al., (27) where a major fraction of

the Ca. Nealsonbacteria were attached to methanogens with weaker fluorescence. *Ca.*

Nealsonbacteria's role as either a parasitic or commensal organism is thus still unclear.

SEM and Cryo-EM confirm attachment to *Methanothrix*.

SEM examination of the parent methanogenic benzene-degrading culture prepared with ionic

liquid exchange to preserve cell structure also showed a similar association of very small *Ca.*

Nealsonbacteria-like cells attached to a large filamentous cell structure consisting of a chain of

rod-shaped, blunt-ended cells (Figure 4b). This filament was enclosed in a tubular sheath and the

flat ends distinctively match the characteristics of *Methanothrix* (30). The *Ca.* Nealsonbacteria-

like cells are spherical with a diameter of 0.2 μm to 0.8 μm and they were always found in budding

pairs, forming a peanut shape. Cryo-EM was also used to examine the *Ca.* Nealsonbacteria

enrichments. The Cryo-EM images (Figure 4c) show the same association of peanut-shaped cells

with *Methanothrix*. The large black rod-shaped blurry cells at the center of the image are

Methanothrix forming a sheathed chain. Multiple peanut-shaped ultra-small cells were observed

attached to the chain of *Methanothrix* (Figure 4c). The *Ca.* Nealsonbacteria-like cells were always

found in pairs and the one attached to the methanogen was usually bigger. A zoomed-in image

(Figure 4c; right side) reveals that *Ca.* Nealsonbacteria cells have a very interesting membrane

structure with a thick sheath and an apparent S-layer. Additional supporting FISH, SEM, Cryo-

EM images are provided in Figures S5 and S6. The close association of *Ca.* Nealsonbacteria and

Methanothrix may facilitate efficient nutrient exchange between these organisms. Given that

Methanothrix are obligate acetoclastic methanogens, *Ca.* Nealsonbacteria may provide acetate to

Methanothrix. The physical association also enables potential direct interspecies electron transfer

or DIET (31). Although, e-pili genes were not found in the *Ca.* Nealsonbacteria genome (see

below), DIET or direct acetate exchange could happen by other mechanisms. Clearly the possible

syntrophy between the *Ca.* Nealsonbacteria and *Methanothrix* requires further investigation.

A co-occurrence network finds *Ca.* Nealsonbacteria variants positively correlated with *Methanothrix.*

A co-occurrence network inference (CoNet) analysis was conducted using all 16S rRNA amplicon sequencing data from the two donor trials (Figure S7a). *Ca.* Nealsonbacteria ASVa was found to correlate positively with its putative host *Methanothrix* ASV89, as well as to *Ca.* Nealsonbacteria ASVb and to two less abundant microbes (*Sporomusa* and *Ignavibacteriales*) and to correlate negatively with *Spirochaetaceae* ASV977, which was only enriched in the bottles amended with pyruvate. *Ca.* Nealsonbacteria ASVb (enriched only when pyruvate or acetate were added) was positively correlated with *Methanothrix* ASV89 as well as to *Methanothrix* ASV91. ASV91 shares 99.4% pairwise identity with *Methanothrix* ASV89 and could be a specific host for *Ca.* Nealsonbacteria ASVb. A heatmap of absolute abundance was generated (Figure S7b) that includes all the ASVs from the co-occurrence network. The rows of heatmap were clustered based on similarity. Interestingly, *Ca.* Nealsonbacteria ASVa clustered with *Methanothrix* ASV89 while *Ca.* Nealsonbacteria ASVb clustered with *Methanothrix* ASV91 (Figure S7b, red rectangles). This clustering suggests that *Ca.* Nealsonbacteria ASVa and ASVb may have different *Methanothrix* hosts.

Impact of *Ca.* Nealsonbacteria on benzene degradation

To further study the role of *Ca.* Nealsonbacteria in benzene degradation, we mixed an enrichment (containing about 10^7 *Ca.* Nealsonbacteria cells/mL) with a "slow" benzene-degrading culture (containing about 10^7 ORM2 cells/mL, degrading at ~0.1 mg/L/day). As a control, we also amended the slow culture with supernatant recovered from a healthy culture (degrading benzene at 10 mg/L/day). The raw data from this experiment are reported in Table S8. We observed a

significant increase in benzene degradation rate in the two replicates that were a 50:50 mix of the

two cultures (Figure 5a), even though mixing diluted the benzene degrader (ORM2) by 50%.

Therefore, adding a *Ca.* Nealsonbacteria enrichment enhanced benzene degradation. Slow benzene

culture amended with supernatant from a healthy culture as well as the slow culture-only controls

all showed overall slower benzene degradation (Figure 5a). Benzene biodegradation rates were

highest in cultures with the highest measured ORM2 concentrations at Day 217, except for one

culture replicate (slow culture #1) with abundant ORM2 but no benzene degradation activity

(Figure 5b). The two vials amended with the *Ca.* Nealsonbacteria enrichment showed the highest

cell concentrations for both ORM2 and *Ca.* Nealsonbacteria. Perhaps *Ca.* Nealsonbacteria

supplied ORM2 or *Methanothrix* with specific limiting metabolites, nutrients or cofactors recycled

from lysed biomass or simply removed inhibitory compounds. Further studies using isotope-

labeled cell lysate may help to clarify these complex interactions.

The genome of *Ca.* Nealsonbacteria DGGOD1a is consistent with a symbiotic epibiotic

lifestyle

We assembled and closed the genome of *Ca.* Nealsonbacteria DGGOD1a (Genbank: CP092821

and IMG ID: 2791354853) corresponding to ASVa using both Illumina and PacBio sequence data.

The details of assembly are summarized in Supplemental Text S4 and reported in a genome

announcement (55). The quality of the genome assembly was validated with read-mapping and a

GC skew plot (Figure S8) clearly showing origin and terminus. The complete genome is 1.16 Mbp

in length with 1173 predicted coding sequences (CDS) including 18 pairs of perfect repeats which

are over 400 bp in length. The majority of these repeats are predicted transposases, and in fact, this

genome has a total of 39 predicted transposases which comprise 3.4% of protein coding genes.

Transposases are involved in genomic rearrangements, gene duplication and for promoting

horizontal gene transfer (32). A total of nine regions were predicted as genomic islands using

Islandviewer4 (31) of which seven contain transposases (Table S9). High numbers of transposases

are found in organisms under stress or in extreme environments (32) which may explain the

unusual high decay rate sometimes seen in our methanogenic benzene degrading culture (19).

Other studies have also shown a high transposase content in bacteria with symbiotic or pathogenic

lifestyles (33-35). The transposases in the genome of *Ca.* Nealsonbacteria may participate in DNA

exchange with the host *Methanothrix.*

Like other CPR bacteria, the genome reveals that most metabolic pathways and biosynthetic

capacities annotated by automatic pipelines are predicted to be incomplete, notably of nucleotides,

lipids, vitamins and most amino acids (Figure 6). As inferred from genomes of other members of

the CPR group (1, 5, 7), DGGOD1a can use the pentose phosphate pathway to bypass glycolysis.

Also, there is no lipid biosynthesis pathway and the pathways for the biosynthesis of most amino

acids and vitamins are incomplete. The only vitamin DGGOD1a is predicted to make is riboflavin.

The large ribosomal RNA subunit L1 is missing; a common feature of members of the OD1-L1

group (1, 5). Interestingly, eight toxin and five antitoxin gene families were found in the genome.

Some were found located within two predicted genomic islands and may contribute to their

maintenance (36). The toxin-antitoxin systems are stress-response elements that could also help

cells adapt to unfavorable growth conditions (36). These systems are common in free-living

microbes but tend to be lost from host-associated prokaryotes due to the relatively constant

environment of host-associated organisms (37). The high toxin-antitoxin genes copies in *Ca.*

Nealsonbacteria indicates that it is possibly still undergoing reductive evolution (36, 38). Twelve

type IV pilus assembly proteins, two cell membrane proteins annotated as holins, and type II

secretion system genes were found in the genome, which are consistent with an epibiont lifestyle.

Epibionts are known to use type IV pili to attach to the host membrane (39). The holin protein

family form pores in cytoplasmic membranes and may be used to permeabilize the host cell

membrane and perhaps kill the host (40). Type II secretion systems may be involved in nutrient

uptake or nutrient exchange between the *Ca.* Nealsonbacteria and its host and may also participate

in DIET (41). Surprisingly, a type V CRISPR-Cas system that contains 49 spacers was found in

the genome with Cas12a as the effector and the complex of cas4, cas1 and cas2 as the spacer

acquisition machinery. It is rare to see a CRISPR-Cas system in such a reduced genome, and only

2.4% of organisms from the Parcubacteria and Microgenomates superphyla have a CRISPR-Cas

system (42). The intact CRISPR-Cas system suggests exposure to phage and extracellular DNA.

Considering that *Ca.* Nealsonbacteria lack the ability to synthesize nucleotides, perhaps the

CRISPR-Cas system is involved in the degradation of host DNA for recycling nucleotides. No

clues were gleaned from a search of CRISPR spacer targets against the OR consortium

metagenome and all the sequences in the NCBI database, as these searches only returned hits to

the CRISPR matrix itself.

We were not able to assign function to a large proportion (30%) of the predicted proteins in the

genome, despite running multiple prediction pipelines and manual curation (Table S10). One

interesting feature is a putative NADP-reducing hydrogenase. This protein has been found in the

bacterium *Desulfovibrio fructosovorans* as an iron-sulfur protein that exclusively functions as a

hydrogen dehydrogenase (43) and may mean that Ca. Nealsonbacteria uses hydrogen produced by

other fermentation reactions. The genome also encoded two putative polysaccharide deacetylases

belonging to carbohydrate esterase family 4 (CE4). The members of this family can hydrolyze the

acetyl group from *N*-acetylglucosamine (44-47) or *O*-acetylxylose residues (48, 49) which exist in

extracellular polymeric substances (EPS). The released acetyl group could be used by the obligate

acetoclastic methanogenic archaeon *Methanothrix*. Three genes encoded putative peptidoglycan-

binding proteins were also found in the genome. The proteins belong to peptidoglycan-binding

domain could involve in cell wall synthesis (50) and binding (51) which may responsible for the

complex membrane structure we found by Cryo-EM.

Implications

Ca. Nealsonbacteria DGG1a likely contributes to recycling biomass including extracellular

polymeric substances in a methanogenic benzene degrading enrichment culture, and in so doing

enhances the growth of benzene fermenters. A role in microbial biomass recycling is supported by

substantial enrichment of this phylotype on lysed biomass. The finding of a close association of

dividing *Ca.* Nealsonbacteria cells with *Methanothrix* is now the second report of a CPR bacterium

that is an episymbiont of an archaeon (27), in this study shown in a stable laboratory enrichment

culture where further experimentation is possible. *Ca.* Nealsonbacteria were observed to be

beneficial for the growth of benzene-metabolizing *Desulfobacterota* ORM2 from yet another

uncultivated lineage. *Ca.* Nealsonbacteria may recycle key cofactors required to initiate benzene

ring activation. These results guide future experiments into specific relationships between *Ca.*

Nealsonbacteria and benzene degradation in these enrichment cultures.

The visualization of this unusual cross-kingdom episymbiosis of *Ca.* Nealsonbacteria on the

surface of *Methanothrix* implies metabolite exchange or direct electron transfer between the two

cell types. The type IV pili encoded in the genomes of many *Ca.* Nealsonbacteria (5, 7, 52) are

possibly used for attachment to the host and as well as a conduit for electron transfer. Perhaps

these tiny microbes revealed in many metagenomic surveys of natural and engineered anaerobic

communities have analogous roles? *Ca.* Nealsonbacteria may be particularly relevant where

energy input is limited and recycling is vital, such as in deep subsurface environments.

Experimental Procedures

Chemicals and microbial cultures. All chemicals were obtained from Sigma-Aldrich (Oakville,

ON, Canada) at the highest purity available unless specified otherwise. The OR consortium is

maintained in 100 mL to 2 L batch culture bottles at the University of Toronto and in larger vessels

(>100 L) at SiREM laboratories (Guelph, Ontario, Canada); the scaled-up culture is referred to as

DGG-B (17). Two OR consortium subculture bottles (or lineages) were used in the experiments

described herein: a bottle called OR-1bBig was the inoculum for *Ca.* Nealsonbacteria enrichment

Donor Trials #1 & #2 and a bottle called OR-p5 was the inoculum for the augmentation

experiment. All cultures were grown in an iron sulfide (FeS)-reduced, bicarbonate-based mineral

medium (MM medium) amended regularly (approx. 1 per month) with benzene targeting aqueous

concentration of 10 mg/L per bottle (17, 18).

Preparation of electron donor solutions for *Ca.* Nealsonbacteria enrichment trials. Two

crude cell lysate solutions from the OR consortium OR-1bBig and a culture of *Escherichia coli*

were prepared using tangential flow filtration and French press as described in Text S1. Next, we

prepared two solutions of specific biomass components, nucleic acids (salmon sperm DNA;

Sigma-Aldrich,   Deoxyribonucleic   acid   sodium   salt)   and   a   membrane   lipid   (L-α-

phosphatidylethanolamine; Sigma-Aldrich, Type II, ≥97%, lyophilized powder from egg yolk).

Acetate, pyruvate and hydrogen were also tested as donors as they are intermediates of benzene

degradation. Stock solution concentrations are listed in Table S1a.

Ca. **Nealsonbacteria Enrichment – Donor Trial #1.** To test if *Ca.* Nealsonbacteria DGGOD1a

could grow on crude lysed biomass or specific sub-cellular components, 25 mL culture tubes with

9 mL MM medium were inoculated with 1 mL OR-1bBig culture and amended with 7 different

electron donor solutions (Table 14a) at approximately the same electron equivalents, targeting ~30

mg COD/L. Each condition was tested in quadruplicate and incubated in a Coy anaerobic glove

box for 7 months (supplied with 10% H_2, 10% CO_2, and 80% N_2). DNA samples were collected

at 2, 4, and 7 months after inoculation by sacrificing an entire replicate from each treatment.

Ca. **Nealsonbacteria enrichment – Donor Trial #2.** After observing successful growth and

enrichment of *Ca.* Nealsonbacteria with crude cell lysate from the benzene culture in Donor Trial

#1, a second experiment was set up to test the impact of various biomass fractions of cell lysate.

Culture (500 mL) was concentrated by tangential flow and cells were lysed using the same French

press method described earlier (Text S1). The cell lysate was then divided into four fractions, as

seen in Figure S3. One fraction was used "as is" (BL for Benzene culture Lysate). The remaining

lysate was centrifuged (at 13,000 × g for 20 mins) to create a pellet fraction and a supernatant

fraction. The supernatant fraction was further separated by tangential flow filtration with 50kDa

cut-off membrane (Pellicon® XL cassette with Durapore® membrane) to create a permeate fraction

(< 50 kDa) and a retentate fraction (>50 kDa). Each of these fractions was tested separately (Table

S1b). Two additional treatments were included in Donor Trial #2. One consisted of pyruvate plus

the antibiotics kanamycin and vancomycin (1 mM) to inhibit bacteria without inhibiting

methanogens. A second additional treatment was amended with *E. coli* extract instead of benzene

culture extract. We ensured that the *E. coli* pellet was well washed with MM medium to remove

any residual *E. coli* rich growth medium that could confound results. Controls without substrate

(starved control; SC) and crude benzene culture lysate without inoculum (no inoculum; NI) were

included. All treatments were conducted in biological duplicates except the starved control which

was a singleton.

Ca. **Nealsonbacteria augmentation trial.** This experiment made use of an inoculum from culture

bottle "OR-p5" with a very slow benzene degradation rate over years (~0.1 mg/L/day). Eight 25-

mL culture vials were each filled with a 10 mL aliquot of this slow culture (see Text S2 for details).

Subsequently, 5 mL of OR-p5 culture was removed from vials #1-2 and replaced with 5 mL of *Ca.*

Nealsonbacteria enrichment culture BL (replicate #1) from Donor Trial #2. In vials #3-5, 5 mL of

culture was removed and replaced with 5 mL of culture supernatant from a very active benzene

culture (degradation rate of ~10 mg/L/day), prepared by centrifugation at $13,000 \times g$ for 20 min.

The remaining vials (#6-8) of slow culture were left as is. All vials were amended with neat

benzene targeting 15 mg/L aqueous concentration and reamended whenever concentrations

decreased below 5 mg/L. Benzene and methane concentrations were monitored by GC/FID (see

below). DNA samples were collected on day 217 for qPCR analysis.

DNA extraction, quantitative PCR, and Illumina amplicon sequencing and statistical

analysis. DNA from Donor Trial #1 was extracted from 10 mL culture samples; otherwise, DNA

was extracted from 1 mL culture samples. Cells were harvested by centrifugation at $13,000 \times g$ for

20 min at 4°C and pellets resuspended in residual (~0.05 mL) supernatant. DNA was extracted

using the DNeasy PowerSoil Kit (QIAGEN) following the manufacturer's protocol. Real-time

quantitative polymerase chain reaction (qPCR) assays were performed to track the gene copy

numbers of Bacteria and Archaea using universal 16S rRNA gene primers; abundances of *Ca.*

Nealsonbacteria and *Thermodesulfobacteria* ORM2 were also tracked using specific 16S rRNA

gene-targeting primers (Table S11). qPCR reactions were performed using a CFX96 real-time

thermal cycler (Bio-Rad Laboratories) using the following thermocycling conditions: an initial

denaturation step at 98 °C for 2 min, 40 cycles of 98 °C for 5 s and Tm (see Table S11) for 10 s,

followed by melt curve analysis (65-95 °C with an increase of 0.5 °C every 5 s). qPCR results

were processed with CFX Manager software (Bio-Rad Laboratories). Standard curves for all the

qPCR studies are reported in Table S12. 16S rRNA gene amplicon sequencing with primers

targeting the V6-V8 region was conducted at the Genome Quebec Innovation Centre (McGill

University) as previously described (17). Raw reads were processed into amplicon sequence

variants (ASVs) and analyzed by nonmetric dimensional scaling (NMDS) and Co-occurrence

Network Inference (CoNet) analysis (53) as described in Text S3.

Metagenome analysis. The OR consortium/DGG culture was sequenced using a combination of

Illumina paired-end (~150 bp reads, ~300 bp insert) and PacBio long-read library (~7000 bp

reads). Metagenomes were submitted to IMG under sequencing project Gp0324998. Detailed

description of assembly, binning and functional and taxonomic annotation can be found in Text

S4.

Epifluorescence microscopy and fluorescence in situ hybridization (FISH). The FISH probe

ARCH915 was used for visualizing all Archaea (54). Eight FISH probes (Table S6) targeting the

16S rRNA gene of *Ca.* Nealsonbacteria were designed and tested in this study but none yielded a

signal distinguishable from background noise. The protocol for epifluorescence microscopy is

provided in Text S5 and details into the design and testing of FISH probes in Text S6.

Scanning Electron Microscopy with Ionic Liquid and Cryo-EM. To maintain the wet native

state of microbial cells and to eliminate artifacts that may be introduced by sample drying and

coating, culture samples were prepared with an ionic liquid and examined by SEM, as detailed in

Text S7. Cryo-EM was also used to explore intracellular structure and spatial associations. Sample

preparation and Cryo-EM conditions are described in Text S8.

Analytical procedures. Benzene and methane were measured in culture bottles and vials by

injecting 0.3 mL headspace samples into a Hewlett-Packard 5890 Series II GC fitted with a GSQ

column (30 m x 0.53 mm I.D. PLOT column; J&W Scientific, Folsom, CA) as previously

described (18). External standards were used for calibration.

Accession numbers. The closed genome sequence of *Candidatus* Nealsonbacteria DGGOD1a is

available at the Joint Genome Institute (IMG) under Taxon ID 2791354853 and at NCBI under

accession no. CP092821. Amplicon sequences (16S rRNA) were uploaded to the NCBI SRA under

BioProject accession number PRJNA830497. The Genome is also available in ggkbase with the

following link: https://ggkbase.berkeley.edu/organisms/410827/contigs/251537015.

Supporting information

505   The PDF file contains: 1) supplementary Figures S1 to S6, including a phylogenomic tree of *Ca.*

506   Nealsonbacteria, fractionation steps used to preparer biomass, NDMS plots, *Ca.* Nealsonbacteria

507   growth curves and additional microscopic images; 2) supporting texts (S1 to S8) providing

508   additional method details; and 3) supplementary Tables S9, S10, S11 listing primers, probes and

509   qPCR standard curves. The excel file contains larger data Tables S1 and S5-S12.

Acknowledgements

This study was funded by Genomic Application Partnership Program (GAPP) grants awarded to

E.A.E. (Project IDs OGI-102 and OGI-173), which were supported by Genome Canada, Ontario

Genomics, the Government of Ontario, Mitacs Canada, SiREM, Alberta Innovates, Federated Co-

operatives Limited, and Imperial Oil. The authors would like to thank SiREM (Guelph, Ontario

N1G 3Z2, Canada) for supplying culture for testing. We would also like to acknowledge the

laboratories of Dr. Neil Thomson (University of Waterloo), Dr. Ania Ulrich (University of

Alberta), SiREM, and Innotech Alberta (Edmonton, Alberta T6N 1E4, Canada) for providing

feedback and intellectual contributions to this study during weekly GAPP consortium meetings.

References

1.   Brown CT, Hug LA, Thomas BC, Sharon I, Castelle CJ, Singh A, Wilkins MJ, Wrighton KC,
Williams KH, Banfield JF. 2015. Unusual biology across a group comprising more than 15% of
domain Bacteria. Nature 523:208-11.

2.   Hug LA, Baker BJ, Anantharaman K, Brown CT, Probst AJ, Castelle CJ, Butterfield CN,
Hernsdorf AW, Amano Y, Ise K, Suzuki Y, Dudek N, Relman DA, Finstad KM, Amundson R,
Thomas BC, Banfield JF. 2016. A new view of the tree of life. Nat Microbiol 1:16048.

3.   Chen L-X, Anantharaman K, Shaiber A, Eren AM, Banfield JF. 2020. Accurate and complete
genomes from metagenomes. Genome Research 30:315-333.

4.   Méheust R, Burstein D, Castelle CJ, Banfield JF. 2019. The distinction of CPR bacteria from other
bacteria based on protein family content. Nature Communications 10.

5.   Castelle CJ, Brown CT, Thomas BC, Williams KH, Banfield JF. 2017. Unusual respiratory
capacity and nitrogen metabolism in a Parcubacterium (OD1) of the Candidate Phyla Radiation. Sci
Rep 7:40101.

6.   Wrighton KC, Thomas BC, Sharon I, Miller CS, Castelle CJ, VerBerkmoes NC, Wilkins MJ,
Hettins RL, Lipton MS, Williams KH, Long PE, Banfield JF. 2012. Fermentation, hydrogen, and
sulfur metabolism in multiple uncultivated bacterial phyla. Science 337:1661-1665.

7.   Nelson WC, Stegen JC. 2015. The reduced genomes of Parcubacteria (OD1) contain signatures of a
symbiotic lifestyle. Front Microbiol 6:713.

8.   Huddy RJ, Sachdeva R, Kadzinga F, Kantor RS, Harrison STL, Banfield JF. 2021. Thiocyanate and
Organic Carbon Inputs Drive Convergent Selection for Specific Autotrophic Afipia and
Thiobacillus Strains Within Complex Microbiomes. Frontiers in Microbiology 12.

9.   He C, Keren R, Whittaker ML, Farag IF, Doudna JA, Cate JHD, Banfield JF. 2021. Genome-
resolved metagenomics reveals site-specific diversity of episymbiotic CPR bacteria and DPANN
archaea in groundwater ecosystems. Nat Microbiol 6:354-365.

10.   Alvarado LEV, Fakra SC, Probst AJ, Giska JR, Jaffe AL, Oltrogge LM, West-Roberts J, Rowland
J, Manga M, Savage DF, Greening C, Baker BJ, Banfield JF. 2022. Autotrophic biofilms sustained
by deeply-sourced groundwater host diverse CPR bacteria implicated in sulfur and hydrogen
metabolism doi:10.1101/2022.11.17.516901. Cold Spring Harbor Laboratory.

11.   He X, McLean JS, Edlund A, Yooseph S, Hall AP, Liu SY, Dorrestein PC, Esquenazi E, Hunter
RC, Cheng G, Nelson KE, Lux R, Shi W. 2015. Cultivation of a human-associated TM7 phylotype
reveals a reduced genome and epibiotic parasitic lifestyle. Proc Natl Acad Sci U S A 112:244-249.

12.   Tian R, Ning D, He Z, Zhang P, Spencer SJ, Gao S, Shi W, Wu L, Zhang Y, Yang Y, Adams BG,
Rocha AM, Detienne BL, Lowe KA, Joyner DC, Klingeman DM, Arkin AP, Fields MW, Hazen
TC, Stahl DA, Alm EJ, Zhou J. 2020. Small and mighty: adaptation of superphylum
Patescibacteria to groundwater environment drives their genome simplicity. Microbiome 8:51.

13.   Jaffe AL, Castelle CJ, Matheus Carnevali PB, Gribaldo S, Banfield JF. 2020. The rise of diversity
in metabolic platforms across the Candidate Phyla Radiation. BMC Biol 18:69.

14.   Nales M, Butler BJ, Edwards EA. 1998. Anaerobic Benzene Biodegradation: A Microcosm Survey.
Bioremediation Journal 2:125-144.

15.   Ulrich AC, Edwards EA. 2003. Physiological and molecular characterization of anaerobic benzene-
degrading mixed cultures. Environ Microbiol 5:92-102.

16.   Luo F. 2016. Characterization of the Microbial Community Composition and Benzene Activation
Mechanisms in Anaerobic Benzene-Degrading Enrichment Cultures. PhD. University of Toronto,
TSpace.

17.   Toth CRA, Luo F, Bawa N, Webb J, Guo S, Dworatzek S, Edwards EA. 2021. Anaerobic Benzene
Biodegradation Linked to the Growth of Highly Specific Bacterial Clades. Environ Sci Technol
55:7970-7980.

18.   Luo F, Devine CE, Edwards EA. 2016. Cultivating microbial dark matter in benzene-degrading
methanogenic consortia. Environ Microbiol 18:2923-2936.

19.   Guo S, Toth CRA, Luo F, Chen X, Xiao J, Edwards EA. 2022. Transient Oxygen Exposure Causes
Profound and Lasting Changes to a Benzene-Degrading Methanogenic Community. BioRxiv
doi:10.1101/2022.04.11.487956.

20. Griebler C, Lueders T. 2009. Microbial biodiversity in groundwater ecosystems. Freshw Biol
54:649-677.

21. Kocur CMD, Lomheim L, Molenda O, Weber KP, Austrins LM, Sleep BE, Boparai HK, Edwards
EA, O'Carroll DM. 2016. Long-Term Field Study of Microbial Community and Dechlorinating
Activity Following Carboxymethyl Cellulose-Stabilized Nanoscale Zero-Valent Iron Injection.
Environmental Science & Technology 50:7658-7670.

22. Suzuki S, Ishii SI, Wu A, Cheung A, Tenney A, Wanger G, Kuenen JG, Nealson KH. 2013.
Microbial diversity in The Cedars, an ultrabasic, ultrareducing, and low salinity serpentinizing
ecosystem. Proceedings of the National Academy of Sciences 110:15336-15341.

23. Anantharaman K, Brown CT, Hug LA, Sharon I, Castelle CJ, Probst AJ, Thomas BC, Singh A,
Wilkins MJ, Karaoz U, Brodie EL, Williams KH, Hubbard SS, Banfield JF. 2016. Thousands of
microbial genomes shed light on interconnected biogeochemical processes in an aquifer system.
Nat Commun 7:13219.

24. Thiel J, Byrne JM, Kappler A, Schink B, Pester M. 2019. Pyrite formation from FeS and H2S is
mediated through microbial redox activity. Proc Natl Acad Sci U S A 116:6897-6902.

25. Ulrich AC, Edwards EA. 2003. Physiological and molecular characterization of anaerobic benzene-
degrading mixed cultures. Environmental Microbiology 5:92-102.

26. Luef B, Frischkorn KR, Wrighton KC, Holman HY, Birarda G, Thomas BC, Singh A, Williams
KH, Siegerist CE, Tringe SG, Downing KH, Comolli LR, Banfield JF. 2015. Diverse uncultivated
ultra-small bacterial cells in groundwater. Nat Commun 6:6372.

27. Kuroda K, Yamamoto K, Nakai R, Hirakata Y, Kubota K, Nobu Masaru K, Narihiro T. 2022.
Symbiosis between Candidatus Patescibacteria and Archaea Discovered in Wastewater-Treating
Bioreactors. mBio 0:e01711-22.

28. Miyoshi T, Iwatsuki T, Naganuma T. 2005. Phylogenetic characterization of 16S rRNA gene
clones from deep-groundwater microorganisms that pass through 0.2-micrometer-pore-size filters.
Appl Environ Microbiol 71:1084-1088.

29. Poulsen LK, Ballard G, Stahl DA. 1993. Use of rRNA fluorescence in situ hybridization for
measuring the activity of single cells in young and established biofilms. Appl Environ Microbiol
59:1354-1360.

30. Beveridge TJ, Patel GB, Harris BJ, Sprott GD. 1986. The ultrastructure of *Methanothrix concilii*, a
mesophilic aceticlastic methanogen. Can J Microbiol 32:703-710.

31. Holmes DE, Shrestha PM, Walker DJF, Dang Y, Nevin KP, Woodard TL, Lovley DR. 2017.
Metatranscriptomic evidence for direct interspecies electron transfer between Geobacter and
Methanothrix species in methanogenic rice paddy soils. Applied and Environmental Microbiology
83.

32. Vigil-Stenman T, Ininbergs K, Bergman B, Ekman M. 2017. High abundance and expression of
transposases in bacteria from the Baltic Sea. The ISME Journal 11:2611-2623.

33. Vigil-Stenman T, Larsson J, Nylander JAA, Bergman B. 2015. Local hopping mobile DNA
implicated in pseudogene formation and reductive evolution in an obligate cyanobacteria-plant
symbiosis. BMC Genomics 16:193.

34. Walker A, Langridge G. 2008. Does my genome look big in this? Nature Reviews Microbiology
6:878-879.

35.   Ran L, Larsson J, Vigil-Stenman T, Nylander JAA, Ininbergs K, Zheng W-W, Lapidus A, Lowry
S, Haselkorn R, Bergman B. 2010. Genome Erosion in a Nitrogen-Fixing Vertically Transmitted
Endosymbiotic Multicellular Cyanobacterium. PLoS ONE 5:e11486.

36.   Van Melderen L, Saavedra De Bast M. 2009. Bacterial toxin-antitoxin systems: more than selfish
entities? PLoS Genet 5:e1000437.

37.   Pandey DP, Gerdes K. 2005. Toxin–antitoxin loci are highly abundant in free-living but lost from
host-associated prokaryotes. Nucleic Acids Research 33:966-976.

38.   Magnuson RD. 2007. Hypothetical functions of toxin-antitoxin systems. J Bacteriol 189:6089-
6092.

39.   Craig L, Pique ME, Tainer JA. 2004. Type IV pilus structure and bacterial pathogenicity. Nature
Reviews Microbiology 2004 2:5 2:363-378.

40.   Saier MH, Reddy BL. 2015. Holins in Bacteria, Eukaryotes, and Archaea: Multifunctional
Xenologues with Potential Biotechnological and Biomedical Applications. Journal of Bacteriology
197:7-17.

41.   Peabody CR, Chung YJ, Yen M-R, Vidal-Ingigliardi D, Pugsley AP, Saier MH. 2003. Type II
protein secretion and its relationship to bacterial type IV pili and archaeal flagella. Microbiology
149:3051-3072.

42.   Chen L-X, Al-Shayeb B, Méheust R, Li W-J, Doudna JA, Banfield JF. 2019. Candidate Phyla
Radiation Roizmanbacteria From Hot Springs Have Novel and Unexpectedly Abundant CRISPR-
Cas Systems. Frontiers in Microbiology 10.

43.   de Luca G, de Philip P, Rousset M, Belaich JP, Dermoun Z. 1998. The NADP-Reducing
Hydrogenase ofDesulfovibrio fructosovorans:Evidence for a Native Complex with Hydrogen-
Dependent Methyl-Viologen-Reducing Activity. Biochemical and Biophysical Research
Communications 248:591-596.

44.   Arnaouteli S, Giastas P, Andreou A, Tzanodaskalaki M, Aldridge C, Tzartos SJ, Vollmer W,
Eliopoulos E, Bouriotis V. 2015. Two Putative Polysaccharide Deacetylases Are Required for
Osmotic Stability and Cell Shape Maintenance in Bacillus anthracis. Journal of Biological
Chemistry 290:13465-13478.

45.   Caufrier F, Martinou A, Dupont C, Bouriotis V. 2003. Carbohydrate esterase family 4 enzymes:
substrate specificity. Carbohydrate Research 338:687-692.

46.   Psylinakis E, Boneca IG, Mavromatis K, Deli A, Hayhurst E, Foster SJ, Vårum KM, Bouriotis V.
2005. Peptidoglycan N-Acetylglucosamine Deacetylases from Bacillus cereus, Highly Conserved
Proteins in Bacillus anthracis. Journal of Biological Chemistry 280:30856-30863.

47.   Vollmer W, Tomasz A. 2000. The pgdA Gene Encodes for a PeptidoglycanN-Acetylglucosamine
Deacetylase in Streptococcus pneumoniae. Journal of Biological Chemistry 275:20496-20501.

48.   Tsigos I, Martinou A, Kafetzopoulos D, Bouriotis V. 2000. Chitin deacetylases: new, versatile tools
in biotechnology. Trends in Biotechnology 18:305-312.

49.   Kafetzopoulos D, Martinou A, Bouriotis V. 1993. Bioconversion of chitin to chitosan: purification
and characterization of chitin deacetylase from Mucor rouxii. Proceedings of the National
Academy of Sciences 90:2564-2568.

50. Dörr T, Lam H, Alvarez L, Cava F, Davis BM, Waldor MK. 2014. A Novel Peptidoglycan Binding
Protein Crucial for PBP1A-Mediated Cell Wall Biogenesis in Vibrio cholerae. PLOS Genetics
10:e1004433.

51. Siewering K, Jain S, Friedrich C, Webber-Birungi MT, Semchonok DA, Binzen I, Wagner A,
Huntley S, Kahnt J, Klingl A, Boekema EJ, Søgaard-Andersen L, van der Does C. 2014.
Peptidoglycan-binding protein TsaP functions in surface assembly of type IV pili. Proceedings of
the National Academy of Sciences 111:E953-E961.

52. Kantor RS, Wrighton KC, Handley KM, Sharon I, Hug LA, Castelle CJ, Thomas BC, Banfield JF.
2013. Small genomes and sparse metabolisms of sediment-associated bacteria from four candidate
phyla. mBio 4:e00708-13.

53. Faust K, Raes J. 2016. CoNet app: inference of biological association networks using Cytoscape.
F1000Research 5:1519.

54. Raskin L, Poulsen LK, Noguera DR, Rittmann BE, Stahl DA. 1994. Quantification of
methanogenic groups in anaerobic biological reactors by oligonucleotide probe hybridization. Appl
Environ Microbiol 60:1241-1248.

55. Toth CRA, Molenda O, Nesbø CL , Luo F, Devine C, Guo S, Chen X, Edwards EA. 2023.
Metagenomic and Genomic Sequences from a Methanogenic Benzene-Degrading Consortium.
Microbiology Resource Announcements – MRA01342-22 - Submitted Dec 20, 2022. In review

  Figure and Legends

678
Figure 1. Results of Growth Trial #1. a) Absolute *Ca.* Nealsonbacteria abundance measured by
qPCR in copies/mL, and b) microbial community profiles (relative abundance) as a function of
substrate after 4 months of incubation, with total abundance shown above bars. In panel a) one
DNA sample was collected on day 1 to represent initial conditions. Subsequent DNA samples were
collected by sacrificing one entire bottle at each time point for each treatment due to the limited
culture volume. In panel b) ASVs with relative abundances < 1% were combined as "Other". ASVs
with the same taxonomy at the genus level were combined. The number of combined ASVs for
each taxon is shown in brackets. Absolute abundance was obtained by adding the qPCR results for
total bacteria, total archaea and *Ca.* Nealsonbacteria together. "Parent Benzene culture" column is
a 10% transfer of the parent culture at time zero.

690
691
Figure 2. *Ca.* Nealsonbacteria strain abundance over 7 months of incubation. Figure shows
treatments amended with benzene culture lysate, acetate, or pyruvate from Donor Trial #1. DNA
samples were collected by sacrificing one entire bottle at each time point for each treatment due
to the limited culture volume. Absolute abundance of *Ca.* Nealsonbacteria strains ASVa and ASVb
were calculated by multiplying the relative abundance of these ASVs (obtained from amplicon
sequencing) by the absolute abundance of total bacterial.
698
699
700

Figure 3. Heatmap of absolute abundance of top 20 ASVs in growth Trial #2. Top 20 ASVs were selected based on the sum of absolute abundance across all DNA samples collected. The ASVs under *Desulfosporosinus* are combined. Rows were clustered using a hierarchical clustering algorithm "hclust" in R based on similarity of absolute abundance. The columns represent different treatment bottles: INOC: Bottle with only 10% inoculum at time 0; NI: no inoculum (lysate only); Py&An: pyruvate & antibiotics; Lysate: French pressed benzene culture; Perm: lysate supernatant permeate through 50 kDa membrane; Ret: lysate supernatant retentate; Pellet: lysate pellet after centrifugation at 13,000 x g for 20 min; EC lys: French pressed lysate of *E. coli* culture; ST: Starved culture. The bracketed ASV numbers in the row names represent ordinal ASV number (see Table S3).

714
715
Figure 4. Various microscopic images of *Ca.* Nealsonbacteria in enrichment cultures. (a)
Epifluorescence microscopy showing archaea (red) and DAPI (yellow). Images are false colored
and overlaid using ImageJ. Images on the right are zoomed in views of green rectangular area in
the left image highlighting archaeal filaments with and without attached cells. (b) SEM images
of methanogenic benzene-degrading culture prepared by ionic liquid exchange. A *Methanothrix*
filament is marked with green arrow. *Ca.* Nealsonbacteria-like cells are marked with a green
circle. (c) Cryto-TEM images of *Ca.* Nealsonbacteria enrichments. *Ca.* Nealsonbacteria-like
cells are marked by green arrows and the right-side shows a zoomed in view of two cells that
appear to be budding. Additional microscopic images are shown in Figures S5 and S6.
725

(a) Cumulative benzene degraded

(b) qPCR results on Day 217

727
728 **Figure 5. Impact of adding *Ca.* Nealsonbacteria enrichment to a slow benzene-degrading**
729 **culture**. a) Cumulative benzene consumption and b) qPCR results tracking *Ca.* Nealsonbacteria
730 and ORM2 on Day 217. The qPCR results are sorted by descending order of benzene degradation
731 rate. The biological replicates are plotted separately to show the differences.

737
738
739 **Figure 6. A summary of key findings from the genome of *Ca.* Nealsonbacteria DGGOD1a**.
740 Detailed annotations for each ORF are in Table S10.
741
742
743
744
AUTHOR CONTRIBUTIONS

All authors made research and substantial intellectual contributions to the completion of this
study. X.C. and E.A.E. conceived and designed the study, and X.C carried out all experiments,
data analysis and interpretation, and drafted and revised the manuscript. O.M. assembled the
OD1 genome with help of C.N. and C.T.B. L.M performed the Cryo-TEM data acquisition and
analysis. C.R.A.T. helped with data analysis and revised the manuscript. S.G. helped to maintain
the culture, performed qPCR data acquisition and analysis and revised the manuscript. F.L.
enriched the culture, extracted DNA and performed the initial metagenome analysis. J.H. assisted
with ionic liquid-SEM data acquisition and analysis. C.H. performed the OD1 taxonomy
classification, phylogenetic tree and OD1 genome annotation. J.H.D.C. and J.F.B. provided
critical research guidance, and manuscript revisions.
756

34