Candidatus Nealsonbacteria (OD1) in a methanogenic benzene-degrading enrichment culture is likely an ectosymbiotic biomass recycler

Xu Chen[1], Olivia Molenda[1], Christopher T. Brown[2,3†], Courtney R. A. Toth[1], Shen Guo[1], Fei Luo[1††], Jane Howe[4], Camilla L. Nesbø[1], Christine He[5], Elizabeth A. Montabana[2,6], Jamie H. D. Cate[2,6,7], Jillian F. Banfield[5,6,8,9], and Elizabeth A. Edwards[1,*]

[1]Department of Chemical Engineering and Applied Chemistry, University of Toronto, Toronto, Ontario, M5S 3E5, Canada.
[2]Department of Molecular and Cell Biology, University of California, Berkeley, CA, USA
[3]California Institute for Quantitative Biosciences, University of California, Berkeley, CA, USA
[4]Department of Materials Science and Engineering, University of Toronto, 184 College Street, Toronto, Ontario M5S 3E4, Canada
[5]Innovative Genomics Institute, University of California, Berkeley, CA, USA
[6]Molecular Biophysics and Integrated Bioimaging, Lawrence Berkeley National Laboratory, Berkeley, CA, USA
[7]Department of Chemistry, University of California, Berkeley, CA, USA
[8]Department of Earth and Planetary Science, University of California, Berkeley, CA, USA;
[9]Environmental Science, Policy and Management, University of California, Berkeley, CA, USA;
†Current Address: Metagenomi, Inc., 1545 Park Avenue, Emeryville, CA, USA
††Current Address: Liven Proteins Corporation, 200 College Street, Toronto, ON, Canada.

Keywords: Candidate Phyla Radiation, Nealsonbacteria, genomics, genome-resolved metagenomics, fluorescent in situ hybridization, CryoEM, methanogenic benzene biodegradation, biomass recycling

*Corresponding author mailing address: University of Toronto, 200 College St., Toronto, Ontario, M5S 3E5, Canada. Phone: (416) 946-0996. Fax: (416) 978-8605.
E-mail: elizabeth.edwards@utoronto.ca;

Abstract

The Candidate Phyla Radiation (CPR, or superphylum Patescibacteria) is a very large group of bacteria with few cultivated representatives first discovered by culture-independent metagenomic analyses. Within the CPR, the candidate phylum Parcubacteria (previously OD1) is prevalent in anoxic lake sediments and groundwater. We identified a bacterium belonging to the Parcubacteria in a methanogenic benzene-degrading enrichment culture originally derived from oil-contaminated sediments. *Candidatus* Nealsonbacteria DGGOD1a is the only bacterium other than a previously identified benzene-degrading fermenter (*Deltaproteobacteria* Candidate Sva0485 clade ORM2) consistently and abundantly detected in all active benzene-degrading transfers of this culture. Therefore, we hypothesized that DGGOD1a must serve an important role in sustaining anaerobic benzene metabolism in the consortium. Growth experiments using a variety of possible substrates suggested that it is involved in biomass recycling. Microscopic observations supported by molecular analyses and a closed genome revealed an epibiont lifestyle with very small *Ca.* Nealsonbacteria DGGOD1a closely associated with much larger *Methanosaeta*. The images reveal a first example of cross-domain episymbiosis that may apply to other *Ca.* Nealsonbacteria found in diverse environments.

Introduction

A methanogenic benzene degrading enrichment culture referred to as consortium OR (for Oil Refinery) was derived from contaminated sediments in 1995 and has since been maintained with only benzene as added electron donor and carbon source. Multiple bottles and lineages of the culture have been maintained with periodic transfers in anaerobic pre-reduced minimal medium (1). This culture harbors benzene-degrading fermenting organisms (Deltaproteobacterium ORM2), Parcubacteria (OD1, further classified as *Candidatus Nealsonbacteria* in this study), acetogens, methanogens and other less numerous microbes which together enable anaerobic benzene degradation at rates from 0.2 to 10 mg benzene/L/day (2-4). The mechanism for anaerobic benzene biodegradation in the OR consortium is still poorly understood. A recent study describes the current state of knowledge (3), revealing coordinated metabolic interaction between fermenters, acetoclastic methanogens, and hydrogenotrophic methanogens. The chief benzene-degrading fermenter was identified as a member of the Candidate clade Sva0485 in the Deltaproteobacteria, present in at least two clonal variants denoted ORM2a and ORM2b. Interestingly, besides the chief benzene degrader, a member of *Ca.* Nealsonbacteria is the only other microbe consistently found in all different consortium OR lineages (2). However, no direct correlation was found between *Ca.* Nealsonbacteria and benzene degradation rate. Given *Ca.* Nealsonbacteria's persistence and the high biomass turnover rates in anoxic hydrocarbon- and organohalide-contaminated habitats (5), perhaps as a result of exposure to toxic concentrations of hydrocarbons, we hypothesized that this microorganism might play a role in biomass recycling. The OR consortium provided us an opportunity to study *Ca.* Nealsonbacteria's metabolic potential *in vivo*.

The Parcubacteria belong to the superphylum referred to as the Candidate Phyla

Radiation (CPR) which was first discovered by culture-independent metagenomic analysis (6).

The CPR, believed to comprise over 15% of the Bacteria domain, is prevalent in lakes, sediment,

and groundwater (6, 7). The discovery of CPR recast the tree of life and further emphasizes the

need for better functional characterization. To date, Parcubacteria remain uncultured and are only

available as metagenome assembled genomes (MAGs), with 14 closed MAGs or cMAGs

publicly available as of Dec. 9th, 2021 (8-12). Parcubacteria genomes are extremely small (~1

Mb) and include a large portion of hypothetical proteins which lack homologs from well-studied

microbes. The reduced genomes of Parcubacteria betray limited metabolic ability, lacking

complete pathways for the biosynthesis of vitamins, lipids amino acids, and nucleotides (11, 13).

The lack of complete biosynthetic pathways or electron transport system suggests a fermentative

symbiotic lifestyle where Parcubacteria obtain nutrients, metabolites, and energy directly from

their host or partner cells, a process known as ectosymbiosis. The recent example of an enriched

aerobic co-culture of a member of the CPR (Saccharibacteria) together with *Actinomyces*

odontolyticus XH001 revealed a symbiont or parasite lifestyle (14). Saccharibacteria genomes

shares similarities with the Parcubacteria in that they are small and lack complete biosynthetic

pathways. Metagenomic analyses suggests that Parcubacteria may also be associated with a host

microbe (11, 12, 15, 16) but experimental evidence has yet to be gleaned because of the

difficulty in cultivating and enriching these tiny microorganisms.

Using both Illumina short reads and PacBio long reads, we successfully reconstructed the

fully closed genome sequence of the uncultured Parcubacteria DGGOD1a in our methanogenic

benzene-degrading culture. A maximum likelihood concatenated ribosomal protein tree (Figure

S1) places the genome within the candidate group Nealsonbacteria, which was first defined by

Anantharaman et al. (17). This study aimed to identify substrates for the *Ca.* Nealsonbacteria

DGGOD1a in benzene-degrading OR consortium, and to visualize and enumerate these cells in

the mixed culture using various microscopic techniques. We tested various potential substrates

for *Ca.* Nealsonbacteria and found that most significant enrichment occurred when the culture

was amended with crude cell lysate, increasing relative abundance from 5% to ~30% and

absolute abundance up to 10 times. Although specific components present in lysed cells used to

enrich *Ca.* Nealsonbacteria were not identified, the results suggest a potential role as a biomass

recycler. Fluorescence *in situ* hybridization (FISH) and electron microscopy (EM) revealed *Ca.*

Nealsonbacteria as an epibiont on the surface *Methanosaeta*, a methanogenic archaeon essential

for completing the biotransformation of benzene to methane. These discoveries begin to shed

light on the uncharacterized majority of the CPR.

Materials and Methods

Chemicals and microbial cultures. All chemicals were obtained from Sigma-Aldrich (Oakville, ON, Canada) at the highest purity available unless specified otherwise. The OR consortium is maintained in 100 mL to 2 L batch culture bottles at the University of Toronto and in larger vessels (>100 L) at SiREM laboratories (Guelph, Ontario, Canada); the scaled-up culture is referred to as DGG-B (3). Two OR consortium subculture bottles (or lineages) were used in the experiments described herein: a bottle called OR-1bBig was the inoculum for *Ca.* Nealsonbacteria enrichment Donor Trials #1 & #2 and a bottle called OR-p5 was the inoculum for the augmentation experiment. All cultures were grown in an iron sulfide (FeS)-reduced, bicarbonate-based mineral medium (MM medium) amended regularly (approx. 1 per month) with 10 mg/L benzene (2, 3).

Preparation of electron donor solutions for *Ca.* Nealsonbacteria enrichment trials. Two crude cell lysate solutions from the OR consortium and a culture of *Escherichia coli* were prepared using tangential flow filtration and French press as described in Text S1. Next, we prepared two solutions of specific biomass components, nucleic acids (salmon sperm DNA; Sigma-Aldrich, Deoxyribonucleic acid sodium salt) and a membrane lipid (L-α-phosphatidylethanolamine; Sigma-Aldrich, Type II, ≥97%, lyophilized powder from egg yolk). Acetate, pyruvate and hydrogen were also tested as donors as they are intermediates of benzene degradation. Stock solution concentrations are listed in Table S1a.

***Ca.* Nealsonbacteria Enrichment – Donor Trial #1.** To test if *Ca.* Nealsonbacteria DGGOD1a could grow on crude lysed biomass or specific sub-cellular components, 25 mL culture tubes with 9 mL MM medium were inoculated with 1 mL OR-1bBig culture and amended with 7 different

electron donor solutions (Table S1a) at approximately the same electron equivalents, targeting ~30

mg COD/L. Each condition was tested in quadruplicate and incubated in a Coy anaerobic glove

box for 7 months (supplied with 10% H_2, 10% CO_2, and 80% N_2). DNA samples were collected

at 2, 4, and 7 months after inoculation by sacrificing an entire replicate from each treatment.

***Ca.* Nealsonbacteria enrichment – Donor Trial #2.** After observing successful growth and

enrichment of *Ca.* Nealsonbacteria with crude cell lysate from the benzene culture in Donor Trial

#1, a second experiment was set up to test the impact of various biomass fractions of cell lysate.

Culture (500 mL) was concentrated by tangential flow and cells were lysed using the same French

press method described earlier (Text S1). The cell lysate was then divided into four fractions, as

seen in Figure S2. One fraction was used "as is" (BL for <u>B</u>enzene culture <u>L</u>ysate). The remaining

lysate was centrifuged (at 13,000 × g for 20 mins) to create a pellet fraction and a supernatant

fraction. The supernatant fraction was further separated by tangential flow filtration with 50kDa

cut-off membrane (Pellicon® XL cassette with Durapore® membrane) to create a permeate fraction

(< 50 kDa) and a retentate fraction (>50 kDa). Each of these fractions was tested separately (Table

S1b). Two additional treatments were included in Donor Trial #2. One consisted of pyruvate plus

the antibiotics kanamycin and vancomycin (1 mM) to inhibit bacteria without inhibiting

methanogens. A second additional treatment was amended with *E. coli* extract instead of benzene

culture extract. We ensured that the *E. coli* pellet was well washed with MM medium to remove

any residual *E. coli* rich growth medium that could confound results. Controls without substrate

(starved control; SC) and crude benzene culture lysate without inoculum (no inoculum; NI) were

included. All treatment were conducted in biological duplicates except the starved control which

was a singleton.

***Ca.* Nealsonbacteria augmentation trial.** This experiment made use of an inoculum from bottle OR-p5 with a very slow benzene degradation rate (~0.1 mg/L/day). Eight 25-mL culture vials were each filled with a 10 mL aliquot of this slow culture (see Text S2 for details). Subsequently, 5 mL of OR-p5 culture was removed from vials #1-2 and replaced with 5 mL of *Ca.* Nealsonbacteria enrichment culture BL (replicate #1) from Donor Trial #2. In vials #3-5, 5 mL of culture was removed and replaced with 5 mL of culture supernatant from a very active benzene culture (degradation rate of ~10 mg/L/day), prepared by centrifugation at 13,000 × g for 20 min. The remaining vials (#6-8) of slow culture were left as is. All vials were amended with neat benzene targeting 15 mg/L (liquid concentration) and reamended whenever concentrations decreased below 5 mg/L. Benzene and methane concentrations were monitored by GC/FID (see below). DNA samples were collected on day 217 for qPCR analysis.

DNA extraction, quantitative PCR, and Illumina amplicon sequencing. DNA from Donor Trial #1 was extracted from 10 mL culture samples; otherwise, DNA was extracted from 1 mL culture samples. Cells were harvested by centrifugation at 13,000 × g for 20 min at 4°C and pellets resuspended in residual (~0.05 mL) supernatant. DNA was extracted using the DNeasy PowerSoil Pro Kit (QIAGEN) following the manufacturer's protocol. Real-time quantitative polymerase chain reaction (qPCR) assays were performed to track the gene copy numbers of Bacteria and Archaea using universal 16S rRNA gene primers; abundances of *Ca.* Nealsonbacteria and Sva0485 clade Deltaproteobacteria (ORM2) were also tracked using specific 16S rRNA gene primers (Table S2). Standard curves for all the qPCR studies are reported in Table S3. 16S rRNA gene amplicon sequencing with primers targeting the V6-V8 region was conducted at the Genome

Quebec Innovation Centre (McGill University) as previously described (3). Raw reads were

processed into amplicon sequence variants (ASVs) and analyzed as described in Text S3.

Metagenome analysis. The OR consortium/DGG culture was sequenced using a combination of

Illumina paired-end (~150 bp reads, ~300 bp insert) and PacBio long-read library (~7000 bp

reads). Metagenomes were submitted to IMG under sequencing project Gp0324998. Detailed

description of assembly, binning and functional and taxonomic annotation can be found in Text

S4.

Epifluorescence microscopy and fluorescence in situ hybridization (FISH). The FISH probe

ARCH915 was used for visualizing all Archaea (18). Eight FISH probes (Table S4) targeting the

16S rRNA gene of *Ca.* Nealsonbacteria were newly designed but none yielded a signal

distinguishable from background noise. The protocol for epifluorescence microscopy is provided

in Text S5 and details into the design and testing of FISH probes in Text S6. A LIVE/DEAD™

BacLight™ Bacterial Viability kit (ThermoFisher) was used to determine the ratio of viable cells

in consortium OR (Text S5).

Scanning Electron Microscopy with Ionic Liquid and Cryo-EM. To maintain the wet native

state of microbial cells and to eliminate artifacts that may be introduced by sample drying and

coating, culture samples were prepared with an ionic liquid and examined by SEM, as detailed in

Text S7. Cryo-EM was also used to explore intracellular structure and spatial associations. Sample

preparation and Cryo-EM conditions are described in Text S8.

Analytical procedures. Benzene and methane were measured in culture bottles and vials by

injecting 0.3 mL headspace samples into a Hewlett-Packard 5890 Series II GC fitted with a GSQ

column (30 m x 0.53 mm I.D. PLOT column; J&W Scientific, Folsom, CA) as previously

described (2). External standards were used for calibration.

Accession numbers. The closed genome sequence of *Candidatus* Nealsonbacteria DGGOD1a is

available at the Joint Genome Institute (IMG) under Taxon ID 2791354853 and at NCBI under

accession no. CP092821. A Genbank accession number is being generated for amplicon sequences

and will be added when available.

180
181   **Results and Discussion**
182
qPCR reveals abundance of *Ca.* Nealsonbacteria and Live/Dead staining confirms

abundance of dead cells

A qPCR survey of 15 different methanogenic benzene-degrading subculture bottles was

conducted to measure the abundance of the two known strains of *Deltaproteobacteria* ORM2

(ORM2a and ORM2b) and *Ca.* Nealsonbacteria (formerly referred to as OD1), as well as total

bacteria and total archaea. The culture bottle named "OR-1bBig" was selected as the parent

culture for this study because it already had a high relative abundance of *Ca.* Nealsonbacteria

(~5% based on 16S rRNA amplicon sequencing results) and sufficient culture volume (1.5 L).

Microscopy using live/dead staining (Figure S3) revealed large proportions of dead cells in all

subculture bottles. Detailed cell counts were conducted for two OR subcultures (~20 microscopic

images per sample) yielding ~46-62% dead cells (Table S5). This is not all surprising given that

these cultures are maintained as fed-batch, with infrequent transfers into fresh medium and

correspondingly long residence times. The products of lysed cells are clearly a source of electron

donor in addition to benzene in these bottles. This led us to hypothesize that components from

lysed cells might serve as substrates for *Ca.* Nealsonbacteria.

***Ca.* Nealsonbacteria found most enriched in vials amended with lysed cells.**

In Donor Trial #1, we found that *Ca.* Nealsonbacteria became most enriched in vials amended with

lysed benzene culture (Figure 1a). In these vials, relative and absolute abundances of *Ca.*

Nealsonbacteria increased ~6-10 times over 4 months, reaching a maximum 30% relative

abundance and 5.1×10^6 copies/mL (Figure 1b; Table S6a). Thus *Ca.* Nealsonbacteria may grow

on components in lysed biomass. None of the other tested substrates, including salmon sperm

DNA, acetate, and a long-chain fatty acid resulted in noticeable enrichment of *Ca.* Nealsonbacteria

relative to benzene-amended controls. Some increases (5x) were measured in bottles amended with

pyruvate (up to 9.8×10^5 copies/mL), but the abundances of other microbes increased even more

such that the average relative abundance of *Ca.* Nealsonbacteria dropped from ~5% to 1% (Figure

1b; Table S7). It is not clear whether the growth of *Ca.* Nealsonbacteria resulted from the direct

utilization of pyruvate or as an indirect consequence of the increases in biomass and corresponding

lysed cells. Either way, sodium pyruvate was not selective for *Ca.* Nealsonbacteria.

Other observations were gleaned from this first trial from looking at the entire microbial. First, we

clearly see that relative abundances of the chief benzene degrader (ORM2) dropped significantly

when benzene was absent (Figure 1b). Considering the vials amended with cell lysate, the only

two bacterial ASVs that co-enriched with *Ca.* Nealsonbacteria belong to the *Syntrophaceae* and

Spirochaetaceae. The *Syntrophaceae* ASV was found to share 100% 16S RNA gene sequence

identity to an uncultured *Syntrophaceae* (MH665869) from a methanogenic pyrite forming culture

fed with FeS, H$_2$S, and CO$_2$ (19). It is also distantly related (96.4%) to *Smithella propionica*

(AF126282), a syntrophic propionate degrader (20). The *Spirochaetaceae* ASV we identified is

distantly related (95.1%) to a spirochaetes (KJ535432) from a methanogenic bioreactor (21).

Spirochaetes are also thought to be involved in biomass recycling and may use protein and

polysaccharides to produce ethanol and acetate (22).

Nonmetric dimensional scaling (NMDS) was applied to further analyze Donor Trial #1 ASV data

(Figure S4a), displaying the impact of adding different electron donors on microbial community

composition. Methane production in each treatment indicated that the added COD was largely

consumed within 4 months (Table S8). Therefore, data collected after 7 months are long-starved

and marked as "+" in NMDS analysis. The "starved control" (unfed culture) community is very

close to the communities from vials amended with cell lysate indicating that cell decay in the OR

culture leads to a similar community as the vials amended with benzene culture cell lysate – which

makes intuitive sense. Interestingly, we found that communities after 7 months of incubation did

not converge, even though they should all be enriching on dead biomass after such a long

incubation time, prompting consideration that different sources of cell biomass might enrich for

different recyclers. In fact, we expected to see all vials shift to a scavenger-dominated microbial

community after 7 months of incubation, and we were hoping to see the enrichment of *Ca.*

Nealsonbacteria in every vial after prolonged starvation. However, such enrichment was only

clearly observed in acetate and pyruvate-amended vials (Figure 2). It is possible that there was

insufficient biomass produced in the other vials to detect enrichment. Indeed, total cell

concentrations were highest in acetate and sodium pyruvate amended vials. Interestingly, we

observed two different *Ca.* Nealsonbacteria ASVs (labeled a and b) in the acetate- and pyruvate-

amended cultures, where *Ca.* Nealsonbacteria ASVb became more abundant than *Ca.*

Nealsonbacteria ASVa – the strain originally detected in benzene cultures (Figure 2) (2). These

two *Ca.* Nealsonbacteria ASVs show only 91.2% 16S rRNA sequence identity. This result

intriguingly suggests that different strains may exhibit a specific preference for the type of lysed

cell material they can metabolize, or that the a and b variants have different hosts, because the

acetate- and pyruvate-amended cultures selected for very different communities.

To further investigate the identity of specific substrates for *Ca.* Nealsonbacteria, a second trial

(Donor Trial #2) was conducted where benzene culture lysate and also *E. coli* cell lysate were

compared. Moreover, the lysed benzene culture was further separated into three sub-fractions by

centrifugation and tangential flow filtration with a 50kDa cut off (Figure S2) and each fraction was

amended to the benzene culture separately. The pyruvate treatment was repeated with added

antibiotics (1 mM each kanamycin and vancomycin) to try to increase the selectivity of pyruvate.

The results of Donor Trial #2 (raw data in Table S6b) were similar to those of Donor Trial #1, with

cell lysate resulting in significant ($>10\times$) growth (Figure S5) regardless of source. The sub-

fractions of benzene culture lysate also resulted in enrichment although we observed considerable

variability between duplicates (Figure S5). The *E. coli* crude lysate appeared most stimulatory, but

it was also provided at $10\times$ the concentration of the lysate from the benzene culture based on

measured COD (Table S1c). The fact that of the three fractions, the >50 kDa fraction (supernatant

retentate) was most effective after 4 months suggests that larger molecules are important; perhaps

we should have tested proteins. An NMDS analysis was also conducted on amplicon data from

Donor Trial #2 (Figure S4b). These results show that the microbial communities enriched on each

fraction of dead biomass are a bit different after the first two months (triangles), but that they

converged after four months (squares). Perhaps after four months, only larger biomolecules

remain.

To look more closely at the microbial composition in each treatment, a heatmap of absolute

abundance (Figure 3) was generated showing the top 18 ASVs in Donor Trial #2, with raw data

reported in Table S9. *Ca.* Nealsonbacteria ASVa was the abundant sequence variant detected, and

Methanosaeta and *Spirochaetaceae* were the only other two microbes consistently enriched with

Ca. Nealsonbacteria in all cell lysate amended vials. Vials amended with lysed *E. coli* enriched

some ASVs belonging to *Clostridium* and *Bacteroides* that were not enriched in other treatments

(Figure 3). The vials amended with antibiotics and pyruvate successfully inhibited the growth of

most bacteria including *Ca.* Nealsonbacteria. However, it is interesting to note that the culture's

benzene fermenter (ORM2) seems to have persisted in those treatments. The growth of

Methanomethylovorans and *Methanobacterium* in the presence of antibiotics confirms their

resistance and points to a possible means of enriching for ORM2. Taken together, the two donor

trials have confirmed that compound(s) present in cell lysate can be used to enrich *Ca.*

Nealsonbacteria. The identity of the compounds remains elusive, but the most parsimonious

answer would be certain higher MW proteins and amino acid products, as they would be found in

all of the fractions tested. Further studies will focus on these substrates.

Microscopy reveals *Ca.* Nealsonbacteria as an epibiont of *Methanosaeta*.

Seeing is believing, and to date, *Ca.* Nealsonbacteria has never been successfully visualized. Eight

specific FISH probes were designed targeting the 16S rRNA gene of *Ca.* Nealsonbacteria

DGGOD1a. However, none of the probes yielded a distinguishable signal from background noise.

This problem was also reported in a previous study with a different set of FISH probes to

Parcubacteria (23). Although FISH was unsuccessful for *Ca.* Nealsonbacteria, we found that using

a combination of a FISH probe targeting all Archaea and DAPI staining could distinguish the

archaea (methanogens) from other microbes, i.e., bacteria. Given that *Ca.* Nealsonbacteria was the

most abundant bacterial phylotype in several of our enrichment cultures (Figure 3), we aimed to

positively identify these cells based on their known abundance. We selected cell lysate enrichment

culture (Lysate replicate #2 at 2 months) because *Ca.* Nealsonbacteria abundance was above 10^6

cells/mL in this culture and all other bacterial phylotypes detected were more than an order of

magnitude less abundant (Figure 3). A representative image of this culture is shown in Figure 4(a).

This method was used to examine both the parent culture and the *Ca.* Nealsonbacteria enrichments.

In the *Ca.* Nealsonbacteria enrichments, we observed many DAPI-stained ultra-small (~0.2 μm)

cells associated with a single filamentous structure that stained with the archaeal FISH probe, and

therefore must be a filamentous methanogen. The ultra-small cell size matches the typical size of

CPR members (11, 15) and is most likely *Ca.* Nealsonbacteria. Using the approach, we counted

the number of small cells attached to long archaeal filaments. This cell count was conducted

thoroughly on two independent samples from *Ca.* Nealsonbacteria enrichment used for

microscopy. Twenty fields of view were imaged and counted for each sample. These small cells

were by far the most numerous bacteria in all views, and thus must correspond to *Ca.*

Nealsonbacteria, as this bacterial ASV was most numerous as well. The cell counts obtained were

2.3×10^6 and 4.7×10^6 cells/mL in the two samples (Table S10). Enumeration of gene copies by

qPCR in samples of DNA extracted from the same vials using primers specific for *Ca.*

Nealsonbacteria gave a result of 8.3×10^6 copies/mL which is of the same order of magnitude as

cell counts. Thus, we are confident that the tiny cells attached to Archaea are *Ca.* Nealsonbacteria.

Interestingly, the archaeal FISH probe was observed to bind more strongly to methanogens that

had many *Ca.* Nealsonbacteria cells attached, compared to naked methanogens (Figure 4a; bottom

right). Naked methanogens stained brightly with DAPI but only faintly with the archaeal FISH

probe. Since the FISH probe intensity indicates metabolic activity, perhaps methanogens with *Ca.*

Nealsonbacteria are more active and have a higher number of ribosomes per cell (24). *Ca.*

Nealsonbacteria's role as either a parasite or commensal organism is still unknown. Perhaps

methanogens without *Ca.* Nealsonbacteria are dying or non-viable, or that *Ca.* Nealsonbacteria

specifically target the most active cells.

SEM examination of the parent methanogenic benzene-degrading culture also showed a similar

association of very small *Ca.* Nealsonbacteria-like cells attached to a large filamentous cell

structure consisting of a chain of rod-shaped, blunt-ended cells. This filament was enclosed in a

tubular sheath and the flat ends distinctively match the characteristics of *Methanosaeta* (25),

shown in Figure 4b. The *Ca.* Nealsonbacteria-like cells are spherical with a diameter of 0.2 µm to

0.8 µm and they are always found in budding pairs, forming a peanut shape. Cryo-EM was also

used to exam the *Ca.* Nealsonbacteria enrichments. The Cryo-EM images (Figure 4c) show the

same association of peanut-shaped cells with *Methanosaeta*. The large black rod-shaped blurry

cells at the center of the image are *Methanosaeta* forming a sheathed chain. Multiple peanut-

shaped ultra-small cells were observed attached to the chain of *Methanosaeta* (Figure 4c). The *Ca.*

Nealsonbacteria-like cells are always found in pairs and the one attached to the methanogen is

usually bigger than the second one. A zoomed-in image (Figure 4c; right side) reveals the detailed

cell structure of this *Ca.* Nealsonbacteria-like cell. It has a very interesting membrane structure

with a thick sheath and an apparent S-layer. This thick sheath might explain why FISH probes

failed, as they may not be able to enter the cells. Additional FISH, SEM, Cryo-EM images are

provided in Figures S6 and S7.

We had attempted to enrich *Ca.* Nealsonbacteria by filtration through a 0.22-μm filter. However,

qPCR results of the filtrate showed no such enrichment: both the total bacterial and the *Ca.*

Nealsonbacteria concentrations decreased 100 fold (data not shown). This is in contrast to other

studies where 0.22-μm filters were successful at enriching ultra-small microbes (6, 23, 26).

Perhaps the strong attachment of *Ca.* Nealsonbacteria to its host caused this approach to fail. The

majority of *Ca.* Nealsonbacteria were seen associated with *Methanosaeta* while very few were

observed free-living, perhaps indicating that it is an obligate epibiont.

Impact of *Ca.* Nealsonbacteria on benzene degradation

To further study its role in benzene degradation, we decided to mix a *Ca.* Nealsonbacteria

enrichment (containing about 10^7 *Ca.* Nealsonbacteria cells/mL) with a "slow" benzene-degrading

culture bottle (containing about 10^7 ORM2 cells/mL). Raw qPCR and monitoring data for this

experiment are reported in Table S11. We observed a significant enhancement of benzene

degradation rate in the two replicates that were a 50:50 mix of the two cultures (Figure 5a), even

though mixing diluted the benzene degrader (ORM2) by 50%. Benzene culture amended with

supernatant from a healthy culture (degrading benzene at 10 mg/L/day benzene) as well as the

(slow) culture-only controls all showed variable, but overall slower benzene degradation (Figure

5a). We tried to see if the rate of benzene degradation would relate to the abundance of *Ca.*

Nealsonbacteria and ORM2 (Figure 5b). In general, benzene biodegradation rates were highest in

cultures with the highest measured ORM2 concentrations at Day 217, except for one culture

replicate (slow culture #1) with abundant ORM2 but no benzene degradation activity. In hindsight

we should have monitored these cultures by qPCR more often. The two vials amended with the

Ca. Nealsonbacteria enrichment do show highest cell concentrations for both ORM2 and *Ca.*

Nealsonbacteria. Therefore, *Ca.* Nealsonbacteria did enhance benzene degradation, maybe by

supplying ORM2 with nutrients or co-factors recycled from lysed biomass or some specific

metabolites. Further studies using isotope-labeled cell lysate and additional replicates may help to

investigate these complex interactions.

_Ca._ Nealsonbacteria DGGOD1a's genome matches with symbiotic epibiotic lifestyle

We successfully assembled and closed the genome of *Ca.* Nealsonbacteria DGGOD1a (Genbank:

CP092821 and IMG ID: 2791354853). The complete genome is tiny: a mere 1.16 Mbp with 1145

predicted coding regions, that includes 18 pairs of perfect repeats over 400 bp. Interestingly, the

majority of these repeats are predicted transposases, and in fact, this genome has an

overrepresentation of predicted transposases, a total of 34, which is the most found in any

Parcubacteria genome and possibly of the highest number of transposases per base pair of genomes

sequenced to date. Most of metabolic pathways annotated by automatic pipelines are incomplete

(Table S12). Key findings are summarized in Figure 6. As inferred from genomes of other

members of the CPR group, it can use the pentose phosphate pathway to bypass glycolysis. Also,

basic transport pathways, and pathways for the biosynthesis of most amino acids, vitamins, and

lipids are incomplete. The large ribosomal RNA subunit L1 is missing; a common feature of

members of the OD1-L1 group (6, 11). It is also worth mentioning that two toxin and five antitoxin

genes were found in the genome which could participate in its symbiotic or parasitic lifestyle (27,

28). Many pili, cell surface, and secretion genes were found in this genome which could also

support an epibiont lifestyle. A CRISPR system with the complex of cas4, cas1, and cas2 was also

found with 49 spacers. Unfortunately, a spacer target search against all available OR consortium

metagenomic DNA did not return a single hit other than the CRISPR matrix itself.

Implications

This study revealed that *Ca.* Nealsonbacteria likely contributes to biomass recycling in a

methanogenic benzene degrading enrichment culture, and in so doing enhances the growth of

benzene fermenters. Although we were unable to target *Ca.* Nealsonbacteria cells using specific

FISH probes, the substantial enrichment of this phylotype on lysed biomass provided a culture

background in which to be certain that tiny DAPI-stained cells observed by epifluorescence were

in fact *Ca.* Nealsonbacteria. Once identified, we could observe their association with

Methanosaeta, which was also consistent with relative abundance from amplicon sequencing data.

SEM and CryoEM further confirmed and refined these observations.

Ca. Nealsonbacteria grew and became enriched when provided with cell lysate in the

absence of benzene in this methanogenic benzene-degrading culture. This result strongly supports

a biomass recycling role for *Ca.* Nealsonbacteria in this and likely in other anaerobic environments

where *Ca.* Nealsonbacteria are found. Diverse microscopic observations combined with a highly

reduced genome corroborate a reliance on a host which in this culture was *Methanosaeta* in a clear

example of an unusual cross-kingdom episymbiosis. The observed direct attachment or connection

between *Ca.* Nealsonbacteria to the surface of *Methanosaeta* implies metabolite exchange or direct

electron transfer between the two cell types. Perhaps the pili encoded in the genome of *Ca.*

Nealsonbacteria are used for attachment to the host and possibly as a conduit electron transfer. *Ca.*

Nealsonbacteria were also found to be beneficial for the growth of benzene-metabolizing

Deltaproteobacteria ORM2 from Candidate clade Sva0485. Perhaps *Ca.* Nealsonbacteria help

recycle nutrients or cofactors required to initiate and sustain higher benzene biodegradation rates.

These results guide future experiments into specific relationships between *Ca.* Nealsonbacteria

and benzene degradation in these enrichment cultures, as well as to suggest possible analogous

roles in other anaerobic communities, particularly where energy input is limited, such as deep

subsurface environments.

References

1. Ulrich AC, Edwards EA. Physiological and molecular characterization of anaerobic

benzene-degrading mixed cultures. Environ Microbiol. 2003;5:92-102.

2. Luo F, Devine CE, Edwards EA. Cultivating microbial dark matter in benzene-degrading

methanogenic consortia. Environ Microbiol. 2016;18:2923-36.

3. Toth CRA, Luo F, Bawa N, Webb J, Guo S, Dworatzek S, et al. Anaerobic benzene

biodegradation linked to the growth of highly specific bacterial clades. Environ Sci

Technol. 2021;55(12):7970-80.

4. Guo S, Toth CRA, Luo F, Chen X, Xiao J, Edwards EA. Transient oxygen exposure causes

profound and lasting changes to a benzene-degrading methanogenic community. BioRxiv.

2022.

5. Griebler C, Lueders T. Microbial biodiversity in groundwater ecosystems. Freshw Biol.

2009;54(4):649-77.

6.   Brown CT, Hug LA, Thomas BC, Sharon I, Castelle CJ, Singh A, et al. Unusual biology

across a group comprising more than 15% of domain Bacteria. Nature.

2015;523(7559):208-11.

7.   Hug LA, Baker BJ, Anantharaman K, Brown CT, Probst AJ, Castelle CJ, et al. A new view

of the tree of life. Nat Microbiol. 2016;1:16048.

8.   Kantor RS, Wrighton KC, Handley KM, Sharon I, Hug LA, Castelle CJ, et al. Small

genomes and sparse metabolisms of sediment-associated bacteria from four candidate

phyla. mBio. 2013;4(5):e00708-13.

9.   Rinke C, Schwientek P, Sczyrba A, Ivanova NN, Anderson IJ, Cheng JF, et al. Insights

into the phylogeny and coding potential of microbial dark matter. Nature.

2013;499(7459):431-7.

10.   Nelson WC, Stegen JC. The reduced genomes of Parcubacteria (OD1) contain signatures

of a symbiotic lifestyle. Front Microbiol. 2015;6:713.

11.   Castelle CJ, Brown CT, Thomas BC, Williams KH, Banfield JF. Unusual respiratory

capacity and nitrogen metabolism in a Parcubacterium (OD1) of the Candidate Phyla

Radiation. Sci Rep. 2017;7:40101.

12.   Tian R, Ning D, He Z, Zhang P, Spencer SJ, Gao S, et al. Small and mighty: adaptation of

superphylum *Patescibacteria* to groundwater environment drives their genome simplicity.

Microbiome. 2020;8:51.

13.   Wrighton KC, Thomas BC, Sharon I, Miller CS, Castelle CJ, VerBerkmoes NC, et al.

Fermentation, hydrogen, and sulfur metabolism in multiple uncultivated bacterial phyla.

Science. 2012;337(6102):1661-5.

14.   He X, McLean JS, Edlund A, Yooseph S, Hall AP, Liu SY, et al. Cultivation of a human-
associated TM7 phylotype reveals a reduced genome and epibiotic parasitic lifestyle. Proc
Natl Acad Sci U S A. 2015;112(1):244-9.

15.   He C, Keren R, Whittaker ML, Farag IF, Doudna JA, Cate JHD, et al. Genome-resolved
metagenomics reveals site-specific diversity of episymbiotic CPR bacteria and DPANN
archaea in groundwater ecosystems. Nat Microbiol. 2021;6(3):354-65.

16.   Jaffe AL, Castelle CJ, Matheus Carnevali PB, Gribaldo S, Banfield JF. The rise of diversity
in metabolic platforms across the Candidate Phyla Radiation. BMC Biol. 2020;18:69.

17.   Anantharaman K, Brown CT, Hug LA, Sharon I, Castelle CJ, Probst AJ, et al. Thousands
of microbial genomes shed light on interconnected biogeochemical processes in an aquifer
system. Nat Commun. 2016;7:13219.

18.   Raskin L, Poulsen LK, Noguera DR, Rittmann BE, Stahl DA. Quantification of
methanogenic groups in anaerobic biological reactors by oligonucleotide probe
hybridization. Appl Environ Microbiol. 1994;60(4):1241-8.

19.   Thiel J, Byrne JM, Kappler A, Schink B, Pester M. Pyrite formation from FeS and H_2S is
mediated through microbial redox activity. Proc Natl Acad Sci U S A. 2019;116(14):6897-
902.

20.   Liu Y, Balkwill DL, Aldrich HC, Drake GR, Boone DR. Characterization of the anaerobic
propionate-degrading syntrophs *Smithella propionica* gen. nov., sp. nov. and
Syntrophobacter wolinii. International Journal of Systematic Bacteriology. 1999;49:545-
56.

21. Nobu MK, Narihiro T, Rinke C, Kamagata Y, Tringe SG, Woyke T, et al. Microbial dark

matter ecogenomics reveals complex synergistic networks in a methanogenic bioreactor.

ISME J. 2015;9(8):1710-22.

22. Dong X, Greening C, Brüls T, Conrad R, Guo K, Blaskowski S, et al. Fermentative

Spirochaetes mediate necromass recycling in anoxic hydrocarbon-contaminated habitats.

ISME J. 2018;12:2039-50.

23. Luef B, Frischkorn KR, Wrighton KC, Holman HY, Birarda G, Thomas BC, et al. Diverse

uncultivated ultra-small bacterial cells in groundwater. Nat Commun. 2015;6:6372.

24. Poulsen LK, Ballard G, Stahl DA. Use of rRNA fluorescence in situ hybridization for

measuring the activity of single cells in young and established biofilms. Appl Environ

Microbiol. 1993;59(5):1354-60.

25. Beveridge TJ, Patel GB, Harris BJ, Sprott GD. The ultrastructure of *Methanothrix concilii*,

a mesophilic aceticlastic methanogen. Can J Microbiol. 1986;32(9):703-10.

26. Miyoshi T, Iwatsuki T, Naganuma T. Phylogenetic characterization of 16S rRNA gene

clones from deep-groundwater microorganisms that pass through 0.2-micrometer-pore-

size filters. Appl Environ Microbiol. 2005;71(2):1084-8.

27. Magnuson RD. Hypothetical functions of toxin-antitoxin systems. J Bacteriol.

2007;189(17):6089-92.

28. Van Melderen L, Saavedra De Bast M. Bacterial toxin-antitoxin systems: more than selfish

entities? PLoS Genet. 2009;5(3):e1000437.

Figures and Figure Legends

Figure 1. Results of Growth Trial #1. a) Absolute *Ca.* Nealsonbacteria abundance measured by qPCR in copies/mL, and b) microbial community profiles (relative abundance) as a function of substrate after 4 months of incubation, with total abundance shown above bars. In panel a) one DNA sample was collected on day 1 to represent initial conditions. Subsequent DNA samples were collected by sacrificing one entire bottle at each time point for each treatment. In panel b) ASVs with relative abundance below 1% were combined as "other". ASVs with the same taxonomy at the genus level were combined. Total abundance was obtained by adding the qPCR results for general bacteria, general archaea and *Ca.* Nealsonbacteria together. "Parent Benzene culture" column is a 10% transfer of the parent culture at time zero.

Figure 2. *Ca.* **Nealsonbacteria strain abundance over 7 months of incubation.** Figure shows treatments amended with benzene culture lysate, acetate, or pyruvate from Donor Trial #1. Absolute abundance of *Ca.* Nealsonbacteria strains ASVa and ASVb were calculated by multiplying the relative abundance of these ASVs (obtained from amplicon sequencing) by the estimate of total bacterial abundance.

Figure 3. Heatmap of absolute abundance of top 18 ASVs in growth Trial #2. Rows were clustered using a hierarchical clustering algorithm "hclust" in R based on similarity of absolute abundance. The columns represent different treatment bottles: INOC: Bottle with only 10% inoculum at time 0; NI: no inoculum (lysate only); Py&An: pyruvate & antibiotics; Lysate: French pressed benzene culture; Perm: lysate supernatant permeate through 50 kDa membrane; Ret: lysate supernatant retentate; Pellet: lysate pellet after centrifugation at 13,000 x g for 20 min; EC lys: French pressed lysate of *E. coli* culture; ST: Starved culture.

518
Figure 4. Various microscopic images of *Ca.* Nealsonbacteria in enrichment cultures. (a)
Epifluorescence microscopy showing archaea (red) and DAPI (yellow). Images are false colored
and overlayed using ImageJ. Images on the right are zoomed in views of red rectangular area in
left image highlighting archaeal filaments with and without attached cells. (b) SEM images of
methanogenic benzene-degrading culture prepared by ionic liquid exchange. *Methanosaeta*
filament shown with red arrow. *Ca.* Nealsonbacteria-like cell is marked with a red circle. (c)
Cryto-TEM images of *Ca.* Nealsonbacteria enrichments. *Ca.* Nealsonbacteria-like cells are
marked by red arrows and the right-side shows a zoomed in view of two cells that appear to be
budding. Additional microscopic images are shown in Figures S6 and S7.

(a) Cumulative Benzene degraded

(b) qPCR results on Day 217

Figure 5. Impact of adding *Ca*. Nealsonbacteria enrichment to a slow benzene degrading culture. a) Cumulative benzene consumption and b) qPCR results tracking *Ca*. Nealsonbacteria and ORM2 on Day 217. The qPCR results are sorted by descending order of benzene degradation rate.

535
536 **Figure 6. A summary of key findings from the genome of *Ca.* Nealsonbacteria DGGOD1a**.
537 Detailed annotations for each ORF are in Table S12.