Latent dynamics of primary sensory cortical population activity is structured by fluctuations in the local field potential

Audrey Sederberg[1,2,3,4], Aurélie Pala[1,5], Garrett B Stanley[1]

[1]Wallace H Coulter Department of Biomedical Engineering, Georgia Institute of Technology and Emory University, Atlanta, GA, USA, [2]Department of Neuroscience, University of Minnesota, Minneapolis, MN, [3]School of Psychology, Georgia Institute of Technology, Atlanta, GA, USA, [4]School of Physics, Georgia Institute of Technology, Atlanta, GA, USA, [5]Department of Biology, Emory University, Atlanta, GA

Running headline: Multiscale dynamics of cortical states

With 38 text pages, 6 figures, and 2 tables.

*Correspondence: Audrey Sederberg

School of Psychology and School of Physics

Georgia Institute of Technology

Atlanta, GA 30332

Phone: 404-385-4470

Email: audrey.sederberg@gatech.edu

Keywords: **Cortical state; LFP; population spiking; latent dynamics**

Abstract (269 words)

As emerging technologies enable measurement of precise details of the activity within microcircuits at ever-increasing scales, there is a growing need to identify the salient features and patterns within the neural populations that represent physiologically and behaviorally relevant aspects of the network. Accumulating evidence from recordings of large neural populations suggests that neural population activity frequently exhibits relatively low-dimensional structure, with a small number of variables explaining a substantial fraction of the structure of the activity. While such structure has been observed across the brain, it is not known how reduced-dimension representations of neural population activity relate to classical metrics of "brain state," typically described in terms of fluctuations in the local field potential (LFP), single-cell activity, and behavioral metrics. Here, we relate the latent dynamics of spiking activity of populations of neurons in the whisker area of primary somatosensory cortex of awake mice to classic measurements of cortical state in S1. We found that a hidden Markov model fit the population spiking data well with a relatively small number of states, and that putative inhibitory neurons played an outsize role in determining the latent state dynamics. Spiking states inferred from the model were more informative of the cortical state than a direct readout of the spiking activity of single neurons or of the population. Further, the spiking states predicted both the trial-by-trial variability in sensory responses and one aspect of behavior, whisking activity. Our results show how classical measurements of brain state relate to neural population spiking dynamics at the scale of the microcircuit and provide an approach for quantitative mapping of brain state dynamics across brain areas.

Author Summary (126 words)

Brain states have long been known to strongly shape sensory perception, decision making, cognition, and movement. Brain state during wakefulness changes constantly, classically assessed using changes in the spectral features of the local field potential (LFP) and behavioral

measures. However, the connection between these classical measurements of brain state and

the collective dynamics of populations of neurons is unclear. Here we fit a latent-variable model

to population spiking activity, finding that latent variables inferred under the model are highly

predictive of cortical state changes and that the latent dynamics are profoundly shaped by

inhibitory cell activity. Our approach connects the activity patterns of ensembles of neurons to a

classical measurement of brain state and opens new avenues for investigating brain state

dynamics across diverse cortical areas.

Keywords: Cortical state; latent dynamics; local field potential; population spiking

____

Introduction (552 words)

Since the earliest electrophysiological measurements, it has been widely acknowledged that the

brain undergoes gross changes in activity across a wide range of conditions related to sleep,

arousal, attention, behavior, etc., collectively referred to as reflecting "brain state." These

changes are qualitatively apparent in measures such as the local field potential (LFP) and

electroencephalogram (EEG) and have been conventionally quantified through changes in the

spectral content across distinct frequency bands (1–5). Indeed, these measures have been

shown to be strongly correlated with externally observable behavior, and thus capture, to some

extent, behaviorally relevant changes in intrinsic brain activity. Although it has been shown in

numerous studies that these measures of brain state are predictive of general levels of

excitability, it remains unclear how they relate to the spiking activity of neural populations.

Descriptions of cortical state derive from decades of observations of behavior and careful circuit

dissections of fluctuations in neural activity and behavior (6). Multiple studies have shown

fluctuations in metrics of cortical activity are related to fluctuations in cognitive and sensorimotor

behaviors in humans (7,8) as well as in animal models (9–13). A key model system for such

studies is the mouse somatosensory cortex, in which states of wakefulness have been

extensively characterized using the LFP and the activity patterns of single neurons (5,14–21). In

the somatosensory cortex, "state" changes in spontaneous activity may be described on a

continuum, from "deactivated" to "activated" (22), with the deactivated state characterized by

local field potential (LFP) activity dominated by low-frequency components (1-10 Hz). The

cortical state prior to sensory stimulation is predictive of trial-by-trial variation in the sensory

response (16,23), and LFP state is informative of behavioral metrics, including whisking activity

(5). Studies of single neurons have revealed cell-type-specific changes in firing rates with

fluctuations in activation level (15,24,25), suggesting that the conventional notion of cortical state

may be directly reflected in correlated fluctuations across the neuronal population, but this has

not been studied in detail.

A picture that is emerging from analysis of simultaneous recordings of large populations of

neurons is that, in many systems, a small number of latent variables explain much of the

spatiotemporal structure of neural activity (26–29), and that frequently these latent variables are

related to observed behavior (28,29). Analytic approaches based on inferring these latent

dynamical variables can predict future neural activity as well as improve decoder performance

in brain-machine interface applications (30). However, the connection between inferred latent

variables and established measures of brain state is not well understood.

Here, we use unsupervised learning to model spontaneous spiking activity recorded in mouse

somatosensory cortex during wakefulness and relate the population spiking states inferred from

the model to classical LFP-based measures of state in mouse S1. We 'open up the black box'

of the model to examine the structure of learned states, finding that putative inhibitory neurons

play a determinative role in defining the population spiking state. We further examined the

relationship between spiking states and classic signatures of cortical state in S1 of awake mice,

finding that spiking states are highly informative of whisking activity as well as predictive of

fluctuations in sensory responses. Our results demonstrate how unsupervised statistical

modeling of populations of individual neuron spiking activity dynamics extends classical metrics

of brain state.

Results

Latent state model of spiking activity

We developed a data-driven modeling framework to relate latent states of spontaneous spiking

activity to LFP and whisking behavior characteristics and trial-by-trial variations in sensory-

evoked responses. Recordings of the LFP and single-neuron spiking activity were acquired

using linear multi-electrode arrays (Neuronexus A1x32-5mm-25-177-A32) in the primary

somatosensory cortex (S1) of the awake, head-fixed mouse while monitoring whisking activity

with videography (see Methods). We analyzed spiking activity from an average of 26 neurons in

each recording (range 9 to 32, N= 6 recordings, see Table 1; see Methods for spike sorting

metrics) distributed across layers 2 through 5 and categorized as FS (fast-spiking) inhibitory

neurons and RS (regular-spiking) putative excitatory neurons as a function of their spike

waveform.

Spontaneous spiking activity in a population of FS (blue) and RS (red) neurons (10-second

sample, Figure 1A) was used to fit a Poisson-emission hidden Markov model (HMM, see

Methods), which has previously been used to capture coordinated firing rate changes across

populations of neurons (31–34). The fitted parameters, estimated on half of the recording

(Methods), are comprised of firing rates $\lambda_n(s^{\mathrm{spk}})$ (spikes per 40-ms time bin) for each state s^{spk}

and single neuron n (Figure 1C) and the transition probabilities A_{ij} between states (Figure 1D).

In Figure 1, the states are ordered by the average firing rate across cells, and the cells are

grouped with the FS neurons (blue) and RS neurons (red) together.

Under the fitted model, we computed the probability that the population is in a particular state at

each point in time, shown over the 10 second epoch for this example (Figure 1B). To get a

qualitative sense of the structure captured by the model, note the "state 6" segments in the

population raster shown in Figure 1A: these periods are marked by higher rates in the FS

neurons (cells 26 to 32) and a few of the RS neurons (Figure 1C). Some spike-states have

overall lower firing rates than others (e.g., state 1 with 0.13 spikes/bin versus state 7 with 0.54

spikes/bin), but the distinction between states also derives from differences in the pattern of

firing, as quantified by the Bhattacharyya coefficient, a generalized measure of differences

between multinomial distributions, which ranges from 0 (no overlap) to 1 (full overlap). For

instance, state 4 and state 5 have similar average firing rates (0.29 spikes/bin and 0.36

spikes/bin, respectively), but are highly distinguishable based on spiking pattern (Bhattacharyya

coefficient of $2.8 \cdot 10^{-13}$; see Methods).

To assess model fit quality, we computed the likelihood on the reserved test set for models with

one to fifteen states. For all recordings, a multi-state model had a higher likelihood than either

the 1-state model or the shuffled model (Figure 1E, Table 1). However, it was not possible to

identify a clear best choice for the number of states based on a cross-validated likelihood

estimate. Thus, in subsequent analyses, we continue to show examples from the seven-state

model, as after this point, the gain from additional states is minimal. All subsequent analyses

were carried out for models fit with 2 to 10 states, with results shown in summary for different

numbers of states.

Figure 1. Multi-state dynamics of population spiking activity are primarily driven by FS cells.
A: Spontaneous spiking activity simultaneously recorded across 32 cells in awake mouse S1. Spikes from FS cells in blue; from RS cells in red.
B: States inferred by a Hidden Markov Model (HMM) during the spontaneous segment in (A). Each row is a state, and each column is a time bin. The shading (white to black) indicates the inferred probability of being in each state at each point in time.
C and D: Parameters of the HMM underlying inference in panel (B).
C: Firing rate parameters ($\lambda_n(s^{\text{spk}})$) across the population of cells, including FS cells (blue) and RS cells (red). Bar length is the rate in spikes per 40-ms bin.
D: Matrix of state transition probabilities (A_{ij}). States in B, C, and D have been ordered based on average state-dependent firing rate across all cells.
E: Gain in the log-likelihood relative to the one-state model for each recording (bold, example in A-D). All recordings were better fit by a model with multiple states than by the one-state (constant rate) model. Error bars are suppressed for clarity and are approximately 0.15 based on the variance across segments of the data (Methods).
F: Impact of individual cells on the fit to the population was assess by shuffling spike counts across time for each single neuron, then recomputing the likelihood using the single-neuron shuffle. Single-neuron shuffle log likelihood is plotted separately for FS neurons (blue) and RS neurons (red). Top horizontal line: full-population (no shuffle) likelihood for the 7-state model (shown, A-D).
G: Average single-neuron shuffle loss in the log-likelihood for all recordings. Shuffling an FS neuron decreased the log-likelihood more than shuffling an RS neuron.

Recording	Data (1-state model)	Data (7-state model)	Simulation (7-state)	Shuffled (7-state)
1-1	-428549	-399075	-401893	-437928
1-2	-295463	-279170	-269299	-287421
2-1	-426533	-400726	-401244	-431003
3-1	-255510	-220216	-217860	-252243
3-2	-508583	-469814	-448926	-500754
4-1	-198327	-185925	-184549	-200262

Table 1. Log likelihood calculations under Poisson-HMM. Comparison of data (or simulated data) log-likelihoods under the Poisson-HMM, for the recorded data (1-state), recorded data (7-state), simulated data (7-state), shuffled simulated data (7-state). Simulation length is matched to recorded data length, so comparisons can be made within each row, but not directly across rows.

We next examined the contribution of individual neurons to the model fit using a shuffle test. For

each neuron, we shuffled the spike counts across time, thus creating a surrogate dataset

preserving structure in all neurons except the shuffled cell. We recomputed the data likelihood

using this surrogate set and compare the single-neuron shuffle likelihood to the original likelihood

computed in Figure 1E. In the example recording (Figure 1F), we find that likelihood dropped

more when an FS neuron was shuffled than when an RS neuron was shuffled, and this trend

held across all recordings ($p < 0.01$, $z = 7.0$, hierarchical bootstrap over N=6 recordings, with

10^4 shuffles, Figure 1G). This effect was not entirely explained by the higher firing rates of FS

neurons relative to RS neurons, as we observed the same difference between FS and RS

neurons when comparing the coefficient of variation of state-dependent firing rates extracted

from model parameters ($\lambda_n(s^{spk})$; $p_{boot}<0.01$; $z = 3.1$, hierarchical bootstrap). We conclude that

FS neurons played a larger role within the model than RS neurons.

Finally, to assess overall model fit quality, we compared the likelihood computed for the test set

to an idealized case, in which observations were generated from the fitted model. Using models

$(A_{ij}, \lambda_n(s^{\mathrm{spk}})$, with 1 to 15 states) that were fit to the data as previously described, we simulated

observations, matched to the data length of the original recording. Qualitatively, we found that a

seven-state model (Figure 2A; parameters from Figure 1) exhibits coordinated fluctuations in

activity observed across multiple single neurons. Such co-fluctuations and temporal modulation

are missing in the one-state model (Figure 2B). In the simulation, the sequence of states

(simulation ground truth, Figure 2C, top) was known, and was accurately inferred from the

simulated observations (inferred states, Figure 2C, bottom). Thus, qualitatively, the multi-state

Poisson-HMM captures the structure across single neurons and across time observed in the

recorded data.

To quantitatively assess model fits, we compared likelihood of decoded states in the data to that

from ideal simulations and null tests generated by shuffling data. We first compared the likelihood

of observations from simulated spontaneous S1 spiking (from the known model) to that of the

recorded data (using a model fit on the reserved training set). Simulation length was matched to

that of the recorded data (see Methods). We plot the log-likelihood (normalized by the number

of time bins times the number of neurons) for data (filled black circles) and simulation (open

purple circles) as a function of the number of states in the model (Figure 2D). For this recording,

the log-likelihood of the data matched that of the simulated data for models with five states and

more, indicating a fit that is as good as if the model were the "true" generating model for the

data.

Figure 2. Neural data was fit by the HMM as well as surrogate data generated directly from the model.

A: Seven-state model simulation for 32 cells and 10 s duration using model fit (parameters shown in Figures 1C and 1D). A sequence of states was generated from the transition matrix (Figure 1D) and observations drawn according to the emission distributions (Poisson spiking, rates in Figure 1C).

B: One-state model simulation for 32 cells and 10 s duration.

C: Top: the ground-truth state sequence used to generate spiking pattern (B). Bottom: inferred state probabilities, where dark purple indicates high probability.

D: Log-likelihood of test-set (reserved) data under models fit with 1 to 15 states (filled black circles). Log-likelihood of simulated data, using the fitted model and matching data length (open purple symbols; error bars are the standard deviation across repetitions of the simulation).

Relationship of LFP and spiking states

We next quantified the relationship between spiking states and the LFP, using an established

metric for defining 'states' of the LFP (22). Here LFP state is determined by calculating a power

spectrogram of the LFP (Methods) and comparing the power in low-frequency (LF, 1-10 Hz)

components to the power in high-frequency (HF, 30-90 Hz) components (Fig. 3B). We refer to

these as "LFP state L" (LF-dominant, with high LF/HF, green, Figure 3A, B) and "LFP state H"

(HF-dominant, with low LF/HF, black, Figure 3A, B). The spiking states inferred during a segment

Figure 3. LFP states are coordinated with distinct states of spiking activity.

A: Example period of spontaneous activity in which LFP state transitioned from state H to state L, defined using Fourier amplitudes spectrum features (B).

B: Fourier spectrum of the LFP, showing the LF frequency band (1-10 Hz) and HF frequency band (30-90 Hz). Division into states (L, green; H, black) is by comparing average amplitude in LF to that in HF ("LF/HF"), following previous studies (22).

C - D: Inferred spiking states (colors) with overlaid spike raster during segment 1 (LFP state H, panel C) and segment 2 (LFP state L, panel D). White indicates state probability was less than 80%.

E: Distribution of spiking states observed during LFP state H (left) and during LFP state L (right). Spiking states 1 and 2 are more likely during LFP state H, while spiking states 5, 6, and 7 are more likely during LFP state L.

F: The transition matrix (A_{ij}, reproduced from Fig. 1D) ordered by firing rate.

G: Transition matrix re-ordered using relative fraction of time the LFP state was H or L while spiking activity was inferred to be in state i. Reordered transition matrix (right) has nearly block-diagonal structure.

H: Directed graph illustrating transition matrix. Common transitions ($A_{ij} > 0.05$) between states (circles, color as in B and C) are indicated by directed arrows between nodes (states) in the graph. Self-connections (A_{ii}) not shown.

of LFP state H (Figure 3C) and LFP state L (Figure 3D) are shown with the population spike

raster overlaid. Transparent overlay (graded from black to green) represents distinct spiking

states during the recording intervals for which spike-state was assigned with >80% probability.

This example illustrates a tendency toward observing spiking states 1 and 2 during LFP state H

(Figure 3C) and a tendency toward observing spiking states 5, 6, and 7 during LFP state L

(Figure 3D).

To quantify the tendency of specific spiking states to occur more often during LFP state H or L,

we divided the recording based on LFP state, then computed the histogram of inferred spiking

state restricted to state H (Fig. 3E, left) or state L (Fig. 3E, right). The color map used for spiking

states in Fig. 3 was ordered based on the LFP, using the ratio of time each spiking state was in

LFP state H to LFP state L. This ordering differs from that used in Figure 1, in which states were

sorted by average firing rate, which produced a transition matrix with many off-diagonal elements

(Figure 3F). In contrast, LFP-based sorting revealed nearly block-diagonal structure in the

spiking state transition matrix (Figure 3G). We visualized the spiking state transition matrix as a

directed graph (Figure 3H), with a node for each state and in which the thickness of the arrows

between spiking states represents the probability of a transition. Spiking state 1 is directly

connected to spiking state 2 (both more likely in LFP state H) but moving from spiking states 1

and 2 to states 5, 6, and 7(more likely in LFP state L) requires multiple steps. This indicates that

while traversing the graph according to the transition matrix probabilities, the system will tend to

stay in LFP-H spiking states or LFP-L spiking states for long periods of time.

Based on the tendency for spiking states to align with one LFP state or the other, we expected

that using the state representation of spiking activity would predict the LFP state (H or L). Thus,

we built a binary classifier (Figure 4) to predict the LFP state (H or L) based on either direct

spiking metrics (single neuron spike count, summed population spike count, or population

vector) or on the inferred spiking states obtained using the HMM. Because LF/HF required a 1-

second window to compute, for this analysis 40-ms time bins were sampled at 1 Hz across the

experiment, then divided randomly into training and test sets, balancing the number of

observations associated with each LFP state so that chance-level performance was 50%. For

decoding, we used an SVM classifier with a linear kernel (see Methods).

To examine what aspects of population activity were informative of LFP state, we built a series

of decoders. In the simplest decoder, the predictor was the summed spike counts across the full

population (single recording example, "summed population rate", Figure 4A, dashed black line),

which was a poor predictor of LFP state (accuracy, 54% ± 5%, range across 6 recordings 46%

to 59%). To utilize the structure in the population firing pattern, we next constructed a linear

classifier based on the population rate vector. The population vector classifier performed better

Figure 4. Decoding LFP state from spiking activity.

A. Decoding accuracy (chance level, 50%) of determining LFP state (H or L) from inferred spiking states (purple), population vector (solid black), and summed population rate (dashed) for an example recording (1-1).
B. Decoding accuracy, using single neuron activity (FS, left/blue; RS right/red), across all neurons.
C. Across 5 of 6 recordings, decoding accuracy is higher from inferred spiking states than from the population vector.

than the summed-activity classifier (Figure 4A, "population vector", solid black line), with

performance in this example recording of 66% ± 3% (across cross-validation folds). Finally, we

predicted LFP state using as input the spiking state inferred under models having from 2 to 10

spiking states (Figure 4A, "spiking states", solid purple line). The highest decoding performance

was achieved by models with at least 4 states, above which decoding performance leveled off.

To determine whether the accuracy of decoding LFP state was explained by a minority of high-

performing cells, we used as predictors spike counts from single FS or RS neurons (Figure 4B).

While a few FS neurons were informative of LFP state (accuracy of best FS in each recording,

58 ± 7%, range 53% to 70%), most neurons decoded LFP state poorly (average decoding

accuracy: 50% ± 2% (RS), N = 119; 52% ± 4% (FS), N = 39). Across five of six recordings, the

best-performing decoders used inferred spiking states (Figure 4C), achieving an average

accuracy of 71% ± 10% (n = 6 recordings, range 61% to 85%). Figure 4C shows the decoding

accuracy using 7 spiking states. In comparison, decoding with the population vector was lower

by 14 percentage points (n = 6 recordings, 57% ± 7%).

Spiking states are predictive of trial-by-trial variability in sensory-evoked responses

Given previous reports of the predictive nature of ongoing LFP activity on sensory evoked

responses in this pathway (16), the relationship between spiking states and the LFP suggests

similar predictive capabilities through the latent spiking states. We now directly examine the

extent to which latent spiking states are predictive of trial-by-trial variability in sensory-evoked

responses in single neurons.

Between periods of spontaneous activity, we recorded sensory-evoked responses to simple

punctate single-whisker stimuli as described previously (16,35). We first established that the

state of pre-stimulus activity, classified under the spiking state model developed for spontaneous

activity (Figure 1), was predictive of single-trial evoked responses. We quantified the sensory-

evoked responses, conditioned on the pre-stimulus state, and identified neurons with state-

dependent sensory-evoked responses at a statistically significant level (see Methods). Briefly,

using one second (1 s) of pre-stimulus activity, we classified each trial based on the pre-stimulus

spiking state that was the most probable state immediately prior to stimulus delivery. Trials were

partitioned based on pre-stimulus spiking states, with population rate parameters plotted in

Figure 5A and sensory responses plotted in Figure 5B. For comparison, black-outlined bars

show the peri-stimulus time histogram (PSTH) averaged across all trials (Figure 5B).

Qualitatively, in some states (state 1, Figure 5A, B) the sensory response was less than in other

states (e.g., state 4, 5), both in terms of the absolute number of spikes in the post-stimulus

window and the size of the evoked (i.e., subtracting the pre-stimulus spike count) response

(Figure 5B).

To quantify significance of apparent state-dependence in sensory responses of single neurons,

we used a shuffle test. A distribution of average response size by state was generated by

randomly assigning trials to each state. The empirical average state-dependent sensory

response was then compared to the shuffle distribution, represented as the percentile of the

magnitude of the state-dependent average evoked response within the boot-strapped

distribution (FS neurons, Fig. 5C, and RS neurons, Figure 5E; in a 7-state model). For reference,

if there were no dependence of sensory response on the spiking state, the distribution of

percentiles would be uniform (dashed black lines, Figure 5C, 5E). Based on this percentile value,

Figure 5. Population spiking states are predictive of variability in sensory responses in single neurons.

A: Patterns of firing rates across neurons in each state. Bars represent the model rate parameter (spikes per second) in single neurons in a 40-ms bin (as in Figure 1). State colors as in Figure 3.

B: Responses to whisker stimulation (punctate deflection at time indicated by gray line) when trials are sorted by the decoded pre-stimulus state. The single neuron shown as an example in B is marked in panel A. PSTH bin width is 5 ms. Significant state-dependence is indicated by a purple star (facilitated) or yellow star (suppressed). This is defined relative to the sensory response averaged across all trials (black outline) based on p < 0.01 in a 2-sided shuffle test (Methods).

C: Distribution of percentile values for state-averaged responses for FS neurons from all recordings for pre-stimulus-sorted (color). Black dashed line shows uniform (chance-level) distribution of percentile values.

D: Number of FS neurons with detected state-dependent responses using models with increasing numbers of states. Yellow/purple indicate suppressed/facilitated. Trial-shuffle control is shown in light colors.

E: Same as C, for RS neurons from all recordings.

F: Same as D, for RS neurons from all recordings.

G: State transition graph (as in Figure 3), with purple/yellow indicating presence of facilitated / suppressed evoked responses in that state. Thicker outline represents more neurons with significant state dependence. Table 2 reports counts of facilitated/suppressed neurons for each recording.

H: Cumulative count of observed values of ratio of time spent in LFP state L to time spent in LFP state H for spiking states identified as suppressed (yellow) and facilitated (purple). Note the log scale, so negative (positive) values are states more frequently associated with LFP state H (state L). States with suppressed responses tended to be associated with LFP state H. Totals do not match the neuron counts in D and F because some individual neurons had facilitated (or suppressed) responses in more than one state.

state-dependent responses were categorized as "suppressed" (percentile <0.5; yellow star) or

"facilitated" (percentile >99.5; purple star), or not significantly different from the distribution of

responses across all trials. Across the population of neurons, more than half of neurons (for 7-

state mode: 91 of 158 total, range 8 to 24 per recording, from N = 6 recordings) had at least one

state-dependent sensory-evoked response, exceeding the false-positive rate expected by our

criterion for rejection (1%).

Recording (total count)	7-state model facilitated count	7-state model suppressed count
1-1 (32)	12	14
1-2 (32)	9	12
2-1 (28)	5	8
3-1 (11)	7	12
3-2 (36)	19	34
4-1 (19)	9	15

Table 2. Number of state-dependent evoked responses detected in each recording, under the 7-state model.

The organization of state-dependence across cells was non-random, with state-dependent

responses more common in the FS population than the RS population. For instance, using the

7-state model, 58% of all FS neurons were facilitated in at least one state (range across

individual recordings: 37% to 80%, N = 6 recordings) and 70% of all FS neurons were

suppressed in at least one state (range: 37% to 100%, N = 6 recordings), while 21% of RS

neurons were facilitated (range: 9% to 31%) and 39% of RS neurons were suppressed (range:

23% to 54%). For this analysis, cells were only counted once, so if there were two states with

suppressed responses (as in Figure 5B), the neuron contributed once to the count in Figure 5D.

We note that, while we show the seven-state model for illustration, this choice was not critical

for the detection of state-dependent responses. Models with four or more states show a relatively

constant proportion of neurons with state-dependent responses (Figure 5D, F). We verified that

this is not simply a consequence of limited trial numbers by repeating the analysis for trial-

shuffled controls with matched trial counts per state (gray, "trial-shuffled", Figure 5D, F).

Finally, we examined the extent to which specific pre-stimulus spiking states were associated

with facilitated or suppressed responses. As in Figure 3, we show state transition graphs,

annotated to show the organization of facilitatory and suppressive pre-stimulus states. Rings

around each state indicate the fraction of single neurons with either facilitated (Figure 5G, purple)

or suppressed (Figure 5G, yellow) responses, relative to the average evoked response across

all trials, where thicker rings represent more neurons with significant state dependence.

Suppressed states (yellow) were consistently found in the LFP-state-H associated spiking states

(black), and a subset of LFP-state-L spiking states (lighter green) were associated with facilitated

responses (purple). Across all recordings (Figure 5H, using 7 spiking states), facilitated spiking

states were more likely to be observed during LFP state L (median time ratio, 1.6), while

suppressed spiking states were more likely to be observed during LFP state H (median time

ratio, 0.6).

Spiking states track whisking activity.

During wakefulness, LFP state (quantified by LF/HF) is correlated with whisking behavior (5).

Thus, we expect that inferred spiking states are informative of whisking behavior as well,

illustrated by aligned trajectories of LFP (Figure 6A), spiking states (Figure 6B), and whisking

behavior (Figure 6C) over an example 25-second period. As in Fig. 3 and 5, spiking states are

ordered based on the LFP state observed but utilizing a color map to enable easier discrimination

between spiking states. Spiking states 1 and 2 are those for which low LF/HF is most likely. In

the example shown in Fig. 6B-C, spiking states 1 and 2 (dark blue shades) appear during bouts

of whisking (Fig. 6C), but rarely otherwise, suggesting that spiking state dynamics were closely

tied to whisking behavior. Across the full recording, the onset of whisking coincided with an

abrupt transition of the inferred spiking states into spiking states 1 or 2 (post-onset state is 1 or

2 in 25 of 38 detected onsets; Fig. 6D). By comparison, we observed a much slower decrease

in LF/HF (Fig. 6E).

To quantify this observation across datasets, we calculated mutual information (MI) between

whisking activity and (a) LFP states (based on $\mathrm{LF/HF}$) and (b) the inferred spiking states (see

Methods). For the example recording in Figure 6D, we found that the spiking states inferred

using a model with at least three states were more informative of whisking/non-whisking

behavior (0.25 ± 0.02 bits/bin, whisking-spike states, bin of 40 ms) than the LFP state was (0.15

± 0.02 bits/bin, Figure 6G). In all recordings, inferred spiking states (with at least 4 states,

showing 4, 7, and 10 states in Figure 6E, "whisk - spiking states") were more informative of

whisking activity than the LFP state ("whisk – LFP"), as shown in Figure 6H. We conclude that

modeling the latent dynamics of spontaneous population spiking activity provides a useful metric

of cortical state in awake mouse S1 that is related to traditional LFP-based metrics and contains

information about the representation of sensory information and changes in behavior with high

temporal resolution.

Figure 6. Spiking state dynamics are related to whisking activity.

A: LFP state metric ($\log_{10}$ (LF/HF) − 1) during spontaneous activity. Gray shaded regions show where LF/HF < 10 and white for LF/HF > 10.

B: Spiking activity across 15 neurons and the inferred latent state during the same period of spontaneous activity. Colormap (differing from Figure 3, 5) was chosen for higher contrast between states.

C: Whisking activity, quantified from motion energy (frame-to-frame changes, see Methods) in videography.

D-F: Onset of whisking is accompanied by a gradual decrease in LF/HF and rapid change of spiking states.

D: Onset times were identified using whisking energy, requiring a period of no whisking over 400 ms followed by crossing a threshold.

E: LFP state metric ($\log_{10}$ (LF/HF) − 1) gradually decreases at the onset of whisking. Gray lines: individual events. Solid black is the average.

F: Spiking state dynamics before and after onset of whisking, for 38 detected whisking events. State transition graph is shown for reference.

G: For this example, information between spiking state representation and whisking activity, for state models with 2 to 10 states. Information carried by comparing LF/HF to 10 is marked by the black line.

H: Information about whisking activity calculated on all six recordings for 4-state, 7-state, and 10-state models are all higher than the information carried by LF/HF > 10. Error bars estimated by resampling half of the data and repeating information calculation (50 repeats) and are typically smaller than the markers.

Discussion (1258 words)

Here we applied a latent state modeling framework to recordings from the awake mouse

somatosensory cortex and compared the latent states inferred from population spiking activity

to standard metrics of brain state. We found that latent spiking states captured structure in

population spiking activity and were closely linked to slow-changing LFP states, predictive of

fluctuations in single-trial sensory-evoked responses, and informative of whisking activity. Fast-

spiking neurons had higher impact on the latent state than regular-spiking neurons, reflecting a

large role of inhibitory cell activity in determining state.

Latent dynamics of population spiking activity in mouse S1

The latent-state representation of population activity inferred using a Poisson-HMM was highly

informative of the state of wakefulness, quantified by the standard measures of LF/HF and

whisking behavior. Spiking states were even more informative of whisking activity than the LF/HF

metric. Moreover, LFP states (based on LF/HF) were more accurately decoded from the latent

state representation of spiking activity than they were from more direct measures, such as the

summed firing rate across the population or even the population rate vector. The improvement

of the inferred spiking states from the HMM over the population rate vector likely arises due to

the temporal information embedded in the inferred spiking state, indicating the importance of

accounting for correlations in time for analysis of spiking data.

We also found that inferred spiking states were predictive of single-trial sensory-evoked

responses. In past work, we built a framework for predicting single-trial sensory-evoked LFP

responses in awake mouse S1, in which features of the pre-stimulus LFP recording were

identified based on their ability to predict response characteristics (16). There, we found that our

most predictive LFP features were equivalent to LF/HF, which had been used previously as a

metric of state in S1 (5,22). Thus, in the current study, we expected that pre-stimulus spiking

states would also be predictive of single-trial sensory-evoked responses in single neurons, and

that is what we found (Figure 5). The tendency of the "suppressed response" pre-stimulus states

to align with low LF/HF is consistent with the predictive features examined in our previous study,

which found smaller LFP responses in a low-LF/HF state. Thus, the Poisson-HMM framework

produces a representation of population dynamics that captures enough of the spatial (across

cells) and temporal (across time) structure in cortical activity to enable accurate decoding of the

LFP state and whisking behavior and to predict single-trial fluctuations in sensory responses in

single neurons.

FS neurons and population spiking state in mouse S1

We found that activity of FS neurons, which are more likely to be inhibitory cells (36), was more

strongly linked to the overall population spiking state than the activity of RS neurons, beyond

any dependence based on firing rate alone. For instance, model fit quality degraded substantially

more when the temporal structure of FS neuron activity was disrupted compared to that of RS

neuron activity. Further, a larger fraction of FS neurons than RS neurons exhibited state-

dependent sensory response patterns. Our general finding that spiking activity in inhibitory

neurons is tightly linked to population spiking state and, by extension, the LFP, is consistent with

past work focused on single-neuron contributions to the LFP that suggested that LFP primarily

reflects inhibitory activity (37). There is variability across single neurons within FS and RS

classifications, however: for example, not all FS neurons are tightly linked to the latent state.

Understanding the circuit dynamics that underlie the emergence of population spiking states will

require finer description of the cell-type specific roles, possibly enabled by new datasets with

more specific cell-type identity information (38) and by computational models permitting cell-type

classification based on single-neuron spiking statistics (39).

Insights from a simple latent-state model

In the Poisson-HMM used here to model spiking states, all cells were assumed to be linked to a single hidden state at any moment of time, and this state is assumed to have Markovian dynamics, with dependence only on the current state. While these assumptions limit the flexibility of the model, inferred spiking states from the model were successful in predicting LFP state, sensory-evoked responses, and whisking activity. Several previous studies have used the Poisson-HMM framework to describe population activity patterns and dynamics (10,33,40,41). For instance, the Poisson-HMM captured the structure of fast sequences of population activity patterns in rat gustatory cortex, demonstrating step-like rather than ramping behavior during a sensory-decision-making task (40). We used a coarser time bin, lasting 40 ms rather than $1-2$ ms in the Jones study. We chose this bin size because our analysis concerned spontaneous activity over relatively long (minutes) segments of data. Thus, our aim in using the Poisson-HMM is to obtain a relatively simple representation of spatial and temporal dynamics. The same model was also used to capture spontaneous activity states in the five seconds preceding a sensory-evoked response (33), and there it was argued that the identified states reflect underlying attractor dynamics. Due to the prevalence of FS neurons in the spiking patterns that determined spiking states, a different interpretation of states may be more appropriate in our study. An intriguing idea, raised by a recent modeling study, is that PV interneurons are poised to switch the cortical network in and out of an inhibition-stabilized network (ISN) dynamical state (42). Such state switches would be reflected in differences in sensory-evoked responses to well-controlled stimuli, as we observed, suggesting that changes in ISN dynamics may parsimoniously account for the state-dependent patterns of sensory-evoked responses we observed.

Limitations of Poisson-HMM for spiking dynamics

While the Poisson-HMM has the advantage of a small number of parameters and simple state

dynamics, the assumption that all neurons are in the same state likely breaks down in

populations spanning larger regions of cortex. An alternative approach, more commonly used in

non-sensory brain areas (motor, prefrontal, hippocampus) is to embed neuronal activity patterns

in a low-dimensional manifold upon which population activity smoothly evolves. For example,

latent structure in task-driven (non-spontaneous) population dynamics inferred utilizing a

dynamical system improved decoding of movement intention (43), and other studies have

identified low-dimensional structure in prefrontal cortex (44) and hippocampus (45). Responses

in visual cortex to visual inputs appear to live in a distinct subspace from spontaneous activity

(28). Determining which latent state model to use in which recording would be clarified by

developing rigorous "null" computational models, in which latent dynamics and population

responses can be carefully controlled (46,47).

Future directions

Understanding of the dynamics of local brain state has advanced tremendously through careful

dissection of modulatory circuit mechanisms in primary sensory areas. Experiments utilizing

patch-clamp electrophysiology in awake mice have revealed state-dependent changes in the

activity of single neurons of specific cell classes (15,25), shared patterns of variability across

small numbers of cells (48), and control of excitability through specific cell type activation (49).

In the large-scale multi-single-neuron recordings enabled by modern high-density multielectrode

recordings, such fine-scale anatomical information is rarely available. On the other hand, highly

parallel recording methods reveal functional relationships across the population that are not

observable from recordings of single neurons, and may carry rich information on the functional

roles of specific anatomical cell types. The latent dynamics uncovered by these approaches may

418   also provide functionally meaningful and pragmatic targets for control of circuit and network

419   activity as tools for applying control theory to neural circuits continue to emerge (50,51). A

420   clearer understanding, through quantitative frameworks such as that presented here, of these

421   features of cortical dynamics will ultimately provide an invaluable tool to map individual-specific

422   differences in the neural signatures of behavior, perception, and cognition.

Materials and Methods

Experimental data collection

Data presented here were previously reported in (35) and (16) and is available. All procedures were approved by the Institutional Animal Care and Use Committee at the Georgia Institute of Technology and agreed with guidelines established by the National Institutes of Health. Four nine- to twenty-six-week-old male C57BL/6J mice were used in this study. Mice were maintained under 1-2% isoflurane anesthesia while being implanted with a custom-made head-holder and a recording chamber. The location of the barrel column targeted for recording was functionally identified through intrinsic signal optical imaging (ISOI) under 1-1.25% isoflurane anesthesia. Recordings were targeted to B1, B2, C1, C2, and D2 barrel columns. Mice were habituated to head fixation, paw restraint and whisker stimulation for 3-6 days before proceeding to electrophysiological recordings.

Electrophysiology

Local field potential and spiking activity were recorded using silicon probes (A1x32-5mm-25-177, NeuroNexus, USA) with 32 recording sites along a single shank covering 775 μm in depth. The probe was coated with DiI (Invitrogen, USA) for post hoc identification of the recording site. The probe contacts were coated with a poly(3,4-160 ethylenedioxythiophene) (PEDOT) polymer (52) to increase signal-to-noise ratio. Contact impedance measured between 0.3 MOhm and 0.5 MOhm. The probe was inserted with a 35° angle relative to the vertical, until a depth of about 1000 μm. Continuous signals were acquired using a Cerebus acquisition system (Blackrock Microsystems, USA). Signals were amplified, filtered between 0.3 Hz and 7.5 kHz and digitized at 30 kHz.

Recording of spontaneous and sensory-evoked activity

Mechanical stimulation was delivered to a single contralateral whisker corresponding to the barrel column identified through ISOI using a galvo motor (Cambridge Technologies, USA). The

galvo motor was controlled with millisecond precision using a custom software written in Matlab

and Simulink Real-Time (Mathworks, USA). The whisker stimulus followed a sawtooth waveform

(16 ms duration) of various velocities (1000 deg/s, 500 deg/s, 250 deg/s, 100 deg/s) delivered

in the caudo-rostral direction (53). Whisker stimuli of different velocities were randomly

presented in blocks of 21 stimuli, with a pseudo-random inter-stimulus interval of 2 to 3 seconds

and an inter-block interval of a minimum of 20 seconds. The total number of trials across all

velocities presented during a recording session ranged from 196 to 616. Recordings were

acquired in 10-minute segments, and the end of each 10-minute segment was 1 minute of

spontaneous activity. All spontaneous segments of 10 seconds or longer were analyzed.

LFP pre-processing

For analysis, the LFP was down-sampled to 2 kHz. Using the same quality criterion as previously

(16), we screened for channels that drifted or had excessive 60-Hz line noise. Only layer-4 LFP

was analyzed in this study. Layer 4 was determined using the sensory-evoked MUA activity and

current source density profiles as described previously (16,35).

Spike sorting and single unit quality criteria.

Spike sorting was performed using Kilosort2 (54) with manual curation of spike clusters in Phy

(55). The quality criteria for identifying single unit activity were the same as in (35), summarized

by these six requirements: (i) units had more than 500 spikes, (ii) spike waveform SNR was

greater than 5, (iii) the trough-to-peak amplitude of the unit had a coefficient of variation less

than 0.2 averaging over 120-s windows, (iv) firing rate (over 120-s windows) had a coefficient of

variation of less than 1, (v) fewer that 1% of inter-spike intervals were shorter than 2 ms, and (vi)

the Maholonobis distance between the unit waveform and the closest waveform that was *not*

assigned to that unit was larger than 55 in the space of the first three principal components.

Units were classified as RS or FS based on the time elapsed from the waveform trough to

waveform peak, with values larger than 0.5 ms classified as RS units and those smaller than 0.4

as FS units (14,22). Intermediate waveforms (about 3% of units) were not included in the

analysis. Overall, we analyzed 39 FS units and 119 RS units across six recordings (average of

26 units per recording, range 19 to 36) in four different mice. Units were distributed across layers

2 to 5, with the large majority in layer 5 (L2/3, 13; L4, 24; L5, 117; L6, 4), precluding layer-

dependent analyses in this study.

Whisking quantification.

Video recordings of whisking activity were acquired under infrared illumination at 25 Hz using a

camera (6.8 pixels/mm, HQCAM) positioned below the mouse's head, acquiring a view of

whiskers as well as of the galvanometers used for stimulus control. For each movie, the portion

of the field of view including the whiskers was extracted. Whisking activity was extracted by first

identifying motion components using non-negative matrix factorization of the square difference

between frames. Up to 20 components were identified and these were manually scored as

representing whisker motion, nose motion, or galvanometer motion. Galvo traces were used to

verify temporal alignment of movies and electrophysiology. Whisking activity was quantified as

the sum over whisker-motion components.

Whisking onset analysis

To identify the transition from non-whisking to whisking behavior (analyses in Fig. 6D-F), we

analyzed the whisking motion time series, obtained by summing over projections of the whisking

moving into NMF modes described above. A fixed threshold for whisking was identified for each

recording. To identify a transition, we required 400 ms of non-whisking followed by a minimum

of two frames of whisking behavior. During the 400 ms of non-whisking behavior, at least 90%

of frames (9/10 frames) had to be below the detection threshold. Transition points occurring

within 500 ms of the beginning or end of a recording segment were discarded.

Experimental design and statistical analysis

Multiple statistical analyses were used in this study and are described in detail below. All

statistical analyses were conducted in MATLAB 2019b and 2022a (Mathworks).

Cortical State and Spiking State Model Fitting and Validation

Cortical state characterization.

To describe cortical state in S1 of awake mice, we calculated LF/HF and separated recordings

into periods with high (greater than 10) and low values of LF/HF, following a convention used in

earlier studies (5,22). Specifically, we compute the spectrogram at 1-s intervals and defining LF

(HF) as the average Fourier amplitude between 1 and 10 Hz (30 and 90 Hz, excluding 58 to 62

Hz).

Hidden Markov Model of cortical spiking activity

To describe cortical spiking states, we fit a Hidden Markov Model with Poisson emission

distribution (31,33,34,40). This model assumes that the next state $s^{\mathrm{spk}}(t + \Delta) \in [1,2, \dots N_S]$ only

depends on the current state $s^{\mathrm{spk}}(t)$, and this is characterized by the transition matrix A:

$$A_{ij} = P(s^{\mathrm{spk}}(t + \Delta) = j | s^{\mathrm{spk}}(t) = i)$$

511 We assuming firing is Poisson and conditionally independent across neurons. Specifically, the

512 distribution of spike counts $\{y_n\}$ over a time bin of duration Δ ($\Delta = 40\mathrm{ms}$) across the population

513 ($n = 1 \dots N_C$ neurons) while in spiking state s^{spk}, is given by

517 $$P(\{y_n\} | s^{\mathrm{spk}}) = \prod_{n=1}^{N_C} \frac{\left(\lambda_n(s^{\mathrm{spk}})\Delta\right)^{k_j} e^{-\lambda_n(s^{\mathrm{spk}})\Delta}}{y_n!}$$

where each state $s^{\mathrm{spk}} \in \{1, 2, \dots N_S\}$ is characterized by rate parameters $\lambda_n(s^{\mathrm{spk}})$. The joint

probability across a set of C simultaneously recorded cells is

$$P(\boldsymbol{y} = \boldsymbol{k} | s^{\mathrm{spk}}(t) = m) = \prod_{i=1}^{C} P(y_i = k_i | s^{\mathrm{spk}}(t) = m)$$

Thus, each state is defined by a vector of firing rates $\lambda_n(s^{\text{spk}})$. We binned spike counts at a

temporal resolution of 40 ms to match the resolution (25 Hz) of video monitoring of whisking

activity. In other works, the temporal resolution of spiking states has ranged from 1 ms (40) to

500 ms (31).

The model was implemented by adding a conditionally independent Poisson observation model

to the pmtk-3 (probabilistic models toolbox, https://github.com/probml/pmtk3). To fit the model,

we split each recording into two halves as follows. A recording consisted of a set of 8 to 12

segments, each of 10 minutes duration. Each segment had 10 periods of spontaneous activity

(duration, 10 s to 70 s) with no controlled whisker stimulation, interspersed with periods of

controlled whisker stimulation during which evoked responses were obtained. Model parameters

were estimated on a training set consisting of spontaneous periods from half of the segments

(1st, 3rd, 5th, etc.). Using the estimated model for each recording, states $s_i(t)$ at time t in each

spontaneous period in the test set (the remaining half of long segments: 2nd, 4th, 6th, etc.) were

decoded by estimating $p(s_i(t)|\{y(1{:}T))$ using standard methods (forward-backward algorithm).

Reported likelihood values are computed across all spontaneous periods in the test set.

We used two controls to interpret the data likelihood under the fitted model. First, to estimate the

ceiling of model performance, we simulated an HMM using the fitted state transition matrix and

drew observations from the emission distribution at each point in time according to the simulated

state. We then computed the likelihood of these simulated observations under the model.

Second, we verified that we could not obtain the same result in shuffled data. Thus, we shuffled

the simulated observations in time and recomputed the likelihood of shuffled observations under

the fitted model. Likelihood calculations for all recordings are listed in Table 1.

For the analysis of pre-stimulus state, we extracted the sensory-evoked from all segments, and

using the 500-ms pre-stimulus time window, we inferred HMM state using the model previously

fit on the spontaneous periods. The pre-stimulus state was assigned to the state with the highest

probability in the final pre-stimulus time bin (-40 ms to 0 ms pre-stimulus).

Quantifying differences between spiking states

To analyze the differences between states identified by the spiking state model, Bhattacharyya

coefficients and distances were computed for pairs of spiking state emission distributions. The

Bhattacharyya coefficient between two (potentially non-Gaussian) distributions quantifies their

overlap and determines the Bhattacharyya distance D_{BC} through the relationship $BC =$

$\exp\left(-D_{BC}\right)$ (56). For each cell i and pair of states (a, b), the state-dependent emission

distributions are Poisson distributions of rates $\lambda_i(a)$ and $\lambda_i(b)$, for which the Bhattacharyya

distance is

558
$$D_{BC}(a, b\ ; i) = \frac{1}{2}\left(\sqrt{\lambda_i(a)} - \sqrt{\lambda_i(b)}\right)^2 \Delta t$$

555   and Δt is the time bin (40 ms). Because we have assumed conditional independence within each

556   state, the full distribution across cells is a product distribution, and the distances add across

557   cells. The Bhattacharyya coefficient between states a and b is then

559
$$BC(a, b) = \exp\left(-\sum_i D_{BC}(a, b; i)\right)$$

This ranges from 0 to 1, where 1 indicates complete overlap ($\lambda_i(a) = \lambda_i(b)$) and 0 is no overlap.

Statistical tests for cell type differences across recordings

To compare the effects of FS and RS neurons on model fit quality, we compare two metrics.

First, we compare the loss in the likelihood relative to the original data in single-neuron-shuffle

surrogate datasets. For each single neuron, we generate a surrogate dataset from the reserved

test-set population raster, in which spike counts are shuffled in time *for that neuron only*. This

surrogate preserves the overall firing rates of all neurons in the population, enabling direct

comparison to the original data likelihood. For each neuron, we obtain the loss in the log

likelihood per time (T, total number of bins) by estimating the likelihood of the new state

sequence inferred from the surrogate dataset. The difference between single-neuron shuffle

surrogate log likelihood and the original log likelihood is taken as an estimate of how correlated

each neuron was with the population state.

The second metric we computed was the coefficient of variation (CV) of the rate parameters

($\lambda_i(m)$) across states m for each neuron i. This metric is an estimate of how much the rate for a

neuron varied with the learned states; higher CVs indicate that the rates vary more across states.

For both statistics, we assess the overall difference between RS and FS neurons across

recordings using a hierarchical bootstrap test and report *p*-values and *z*-scores (57).

Analyses of inferred spiking states

Ordering spiking states by LFP states
We ordered spiking states based on whether LFP state H or LFP state L was more likely to be

observed during the periods in which that spiking state was inferred (Figure 3). Specifically, we

computed $P(x|s_i)$, where s_i is the spiking state and x is the LFP state (either H, if LF/HF < 10, or

else L), and sorted states by the ratio: $P(H|s_i)/P(L|s_i)$.

Decoding analysis of LFP-spiking state relationships
Multiple classifiers were fit in Figure 4. For all classifiers, the target response is whether $\mathrm{LF/HF} >$

10. Observations were balanced so that 50% accuracy is chance and divided into test and

training sets. The inputs to the different types of classifiers are (i) summed spike count across

the population, (ii) the population spike count vector, (iii) the inferred spiking state, or (iv) the

single neuron spike counts. The classifier was a linear SVM.

Spiking state-dependent analysis of sensory-evoked responses

The evoked response was quantified as the difference between the spike count in the post-

stimulus window $[5\text{ms}, 25\text{ms}]$ and the pre-stimulus window $[-20\text{ms}, 0\text{ms}]$. Evoked spike counts

were computed for each trial, and we wanted to know if the average evoked response in each

state was higher or lower than would be expected from the null hypothesis that evoked

responses have no dependence on pre-stimulus spiking state. The number of trials assigned to

each state was not identical, thus we determined significance using a shuffle test. Specifically,

we randomly assigned trials to states matching the number of trials that were classified to that

state and computed the average evoked rate by state in each random draw. This was repeated

200 times for each recording and for each number of spiking states (2 to 10) to generate the

shuffled distribution of average evoked response. The empirical mean state-dependent evoked

spike count was compared to this distribution. Significance was determined by a two-tailed 1%

cut-off, so 'facilitated' responses were those in which the empirical mean state-dependent

evoked spike count scored above the 99.5% percentile of all shuffled mean evoked responses.

Information calculation between whisking activity and spiking state identity

Mutual information quantifies the difference between the joint distribution $P(X, Y)$ of quantities X

and Y compared to the distribution $P(X)P(Y)$ in which X and Y are independent, providing a

generalized measure of correlation (including non-linear correlation) between quantities X and

Y. Specifically, the mutual information (in bits) is $\langle \log_2 \frac{P(X,Y)}{P(X)P(Y)} \rangle$, where angle brackets denote an

average over the joint distribution $P(X, Y)$. For each calculation of mutual information (Fig. 6), X

is the whisking energy, computed in each 40-ms time bin (i) from videography and discretized

into four equipartition bins ($W_i \in \{0,1,2,3\}$). To compute information between whisking and

spiking states ("Y" is the spiking state), we identified the maximum-likelihood spiking state

$(s_{\text{spk}}(i) \in \{0, 1, \dots N_S\})$ in each time bin and estimated the mutual information between W_i and

$s_{spk}(i)$. To compute information between whisking energy and LFP state ("Y" is LFP state), we

assigned LFP state to L or H according to whether $r{>}0$ or $r{<}0$. Error bars were estimated by

resampling over half the data.

Code accessibility

Custom MATLAB code to generate figures and perform analyses can be accessed from GitHub

following publication.

References

1. Berger H. Über das Elektrenkephalogramm des Menschen. Arch Für Psychiatr Nervenkrankh. 1929 Dec 1;87(1):527–70.

2. Loomis AL, Harvey EN, Hobart GA. Cerebral states during sleep, as studied by human brain potentials. J Exp Psychol. 1937 Aug;21(2):127–44.

3. Başar E. EEG — Dynamics and Evoked Potentials in Sensory and Cognitive Processing by the Brain. In: Başar E, editor. Dynamics of Sensory and Cognitive Processing by the Brain. Berlin, Heidelberg: Springer Berlin Heidelberg; 1988. p. 30–55.

4. Steriade M, McCormick D, Sejnowski T. Thalamocortical oscillations in the sleeping and aroused brain. Science. 1993 Oct 29;262(5134):679–85.

5. Fernandez LMJ, Comte J christophe, Le Merre P, Lin JS, Salin PA, Crochet S. Highly Dynamic Spatiotemporal Organization of Low-Frequency Activities During Behavioral States in the Mouse Cerebral Cortex. Cereb Cortex. 2016 Oct 14;(January 2018):5444–62.

6. McCormick DA, Nestvogel DB, He BJ. Neuromodulation of Brain State and Behavior. Vol. 43, Annual Review of Neuroscience. 2020. p. 391–415.

7. He BJ, Zempel JM. Average Is Optimal: An Inverted-U Relationship between Trial-to-Trial Brain Activity and Behavioral Performance. PLoS Comput Biol. 2013;9(11).

8. Monto S, Palva S, Voipio J, Palva JM. Very Slow EEG Fluctuations Predict the Dynamics of Stimulus Detection and Oscillation Amplitudes in Humans. J Neurosci. 2008;28(33):8268–72.

9. Salkoff DB, Zagha E, McCarthy E, McCormick DA. Movement and Performance Explain Widespread Cortical Activity in a Visual Detection Task. Cereb Cortex. 2020;30(1):421–37.

10. Engel TA, Steinmetz NA, Gieselmann MA, Thiele A, Moore T, Boahen K. Selective modulation of cortical state during spatial attention. Science. 2016;354(6316):1140–4.

11. Speed A, Del Rosario J, Burgess CP, Haider B. Cortical State Fluctuations across Layers of V1 during Visual Spatial Perception. Cell Rep. 2019;26(11):2868-2874.e3.

12. Beaman CB, Eagleman SL, Dragoi V. Sensory coding accuracy and perceptual performance are improved during the desynchronized cortical state. Nat Commun. 2017;8(1):1–14.

13. Pinto L, Goard MJ, Estandian D, Xu M, Kwan AC, Lee SH, et al. Fast modulation of visual perception by basal forebrain cholinergic neurons. Nat Neurosci. 2013;16(12):1857–63.

14. Poulet JFA, Petersen CCH. Internal brain state regulates membrane potential synchrony in barrel cortex of behaving mice. Nature. 2008;454(7206):881–5.

15. Pala A, Petersen CC. State-dependent cell-type-specific membrane potential dynamics and unitary synaptic inputs in awake mice. eLife. 2018 Jul 27;7:1–13.

16. Sederberg AJ, Pala A, Zheng HJV V, He BJ, Stanley GB. State-aware detection of sensory stimuli in
the cortex of the awake mouse. PLOS Comput Biol. 2019 May 31;15(5):e1006716.

17. Petersen CCH, Hahn TTG, Mehta M, Grinvald A, Sakmann B. Interaction of sensory responses with
spontaneous depolarization in layer 2/3 barrel cortex. Proc Natl Acad Sci U S A.
2003;100(23):13638–43.

18. Senzai Y, Fernandez-Ruiz A, Buzsáki G. Layer-Specific Physiological Features and Interlaminar
Interactions in the Primary Visual Cortex of the Mouse. Neuron. 2019 Feb;101(3):500-513.e5.

19. Vinck M, Batista-brito R, Cardin JA, Vinck M, Batista-brito R, Knoblich U, et al. Arousal and
Locomotion Make Distinct Contributions to Cortical Activity Patterns and Visual Encoding Article
Arousal and Locomotion Make Distinct Contributions to Cortical Activity Patterns and Visual
Encoding. Neuron. 2015;86(3):740–54.

20. Bennett C, Arroyo S, Hestrin S. Subthreshold Mechanisms Underlying State-Dependent Modulation
of Visual Responses. Neuron. 2013 Oct;80(2):350–7.

21. Eggermann E, Kremer Y, Crochet S, Petersen CCH. Cholinergic Signals in Mouse Barrel Cortex during
Active Whisker Sensing. Cell Rep. 2014;9(5):1654–60.

22. Poulet JFA, Crochet S. The cortical states of wakefulness. Front Syst Neurosci. 2019;12(January):1–
18.

23. Haslinger R, Ulbert I, Moore CI, Brown EN, Devor A. Analysis of LFP phase predicts sensory
response of barrel cortex. J Neurophysiol. 2006;96(3):1658–63.

24. Gentet LJ, Kremer Y, Taniguchi H, Huang ZJ, Staiger JF, Petersen CCH. Unique functional properties
of somatostatin-expressing GABAergic neurons in mouse barrel cortex. Nat Neurosci.
2012;15(4):607–12.

25. Gentet LJ, Avermann M, Matyas F, Staiger JF, Petersen CCH. Membrane Potential Dynamics of
GABAergic Neurons in the Barrel Cortex of Behaving Mice. Neuron. 2010;65(3):422–35.

26. Köster U, Sohl-Dickstein J, Gray CM, Olshausen BA. Modeling Higher-Order Correlations within
Cortical Microcolumns. PLoS Comput Biol. 2014;10(7):1–12.

27. Okun M, Steinmetz NA, Cossell L, Iacaruso MF, Ko H, Barthó P, et al. Diverse coupling of neurons to
populations in sensory cortex. Nature. 2015;521(7553):511–5.

28. Stringer C, Pachitariu M, Steinmetz N, Reddy CB, Carandini M, Harris KD. Spontaneous behaviors
drive multidimensional, brainwide activity. Science. 2019 Apr 19;364(6437):eaav7893.

29. Musall S, Kaufman MT, Juavinett AL, Gluf S, Churchland AK. Single-trial neural dynamics are
dominated by richly varied movements. Nat Neurosci. 2019 Oct;22(10):1677–86.

30. Pandarinath C, O'Shea DJ, Collins J, Jozefowicz R, Stavisky SD, Kao JC, et al. Inferring single-trial
neural population dynamics using sequential auto-encoders. Nat Methods. 2018 Oct;15(10):805–
15.

31.   Abeles M, Bergman H, Gat I, Meilijson I, Seidemann E, Tishby N, et al. Cortical activity flips among
689   quasi-stationary states. Proc Natl Acad Sci. 1995 Sep 12;92(19):8616–20.

690   32.   Fontanini A, Katz DB. Behavioral States, Network States, and Sensory Response Variability. J
691   Neurophysiol. 2008;100(3):1160–8.

692   33.   Mazzucato L, Fontanini A, La Camera G. Dynamics of Multistable States during Ongoing and Evoked
693   Cortical Activity. J Neurosci. 2015 May 27;35(21):8214–31.

694   34.   Radons G, Becket D, Diilfer B. Analysis , classification , and coding of multielectrode spike trains
695   with hidden Markov models. 1994;373:359–73.

696   35.   Pala A, Stanley GB. Ipsilateral Stimulus Encoding in Primary and Secondary Somatosensory Cortex
697   of Awake Mice. J Neurosci. 2022 Mar 30;42(13):2701–15.

698   36.   Barthó P, Hirase H, Zugaro M, Harris KD, Zu- M, Monconduit L, et al. Characterization of Neocortical
699   Principal Cells and Interneurons by Network Interactions and Extracellular Features. J
700   Neurophysiol. 2004;92(1):600–8.

701   37.   Teleńczuk B, Dehghani N, Le Van Quyen M, Cash SS, Halgren E, Hatsopoulos NG, et al. Local field
702   potentials primarily reflect inhibitory neuron activity in human and monkey cortex. Sci Rep. 2017
703   Jan 11;7(1):40211.

704   38.   de Vries SEJ, Lecoq JA, Buice MA, Groblewski PA, Ocker GK, Oliver M, et al. A large-scale
705   standardized physiological survey reveals functional organization of the mouse visual cortex. Nat
706   Neurosci. 2020;23(1):138–51.

707   39.   Schneider A, Azabou M, McDougall-Vigier L, Parks DF, Ensley S, Bhaskaran-Nair K, et al.
708   Transcriptomic cell type structures in vivo neuronal activity across multiple timescales. Cell Rep.
709   2023 Apr;42(4):112318.

710   40.   Jones LM, Fontanini A, Sadacca BF, Miller P, Katz DB. Natural stimuli evoke dynamic sequences of
711   states in sensory cortical ensembles. Proc Natl Acad Sci U S A. 2007;104(47):18772–7.

712   41.   Prentice JS, Marre O, Ioffe ML, Loback AR, Tkačik G, Berry MJ. Error-Robust Modes of the Retinal
713   Population Code. Stevenson IH, editor. PLOS Comput Biol. 2016 Nov 17;12(11):e1005148.

714   42.   Bryson A, Berkovic SF, Petrou S, Grayden DB. State transitions through inhibitory interneurons in a
715   cortical network model. PLoS Comput Biol. 2021;17(10 October):1–29.

716   43.   Pandarinath C, Gilja V, Blabe CH, Nuyujukian P, Sarma AA, Sorice BL, et al. Neural population
717   dynamics in human motor cortex during movements in people with ALS. eLife. 2015;4:e07436.

718   44.   Mante V, Sussillo D, Shenoy K V, Newsome WT. Context-dependent computation by recurrent
719   dynamics in prefrontal cortex. Nature. 2013;503(7474):78–84.

720   45.   Nieh EH, Schottdorf M, Freeman NW, Low RJ, Lewallen S, Koay SA, et al. Geometry of abstract
721   learned knowledge in the hippocampus. Nature. 2021;595(7865):80–4.

46. Morrell MC, Sederberg AJ, Nemenman I. Latent Dynamical Variables Produce Signatures of
Spatiotemporal Criticality in Large Biological Systems. Phys Rev Lett. 2021;126(11):118302.

47. Sederberg A, Nemenman I. Randomly connected networks generate emergent selectivity and
predict decoding properties of large populations of neurons. PLOS Comput Biol. 2020 May
7;16(5):e1007875.

48. Arroyo S, Bennett C, Hestrin S. Correlation of Synaptic Inputs in the Visual Cortex of Awake,
Behaving Mice. Neuron. 2018;99(6):1289-1301.e2.

49. Fu Y, Tucciarone JM, Espinosa JS, Sheng N, Darcy DP, Nicoll RA, et al. A Cortical Circuit for Gain
Control by Behavioral State. 2014;1139–52.

50. Bolus MF, Willats AA, Rozell CJ, Stanley GB. State-space optimal feedback control of
optogenetically driven neural activity. J Neural Eng. 2021;18(3).

51. Weiss DA, Borsa AM, Pala A, Sederberg AJ, Stanley GB. A machine learning approach for real-time
cortical state estimation. J Neural Eng. 2024 Feb;21(1):016016.

52. Ludwig KA, Uram JD, Yang J, Martin DC, Kipke DR. Chronic neural recordings using silicon
microelectrode arrays electrochemically deposited with a poly(3,4-ethylenedioxythiophene)
(PEDOT) film. J Neural Eng. 2006;3(1):59–70.

53. Wang Q, Webber RM, Stanley GB. Thalamic synchrony and the adaptive gating of information flow
to cortex. Nat Neurosci. 2010;13(12):1534–41.

54. Pachitariu M, Steinmetz N, Kadir S, Carandini M, Harris K. Fast and accurate spike sorting of high-
channel count probes with KiloSort. Adv Neural Inf Process Syst. 2016;(Nips):4455–63.

55. Rossant C, Harris KD. Hardware-accelerated interactive data visualization for neuroscience in
Python. Front Neuroinformatics. 2013;7(DEC):1–9.

56. Bhattacharyya A. On a Measure of Divergence between Two Multinomial Populations. Sankhya
Indian J Stat. 1946;7(4):401–6.

57. Saravanan V, Berman GJ, Sober SJ. Application of the hierarchical bootstrap to multi-level data in
neuroscience. Neurons Behav Data Anal Theory. 2020;3(5):1–25.

748