Glucagon-Like Peptide-1 Receptor Activation Stimulates PKA-Mediated Phosphorylation

of Raptor and this Contributes to the Weight Loss Effect of Liraglutide

Running title: GLP-1R agonists and PKA phosphorylation of Raptor

Thao D. V. Le[1], Dianxin Liu[2.], Blair J. Ellis[1], Sheila Collins[1,2*], Julio E. Ayala[1*]

[1]Department of Molecular Physiology and Biophysics, Vanderbilt University School of Medicine,

2215 Garland Avenue, Nashville, TN 37232, USA

[2]Department of Medicine, Vanderbilt University Medical Center, 2215 Garland Avenue,

Nashville, TN 37232, USA

*Corresponding Authors

Email: julio.e.ayala@vanderbilt.edu

Email: sheila.collins@vumc.org

Abstract

Objective: The canonical target of the glucagon-like peptide-1 receptor (GLP-1R), Protein Kinase A (PKA), has been shown to stimulate mechanistic Target of Rapamycin Complex 1 (mTORC1) by phosphorylating the mTOR-regulating protein Raptor at Ser^{791} following β-adrenergic stimulation. The objective of these studies is to test whether GLP-1R agonists similarly stimulate mTORC1 via PKA phosphorylation of Raptor at Ser^{791} and whether this contributes to the weight loss effect of the therapeutic GLP-1R agonist liraglutide.

Methods: We measured phosphorylation of the mTORC1 signaling target ribosomal protein S6 in Chinese Hamster Ovary cells expressing GLP-1R (CHO-Glp1r) treated with liraglutide in combination with PKA inhibitors. We also assessed liraglutide-mediated phosphorylation of the PKA substrate RRXS*/T* motif in CHO-Glp1r cells expressing Myc-tagged wild-type (WT) Raptor or a PKA-resistant ($Ser^{791}Ala$) Raptor mutant. Finally, we measured the body weight response to liraglutide in WT mice and mice with a targeted knock-in of PKA-resistant $Ser^{791}Ala$ Raptor.

Results: Liraglutide increased phosphorylation of S6 and the PKA motif in WT Raptor in a PKA-dependent manner. Liraglutide failed to stimulate phosphorylation of the PKA motif in $Ser^{791}Ala$ Raptor. $Ser^{791}Ala$ Raptor knock-in mice were resistant to liraglutide-induced weight loss.

Conclusion: GLP-1R agonists promote PKA-mediated phosphorylation of Raptor at Ser^{791}, and this facilitates liraglutide-induced weight loss.

Introduction

Glucagon-like peptide-1 (GLP-1) receptor (GLP-1R) agonists reduce food intake and

body weight and are approved for weight loss[1]. Elucidating the mechanisms through which

GLP-1R agonists regulate body weight could enhance the therapeutic potential of these

compounds or identify novel targets for anti-obesity drug development. We have shown that the

GLP-1R agonist Exendin-4 (Ex4) stimulates mechanistic Target of Rapamycin (mTOR)

Complex-1 (mTORC1) signaling and that inhibition of mTORC1 signaling in the ventromedial

hypothalamus (VMH) with rapamycin blocks the anorectic effect induced by Ex4 in this

nucleus[2,3]. The present studies aim to define the mechanism(s) by which GLP-1R agonism

stimulates mTORC1 signaling and reduces body weight.

mTOR is a Ser/Thr protein kinase that forms two complexes: mTORC1 and mTOR

complex 2 (mTORC2)[4]. The rapamycin-sensitive mTORC1 has unique accessory proteins

including Raptor (regulatory associated protein of mTOR). mTORC1 activity is stimulated by

growth factors, nutrients, and hormones by promoting PI3K/Akt-mediated phosphorylation of

tuberous sclerosis complex 1/2 (TSC1/2)[5]. Activation of mTORC1 by GLP-1R agonists seemed

unlikely since the GLP-1R is a $G_{\alpha}s$-coupled receptor that signals via protein kinase A (PKA)[6].

However, activation of the $G_{\alpha}s$-coupled β-adrenergic receptor stimulates mTORC1 via Raptor

phosphorylation by PKA[7], suggesting that GLP-1R activation can similarly stimulate mTORC1

via PKA phosphorylation of Raptor.

Here we demonstrate that GLP-1R activation induces mTORC1 signaling and Raptor

phosphorylation in a PKA-dependent manner and show that this contributes to weight loss by

the therapeutic GLP-1R agonist liraglutide. This reveals a novel weight-lowering GLP-1R-PKA-

mTORC1 axis and broadly suggests that other $G_{\alpha}s$-coupled receptors may use this PKA-

mTORC1 pathway for certain biological responses.

Materials and Methods

Cell culture

Chinese Hamster Ovary cells expressing human GLP-1R (CHO-GLP-1R; DiscoveRx,

Fremont, CA) were maintained at 37°C (5% CO2, humidified) in Ham's F12 media containing

10mM glucose, 1mM pyruvate, and 1mM glutamine (Cat#11765054, Gibco) supplemented with

10% fetal bovine serum (FBS; Cat#16140071, Gibco), 10ml/l penicillin-streptomycin (Cat#91,

Hyclone), and 0.8mg/ml Geneticin G418 Sulfate (Cat#10131027, Gibco). β-TC3 cells[8] were

gifted by Dr. Richard O'Brien and cultured in DMEM media containing 25mM glucose, 1mM

pyruvate, and 1mM glutamine (Cat#11995065, Gibco) supplemented with 15% FBS, 2% horse

serum (Cat#26050088, Gibco) and 10ml/l penicillin-streptomycin. All cells were authenticated by

STR profiling (ATCC) and were verified to be free of mycoplasma. Cells were washed twice with

serum-free Ham's F12 (CHO-GLP-1R) or DMEM (β-TC3) media containing 5.5mM glucose for

3h and were treated with vehicle (1X PBS) or liraglutide (Victoza, Novo Nordisk) for 1h with or

without 30-min pre-treatment with vehicle (DMSO), PKA inhibitors H89 (Cat#2910, Tocris

Bioscience) or KT5720 (Cat#1288, Tocris Bioscience), or the Akt inhibitor MK-2207

(Cat#S1078, Selleck Chemicals LLC) at concentrations stated in the figure legends.

Immunoblotting

Cells were washed once with ice-cold 1X PBS and lysed in RIPA buffer (Cat#R0278,

Sigma-Aldrich) containing 1X protease and phosphatase inhibitors. After centrifugation of cell

homogenates (17,000xg, 10min, 4°C), supernatant protein concentration was determined by

BCA protein assay (Pierce BCA Protein Assay Kit, Cat#23225, Thermo Scientific). Twenty

μg/lane was run (200V, 45min) in 4%–20% Tris-glycine gels (Invitrogen) and transferred (100V,

1h, 4°C) onto PVDF membranes (Bio-Rad). Blots were blocked in buffer containing 5% BSA

(Cat#A9418, Sigma-Aldrich) in 1X Tris-Buffered Saline and 0.1% Tween®-20 for 1h and

incubated in primary antibodies (phospho-S6 Ser$^{235/236}$, Cat#4858, RRID:AB_916156; total-S6,

Cat#2317, RRID:AB_2238583; p-CREB, Cat#9198, RRID:AB_2561044; total-CREB, Cat#9104,

RRIS:AB_490881; phospho-Akt Thr[308], Cat#9275, RRID:AB_329828; total-Akt, Cat#9272,

RRID:AB_329827; and Myc tag, Cat#2276, RRID:AB_2314825; Cell Signaling Technology)

overnight at 4°C and secondary antibodies (IRDye 800CW Goat anti-Rabbit IgG, Cat#926-

32211, RRID:AB_621843 and IRDye® 680RD Goat anti-Mouse IgG, Cat#926-68070,

RRID:AB_10956588 LI-COR Biosciences) for 1h at room temperature. Blots were imaged with

an Odyssey Imaging System and full lanes were analyzed using ImageJ.

Immunoprecipitation

Myc-tagged wild-type (WT) and PKA-resistant mutant (Ser[791]Ala) Raptor vectors[7] were

transfected into CHO-GLP-1R cells using lipofectamine-3000 (Invitrogen). 48h later, cells were

washed 2X and maintained in serum-free media for 3h. Cells were treated with DMSO, H89 or

MK-2207 for 30min and treated with 1X PBS, liraglutide, forskolin, or insulin for 1h at

concentrations stated in the figure legend. Cells were washed once with ice-cold 1X PBS and

lysed in lysis buffer (25mM Tris HCl, 131 150mM NaCl, 1μM EDTA, 1μM EGTA, 2.5mM sodium

pyrophosphate, 100mM sodium fluoride, 1mM sodium orthovanadate) containing 1X protease

and phosphatase inhibitors. After centrifugation of cell homogenates (17,000xg, 10min, 4°C)

supernatant protein concentration was determined by BCA protein assay. 200μg protein was

incubated with anti–c-Myc affinity gel (EZview™ Red Anti-c-Myc Affinity Gel, #E6654, Sigma-

Aldrich) overnight. Beads were washed in lysis buffer and immunoblotting was done as

described above using a PKA substrate monoclonal antibody (Cat#9624, Cell Signaling

Technology).

Animals

Mice were kept on a 12h/12h light/dark cycle with *ad libitum* access to water and chow

(5L0D, LabDiet) from weaning and were studied at 12–16 weeks old. Procedures were

approved by the Institutional Animal Care and Use Committee at Vanderbilt University (Protocol

#M1500158). Ser^{791}Ala Raptor knock-in mice were generated by inGenious Targeting

Laboratory Inc. using the targeting vector shown in Supp. Fig. 1A. A two base pair point

mutation was introduced into a second inverted exon 20 of the Raptor coding sequence.

Raptor$^{fl/fl}$ mice (013188, The Jackson Laboratory) were crossed to mice expressing the

Ingenious construct, and offspring were then crossed to CMV-*Cre* (006054, The Jackson

Laboratory) mice to generate hemizygous wild-type mice or hemizygous mice with global knock-

in of Raptor Ser^{791}Ala. This strategy was chosen because the Ingenious construct in

homozygosity prior to Cre recombinase inadvertently created a hairpin loop in the transcript

between the two exon 20 regions that was spliced out. Genotyping and cDNA sequencing

confirmed expression of Ser^{791}Ala Raptor in place of endogenous Raptor (Supp. Fig. 1B). Male

mice were switched to low-fat diet (D12450J, Research Diets, Inc.) for 4 weeks prior to initiation

of dosing. Mice were maintained on the low-fat diet and were administered vehicle (0.9% saline)

or liraglutide (200μg/kg body weight, BID) subcutaneously for 14 days. Body weight was

measured daily, and body composition was obtained by NMR (Minispec 235 LF90II-TD NMR

Analyzer, Bruker) prior to and on the last day of dosing. 4h fasting blood glucose and insulin

levels were measured using a Contour Next EZ glucose meter (Cat#BD15789, Bayer) and an

Ultra Sensitive Mouse Insulin ELISA kit (Crystal Chem), respectively. A separate cohort of male

mice was acclimated to a Promethion System (Sable Systems International, Inc.) for one week

prior to undergoing dosing with vehicle or liraglutide as described above for 10 days. Food

intake and energy expenditure were measured continuously.

Data Analysis

Data were analyzed using Prism-9 (GraphPad Software, Inc.). One-way ANOVA or

mixed-effects analysis followed by Holm-Sidak's multiple comparisons was used when

appropriate and indicated in the figure legends. Sample size was determined based on the

variability of the primary measurements of interest (phosphorylation of S6 for cell culture studies

and body weight for *in vivo* studies) and the objective of detecting significance (p<0.05) with

90% certainty. Treatments were randomized and blinded to the experimenter when possible.

Results

GLP-1R agonists activate mTORC1 in a PKA-dependent manner

Similar to Ex4[2,3], liraglutide increases phosphorylation of the mTORC1 signaling target

ribosomal protein S6 in CHO-GLP-1R cells, and this is blocked by the mTORC1 inhibitor

rapamycin (Fig. 1A; Supp. Fig. 2A). The PKA inhibitor H89 also blocks S6 phosphorylation by

liraglutide or the adenylyl cyclase/PKA activator forskolin (Fig. 1B; Supp. Fig. 2B). H89 has off-

target effects inhibiting the S6 kinase p70S6K[9], likely explaining why H89 inhibits S6

phosphorylation by insulin (Fig. 1B). KT5720, a PKA inhibitor with minimal off-target effects on

p70S6K[9], also reduces liraglutide- and forskolin-induced phosphorylation of S6 and the PKA

target CREB in a dose-dependent manner (Fig. 1C; Supp. Fig. 2C). Neither dose of KT5720

affected S6 phosphorylation by insulin (Fig. 1C). Two doses of the pan-Akt inhibitor MK-2206

that block insulin-stimulated Akt phosphorylation did not affect stimulation of mTORC1 activity

by liraglutide (Fig. 1D; Supp. Fig. 2D). These data suggest that liraglutide stimulates mTORC1

signaling in a PKA-dependent and Akt-independent manner.

GLP-1R activation induces PKA phosphorylation of Raptor at Ser^{791}

The catalytic subunit of PKA directly phosphorylates Ser^{791} of Raptor[7]. We tested

whether liraglutide promotes PKA phosphorylation of Raptor at Ser^{791} by expressing Myc-tagged

wild-type (WT) Raptor or Myc-tagged mutant Raptor with alanine replacing serine at the 791st

position (Ser^{791}Ala Raptor) in CHO-GLP-1R cells. Using an antibody against the PKA motif

RRXS*/T*, immunoblotting of immunoprecipitated Myc-proteins shows that liraglutide and

forskolin increase PKA motif phosphorylation on Myc-WT Raptor in a PKA-dependent manner

(blocked by H89), but liraglutide fails to increase PKA motif phosphorylation on Myc-Ser^{791}Ala

Raptor (Fig. 2A and 2B; Supp. Fig. 3A). This was also observed in β-TC3 cells, an immortalized

β-cell line endogenously expressing the GLP-1R (Fig. 2C; Supp. Fig. 3C). A high concentration

of the Akt inhibitor MK-2206 (20μM) did not affect liraglutide-induced PKA motif phosphorylation

of Raptor in CHO-GLP-1R (Figs. 2A and 2B; Supp. Figs. 3A and 3B) or β-TC3 (Fig. 2C; Supp.

Fig. 3C) cells. Thus, liraglutide stimulates PKA phosphorylation of Raptor at Ser^{791}.

PKA phosphorylation of Raptor at Ser^{791} mediates GLP-1R agonist-induced weight loss

We tested the physiological role of PKA phosphorylation of Raptor at Ser^{791} by

measuring body weight and glycemic responses to liraglutide in wild-type (WT) and knock-in

mice in which endogenous Raptor is replaced by a PKA-resistant Raptor expressing alanine

instead of serine at position 791 in all tissues (CMV-Ser^{791}Ala Raptor). WT and CMV-Ser^{791}Ala

Raptor knock-in mice lost weight in response to liraglutide treatment, but liraglutide-induced

weight loss was attenuated in the CMV-Ser^{791}Ala Raptor knock-in mice (Figs. 3A – 3D). This

was primarily due to a lesser reduction in fat mass in liraglutide-treated CMV-Ser^{791}Ala Raptor

knock-in compared to WT mice (Figs. 3E- F). There was no significant difference between

genotypes in the ability of liraglutide to reduce lean and free fluid mass (Figs. 3G – 3J).

Metabolic cage experiments demonstrate that food intake was equally suppressed in WT and

CMV-Ser^{791}Ala Raptor knock-in mice during the first day of liraglutide treatment. However, food

intake returned to pre-liraglutide treatment levels in CMV-Ser^{791}Ala Raptor knock-in mice by the

second day of treatment whereas they did not return to pre-liraglutide treatment levels in WT

mice until the fifth day of dosing (Figs. 3K – 3L). Energy expenditure (EE) decreased in

response to liraglutide in both genotypes and remained significantly lower in CMV-Ser^{791}Ala

Raptor knock-in mice compared to their vehicle counterparts for up to eight days of dosing

(Figs. 3M – 3N). In WT mice, EE only remained lower in liraglutide-treated mice for three days

(Figs. 3M – 3N). These data suggest that phosphorylation of Raptor at Ser^{791} is required for the

full body weight-lowering effect of liraglutide via regulation of both food intake and EE.

There was no difference between genotypes in fasting blood glucose prior to liraglutide

treatment (Supp. Fig. 2A). Two weeks of liraglutide treatment lowered fasting glucose to similar

degrees in both genotypes, and there was no difference in fasting plasma insulin levels (Supp

Figs. 2B – 2C).

Discussion

GLP-1R-stimulated mTORC1 signaling in the VMH reduces food intake[3]. This is

consistent with studies showing that activation of hypothalamic mTORC1 reduces food intake

and mediates weight-lowering by leptin[10-12]. We show that PKA phosphorylates the mTORC1-

regulatory protein Raptor at Ser^{791} in response to liraglutide and that this is responsible for

~50% of liraglutide-induced weight loss. Given the complexity of signaling mechanisms

downstream of the GLP-1R[13], assigning specific pathways to a desirable phenotype could be

leveraged towards more effective weight loss therapeutics.

To our knowledge, our study is the first to describe a PKA-mTORC1 interaction and its

physiological relevance in the context of GLP-1R signaling. PKA is reported to either stimulate[7]

or inhibit[14] mTORC1 activity. Similar to our study, Liu *et al.*[7] show PKA-mediated stimulation of

mTORC1 in cells incubated in serum-free media. This contrasts inhibition of mTORC1 in cells

cultured in serum-supplemented media reported by Jewell *et al.*[14]. Since mTORC1 is a cellular

energy sensor, PKA may exert differential actions on mTORC1 depending on nutrient

availability or cellular signaling and energy status, as was also shown previously by Liu and

colleagues[7].

One limitation of our *in vivo* studies is that PKA-resistant Raptor is expressed in all

tissues. The brain is a key target for the weight-lowering effects of GLP-1R agonists[15-17].

Regulation of energy balance by mTORC1 appears to be most critical in hypothalamic regions

such as the arcuate (ARC), paraventricular hypothalamus (PVH), VMH, and suprachiasmatic

nucleus[12,18,19], where the GLP-1R is highly expressed[20,21]. The ARC mediates weight loss

effects of liraglutide[22] and is engaged by systemic liraglutide[23]. Hindbrain regions are also

engaged by systemic liraglutide and mediate its weight-lowering effects[23,24]. Inhibition of PKA

attenuates the anorectic effect of hindbrain-targeted Ex4 in rats[25]. Future studies expressing

Ser[791]Ala Raptor in a brain region/cell-type specific manner will enable identification of neuronal

population(s) where a PKA-mTORC1 interaction mediates the anorectic effect of GLP-1R

agonists.

Our results suggest that liraglutide does not regulate mTORC1 via Akt. Inhibition of Akt

activity in the hindbrain[26] or hypothalamus[27] attenuates the anorectic effect of Ex4, so identifying

targets of Akt mediating the effects of GLP-1R agonists requires further study. AMP-activated

protein kinase (AMPK) is also implicated in the metabolic actions of GLP-1R agonists. Studies

from our lab and others show that GLP-1R agonists reduce food intake by inhibiting

AMPK[3,25,28,29]. Since AMPK suppresses mTORC1 by phosphorylating TSC2[30] or Raptor[31], GLP-

1R activation could stimulate mTORC1 by reducing AMPK activity. AMPK can also be inhibited

by mTORC1 signaling[32], so GLP-1R-mediated inhibition of AMPK may be downstream of

mTORC1.

Given the variability in response and adverse effects (e.g. nausea) of current GLP-1

based drugs, the identification of therapeutically relevant signaling profiles downstream of the

GLP-1R holds promise for improving efficacy and reducing side effects of these drugs. Our

studies provide a framework for future investigations into the significance of the PKA-mTOR

interaction to other beneficial metabolic effects of GLP-1R agonists.

Acknowledgements

We thank Wei Zhang for excellent technical assistance and mouse colony management. This

work was supported by NIH R01 DK097361 (JEA) and NIH R01 DK116625 (SC).

References

Drucker, D. J. GLP-1 physiology informs the pharmacotherapy of obesity. *Mol Metab* **57**, 101351, (2022),PMC8859548.

Brown, J. D., McAnally, D., Ayala, J. E., Burmeister, M. A., Morfa, C., Smith, L. & Ayala, J. E. Oleoylethanolamide modulates glucagon-like peptide-1 receptor agonist signaling and enhances exendin-4-mediated weight loss in obese mice. *Am J Physiol Regul Integr Comp Physiol* **315**, R595-R608, (2018),PMC6230892.

Burmeister, M. A., Brown, J. D., Ayala, J. E., Stoffers, D. A., Sandoval, D. A., Seeley, R. J. & Ayala, J. E. The Glucagon-Like Peptide-1 Receptor in the Ventromedial Hypothalamus Reduces Short-Term Food Intake in Male Mice by Regulating Nutrient Sensor Activity. *Am J Physiol Endocrinol Metab*, ajpendo 00113 02017, (2017)

Saxton, R. A. & Sabatini, D. M. mTOR Signaling in Growth, Metabolism, and Disease. *Cell* **168**, 960-976, (2017),PMC5394987.

Kim, D. H., Sarbassov, D. D., Ali, S. M., King, J. E., Latek, R. R., Erdjument-Bromage, H., Tempst, P. & Sabatini, D. M. mTOR interacts with raptor to form a nutrient-sensitive complex that signals to the cell growth machinery. *Cell* **110**, 163-175, (2002)

Thorens, B. Expression cloning of the pancreatic beta cell receptor for the gluco-incretin hormone glucagon-like peptide 1. *Proc Natl Acad Sci U S A* **89**, 8641-8645, (1992)

Liu, D., Bordicchia, M., Zhang, C., Fang, H., Wei, W., Li, J. L., Guilherme, A., Guntur, K., Czech, M. P. & Collins, S. Activation of mTORC1 is essential for beta-adrenergic stimulation of adipose browning. *J Clin Invest* **126**, 1704-1716, (2016),PMC4855937.

Efrat, S., Linde, S., Kofod, H., Spector, D., Delannoy, M., Grant, S., Hanahan, D. & Baekkeskov, S. Beta-cell lines derived from transgenic mice expressing a hybrid insulin gene-oncogene. *Proc Natl Acad Sci U S A* **85**, 9037-9041, (1988),PMC282658.

9   Davies, S. P., Reddy, H., Caivano, M. & Cohen, P. Specificity and mechanism of action
of some commonly used protein kinase inhibitors. *Biochem J* **351**, 95-105,
(2000),PMC1221339.

10   Blouet, C., Ono, H. & Schwartz, G. J. Mediobasal hypothalamic p70 S6 kinase 1
modulates the control of energy homeostasis. *Cell Metab* **8**, 459-467,
(2008),PMC2637401.

11   Cota, D., Matter, E. K., Woods, S. C. & Seeley, R. J. The role of hypothalamic
mammalian target of rapamycin complex 1 signaling in diet-induced obesity. *J Neurosci*
28, 7202-7208, (2008),PMC2597379.

12   Cota, D., Proulx, K., Smith, K. A., Kozma, S. C., Thomas, G., Woods, S. C. & Seeley, R.
J. Hypothalamic mTOR signaling regulates food intake. *Science* **312**, 927-930, (2006)

13   Fletcher, M. M., Halls, M. L., Christopoulos, A., Sexton, P. M. & Wootten, D. The
complexity of signalling mediated by the glucagon-like peptide-1 receptor. *Biochem Soc*
Trans **44**, 582-588, (2016)

14   Jewell, J. L., Fu, V., Hong, A. W., Yu, F. X., Meng, D., Melick, C. H., Wang, H., Lam, W.
M., Yuan, H. X., Taylor, S. S. & Guan, K. L. GPCR signaling inhibits mTORC1 via PKA
phosphorylation of Raptor. *Elife* **8**, (2019),PMC6529218.

15   Adams, J. M., Pei, H., Sandoval, D. A., Seeley, R. J., Chang, R. B., Liberles, S. D. &
Olson, D. P. Liraglutide Modulates Appetite and Body Weight Through Glucagon-Like
Peptide 1 Receptor-Expressing Glutamatergic Neurons. *Diabetes* **67**, 1538-1548,
(2018),PMC6054439.

16   Sisley, S., Gutierrez-Aguilar, R., Scott, M., D'Alessio, D. A., Sandoval, D. A. & Seeley, R.
J. Neuronal GLP1R mediates liraglutide's anorectic but not glucose-lowering effect. *J*
Clin Invest **124**, 2456-2463, (2014),4038572.

17 Varin, E. M., Mulvihill, E. E., Baggio, L. L., Koehler, J. A., Cao, X., Seeley, R. J. &
Drucker, D. J. Distinct Neural Sites of GLP-1R Expression Mediate Physiological versus
Pharmacological Control of Incretin Action. *Cell Rep* **27**, 3371-3384 e3373, (2019)

18 Inhoff, T., Stengel, A., Peter, L., Goebel, M., Tache, Y., Bannert, N., Wiedenmann, B.,
Klapp, B. F., Monnikes, H. & Kobelt, P. Novel insight in distribution of nesfatin-1 and
phospho-mTOR in the arcuate nucleus of the hypothalamus of rats. *Peptides* **31**, 257-
262, (2010),PMC4043136.

19 Proulx, K., Cota, D., Woods, S. C. & Seeley, R. J. Fatty acid synthase inhibitors
modulate energy balance via mammalian target of rapamycin complex 1 signaling in the
central nervous system. *Diabetes* **57**, 3231-3238, (2008),PMC2584128.

20 Jensen, C. B., Pyke, C., Rasch, M. G., Dahl, A. B., Knudsen, L. B. & Secher, A.
Characterization of the Glucagonlike Peptide-1 Receptor in Male Mouse Brain Using a
Novel Antibody and In Situ Hybridization. *Endocrinology* **159**, 665-675, (2018)

21 Cork, S. C., Richards, J. E., Holt, M. K., Gribble, F. M., Reimann, F. & Trapp, S.
Distribution and characterisation of Glucagon-like peptide-1 receptor expressing cells in
the mouse brain. *Mol Metab* **4**, 718-731, (2015),PMC4588458.

22 Secher, A., Jelsing, J., Baquero, A. F., Hecksher-Sorensen, J., Cowley, M. A., Dalboge,
L. S., Hansen, G., Grove, K. L., Pyke, C., Raun, K., Schaffer, L., Tang-Christensen, M.,
Verma, S., Witgen, B. M., Vrang, N. & Bjerre Knudsen, L. The arcuate nucleus mediates
GLP-1 receptor agonist liraglutide-dependent weight loss. *J Clin Invest* **124**, 4473-4488,
(2014),PMC4215190.

23 Salinas, C. B. G., Lu, T. T., Gabery, S., Marstal, K., Alanentalo, T., Mercer, A. J.,
Cornea, A., Conradsen, K., Hecksher-Sorensen, J., Dahl, A. B., Knudsen, L. B. &
Secher, A. Integrated Brain Atlas for Unbiased Mapping of Nervous System Effects
Following Liraglutide Treatment. *Sci Rep* **8**, 10310, (2018),PMC6037685.

24   Fortin, S. M., Lipsky, R. K., Lhamo, R., Chen, J., Kim, E., Borner, T., Schmidt, H. D. &

Hayes, M. R. GABA neurons in the nucleus tractus solitarius express GLP-1 receptors

and mediate anorectic effects of liraglutide in rats. *Sci Transl Med* **12**,

(2020),PMC7211411.

25   Hayes, M. R., Leichner, T. M., Zhao, S., Lee, G. S., Chowansky, A., Zimmer, D., De

Jonghe, B. C., Kanoski, S. E., Grill, H. J. & Bence, K. K. Intracellular signals mediating

the food intake-suppressive effects of hindbrain glucagon-like peptide-1 receptor

activation. *Cell metabolism* **13**, 320-330, (2011),3108145.

26   Rupprecht, L. E., Mietlicki-Baase, E. G., Zimmer, D. J., McGrath, L. E., Olivos, D. R. &

Hayes, M. R. Hindbrain GLP-1 receptor-mediated suppression of food intake requires a

PI3K-dependent decrease in phosphorylation of membrane-bound Akt. *Am J Physiol*

Endocrinol Metab **305**, E751-759, (2013),PMC3761195.

27   Yang, Y., Choi, P. P., Smith, W. W., Xu, W., Ma, D., Cordner, Z. A., Liang, N. C. &

Moran, T. H. Exendin-4 reduces food intake via the PI3K/AKT signaling pathway in the

hypothalamus. *Sci Rep* **7**, 6936, (2017),PMC5537284.

28   Burmeister, M. A., Ayala, J., Drucker, D. J. & Ayala, J. E. Central glucagon-like peptide 1

receptor-induced anorexia requires glucose metabolism-mediated suppression of AMPK

and is impaired by central fructose. *American journal of physiology. Endocrinology and*

metabolism **304**, E677-685, (2013)

29   Sandoval, D., Barrera, J. G., Stefater, M. A., Sisley, S., Woods, S. C., D'Alessio, D. D. &

Seeley, R. J. The anorectic effect of GLP-1 in rats is nutrient dependent. *PLoS One* **7**,

e51870, (2012),3524167.

30   Inoki, K., Zhu, T. & Guan, K. L. TSC2 mediates cellular energy response to control cell

growth and survival. *Cell* **115**, 577-590, (2003)

31   Gwinn, D. M., Shackelford, D. B., Egan, D. F., Mihaylova, M. M., Mery, A., Vasquez, D.

S., Turk, B. E. & Shaw, R. J. AMPK phosphorylation of raptor mediates a metabolic

checkpoint. *Mol Cell* **30**, 214-226, (2008),PMC2674027.

32   Dagon, Y., Hur, E., Zheng, B., Wellenstein, K., Cantley, L. C. & Kahn, B. B. p70S6

Kinase Phosphorylates AMPK on Serine 491 to Mediate Leptin's Effect on Food Intake.

Cell metabolism, (2012)

Fig 1. The GLP-1R agonist liraglutide increases mTOR activity in a PKA- dependent

manner (Figure 1-Source data 1). (A) Immunoblot of pS6, S6, pCREB, and CREB and

quantification of pS6 expression (pS6 / S6) in CHO-K1 cells transfected with the a vector

expressing the *hGLP1R* and treated with DMSO or the mTORC1 inhibitor rapamycin (100 nM)

for 30 min followed by treatment with 1X PBS (Veh) or liraglutide (Lira, 10 nM) for 1 hr. (B-C)

Immunoblot of pS6, S6, pCREB, and CREB and quantification of pS6 expression (pS6/S6) in

CHO-K1 cells stably expressing the *hGLP1R* treated with DMSO (Veh) or either of the PKA

inhibitors, H89 (10 µM) (B) or KT5720 (10 µM or 50 µM) (C), for 30 min followed by treatment

with vehicle (PBS), Lira (10 nM), forskolin (Fsk, 10 µM), or insulin (Ins, 10 mg/mL) for 1 hr.

Quantification of pCREB expression (pCREB/CREB) is shown in Supp. Figs. 2B and 2C. (D)

Immunoblot of pS6, S6, pAkt, and Akt and quantification of pS6 expression (pS6/S6) in CHO-K1

cells stably expressing the *hGLP1R* treated with DMSO or the pan-Akt inhibitor MK-2206 (5 µM

or 20 µM) for 30 min followed by treatment with vehicle (PBS), Lira (10 nM), Fsk (10 µM), or Ins

(10 mg/mL) for 1 hr. Quantification of pAkt expression (pAkt/Akt) is shown in Supp. Fig. 2D.

Analysis was done using one-way ANOVA followed by Holm-Sidak's multiple comparisons.

Data are normalized to the respective control (Veh or Veh and inhibitor). Absolute data are

shown in Supp. Fig. 2. All data are shown as mean ± SEM, ns not significant, * $p < 0.05$, ** $p <$

0.01, *** $p < 0.001$, **** $p < 0.0001$, n = 3-4 biological replicates.

Fig 2. PKA phosphorylates Raptor at S791 upon GLP-1R activation (Figure 2-Source data

1). (A) Immunoblot and quantification of phospho-PKA substrate (pRRXS*/T*) and Myc in CHO-

K1 cells stably expressing the *hGLP1R* transfected with Myc-wild-type (WT) Raptor or Myc-

Ser^{791}Ala Raptor and treated with vehicle (DMSO) or H89 (10 µM) for 30 min followed by

treatment with Lira (10 nM), forskolin (Fsk, 10 µM), or insulin (Ins, 10 mg/mL) for 1 hr. (B)

Immunoblot and quantification of phospho-PKA substrate and Myc in CHO-K1 cells stably

expressing the *hGLP1R* transfected with Myc-wild-type (WT) Raptor and treated with vehicle (DMSO) or MK-2206 (20 µM) for 30 min followed by treatment with Lira (10 nM), Fsk (10 µM), or Ins (10 mg/mL) for 1 hr. (C) Immunoblot and quantification of phospho-PKA substrate and Myc in β-TC3 cells transfected with Myc-wild-type (WT) Raptor or Myc-Ser[791]Ala Raptor and treated with vehicle (DMSO) or MK-2206 (10 µM) for 30 min followed by treatment with Lira (10 nM), forskolin (Fsk, 10 µM), or insulin (Ins, 10 mg/mL) for 1 hr. Analysis was done using one-way ANOVA followed by Holm-Sidak's multiple comparisons. Data are shown as mean ± SEM, no symbol = not significant, * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, n = 3-4 biological replicates.

Fig. 3. PKA phosphorylation of Raptor at S791 facilitates the weight-lowering effects of liraglutide (Figure 3-Source data 1). Body weight, body composition, and energy balance parameters in male wild-type (WT) and CMV-Ser[791]Ala Raptor knock-in mice given vehicle (0.9% saline) or liraglutide (200 µg/kg) subcutaneously twice a day for 14 days. Absolute body weight (A), total mass (C), fat mass (E), lean mass (G), and free fluid mass (I). Change in body weight (B), total mass (D), fat mass (F), lean mass (H) and free fluid mass (J) from baseline (Day 0). Absolute daily food intake (K) and EE (M) and change in daily food intake (L) and EE (N) relative to baseline (Day 0) during a 10-day treatment period in metabolic cages. Analysis was done using mixed-effects analysis followed by Holm-Sidak's multiple comparisons between liraglutide-treated WT and CMV-Ser[791]Ala Raptor mice. Data are shown as mean ± SEM, ns not significant, * $p < 0.05$, ** $p < 0.01$, **** $p < 0.0001$, n = 13-14 (A-D), 9-12 (E-J), or 4 (K-N) mice per group.

Supp. Fig. 1. Generation of CMV-Ser^{791}Ala Raptor knock-in mice. (Supplementary Figure

1-Source data 1). (A) Targeting vector used by Ingenious Targeting Laboratory, Inc. to

introduce a *Cre*-inducible two base pair point mutation in an inverted exon 20 of the Raptor

coding sequence. (B) Verification of correct target sequence expression by genomic DNA

digestion. Wild-type (Control) DNA contains a SacI sequence that is absent in the Ser^{791}Ala

point mutant (red). SacI digestion of a subcloned 911 base pair (bp) genomic DNA fragment

obtained from liver tissue of control mice produces two products at 371bp and 540bp. The

corresponding 989bp genomic DNA sequence obtained from CMV-Ser^{791}Ala Raptor knock-in

mice cannot be digested by SacI.

Supp. Fig. 2. Absolute pS6, pCREB, and pAkt expression in CHO-K1 cells stably

expressing the *hGLP1R* and treated with liraglutide. (Supplementary Figure 2-Source

data 1). (A) Quantification of pS6 expression (pS6 / S6) in CHO-K1 cells transfected with the a

vector expressing the *hGLP1R* and treated with DMSO or the mTORC1 inhibitor rapamycin

(100 nM) for 30 min followed by treatment with 1X PBS (Veh) or liraglutide (Lira, 10 nM) for 1 hr.

(B-C) Quantification of pS6 (pS6/S6) and pCREB (pCREB/CREB) expression in CHO-K1 cells

stably expressing the *hGLP1R* treated with DMSO (Veh) or either of the PKA inhibitors, H89 (10

µM) (B) or KT5720 (10 µM or 50 µM) (C), for 30 min followed by treatment with vehicle (PBS) or

Lira (10 nM) for 1 hr. Quantification of pS6 and pAkt (pAkt/Akt) in CHO-K1 cells stably

expressing the *hGLP1R* treated with DMSO or the pan-Akt inhibitor MK-2206 (5 µM or 20 µM)

for 30 min followed by treatment with vehicle (PBS), Lira (10 nM) or Ins (10 mg/mL) for 1 hr.

Analysis was done using one-way ANOVA followed by Holm-Sidak's multiple comparisons.

Data are presented as absolute densitometry values and are shown as mean ± SEM, ns not

significant, * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$, n = 3-4 biological replicates.

Supp. Fig. 3. Blood glucose prior to drug treatment and blood glucose and insulin post-

treatment in male wild-type (WT) and Ser^{791}Ala Raptor knock-in mice (Supplementary

Figure 3-Source data 1). (A-D) Total (A), fat (B), lean (C), and free body fluid (D)

measurements in WT and CMV-Ser^{791}Ala Raptor mice prior to vehicle or liraglutide

administration. (E-F) 5-hr fasting blood glucose in male WT and CMV-Ser^{791}Ala Raptor mice

before drug treatment (E) and on treatment day 14 (F). (G) 5-hr fasting insulin in male WT and

CMV-Ser^{791}Ala Raptor mice before on treatment day 14. Analysis was done using one-way

ANOVA followed by Holm-Sidak's multiple comparisons. Data are shown as mean ± SEM, ns

not significant, n = 4-8 mice per group.

Figure 1

Figure 2

Figure 3

Figure 3 (contd.)

436
437

Supplementary Figure 1

A

B

Supplementary Figure 2

Supplementary Figure 3

440