1 **Title: Why the architecture of environmental fluctuation matters for fitness**

3 Authors: John S. Park[1*], Anja Felmy[1,2]

4 [1]Department of Biology, University of Oxford; *john.park@biology.ox.ac.uk*

5 [2]Department of Biology, Lund University; *anja.felmy@biol.lu.se*

7 *Corresponding author

Number of words in abstract: 269

Number of words in main text: 5723

Number of figures: 4

Number of tables: 1

Number of text boxes: 0

Abstract

The physical environment provides the very stage upon which the eco-evolutionary play unfolds. How fluctuations in the environment affect demographic fitness is thus central to selection predictions, life history analyses, and viability of populations. Treatment of fluctuating environments typically leverages the mathematics of random variability. However, environmental fluctuations in nature are almost always combinations of random and non-random components. For example, some fluctuations contain feedbacks which generate autocorrelation (*e.g.* disturbances such as floods, fires, and hurricanes), while others are driven by geophysical forces that create fixed cyclicality (*e.g.* seasonal, tidal, and diel). Despite theoretical developments, the consideration of non-random characteristics of fluctuations is still rare in empirical work on natural populations, mostly due to convention and partially due to difficulties in measuring and analyzing timeseries of environmental fluctuations. We show why non-randomness matters for fitness. Using a simple demographic model, we systematically compare four major categories of fluctuating environments: stochastic, positively autoregressive, negatively autoregressive, and periodic with error ("Noisy Clock"). The architectures of fluctuations influence the fitness of structured populations even when the modelled environments only differ in the timing of fluctuations, and not in their overall frequency. Importantly, we highlight two quantitative mechanisms through which fitness depends on fluctuation architecture—the consecutiveness of deviations from the environmental mean, and Jensen's Inequality acting on nonlinear biological parameters—both relevant features in virtually all populations inhabiting variable environments. Our goal is to argue that non-random structures of environmental variability should be more seriously considered in empirical work. Such an

37 endeavor would tap into the rich diversity of variable environments in nature to expand our

38 understanding of the commensurate diversity of population dynamics.

39 *Keywords:* fluctuating environments, life history, evolutionary demography, stochastic fitness,

40 nonlinearity, Jensen's Inequality

Introduction

Estimating population growth—or fitness—is a centerpiece of life history theory, evolutionary demography, and conservation biology. Reliable estimates of fitness have only become more important as climate change appears to increase variability in many systems, rendering the demographic consequences more varied and complex (Pike et al. 2004, Boyce et al. 2006, Davison et al. 2019). The potential for environmental variability to profoundly affect fitness outcomes is well-known. Yet, why and how types of environmental variability differ in their impact on fitness is less clear, especially in empirical work. Here we provide a synthesis of why and how the structure of environmental variability influences fitness. We first outline the history of conceptual developments, and then systematically compare the fitness consequences in four major categories, or 'architectures', of environmental variability. Lastly, we highlight two specific mechanisms that underlie fluctuations' influence on fitness. We aim to spur a more careful consideration of the architecture of environmental variability in future empirical work.

There is a wealth of sophisticated theory on environmental impacts on demography and fitness. Cole (1954) first conceived of the fitness of life history strategies as directly linked to the strategies' influence on the population growth rate of asexually reproducing organisms. Variations among genotypes in the age distribution of survival and reproduction shape the fitness landscapes of life histories (Gadgil and Bossert 1970, Stearns 1976), upon which natural selection can act. Evolutionary demography (Keyfitz 1968, Caswell 2001) leverages these concepts to analyze how the trajectories and sensitivities of population growth are influenced by population structures (*e.g.* compositions of ages, stages, or sizes) consisting of individuals on variable courses of life histories.

Early theoretical work such as above began with assumptions of constant environments for

simplicity, and because it was necessary to lay the groundwork for comparing species in a

systematic manner (Fisher 1958, MacArthur 1962). Developments since (*e.g.* Levins 1968,

Schaffer 1974) have largely focused on extending analyses to include stochastic environment

assumptions to reflect the complexities of nature (Lande et al. 2003, Tuljapurkar et al. 2003), and

to spotlight the increasing environmental variability under climate change (Easterling et al. 2000,

Drake 2005, Boyce et al. 2006, Morris et al. 2008, Lawson et al. 2015). Tuljapurkar *et al.* (2009)

demonstrated why the expected optimal life histories differ between constant and stochastic

environments, by decomposing the effects of life history speed, dispersion, and temporal

correlation on fitness. Over time, 'constant' and 'stochastic' settled as the two dominant

archetypical descriptions of environments, with the latter becoming the modern standard.

Stochasticity indeed offers a powerful suite of tools to deal with many kinds of variable

environments, afforded by mature statistical theories of probability distributions. In short, a

variable environment can typically be summarized by the first two moments of a distribution, *i.e.*

the mean and variance, whose derivations lead to tractable analytical solutions for biological

predictions in that environment (Lawson et al. 2015), *e.g.* bet-hedging strategies (Cohen 1966,

Childs et al. 2010, Gremer and Venable 2014). Higher-order moments of stochastic

environments can be used to analyze the effects of, *e.g.*, intra- and inter-annual variability and

temporal autocorrelation on fitness (Tuljapurkar et al. 2009). There is substantial theoretical

evidence that the 'color' of noise (negative to zero to positive autocorrelation) should influence

fitness and extinction risk (Cohen 1966, Lande and Orzack 1988, Vasseur and Yodzis 2004,

Halley 2005, Tuljapurkar and Haridas 2006, Reuman et al. 2008, Engen et al. 2013, Chevin et al.

2017). Overall, the subfield of 'stochastic demography' uses stochastic environmental

formulation at its core to ask ecological and evolutionary questions (Tuljapurkar 1990, Fieberg

and Ellner 2001, Engen et al. 2009, Lande et al. 2017, Chevin et al. 2017, Schreiber and Moore

2018, Crewe et al. 2018, Davison et al. 2019).

Mathematically, stochasticity is specific: it describes a process that can be captured with a

random probability distribution. However, many real categories of environmental fluctuations

have non-random components. For example, fundamental geophysical fixtures such as the

rotation of Earth (governing diel cycles), its axial tilt and revolution around the Sun (seasonal

cycles), and the revolution of the Moon around Earth (tidal cycles) set periodic oscillations in

most terrestrial and aquatic ecosystems. Analogous to driven harmonic oscillators in physics,

such environments are governed by fixed (at least on ecological and evolutionary time scales)

external drivers of fluctuations. Distinct from stochastic variability, geophysically-driven

fluctuations are thus not emergent products of the internal ecological and evolutionary dynamics

of the system such as in population cycles or ecological succession.

The particular sequence of environmental fluctuations is biologically important, over and above

the overall amount of environmental variability. Stearns (1976, Fig. 7) outlined the possible

outcomes of environmental fluctuation frequency and predictability for the age distribution of

reproduction in a population. Later, Tuljapurkar (1985) analyzed how the relative scaling

between the period of environmental cycles and generation time influences population growth

rate. Recent work demonstrated that environmental periodicity may itself be a selective force that

determines both optimal (Park 2019) and individual variation in life histories (Park and Wootton

2021).

Empirically speaking, climate change is rapidly perturbing seasonality and fluctuation regimes,

with far-reaching consequences such as phenological shifts (Walther et al. 2002, Westerling et al.

2006, Marlon et al. 2009). Importantly, the mean and variance of environmental fluctuations can

change independently in disturbance regimes (Lytle 2001, Lytle et al. 2008, Denny et al. 2009),

which is a growing focus in climate change biology (Jentsch et al. 2007, IPCC 2012, Thornton et

al. 2014). Thus, teasing apart the non-random and random components that constitute

environmental variability—instead of collapsing all environmental variability to stochasticity—is

crucial not only for more accurate predictions of population growth (Tuljapurkar and Haridas

2006), but for systematizing the variety of causal drivers of population dynamics and life history

selection.

We focus on four broadly representative architectures of fluctuating environments and use a

simple demographic model to show the architectures' contrasting effects on demographic fitness.

The four environments solely differ in the exact timing sequence of events. Events here represent

ecological disturbances (Lytle 2001) such as fires, floods, hurricanes, avalanches, droughts, and

heatwaves, or, in seasonal systems, onsets of winter that can largely impede biological activity

(Inouye 2008). A particularly novel consideration is the fourth category of fluctuations:

inherently periodic environments such as seasonal or tidal ones. These externally driven cycles

with fixed periodicity are fundamentally different in their memory process compared to

internally produced cycles (*e.g.* autocorrelated): in the former, environmental states gravitate to

an external schedule, whereas in the latter, environments depend on former states of themselves.

We then proceed to identify two general quantitative mechanisms that cause divergences in

fitness estimates among all four categories. The first mechanism involves the pattern of

consecutive deviations from the environmental mean. The second involves Jensen's Inequality

acting on nonlinear life history traits, or vital rates; the prevalence of such nonlinearity in nature

is likely universal. Both are very general properties of populations in variable environments. We

do not claim that these are the only mechanisms at play, nor that our list of fluctuation categories

is exhaustive. We also do not generate new theory. Instead, our goal is to show how the two

mechanisms influence fitness even in simple situations, and to systematically highlight the role

of non-random variability beyond the conventional archetype of random variability. Finally, we

discuss the general relevance of our findings and identify key extensions and empirical tests that

will be important moving forward.

Four categories of fluctuating environments

We start by setting up illustrative examples of four major categories of fluctuating environments

(Fig. 1): stochastic (white noise), positively autoregressive (red noise), negatively autoregressive

(blue noise), and "Noisy Clock" (fundamentally periodic, but with error). For all four categories,

we focus on the parameter τ_i which is the *interval* between events. Again, these events represent

ecological disturbances such as fires, or the onset of winter in seasonal systems with the

assumption that winters are harsh enough to perturb or halt biological activity. In seasonal

systems, note that celestial seasons are perfectly periodic, but the meteorological onset of winter

(*e.g.* temperature, frost) and arrival of spring are variable, making the intervening warm season

(the interval) also variable in length ('climatological growing seasons', *sensu* Linderholm 2006).

We derive the mean and variance of the distribution of intervals between events from each

category to make them exactly analytically equivalent, before comparing the fitness

consequences in environments characterized by the different fluctuations.

Figure 1. Four categories of fluctuating environments. Each fluctuation architecture is simulated by generating a sequence of event intervals on a timeline. Events, denoted by the tick marks, may represent ecological disturbances or winters that bring harsh conditions for biological activity. A) A stochastic environment is generated by drawing a sequence of intervals from a random (Normal) probability distribution. B) A positively autoregressive environment is generated with positive linear dependency of subsequent intervals on immediately preceding ones (an AR(1) process), with random error; C) the negatively autoregressive environment is only different from B in that the linear dependency on the past interval is negative. D) A Noisy Clock environment assumes a series of unseen "due times" that are perfectly periodic (dashed blue tick marks), but the actual events occur as a result of error around those due times (blue shaded regions); the gaps between realized events form the sequence of realized intervals.

Category I: Stochastic (random), without feedback

166 We generate a fluctuating environment with a random sequence of intervals   between

167 environmental disturbance events:

$$\tau_i \sim Normal(\mu, \sigma^2) \tag{1}$$

where μ is the desired interval mean and σ^2 the variance. We do this instead of generating a

series of events via a Poisson process, which is a more common way of generating random

intervals on a timeline, to make this case comparable with the other three cases with respect to

their error distributions and because below we will manipulate the amount of noise (σ^2) while

holding the mean interval constant (which is not possible with Poisson distributions because their

mean and variance are equal). While using a normal distribution does not theoretically ensure

negative values cannot be drawn—which would be nonsensical for interval length—for

simplicity we have set the mean (=100) and variances (ranging from 0 to 300) of our simulations

such that the probability of a negative value is exceedingly low ($< 3.9 \times 10^{-9}$); we then

confirmed that negative values were in fact not drawn in our simulations.

Importantly, note that this manner of generating a fluctuating environment does not consider the

source or generative nature of the fluctuations, nor the length of preceding intervals (*i.e.* there is

no memory of past fluctuations), but only assumes the long-run statistical distribution of

variability (Fig. 1A). While measured environmental variability may in reality always have a

physical source, and all systems may have some memory process to varying degrees, some

systems are so complex and noisy that they are well approximated by pure randomness. Random

environments may be experienced, for example, by mice living in dryland ecosystems with

variable resource availability (Noble et al. 2019), by weevil species using strongly fluctuating

oak acorns as egg-laying sites (Pélisson et al. 2012), and by annual herbs growing in or near

vernal pools created by stochastic winter precipitation events (Emery and La Rosa 2019).

Category II: Positively autoregressive

We generate a positively autocorrelated environment with a simple AR(1) process:

$$\tau_i = \mu + \varphi\tau_{i-1} + \varepsilon_i \tag{2}$$

where μ is the imposed mean interval between environmental disturbances, and the interval τ_i is

positively correlated by strength of autocorrelation φ with the interval immediately preceding it

(τ_{i-1}). The i.i.d. (independent identically distributed) noise ε_i is drawn from $N(0, \sigma_\varepsilon^2)$. Positively

autocorrelated environments are characterized by a 'compounding' behavior where a longer or

shorter interval than the mean is likely to beget another longer or shorter interval in the

immediate future, respectively (Fig. 1B). Examples of positively autocorrelated fluctuations in

nature include some fire regimes (Caswell and Kaye 2001), oceanic temperatures (Steele 1985,

Halley 2005), and flood regimes (Sabo and Post 2008).

Category III: Negatively autoregressive

The negative counterpart to the positively autoregressive environment is one with a negative

autoregressive term $(-\varphi)$ and all other terms identical to Eq. 2:

$$\tau_i = \mu - \varphi\tau_{i-1} + \varepsilon_i \tag{3}$$

Negatively autocorrelated environments are characterized by a 'compensating' or self-correcting

behavior where an interval between disturbances that is longer or shorter than the mean is likely

to be counterbalanced with a shorter or longer interval in the immediate future, respectively (Fig.

1C). Real examples of negatively autocorrelated environments in nature, though much more

rarely considered than positively autocorrelated ones (Metcalf and Koons 2007), include

terrestrial plant systems with delayed density-dependence driven by nutrient cycling

(Gonzalez☐Andujar et al. 2006), and tree masting (Hacket-Pain and Bogdziewicz 2021).

Category IV: 'Noisy Clock'

The last and least explicitly considered category of fluctuations, but one that is ubiquitous in

nature, includes environments with a fundamental external driver of cyclicality, whose

deterministic periodicity is not affected by the realized noise. We henceforth refer to this case as

the 'Noisy Clock' to describe the clockwork (deterministic) nature of fluctuations that underlies

the noise. Here, we first punctuate the timeline with a series of deterministic 'due' times of

events much like the celestial periodicity of Earth's revolution that determines the boundaries of

the annual cycle. Intervals between disturbances are not generated by counting time from the

preceding disturbance events, but instead, imperfectly timed disturbances events occur around

each deterministic due time with i.i.d. noise (Fig. 1D). Then the intervals are the gaps between

subsequent realized events:

$$\tau_i = [\mu(i + 1) + \varepsilon_{i+1}] - [\mu i + \varepsilon_i] \tag{4}$$

where $\varepsilon_i \sim N(0, \sigma_\varepsilon^2)$ and $i \in \mathbb{Z}_{>0}$, meaning clocked due times are positive integer multiples of μ,

the same mean interval as in categories I, II, and III. Counting time from preceding events

(which is the case for categories I, II, and III) as opposed to in reference to an external clock

would be an unrealistic way to model fundamentally cyclical systems such as seasonal ones. For

instance, if one were to model a winter event (*e.g.* first snowfall), but stochastically modelled

the length of intervening summer seasons with no regard to due times, then after some additive

sequence of stochastic lengths it would be possible to have winters starting in July when lined

up against a deterministic calendar. This would be unrealistic under any reasonable geophysical

stationarity assumptions. Adding error to due times has the effect of gravitating noisy events

back to realistic due times set by the underlying forces of harmonic oscillations in the

environment. Real fluctuations with clocked due times driven by geophysical forces—but with

error in the measured proxy variables—are exemplified by a myriad of diel (*e.g.* hunting

behavior of African wild dogs; Nouvellet *et al.* 2012), tidal (*e.g.* foraging of intertidal snails;

Hayford *et al.* 2018), and seasonal environments (*e.g.* advancement of egg-laying with warming

springs in numerous bird species; Dunn & Møller 2014). Despite widespread attention to

seasonal environments, particularly with respect to global phenological shifts (Walther et al.

2002), the theoretical foundation of demographic dynamics and life history evolution in cyclical

environments is not yet well understood (Lande et al. 2017).

Comparing apples to apples: the same normal distribution

Probability distributions that generate timeseries can be described by their moments, the first of

which are the man, variance, skewness, and kurtosis. Conventionally, a fluctuating environment

under study is analyzed or modelled using the first two moments – the mean and variance of

event timing or of some continuous parameter (*e.g.* temperature). However, fluctuating

environments often also differ in their higher-order structures such as autocorrelation in the

timing of environmental disturbances and intervals between them. Here, we additionally consider

a structure that is neither purely stochastic nor simply autocorrelated, but one that has a

fundamental deterministic periodicity (Noisy Clock). To compare them systematically, we first

derive input parameters that would make the mean and variance of the four fluctuating

environments (Eqs. 1-4) analytically equivalent (Table 1; Fig. S1). We set mean interval (μ) to

equal 100 time steps, and equivalently vary the variance across the four environments to see its

effect on fitness. Then we explore why even given this equivalency, fitness estimates diverge

among the environments. Our goal is to demonstrate that the conventional assumption that any

fluctuating environment is sufficiently well-characterized as a random environment with mean

and variance of the fluctuation—while justified in some cases—can subtly, but consistently, lead

to inaccurate estimates of fitness.

	Fluctuation	Expected Value ($E[\tau]$)	Variance ($Var[\tau]$)	**Input parameters in terms of μ_s and σ_s^2**
I	$\tau_i \sim Normal(\mu, \sigma^2)$	μ	σ^2	$\mu = \mu_s;$ $\sigma^2 = \sigma_s^2$
II	$\tau_i = \mu + \varphi\tau_{i-1} + \varepsilon_i,$ $where\ \varepsilon_i \sim N(0, \sigma_\varepsilon^2)$	μ	$\dfrac{\sigma_\varepsilon^2}{1 - \varphi^2}$	$\mu = \mu_s;$ $\sigma_\varepsilon^2 = \sigma_s^2(1 - \varphi^2)$
III	$\tau_i = \mu - \varphi\tau_{i-1} + \varepsilon_i,$ $where\ \varepsilon_i \sim N(0, \sigma_\varepsilon^2)$	μ	$\dfrac{\sigma_\varepsilon^2}{1 - \varphi^2}$	$\mu = \mu_s;$ $\sigma_\varepsilon^2 = \sigma_s^2(1 - \varphi^2)$
IV	$\tau_i = [\mu(i + 1) + \varepsilon_{i+1}] - [\mu i + \varepsilon_i],$ $where\ \varepsilon_i \sim N(0, \sigma_\varepsilon^2)\ and\ i \in \mathbb{Z}_{>0}$	μ	$2 \times \sigma_\varepsilon^2$	$\mu = \mu_s;$ $\sigma_\varepsilon^2 = \dfrac{1}{2}\sigma_s^2$

Table 1. Expressions of the expected value and variance of the four fluctuations, and the input
parameters in terms of the desired sample moments (terms with subscript *s*) that will force τ_i's of
all four fluctuations to be equivalently distributed as the normal distribution of category I.

A simple demographic model in fluctuating environments

Upon the environmental fluctuation backgrounds (Fig. 2A), we model a simple population that

proceeds through time. There are 2 time indices: t denotes the discrete time steps, here imagined

as days (but scalable depending on system), and τ_i indexes the interval lengths (measured in time

steps) between subsequent disturbance events, the sequence of which is determined by the

267   categories of fluctuation (Eqs. 1-4). J_t and A_t are numbers of juveniles and adults at time t

268   respectively, f is the per capita rate of reproduction, ℓ is the rate of juveniles maturing into

269   adults, and S_J and S_A are rates of juvenile and adult survival. The population dynamic is then

270   given by the system of equations:

$$
\begin{aligned}
J_{t+1} &= (1-\ell)S_J J_t + fA_t \\
A_{t+1} &= S_A A_t + \ell J_t
\end{aligned} \quad ,
$$

where S_J

$$
= \begin{cases} S_J(\tau_i) = \dfrac{1}{1 + e^{-0.3(\tau_i - \mu)}}, & if \ t \in \left\{ t_0 + \tau_1, t_0 + \tau_1 + \tau_2, t_0 + \displaystyle\sum_{i=1}^{k} \tau_i, \dots \right\}, \\ \qquad\qquad 0.99, & \text{otherwise} \end{cases}
$$

(5)

We focus on juvenile survival S_J as the vital rate that actively interacts with the disturbances that

arrive intermittently into the system. This is because in many natural populations of plants or

animals juvenile mortality is known to be especially sensitive to harsh environmental conditions

such as disturbances, and particularly to how frequently they arrive (Gaillard et al. 1998, 2000,

Higgins et al. 2000, Lytle 2002, McAdam and Boutin 2003, Staver et al. 2009, Park 2019). To

reflect this in Eq. 5, juvenile survival S_J is determined by a logistic response function $S_J(\tau_i)$ (Fig.

2B) when disturbance occurs (*i.e.* when time step t is an element of the predetermined sequence

of disturbance timings, which is the cumulative sum of successive interval lengths $\sum_{i=1}^{k} \tau_i$, where

k is the total number of disturbance intervals simulated), and is 0.99 otherwise (in between

disturbances). The shape parameter 0.3 in the denominator is the steepness of the logistic curve,

the inflection point is at $S_J = 0.5$, and the difference $\tau_i - \mu$ determines how far above or below

$S_J(\tau_i)$ is from mean S_J. Verbally, this means that if two disturbance events happen in quick

succession, juvenile survival will be low, and if a long time passes between events, it will be

high, in a nonlinear fashion. Biologically, this function signifies the time needed for juveniles to

sufficiently develop, grow, or learn to endure the next impending perturbation in the

environment. Though not always logistic, nonlinearity in vital rate response functions in between

perturbations in the environment is likely common (Caswell 2008, Childs et al. 2010, Sletvold et

al. 2013, Taylor et al. 2014, Ehrlén et al. 2016, Shriver 2016). For example, Morris et al. (2006)

showed that, across five species of perennial plants, vital rates including seedling survival

(equivalent to S_J here), survival at other stages, growth, germination, and seed production

(equivalent to f here) changed nonlinearly as a function of time since last hurricane or fire

disturbance. Similarly, Bürkli & Jokela (2017) showed that embryonic development success

(early-stage juvenile survival) of a freshwater snail changed nonlinearly across the species'

reproductive season. The exact shape of vital rate response functions is another key point to be

tailored for system-specific inquiry, whether with empirical measurements or as realistic

assumptions built into models.

The model is otherwise kept simple to focus on how the different categories of environmental

fluctuation affect fitness through just one vital rate. Time-step values for all parameters are

biologically realistic for daily rates (*e.g.* Least Killifish, a small live-bearing freshwater fish

(Felmy et al. 2021)): we hold adult survival S_A constant at 0.99 regardless of disturbance, and

maturation rate ℓ at 0.01 (thus birth-to-birth generation time $\approx$ mean disturbance cycle length,

100). Rate of reproduction f is logistically density-dependent, where $f(N) = 0.1 * (1 - \frac{N}{K})$,

where N is total population size at t and carrying capacity K=1000. Structural complexity such

as the number of developmental stages (*i.e.* the number of equations in the system, or the number

of vital rate terms), or whether multiple vital rates interact with disturbance frequency, would be

an obvious area of customization for system-specific inquiry. The downstream results remain

qualitatively unchanged when f, ℓ, and S_A are allowed to vary within a reasonable range (Fig. S2).

As we will argue, the key mechanisms should influence fitness regardless of demographic

nuance.

Time-averaged fitness

In each of the four architectures of environmental fluctuations, we ask what the long-run fitness

of a life history (the set of vital rate parameters defined above) would be. For fitness, we use the

standard

$$log_e \lambda_s = \lim_{t \to \infty} \left(\frac{1}{t}\right) log_e[N(t)/N(0)] \tag{6}$$

which gives the long-run growth rate in fluctuating environments (Tuljapurkar 1990) where $N(t)$

is the total population size at time *t*, including both juvenile and adult stages. This metric can be

computed from numerical simulations over a long time course of the environment (Caswell

2001), and robustly approximates fitness even under considerably large fluctuations (Lande et al.

2003, Tuljapurkar et al. 2003, 2009). We simulate demographic dynamics (Eq. 5) in each

fluctuating environment (Fig. 1) for 500 event intervals with mean interval (μ) = 100 time steps

($\sim$50,000 time steps), calculate $log_e \lambda_s$, and repeat these calculations over 500 iterations. Lastly,

we compute all of the above across a spectrum of variance in event intervals (0 to 300), to assess

how the amount of fluctuation noise affects the divergence of fitness estimates.

Figure 2. The interaction of environmental fluctuation architecture and nonlinearity in a vital rate creates divergence in fitness estimates. A) Diagrammatic representation of the four architectures of fluctuation, through which the demographic dynamics proceed. B) Juvenile survival rate ( ) increases nonlinearly as a function of the length of the time passed since the previous disturbance events. Dashed lines indicate the means of each axis. C) When passed through B, the different architectures of A result in divergent fitness outcomes. Panels are split for better visualization. Each colored line shows the mean time-averaged fitness ( ) over 500 iterations with increasing , after 500 event intervals (. Bands show over the iterations. Grey lines in each panel show the mean lines of the other 3 environments. Fitness is driven apart further as (amount of noise, or unpredictability, in the environment) increases.

Fitness divergence across categories of fluctuating environments

Our results show that fitness estimates of the same population model differ substantially among the four architectures of fluctuating environments (Fig. 2C), despite the analytical equivalency of the environments' distributions of intervals between disturbance events (Table 1; Fig. S1). Across the spectrum of environmental variance, was overall highest for the negatively

342   autoregressive environment, slightly lower for the cyclically driven but noisy environment

343   (Noisy Clock), still lower for the stochastic environment without autocorrelation, and by far the

344   lowest for the positively autoregressive environment. Importantly, all above differences in fitness

345   among the four environments are amplified as σ^2 is increased.

Mechanisms of fitness divergence: Consecutiveness of deviations and Jensen's Inequality

The central question that arises from Fig. 2C is why fitness estimates diverge between the four

architectures of fluctuating environments, despite the environments' distributions of event

intervals being forced to be analytically equivalent (Table 1). There are two mechanisms that

work in concert to drive the divergence: (1) *Consecutiveness of deviations from the*

environmental mean, and (2) *Jensen's Inequality in vital rate fluctuations.*

Figure 3. Mechanisms of fitness divergence between the architectures of fluctuating
environments. A) Despite analytically equivalent distributions of event intervals (Table 1),
environments have very different distributions of consecutive runs to one side of (below or above)
the mean interval depending on the nature of the generating function. Negatively autoregressive
and Noisy Clock environments inherently exhibit compensatory or self-correcting behavior,

resulting in shorter tails in the consecutive run distribution. The opposite is true of the positively
autoregressive case. The stochastic environment exhibits an intermediate exponential distribution.
B) Jensen's Inequality exists in any convex/cave fluctuating variable, and introduces an
asymmetry between environmental and demographic fluctuations. Nonlinear responses in vital
rates as a function of interval length between ecological events such as disturbances are likely
common.

(1) Consecutiveness of deviations from the environmental mean

In fluctuating environments with a mean interval between environmental disturbances,

deviations to one side of the mean interval (*i.e.* below or above) can have short consecutive runs

(*e.g.* flipping equally and quickly between below and above the mean) or long consecutive runs

(*e.g.* many deviations below or above the mean in a row) even if the long-term overall

distribution of deviations is normal. The four categories of environmental fluctuations have

different levels of compounding or compensatory behavior of fluctuations, creating different

distributions of consecutiveness (Fig. 3A). Negatively autoregressive and Noisy Clock

environments both tend to self-correct deviations back to the mean, driven by the $-\varphi$ in the

former, and by the deterministic 'due times' in the latter. Thus, these environments have short

tails in their distributions of consecutive deviation length. In contrast, the positively

autoregressive environment by definition repeats deviations to one side of the mean, resulting in

a long tail. The stochastic environment exhibits an intermediate exponential distribution of

consecutive deviations.

Why does the distribution shape of consecutive deviations matter for fitness, if over time, there is

an identical symmetrical accumulation of negative and positive deviations (*i.e.* normal

distribution of intervals between disturbances; Table 1 and Fig. S1)? Long-run fitness $log_e \lambda_s$ is a

multiplicative, or geometric, fitness measure. Consecutive runs of positive deviations of a vital

rate (*e.g.* 'good' years) will never evenly compensate for equally long consecutive runs of

negative deviations (*e.g.* 'bad' years) with respect to population growth (Haccou and Vatutin

2003, Cuddington and Hastings 2016). To illustrate this point, imagine an extreme scenario in

which a population of organisms relies on fog for moisture and can only survive for one day

without it. In environment A, a sunny day occurs on average every 100 days and is always

followed by a foggy one. Hence, the population persists. But in environment B, sunny days

always come in consecutive pairs, occurring every 200 days on average. Although sunny days

are equally common in both environments over the long run, a population inhabiting

environment B will soon be extinct. Therefore, in structured populations, the distribution of the

lengths of consecutive deviations should influence long-run fitness (longer tail = lower fitness).

(2) Jensen's Inequality in vital rate fluctuations

Fluctuations below and above the mean disturbance interval also have asymmetrical

consequences for the corresponding nonlinear demographic parameter; here that parameter is

juvenile survival S_J, which has a direct effect on fitness $log_e \lambda_s$. This asymmetry is referred to as

Jensen's Inequality (Jensen 1906, Ruel and Ayres 1999), which states that for a convex or

concave function $f(x)$, $E[f(x)] \leq or \geq f(E[x])$. Verbally, this means that symmetrical

fluctuation of a variable x about its mean along the x-axis does not translate equivalently to

symmetrical fluctuation about its mean along the y-axis. If we take fluctuation along the x-axis

as that of the environment, and fluctuation along the y-axis as that of demography, Jensen's

Inequality introduces a break in the environment-demography link in fluctuating environments

(Fig. 3B).

The direction of that asymmetry (*i.e.* sign of the inequality) and its magnitude depend on the

convex/cavity of the region of the nonlinear function where the mean lies. We demonstrate the

consequence of this dependency by manipulating the convex/cavity of the nonlinear

demographic parameter S_J around the mean. Consequently, the mean S_J, which was 0.5 in the

prior demonstration (Fig. 2C), will now equal 0.1, 0.3, 0.5, 0.7, or 0.9 (Fig. 4A). When mean

juvenile survival is low and its response function convex around the mean (Fig. 4A, panels

towards the left), fitness increases with increasing environmental variability σ^2 (corresponding

panels in Fig. 4B). This is because deviations to the right of the mean interval over-compensate

for deviations to the left in terms of survival S_J and thus boosts long-term growth of the

population. Conversely, when mean juvenile survival is high and its response function concave

around the mean (Fig. 4A, panels towards the right), fitness curves are pulled down with

increasing σ^2. The reverse explanation of the former explains this behavior. Note that, when

mean interval is exactly at the inflection point of the response function ($S_J = 0.5$; Fig. 4A center

panel), Jensen's Inequality does not come into play, as deviations to either side of the mean

interval equally compensate one another in terms of their effects on survival. Regardless of the

convex/cavity of the response function of juvenile survival, as σ^2 increases fitness curves in all

four environments begin to stabilize; this occurs because with higher environmental variance,

fluctuations begin to sample from wider breadths of the logistic response function which has a

combination of convex and concave regions, weakening the role of Jensen's Inequality.

Nevertheless, the relative ordering of fitness across fluctuation architectures remains stable (Fig.

4B, all panels), demonstrating the role that the consecutiveness of deviations from the

environmental mean plays in setting and maintaining the divergence of fitness between

fluctuation architectures.

Figure 4. Demonstration of the effect of Jensen's Inequality on fitness divergence across architectures of environmental fluctuation. A) We shift the response function of juvenile survival , and thus mean (at mean event interval), to manipulate the nonlinear shape around the mean event interval (dashed line), which is held constant. Going from left to right, the response function is more convex to more concave around the mean interval, highlighted in the yellow circles. This flips the sign of Jensen's Inequality, so that the overall effect of environmental fluctuations on fitness would generally become more positive to negative. B) The corresponding fitness plots demonstrate the sign flip in the effect of Jensen's Inequality as a result of the convex/cavity around the mean. Colored lines are means over all iterations, and bands are ± 1 sd. Colors correspond to the four architectures in previous figures. The middle panel is equivalent to a composite of Fig. 2C.

Summary and Future Outlook

Non-random temporal environmental variability abounds in nature, from autocorrelated fluctuations to externally driven cycles (Sabo and Post 2008, Dunn and Møller 2014, Hacket-Pain and Bogdziewicz 2021). Yet the rich diversity of non-random fluctuations is often subsumed under the convention of stochasticity (described by random probability distributions) in ecological and evolutionary models, for its alluring mathematical properties and 'good enough' approximations of nature. We have shown that for fitness, the architecture of non-random components of variability matters, even when long-term environmental variabilities are

distributed equivalently.

In itself, incorporating non-randomness in environmental models is not new to many researchers

working on life history theory or demography. Developments in periodic matrix models (Caswell

and Trevisan 1994), the seasonal forcing of disease dynamics (Altizer et al. 2006), bet-hedging

strategies (Venable 2007), and the effect of temporal autocorrelation on life history evolution and

population persistence (Pike et al. 2004, Metcalf and Koons 2007, Tuljapurkar et al. 2009) are

some examples of considering non-random fluctuations. The novelty of our results is two-fold:

we systematically show that random and non-random fluctuations lead to consistent differences

in demographic fitness; then, we highlight two simple but general mechanisms that drive this

divergence. These mechanisms are prevalent, and their treatment should open new doors for

investigating demography and selection dynamics in the diverse—and changing—fluctuation

architectures in nature.

Our results corroborate a message that has repeatedly emerged out of the theoretical literature of

colored noise and extinction risk, namely that "red" environments (positively autocorrelated)

increase extinction risk while "blue" environments (negatively autocorrelated) do the opposite

(Lawton 1988, Halley and Kunin 1999, van de Pol et al. 2011, Halley et al. 2018). Although we

focused on long-run fitness as opposed to extinction risk *per se,* the link to population

persistence is obvious: in positively autocorrelated environments, the long-run growth rate was

not only lowest by far, but also showed the highest amount of variance (Fig. 2C), increasing the

likelihood of a population crash proving fatal. In contrast, negatively autocorrelated fluctuations

led to the highest, most constant long-run fitness (Fig. 2C). Importantly, we further distinguish

autocorrelated environments from Noisy Clock environments (*e.g.* seasonal, tidal, and diel).

Autocorrelated environments rely on the memory of past events, whereas Noisy Clock

environments, in their simplest form, are memoryless and instead anchored to an external clock

such as Earth's spin and revolution around the sun. This subtle difference—invisible through

distributions of intervals between environmental disturbances—is visible in the distribution of

consecutive deviations from the fluctuation mean (Fig. 3A), and thus long-run fitness. In our

simulations, fitness in Noisy Clock environments consistently fell between that of red and blue

environments (Figs. 2C & 4B), suggesting that modelling a seasonal environment simply as an

autocorrelated one might lead to erroneous calculations of fitness. This result further suggests

that Noisy Clock environments, due to their locking onto an external clock, stabilize population

growth compared to other types of fluctuations; mathematical proof of this notion would be

interesting and broadly relevant as Noisy Clock-like environments are ubiquitous.

In reality, environmental fluctuations will seldom neatly fall into only one of the architectural

categories simulated here. For example, a coastal system that is strongly perturbed by periodic

tidal disturbances will also be influenced by seasonal fluctuations (Denny et al. 2009). How

resonances of non-random fluctuations constructively or destructively interfere to produce a

variety of fluctuation superpositions in different ecosystems might explain a wider breadth of

observed life history diversity and population trajectories. Even if an environment is

predominantly driven by one type of fluctuation architecture, the equivalence of the interval

distributions as we showed here can make it difficult to place a given environment into one

fluctuation category. Empiricists could consider plotting the distribution of consecutive

deviations from the measured mean fluctuation interval, as we did in Fig. 3A. The shape of this

distribution, compared against the null of a distribution of consecutive deviations in a purely

random environment (Fig. 3A, "stochastic") might give an indication of whether an environment

is fluctuating positively autoregressively (long tail in the distribution of consecutive deviations),

or negatively autoregressively / in a Noisy Clock fashion (short tail). Climate scientists and

population ecologists regularly use spectral analyses to detect dominant periodicities and lag

effects in time series of disturbance events or continuous environmental variables (Priestley 1981,

Chatfield 2003, Sabo and Post 2008), though often to remove temporal autocorrelation so that

long-term trends can be detected (Pyper and Peterman 1998, Brown et al. 2011). We argue that

consideration of such methods should become more common for life history and evolutionary

demography analyses in nature. Our demonstrations suggest that interactions between the

architecture of environmental fluctuation and nonlinear biological vital rates (*e.g.* survival,

growth, reproduction) will alter fitness estimates in a broad range of situations.

Notably, these fitness differences arise from environmental sensitivity in just one vital rate—

juvenile survival—while adult survival and fecundity were modelled as environment-

independent. For generality, we strived to simplify our life history model, but its degree of

complexity is likely another fruitful avenue of expansion and customization for specific

empirical tests. For instance, many species have more than two distinct life stages, for example

amphibians (*e.g.* egg, larva, adult) and insects (*e.g.* egg, nymph, pupa, adult). The number and

interconnectedness of stages via growth, survival, and reproduction might have other complex

and interesting influences on how fitness differs between fluctuation architectures. The nonlinear

response function of a vital rate can take many different shapes depending on the proximate

developmental, physiological, or behavioral mechanisms, from quadratic to self-excitatory

threshold to polynomial functions (Ellis and Post 2004, Stenseth et al. 2004, Nevoux et al. 2008).

However, regardless of life history complexity or functional response shape, Jensen's Inequality

must be in play if there is any nonlinear fitness-related vital rate at any life history stage. If so,

the effect should propagate through the entire population dynamic, influence fitness, and drive

apart fitness estimates in the different architectures of environmental fluctuations as variability

increases. What would vary between systems is the magnitude of that effect depending on how

nonlinear the vital rates are, and how sensitive fitness is to those vital rates. Another open

question is if interactions between multiple nonlinear vital rates would amplify or dampen the

effect of the mechanisms, to exacerbate or lessen fitness differences between environmental

fluctuation architectures.

Perturbation events or periods of harsh conditions are often stark enough to justify focusing on

the discrete intervals between them, as we did here, and as is common in disturbance ecology.

However, the amplitude, or magnitude, of a continuous environmental variable can itself

fluctuate. A prominent example is temperature variability, of particular interest in the context of

climate change (Post and Stenseth 1999, AghaKouchak et al. 2020). In the modeling framework

we present, variable amplitude can be simulated by allowing the magnitudes of subsequent

events to vary randomly or non-randomly, for example in the way events incur mortality or

influence the nonlinear vital rate response function following each event. Such analyses might

bridge the distinct modelling approaches that either focus on refractory periods between events

such as temperature extremes, and those that focus on continuous fluctuations in such variables.

Our general conclusion is that careful and appropriate treatments of the generating processes of

environmental fluctuations matter for estimating demographic fitness. In conjunction, while

nonlinear change in vital rates within seasons or disturbance phases might be arduous to measure

given the temporal sampling required, we argue that more often than not linearity is an

assumption made for convenience, not for truthfulness. At least slight nonlinearity is likely a

dominant feature of many vital rates, and even if relatively weak, nonlinearity should unlock the

synergistic role of the two mechanisms discussed here to drive differences in fitness estimates

depending on the architecture of an environment's temporal fluctuations. Such analyses might

significantly expand—as moving from constant to stochastic assumptions did (Tuljapurkar et al.

2009)—our understanding of evolutionary demography so far largely based on stochastic

assumptions.

References

AghaKouchak, A. et al. 2020. Climate Extremes and Compound Hazards in a Warming World. -
Annu. Rev. Earth Planet. Sci. 48: 519–548.

Altizer, S. et al. 2006. Seasonality and the dynamics of infectious diseases. - Ecol. Lett. 9: 467–
484.

Boyce, M. S. et al. 2006. Demography in an increasingly variable world. - Trends Ecol. Evol. 21:
141–148.

Brown, C. J. et al. 2011. Quantitative approaches in climate change ecology. - Glob. Change Biol.
17: 3697–3713.

Bürkli, A. and Jokela, J. 2017. Increase in multiple paternity across the reproductive lifespan in a
sperm-storing, hermaphroditic freshwater snail. - Mol. Ecol. 26: 5264–5278.

Caswell, H. 2001. Matrix population models: Construction, analysis, and interpretation. 2nd edn
sinauer associates. - Inc Sunderland MA in press.

Caswell, H. 2008. Perturbation analysis of nonlinear matrix population models. - Demogr. Res.
18: 59–116.

Caswell, H. and Trevisan, M. C. 1994. Sensitivity Analysis of Periodic Matrix Models. -
Ecology 75: 1299–1303.

Caswell, H. and Kaye, T. N. 2001. Stochastic demography and conservation of an endangered
perennial plant (Lomatium bradshawii) in a dynamic fire regime. - In: Advances in
Ecological Research. Academic Press, pp. 1–51.

Chatfield, C. 2003. The Analysis of Time Series: An Introduction, Sixth Edition. - Chapman and
Hall/CRC.

Chevin, L.-M. et al. 2017. Stochastic Evolutionary Demography under a Fluctuating Optimum
Phenotype. - Am. Nat. 190: 786–802.

Childs, D. Z. et al. 2010. Evolutionary bet-hedging in the real world: empirical evidence and
challenges revealed by plants. - Proc. R. Soc. B Biol. Sci. 277: 3055–3064.

Cohen, D. 1966. Optimizing reproduction in a randomly varying environment. - J. Theor. Biol. 12: 119–129.

Cole, L. C. 1954. The Population Consequences of Life History Phenomena. - Q. Rev. Biol. 29: 103–137.

Crewe, P. et al. 2018. Defining fitness in an uncertain world. - J. Math. Biol. 76: 1059–1099.

Cuddington, K. and Hastings, A. 2016. Autocorrelated environmental variation and the establishment of invasive species. - Oikos 125: 1027–1034.

Davison, R. et al. 2019. Stochastic effects contribute to population fitness differences. - Ecol. Model. 408: 108760.

Denny, M. W. et al. 2009. On the prediction of extreme ecological events. - Ecol. Monogr. 79: 397–421.

Drake, J. M. 2005. Population effects of increased climate variation. - Proc. R. Soc. B Biol. Sci. 272: 1823–1827.

Dunn, P. O. and Møller, A. P. 2014. Changes in breeding phenology and population size of birds. - J. Anim. Ecol. 83: 729–739.

Easterling, D. R. et al. 2000. Climate Extremes: Observations, Modeling, and Impacts. - Science 289: 2068–2074.

Ehrlén, J. et al. 2016. Advancing environmentally explicit structured population models of plants. - J. Ecol. 104: 292–305.

Ellis, A. M. and Post, E. 2004. Population response to climate change: linear vs. non-linear modeling approaches. - BMC Ecol. 4: 2.

Emery, N. C. and La Rosa, R. J. 2019. The Effects of Temporal Variation on Fitness, Functional Traits, and Species Distribution Patterns. - Integr. Comp. Biol. 59: 503–516.

Engen, S. et al. 2009. Reproductive Value and the Stochastic Demography of Age☐Structured Populations. - Am. Nat. 174: 795–804.

Engen, S. et al. 2013. Estimating the effect of temporally autocorrelated environments on the demography of density-independent age-structured populations. - Methods Ecol. Evol. 4: 573–584.

Felmy, A. et al. 2021. Ancestral ecological regime shapes reaction to food limitation in the Least Killifish, Heterandria formosa. - Ecol. Evol. 11: 6391–6405.

Fieberg, J. and Ellner, S. P. 2001. Stochastic matrix models for conservation and management: a comparative review of methods. - Ecol. Lett. 4: 244–266.

Fisher, R. A. 1958. The genetical theory of natural selection. - Dover.

Gadgil, M. and Bossert, W. H. 1970. Life Historical Consequences of Natural Selection. - Am. Nat. 104: 1–24.

Gaillard, J.-M. et al. 1998. Population dynamics of large herbivores: variable recruitment with constant adult survival. - Trends Ecol. Evol. 13: 58–63.

Gaillard, J.-M. et al. 2000. Temporal Variation in Fitness Components and Population Dynamics of Large Herbivores. - Annu. Rev. Ecol. Syst. 31: 367–393.

Gonzalez□Andujar, J. L. et al. 2006. Population Cycles Produced by Delayed Density Dependence in an Annual Plant. - Am. Nat. 168: 318–322.

Gremer, J. R. and Venable, D. L. 2014. Bet hedging in desert winter annual plants: optimal germination strategies in a variable environment. - Ecol. Lett. 17: 380–387.

Haccou, P. and Vatutin, V. 2003. Establishment success and extinction risk in autocorrelated environments. - Theor. Popul. Biol. 64: 303–314.

Hacket-Pain, A. and Bogdziewicz, M. 2021. Climate change and plant reproduction: trends and drivers of mast seeding change. - Philos. Trans. R. Soc. B Biol. Sci. 376: 20200379.

Halley, J. M. 2005. Comparing aquatic and terrestrial variability: at what scale do ecologists communicate? - Mar. Ecol. Prog. Ser. 304: 274–280.

Halley, J. M. and Kunin, W. E. 1999. Extinction Risk and the 1/f Family of Noise Models. - Theor. Popul. Biol. 56: 215–230.

Halley, J. M. et al. 2018. How survival curves affect populations' vulnerability to climate change. - PLOS ONE 13: e0203124.

Hayford, H. A. et al. 2018. Radio tracking detects behavioral thermoregulation at a snail's pace. - J. Exp. Mar. Biol. Ecol. 499: 17–25.

Higgins, S. I. et al. 2000. Fire, resprouting and variability: a recipe for grass–tree coexistence in savanna. - J. Ecol. 88: 213–229.

Inouye, D. W. 2008. Effects of Climate Change on Phenology, Frost Damage, and Floral Abundance of Montane Wildflowers. - Ecology 89: 353–362.

IPCC 2012. Managing the risks of extreme events and disasters to advance climate change adaptation: special report of the Intergovernmental Panel on Climate Change (CB Field, Ed.). - Cambridge University Press.

Jensen, J. L. W. V. 1906. Sur les fonctions convexes et les inégalités entre les valeurs moyennes. - Acta Math. 30: 175–193.

Jentsch, A. et al. 2007. A new generation of climate-change experiments: events, not trends. - Front. Ecol. Environ. 5: 365–374.

Keyfitz, N. 1968. Introduction to the Mathematics of Population.

Lande, R. and Orzack, S. H. 1988. Extinction dynamics of age-structured populations in a
fluctuating environment. - Proc. Natl. Acad. Sci. 85: 7418–7421.

Lande, R. et al. 2003. Stochastic Population Dynamics in Ecology and Conservation. - Oxford
University Press.

Lande, R. et al. 2017. Evolution of stochastic demography with life history tradeoffs in density-
dependent age-structured populations. - Proc. Natl. Acad. Sci. 114: 11582–11590.

Lawson, C. R. et al. 2015. Environmental variation and population responses to global change. -
Ecol. Lett. 18: 724–736.

Lawton, J. H. 1988. More time means more variation. - Nature 334: 563–563.

Levins, R. 1968. Evolution in changing environments Princeton University Press. - Princet. N. J.
in press.

Linderholm, H. W. 2006. Growing season changes in the last century. - Agric. For. Meteorol.
137: 1–14.

Lytle, D. A. 2001. Disturbance Regimes and Life History Evolution. - Am. Nat. 157: 525–536.

Lytle, D. A. 2002. Flash Floods and Aquatic Insect Life-History Evolution: Evaluation of
Multiple Models. - Ecology 83: 370–385.

Lytle, D. A. et al. 2008. Evolution of aquatic insect behaviours across a gradient of disturbance
predictability. - Proc. R. Soc. B Biol. Sci. 275: 453–462.

MacArthur, R. H. 1962. Some generalized theorems of natural selection. - Proc. Natl. Acad. Sci.
U. S. A. 48: 1893.

Marlon, J. R. et al. 2009. Wildfire responses to abrupt climate change in North America. - Proc.
Natl. Acad. Sci. 106: 2519–2524.

McAdam, A. G. and Boutin, S. 2003. Variation in Viability Selection Among Cohorts of
Juvenile Red Squirrels (tamiasciurus Hudsonicus). - Evolution 57: 1689–1697.

Metcalf, C. j. e and Koons, D. n 2007. Environmental uncertainty, autocorrelation and the
evolution of survival. - Proc. R. Soc. B Biol. Sci. 274: 2153–2160.

Morris, W. F. et al. 2006. Sensitivity of the population growth rate to demographic variability
within and between phases of the disturbance cycle. - Ecol. Lett. 9: 1331–1341.

Morris, W. F. et al. 2008. Longevity Can Buffer Plant and Animal Populations Against Changing
Climatic Variability. - Ecology 89: 19–25.

Nevoux, M. et al. 2008. Nonlinear impact of climate on survival in a migratory white stork
670   population. - J. Anim. Ecol. 77: 1143–1152.

671   Noble, J. D. et al. 2019. Foraging strategies of individual silky pocket mice over a boom–bust
672   cycle in a stochastic dryland ecosystem. - Oecologia 190: 569–578.

673   Nouvellet, P. et al. 2012. Noisy clocks and silent sunrises: measurement methods of daily
674   activity pattern. - J. Zool. 286: 179–184.

675   Park, J. S. 2019. Cyclical environments drive variation in life-history strategies: a general theory
676   of cyclical phenology. - Proc. R. Soc. B 286: 20190214.

677   Park, J. S. and Wootton, J. T. 2021. Slower environmental cycles maintain greater life-history
678   variation within populations. - Ecol. Lett. 24: 2452–2463.

679   Pélisson, P.-F. et al. 2012. Contrasted breeding strategies in four sympatric sibling insect species:
680   when a proovigenic and capital breeder copes with a stochastic environment. - Funct.
681   Ecol. 26: 198–206.

682   Pike, N. et al. 2004. The effect of autocorrelation in environmental variability on the persistence
683   of populations: an experimental test. - Proc. R. Soc. Lond. B Biol. Sci. 271: 2143–2148.

684   Post, E. and Stenseth, N. Chr. 1999. Climatic Variability, Plant Phenology, and Northern
685   Ungulates. - Ecology 80: 1322–1339.

686   Priestley, M. B. 1981. Spectral analysis and time series: probability and mathematical statistics. -
687   Academic Press.

688   Pyper, B. J. and Peterman, R. M. 1998. Comparison of methods to account for autocorrelation in
689   correlation analyses of fish data. - Can. J. Fish. Aquat. Sci. 55: 2127–2140.

690   Reuman, D. C. et al. 2008. Colour of environmental noise affects the nonlinear dynamics of
691   cycling, stage-structured populations. - Ecol. Lett. 11: 820–830.

692   Ruel, J. J. and Ayres, M. P. 1999. Jensen's inequality predicts effects of environmental variation.
693   - Trends Ecol. Evol. 14: 361–366.

694   Sabo, J. L. and Post, D. M. 2008. Quantifying Periodic, Stochastic, and Catastrophic
695   Environmental Variation. - Ecol. Monogr. 78: 19–40.

696   Schaffer, W. M. 1974. Optimal Reproductive Effort in Fluctuating Environments. - Am. Nat. 108:
697   783–790.

698   Schreiber, S. J. and Moore, J. L. 2018. The structured demography of open populations in
699   fluctuating environments. - Methods Ecol. Evol. 9: 1569–1580.

700   Shriver, R. K. 2016. Quantifying how short-term environmental variation leads to long-term
701   demographic responses to climate change. - J. Ecol. 104: 65–78.

Sletvold, N. et al. 2013. Climate warming alters effects of management on population viability of
threatened species: results from a 30-year experimental study on a rare orchid. - Glob.
Change Biol. 19: 2729–2738.

Staver, A. C. et al. 2009. Browsing and fire interact to suppress tree density in an African
savanna. - Ecol. Appl. 19: 1909–1919.

Stearns, S. C. 1976. Life-History Tactics: A Review of the Ideas. - Q. Rev. Biol. 51: 3–47.

Steele, J. H. 1985. A comparison of terrestrial and marine ecological systems. - Nature 313: 355–
358.

Stenseth, N. Chr. et al. 2004. Modelling non–additive and nonlinear signals from climatic noise
in ecological time series: Soay sheep as an example. - Proc. R. Soc. Lond. B Biol. Sci.
271: 1985–1993.

Taylor, C. et al. 2014. Nonlinear Effects of Stand Age on Fire Severity. - Conserv. Lett. 7: 355–
370.

Thornton, P. K. et al. 2014. Climate variability and vulnerability to climate change: a review. -
Glob. Change Biol. 20: 3313–3328.

Tuljapurkar, S. 1985. Population dynamics in variable environments. VI. Cyclical environments.
- Theor. Popul. Biol. 28: 1–17.

Tuljapurkar, S. 1990. Population Dynamics in Variable Environments. - Springer-Verlag.

Tuljapurkar, S. and Haridas, C. V. 2006. Temporal autocorrelation and stochastic population
growth. - Ecol. Lett. 9: 327–337.

Tuljapurkar, S. et al. 2003. The Many Growth Rates and Elasticities of Populations in Random
Environments. - Am. Nat. 162: 489–502.

Tuljapurkar, S. et al. 2009. From stochastic environments to life histories and back. - Philos.
Trans. R. Soc. B Biol. Sci. 364: 1499–1509.

van de Pol, M. et al. 2011. Poor environmental tracking can make extinction risk insensitive to
the colour of environmental noise. - Proc. R. Soc. B Biol. Sci. 278: 3713–3722.

Vasseur, D. A. and Yodzis, P. 2004. The Color of Environmental Noise. - Ecology 85: 1146–
1152.

Venable, D. L. 2007. Bet Hedging in a Guild of Desert Annuals. - Ecology 88: 1086–1090.

Walther, G.-R. et al. 2002. Ecological responses to recent climate change. - Nature 416: 389–395.

Westerling, A. L. et al. 2006. Warming and Earlier Spring Increase Western U.S. Forest Wildfire
Activity. - Science 313: 940–943.