Why the architecture of environmental fluctuation matters for fitness

Article type: Synthesis & Perspective

Short running head: Architecture of fluctuations and fitness

Keywords: fluctuating environments, life history, evolutionary demography, stochastic fitness, nonlinearity, Jensen's Inequality

Data accessibility statement: No new data were used in this study.

Number of words in abstract: 191

Number of words in main text: 5246

Number of figures: 4

Number of tables: 1

Number of text boxes: 0

Number of references: 93

Abstract

The physical environment provides the very stage upon which the eco-evolutionary play unfolds. How environmental fluctuations affect fitness is thus central to demographic projections, selection predictions, life history analyses, and the conservation of populations. Modelling efforts have mostly dealt with fluctuating environments using stochasticity. However, environmental fluctuations in nature are combinations of non-random components and stochastic noise. For example, some fluctuations contain feedbacks (*e.g.* disturbances such as floods, fires, and hurricanes), and others are driven by geophysical forces that create fixed cyclicality (*e.g.* seasonal, tidal, and diel). Here we systematically compare and explain why the 'architecture' of fluctuations—both how they are generated in nature and constructed theoretically—influences fitness of structured populations. Fitness is sensitive to the architectures even if they appear indistinguishable in their long-term statistical distributions. Importantly, we highlight two quantitative mechanisms through which fitness depends on fluctuation architecture— consecutiveness of deviations from the environmental mean, and Jensen's Inequality acting on nonlinear biological parameters—both arguably relevant features in virtually all populations inhabiting variable environments. The fitness divergence we demonstrate and explain is a useful step towards developing evolutionary demographic theory for the prevalent categories of non-random environments.

Introduction

Predicting population growth—or fitness—is a centerpiece of life history theory, evolutionary demography, and conservation biology. Cole (1954) first conceived of the fitness of life history strategies as directly linked to the strategies' influence on population growth rate. Variations in the age distribution of survival and reproduction shape the fitness landscapes of life histories (Gadgil and Bossert 1970; Stearns 1976), upon which natural selection can act. Classical demography (Keyfitz 1968; Caswell 2001) leverages these concepts to analyze how the trajectories and sensitivities of population growth are influenced by population structures (*e.g.* in age, stage, or size) and their variations.

Early theories began with assumptions of constant environments for simplicity, and because it was necessary to lay the groundwork for comparing species in a systematic manner (Fisher 1958; MacArthur 1962). Developments since (Levins 1968; Schaffer 1974) have largely focused on extending analyses to include stochastic environment assumptions to reflect the complexities of nature (Lande et al. 2003; Tuljapurkar et al. 2003), and to spotlight increasing environmental variability under climate change (Easterling et al. 2000; Drake 2005; Boyce et al. 2006; Morris et al. 2008; Lawson et al. 2015). Seminal work by Tuljapurkar *et al.* (2009) demonstrated how and why the expected optimal life histories differ between constant and stochastic environments in a general sense, by decomposing the effects of life history speed, dispersion, and temporal correlation on fitness in the two archetypical environmental formulations.

Stochasticity indeed offers a powerful suite of tools to model an extensive spectrum of environments, principally afforded by mature statistical theories of probability distributions. In short, a variable environment can typically be summarized by moments of distributions, *e.g.* the

mean and variance of a normal or lognormal distribution, whose derivations lead to tractable

analytical solutions for biological predictions in that environment (Lawson et al. 2015), *e.g.* bet-

hedging strategies (Cohen 1966; Childs et al. 2010). Higher order moments of stochastic

environments can be decomposed with Taylor expansion methods to analyze the effects of, *e.g.*,

intra- and inter-annual variability and temporal autocorrelation on fitness (Tuljapurkar et al. 2009).

Indeed, there is substantial theoretical and empirical evidence that the 'color' of noise (negative to

zero to positive autocorrelation) influences fitness and extinction risk (Cohen 1966; Lande and

Orzack 1988; Vasseur and Yodzis 2004; Halley 2005; Tuljapurkar and Haridas 2006; Reuman et

al. 2008; Engen et al. 2013; Chevin et al. 2017). In summary, modern 'stochastic demography' is

now a rich discipline that places stochastic environmental formulation at its core to ask ecological

and evolutionary questions (Tuljapurkar 1990; Fieberg and Ellner 2001; Engen et al. 2009; Chevin

et al. 2017; Lande et al. 2017; Crewe et al. 2018; Schreiber and Moore 2018; Davison et al. 2019).

Mathematically, however, stochasticity is specific: it describes a process that can be captured with

a random probability distribution.

Yet, many common categories of environmental fluctuations have non-random components. For

example, fundamental geophysical fixtures such as the rotation of Earth (governing diel cycles),

its axial tilt and revolution around the Sun (seasonal cycles), and the revolution of the Moon around

Earth (tidal cycles) set sinusoidal oscillations in most terrestrial and aquatic ecosystems.

Analogous to driven harmonic oscillators in physics, such environments are governed by

permanent (at least on ecological and evolutionary time scales) external drivers of fluctuations.

Distinct from stochastic variability, geophysically-driven fluctuations are, due to the external

driver, thus not emergent products of the internal ecological and evolutionary dynamics of the

system such as in population cycles or ecological succession.

Here we envision four broadly representative architectures of the timing patterns of events in an

environment. In all four scenarios, events represent ecological disturbances (Lytle 2001) such as

fires, floods, hurricanes, avalanches, droughts, and heatwaves, or, in seasonal systems, onsets of

winter that can largely impede biological activity (Inouye 2008). Stearns (1976, Fig. 7) outlined

the possible outcomes of environmental fluctuation frequency and predictability for the age

distribution of reproduction. Tuljapurkar (1985) analyzed how the relative scaling between the

period of environmental cycles and generation time influences population growth rate. Recent

theoretical work demonstrated that environmental periodicity is a distinct selective force that

determines both optimal (Park 2019) and individual variation in life histories (Park and Wootton

2021). Further, perturbations to seasonality and disturbance regimes are among the most

widespread signatures of climate change, with far-reaching consequences such as phenological

shifts (Walther et al. 2002; Westerling et al. 2006; Marlon et al. 2009). Importantly, mean and

variability can change independently in disturbance regimes (Lytle 2001; Lytle et al. 2008; Denny

et al. 2009), which is a growing focus in climate change biology (Jentsch et al. 2007; IPCC 2012;

Thornton et al. 2014). Thus, teasing apart the non-random and random components that constitute

environmental variability—instead of collapsing all environmental variability to stochasticity—is

crucial not only for more accurate predictions of population growth (Tuljapurkar and Haridas

2006), but for understanding fundamentally different causal drivers of population dynamics and

life history selection.

We investigate why long-run fitness differs between stochastic *vs*. non-randomly fluctuating

environments in a general sense. A particularly novel consideration is the treatment of inherently

periodic environments such as seasonal or tidal ones. These externally driven cycles with fixed

periodicity are fundamentally different in their memory process compared to internally produced

cycles (*e.g.* autocorrelated): in the former, environmental states gravitate to an external schedule,

whereas in the latter they depend on former states of themselves. We systematically demonstrate

that this difference translates to consistent fitness differences. We identify two general quantitative

mechanisms that cause divergences in fitness estimates across the four categories. The first

mechanism involves the pattern of consecutiveness of positive or negative deviations from the

mean environment; this pattern arises from the generating functions of fluctuations. The second

involves Jensen's Inequality acting on nonlinear life history traits, or vital rates; the prevalence of

such nonlinearity in nature is likely universal. Both are very general properties of populations in

variable environments, and we elucidate how they operate in concert to drive fitness estimates to

diverge from purely stochastic models. Finally, we discuss the general relevance of our findings

and identify key theoretical extensions and empirical tests that will be important moving forward.

The generating functions of fluctuating environments

We start by setting up four generating functions of environmental fluctuations (Fig. 1): stochastic

("white noise"), positively autoregressive ("red noise"), negatively autoregressive ("blue noise"),

and periodic. We mean generating functions in the mathematical sense of rules underlying discrete

encodings of sequences of events on a timeline. For all four generating functions, our focal

parameter τ_i is the *interval* between events. Again, these events represent ecological disturbances

such as fires, or the onset of winter in seasonal systems with the simplifying assumption that

winters are harsh enough to perturb or halt biological activity. In seasonal systems, note that

celestial seasons are perfectly periodic, but the meteorological (*e.g.* temperature, frost) onset of

winter and arrival of spring are variable, making the intervening warm season (the interval) also

variable in length ('climatological growing seasons', *sensu* Linderholm 2006). We derive the mean

and variance of interval distributions from each generating function to then make them exactly

analytically equivalent, before comparing the fitness consequences in environments characterized

by the different fluctuations.

Generating function I: Stochastic (random) without feedback

We generate a fluctuating environment with a random sequence of event intervals τ_i:

$$\tau_i \sim Lognormal(\mu, \sigma^2) \tag{1}$$

where μ is the imposed interval mean and σ^2 its variance. We do this instead of generating a series

of events via a Poisson process, which is a more common way of generating random events on a

timeline, to make this case comparable with the other three cases with respect to their error

distributions and because below we will manipulate the strength of noise (σ^2) while holding the

mean interval constant (which is not possible with Poisson distributions because their mean and

variance are equal). We use the lognormal distribution because intervals cannot have negative

values. Note that this manner of generating a fluctuating environment does not consider the source

or nature of the noise, nor the length of preceding intervals (*i.e.* there is no memory of past

fluctuations), but only assumes the long-run statistical distribution of variability (Fig. 1A). While

measured environmental variability may in reality always have a physical source, and all systems

may have some memory process to varying degrees, some systems are so complex and noisy that

they are well approximated by randomness. Random environments may be experienced, for

example, by mice living in dryland ecosystems with variable resource availability (Noble et al.

2019), by weevil species using strongly fluctuating oak acorns as egg-laying sites (Pélisson et al.

2012), and by annual herbs growing in or near vernal pools created by stochastic winter

precipitation events (Emery and La Rosa 2019).

Generating function II: Positively autoregressive

134   We generate a positively autocorrelated environment with a simple AR(1) process:

$$\tau_i = \mu + \varphi\tau_{i-1} + \varepsilon_i \tag{2}$$

where μ is the desired interval mean, and the interval τ_i is positively correlated by strength of

autocorrelation φ with the interval immediately preceding it (τ_{i-1}). The i.i.d. (independent

identically distributed) noise ε_i is drawn from $N(0, \sigma_\varepsilon^2)$. Positively autocorrelated environments

are characterized by a 'compounding' behavior where a longer or shorter interval than the mean is

likely to beget another longer or shorter interval in the immediate future, respectively (Fig. 1B).

Examples of positively autocorrelated fluctuations in nature include some fire regimes (Caswell

and Kaye 2001), oceanic temperatures (Steele 1985; Halley 2005), and flood regimes (Sabo and

Post 2008).

Generating function III: Negatively autoregressive

145   The negative counterpart to the positively autoregressive environment is one with a negative

146   autoregressive term $(-\varphi)$ and all other terms identical to Eq. 2:

$$\tau_i = \mu - \varphi \tau_{i-1} + \varepsilon_i \tag{3}$$

Negatively autocorrelated environments are characterized by a 'compensating' or self-correcting

behavior where a longer or shorter interval than the mean is likely to be counterbalanced with a

shorter or longer interval in the immediate future, respectively (Fig. 1C). Real examples of

negatively autocorrelated environments in nature, though much more rarely considered than

positively autocorrelated ones (Metcalf and Koons 2007), include terrestrial plant systems with

delayed density-dependence driven by nutrient cycling (Gonzalez-Andujar et al. 2006), and tree

masting (Hacket-Pain and Bogdziewicz 2021).

Generating function IV: 'Noisy Clock'

The last generating function describes a fluctuating environment with a fundamental external

driver of cyclicality, whose deterministic periodicity is not affected by the realized noise. We

henceforth refer to this case as the 'Noisy Clock' to describe the clocked (deterministic) nature of

fluctuations that underlies the noise. Here, we first punctuate the timeline with a series of

deterministic 'due' times of events much like the celestial periodicity of Earth's revolution that

determines the boundaries of the annual cycle. These due times are indexed here by n. Intervals

are not generated by counting time from the preceding event, but imperfectly timed events occur

around each due time with i.i.d. noise (Fig. 1D) Then the intervals are computed by measuring

gaps between subsequent realized events:

$$\tau_i = [\mu(i+1) + \varepsilon_{i+1}] - [\mu i + \varepsilon_i] \tag{4}$$

where $\varepsilon_i \sim N(0, \sigma_\varepsilon^2)$ and $i \in \mathbb{Z}_{>0}$, meaning clocked due times are positive integer multiples of μ, the

same mean interval as in generating functions I, II, and III. Counting time from preceding events

(which is the case for generating functions I, II, and III) as opposed to in reference to an external

clock would be an unrealistic way to model fundamentally cyclical systems such as seasonal ones.

For instance, if one were to model a winter event (*e.g.* first snowfall), but stochastically modelled

the length of intervening summer seasons with no regard to due times, then after some additive

sequence of stochastic lengths it would be possible to have winters starting in July when lined up

against a deterministic calendar. This would be unrealistic under any reasonable geophysical

stationarity assumptions. Adding error to due times has the effect of gravitating noisy events back

to realistic due times set by the underlying forces of harmonic oscillations in the environment.

Real fluctuations with clocked due times driven by geophysical forces—but with error in the

measured proxy variables—are exemplified by a myriad of diel (*e.g.* hunting behavior of African

wild dogs; Nouvellet *et al.* 2012), tidal (*e.g.* foraging of intertidal snails; Hayford *et al.* 2018), and

seasonal environments (*e.g.* advancement of egg-laying with warming springs in numerous bird

species; Dunn & Møller 2014). Despite widespread attention to seasonal environments,

particularly with respect to global phenological shifts (Walther et al. 2002), the theoretical

foundation of demographic dynamics and life history evolution in cyclical environments is not

yet well understood (Lande et al. 2017).

Comparing apples to apples: equivalent distributions of intervals

To systematically compare the fitness consequences in the four scenarios of fluctuating

environments, we take the parametric moments of the generating functions and derive input

parameters that make their interval distributions analytically equivalent (Table 1). All distributions

will be normal due to the central limit theorem (Fig. S1). In other words, by using these input

parameters all four environments' distributions of intervals will have the same mean and variance,

despite the environments' differences in fluctuation architecture. The purpose of establishing this

interval distribution equivalency is to demonstrate that long-term statistical summaries of real

environments (particularly those coming from sparse data) might lead, incorrectly, to the

complacency that one can model any actual environment as a randomly noisy environment that

can, consequently be drawn from a random probability distribution. While this simplification

might be acceptable in some cases, we will show that long-run fitness is substantially and

consistently influenced by the underlying architecture of fluctuations even if their interval

distributions are identical.

For our simulations, we set up all four generating functions to have a sample mean interval length

(μ_s) of 365 time steps (*i.e.* days) to emulate an annual event process.

A simple demographic model in fluctuating environments

Upon the environmental fluctuation backgrounds that have fundamentally different generating

functions but equivalent interval distributions, we model a simple population that proceeds through

time. There are two time indices: t which denotes time steps, here representing days, and τ_i which

denotes the intervals between intermittent disturbance events set by the generating functions. J_t

and A_t are numbers of juveniles and adults at time t respectively, f is rate of reproduction, and S_J

[0,1] and S_A [0,1] are rates of juvenile and adult survival when disturbances happen, respectively.

The population dynamic is then described in a system of discrete difference equations consisting

of juveniles and reproducing adults:

$$J_{t+1} = A_t f$$
$$A_{t+1} = A_t S_A + J_t S_J \text{ ,}$$

$$where \ S_J = \begin{cases} S_J(\tau) = \dfrac{1}{1 + e^{-0.1(\tau-365)}}, & \text{if } t \in y_k = \sum_{i=1}^{k} \tau_i, \\ 1, & \text{otherwise} \end{cases} \quad (5)$$

We focus on juvenile survival S_J as the vital rate that dynamically interacts with the disturbance

events that arrive intermittently into the system. This is because, in many natural populations of

plants or animals, juvenile mortality is especially sensitive to harsh environmental conditions

(disturbance or winter) (Gaillard et al. 1998, 2000; Higgins et al. 2000; Lytle 2002; McAdam and

Boutin 2003; Staver et al. 2009; Park 2019). In Eq 5, juvenile survival S_J is determined by a

response function $S_J(\tau)$ when disturbance occurs (*i.e.* when time step t is an element of the

predetermined sequence of disturbance timings, which is the cumulative sum of successive interval

lengths $\sum_{i=1}^{k} \tau_i$, where k is number of intervals simulated), and is 1 otherwise. Here we have set

the response function $S_J(\tau)$ to be a nonlinearly increasing function that depends on how much time

has passed since the last event (τ), with juvenile survival being 50% (*i.e.* $S_J = 0.5$) at the mean

disturbance interval (Fig. 2B). Verbally, this means that if two disturbance events happen in quick

succession, juvenile survival will be low, and if a long time passes between events, it will be high;

and this change in juvenile survival as a function of interval length is nonlinear. Biologically, this

function signifies the time needed for juveniles to sufficiently develop, grow, or learn to endure

the next impending perturbation in the environment. Though not always logistic, nonlinearity in

vital rate response functions in between perturbations in the environmental state is likely common

(Caswell 2008; Childs et al. 2010; Sletvold et al. 2013; Taylor et al. 2014; Ehrlén et al. 2016;

Shriver 2016). For example, Morris et al. (2006) showed that, across five species of perennial

plants, vital rates including seedling survival (equivalent to S_J here), survival at other stages,

growth, germination, and seed production (equivalent to f here) changed nonlinearly as a function

of time since last hurricane or fire disturbance. Similarly, Bürkli & Jokela (2017) showed that

embryonic development success (early-stage juvenile survival) of a freshwater snail changed

nonlinearly across the species' reproductive season. The exact shape of vital rate response

functions is another key point to be tailored for system-specific inquiry, whether with empirical

measurements or as realistic assumptions built into models.

The model is otherwise kept simple for generality, and to focus on the effect that the different

generating functions have on fitness. Structural complexity such as the number of developmental

stages (*i.e.* the number of equations in the system, or the number of vital rate terms) would be an

obvious area of customization for system-specific inquiry. We also hold f constant at 0.3, and S_A

constant at 0.9 when disturbance occurs irrespective of time since last disturbance, and 1 otherwise.

These values are arbitrary but biologically realistic; exact values did not influence the downstream

qualitative results. As we will show, the important mechanisms should influence fitness regardless

of demographic nuance.

Time-averaged fitness varies across the fluctuating environments

Within each of the four architectures of environmental fluctuations, we ask what the long-run

fitness of a life history would be. For fitness, we use:

$$log_e\lambda_s = \lim_{t\to\infty} \left(\frac{1}{t}\right) log_e[N(t)/N(0)] \qquad\qquad (6)$$

which gives the long-run growth rate in fluctuating environments (Tuljapurkar 1990) where $N(t)$

is the total population size at time *t*, including both juvenile and adult stages. This metric can be

computed from numerical simulations over a long time course of the environment (Caswell 2001),

and robustly approximates fitness even under considerably large fluctuations (Lande et al. 2003;

Tuljapurkar et al. 2003, 2009). We simulate demographic dynamics (Eq. 5) in each fluctuating

environment (Fig. 1) for 1000 event intervals, calculate $log_e\lambda_s$, and repeat these calculations over

500 iterations. Lastly, we compute all of the above across a spectrum of the generating functions'

variance, to assess how the amount of fluctuation noise affects the divergence of fitness estimates.

Our results show that fitness estimates of the same life history model differ substantially among

the four architectures of fluctuating environments (Fig. 2C), despite the analytical equivalency of

the environments' distributions of event intervals (Table 1). The long-run growth rate $log_e\lambda_s$ was

overall highest for the negatively autoregressive environment, slightly lower for the cyclically

driven but noisy environment (Noisy Clock), still lower for the stochastic environment without

autocorrelation, and by far the lowest for the positively autoregressive environment. Mean fitness

(thicker lines, Fig. 2C) was roughly inversely correlated with amount of variance (spread of dots,

Fig. 2C): the lower the mean, the greater the spread. Furthermore, the four environments exhibited

unequal responses of fitness to an increase in the amount of noise (σ^2 of event intervals): while

increasing σ^2 slightly increased fitness in the negatively autoregressive and slightly decreased

fitness in the Noisy Clock environment, it considerably reduced fitness in stochastic and positively

autoregressive environments (Fig. 2C). Importantly, all above differences in fitness among the

four environments are amplified as σ^2 is increased. The divergence in fitness also remains largely

unchanged when the rate of reproduction f and adult survival S_A are allowed to vary (Fig. S2).

Mechanisms of fitness divergence: Consecutiveness of deviations and Jensen's Inequality

The central question that arises from Fig. 2C is why fitness estimates diverge between the four

architectures of fluctuating environments, despite the environments' distributions of event

intervals being forced to be analytically equivalent (Table 1). There are two mechanisms that work

in concert to drive the divergence: (1) *Consecutiveness of deviations from the environmental mean,*

and (2) *Jensen's Inequality in vital rate fluctuations.*

(1) Consecutiveness of deviations from the environmental mean

In fluctuating environments with a mean interval, deviations to one side of the mean interval (*i.e.*

below or above) can have short consecutive runs (*e.g.* flipping equally and quickly between below

and above the mean) or long consecutive runs (*e.g.* many deviations below or above the mean in a

row) even if the long-term overall distribution of deviations is normal. The four generating

functions have different levels of compounding or compensatory behavior of fluctuations, creating

different distributions of consecutiveness (Fig. 3A). Negatively autoregressive and Noisy Clock

environments both tend to self-correct deviations back to the mean, driven by the $-\varphi$ in the former,

and by the deterministic 'due times' in the latter. Thus, these environments have short tails in their

distributions of consecutive deviation length. In contrast, the positively autoregressive

environment by definition repeats deviations to one side of the mean, resulting in a long tail. The

stochastic environment exhibits an intermediate exponential distribution of consecutive deviations.

Why does the distribution shape of consecutive deviations matter for fitness, if over time, there is

an identical symmetrical accumulation of negative and positive deviations (*i.e.* normal distribution

of intervals; Table 1)? Long-run fitness $log_e\lambda_s$ is a multiplicative, or geometric, fitness measure.

Consecutive runs of positive deviations of a vital rate (*e.g.* 'good' years) will never evenly

compensate for equally long consecutive runs of negative deviations of the same magnitude (*e.g.*

'bad' years) with respect to population growth (Haccou and Vatutin 2003; Cuddington and

Hastings 2016). Therefore, in structured populations, the distribution of the lengths of consecutive

deviations should influence long-run fitness (longer tail = lower fitness).

(2) Jensen's Inequality in vital rate fluctuations

Fluctuations below and above the mean interval also have asymmetrical consequences for the

corresponding nonlinear demographic parameter; here that parameter is juvenile survival S_J, which

has a direct effect on fitness $log_e\lambda_s$. This asymmetry is referred to as Jensen's Inequality (Jensen

1906; Ruel and Ayres 1999), which states that for a convex or concave function $f(x)$, $E[f(x)] \leq$

$or \geq f(E[x])$. Verbally, this means that symmetrical fluctuation of a variable x about its mean

along the x-axis does not translate equivalently to symmetrical fluctuation about its mean along

the y-axis. If we take fluctuation along the x-axis as that of the environment, and fluctuation along

the y-axis as that of demography, Jensen's Inequality introduces a break in the environment-

demography link in fluctuating environments (Fig. 3B).

The direction of that asymmetry (*i.e.* sign of the inequality) and its magnitude depend on the

convex/cavity of the region of the nonlinear function where the mean lies. We demonstrate the

consequence of this dependency by manipulating the convex/cavity of the nonlinear demographic

parameter S_J around the mean (Fig. 4A). When mean juvenile survival is low and its response

function convex around the mean (Fig. 4A, panels towards the left), fitness increases with

increasing environmental variability σ^2 (corresponding panels in Fig. 4B). Conversely, when

mean juvenile survival is high and its response function concave around the mean (Fig. 4A, panels

towards the right), fitness curves are pulled down with increasing σ^2. Regardless of the

convex/cavity of the response function of juvenile survival, as σ^2 increases fitness curves in all

four environments begin to stabilize; this occurs because with higher environmental variance,

fluctuations begin to sample from wider breadths of the logistic response function which has a

combination of convex and concave regions, weakening the relative role of Jensen's Inequality.

Nevertheless, the relative ordering of fitness across fluctuation architectures remains stable (Fig.

4B, all panels), demonstrating the interactive role that the consecutiveness of deviations from the

environmental mean and the geometric nature of $log_e\lambda_s$ play in setting the initial divergence of

fitness between fluctuation architectures, and maintaining it.

Summary and Future Outlook

Non-random temporal environmental variability abounds in nature, from autocorrelated

fluctuations to externally driven cycles (Sabo and Post 2008; Dunn and Møller 2014; Hacket-Pain

and Bogdziewicz 2021).Yet the rich diversity of non-random fluctuations is often absorbed into

stochasticity (described by random probability distributions) in ecological and evolutionary

models, for its alluring mathematical properties and 'good enough' approximations of nature. We

have shown that for fitness, the architecture of non-random components of variability matters,

even when long-term environmental variabilities are distributed equivalently.

In itself, incorporating non-randomness in environmental models is not new to many researchers

working on life history theory or demography. Developments in periodic matrix models (Caswell

and Trevisan 1994), the seasonal forcing of disease dynamics (Altizer et al. 2006), bet-hedging

strategies (Venable 2007), and the effect of temporal autocorrelation on life history evolution and

population persistence (Pike et al. 2004; Metcalf and Koons 2007; Tuljapurkar et al. 2009) are just

some examples. The novelty of our results is two-fold: we show for the first time that a systematic

comparison of random and non-random fluctuations reveals consistent differences in demographic

fitness; then, we highlight two simple mechanisms that drive this divergence. These mechanisms

are prevalent, and their treatment should open new doors for investigating demography and

selection dynamics in the diverse—and changing—fluctuation architectures in nature.

Our results corroborate a message that has repeatedly emerged out of the theoretical literature of

colored noise and extinction risk, namely that "red" environments (positively autocorrelated)

increase extinction risk while "blue" environments (negatively autocorrelated) do the opposite

(Lawton 1988; Halley and Kunin 1999; van de Pol et al. 2011; Halley et al. 2018). Although we

focused on long-run fitness as opposed to extinction risk *per se,* the link to population persistence

is obvious: in positively autocorrelated environments, the long-run growth rate was not only lowest

by far, it also showed the highest amount of variance (Fig. 2C), increasing the likelihood of a

population crash proving fatal. In contrast, negatively autocorrelated fluctuations led to the highest,

most constant long-run fitness (Fig. 2C). Importantly, we further distinguish autocorrelated

environments from Noisy Clock environments (*e.g.* seasonal, tidal, and diel). Autocorrelated

environments rely on the memory of past events, whereas Noisy Clock environments, in their

simplest form, are memoryless and instead anchored to an external clock such as Earth's spin and

revolution around the sun. This subtle difference—invisible through distributions of fluctuation

intervals—translates to fundamental differences in the distribution of consecutive deviations from

the fluctuation mean (Fig. 3A), and thus long-run fitness. In our simulations, fitness in Noisy Clock

environments consistently fell between that of red and blue environments (Figs. 2C & 4B),

suggesting that modelling a seasonal environment simply as an autocorrelated one might lead to

erroneous calculations of fitness. This result further suggests that Noisy Clock environments, due

to their locking onto an external clock, stabilize population growth compared to other types of

fluctuations; mathematical proof of this notion would be interesting and broadly relevant as Noisy

Clock-like environments are ubiquitous.

In reality, environmental fluctuations will seldom neatly fall into only one of the architectural

categories simulated here. For example, a coastal system that is strongly perturbed by periodic

tidal disturbances will also be influenced by seasonal fluctuations (Denny et al. 2009). How

resonances of non-random fluctuations constructively or destructively interfere to produce a

variety of fluctuation superpositions in different ecosystems might explain a wider breadth of

observed life history diversity and population trajectories. Even if an environment is

predominantly driven by one type of fluctuation architecture, the equivalence of the interval

distributions as we showed here can make it difficult to place a given environment into one

fluctuation category. Empiricists could consider plotting the distribution of consecutive deviations

from the measured mean fluctuation interval, as we did in Fig. 3A. The shape of this distribution,

compared against the null of a distribution of consecutive deviations simulated with a purely

random fluctuation assumption (Fig. 3A, "stochastic") might give an indication of whether an

environment is fluctuating positively autoregressively (long tail in the distribution of consecutive

deviations), or negatively autoregressively / in a Noisy Clock fashion (short tail). Climate scientists

and population ecologists regularly use spectral analyses to detect dominant periodicities and lag

effects in time series of disturbance events or continuous environmental variables (Priestley 1981;

Chatfield 2003; Sabo and Post 2008), though often to remove temporal autocorrelation so that

long-term trends can be detected (Pyper and Peterman 1998; Brown et al. 2011). We argue that

consideration of such methods should become more common for life history and evolutionary

demography analyses in nature. Our demonstrations suggest that interactions between the

architecture of environmental fluctuation and nonlinear biological vital rates (*e.g.* survival, growth,

reproduction) will alter fitness estimates in a broad range of situations.

Notably, these fitness differences arise from environmental sensitivity in just one vital rate—

juvenile survival—while adult survival and fecundity were modelled as environment-independent.

For generality, we strived to simplify our life history model, but its degree of complexity is likely

another fruitful avenue of expansion and customization for specific empirical tests. For instance,

many species have more than two life stages, for example amphibians (*e.g.* egg, larva, adult) and

insects (*e.g.* egg, nymph, pupa, adult). The number and interconnectedness of stages via growth,

survival, and reproduction might have other complex and interesting influences on how fitness

differs between fluctuation architectures. The nonlinear response function of a vital rate can take

many different shapes depending on the proximate developmental, physiological, or behavioral

mechanisms, from quadratic to self-excitatory threshold to polynomial functions (Ellis and Post

2004; Stenseth et al. 2004; Nevoux et al. 2008). However, regardless of life history complexity or

functional response shape, the mechanisms we elucidated here (consecutive patterns in

fluctuations and Jensen's Inequality) must be in play if there is any nonlinear fitness-related vital

rate at any life history stage. If so, the effect should propagate through the entire population

dynamic, influence fitness, and drive apart fitness estimates in the different architectures of

environmental fluctuations as variability increases. What would vary between systems is the

magnitude of that effect depending on how nonlinear the vital rates are, and how sensitive fitness

is to those vital rates. Another open question is if interactions between multiple nonlinear vital

rates would amplify or dampen the effect of the mechanisms, to exacerbate or lessen fitness

differences between environmental fluctuation architectures.

Perturbation events or periods of harsh conditions are often stark enough to justify focusing on the

discrete intervals between them, as we did here, and as is common in disturbance ecology theory.

However, the amplitude, or magnitude, of a continuous environmental variable can itself fluctuate.

A prominent example is temperature variability, of particular interest in the context of climate

change (Post and Stenseth 1999; AghaKouchak et al. 2020). In the theoretical framework we

present (Fig. 1), variable amplitude can be simulated by allowing the magnitudes of subsequent

events to vary randomly or non-randomly, for example in the way events incur mortality or

influence the nonlinear vital rate response function following each event. Such analyses might

bridge the modelling philosophies that focus on refractory periods between events such as

temperature extremes and those that focus on continuous fluctuations in such variables.

Our general conclusion is that careful and appropriate treatments of the generating processes of

environmental fluctuations matter for estimating demographic fitness. In conjunction, while

nonlinear change in vital rates within seasons or disturbance phases might be arduous to measure

given the temporal sampling required, we argue that more often than not linearity is an assumption

made for simplicity, not inspired by nature. At least slight nonlinearity is likely a dominant feature

of many vital rates, and even if relatively weak, nonlinearity should unlock the synergistic role of

the two mechanisms discussed here to drive differences in fitness estimates depending on the

architecture of an environment's temporal fluctuations. Such analyses might significantly

improve—as moving from constant to stochastic assumptions did (Tuljapurkar et al. 2009)—our

understanding of evolutionary demography so far largely based on stochastic assumptions.

References

AghaKouchak, A., F. Chiang, L. S. Huning, C. A. Love, I. Mallakpour, O. Mazdiyasni, H.
Moftakhari, et al. 2020. Climate Extremes and Compound Hazards in a Warming World.
Annual Review of Earth and Planetary Sciences 48:519–548.

Altizer, S., A. Dobson, P. Hosseini, P. Hudson, M. Pascual, and P. Rohani. 2006. Seasonality
and the dynamics of infectious diseases. Ecology Letters 9:467–484.

Boyce, M. S., C. V. Haridas, C. T. Lee, and the NCEAS Stochastic Demography Working
Group. 2006. Demography in an increasingly variable world. Trends in Ecology & Evolution
21:141–148.

Brown, C. J., D. S. Schoeman, W. J. Sydeman, K. Brander, L. B. Buckley, M. Burrows, C. M.
Duarte, et al. 2011. Quantitative approaches in climate change ecology. Global Change
Biology 17:3697–3713.

Bürkli, A., and J. Jokela. 2017. Increase in multiple paternity across the reproductive lifespan in
a sperm-storing, hermaphroditic freshwater snail. Molecular Ecology 26:5264–5278.

Caswell, H. 2001. Matrix population models: Construction, analysis, and interpretation. 2nd edn
sinauer associates. Inc., Sunderland, MA.

———. 2008. Perturbation analysis of nonlinear matrix population models. Demographic
Research 18:59–116.

Caswell, H., and T. N. Kaye. 2001. Stochastic demography and conservation of an endangered
perennial plant (Lomatium bradshawii) in a dynamic fire regime. Pages 1–51 *in*Advances in
Ecological Research (Vol. 32). Academic Press.

Caswell, H., and M. C. Trevisan. 1994. Sensitivity Analysis of Periodic Matrix Models. Ecology
75:1299–1303.

Chatfield, C. 2003. The Analysis of Time Series: An Introduction, Sixth Edition (6th ed.).
Chapman and Hall/CRC, New York.

Chevin, L.-M., O. Cotto, and J. Ashander. 2017. Stochastic Evolutionary Demography under a
Fluctuating Optimum Phenotype. The American Naturalist 190:786–802.

Childs, D. Z., C. J. E. Metcalf, and M. Rees. 2010. Evolutionary bet-hedging in the real world:
empirical evidence and challenges revealed by plants. Proceedings of the Royal Society B:
Biological Sciences 277:3055–3064.

Cohen, D. 1966. Optimizing reproduction in a randomly varying environment. Journal of
Theoretical Biology 12:119–129.

Cole, L. C. 1954. The Population Consequences of Life History Phenomena. The Quarterly
Review of Biology 29:103–137.

Crewe, P., R. Gratwick, and A. Grafen. 2018. Defining fitness in an uncertain world. Journal of
Mathematical Biology 76:1059–1099.

Cuddington, K., and A. Hastings. 2016. Autocorrelated environmental variation and the
establishment of invasive species. Oikos 125:1027–1034.

Davison, R., M. Stadman, and E. Jongejans. 2019. Stochastic effects contribute to population
fitness differences. Ecological Modelling 408:108760.

Denny, M. W., L. J. H. Hunt, L. P. Miller, and C. D. G. Harley. 2009. On the prediction of
extreme ecological events. Ecological Monographs 79:397–421.

Drake, J. M. 2005. Population effects of increased climate variation. Proceedings of the Royal
Society B: Biological Sciences 272:1823–1827.

Dunn, P. O., and A. P. Møller. 2014. Changes in breeding phenology and population size of
birds. Journal of Animal Ecology 83:729–739.

Easterling, D. R., G. A. Meehl, C. Parmesan, S. A. Changnon, T. R. Karl, and L. O. Mearns.
2000. Climate Extremes: Observations, Modeling, and Impacts. Science 289:2068–2074.

Ehrlén, J., W. F. Morris, T. von Euler, and J. P. Dahlgren. 2016. Advancing environmentally
explicit structured population models of plants. Journal of Ecology 104:292–305.

Ellis, A. M., and E. Post. 2004. Population response to climate change: linear vs. non-linear
modeling approaches. BMC Ecology 4:2.

Emery, N. C., and R. J. La Rosa. 2019. The Effects of Temporal Variation on Fitness, Functional
Traits, and Species Distribution Patterns. Integrative and Comparative Biology 59:503–516.

Engen, S., R. Lande, B. Sæther, and F. S. Dobson. 2009. Reproductive Value and the Stochastic
Demography of Age-Structured Populations. The American Naturalist 174:795–804.

Engen, S., B.-E. Sæther, K. B. Armitage, D. T. Blumstein, T. H. Clutton-Brock, F. S. Dobson,
M. Festa-Bianchet, et al. 2013. Estimating the effect of temporally autocorrelated
environments on the demography of density-independent age-structured populations.
Methods in Ecology and Evolution 4:573–584.

Fieberg, J., and S. P. Ellner. 2001. Stochastic matrix models for conservation and management: a
comparative review of methods. Ecology Letters 4:244–266.

Fisher, R. A. 1958. The genetical theory of natural selection. Dover, New York.

Gadgil, M., and W. H. Bossert. 1970. Life Historical Consequences of Natural Selection. The
American Naturalist 104:1–24.

Gaillard, J.-M., M. Festa-Bianchet, and N. G. Yoccoz. 1998. Population dynamics of large herbivores: variable recruitment with constant adult survival. Trends in Ecology & Evolution 13:58–63.

Gaillard, J.-M., M. Festa-Bianchet, N. G. Yoccoz, A. Loison, and C. Toïgo. 2000. Temporal Variation in Fitness Components and Population Dynamics of Large Herbivores. Annual Review of Ecology and Systematics 31:367–393.

Gonzalez-Andujar, J. L., C. Fernandez-Quintanilla, and L. Navarrete. 2006. Population Cycles Produced by Delayed Density Dependence in an Annual Plant. The American Naturalist 168:318–322.

Haccou, P., and V. Vatutin. 2003. Establishment success and extinction risk in autocorrelated environments. Theoretical Population Biology, Understanding the role of environmental vatiation in population and community dynamics 64:303–314.

Hacket-Pain, A., and M. Bogdziewicz. 2021. Climate change and plant reproduction: trends and drivers of mast seeding change. Philosophical Transactions of the Royal Society B: Biological Sciences 376:20200379.

Halley, J. M. 2005. Comparing aquatic and terrestrial variability: at what scale do ecologists communicate? Marine Ecology Progress Series 304:274–280.

Halley, J. M., K. S. V. Houtan, and N. Mantua. 2018. How survival curves affect populations' vulnerability to climate change. PLOS ONE 13:e0203124.

Halley, J. M., and W. E. Kunin. 1999. Extinction Risk and the 1/f Family of Noise Models. Theoretical Population Biology 56:215–230.

Hayford, H. A., M. J. O'Donnell, and E. Carrington. 2018. Radio tracking detects behavioral thermoregulation at a snail's pace. Journal of Experimental Marine Biology and Ecology 499:17–25.

Higgins, S. I., W. J. Bond, and W. S. W. Trollope. 2000. Fire, resprouting and variability: a recipe for grass–tree coexistence in savanna. Journal of Ecology 88:213–229.

Inouye, D. W. 2008. Effects of Climate Change on Phenology, Frost Damage, and Floral Abundance of Montane Wildflowers. Ecology 89:353–362.

IPCC. 2012. Managing the risks of extreme events and disasters to advance climate change adaptation: special report of the Intergovernmental Panel on Climate Change. (C. B. Field, ed.). Cambridge University Press, New York.

Jensen, J. L. W. V. 1906. Sur les fonctions convexes et les inégalités entre les valeurs moyennes. Acta Mathematica 30:175–193.

Jentsch, A., J. Kreyling, and C. Beierkuhnlein. 2007. A new generation of climate-change experiments: events, not trends. Frontiers in Ecology and the Environment 5:365–374.

Keyfitz, N. 1968. *Introduction to the Mathematics of Population*.

Lande, R., P. of B. S. Engen, S. Engen, B.-E. Sæther, and P. of P. E. B.-E. Saether. 2003.
Stochastic Population Dynamics in Ecology and Conservation. Oxford University Press.

Lande, R., S. Engen, and B.-E. Sæther. 2017. Evolution of stochastic demography with life
history tradeoffs in density-dependent age-structured populations. Proceedings of the
National Academy of Sciences 114:11582–11590.

Lande, R., and S. H. Orzack. 1988. Extinction dynamics of age-structured populations in a
fluctuating environment. Proceedings of the National Academy of Sciences 85:7418–7421.

Lawson, C. R., Y. Vindenes, L. Bailey, and M. van de Pol. 2015. Environmental variation and
population responses to global change. Ecology Letters 18:724–736.

Lawton, J. H. 1988. More time means more variation. Nature 334:563–563.

Levins, R. 1968. Evolution in changing environments Princeton University Press. Princeton,
New Jersey.

Linderholm, H. W. 2006. Growing season changes in the last century. Agricultural and Forest
Meteorology 137:1–14.

Lytle, D. A. 2001. Disturbance Regimes and Life-History Evolution. The American Naturalist
157:525–536.

———. 2002. Flash Floods and Aquatic Insect Life-History Evolution: Evaluation of Multiple
Models. Ecology 83:370–385.

Lytle, D. A., M. T. Bogan, and D. S. Finn. 2008. Evolution of aquatic insect behaviours across a
gradient of disturbance predictability. Proceedings of the Royal Society B: Biological
Sciences 275:453–462.

MacArthur, R. H. 1962. Some generalized theorems of natural selection. Proceedings of the
National Academy of Sciences of the United States of America 48:1893.

Marlon, J. R., P. J. Bartlein, M. K. Walsh, S. P. Harrison, K. J. Brown, M. E. Edwards, P. E.
Higuera, et al. 2009. Wildfire responses to abrupt climate change in North America.
Proceedings of the National Academy of Sciences 106:2519–2524.

McAdam, A. G., and S. Boutin. 2003. Variation in Viability Selection Among Cohorts of
Juvenile Red Squirrels (tamiasciurus Hudsonicus). Evolution 57:1689–1697.

Metcalf, C. j. e, and D. n Koons. 2007. Environmental uncertainty, autocorrelation and the
evolution of survival. Proceedings of the Royal Society B: Biological Sciences 274:2153–
2160.

Morris, W. F., C. A. Pfister, S. Tuljapurkar, C. V. Haridas, C. L. Boggs, M. S. Boyce, E. M.
Bruna, et al. 2008. Longevity Can Buffer Plant and Animal Populations Against Changing
Climatic Variability. Ecology 89:19–25.

Morris, W. F., S. Tuljapurkar, C. V. Haridas, E. S. Menges, C. C. Horvitz, and C. A. Pfister.
2006. Sensitivity of the population growth rate to demographic variability within and
between phases of the disturbance cycle. Ecology Letters 9:1331–1341.

Nevoux, M., J.-C. Barbraud, and C. Barbraud. 2008. Nonlinear impact of climate on survival in a
migratory white stork population. Journal of Animal Ecology 77:1143–1152.

Noble, J. D., S. L. Collins, A. J. Hallmark, K. Maldonado, B. O. Wolf, and S. D. Newsome.
2019. Foraging strategies of individual silky pocket mice over a boom–bust cycle in a
stochastic dryland ecosystem. Oecologia 190:569–578.

Nouvellet, P., G. S. A. Rasmussen, D. W. Macdonald, and F. Courchamp. 2012. Noisy clocks
and silent sunrises: measurement methods of daily activity pattern. Journal of Zoology
286:179–184.

Park, J. S. 2019. Cyclical environments drive variation in life-history strategies: a general theory
of cyclical phenology. Proceedings of the Royal Society B 286:20190214.

Park, J. S., and J. T. Wootton. 2021. Slower environmental cycles maintain greater life-history
variation within populations. Ecology Letters 24:2452–2463.

Pélisson, P.-F., M.-C. Bel-Venner, B. Rey, L. Burgevin, F. Martineau, F. Fourel, C. Lecuyer, et
al. 2012. Contrasted breeding strategies in four sympatric sibling insect species: when a
proovigenic and capital breeder copes with a stochastic environment. Functional Ecology
26:198–206.

Pike, N., T. Tully, P. Haccou, and R. Ferrière. 2004. The effect of autocorrelation in
environmental variability on the persistence of populations: an experimental test.
Proceedings of the Royal Society of London. Series B: Biological Sciences 271:2143–2148.

Post, E., and N. Chr. Stenseth. 1999. Climatic Variability, Plant Phenology, and Northern
Ungulates. Ecology 80:1322–1339.

Priestley, M. B. 1981. Spectral analysis and time series: probability and mathematical statistics.
Academic Press.

Pyper, B. J., and R. M. Peterman. 1998. Comparison of methods to account for autocorrelation in
correlation analyses of fish data. Canadian Journal of Fisheries and Aquatic Sciences
55:2127–2140.

Reuman, D. C., R. F. Costantino, R. A. Desharnais, and J. E. Cohen. 2008. Colour of
environmental noise affects the nonlinear dynamics of cycling, stage-structured populations.
Ecology Letters 11:820–830.

Ruel, J. J., and M. P. Ayres. 1999. Jensen's inequality predicts effects of environmental
variation. Trends in Ecology & Evolution 14:361–366.

Sabo, J. L., and D. M. Post. 2008. Quantifying Periodic, Stochastic, and Catastrophic
Environmental Variation. Ecological Monographs 78:19–40.

Schaffer, W. M. 1974. Optimal Reproductive Effort in Fluctuating Environments. The American
Naturalist 108:783–790.

Schreiber, S. J., and J. L. Moore. 2018. The structured demography of open populations in
fluctuating environments. Methods in Ecology and Evolution 9:1569–1580.

Shriver, R. K. 2016. Quantifying how short-term environmental variation leads to long-term
demographic responses to climate change. Journal of Ecology 104:65–78.

Sletvold, N., J. P. Dahlgren, D.-I. Øien, A. Moen, and J. Ehrlén. 2013. Climate warming alters
effects of management on population viability of threatened species: results from a 30-year
experimental study on a rare orchid. Global Change Biology 19:2729–2738.

Staver, A. C., W. J. Bond, W. D. Stock, S. J. van Rensburg, and M. S. Waldram. 2009. Browsing
and fire interact to suppress tree density in an African savanna. Ecological Applications
19:1909–1919.

Stearns, S. C. 1976. Life-History Tactics: A Review of the Ideas. The Quarterly Review of
Biology 51:3–47.

Steele, J. H. 1985. A comparison of terrestrial and marine ecological systems. Nature 313:355–
358.

Stenseth, N. Chr., K.-S. Chan, G. Tavecchia, T. Coulson, A. Mysterud, T. Clutton-Brock, and B.
Grenfell. 2004. Modelling non–additive and nonlinear signals from climatic noise in
ecological time series: Soay sheep as an example. Proceedings of the Royal Society of
London. Series B: Biological Sciences 271:1985–1993.

Taylor, C., M. A. McCarthy, and D. B. Lindenmayer. 2014. Nonlinear Effects of Stand Age on
Fire Severity. Conservation Letters 7:355–370.

Thornton, P. K., P. J. Ericksen, M. Herrero, and A. J. Challinor. 2014. Climate variability and
vulnerability to climate change: a review. Global Change Biology 20:3313–3328.

Tuljapurkar, S. 1985. Population dynamics in variable environments. VI. Cyclical environments.
Theoretical Population Biology 28:1–17.

———. 1990. Population Dynamics in Variable Environments. Springer-Verlag.

Tuljapurkar, S., J.-M. Gaillard, and T. Coulson. 2009. From stochastic environments to life
histories and back. Philosophical Transactions of the Royal Society B: Biological Sciences
364:1499–1509.

Tuljapurkar, S., and C. V. Haridas. 2006. Temporal autocorrelation and stochastic population
growth. Ecology Letters 9:327–337.

Tuljapurkar, S., C. C. Horvitz, and J. B. Pascarella. 2003. The Many Growth Rates and
Elasticities of Populations in Random Environments. The American Naturalist 162:489–502.

van de Pol, M., Y. Vindenes, B.-E. Sæther, S. Engen, B. J. Ens, K. Oosterbeek, and J. M.
Tinbergen. 2011. Poor environmental tracking can make extinction risk insensitive to the
colour of environmental noise. Proceedings of the Royal Society B: Biological Sciences
278:3713–3722.

Vasseur, D. A., and P. Yodzis. 2004. The Color of Environmental Noise. Ecology 85:1146–
1152.

Venable, D. L. 2007. Bet Hedging in a Guild of Desert Annuals. Ecology 88:1086–1090.

Walther, G.-R., E. Post, P. Convey, A. Menzel, C. Parmesan, T. J. C. Beebee, J.-M. Fromentin,
et al. 2002. Ecological responses to recent climate change. Nature 416:389–395.

Westerling, A. L., H. G. Hidalgo, D. R. Cayan, and T. W. Swetnam. 2006. Warming and Earlier
Spring Increase Western U.S. Forest Wildfire Activity. Science 313:940–943.

Figures & Legends

Figure 1. Four generating functions of fluctuating environments. Each fluctuation architecture

is simulated by generating a sequence of event intervals on a timeline. Events, denoted by the tick

marks, may represent ecological disturbances or winters that bring harsh conditions for biological

activity. A) A stochastic environment is generated by drawing a sequence of intervals from a

random probability distribution. B) A positively autoregressive environment is generated with

positive linear dependency of subsequent intervals on immediately preceding ones (an AR(1)

process), with random error; C) the negatively autoregressive environment is only different from

B in that the linear dependency on the past interval is negative. D) A Noisy Clock environment

assumes a series of unseen "due times" that are perfectly periodic (dashed blue tick marks), but

653   the actual events occur as a result of error around those due times (blue shaded regions); the gaps

654   between realized events form the sequence of realized intervals.

Figure 2. The interaction of environmental fluctuation architecture and nonlinearity in a vital rate creates divergence in fitness estimates. A) Diagrammatic representation of the four architectures of fluctuation, through which the demographic dynamics proceed. B) Nonlinear response function of a vital rate relative to time since last event. Here, juvenile survival rate (S_J) increases nonlinearly as the length of the interval between subsequent disturbance events increases. Dashed lines indicate the means of each axis, held constant in all modelled environments. C) Time-averaged fitness estimate*s* ($log_e\lambda_s$) are driven apart by the generating functions of environmental fluctuations as the amount of noise, or unpredictability, in the environment increases (modelled with variance of event intervals). Each colored point depicts a single computation of $log\lambda_s$ after 1000 event intervals, corresponding to a given fluctuating environment and a given σ^2, with points

666   corresponding to the other three simulated environments shown in gray for comparison. The

667   thicker lines correspond to the means of 500 iterations for each environment (architecture $\times$ noise

668   level). The divergence in fitness estimates is largely consistent over the iterations.

Figure 3. Mechanisms of fitness divergence between the architectures of fluctuating environments. A) Despite analytically equivalent distributions of event intervals (Table 1), environments have very different distributions of consecutive runs to one side of (below or above) the mean interval depending on the nature of the generating function. Negatively autoregressive and Noisy Clock environments inherently exhibit compensatory or self-correcting behavior, resulting in shorter tails in the consecutive run distribution. The opposite is true of the positively autoregressive case. The stochastic environment exhibits an intermediate exponential distribution. B) Jensen's Inequality exists in any convex/cave fluctuating variable, and introduces asymmetry between environmental and demographic fluctuations. Nonlinear functions in vital rates with respect to intervals between ecological events such as disturbances are likely common.

Figure 4. Demonstration of the effect of Jensen's Inequality on fitness divergence across architectures of environmental fluctuation. A) We morph the response function of juvenile survival S_J to manipulate its nonlinear shape around the mean event interval (dashed line), which is held constant. Going from left to right, the response function is more convex to more concave around the mean interval, highlighted in the yellow circles. This flips the sign of Jensen's Inequality, so that the overall effect of environmental fluctuations on fitness would generally become more positive to negative. B) The corresponding fitness plots demonstrate the sign flip in the effect of Jensen's Inequality as a result of the convex/cavity around the mean. Thick lines are means over all iterations, and bands are ± 1 sd. Colors correspond to the four architectures in previous figures. The middle panel is equivalent to a composite of Fig. 2C.

691 **Tables**

	Generating Function for interval τ_i	Expected Value	Variance	Generating function parameters in terms of desired sample μ_s and σ_s^2
I	$\tau_i \sim Lognormal(\mu, \sigma^2)$	$E[\tau]$ $= e^{\mu + \frac{1}{2}\sigma^2}$	$Var[\tau]$ $= (e^{\sigma^2} - 1)(e^{2\mu + \sigma^2})$	$\mu = log_e\left(\frac{\mu_s^2}{\sqrt{\mu_s^2 + \sigma_s^2}}\right)$ $\sigma^2 = log_e\left(1 + \frac{\sigma_s^2}{\mu_s^2}\right)$
II	$\tau_i = \mu + \varphi\tau_{i-1} + \varepsilon_i,$ where $\varepsilon_i \sim N(0, \sigma_\varepsilon^2)$	$E[\tau] = \mu$	$Var[\tau] = \frac{\sigma_\varepsilon^2}{1 - \varphi^2}$	$\mu = \mu_s$ $\sigma_\varepsilon^2 = \sigma_s^2(1 - \varphi^2)$
III	$\tau_i = \mu - \varphi\tau_{i-1} + \varepsilon_i,$ where $\varepsilon_i \sim N(0, \sigma_\varepsilon^2)$	$E[\tau] = \mu$	$Var[\tau] = \frac{\sigma_\varepsilon^2}{1 - \varphi^2}$	$\mu = \mu_s$ $\sigma_\varepsilon^2 = \sigma_s^2(1 - \varphi^2)$
IV	$\tau_i = [\mu(i + 1) + \varepsilon_{i+1}] - [\mu i + \varepsilon_i],$ where $\varepsilon_i \sim N(0, \sigma_\varepsilon^2)$ and $i \in \mathbb{Z}_{>0}$	$E[\tau] = \mu$	$Var[\tau] = \sigma_\varepsilon^2 + \sigma_\varepsilon^2$	$\mu = \mu_s$ $\sigma_\varepsilon^2 = \frac{1}{2}\sigma_s^2$

692 **Table 1.** Expressions of parametric moments of the four generating functions of environmental

693 fluctuations, and the derivations for input moments in terms of the desired (imposed) sample mean

694 and variance to make all distributions analytically equivalent across the four generating functions.