A natural variation-based screen in mouse cells reveals USF2 as a

regulator of the DNA damage response and cellular senescence

Taekyu Kang[1,2], Emily C. Moore[3], Emily E. K. Kopania[3], Christina D. King[1], Birgit

Schilling[1], Judith Campisi[1], Jeffrey M. Good[3], and Rachel B. Brem[1,2*]

[1]Buck Institute for Research on Aging, Novato, CA; [2]Department of Plant and Microbial Biology,

UC Berkeley, Berkeley, CA; [3]Division of Biological Sciences, University of Montana, Missoula,

MT

*Correspondence: rbrem@berkeley.edu

Abstract

Cellular senescence is a program of cell cycle arrest, apoptosis resistance, and cytokine release induced by stress exposure in mammalian cells. Landmark studies in laboratory mice have characterized a number of master senescence regulators, including p16^{INK4a}, p21, NF-kB, p53, and C/EBPβ. To discover other molecular players in senescence, we developed a screening approach to harness the evolutionary divergence between mouse species. We found that primary cells from the Mediterranean mouse *Mus spretus,* when treated with DNA damage to induce senescence, produced less cytokine and had less-active lysosomes than cells from laboratory *M. musculus.* We used allele-specific expression profiling to catalog senescence-dependent *cis*-regulatory variation between the species at thousands of genes. We then tested for correlation between these expression changes and interspecies sequence variants in the binding sites of transcription factors. Among the emergent candidate senescence regulators, we chose a little-studied cell cycle factor, USF2, for molecular validation. In acute irradiation experiments, cells lacking USF2 exhibited compromised DNA damage repair and response. Longer-term senescent cultures without USF2 mounted an exaggerated senescence regulatory program—shutting down cell cycle and DNA repair pathways, and turning up cytokine expression, more avidly than wild-type. We interpret these findings under a model of pro-repair, anti-senescence regulatory function by USF2. Our study affords new insights into the mechanisms by which cells commit to senescence, and serves as a validated proof of concept for natural variation-based regulator screens.

Introduction

Mammalian cells of many types, upon exposure to stress, can enter a senescence program, in which they stop dividing, become refractory to apoptosis, and release soluble inflammation and tissue remodeling factors termed the senescence-associated secretory phenotype (SASP) (1–4). The resulting acute immune response can clear debris, promote wound healing, and/or suppress tumorigenesis (5–8). However, during aging, senescent cells can remain long past any initial triggering event, resulting in chronic inflammation that damages the surrounding tissue (5,9–12). Landmark work has revealed the benefits of eliminating senescent cells to treat age-related pathologies and boost median lifespan (7,8,13,14).

Establishment of the senescent state and the activity of senescent cells hinge in large part on gene regulatory events. Finding molecular players that control this process is an active area of research. Now-classic work has implicated p16[INK4a] and p21 in the repression of pro-cell cycle genes and promotion of growth arrest (15) after DNA damage, and NF-kB, p53, and C/EBPβ (16) as regulators of the SASP. However, given the complexity of the senescence program, many more regulators likely remain to be identified. Indeed, bioinformatic approaches have identified dozens of other transcription factor candidates in senescence (17–24), many of which remain unvalidated (but see (18–21) for recent discoveries of the roles of DLX2, FOXO3 and AP-1 in senescence).

We set out to develop a novel approach to survey transcription factors that play a role in cellular senescence, with the potential for increased specificity relative to traditional genomic screens. Our strategy took advantage of the natural genetic variation in senescence gene expression, and transcription factor binding sites, across mouse species. Among the top hits from this

analysis, we chose the under-studied factor USF2 for validation experiments, focusing on gene

regulation and cellular phenotypes during senescence induction and maintenance.

Results

A dampened senescence program in primary fibroblasts of *Mus spretus*

We initiated our study on the premise that senescent phenotypes could differ among mouse

lineages, and that these changes could serve as a tool to uncover novel regulators. Toward this

end, we first characterized the dependence of senescent phenotypes on genotype. We isolated

primary tail fibroblasts from PWK, a wild-derived purebred strain of *Mus musculus,* and from the

wild-derived, purebred STF strain of the Mediterranean mouse species *M. spretus*. We treated

each with ionizing radiation, incubated to allow development of senescence (3,25,26), and

assessed features of the resulting senescent cells, with untreated cells as controls. Such

cultures from both species arrested growth and exhibited a flattened and enlarged morphology

(Figure 1A), as expected if both cultures had entered the senescent state (26). By contrast,

assays of senescence-associated β-galactosidase activity, a senescence biomarker (27),

revealed a marked difference between the cultures, with two to eight-fold lower staining in *M.*

spretus cells relative to *M. musculus* cells (Figure 1B).

To explore further the divergence in senescent phenotypes between cells of our focal species,

we profiled bulk RNA levels in senescent and control primary fibroblast cultures from each

species (Supplemental Table S1). We inspected genes of the SASP immune-stimulatory

program (4), and found that this gene cohort was induced more highly in *M. musculus* cells than

in those of *M. spretus* during senescence (Figure 2A). Likewise, in an unbiased search of Gene

Ontology terms, we identified several suites of immune response and NF-κB signaling genes

that were enriched for senescence-specific differential expression between the cultures. In

each case, the gene groups were more strongly induced during senescence in *M. musculus*

cells than in *M. spretus* cultures (Figure 2B and Supplemental Tables S2 and S3); among

members of the latter we noted *Cxcl1* (28), *Il6* (29), *Ccls* 2,7 and 8 (30), *Mmp13* (31), and other

reported SASP genes (Supplemental Figure S1). These data make clear that, at the level of

mRNA, the senescence regulatory program differs markedly between fibroblasts of our focal

species.

We hypothesized that much of the divergence between *M. musculus* and *M. spretus* cells during

senescence, which we detected in RNA measurements, would result in differential protein

abundance. In pursuing this notion, we focused on proteins secreted into the medium by

senescent cells, owing to the physiological importance of the SASP (30). We collected

conditioned media from senescent and control cultures of primary fibroblasts of each genotype,

and we used it as input into unbiased mass spectrometry to quantify protein abundance. The

results revealed an overall increase in the secretion of common SASP components, including

CCL chemokines, matrix metalloproteases, and serpins, in senescent *M. musculus* cells relative

to those of *M. spretus* (Figure 2C, Supplemental Figure S2, and Supplemental Table S4).

Together, our characterization experiments show that *M. spretus* fibroblasts enter a senescent

growth arrest similar to that of *M. musculus* fibroblasts, but with expression of certain SASP

factors spiking less highly.

A genomic screen for senescence transcription factors using *cis*-regulatory sequence

variations

Having established divergence between *M. musculus* and *M. spretus* primary fibroblasts in

senescence mRNA and protein secretion (Figures 1 and 2), we reasoned that such differences

could be harnessed in an *in silico* screen for senescence regulators. We designed an analysis

focused on gene regulation—in particular, on variation between the species at the binding sites

of transcription factors (Figure 3A). We expected that, at some genes, *cis*-regulatory elements

encoded in the *M. musculus* genome would drive expression during senescence differently than

those in the *M. spretus* genome. We reasoned that if *cis*-acting variation effects were enriched

among the loci bound by a given transcription factor across the genome, the signal could be

interpreted as a signpost for the factor's activity during senescence (Figure 3A). In this way, all

cis-regulatory variants between the species that manifested in cultured primary fibroblasts,

whether of large or small effect size, and regardless of their potential for phenotypic impact,

could contribute to the search for transcription factors relevant for senescence.

As a resource for this approach, we mated PWK *M. musculus* and STF *M. spretus* to yield F1

hybrid animals from which we derived primary tail fibroblasts for culture and irradiation. These

M. musculus x *M. spretus* F1 hybrid fibroblasts, when irradiated, exhibited a flattened

morphology reflecting entry into senescence; senescence-associated β-galactosidase activity

was of a magnitude between those of purebred *M. musculus* and *M. spretus* fibroblasts upon

irradiation (Supplemental Figure S3). We subjected senescent and control F1 hybrid fibroblasts

to RNA-seq profiling, and we used the results to quantify levels of transcripts derived from the

M. musculus and *M. spretus* alleles of each gene in each condition (Supplemental Table S1).

At a given gene, any difference between allele-specific expression in an F1 hybrid can be

attributed to variants inherited from the parent species that perturb gene regulation in *cis* at the

locus, because *trans*-acting factors impinge to the same extent on both alleles (32). Analyzing

the response to senescence induction for a given gene, we found that the allele-specific

expression difference between the alleles in the F1 hybrid was a partial predictor of the

expression divergence between the *M. musculus* and *M. spretus* purebreds, in our primary cell

system (Supplemental Figure S4). The latter trend reflects the joint contributions of *cis-* and

trans-acting variants to total expression divergence between the species, as expected (33).

Separately, to survey overall regulatory programs in F1 hybrid primary fibroblasts, we

formulated the expression level of a given gene in a given condition as the sum of the measured

levels of the *M. musculus* and *M. spretus* alleles. In this analysis, focusing on SASP genes as

we had done for the purebreds (Figure 2A), we found that the expression program of senescent

F1 hybrid cells was, for most genes, intermediate between the low levels seen in *M. spretus*

cells and the high levels in *M. musculus* (Supplemental Figure S5). These data indicate that *M.*

musculus x *M. spretus* F1 hybrid fibroblasts do not exhibit heterosis with respect to senescence-

associated genes, and do manifest extensive, senescence-dependent *cis*-regulatory variation.

We next used the expression measurements from F1 hybrid cells as input into our *in silico*

screen to identify senescence-dependent transcription factor activity. For a given transcription

factor, we collated binding sites detected by chromatin immunoprecipitation upstream of genes

across a panel of tissues (34). At each site, we tabulated the presence or absence of DNA

sequence variants in the respective genomes of *M. musculus* and *M. spretus*. We then tested

whether, across the genome, genes with these binding site variants were enriched for

senescence-associated expression differences between the two alleles in the F1 hybrid. This

test had the capacity for high power to detect even subtle contributions from transcription factors

if they had deep binding-site coverage in the input data; five factors attained genome-wide

significance (Figure 3B and Supplemental Table S5). Among them, PBX1 (35) and CREBBP

(36,37) were previously implicated in cellular senescence, providing a first line of evidence for

the strength of our approach to identify signatures of condition-dependent transcription factor

function. The top-scoring transcription factor in our screen, a basic-helix-loop-helix leucine-

zipper protein called upstream stimulatory factor 2 (USF2; Figure 3B-C), had not been

experimentally characterized in stress response or senescence. However, classic studies had

established USF2 as a regulator of the cell cycle and tumor suppression (38–41). More

recently, USF2 was shown to control cytokine release in immune cells (42). Considering these

known functions, and bioinformatic analyses suggesting a link between USF2 and senescence

programs (21), we chose USF2 for in-depth validation. In detailed genomic tests, single

variants between *M. musculus* and *M. spretus* at USF2 binding sites drove most of the

relationship with allele-specific expression in hybrid senescent cells (Figure 3D). These variants

were over-represented at positions central to, and slightly downstream of, experimentally

determined peaks for USF2 (Figure 3E), highlighting the likely importance of this region in

USF2's mechanisms of binding and regulation.

USF2 modulates cell proliferation and the acute DNA damage response

Our question at this point was whether and how USF2 regulated senescence programs. As

such, we shifted our focus from natural genetic variation to controlled, laboratory-induced

genetic perturbations in a single genetic background. We designed two short hairpin RNAs

(shRNAs) targeting *Usf2*, each in a lentiviral vector under the U6 promoter. Expression

measurements upon transformation of PWK *M. musculus* primary tail fibroblasts confirmed 2.5

and 3-fold knockdown of *Usf2* expression, respectively, from these shRNAs (Supplemental

Figure S6). We developed two experimental workflows using these shRNAs. One approach

aimed to define the role of USF2 in the acute DNA damage response (Figure 4A) and

establishment of senescence (Figure 5A): we infected cells with *Usf2* shRNAs, cultured them

under standard conditions, and then induced senescence by irradiation. In a second approach,

we focused on the function of USF2 in maintenance of senescence, inducing senescence in

wild-type cells and knocking down *Usf2* afterward (Figure 5C).

Given that our first experimental design used *Usf2* knockdown in cells under standard

conditions to study the irradiation response and senescence (Figures 4A and 5A), we began by

characterizing the effects of reduced *Usf2* expression in resting cells. Uptake of the nucleotide

analog EdU, a marker of DNA synthesis, was reduced by 40% in resting primary fibroblasts

expressing the *Usf2* shRNA relative to scrambled shRNA controls (Supplemental Figure S7).

Transcriptional profiling of these resting cultures (Supplemental Table S6), followed by

functional-genomic (Gene Ontology) analyses, revealed enrichment for biosynthesis,

development, cell growth, and DNA repair terms altered upon *Usf2* knockdown relative to

controls (Supplemental Table S7). These data support a role for USF2 in wild-type growth and

housekeeping gene expression programs in untreated primary fibroblasts, consistent with

studies of USF2 in cell growth in other tissues and contexts (26,38–41). Our results in resting

cells also provide a first indication of the potential function of USF2 in the DNA damage

response.

We next directly pursued the importance of USF2 during the acute response to irradiation

(Figure 4A). DNA damage signaling, an inducer of cellular senescence (2,3), drops sharply in

intensity within eight hours and then more gradually over several day after irradiation (43),

culminating in a lower persistent signal (44–47). We focused on the early phase of this process

(six hours after irradiation) in cultures of primary fibroblasts expressing *Usf2* shRNAs or

scrambled shRNA controls. Transcriptional profiling followed by Gene Ontology analyses

identified gene groups enriched for expression changes dependent on condition and USF2

(Figure 4B and Supplemental Tables S6 and S8). Most salient was a trend of pervasive

repression transcriptome-wide six hours after irradiation, which was detected in control cells as

expected (48,49), and was blunted in cells with *Usf2* knocked down. The latter effect was

particularly enriched in the transcriptional machinery, repressors of apoptosis, and several cell

proliferation regulators (Figure 4B-C and Supplemental Figure S8A). We also inspected the few

top-scoring gene groups that were induced by irradiation in control cultures. These gene groups

reached even higher expression levels in *Usf2* knockdown cells; notable among them was the

EGFR pathway (Figure 4B and 4D and Supplemental Figure S8B), a determinant of entry into

senescence in some contexts (50). DNA repair genes, a likely target for changes upon

irradiation, are regulated primarily at the post-transcriptional level (51,52), and we did not detect

effects of *Usf2* knockdown on their transcripts (Supplemental Figure S9).

We hypothesized that *Usf2* knockdown during the acute DNA damage response would also

have cell-physiological effects. Assays of EdU incorporation to report on DNA synthesis

showed effects of *Usf2* knockdown after irradiation to the same degree as in resting cultures

(Supplemental Figure S7). To focus on phenotypes more proximal to DNA damage, we used

the neutral comet assay (53) to measure DNA double-stranded breaks on a per-cell basis. In

this setup, *Usf2* depletion increased comet tail moments by 50% six hours after irradiation, with

an effect that was similar, though of smaller magnitude, in resting cell controls (Figure 4E).

Next, we tracked foci of phosphorylated histone H2AX (γH2AX) in fibroblasts as a marker of

chromatin decondensation, preceding the repair of DNA double-stranded breaks (54). Cells

harboring *Usf2* shRNAs exhibited 30% fewer γH2AX foci than cells of the control genotype, six

hours after irradiation (Figure 4F). These data establish a role for USF2 in the response to

irradiation, with knockdown of this factor compromising cells' ability to mount the classical

transcriptional program under this stress, and to carry out DNA damage repair.

USF2 tunes the commitment to senescence

Having knocked down *Usf2* in naïve PWK primary fibroblasts and irradiated them to study the

acute DNA damage response, we now allowed the irradiated cultures to enter senescence

(Figure 5A). Growth arrest and flattened morphology were indistinguishable between these

cells and controls harboring scrambled shRNAs (see Figure 5G), indicating that wild-type levels

of *Usf2* were not required at the point of irradiation to establish senescence *per se*. To

investigate quantitative characteristics of these senescent cultures, we subjected them to

expression profiling and Gene Ontology enrichment analyses (Supplemental Tables S6 and

S9). Among top-scoring gene groups, the most dramatic effects were in those that dropped in

expression in senescent cultures of the control genotype, which as expected (17,55) included

cell cycle and DNA repair pathways (Figure 5B and Supplemental Table S9). mRNA levels of

the latter were even lower in senescent cells that had been irradiated after *Usf2* knockdown,

showing a reduction of ~20% on average (Figure 5B). Among the genes of this cohort, some of

which declined in expression by >5-fold with *Usf2* knockdown in senescence (Figure 5F), we

noted cell cycle regulators (*Ccna2*, *Cdc20*, *Cdk1*), kinesin components (*Kif2c, Knl1*), the DNA

polymerase *Pole,* and the DNA damage checkpoint ubiquitin ligase *Uhrf1* (Supplemental Figure

S10A). We conclude that genes of the cell cycle and DNA repair machinery are detectable at a

low but non-zero expression level in wild-type senescent cells, and that these pathways are

subject to further reduction when *Usf2* is limiting.

Next, we inspected the impact of *Usf2* knockdown on inflammation and immune recruiting

factors, another set of processes that had emerged from our Gene Ontology analyses of

expression (Supplemental Table S9). Cells of the control genotype induced these pathways

during senescence, as expected (2,3,9,29,56,57); in cells reaching senescence after *Usf2*

knockdown, induction of inflammatory factors was amplified by ~10% on average (Figures 5B

and 5E, and Supplemental Figure S10B). Similarly, in assays of the β-galactosidase

senescence marker, we observed a modest, though robustly significant, increase in cells

subject to *Usf2* knockdown and senescence treatment (Figure 5G). Together, our profiling data

establish that irradiation of primary fibroblasts with reduced *Usf2* expression leads to a

quantitatively perturbed, exaggerated senescent state, with reduced expression of proliferation

and DNA repair pathways, and elevated pro-inflammatory gene expression and β-galactosidase

activity.

We reasoned that the changes in senescence we had seen upon irradiation of *Usf2*-depleted

cells could constitute, in part, effects from increased DNA damage in the knockdown genotype

(Figure 4). To pursue the role of USF2 in senescence more directly, we used our second

experimental paradigm: we irradiated wild-type cells, incubated them for 10 days to allow

establishment of senescence, then expressed shRNAs targeting *Usf2,* followed by 10 additional

days of incubation and transcriptional profiling (Figure 5C). In control cells (harboring a

scrambled shRNA), we observed expression changes for many genes from the resting state

through irradiation and early and late senescence (Supplemental Tables S6 and S10, and

Supplemental Figure S11), attesting to the dynamics of senescence across this timecourse as

expected (17,55). Next we focused on cultures subject to *Usf2* knockdown during late

senescence. The latter manifested quantitative changes largely paralleling those in cells with

Usf2 knocked down at the start of the experiment. Thus, on average, cell cycle and DNA repair

genes, repressed in control cells during senescence, were expressed at even lower levels when

Usf2 was knocked down midway through the senescence time course (Figures 5D and 5F,

Supplemental Tables S6 and S11, and Supplemental Figure S10A), analogous to our findings

upon early knockdown of *Usf2*. Likewise, inflammatory response factors, induced during

senescence in the control setting, were more highly expressed when *Usf2* was depleted after

senescence establishment, as seen in our early-knockdown design (Figures 5D and 5E,

Supplemental Tables S6 and S11, and Supplemental Figure S10B). We noted a handful of

genes for which early *Usf2* knockdown effects were not recapitulated in our paradigm of

knockdown after senescence entry (*e.g. Adamts1* and *Rarres2* in Supplemental Figure S10B).

Overall, however, our analyses establish USF2 as a senescence regulator at least in part

independent of its role in the acute DNA damage response, such that in its absence, cells

commit even more strongly to the senescent state.

Discussion

Complex regulatory networks likely underlie many of the quantitative behaviors of senescent

cells, including kinetics and dependence on cell type and inducer (15,17,25,58,59). Exactly how

these nuances are encoded remains poorly understood. In this study, we pioneered the use of

interspecies genetic divergence to screen for components of the senescence regulatory

machinery. This strategy complements previous studies of transcription factor binding sites in

genes of the senescence program in a single genetic background (17,21). Our approach

harnesses the correlation between interspecies variation in sequence and expression levels, as

an additional line of evidence for senescence-specific regulatory functions by a given factor.

This paradigm parallels similar tools previously used to dissect divergence in expression (60,61)

and transcription factor binding (62,63) in other contexts. Broadly speaking, these methods are

not highly powered for pathways under strong evolutionary constraint, which, by definition, will

not vary enough among species to yield the raw observations that would go into a screening

pipeline. Rather, we expect the natural variation-based approach to work best for discovering

less-constrained modifiers, many of which may confer layers of quantitative regulation onto a

master regulatory pathway.

We focused our experimental validation on one such modifier, the transcription factor USF2. By

tracing USF2's function in proliferation and genome-wide expression in untreated cells, we

extended conclusions from studies of USF2 in tumor suppression and cell cycle regulation (38–

40,42), apoptosis (64) and ERK1/2 signaling (65). In an acute DNA damage setting, we

discovered that USF2 is required for cells to mount DNA repair and downstream DNA damage

responses. And in senescence proper, we showed that USF2 acts as a repressor, such that in

its absence, the senescence program—shutoff of cell proliferation and DNA repair, and

induction of cytokines—is amplified. A compelling model is thus that even long after damage

exposure, cells have access to expression states along a continuum of commitment to

senescence, and that USF2 acts to help determine which state they occupy. If so, USF2 would

take a place among a network of factors, including p53, ING, Rb (66,67), p21 (68,69), and p16

(70), that govern the choice between senescence, apoptosis, and repair and proliferation,

depending on cell type (66) and the amount of damage or stress incurred (67).

As a corollary of these conclusions from expression profiling, we note that cell cycle and DNA

repair genes, classically known to be repressed during senescence (17,55), did not hit a floor of

expression in senescent cultures: we could detect them at even lower expression levels upon

Usf2 knockdown. Since our cultures comprise >99% arrested cells within several days of

irradiation (see Methods), the emerging picture is that the proliferation machinery is maintained

at non-zero levels even in such a population. Any ability of these gene products to reattain

activity could be of particular interest as a potential mediator of the return to proliferation seen

among senescent cells in certain scenarios (71,72).

Our work leaves open the mechanisms by which USF2 exerts its effects in the DNA damage

response and cellular senescence. It is tempting to speculate that USF2 ultimately works in

these processes in concert with its better-studied family member, USF1. Indeed, USF1 has

been implicated in DNA repair (73), inflammation (74,75), immune responses (76), and p53-

mediated cell cycle arrest (77) in contexts other than senescence. In addition, given that USF2

has been implicated in the TGFβ-p53 axis in apoptosis (64) and fibrosis (78), the latter pathway

could mediate some part of the USF2 effects we have seen. Furthermore, regulatory network

reconstruction (Supplemental Table S12) suggests that USF2 acts upstream of several other

transcription factors (KLF3, GLI3, NFIL3) with direct targets in DNA repair, DNA damage

response and senescence pathways.

Alongside our use of *cis*-regulatory variation between mouse species as a screening tool for

senescence genes, we also characterized overall patterns of divergence between *M. spretus*

senescent cells and those of *M. musculus.* Given that the former exhibited lower levels of

SASP mRNAs and proteins, we suggest that the rheostat of the senescence response is at a

higher set point in this species*,* such that at a given level of stress (*e.g.* the irradiation we study

here), cells of this species synthesize and secrete less of the SASP. We further hypothesize

that this down-tuning might be a proximal cause for the lower senescence-associated β-

galactosidase activity we have seen in *M. spretus* cells. Such a link would follow from current

models of the senescent state in which production and secretion of SASP components (79)

leads to proteotoxic stress from insoluble aggregates (80,81), an increase in the number and

size of lysosomes (82), and enhanced β-galactosidase activity (83). Plausibly, any of the

phenotypes we study here in cell culture could have consequences *in vivo*, with potential links to

the stress- and pathogen-resistance phenotypes characterized in *M. spretus* (84–88). Together,

our findings attest to the power of natural variation-based-approaches—to survey the

differences that evolution has wrought over long timescales, and as a tool for molecular

discoveries of relevance in any one genetic background.

Methods

Primary cell extraction and culture

Wild-derived, inbred lines of *Mus musculus* (PWK/PhJ) and *Mus spretus* (STF/Pas) and their

interspecies F1 hybrids (*M. musculus* x *M. spretus*) were maintained in standard conditions

under Montana Institutional Animal Care and Use Committee protocol number 062-1JGDBS-

120418. Animals aged 3 to 5 months were humanely euthanized prior to tissue collection via

CO_2 treatment and cervical dislocation. Five tails from each genotype were collected into chilled

Dulbecco's Modified Eagle Medium (DMEM) and shipped to the Buck Institute for further

processing. Metadata for the mice used in this experiment is provided in (Supplemental Table

S13). Primary tail fibroblasts were extracted from the cuttings essentially as described (89).

Briefly, tail cuttings first were soaked in 70% ethanol for 5 minutes and then laid out in sterile 10

cm tissue culture dishes in a biosafety cabinet for 5 minutes to dry. The tails were then

transferred to a 10-cm dish containing 10 mL of complete medium (RPMI 1640 medium, 10%

fetal bovine serum, 50 µM 2-mercaptoethanol, 100 µM asparagine, 2 mM glutamine, and 1%

penicillin-streptomycin). Hair was removed from the tails using sterile forceps and razor blade,

and the tails were cut into 2-3 mm pieces using sterile surgical scissors. These pieces were

then transferred into a 2 mL cryotube containing a collagenase D–pronase solution and left in a

shaking incubator at 37°C for 90 minutes. Following incubation, the contents of the cryotubes

were placed into a 70 µm cell strainer in a new 10-cm dish containing 10 mL of compete

medium. Using the back end of a sterile 10 mL syringe plunger, the tissue was ground into the

strainer for 5-10 minutes to release the cells into the media. The cell suspension was then

collected into 15 mL conical tubes and centrifuged at 580 x g and 4°C for 5 minutes. The

supernatant was removed and replaced with fresh complete medium, and the centrifugation was

repeated for two additional rounds. After the final spin, the media was replaced with complete

medium supplemented with 250 ng/mL of amphotericin B. The cell suspension was placed in a

10-cm dish and incubated in a 37°C humidified incubator at 3% O_2 and 10% CO_2 for two days

before passage and continued culture in complete medium (DMEM, 10% FBS, 1% penicillin-

streptomycin). This complete medium was used for the remainder of our experiments. For long

term storage, cells were suspended in 5% DMSO in FBS, aliquoted into cryotubes, and placed

in a slow cooling container in a -80°C freezer overnight. The following day, the cryotubes were

moved out of the slow cooling container and placed into long term containers in the freezer. To

thaw cells for continued culture, cryotubes containing frozen aliquots were swirled in a 37°C

water bath until only a thin layer of ice remained in the vial. The contents of the cryotube were

then immediate transferred into a flask containing pre-warmed complete medium and incubated

overnight. The following day the cells were washed twice with PBS to remove any trace of

DMSO and cultured in complete medium. For experiments in wild-type *M. musculus*, *M.*

spretus, and F1 hybrid cells we considered the culture from each individual animal to represent

one biological replicate of the respective genotype. In knockdown experiments (see below) we

carried out infection of two such cultures with each shRNA-bearing virus, which we refer to as

biological replicates. In either setup, we refer to the split cultures from a given biological

replicate as technical replicates in a given assay.

Irradiation treatment

To treat a given cell culture replicate with ionizing radiation (3,25,26) for a cell-biological assay

or omics profiling, we proceeded as follows. The day before irradiation, cells were seeded at

60-70% confluency and incubated in a 37°C humidified incubator at 3% O_2 and 10% CO_2

overnight in complete medium. The next day, a subset of cells was collected and used as input

into the respective experiment as the unirradiated control. The remainder of the culture was

transferred into an X-RAD 320 X-Ray Biological Irradiator and treated with 15 Gy of X-ray

irradiation. Cultures were then placed back into the 37°C humidified incubator at 3% O_2 and

10% CO_2 until sampling at 6 hours for marker assays and RNA-seq focused on acute DNA

damage response, or 7, 10 or 20 days for marker assays and RNA-seq and proteomics focused

on senescence (in which case the medium was replaced 6-8 hours after irradiation, then every

48 hours for the remainder of the experiment), as detailed below.

Senescence marker assays

For a given replicate culture after irradiation and incubation (see above), senescence-

associated β-galactosidase activity was measured using the BioVision Inc. Senescence

Detection Kit (cat. #K320): cells were fixed, permeabilized, and incubated with the staining

solution containing X-gal overnight. Multiple images were taken the following day using a

brightfield microscope, and the image names were randomized before the proportion of β-

galactosidase-positive cells was counted manually to remove potential sources of bias.

Cultures were considered to be senescent if they showed less than 10% EdU incorporation (see

below) and over 90% β-galactosidase-positive cells. In the species comparison of Figure 1 we

subjected two technical replicate cultures of each of three biological replicates per purebred

species to irradiation (see above) followed by β-galactosidase assays at the indicated

timepoints. In Supplemental Figure S3, data for purebreds were from Figure 1 at day 7;

separately, for the interspecies F1 hybrid, we carried out irradiation and β-galactosidase assays

as above for two technical replicates from one biological replicate. In Figure 5G, we carried out

irradiation and β-galactosidase assays as above at day 7 after irradiation for two technical

replicates from each of two biological replicates of *M. musculus* cells infected with lentivirus

harboring the scrambled control and two of each *Usf2* knockdown (see below).

RNA collection and sequencing

For a given replicate culture of a given genotype, either before irradiation or 6 hours, 10 days, or

20 days after irradiation (see above), cells were treated with TRIzol™ and RNA was extracted

using chloroform and ethanol precipitation. The RNA was then further purified using the Qiagen

RNeasy Kit (cat. #74004) for DNase treatment and column cleanup. The purity of the extracted

RNA was verified using a NanoDrop ND-1000 Spectrophotometer; all samples had 260/280 and

260/230 ratios greater than 2.0. Purified RNA in distilled RNase/DNase free water was snap

frozen using dry ice and stored at -80°C. Samples were then either transferred to the QB3

Genomics core at University of California, Berkeley for library prep and sequencing on 150PE

NovaSeq S4 or shipped to Novogene Co. (Sacramento, CA) for the same. Both facilities

provided 25 million paired end reads per sample. For expression profiles of purebred cells and

M. musculus x *M. spretus* F1 hybrid cells, we subjected three biological replicates of a given

genotype to irradiation (see above) followed by RNA-seq at the indicated timepoints. To assess

transcriptional impacts of *Usf2* knockdown we carried out irradiation and RNA isolation as above

for two biological replicates of *M. musculus* cells infected with lentivirus harboring the scrambled

control and two of each *Usf2* knockdown (see below).

Pseudogenome and VCF generation

As publicly available annotations for *M. musculus* and *M. spretus* are in the context of their

reference genomes (GRCm38.96 and SPRET_EiJ_v1.96 respectively), custom

pseudogenomes for strains PWK and STF were generated and used for this study. For the

PWK pseudogenome, variant calls between PWK and the reference genome in the form of a

variant call file (VCF) were downloaded from the Sanger Mouse Genomes Project database

(https://www.sanger.ac.uk/data/mouse-genomes-project/). A pseudogenome using the VCF

and the GRCm38.96 reference genome was created using bcftools v1.9 (90). To generate

shotgun sequence data for the STF pseudogenome, DNA was extracted from *M. spretus* liver

tissue using Qiagen DNeasy spin columns (cat. #69506). The sample was sheared via

sonication (Covaris E220), and prepared using the New England Biolabs NEBNext Ultra DNA

Library kit (cat. #E7370L). The final library was sequenced on a single lane of 150bp PE

Illumina HiSeq X at Novogene, Inc. The latter reads were aligned to the SPRET_EiJ_v1.96

reference genome using bowtie v2.2.3 (91), and a VCF was generated using bfctools mpileup

and filtered for quality, depth, and SNPs using vcfutils (92,93). The VCF was then used with the

reference genome to create a STF pseudogenome using bcftools. The pseudogenomes were

verified by identifying a number of called variants by hand. A VCF of variants between the STF

and PWK pseudogenomes was generated as above aligning the STF whole genome

sequencing reads to the PWK pseudogenome.

RNA-seq processing

In transcriptional profiles of *M. musculus* x *M. spretus* F1 hybrid cells, sequencing reads from a

given replicate were aligned to a concatenated PWK/STF pseudogenome using tophat v2.1.1

(94) allowing for zero mismatches to ensure allele specific mapping. Alignments were then

filtered for uniquely mapped reads using samtools v1.3.1, and gene counts were generated

using HTSeq v0.11.2 (95) and genome annotations (GRCm38.96, SPRET_EiJ_v1.96) for both

species from Ensembl. Counts were then converted to transcripts per million (TPM) using

custom R scripts, and genes were filtered for those showing counts in more than half the

samples sequenced.

In transcriptional profiles of purebred *M. musculus* wild-type cells and *M. musculus* cells infected

with lentiviruses harboring shRNAs, RNA-seq processing was as above but mapping was to the

PWK pseudogenome only; for profiles of purebred *M. spretus* wild-type cells, mapping was to

the STF pseudogenome only.

Data for the average TPM across biological replicates for purebred parents and interspecific F1

hybrid are reported in Supplemental Table S1. Average TPM across biological replicates for

shRNA-treated *M. musculus* cells are reported in Supplemental Table S6.

Gene Ontology enrichment analysis of RNA-seq data

For the comparison of transcriptional profiles of *M. musculus* and *M. spretus* purebred cells, for

each gene in turn we tabulated the average TPM count from each species across replicates,

and then took the $\log_2$ of the ratio of these averages, $r_{sen,true}$. We downloaded Gene Ontology

annotations from the AmiGO 2 (96,97) database and filtered for those with supporting biological

data. For each term, we summed the $r_{sen,true}$ values across all genes of the term, yielding

$s_{sen,true}$. To assess the enrichment for high or low values of this sum, we first took the absolute

value, $|s_{sen,true}|$. We then sampled, from the total set of genes with expression data, a random

set of the same number as that in the true data for the term; we calculated the species

difference $r_{sen,rand}$ for each such gene and the absolute value of the sum over them all, $|s_{sen,rand}|$.

We used as a *p*-value the proportion of 10,000 resampled data sets in which $|s_{sen,true}| > |s_{sen,rand}|$.

For analysis of the impact of *Usf2* knockdown on expression before or 6 hours, 10 days, or 20

days after irradiation (see below), Gene Ontology enrichment tests were as above except that

we took the ratio, for a given gene, between the average expression in purebred *M. musculus*

(PWK) cells infected with lentivirus harboring scrambled shRNA and the analogous quantity

across both *Usf2*-targeting shRNA treatments.

Proteomic analysis of secreted proteins

Conditioned medium preparation. For a given replicate culture, either before irradiation or 10

days after irradiation (see above), cells were washed three times with PBS and incubated with

serum and phenol red free DMEM containing 1% pen-strep for 24 hours. The following day the

conditioned medium was collected and passed through a 0.45 µm filter to remove cellular

debris. The conditioned medium was placed in a -80°C freezer for storage before use as input

into proteomic profiling (see below). For proteomic profiles of purebred cells, we carried out this

procedure for three technical replicate cultures of one biological replicate per species.

Sample concentration. 30 mL of conditioned media for each replicate were concentrated to 400

µL with 15 mL 3 kDa filters (Millipore Sigma, Burlington, MA). Protein concentration was

determined using the Bicinchoninic Acid (BCA) assay (Thermo Fisher Scientific, Waltham, MA).

Protein digestion and desalting. Aliquots of 200 µg protein lysates for each sample were

brought to the same overall volume of 52 µL with water, reduced using 20 mM dithiothreitol in

50 mM triethylammonium bicarbonate buffer (TEAB) at 50°C for 10 min, cooled to room

temperature (RT) and held at RT for 10 min, and alkylated using 40 mM iodoacetamide in 50

mM TEAB at RT in the dark for 30 min. Samples were acidified with 12% phosphoric acid to

obtain a final concentration of 1.2% phosphoric acid. S-Trap buffer consisting of 90% methanol

in 100 mM TEAB at pH ~7.1, was added and samples were loaded onto the S-Trap mini spin

columns. The entire sample volume was spun through the S-Trap mini spin columns at 4,000 x

g and RT, binding the proteins to the mini spin columns. Subsequently, S-Trap mini spin

columns were washed twice with S-Trap buffer at 4,000 x g at RT and placed into clean elution

tubes. Samples were incubated for one hour at 47°C with sequencing grade trypsin (Promega,

San Luis Obispo, CA) dissolved in 50 mM TEAB at a 1:25 (w/w) enzyme:protein ratio.

Afterwards, trypsin solution was added again at the same ratio, and proteins were digested

overnight at 37°C.

Peptides were sequentially eluted from mini S-Trap spin columns with 50 mM TEAB, 0.5%

formic acid (FA) in water, and 50% acetonitrile (ACN) in 0.5% FA. After centrifugal evaporation,

samples were resuspended in 0.2% FA in water and desalted with Oasis 10 mg Sorbent

Cartridges (Waters, Milford, MA). The desalted elutions were then subjected to an additional

round of centrifugal evaporation and re-suspended in 0.2% FA in water at a final concentration

of 1 µg/µL. Subsequently, 8 µL of each sample were diluted with 2% ACN in 0.1% FA to obtain

a concentration of 400 ng/µL. Finally, 1 µL of indexed Retention Time Standard (iRT,

Biognosys, Schlieren, Switzerland) was added to each sample, thus bringing up the total

volume to 20 µL (98).

Mass spectrometric analysis. Reverse-phase HPLC-MS/MS analyses were performed on a

Dionex UltiMate 3000 system coupled online to an Orbitrap Exploris 480 mass spectrometer

(Thermo Fisher Scientific, Bremen, Germany). The solvent system consisted of 2% ACN, 0.1%

FA in water (solvent A) and 80% ACN, 0.1% FA in ACN (solvent B). Digested peptides (400 ng)

were loaded onto an Acclaim PepMap 100 C_{18} trap column (0.1 x 20 mm, 5 µm particle size;

Thermo Fisher Scientific) over 5 min at 5 µL/min with 100% solvent A. Peptides (400 ng) were

eluted on an Acclaim PepMap 100 C_{18} analytical column (75 µm x 50 cm, 3 µm particle size;

Thermo Fisher Scientific) at 300 nL/min using the following gradient: linear from 2.5% to 24.5%

of solvent B in 125 min, linear from 24.5% to 39.2% of solvent B in 40 min, up to 98% of solvent

B in 1 min, and back to 2.5% of solvent B in 1 min. The column was re-equilibrated for 30 min

with 2.5% of solvent B, and the total gradient length was 210 min. Each sample was acquired in

data-independent acquisition (DIA) mode (99–101), in triplicates. Full MS spectra were

collected at 120,000 resolution (Automatic Gain Control (AGC) target: 3e6 ions, maximum

injection time: 60 ms, 350-1,650 *m/z*), and MS2 spectra at 30,000 resolution (AGC target: 3e6

ions, maximum injection time: Auto, Normalized Collision Energy (NCE): 30, fixed first mass 200

m/z). The isolation scheme consisted of 26 variable windows covering the 350-1,650 *m/z* range

with an overlap of 1 *m/z* (100).

DIA data processing and statistical analysis. DIA data was processed in Spectronaut (version

15.6.211220.50606) using directDIA. Data extraction parameters were set as dynamic and non-

linear iRT calibration with precision iRT was selected. Data was searched against the *Mus*

musculus reference proteome with 58,430 entries (UniProtKB-TrEMBL), accessed on

01/31/2018. Trypsin/P was set as the digestion enzyme and two missed cleavages were

allowed. Cysteine carbamidomethylation was set as a fixed modification; methionine oxidation

and protein N-terminus acetylation were set as dynamic modifications. Identification was

performed using 1% precursor and protein q-value. Quantification was based on the peak

areas of extracted ion chromatograms (XICs) of 3 – 6 MS2 fragment ions, specifically b- and y-

ions, with local normalization and q-value sparse data filtering applied (Supplemental Table S4).

In addition, iRT profiling was selected. Senescence specific differential protein secretion

analysis comparing samples derived from unirradiated and senescent *M. musculus* and *M.*

spretus cells was performed using the proteomics data as input for a two-factor ANOVA with

Benjamini-Hochberg multiple testing correction (102). Only those proteins with a corrected *p-*

value < 0.05 were considered significantly differentially secreted between the species during

senescence.

Transcriptomic screen for senescence regulators

To associate expression variation in genes with sequence variation in their upstream binding

sites for a given transcription factor, we proceeded as follows. From RNA-seq profiling of *M.*

musculus x *M. spretus* F1 hybrid cells (see above), we used the TPM counts for each parent

species' allele from each replicate profile from control and senescent conditions as input into a

two-factor ANOVA. A given gene was categorized as exhibiting senescence-associated

differential allele specific expression if the interaction F statistic value from this ANOVA was

among the top 25% of all genes tested. Separately, we used compiled data from chromatin

immunoprecipitation via high-throughput sequencing from the Gene Transcription Regulation

Database (GTRD) (34) to identify all experimentally determined transcription factor (TF) binding

sites located within a 5kb window upstream of the transcriptional start site for each gene in turn

in the *M. musculus* genome; we refer to the downstream gene of each such binding location as

the target of the TF. This calculation used *M. musculus* gene start sites from the Ensembl

GRCm38.96 GFF. Next, for each binding site, we used the VCF between PWK and STF

pseudogenomes (see above) to identify single nucleotide variants between PWK and STF in the

binding site locus. Now, for all the target genes of a given TF, we categorized them as having

sequence variants or not in the respective binding site, and exhibiting senescence-associated

differential allele-specific expression. We eliminated from further consideration any TF with

fewer than 250 target genes in each of the four categories. For all remaining TFs, the 2 x 2

contingency table was used as input into a Fisher's exact test with Benjamini-Hochberg multiple

testing correction.

***Usf2* shRNA vector design, construction and application**

Usf2 knockdown shRNA sequences were obtained from the Broad Institute Genetic

Perturbation Portal (https://portals.broadinstitute.org/gpp/public/). Two shRNA sequences for

Usf2

(CCGGGCAAGACAGGAGCAAGTAAAGCTCGAGCTTTACTTGCTCCTGTCTTGCTTTTTTGAA

T; CCGGACAAGGAGACATAATGCATTTCTCGAG-

AAATGCATTATGTCTCCTTGTTTTTTTGAAT)*,* and, separately, a scrambled control sequence

(CCTAAGGTTAAGTCGCCCTCGCTCGAGCGAGGGCGACTTAACCTTAGG, Addgene cat.

#1864), were each cloned into pLKO.1 puro lentiviral vectors (Addgene cat. #8453). Lentiviral

particles containing each of the shRNA constructs were generated by calcium phosphate co-

transfection of HEK 293T cells with the shRNA pLKO.1 puro vectors and separate

pMDLg/pRRE packaging and pCMV-VSV-G envelope plasmids generously provided by Dr.

Marius Walter of the Verdin Lab at the Buck Institute. The number of viral particles generated

was determined using the Origene One-Wash™ Lentivirus Titer Kit, p24 ELISA (cat.

#TR30038). These particles were used to infect two biological replicates of purebred *M.*

musculus (PWK) primary tail fibroblasts at a multiplicity of infection of 5 with 4 µg/mL of

polybrene, and infected cells were selected by incubating with 2 µg/mL puromycin for 10 days,

changing media and antibiotic every other day. Knockdown of *Usf2* was determined by qPCR,

using *Usf2* qPCR primer sequences chosen through NCBI Primer Blast, filtering for those

spanning an exon-exon junction. The primer pair with the same efficiency (calculated as $10^{(-1/slope)}$

when plotting log concentration of template cDNA versus Ct) as the internal control *Actb*

qPCR primers was chosen: *Usf2* forward 5' TTCGGCGACCACAATATCCAG 3', *Usf2* reverse 5'

TTCGGCGACCACAATATCCAG 3', *Actb* forward 5' CAACCGTGAAAAGATGACCC 3', *Actb*

reverse 5' GTAGATGGGCACAGTGTGGG 3'. *Usf2* expression was calculated using the Delta-

Delta Ct method (103).

Cell proliferation and DNA damage assays

For each of two biological replicates of purebred *M. musculus* (PWK) cells infected with

lentivirus harboring the scrambled control and two of each *Usf2* knockdown, either before

irradiation or 6 hours after irradiation (see above), we measured cell proliferation and DNA

damage response as follows.

For a given replicate, DNA synthesis was measured via 5-ethynyl-2′-deoxyuridine (EdU)

incorporation assays using the Invitrogen™ Click-iT™ Edu Alexa Fluor™ 488 Flow Cytometry

Assay Kit (cat. #C10420). Cells were treated with 5 µM EdU in complete medium and left in the

incubator overnight. The following day cells were fixed, permeabilized, and treated with Alexa

Fluor™ 488 azide before running through a BD LSRFortessa™ Cell Analyzer to identify the

percentage of EdU positive cells.

For a given replicate, we carried out a comet assay to measure levels of DNA double stranded

breaks for a given replicate culture as described (53). Briefly, slides scored with a diamond

tipped scribe were dipped in 1% low melting point agarose and left at 4°C overnight. Cells of all

genotypes tested were mixed into 1% low melting point agarose at 2×10^4 cells/ml, spread over

the prepared slides and left to solidify for 1 hour. Slides were then incubated at 37°C in 2%

sarkosyl, 0.5 M NA$_2$EDTA and 0.5 mg/mL proteinase K pH 8 lysis buffer overnight. The

following day the slides were washed three times in 90 mM Tris, 90 mM boric acid, and 2 mM

NA$_2$EDTA pH 8.5 wash buffer, then subject to electrophoresis at 0.6 V/cm for 25 minutes in

wash buffer. The slides were then washed three times in distilled water and placed in a staining

solution containing 5 µg/mL of propidium iodide for 20 minutes. The slides were washed again

once with distilled water and multiple representative images were taken of each sample using a

Zeiss AxioObserver epi-fluorescent microscope. Images were processed via ZEN Digital

Imaging for Light Microscopy (RRID:SCR_013672) and comet tail moments were analyzed via

OpenComet (104) in ImageJ (105).

For H2AX assays, for a given replicate, cells were cultured in 8-chamber Nunc™ Lab-Tek™ II

Chamber Slides™ (Thermo Fisher cat. # 154453). The day before staining, all cells were

seeded at 60-70% confluence in each chamber. Cells were fixed by incubating with 4%

paraformaldehyde for 5 minutes at 4°C and permeabilized using 0.1% Triton X-100 for 15

minutes at room temperature. The cells were then blocked using 3% bovine serum albumin

(BSA) in PBS for 45 minutes at room temperature, then incubated with 1 µg/mL of primary

antibodies specific to phosphorylated (Ser 139) H2AX (cat. # sc-517348, Santa Cruz

Biotechnology) in 3% BSA overnight at 4°C. The following day the cells were washed in PBS

three times before incubating with 2 µg/mL of Alexa 488 secondary antibodies purchased from

Invitrogen (cat. # A11001) for two hours at room temperature. Cells were washed three times

with PBS then incubated with 0.5 µg/mL DAPI for 5 minutes at room temperature. The cells

were washed once more with PBS before mounting for imaging. Multiple representative

confocal images of each sample were taken using a Zeiss LSM 710 AxioObserver. Images

were processed using ImageJ, taking the average background fluorescence across several

nuclei in each field of vision and only counting foci with fluorescence intensities above the

background threshold.

In addition to the above, we also carried out EdU incorporation assays on two biological

replicates of wild-type primary fibroblasts of the *M. musculus* (PWK) background at five days

after irradiation, and observed 0.5 and 1% EdU staining, respectively.

Multivariate ANOVA of irradiation and senescence timecourse

To identify genes whose expression changed in wild-type cells through irradiation and

senescence, we used RNA-seq profiling data from purebred *M. musculus* (PWK) cells harboring

a scrambled shRNA before and 6 hours, 10 days, and 20 days after irradiation (see

Supplemental Table S6) as input into a multivariate ANOVA test.

MERLIN regulatory network reconstruction

To reconstruct a regulatory network we used RNA-seq profiling data for purebred *M. musculus*

(PWK) cells harboring a scrambled shRNA or a *Usf2*-targeting RNA, before and 6 hours, 10

days and 20 days after irradiation, as input into MERLIN (106) with default settings. Analysis

used a catalog of murine transcription factors from the Gene Transcription Regulation Database

(34).

Data access

All raw and processed sequencing data generated in this study have been submitted to the

NCBI Gene Expression Omnibus (GEO; https://www.ncbi.nlm.nih.gov/geo/) under accession

number GSE201217. Raw data and complete MS data sets have been uploaded to the Mass

Spectrometry Interactive Virtual Environment (MassIVE) repository, developed by the Center for

Computational Mass Spectrometry at the University of California San Diego, and can be

downloaded using the following link:

http://massive.ucsd.edu/ProteoSAFe/status.jsp?task=5e7aa6a2b31f4dcfafa679e9d3f3d9c3

(MassIVE ID number: MSV000089246; ProteomeXchange ID: PXD033182). *Enter the*

username and password in the upper right corner of the page: Username:

MSV000089246_reviewer, Password: winter.

Conflict of interest statement

The authors report no conflicts of interest.

Acknowledgements

The authors thank Mary West for support of cell culture resources, Melissa Chao for help with

data collection, and Herbert Kasler, Marius Walter, and Eric Verdin for helpful discussions and

materials. This work was supported by National Institutes of Health R01 NS116992 and R01

GM120430 to R.B.B and R01 HD094787 to J.M.G. E.E.K.K. was supported by the National

Science Foundation Graduate Research Fellowship Program (DGE-1313190). Any opinions,

findings, and conclusions or recommendations expressed in this material are those of the

author(s) and do not necessarily reflect the views of the National Science Foundation or the

National Institutes of Health.

Author contributions

RBB and TK conceived of the idea of the study; TK carried out fibroblast harvest and culture,

RNA-seq, knockdown, and cell-biological assays and all data analysis. BS and CDK carried out

proteomics profiling. EEKK, ECM, and JMG carried out all mouse mating and husbandry. TK,

JC, and RBB wrote the manuscript with input from all authors.

Figure captions

Figure 1: Senescent *M. spretus* cells exhibit lower β-galactosidase activity. (A)

Representative images of senescent primary fibroblasts from *M. musculus* (above) and *M.*

spretus (below) stained with the β-galactosidase indicator X-Gal seven days after irradiation. (B)

Each trace reports results from a timecourse of X-Gal staining assays of primary senescent

cells of the indicated species as in (A). The *y*-axis reports the proportion of cells stained

positive for senescence-associated β-galactosidase (SABG) activity. In a given column, small

points report biological and technical replicates and large points report their average (*M.*

musculus n = 9, *M. spretus n* = 5). ***, $p < 0.001$, ****, $p < 0.0001$, one-tailed Wilcoxon test

comparing species.

Figure 2: Senescent *M. spretus* cells express fewer secreted factors. (A) Each trace

reports a cumulative distribution of the change, in senescent primary fibroblasts of the indicated

species, in mRNA levels of genes of the senescence associated secretory phenotype (4) with

senescence. The *y*-axis reports the proportion of genes with the expression change on the *x*-

axis, with the latter taken as an average across replicates. (B) Each row shows results from a

test of the genes of the indicated Gene Ontology term for enrichment of expression change

between the species during senescence, with *p*-values from a resampling-based test, corrected

for multiple testing. (C) Annotations are as in (A) except that measurements were of secreted

peptides, and the complete set of peptides with significant differential abundance between

species is shown.

Figure 3: A screen correlating divergence in transcription factor binding sites and *cis-*

regulatory expression during senescence reveals USF2 as a top hit. (A) *M. musculus*

(blue) and *M. spretus* (red) alleles of a gene are expressed differently in interspecific F1 hybrid

cells in a senescence-dependent manner, as a product of a sequence variant (green striped) in

the binding site for a transcription factor (yellow). (B) Each row reports the multiple testing-

corrected *p*-value from a Fisher's Exact Test of target genes of the indicated transcription factor,

quantifying association between species differences in experimentally determined binding sites

(34) and allele-specific expression in primary cells of the *M. musculus* x *M. spretus* F1 hybrid

background before and after senescence induction. Results for all tested factors are listed in

Supplemental Table S5. (C) Shown are the input data for the Fisher's Exact Test in (B) for

USF2. Each bar reports analysis of genes with USF2 binding sites in their upstream regions

that do (right) or do not (left) harbor sequence variants between *M. musculus* and *M. spretus*.

For a given bar, genes are categorized by the presence or absence of senescence-dependent

differential allele-specific expression (Δ ASE). (D) For each trace, the *x*-axis reports the number

of sequence variants between *M. musculus* and *M. spretus* in a given USF2 binding site, and

the *y*-axis reports the proportion of all USF2 target genes bearing the number of variants on the

x, as a kernel density estimate. Colors denote the presence or absence of senescence-

dependent differential allele-specific expression at USF2 target genes as in (C). (E) Data are as

in (D) except that the *x*-axis reports the distance of the variant from the center of the USF2

binding site.

Figure 4: *Usf2* depletion results in more DNA damage but a muted DNA damage response

following irradiation. (A) *M. musculus* primary fibroblasts were infected with a lentivirus

encoding a short hairpin RNA (shRNA, SH) targeting *Usf2* or a scrambled control, and analyzed

before (SH unirradiated) or six hours after (SH→IR) treatment with ionizing radiation. (B) In a

given row, the second column reports the average, across genes of the indicated Gene

Ontology term, of the $\log_2$ of the ratio of expression between cells bearing shRNAs targeting

Usf2 and those with a scrambled control (SCR), six hours after irradiation. The third column

reports significance in a resampling-based test for enrichment of directional differential

expression between *Usf2* knockdown cells and the scrambled control in the respective term,

corrected for multiple testing. The fourth column reports the direction of the change in

expression before and six hours after irradiation in cells of the scrambled control genotype. (C)

Each trace reports a cumulative distribution of the $\log_2$ of the ratio of expression between cells

harboring bearing shRNAs targeting *Usf2* or the scrambled control in genes annotated in

apoptosis, before or six hours after irradiation treatment as indicated. The *y*-axis reports the

proportion of genes with the expression change on the *x*-axis. (D) Data are as in (C), except

that EGFR signaling pathway genes were analyzed. (E) Each column reports tail moments

detected in a comet assay on primary fibroblasts harboring the indicated shRNAs, before or six

hours after irradiation as indicated. In a given column, points report biological and technical

replicates and the bar height reports their average (SCR $n = 5$, *Usf2* KD $n = 20$). ****, $p <$

0.0001, one-tailed Wilcoxon test. (F) Left, each column reports number of γH2AX foci per cell

detected in primary fibroblasts harboring the indicated shRNAs six hours after irradiation. In a

given column, points report biological and technical replicates and the bar height reports their

average (SCR $n = 12$, *Usf2* KD $n = 28$). ***, $p < 0.001$, one-tailed Wilcoxon test. Right,

representative images of the indicated cultures.

Figure 5: *Usf2* knockdown results in an enhanced senescence profile. (A) *M. musculus*

primary fibroblasts were infected with a lentivirus encoding an shRNA (SH) targeting *Usf2* or a

scrambled control, and analyzed before (SH unirradiated) or after (SH→SEN) treatment with

ionizing radiation (IR) to induce senescence (SEN). (B) Data are as in Figure 4B except that

cells were analyzed 10 days after irradiation. (C) *M. musculus* primary fibroblasts were

irradiated, incubated for 10 days to senesce, then infected with shRNAs and analyzed after 10

additional days. (D) Data are as in (B) except that cells from the scheme in (C) were analyzed.

(E) Each trace reports a cumulative distribution of the $\log_2$ of the ratio of expression in cells

bearing shRNAs targeting *Usf2* and those with a scrambled control (SCR), in genes annotated

in the inflammatory response, when shRNAs were administered to a resting culture (SH

unirradiated), to resting cells followed by irradiation (SH→SEN), or after irradiation and

senescence establishment (SEN→SH). The *y*-axis reports the proportion of genes with the

expression change on the *x*-axis, with the latter taken as an average across replicates. (F) Data

are as in (E), except that genes involved in G2/M transition of mitotic cell cycle were analyzed.

(G) Left, each column reports the proportion of senescence associated β-galactosidase (SABG)

positive cells treated with the indicated shRNAs in resting culture (unirradiated) or 7 days after

irradiation (SEN). In a given column, points report biological and technical replicates and the

bar height reports their average (unirradiated $n = 4$, SEN $n = 24$). **, $p < 0.01$, one-tailed

Wilcoxon test. Right, representative images of the indicated cultures.

Supplemental figure captions

Supplemental Figure S1: Senescent *M. spretus* primary fibroblasts display muted mRNA

induction of genes of the senescence-associated secretory phenotype. Each column

reports expression (TPM, transcripts per million) from RNA-seq profiling of primary fibroblasts

from the indicated species for the indicated genes, in control or irradiation-induced senescent

(SEN) cells. In a given column, points report biological replicates and the bar height reports

their average ($n = 3$).

Supplemental Figure S2: Senescent *M. spretus* primary fibroblasts display muted

secretion of proteins of the senescence-associated secretory phenotype. Data are as in

Supplemental Figure S1 except that protein abundance from conditioned medium is shown.

Supplemental Figure S3: Senescent *M. musculus* x *M. spretus* F1 primary fibroblasts

display intermediate activity of senescence-associated β-galactosidase (SABG). Each

column reports the proportion of SABG positive cells for the indicated genotype in control cells

or in senescent cells seven days after irradiation (SEN). In a given column, points report

biological and technical replicates and the bar height reports their average (*M. musculus* $n = 9$,

M. spretus $n = 5$, F1 hybrid $n = 2$).

Supplemental Figure S4: Variation in gene expression during senescence between *M.*

***musculus* and *M. spretus* is controlled by both *cis*- and *trans*-acting elements.** Each point

represents expression of one gene in primary fibroblasts induced to senesce. The *x*-axis

reports the ratio of expression, as an average across biological replicates, measured in

purebred *M. musculus* (Mm) and *M. spretus* (Ms) cells; the *y*-axis reports the ratio of allele-

specific expression from the *M. musculus* and *M. spretus* alleles in cells from the interspecific

F1 hybrid.

Supplemental Figure S5: Senescent *M. musculus* x *M. spretus* F1 primary cells display

intermediate induction in mRNA expression of genes of the senescence-associated

secretory phenotype. Data are as in Figure 2A of the main text except that measurements

from the interspecific F1 hybrid are shown in green.

Supplemental Figure S6: *Usf2*-targeting shRNAs knock down *Usf2* expression. Each set

of bars reports *Usf2* expression measured via qPCR in *M. musculus* primary cells, in one

biological replicate. In a given replicate, each bar reports the fold change (FC) in *Usf2*

expression between cells harboring *Usf2*-targeting shRNA (*Usf2* KD) and those with a

scrambled shRNA (SCR), normalized with respect to the value of the latter.

Supplemental Figure S7: *Usf2* depletion slightly slows growth of primary *M. musculus*

cells independent of irradiation. Each column reports the percentage of EdU incorporation in

primary fibroblasts harboring *Usf2* or scrambled shRNAs, before or six hours after irradiation as

indicated. *, $p < 0.05$, one-tailed Wilcoxon test. In a given column, points report biological and

technical replicates and the bar height reports their average (SCR $n = 11$, *Usf2* KD $n = 22$).

Supplemental Figure S8: *Usf2* knockdown results in less transcriptional repression

following acute DNA damage. In a given panel, each column reports mRNA expression of the

indicated gene in transcripts per million (TPM) in primary fibroblasts harboring *Usf2* or

scrambled shRNAs, before or six hours after irradiation as indicated. (A) Shown are a

representative subset of the genes that were repressed in cells harboring the scrambled shRNA

control following acute DNA damage. (B) Shown are genes in the Gene Ontology term

"epidermal growth factor receptor signaling pathway." Inset reports genes with average TPM

>125 in one or more samples. In a given column, points report biological replicates and the bar

height reports their average (n = 3).

Supplemental Figure S9: Expression of core DNA damage response genes are largely not

affected by *Usf2* knockdown. Data are as in Supplemental Figure S8 except that DNA

damage response genes are shown.

Supplemental Figure S10: *Usf2* knockdown results in an enhanced senescent gene

expression profile. In a given panel, each column reports mRNA expression in transcripts per

million (TPM) in primary fibroblasts harboring *Usf2* or scrambled shRNAs, when shRNAs were

administered to a resting culture (unirradiated), to resting cells followed by irradiation (→SEN),

or after irradiation and senescence establishment (→KD). (A) Genes annotated in inflammation

and immune response that were upregulated during senescence in scrambled shRNA controls.

(B) Genes annotated in cell cycle that were repressed with senescence in scrambled shRNA

controls. In a given column, points report biological replicates and the bar height reports their

average (SCR n = 2, *Usf2* KD n = 4).

Supplemental Figure S11: Expression profiling of control cells during a senescence

timecourse reveals a dynamic expression program with senescence progression. Each

trace reports a cumulative distribution of gene expression in primary *M. musculus* fibroblasts

harboring a scrambled shRNA, at the indicated timepoint after irradiation treatment relative to

unirradiated controls. Included in the distribution are all genes significant in a test for significant

expression change across the timepoints by multivariate ANOVA (Supplemental Table S10).

The *y*-axis reports the proportion of genes with the expression change on the *x*-axis.

Supplemental table captions

Supplemental Table S1: RNA-seq gene expression profiling of resting and senescent

purebred *M. musculus* and *M. spretus* primary cells. Each row reports expression of the

indicated gene in transcripts per million, as an average across biological replicates of primary

fibroblasts from the indicated background cultured in the indicated condition. Columns report

expression in purebred *M. musculus* (Mm) or *M. spretus* (Ms) primary fibroblasts, or expression

from each allele (*M. musculus*, Mm; *M. spretus*, Ms) in primary fibroblasts from the interspecific

F1 hybrid, in unirradiated control (CONT) or 10 days after irradiation (senescent, SEN).

Supplemental Table S2: Functional-genomic enrichment analysis of expression profiles

of unirradiated purebred *M. musculus* and *M. spretus* primary cells. Each row reports the

results of a resampling test for enrichment of expression change between unirradiated primary

fibroblasts from purebred *M. musculus* and *M. spretus* (see Supplemental Table S1) for the

indicated Gene Ontology biological process term. The second column represents the absolute

value of the sum of the ratios of expression in *M. musculus* versus *M. spretus* for all the genes

in the indicated term. The third column reports the number of genes with expression data in the

term. The fourth column reports the raw p-value from resampling analysis, and the final column

reports Benjamini-Hochberg corrected p-values. Only terms with significant enrichment

(corrected $p < 0.05$) are shown.

Supplemental Table S3: Functional-genomic enrichment analysis of expression profiles

of senescent purebred *M. musculus* and *M. spretus* primary cells. Data are as in

Supplemental Table S2 except that *M. musculus* and *M. spretus* cells were analyzed 10 days

after irradiation (see Supplemental Table S1).

Supplemental Table S4: Proteomic profiling of conditioned medium from purebred *M. musculus* and *M. spretus* primary cells via mass spectrometry. Columns report the abundance of the indicated protein via mass spectrometry, as an average over replicates and individual detected peptides, in conditioned medium from *M. musculus* (Mm) or *M. spretus* (Ms) cells, each as an unirradiated control (CONT) or 10 days after irradiation (senescent, SEN).

Supplemental Table S5: A screen correlating divergence in transcription factor binding sites and *cis*-regulatory expression during senescence. Each row reports the results of a Fisher's exact test relating, for the indicated transcription factor, two measures of variation between *M. musculus* and *M. spretus*: sequence variants at the factor's experimentally determined binding sites (34) upstream of genes, and, at these genes, senescence-specific expression differences between the two species' alleles in fibroblasts of the F1 hybrid background. The second and third columns report nominal and Benjamini-Hochberg corrected *p*-values respectively.

Supplemental Table S6: Expression profiling of shRNA-expressing purebred *M. musculus* primary cells across a timecourse of irradiation and senescence. Columns report expression of the indicated gene in transcripts per million, as an average across biological replicates, in purebred primary fibroblasts from *M. musculus* (PWK) subjected to either of two regimes. In one, cells were first infected with lentivirus expressing the indicated short hairpin RNA (shRNA) and then profiled before (CONT) or 6 hours (6HR) or 10 days (SEN.10d) after irradiation. In the other regime, cells were irradiated, then infected with lentivirus expressing the indicated shRNA, and then profiled 20 days later (SEN.20d). Usf2, *Usf2* targeting shRNA; SCR, scrambled control shRNA.

Supplemental Table S7: Functional-genomic enrichment analysis of shRNA-expressing purebred *M. musculus* primary cells in untreated conditions. Data are as in Supplemental Table S2 except that *M. musculus* (PWK) primary cells harboring short hairpin RNAs targeting *Usf2* or a scrambled control were cultured in unirradiated conditions (see Supplemental Table S6); for each gene we formulated the average expression measurement across replicates from each shRNA category in turn, and we then took the ratio between these averages for input into Gene Ontology enrichment tests.

Supplemental Table S8: Functional-genomic enrichment analysis of purebred *M. musculus* primary cell transcriptomes upon *Usf2* knockdown and irradiation. Data are as in Supplemental Table S7 except that cells were infected with lentivirus expressing the indicated short hairpin RNA (shRNA), treated with ionizing radiation, and profiled 6 hours afterward (see Supplemental Table S6).

Supplemental Table S9: Functional-genomic enrichment analysis of purebred *M. musculus* primary cell transcriptomes upon *Usf2* knockdown, irradiation, and senescence establishment. Data are as in Supplemental Table S7 except that cells were infected with lentivirus expressing the indicated short hairpin RNA (shRNA), treated with ionizing radiation, and profiled 10 days afterward (see Supplemental Table S6).

Supplemental Table S10: Expression analysis of purebred *M. musculus* primary cells across a timecourse of irradiation and senescence. Each row reports results from analysis of expression of one gene in *M. musculus* (PWK) primary cells harboring a scrambled short hairpin RNA, before and 6 hours, 10 days, and 20 days after irradiation (see Supplemental Table S6). Second and third columns report the raw and Benjamini-Hochberg corrected *p*-values respectively from a multivariate ANOVA test for variation in expression across the

timecourse. Only genes with significant differential expression across the timecourse (corrected

$p < 0.05$) are shown.

Supplemental Table S11: Functional-genomic enrichment analysis of purebred *M.

musculus **primary cell transcriptomes upon irradiation, senescence establishment, *Usf2***

knockdown, and long-term senescence. Data are as in Supplemental Table S2 except that

M. musculus (PWK) primary cells were irradiated, incubated for 10 days, infected with lentivirus

harboring short hairpin RNAs targeting *Usf2* or a scrambled control, and allowed to maintain

senescence for another 10 days (see Supplemental Table S6); for each gene we formulated the

average expression measurement across replicates from each shRNA category in turn, and we

then took the ratio between these averages for input into Gene Ontology enrichment tests.

Supplemental Table S12: *In silico* regulatory network reconstruction from irradiation and

senescence transcriptomes. Shown are results from regulatory network inference using all

shRNA treated transcriptomes in this study (see Supplemental Table S6) as input into the

MERLIN package (106). Each row reports results from a transcription factor inferred to be a

downstream target of USF2. The second column reports the strength of the inferred regulation

by USF2; negative values indicate repression. In the remaining columns, "Yes" indicates that

one or more genes of the indicated GO term were inferred to be targets of the indicated factor.

1004

1005   **Supplemental Table S13: Mouse metadata.** Shown are characteristics of mice used for tail

1006   cuttings and fibroblast isolation in this study.

1007

1008   Figure 1

1009

1010   Figure 2

GO term	M. mus / M. spr effect size	p-value	SEN direction in M. mus | M. spr
antimicrobial humoral immune response mediated by antimicrobial peptide	0.7051633	0.000000	+ | +
regulation of NIK/NF-kappaB signaling	0.3395436	0.000000	+ | -
immune response	0.7865077	0.000000	+ | +
neutrophil chemotaxis	0.4491496	0.018600	+ | +

1011

1012   Figure 3

1013

1014

Figure 4

GO term (SH→IR)	Usf2 KD effect size	Usf2 KD p-value	IR direction in SCR
positive regulation of transcription by RNA polymerase II	0.1481405	0.0000000	-
negative regulation of cell population proliferation	0.1087087	0.0000000	-
negative regulation of apoptotic process	0.1351093	0.0000000	-
epidermal growth factor receptor signaling pathway	0.3011997	0.0000000	+
positive regulation of ERK1 and ERK2 cascade	0.1060674	0.0087796	-

1016

1017

1018   Figure 5

1019

1020

1021   Supplemental Figure S1

1022

1023 Supplemental Figure S2

1024

1025 Supplemental Figure S3

1026

1027 Supplemental Figure S4

1028

1029

1030 Supplemental Figure S5

1031

1032

1033   Supplemental Figure S6

1034

1035

1036   Supplemental Figure S7

1037

1038

1039 Supplemental Figure S8

1040

1041

1042   Supplemental Figure S9

1043

1044

Supplemental Figure S10

1048   Supplemental Figure S11

1049

References

1. Hayflick L. The limited in vitro lifetime of human diploid cell strains. Exp Cell Res. 1965 Mar 1;37(3):614–36.

2. Campisi J. Senescent cells, tumor suppression, and organismal aging: good citizens, bad neighbors. Cell. 2005 Feb 25;120(4):513–22.

3. Campisi J, Fagagna FDD. Cellular senescence: when bad things happen to good cells. Nat Rev Mol Cell Biol. 2007;

4. Coppé JP, Patil CK, Rodier F, Sun Y, Muñoz DP, Goldstein J, et al. Senescence-Associated Secretory Phenotypes Reveal Cell-Nonautonomous Functions of Oncogenic RAS and the p53 Tumor Suppressor. PLOS Biol. 2008 Dec 2;6(12):e301.

5. Wan M, Gray-Gaillard EF, Elisseeff JH. Cellular senescence in musculoskeletal homeostasis, diseases, and regeneration. Bone Res. 2021 Sep 10;9(1):1–12.

6. Paramos-de-Carvalho D, Jacinto A, Saúde L. The right time for senescence. Bronner ME, editor. eLife. 2021 Nov 10;10:e72449.

7. Baker DJ, Wijshake T, Tchkonia T, LeBrasseur NK, Childs BG, van de Sluis B, et al. Clearance of p16Ink4a-positive senescent cells delays ageing-associated disorders. Nature. 2011 Nov 2;479(7372):232–6.

8. Demaria M, Ohtani N, Youssef SA, Rodier F, Toussaint W, Mitchell JR, et al. An essential role for senescent cells in optimal wound healing through secretion of PDGF-AA. Dev Cell. 2014 Dec 22;31(6):722–33.

9.   Olivieri F, Prattichizzo F, Grillari J, Balistreri CR. Cellular Senescence and Inflammaging

in Age-Related Diseases. Mediators Inflamm. 2018 Apr 17;2018:e9076485.

10.   Davalos AR, Coppe JP, Campisi J, Desprez PY. Senescent cells as a source of

inflammatory factors for tumor progression. Cancer Metastasis Rev. 2010

Jun;29(2):273–83.

11.   Krtolica A, Parrinello S, Lockett S, Desprez PY, Campisi J. Senescent fibroblasts

promote epithelial cell growth and tumorigenesis: a link between cancer and aging. Proc

Natl Acad Sci U S A. 2001 Oct 9;98(21):12072–7.

12.   Parrinello S, Coppe JP, Krtolica A, Campisi J. Stromal-epithelial interactions in aging

and cancer: senescent fibroblasts alter epithelial cell differentiation. J Cell Sci. 2005 Feb

1;118(Pt 3):485–96.

13.   Kang C. Senolytics and Senostatics: A Two-Pronged Approach to Target Cellular

Senescence for Delaying Aging and Age-Related Diseases. Mol Cells. 2019

Dec;42(12):821–7.

14.   Kim EC, Kim JR. Senotherapeutics: emerging strategy for healthy aging and age-related

disease. BMB Rep. 2019 Jan;52(1):47–55.

15.   Campisi J. Aging, cellular senescence, and cancer. Annu Rev Physiol. 2013;75:685–

705.

16.   Salotti J, Johnson PF. Regulation of senescence and the SASP by the transcription

factor C/EBPβ. Exp Gerontol. 2019 Dec;128:110752.

17.   Chan M, Yuan H, Soifer I, Maile TM, Wang RY, Ireland A, et al. Novel insights from a

1092   multiomics dissection of the Hayflick limit. eLife [Internet]. 2022 Feb 1 [cited 2022 Mar

1093   10];11. Available from: https://doi.org/10.7554/eLife.70283

1094   18.   Wang Y, Xu Q, Sack L, Kang C, Elledge SJ. A gain-of-function senescence bypass

1095   screen identifies the homeobox transcription factor DLX2 as a regulator of ATM–p53

1096   signaling. Genes Dev. 2016 Feb 1;30(3):293–306.

1097   19.   Xie Q, Peng S, Tao L, Ruan H, Yang Y, Li TM, et al. E2F Transcription Factor 1

1098   Regulates Cellular and Organismal Senescence by Inhibiting Forkhead Box O

1099   Transcription Factors. J Biol Chem. 2014 Dec 5;289(49):34205–13.

1100   20.   Han R, Li L, Ugalde AP, Tal A, Manber Z, Barbera EP, et al. Functional CRISPR screen

1101   identifies AP1-associated enhancer regulating FOXF1 to modulate oncogene-induced

1102   senescence. Genome Biol. 2018 Aug 17;19(1):118.

1103   21.   Martínez-Zamudio RI, Roux PF, de Freitas JANLF, Robinson L, Doré G, Sun B, et al.

1104   AP-1 imprints a reversible transcriptional programme of senescent cells. Nat Cell Biol.

1105   2020 Jul;22(7):842–55.

1106   22.   Brückmann NH, Bennedsen SN, Duijf PHG, Terp MG, Thomassen M, Larsen M, et al. A

1107   functional genetic screen identifies the Mediator complex as essential for SSX2-induced

1108   senescence. Cell Death Dis. 2019 Nov 6;10(11):1–12.

1109   23.   Tyler EJ, Gutierrez Del Arroyo A, Hughes BK, Wallis R, Garbe JC, Stampfer MR, et al.

1110   Early growth response 2 (EGR2) is a novel regulator of the senescence programme.

1111   Aging Cell. 2021 Mar;20(3):e13318.

24.   Zhang C, Zhang X, Huang L, Guan Y, Huang X, Tian XL, et al. ATF3 drives senescence

by reconstructing accessible chromatin profiles. Aging Cell. 2021 Mar;20(3):e13315.

25.   Casella G, Munk R, Kim KM, Piao Y, De S, Abdelmohsen K, et al. Transcriptome

signature of cellular senescence. Nucleic Acids Res. 2019 Aug 22;47(14):7294–305.

26.   Zhao H, Darzynkiewicz Z. Biomarkers of Cell Senescence Assessed by Imaging

Cytometry. Methods Mol Biol Clifton NJ. 2013;965:83–92.

27.   Dimri GP, Lee X, Basile G, Acosta M, Scott G, Roskelley C, et al. A biomarker that

identifies senescent human cells in culture and in aging skin in vivo. Proc Natl Acad Sci

U S A. 1995 Sep 26;92(20):9363–7.

28.   Kim EK, Moon S, Kim DK, Zhang X, Kim J. CXCL1 induces senescence of cancer-

associated fibroblasts via autocrine loops in oral squamous cell carcinoma. PLOS ONE.

2018 Jan 23;13(1):e0188847.

29.   Coppé JP, Patil CK, Rodier F, Krtolica A, Beauséjour CM, Parrinello S, et al. A Human-

Like Senescence-Associated Secretory Phenotype Is Conserved in Mouse Cells

Dependent on Physiological Oxygen. PLOS ONE. 2010 Feb 12;5(2):e9188.

30.   Coppé JP, Desprez PY, Krtolica A, Campisi J. The senescence-associated secretory

phenotype: the dark side of tumor suppression. Annu Rev Pathol. 2010;5:99–118.

31.   Levi N, Papismadov N, Solomonov I, Sagi I, Krizhanovsky V. The ECM path of

senescence in aging: components and modifiers. FEBS J. 2020 Jul;287(13):2636–46.

32.   Sun W, Hu Y. eQTL Mapping Using RNA-seq Data. Stat Biosci. 2013 May 1;5(1):198–

219.

33.   Wittkopp PJ, Haerum BK, Clark AG. Evolutionary changes in cis and trans gene

1134   regulation. Nature. 2004 Jul 1;430(6995):85–8.

1135   34.   Yevshin I, Sharipov R, Kolmykov S, Kondrakhin Y, Kolpakov F. GTRD: a database on

1136   gene transcription regulation-2019 update. Nucleic Acids Res. 2019 Jan 8;47(D1):D100–

1137   5.

1138   35.   Wang Y, Sui Y, Lian A, Han X, Liu F, Zuo K, et al. PBX1 Attenuates Hair Follicle-Derived

1139   Mesenchymal Stem Cell Senescence and Apoptosis by Alleviating Reactive Oxygen

1140   Species-Mediated DNA Damage Instead of Enhancing DNA Damage Repair. Front Cell

1141   Dev Biol. 2021;9:739868.

1142   36.   Bandyopadhyay D, Okan NA, Bales E, Nascimento L, Cole PA, Medrano EE. Down-

1143   regulation of p300/CBP histone acetyltransferase activates a senescence checkpoint in

1144   human melanocytes. Cancer Res. 2002 Nov 1;62(21):6231–9.

1145   37.   Yang K, Wang F, Zhang H, Wang X, Chen L, Su X, et al. Target Inhibition of CBP

1146   Induced Cell Senescence in BCR-ABL- T315I Mutant Chronic Myeloid Leukemia. Front

1147   Oncol [Internet]. 2021 [cited 2022 Mar 10];10. Available from:

1148   https://www.frontiersin.org/article/10.3389/fonc.2020.588641

1149   38.   Qyang Y, Luo X, Lu T, Ismail PM, Krylov D, Vinson C, et al. Cell-type-dependent activity

1150   of the ubiquitous transcription factor USF in cellular proliferation and transcriptional

1151   activation. Mol Cell Biol. 1999 Feb;19(2):1508–17.

1152   39.   Pawar SA, Szentirmay MN, Hermeking H, Sawadogo M. Evidence for a cancer-specific

1153   switch at the CDK4 promoter with loss of control by both USF and c-Myc. Oncogene.

1154   2004 Aug 12;23(36):6125–35.

40.   Chen N, Szentirmay MN, Pawar SA, Sirito M, Wang J, Wang Z, et al. Tumor-

suppression function of transcription factor USF2 in prostate carcinogenesis. Oncogene.

2006 Jan;25(4):579–87.

41.   Aperlo C, Boulukos KE, Pognonec P. The basic region/helix-loop-helix/leucine repeat

transcription factor USF interferes with Ras transformation. Eur J Biochem. 1996 Oct

1;241(1):249–53.

42.   Hu D, Tjon EC, Andersson KM, Molica GM, Pham MC, Healy B, et al. Aberrant

expression of USF2 in refractory rheumatoid arthritis and its regulation of

proinflammatory cytokines in Th17 cells. Proc Natl Acad Sci. 2020 Dec;117(48):30639–

48.

43.   Redon CE, Dickey JS, Bonner WM, Sedelnikova OA. γ-H2AX as a biomarker of DNA

damage induced by ionizing radiation in human peripheral blood lymphocytes and

artificial skin. Adv Space Res Off J Comm Space Res COSPAR. 2009;43(8):1171–8.

44.   Fumagalli M, Rossiello F, Mondello C, d'Adda di Fagagna F. Stable Cellular Senescence

Is Associated with Persistent DDR Activation. PLoS ONE. 2014 Oct 23;9(10):e110969.

45.   Chen JH, Ozanne SE. Deep senescent human fibroblasts show diminished DNA

damage foci but retain checkpoint capacity to oxidative stress. FEBS Lett. 2006 Dec

11;580(28–29):6669–73.

46.   Bakkenist CJ, Drissi R, Wu J, Kastan MB, Dome JS. Disappearance of the telomere

dysfunction-induced stress response in fully senescent cells. Cancer Res. 2004 Jun

1;64(11):3748–52.

47.   Siddiqui MS, François M, Fenech MF, Leifert WR. Persistent γH2AX: A promising

1177   molecular marker of DNA damage and aging. Mutat Res Rev Mutat Res. 2015

1178   Dec;766:1–19.

1179   48.   Venkata Narayanan I, Paulsen MT, Bedi K, Berg N, Ljungman EA, Francia S, et al.

1180   Transcriptional and post-transcriptional regulation of the ionizing radiation response by

1181   ATM and p53. Sci Rep. 2017 Apr;7(1):43598.

1182   49.   Silva E, Ideker T. Transcriptional Responses to DNA Damage. DNA Repair. 2019

1183   Jul;79:40–9.

1184   50.   Shang D, Sun D, Shi C, Xu J, Shen M, Hu X, et al. Activation of epidermal growth factor

1185   receptor signaling mediates cellular senescence induced by certain pro-inflammatory

1186   cytokines. Aging Cell. 2020 May;19(5):e13145.

1187   51.   Huang RX, Zhou PK. DNA damage response signaling pathways and targets for

1188   radiotherapy sensitization in cancer. Signal Transduct Target Ther. 2020 May 1;5(1):1–

1189   27.

1190   52.   Nickoloff JA, Boss MK, Allen CP, LaRue SM. Translational research in radiation-induced

1191   DNA damage signaling and repair. Transl Cancer Res. 2017 Jul;6(Suppl 5):S875–91.

1192   53.   Olive PL, Banáth JP. The comet assay: a method to measure DNA damage in individual

1193   cells. Nat Protoc. 2006;1(1):23–9.

1194   54.   Podhorecka M, Skladanowski A, Bozko P. H2AX Phosphorylation: Its Role in DNA

1195   Damage Response and Cancer Therapy. J Nucleic Acids. 2010 Aug 3;2010:920161.

1196   55.   Kim YM, Byun HO, Jee BA, Cho H, Seo YH, Kim YS, et al. Implications of time-series

1197   gene expression profiles of replicative senescence. Aging Cell. 2013;12(4):622–34.

56.   Santoro A, Spinelli CC, Martucciello S, Nori SL, Capunzo M, Puca AA, et al. Innate

immunity and cellular senescence: The good and the bad in the developmental and

aged brain. J Leukoc Biol. 2018;103(3):509–24.

57.   Kale A, Sharma A, Stolzing A, Desprez PY, Campisi J. Role of immune cells in the

removal of deleterious senescent cells. Immun Ageing. 2020 Dec;17(1):16.

58.   Basisty N, Kale A, Jeon OH, Kuehnemann C, Payne T, Rao C, et al. A proteomic atlas of

senescence-associated secretomes for aging biomarker development. PLOS Biol. 2020

Jan 16;18(1):e3000599.

59.   Purcell M, Kruger A, Tainsky MA. Gene expression profiling of replicative and induced

senescence. Cell Cycle Georget Tex. 2014;13(24):3927–37.

60.   Villarroel CA, Bastías M, Canessa P, Cubillos FA. Uncovering Divergence in Gene

Expression Regulation in the Adaptation of Yeast to Nitrogen Scarcity. mSystems. 2021

Aug 31;6(4):e0046621.

61.   Veyrieras JB, Kudaravalli S, Kim SY, Dermitzakis ET, Gilad Y, Stephens M, et al. High-

Resolution Mapping of Expression-QTLs Yields Insight into Human Gene Regulation.

PLOS Genet. 2008 Oct 10;4(10):e1000214.

62.   Vierbuchen T, Ling E, Cowley CJ, Couch CH, Wang X, Harmin DA, et al. AP-1

Transcription Factors and the BAF Complex Mediate Signal-Dependent Enhancer

Selection. Mol Cell. 2017 Dec 21;68(6):1067-1082.e12.

63.   Heinz S, Romanoski CE, Benner C, Allison KA, Kaikkonen MU, Orozco LD, et al. Effect

of natural genetic variation on enhancer selection and function. Nature. 2013

Nov;503(7477):487–92.

64. Sato AYS, Antonioli E, Tambellini R, Campos AH. ID1 inhibits USF2 and blocks TGF-β-

induced apoptosis in mesangial cells. Am J Physiol Renal Physiol. 2011

Dec;301(6):F1260-1269.

65. Chi TF, Khoder-Agha F, Mennerich D, Kellokumpu S, Miinalainen Ii, Kietzmann T, et al.

Loss of USF2 promotes proliferation, migration and mitophagy in a redox-dependent

manner. Redox Biol. 2020 Oct 7;37:101750.

66. Vicencio JM, Galluzzi L, Tajeddine N, Ortiz C, Criollo A, Tasdemir E, et al. Senescence,

apoptosis or autophagy? When a damaged cell must decide its path--a mini-review.

Gerontology. 2008;54(2):92–9.

67. Childs BG, Baker DJ, Kirkland JL, Campisi J, van Deursen JM. Senescence and

apoptosis: dueling or complementary cell fates? EMBO Rep. 2014 Nov;15(11):1139–53.

68. Zhang X, Li J, Sejas DP, Pang Q. The ATM/p53/p21 pathway influences cell fate

decision between apoptosis and senescence in reoxygenated hematopoietic progenitor

cells. J Biol Chem. 2005 May 20;280(20):19635–40.

69. Hsu CH, Altschuler SJ, Wu LF. Patterns of Early p21 Dynamics Determine Proliferation-

Senescence Cell Fate after Chemotherapy. Cell. 2019 Jul 11;178(2):361-373.e12.

70. Panneer Selvam S, Roth BM, Nganga R, Kim J, Cooley MA, Helke K, et al. Balance

between senescence and apoptosis is regulated by telomere damage-induced

association between p16 and caspase-3. J Biol Chem. 2018 Jun 22;293(25):9784–800.

71. Lee S, Schmitt CA. The dynamic nature of senescence in cancer. Nat Cell Biol. 2019

Jan;21(1):94–101.

72.   Beauséjour CM, Krtolica A, Galimi F, Narita M, Lowe SW, Yaswen P, et al. Reversal of
human cellular senescence: roles of the p53 and p16 pathways. EMBO J. 2003 Aug
15;22(16):4212–22.

73.   Baron Y, Corre S, Mouchet N, Vaulont S, Prince S, Galibert MD. USF-1 Is Critical for
Maintaining Genome Integrity in Response to UV-Induced DNA Photolesions. PLoS
Genet. 2012 Jan 26;8(1):e1002470.

74.   Song X, Zhu M, Li H, Liu B, Yan Z, Wang W, et al. USF1 promotes the development of
knee osteoarthritis by activating the NF-κB signaling pathway. Exp Ther Med. 2018 Oct
1;16(4):3518–24.

75.   Ruuth M, Soronen J, Kaiharju E, Merikanto K, Perttilä J, Metso J, et al. USF1 deficiency
alleviates inflammation, enhances cholesterol efflux and prevents cholesterol
accumulation in macrophages. Lipids Health Dis. 2018 Dec 13;17:285.

76.   Corre S, Galibert MD. Upstream stimulating factors: highly versatile stress-responsive
transcription factors. Pigment Cell Res. 2005;18(5):337–48.

77.   Bouafia A, Corre S, Gilot D, Mouchet N, Prince S, Galibert MD. p53 Requires the Stress
Sensor USF1 to Direct Appropriate Cell Fate Decision. PLoS Genet. 2014 May
15;10(5):e1004309.

78.   Samarakoon R, Overstreet JM, Higgins SP, Higgins PJ. TGF-β1 → SMAD/p53/USF2 →
PAI-1 transcriptional axis in ureteral obstruction-induced renal fibrosis. Cell Tissue Res.
2012 Jan;347(1):117–28.

79.   Dörr JR, Yu Y, Milanovic M, Beuster G, Zasada C, Däbritz JHM, et al. Synthetic lethal

1262   metabolic targeting of cellular senescence in cancer therapy. Nature. 2013

1263   Sep;501(7467):421–5.

1264   80.   Park JT, Lee YS, Cho KA, Park SC. Adjustment of the lysosomal-mitochondrial axis for

1265   control of cellular senescence. Ageing Res Rev. 2018 Nov;47:176–82.

1266   81.   Brunk UT, Terman A. Lipofuscin: mechanisms of age-related accumulation and influence

1267   on cell function. Free Radic Biol Med. 2002 Sep 1;33(5):611–9.

1268   82.   Kurz DJ, Decary S, Hong Y, Erusalimsky JD. Senescence-associated (beta)-

1269   galactosidase reflects an increase in lysosomal mass during replicative ageing of human

1270   endothelial cells. J Cell Sci. 2000 Oct;113 (Pt 20):3613–22.

1271   83.   Lee BY, Han JA, Im JS, Morrone A, Johung K, Goodwin EC, et al. Senescence-

1272   associated beta-galactosidase is lysosomal beta-galactosidase. Aging Cell. 2006

1273   Apr;5(2):187–95.

1274   84.   Pinheiro I, Dejager L, Petta I, Vandevyver S, Puimège L, Mahieu T, et al. LPS resistance

1275   of SPRET/Ei mice is mediated by Gilz, encoded by the Tsc22d3 gene on the X

1276   chromosome. EMBO Mol Med. 2013 Mar;5(3):456–70.

1277   85.   Vanlaere I, Vanderrijst A, Guénet JL, De Filette M, Libert C. Mx1 causes resistance

1278   against influenza A viruses in the Mus spretus-derived inbred mouse strain SPRET/Ei.

1279   Cytokine. 2008 Apr;42(1):62–70.

1280   86.   Blanchet C, Jaubert J, Carniel E, Fayolle C, Milon G, Szatanik M, et al. Mus spretus

1281   SEG/Pas mice resist virulent Yersinia pestis, under multigenic control. Genes Immun.

1282   2011 Jan;12(1):23–30.

87. Pérez del Villar L, Vicente B, Galindo-Villardón P, Castellanos A, Pérez-Losada J, Muro

A. Schistosoma mansoni experimental infection in Mus spretus (SPRET/EiJ strain) mice.

Parasite. 2013;20:27.

88. Dejager L, Libert C, Montagutelli X. Thirty years of Mus spretus: a promising future.

Trends Genet. 2009 May 1;25(5):234–41.

89. Khan M, Gasser S. Generating Primary Fibroblast Cultures from Mouse Ear and Tail

Tissues. J Vis Exp JoVE [Internet]. 2016 [cited 2022 Mar 23];(107). Available from:

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4781275/

90. Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, et al. Twelve years of

SAMtools and BCFtools. GigaScience. 2021 Feb 16;10(2):giab008.

91. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nat Methods.

2012 Apr;9(4):357–9.

92. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence

Alignment/Map format and SAMtools. Bioinforma Oxf Engl. 2009 Aug 15;25(16):2078–9.

93. Li H. A statistical framework for SNP calling, mutation discovery, association mapping

and population genetical parameter estimation from sequencing data. Bioinforma Oxf

Engl. 2011 Nov 1;27(21):2987–93.

94. Kim D, Pertea G, Trapnell C, Pimentel H, Kelley R, Salzberg SL. TopHat2: accurate

alignment of transcriptomes in the presence of insertions, deletions and gene fusions.

Genome Biol. 2013 Apr 25;14(4):R36.

95. Putri GH, Anders S, Pyl PT, Pimanda JE, Zanini F. Analysing high-throughput

sequencing data in Python with HTSeq 2.0. Bioinformatics. 2022 Mar 21;btac166.

96.   Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, et al. Gene Ontology:

tool for the unification of biology. Nat Genet. 2000 May;25(1):25–9.

97.   Gene Ontology Consortium. The Gene Ontology resource: enriching a GOld mine.

Nucleic Acids Res. 2021 Jan 8;49(D1):D325–34.

98.   Escher C, Reiter L, MacLean B, Ossola R, Herzog F, Chilton J, et al. Using iRT, a

normalized retention time for more targeted measurement of peptides. Proteomics. 2012

Apr;12(8):1111–21.

99.   Gillet LC, Navarro P, Tate S, Röst H, Selevsek N, Reiter L, et al. Targeted Data

Extraction of the MS/MS Spectra Generated by Data-independent Acquisition: A New

Concept for Consistent and Accurate Proteome Analysis*. Mol Cell Proteomics. 2012

Jun 1;11(6):O111.016717.

100.   Bruderer R, Bernhardt OM, Gandhi T, Xuan Y, Sondermann J, Schmidt M, et al.

Optimization of Experimental Parameters in Data-Independent Mass Spectrometry

Significantly Increases Depth and Reproducibility of Results. Mol Cell Proteomics MCP.

2017 Dec;16(12):2296–309.

101.   Collins BC, Hunter CL, Liu Y, Schilling B, Rosenberger G, Bader SL, et al. Multi-

laboratory assessment of reproducibility, qualitative and quantitative performance of

SWATH-mass spectrometry. Nat Commun. 2017 Aug 21;8(1):291.

102.   Benjamini Y, Hochberg Y. Controlling the False Discovery Rate: A Practical and

Powerful Approach to Multiple Testing. J R Stat Soc Ser B Methodol. 1995;57(1):289–

300.

103.   Livak KJ, Schmittgen TD. Analysis of Relative Gene Expression Data Using Real-Time

Quantitative PCR and the 2−ΔΔCT Method. Methods. 2001 Dec 1;25(4):402–8.

104.   Gyori BM, Venkatachalam G, Thiagarajan PS, Hsu D, Clement MV. OpenComet: An

automated tool for comet assay image analysis. Redox Biol. 2014 Jan 1;2:457–65.

105.   Schneider CA, Rasband WS, Eliceiri KW. NIH Image to ImageJ: 25 years of image

analysis. Nat Methods. 2012 Jul;9(7):671–5.

106.   Roy S, Lagree S, Hou Z, Thomson JA, Stewart R, Gasch AP. Integrated Module and

Gene-Specific Regulatory Inference Implicates Upstream Signaling Networks. PLOS

Comput Biol. 2013 Oct 17;9(10):e1003252.

1335

1336