ACSS2 Regulates HIF-2α Degradation through the E3-Ubiquitin Ligase MUL1 in Clear Cell Renal Cell Carcinoma

Zachary A. Bacigalupa[1,2], Whitney A. Brown[1], Evan S. Krystofiak[3], Melissa M. Wolf[1], Rachel A. Hongo[1,2], Madelyn Landis[1,2], Edith K. Amason[1], Kathryn E. Beckermann[1], Jeffrey C. Rathmell[2,4], W. Kimryn Rathmell[1,2]

[1]Department of Medicine, Vanderbilt University Medical Center, Nashville, TN 37232

[2]Vanderbilt Center for Immunobiology, Vanderbilt University Medical Center, Nashville, TN 37232

[3]Cell Imaging Shared Resource, Vanderbilt University, Nashville, TN 37232

[4]Department of Pathology, Microbiology, and Immunology, Vanderbilt University Medical Center, Nashville, TN 37232

Running Title: ACSS2-dependent regulation of HIF-2α in ccRCC

Keywords: ACSS2, ccRCC, HIF-2α, mitochondria, stability

Address Correspondence to:

W. Kimryn Rathmell, M.D., Ph.D.

1161 21st Avenue South, Suite D-3100 Medical Center North

Nashville, TN 37232

Phone: 615-343-8701, Fax: 615-343-2551

Email: Kimryn.Rathmell@VUMC.org

ABSTRACT

Clear cell renal cell carcinoma (ccRCC) is an aggressive kidney cancer driven by *VHL* loss and

aberrant HIF-2α signaling. Acetate metabolism may contribute to this axis by ACSS2-dependent

acetylation of HIF-2α and may provide opportunities to intervention. Here we tested the effects

of pharmacological and genetic manipulation of ACSS2 on HIF-2α, ccRCC cells, and tumors.

ACSS2 inhibition led to HIF-2α degradation and suppressed ccRCC growth *in vitro*, *in vivo*, and

in primary cell cultures of ccRCC patient tumors. This treatment resulted in reduced glucose and

cholesterol metabolism, mitochondrial biogenesis and altered cristae deformation, that are

consistent with loss of HIF-2α. Mechanistically, HIF-2α protein levels are regulated through

proteolytic degradation and we found, in parallel to *VHL*, HIF-2α stability was dependent on

ACSS2 activity to prevent direct interaction with the E3 ligase MUL1. These findings highlight

ACSS2 as a critical upstream regulator of HIF-2α that may be exploited to overcome resistance

to HIF-2α inhibitor therapies.

1 **STATEMENT OF SIGNIFICANCE**

2 We have unveiled ACSS2 as a critical upstream regulator of HIF-2α in ccRCC. Targeting

3 ACSS2 potently promotes HIF-2α degradation via MUL1 to effectively deplete mitochondrial

4 activity and block ccRCC primary tumor models and growth models resistant to HIF-2α inhibitor

5 therapy.

INTRODUCTION

Renal cell carcinoma (RCC) was diagnosed in 76,000 individuals in 2020 in the United States [1]. Clear cell renal cell carcinoma (ccRCC) is the most abundant and aggressive subtype of RCC, accounting for 75% of all RCC cases [2]. ccRCC has been intimately linked to the hereditary condition von Hippel-Lindau (VHL) disease, where patients harbor an inactivating mutation to the *VHL* tumor suppressor gene. In addition to familial associations, 90-95% of ccRCC cases are sporadically derived and the vast majority, upwards of 90%, present with a somatic mutation or suppressive hypermethylation of the promoter to the *VHL* gene [3-8]. The *VHL* gene encodes the E3 ubiquitin ligase pVHL, which functions to regulate the hypoxia inducible factor (HIF) family of transcription factors. Under normal oxygen conditions, prolyl hydroxylases (PHDs) add hydroxyl groups to proline residues on HIF-α proteins, which are then recognized by pVHL and targeted for proteasomal degradation [9]. Hypoxia, or the absence of pVHL, allows HIF-α proteins to accumulate, enter the nucleus and dimerize with HIF-1β/ARNT, and direct a major transcriptional response [10]. In ccRCC, HIF-2α is the major driver of transformation in part due to the loss of the 14q chromosomal arm where the HIF1A locus resides [11, 12]. Indeed, HIF-2α has been shown to directly regulate pathways such as lipid and cholesterol metabolism that fuel the clear cell phenotype [13-17]. Clinically, this has led to the development of small molecule inhibitors targeting HIF-2α interaction with the essential binding partner HIF-1β and HIF-2α transcriptional activity [18-21]. Belzutifan (MK-6482, previously PT-2399) is the first of such molecules to have received FDA approval for the treatment of adult patients with VHL disease who need therapy for ccRCC and other hemangioblastomas [22].

pVHL-mediated degradation of HIF-α proteins remains the most well understood mechanism of HIF-α disposal. However, VHL-independent degradation pathways have been identified, one involving JNK1-mediated stability of the chaperone proteins Hsp90/Hsp70, and another describes an endoplasmic reticulum (ER) stress-directed GSK3β-FBXW1A degradation

axis [23]. More recently, the mitochondrial-tethered E3 ubiquitin ligase, MUL1, was found to

indirectly promote HIF-1α degradation by targeting the HIF-1α degradation complex UBXN7 for

proteasomal destruction [24]. Though MUL1 is best described to target proteins for degradation

via SUMOylation, and K48 or K63 polyubiquitination, it can also aid mitophagy and autophagic

degradation of mitochondrial associated proteins [25]. Localized to the outer mitochondrial

membrane, MUL1 activity is integrated into mitochondrial quality control processes by modifying

proteins such as DRP1 and Mitofusin 1, regulating the function and morphology of mitochondria

[26, 27]. A role for MUL1 to regulate HIF-2α remains uncertain.

Targeting HIF-2α transcriptional activity has proven to be an effective treatment in VHL

syndromic tumors, however, development of resistance makes parallel therapeutic strategies

essential [28]. Resistance to belzutifan may be caused by HIF-2α mutations or the high levels

of HIF-2α that can accumulate in the absence of pVHL and exceed local drug concentration.

HIF-2α acetylation may provide an alternate approach to overcome resistance. Several studies

have described the post-translational addition of an acetyl group to lysines on HIF-2α that

supported protein stability and enhanced downstream signaling [29-32]. In these studies, the

acetylation of HIF-2α, particularly during stress conditions, required the activity of acetyl-CoA

synthetase 2 (ACSS2) [29-31]. ACSS2 is a metabolic enzyme present in both the cytoplasm

and nucleus, which converts acetate into acetyl-CoA in an ATP-dependent reaction. In pre-

clinical cancer models, ACSS2 has proven to be a promising target and has led to the rapid

development of novel small molecule inhibitors [33-38]. ACSS2 has been found to facilitate

epigenetic changes enhancing lysosomal biogenesis and autophagic activity in other biological

models [39], which coupled with its reported interaction with physiologically expressed HIF-2α

make ACSS2 a promising target in the context of ccRCC.

Here we investigate the relationship between ACSS2 activity and constitutive HIF-2α

protein stability in the context of cancer models driven by *VHL* loss. We found that inhibition of

ACSS2 resulted in transcriptional downregulation of genes involved in HIF-2α signaling, as well

as cholesterol biosynthesis and glucose uptake. In addition, we found that ACSS2 inhibition

induced HIF-2α proteasomal degradation even in the absence of pVHL. Mechanistically, ACSS2

inhibition or silencing stimulated the expression of MUL1, which formed a direct complex with

HIF-2α to provide a potential VHL-independent and HIF2-α-specific mechanism to disrupt HIF-

dependent transcription. Electron microscopy revealed ACSS2 inhibition, as well as HIF-2α

silencing, caused massive mitochondrial deformation likely to result in catastrophic loss of

function. Importantly, pharmacological inhibition of ACSS2 effectively blocked cancer cell growth

in clinical samples and the HIF-2α inhibitor resistant 769-P cell line. Together, these findings

demonstrate a critical role for ACSS2 to support ccRCC growth by interfering with direct

regulation of HIF-2α stability by MUL1 to maintain the activity of key metabolic pathways.

ACSS2 is, therefore, a novel therapeutic target for HIF-2α driven ccRCC.

RESULTS

ACSS2 is essential for ccRCC growth and proliferation.

16   ACSS2 has remained largely unexplored as a potential therapeutic target in ccRCC. To

examine the role of ACSS2 in ccRCC, we first performed a dose response, time course study

with a pharmacological inhibitor of ACSS2 in HKC (normal kidney epithelia), 786-O ($VHL^{-/-}$

ccRCC), and A498 ($VHL^{-/-}$ ccRCC) cell lines. HKC cells were affected only at the highest dose,

but ACSS2 inhibition led to a significant reduction in cell growth for the ccRCC cell lines (Figure

1A and 1B, Supplemental Figure 1A). We next analyzed the effect of ACSS2 inhibition on

proliferation by measuring BrdU incorporation. Similarly, while HKC cells were only significantly

affected after 48 hours of 5 μM and 10 μM doses, proliferation was significantly reduced in

ccRCC cell lines at 24-hours with all doses of the inhibitor, (Figure 1C and 1D, Supplemental

Figure 1B). Tumor sphere formation assays also revealed a striking difference in the ability of

786-O cells to form spheres when treated with ACSS2i (Supplemental Figure 1C). To assess

the impact of ACSS2 on ccRCC growth more directly, we transduced the 786-O cell line to

stably express a doxycycline-inducible shRNA construct targeting a control sequence or

ACSS2. After 48 hours, we assessed cell growth via crystal violet staining and observed

significant reduction in cell growth resulting from ACSS2 depletion (Figure 1E). Like the

inhibitor, after 24 hours, 786-O cells expressing shACSS2 displayed significantly reduced BrdU

incorporation (Figure 1F).

With our data confirming ACSS2 is required for ccRCC growth in vitro and that ccRCC

cell lines are more sensitive to ACSS2 inhibition, we next investigated the ability of ACSS2

inhibition to block tumor growth in vivo. Doxycycline induction of shACSS2 in mice bearing

subcutaneous 786-O tumors resulted in a significant reduction in tumor growth (Figure 1G) and

final tumor weight (Figure 1H). Similarly, mice bearing subcutaneous 786-O tumors receiving

daily treatment with 15 mg/kg ACSS2i exhibited a remarkable decrease in tumor growth and

final tumor weight (Figure 1I and 1J). In neither model did targeting ACSS2 result in any

observable toxicity such as weight loss (Supplemental Figure 1D and 1E).

Immunohistochemistry staining for CD31 and Periodic Acid Schiff revealed that targeting

ACSS2 also reduced vascularization and glycogen deposition (Figure 1K-L and Supplemental

Figure 1F-G), which are commonly associated clinical features of ccRCC linked to HIF

signaling. Together, these data identify ACSS2 as a key contributor to tumor growth and a

druggable vulnerability in ccRCC.

HIF-2α gene and protein expression is regulated by ACSS2 activity.

In ccRCC, aberrant HIF-2α signaling configures widespread signaling and metabolic

changes. We hypothesized that these effects would be tied to the VHL-HIF axis. To test the role

of ACSS2 to regulate essential metabolic signaling pathways in ccRCC, we performed

transcriptomic analysis using the NanoString Metabolic Pathways probe set on 786-O cells

transduced with shRNA targeting a control sequence or shACSS2 (Figure 2A). Intriguingly,

differential expression analysis of this dataset identified a HIF-2α RCC Signaling gene set

consisting of *EGLN3*, *EPAS1*, *PDGFB*, *SLC2A1*, and *VEGFA* as the most significantly

downregulated pathway in response to ACSS2 deficiency (Figure 2B), suggesting that HIF-2α

transcriptional activation was disrupted. To confirm these findings, we performed qRT-PCR on

786-O cells treated with 5 µM of ACSS2i for 24 hours (Supplemental Figure 2A) or transduced

with shHIF2A (Supplemental Figure 2B) and observed a significant reduction in *Epas1* gene

expression.

Because HIF-2α is largely regulated at the protein level, we hypothesized ACSS2 may

impact HIF-2α protein. Consistent with this model, we observed a robust decrease in HIF-2α

protein in 786-0 cells transduced with shACSS2 (Figure 2C). HIF-2α protein expression was

also suppressed by pharmacologic ACSS2 inhibition to result in reduced expression of known

HIF-2α targets (Figure 2D). Notably, HIF-2α protein expression was not impacted by ACSS2

inhibition in the HKC cell line (Supplemental 2C). To further test the involvement of ACSS2

activity for the observed changes, we stably overexpressed ACSS2 in 786-O cells and

assessed protein expression, where we observed an increase in the expression of HIF-2α and

its downstream signaling network (Figure 2E). Additionally, a BrdU proliferation assay confirmed

that overexpression of ACSS2 in 786-O cells also resulted in enhanced proliferation (Figure 2F).

Next, we tested whether overexpressing ACSS2 was sufficient to enhance tumorigenic potential

and determine if any observed effect was dependent on HIF-2α utilizing an anchorage-

independent growth assay. 786-O cells were transduced to express V5-ACSS2 or the empty

vector followed by a second transduction to target HIF2A with shRNA (Supplemental Figure

2D). After 21 days of growth, it was determined that ACSS2 overexpression increased the size

and number of colonies, but this increase was fully mitigated when *HIF2A* was targeted with

shRNA (Figure 2G).

ACSS2 activity mediates HIF-2α stability

Previous studies have identified acetylated lysine residues on HIF-2α, which confer

protein stability and yield optimal signaling, that are acquired from acetyl-CoA produced from

ACSS2 [29-32]. To better understand the mechanism by which ACSS2 may regulate HIF-2α

protein expression, we performed a 1-hour incubation with the proteasomal inhibitor MG132

followed by the addition of DMSO or 5 µM ACSS2i for 24 hours. Interestingly, 786-O cells

receiving the MG132 pre-treatment prior to ACSS2i displayed a significant rescue in HIF-2α

expression (Figure 3A and Supplemental 3A) suggesting ACSS2 inhibition enhances HIF-2α

proteasomal degradation, even in *VHL* deficient 786-O cells.

Recently, several studies have linked the E3 ubiquitin ligase, MUL1, to the indirect

regulation of HIF-1α degradation by the E3-ligase co-factor UBXN7 [24]. Consistent with a

potential role for MUL1, analysis of TCGA and CPTAC samples via the UALCAN database

show that MUL1 transcript abundance and protein expression are significantly reduced when

compared to normal adjacent tissue (Supplemental Figure 3B and 3C). Survival analysis

performed on the KIRC dataset assessing the effect of MUL1 expression on patient outcome

identified a correlation between low MUL1 expression and reduced survival (Figure 3B).

Intriguingly, depleting MUL1 via shRNA provided 786-O cells with resistance to treatment with

ACSS2i or PT-2385, enabling successful anchorage-independent growth (Figure 3C and

Supplemental Figure 3D). Considering these findings, we next analyzed the effect of targeting

ACSS2 on MUL1 expression. Pharmacologically or genetically targeting ACSS2 in 786-O cells

increased MUL1 expression (Figure 3D and 3E). In the HKC cell line, we also observed an

increase in MUL1 expression in response to ACSS2 inhibition despite having no impact on HIF-

2α expression, which was possibly due to continued expression of pVHL (Supplemental Figure

3E). To test whether MUL1 could interact with and direct the degradation of HIF-2α, we

immunoprecipitated MUL1 from 786-O cells treated with DMSO or 5 µM ACSS2i for 24 hours

and observed HIF-2α and MUL1 to interact in a complex (Figure 3F). These results show that

ACSS2 is a critical and direct mediator of MUL1-directed HIF-2α protein degradation. Moreover,

these findings highlight ACSS2 and MUL1 as means to target HIF-2α in ccRCC.

ACSS2 activity supports glucose and cholesterol metabolism.

The link between hypoxia signaling and metabolism has been well-characterized and

aberrant HIF-2α signaling has been shown to promote glucose and cholesterol metabolism in

ccRCC. To address the impact of ACSS2 on metabolic pathways, we performed a

transcriptomic analysis. Differential gene expression analysis of 786-O cells treated with DMSO

or 5 µM ACSS2i for 24 hours identified cholesterol metabolism as the most significantly altered

pathway (Figure 4A). Notable changes within the cholesterol metabolism pathway

corresponding to ACSS2 inhibition included the decrease in HMG-CoA synthase (HMGCS1),

which synthesizes HMG-CoA from acetyl-CoA, HMG-CoA reductase (HMGCR), whose catalytic

activity produces mevalonate in the rate-limiting step of cholesterol synthesis, and 7-

dehydrocholesterol reductase (DHCR7) and 24-dehydrocholesterol reductase (DHCR24) which

perform the final reaction in cholesterol biosynthesis (Supplemental Figure 4A). Functionally, we

measured total cholesterol levels in 786-O cells treated with DMSO or 5 µM ACSS2i for 24

hours and found a significant reduction in response to ACSS2 inhibition (Figure 4B).

Given the link between hypoxia signaling and glucose metabolism, we next investigated

the effect of ACSS2 inhibition on *SLC2A1* (Glut1) and hexokinase II (*HKII*) gene expression. To

this end, we performed RT-PCR on 786-O cells treated with DMSO or 5 µM ACSS2i for 24

hours and found the transcripts of *SLC2A1* and *HKII* to be significantly reduced in response to

ACSS2 inhibition (Figure 4C). Next, we tested if the observed mRNA changes correspondingly

altered protein expression. Following 24 hours of treatment with DMSO or ACSS2i, HIF-2α, as

well as GLUT1, HKII, SREBP-1, and SREBP-2 which are more commonly associated with HIF-

1α [40, 41], protein levels were all decreased in response to ACSS2 inhibition (Figure 4D).

Conversely, overexpression of ACSS2 in 786-O cells enhanced expression of HIF-2α, GLUT1,

and HKII (Figure 4E). A time course measurement on glucose present in the media of 786-O

cells treated with DMSO or 5 µM ACSS2i revealed a significant increase in glucose present in

the media after 48 hours of ACSS2i treatment, indicating an impaired ability of the cells to

uptake and consume glucose (Supplemental 4B). Additionally, we quantified the acetate

concentrations in the media of 786-O cells following a 24-hour dose response with ACSS2i and

found that ACSS2 inhibition significantly enhanced the amount of acetate present in the media

(Supplemental Figure 4C). Similarly, 786-O cells transduced with the doxycycline-inducible

shRNA system saw an increase in acetate in the media when shACSS2 was induced, whereas

a significant decrease in acetate was observed when ACSS2 was overexpressed

(Supplemental Figure 4D). These results confirm that altering ACSS2 activity directly impacts

acetate consumption. Furthermore, our results demonstrate that ACSS2 activity transcriptionally

regulates the rate-limiting enzymes in cholesterol synthesis and glucose metabolism, which are

likely to have widespread effects on the cell.

Targeting ACSS2 elicits cancer cell-specific mitochondrial defects.

Mitochondrial quality is intimately linked to their function and ability to facilitate metabolic

processes [42, 43]. Considering the magnitude by which targeting ACSS2 has shown to inhibit

glucose metabolism and cholesterol biosynthesis, we hypothesized that inhibiting ACSS2 would

selectively disrupt mitochondrial homeostasis in cancer cells. To test the effects on

mitochondrial morphology electron microscopy was performed on HKC and 786-O cells treated

with DMSO for 72 hours or with 5 µM ACSS2i for 24, 48, and 72 hours (Figure 5A and 5B).

Strikingly, the mitochondria in 786-O cells began to transform as early as 24 hours to become

smaller and circular, a process not detected in the mitochondria of HKC cells (Figure 5A and

5B). Additionally, the cristae no longer retained their typical morphology and became circular

(Figure 5B). These results were further corroborated when we looked at the expression of

proteins involved in cristae structure such as Opa-1, Mitofusin 1, Mitofusin 2, and Drp1, after

treatment with 5 µM ACSS2i for 48 hours result in a loss of expression specifically in 786-O

cells (Supplemental 5A and 5B). We also observed a cancer cell-specific reduction in the

expression of PGC1β, which has been shown to control mitochondrial biogenesis [44] (Figure

5A and 5B).

The impact of ACSS2 inhibition on mitochondrial biogenesis was next tested. Applying a

colorimetric assay that detects both the nuclear encoded mitochondrial gene SDH-A and the

mitochondrial encoded COX-I, we discovered a significant, dose-dependent decrease in COX-I

after 24 hours of ACSS2i treatment in the 786-O cells. This had no impact, however, on the

HKC cells (Supplemental 5C and 5D). To determine if the observed mitochondrial phenotype

was specific to ACSS2, we performed electron microscopy on 786-O cells where shRNA

targeting ACSS2 had been induced with doxycycline for 48 hours and we observed shACSS2

induction had the same morphological changes that pharmacological inhibition produced (Figure

5C). Additionally, shACSS2 had a similar effect on mitochondrial biogenesis (Supplemental

Figure 5E). However, the reduced expression of cristae structural proteins observed in response

to ACSS2i was not as robust with shACSS2, although PGC1β induction was lost (Supplemental

Figure 5F). Together, these data suggest that ACSS2 activity serves an integral role to maintain

the quality of mitochondria in *VHL*-mutated cancer cells, possibly through a PGC1β-dependent

mechanism.

We next questioned whether mitochondrial maintenance influenced by ACSS2 activity

was dependent on HIF-2α. To address this question, electron microscopy was performed on

786-O cells overexpressing ACSS2 which had been transduced to express shControl or

shHIF2A. Mitochondria from cells overexpressing ACSS2 had a characteristically normal

appearance with defined cristae; however, when shHIF2A was introduced, we observed a return

of the small, circular mitochondrial phenotype with deformed cristae (Figure 5D). Interestingly,

treatment with the HIF-2α inhibitor PT-2385 did not result in a morphological defect in the

mitochondria (Figure 5E), possibly indicating that PT-2385 has lower efficacy than targeting with

shRNA. In sum, these results suggest a role for HIF-2α in maintaining healthy mitochondria and

provide further insight into the scope of the ACSS2-HIF-2α axis.

ACSS2 inhibition selectively impedes cancer cell growth and HIF-2α expression in

ccRCC patient samples.

The HIF-2α inhibitor belzutifan was shown to be highly effective in patients with von

Hippel-Lindau disease and has recently gained FDA approval for the treatment of adult patients

with the disease who need treatment for RCC. As our previous results demonstrated that

targeting ACSS2 is an effective strategy to selectively target HIF-2α, we next moved our

experiments into primary cell cultures isolated from ccRCC patients. A time-course, dose-

response treatment was performed on matching normal adjacent tissue and cancer cells from

three patients and cancer cells again observed to be significantly more sensitive than normal

epithelial cells to ACSS2 inhibition (Figure 6A-B and Supplemental Figure 6A-D). Next, we

explored the effect of ACSS2 inhibition on HIF-2α expression and found that 24 hours of

treatment at all doses of ACSS2i in these clinical specimens was effective at reducing HIF-2α

expression in a cancer cell-specific manner (Figure 6C and 6D). Additionally, HKII expression

increased and both findings were supported by transcriptomic analysis to corroborate our

previous findings that ACSS2 inhibition reduces the expression of genes involved in cholesterol

biosynthesis (Figure 6C-D and Supplemental Figure 6E-F). Lastly, we explored whether ACSS2

inhibition could be an effective means to treat ccRCC that are resistant to HIF-2α inhibitor

therapy. To this end, we performed a dose-response, time-course ACSS2i treatment in the

previously described HIF-2α inhibitor resistant 769-P ccRCC cell line and found that ACSS2i,

when provided as single agent, was able to block the growth of these cells (Figure 6E).

Interestingly, a slight synergistic effect was observed when treating 769-P cells with the ACSS2

inhibitor and PT-2385, suggesting that providing ACSS2i in conjunction with HIF-2α inhibition

could be effective at circumventing resistance to deliver a more effective suppression of HIF-2α

driver signals (Figure 6E). Altogether, these results demonstrate that our in vitro and in vivo

studies translate to ccRCC clinical samples, highlighting ACSS2 as a novel way to target HIF-2α

and major metabolic pathways specifically in cancer cells.

DISCUSSION

ACSS2 has recognized functions in the dynamic interplay of stress responses, including

interaction with HIF-2α in the setting of adaptive responses to oxygen and nutrient limited states

[29, 30, 35, 36, 45]. In clear cell renal cell carcinoma, the pseudohypoxic state is defined by high

level, constitutive HIF-2α signaling that is unimpeded by normal stress signals. We sought to

directly interrogate the interaction of ACSS2-mediated acetylation to maintain the stability of

HIF-2α in the absence of *VHL* and provide a key driver of these tumors [46-49]. Recent work

has advanced a HIF-2α specific transcriptional inhibitor for treatment of *VHL*-associated renal

cell carcinomas although resistance can develop over time[18, 50]. Parallel strategies to block

the HIF-2α node, such as targeting alternate mechanisms of HIF-2α protein stability will be

necessary for adequate disease control.

By applying various strategies to alter ACSS2 activity, we showed that ccRCC cells have

some level of dependence on ACSS2 to support HIF-2α expression, stability, and signaling,

which makes them inherently more vulnerable to ACSS2 inhibition. Importantly, we also

established that ACSS2 inhibition was effective to block the growth of cancer cells derived from

ccRCC patients, as well as the HIF-2α inhibitor resistant 769-P cell line. Our data suggest that

treatment with an ACSS2 inhibitor could have a potent effect at treating ccRCC patients who

have a *VHL* mutation and could possibly have a synergistic effect when used in combination

with HIF-2α inhibition. Novel small molecules targeting ACSS2 are in development [34] and

being characterized in preclinical models of cancer. Indeed, while HIF-2α inhibition with

belzutifan has been revolutionary for VHL syndrome, the effect is slow. Also, although each

cancer in these scenarios displays HIF-2a activation and transcriptional dependencies, simple

inhibition of dimerization does not promote cell death promptly. We show here that acetylation

via ACSS2 is important for HIF-2α stabilization, that inhibiting ACSS2 reduces transcription of

EPAS1 as well as induces protein degradation via a secondary mechanism. This more potent

blockade effectively shuts down hypoxia transcription and selectively impairs mitochondrial

function in a way that proves to be lethal to cells that are dependent on HIF-2α signaling.

Vhl is mutated in over 70% of ccRCC and aberrant HIF-2α signaling persists due to a

lack of an effective degradation mechanism. The identification and characterization of

alternative HIF-2α degradation pathways could manifest into clinically relevant findings.

Recently, the mitochondria-associated E3 ubiquitin ligase MUL1 has been shown to indirectly

regulate the degradation of HIF-1α through its action of UBXN7, and was also found to promote

autophagy and be suppressed in ccRCC cancer cells [24, 51]. Given the high level of homology

between HIF-1α and HIF-2α and the link made to MUL1 and ccRCC, we rationalized

interrogating a link between ACSS2 and MUL1 as a mechanism to alter HIF-2α stability.

Identifying that MUL1 directly interacts with HIF-2α and that ACSS2 inhibition can enhance

MUL1 expression and its interaction with HIF-2α, implicates MUL1 as a potential mechanistic

link as to how ACSS2 activity regulates HIF-2α in a tumor lacking *Vhl* (Supplemental Figure

6G). Beyond direct polyubiquitin-mediated degradation, it has also been demonstrated that

MUL1 can contribute to the degradation of targets such as Mitofusin 1 via mitophagy [25, 27].

Interestingly, it was recently shown that metabolic stress induces the nuclear localization of

ACSS2, where it promotes acetylation and interacts with TFEB to facilitate the transcription of

genes involved in lysosomal biogenesis and autophagy [39]. Taking these findings into

consideration with our own, it would be intriguing to investigate the involvement of mitophagy in

the ACSS2-dependent degradation of HIF-2α.

Acetate is a major precursor to produce acetyl-CoA, particularly under times of metabolic

stress. While citrate-derived acetyl-CoA has been shown to be more heavily utilized in lipid

synthesis [52, 53], acetyl-CoA produced from ACSS2 is thought to have more of a role in post-

translational and epigenetic regulation [29, 31, 38]. Though the findings we describe here

remain focused on the relationship between ACSS2 and HIF-2α, the role of ACSS2 in

supporting histone acetylation and global transcriptional regulation cannot be ignored. It is likely

that manipulating ACSS2 activity causes shifts in epigenetic patterns, which could have a much

more global impact on biological processes and signaling pathways. Future studies will be

aimed at dissecting the role of ACSS2 in epigenetic maintenance in ccRCC.

Exploration into the clinically translatable effects of pharmacologically targeting ACSS2

has yet to commence. The results of our study demonstrate ACSS2 inhibition is a well-tolerated

and effective strategy to ablate HIF-2α expression and signaling in pre-clinical models of

ccRCC. Furthermore, when we employed a cell line known to be resistant to HIF-2α inhibition,

targeting ACSS2 proved capable of overcoming this resistance. This is the first study in ccRCC

to highlight ACSS2 as an integral mediator of HIF-2α protein stability likely by regulating the

post-translational modification landscape and MUL1-directed degradation. These novel methods

of regulating HIF-2α could provide a much-needed approach to complement and enhance the

efficacy of treatment in populations resistant to HIF-2α inhibition.

METHODS

Materials Availability

Pharmacological agents used in this study were acquired commercially and available at

Selleckchem. Mice used in this study were acquired commercially and available at Jackson

Laboratories. Stable cell lines carrying targeted shRNA or with different constructs are available

through establishment of Material Transfer Agreement between Vanderbilt University Medical

Center and requesting institution.

Experimental Models

15  *Cell Cultures*

16  HKC cells were a gift from L.C. Racusen. HEK293T, A498, 786-O, and 769-P cell lines

were obtained from the ATCC. Prior to experimentation, all cell lines were tested for

Mycoplasma contamination using the ATCC Universal Mycoplasma detection kit (ATCC, catalog

number 30-1012K). HEK293T, HKC, and A498 cell lines were cultured using Gibco Dulbecco's

Modified Eagle Medium (DMEM) supplemented with 10% fetal bovine serum (FBS), 1% L-

glutamine, and 10 mL/L penicillin/streptomycin (Sigma). 768-O and 769-P cell lines were

cultured using Gibco RPMI-1640 supplemented with 10% FBS, 1% L-glutamine (Sigma), and 10

mL/L penicillin/streptomycin (Sigma). All cells were maintained at 37°C in a 5% CO_2 incubator.

Animal Studies

6-week old male NOD-scid IL2Rg[null] (NSG) mice were purchased from The Jackson Laboratory (catalog number 005557) and utilized in all tumorigenesis xenograft models. Mice were housed five to a cage in a temperature- and humidity-controlled space (20-25°C, 45%–64% humidity) with regulated water and lighting (12 h light/dark) within the animal facility. All in vivo mouse studies designed and performed in accordance to animal protocols that were approved by the Institutional Animal Care and Use Committee of VUMC. Prior to inoculation, 786-O cells were cultured as described above then trypsinized, counted, and resuspended in growth media and Matrigel at a 1:1 ratio. For all in vivo experiments, NSG mice were subcutaneously injected with 100 µl of the cell suspension providing $5x10^6$ cells. Mice in pharmacological studies were then subjected to daily treatment with vehicle control or 15 mg/kg ACSS2i delivered via intraperitoneal injection. Alternatively, mice which were inoculated with 786-O cells transduced to express the pTRIPZ doxycycline-inducible shRNA system were provided a 200 mg/kg doxycycline rodent diet (Bio-Serv) and allowed to feed ad libitum. Tumor burden was monitored via weekly manual, digital caliper measurement with intermittent measurements taken as needed. Mice were euthanized once the first tumor reached size endpoint (1,000 mm^3) or if discomfort was observed as outlined in the IACUC approved protocol.

Patient Samples

Primary cell cultures were derived from tumors of patients diagnosed with clear cell renal cell carcinoma being treated at Vanderbilt University Medical Center. Informed written consent was collected from all patients whose samples were used in this study and all samples were processed and utilized in accordance with the IRB protocol (151549). Single cell suspensions were generated from the tumors and were subsequently cultured in Gibco RPMI-1640 media supplemented with 10% FBS, 1% L-Glutamine (Sigma), B-27 supplement (Gibco; 10 mL, 50x

stock), and 1x Antibiotic-Antimycotic (Gibco). Fresh growth media was provided every other day

for approximately 10 days until a monolayer of epithelial cells was established, at which point

the primary cells were able to be passaged.

Method Details

Protein Extraction and Western Blotting

Cells were harvested and suspended in RIPA buffer supplemented with 1x Halt protease

inhibitor cocktail (ThermoFisher) and subjected to mechanical needle lysing with a 25G needle

(BD) while being kept on ice. Cell lysates were pelleted via centrifugation at 14,000 x g for 5

minutes at 4°C and supernatants were transferred to fresh Eppendorf tubes for immediate

analysis via SDS-PAGE or preservation at -20°C. Protein concentrations were determined using

the BCA protein quantification method. For SDS-PAGE analysis, 50 µg of protein per sample

was loaded into and run on a 4-20% gradient polyacrylamide gel (Bio-rad), followed by transfer

onto a PVDF membrane. The membranes were then subjected to immunoblotting with the

primary antibodies found in **Supplemental Table 1**. For western blot analysis, β-Actin served as

the loading control.

Immunoprecipitation

Cell lysates were prepared as described above and 500 µg of total protein per sample

was first cleared via incubation with protein A agarose beads at 4°C for 1 hour then spun down

at 3,500 x g for 10 minutes at 4°C. The pre-cleared lysates were then transferred to fresh

Eppendorf tubes and immunoprecipitation commenced via incubation with the primary antibody

overnight at 4°C. The following day, 30 µl of a 50% protein A agarose bead slurry was added to

the immunoprecipitation reactions and continued incubating for 2 hours at 4°C on a rotator. The

immunoprecipitations were then microcentrifuged at 3,500 x g for 10 minutes at 4°C to collect

the beads, which were then washed five times with Ripa buffer supplemented with protease

inhibitors. After the final wash, the pelleted beads were resuspended with 3X SDS sample

buffer, vortexed briefly, then microcentrifuged at 14,000 x g for 1 minute at room temperature.

Samples were then heated at 95°C for 5 minutes, microcentrifuged at 14,000 x g for 1 minute at

room temperature and loaded into a gel for SDS-PAGE to then be analyzed via western blotting.

Lentiviral Transduction

HEK293T cells were used for packaging the vectors in Supplemental Table 2 into

lentivirus. HEK293T cells were allowed to reach 70% confluency and on the day of transfection,

1 hour prior to transfection growth media was replaced with Optimem. For each plate, a

transfection cocktail was was prepared using 20 µg of the experimental plasmid (**Supplemental**

Table 2)**,** 15 µg PAX2, 6.5 µg of pMD2.G, 62 µl of $CaCl_2$, and sterile dH_2O to bring the total

volume to 500 µl, then 500 µl of 2X HBS was added and mixed by pipetting. The transfection

media was then added to the HEK293T cells, gently swirled, and incubated for 5-7 hours at

37°C. Transfection medium was then discarded and replaced with 10-12 mL of fresh growth

media and the cells were allowed to produce viral particles for 48-72 hours. The lentiviral

supernatants were then collected and passed through a 0.45 micron filter to be used

immediately or stored at -80°C for later use.

Lentiviral transduction of 786-O cells was performed by mixing 1 mL lentivirus, 6.5 µl of

10 mg/mL polybrene, and 7 mL growth media and incubating for 24 hours at 37°C. The next

day, transduction media was replaced with fresh growth media and cells were returned to 37°C

incubator for another 24 hours. On the following day, the media of the transduced cells was

supplemented with the antibiotic corresponding to the selectable antibiotic resistance marker in

each plasmid for 24 hours, and the resistant clones were expanded for experimental use.

RNA Isolation and qRT-PCR

Total RNA extracts were isolated and purified using the RNeasy RNA isolation kit

(Qiagen) and converted to cDNA via reverse transcription (Applied Biosystems, High Capacity

cDNA Reverse Transcription kit). For each experiment, 25 ng of cDNA per sample was

combined with Taqman Universal Master Mix (Applied Biosystems) and RNAse-free water in a

96-well reaction plate and qRT-PCR was performed using a Bio-Rad CFX96 thermocycler.

Primers used for qRT-PCR reactions were Epas1 (Hs01026149_m1), HKII (Hs00606086_m1),

SLC2A1 (Hs00892681_m1), and 18S rRNA (Hs99999901_s1). Data analysis was performed

using CFX Manager software (Bio-Rad) and transcript abundance was determined via the ΔΔCt

with normalization to 18S rRNA.

Metabolic Gene Expression

To analyze changes in expression of genes involved in metabolism, we extracted RNA

as described above and performed a gene expression assay using the Nanostring nCounter

Metabolic Pathways panel. The assay was performed by Vanderbilt Advanced Technologies for

Genomics (VANTAGE) core facility following the manufacturer's recommended procedures.

RNA concentrations were determined via Qubit and normalized to 20ng/ul. For each sample

hybridization reaction, we used 100 ng of total RNA and hybridizations were performed for a

total of 20 hours. Analysis of raw data was performed using the nSolver Analysis Software 4.0

(Nanostring). For all analyses, the background threshold parameters were set to the mean of 8

negative control spike-in genes and mRNA expression was normalized to the geometric mean

of 6 positive control spike-in genes and 20 housekeeping genes (ABCF1, AGK, COG7, DHX16,

DNAJC14, EDC3, FCF1, G6PD, MRPS5, NRDE2, OAZ1, POLR2A, SAP130, SDHA, STK11IP,

TBC1D10B, TBP, TLK2, UBB, and USP39).

Data was analyzed by ROSALIND® (https://rosalind.bio/), with a HyperScale

architecture developed by ROSALIND, Inc. (San Diego, CA). Read Distribution percentages,

violin plots, identity heatmaps, and sample MDS plots were generated as part of the QC step.

The limma R library was used to calculate fold changes and p-values and perform optional

covariate correction. Clustering of genes for the final heatmap of differentially expressed genes

was done using the PAM (Partitioning Around Medoids) method using the fpc R library that

takes into consideration the direction and type of all signals on a pathway, the position, role and

type of every gene, etc. Hypergeometric distribution was used to analyze the enrichment of

pathways, gene ontology, domain structure, and other ontologies. The topGO R library, was

used to determine local similarities and dependencies between GO terms in order to perform

Elim pruning correction. Several database sources were referenced for enrichment analysis,

including Interpro, NCBI, MSigDB, REACTOME, WikiPathways. Enrichment was calculated

relative to a set of background genes relevant for the experiment.

Crystal Violet Cell Growth Assay

To assess the impact of the experimental conditions on cell growth, 2.5×10^4 or 1×10^5

cells were seeded into 12-well or 6- well plates, respectively, and allowed to adhere for 24

hours. On the following day, growth media was supplemented with vehicle, 1 µg/mL

doxycycline, or various doses of the ACSS2 inhibitor and the effect on growth was monitored

over time by staining with a 1% crystal violet solution prepared in 20% methanol and 80% dH_2O.

After imaging, the crystal violet stain was then stripped using a 1% deoxycholate solution, which

was transferred to a 96-well plate and read on a Promega GloMax plate-reader at 560 nm to quantify the crystal violet staining.

BrdU Cell Proliferation Assay

To determine the impact of experimental conditions on proliferation, we used a BrdU cell proliferation ELISA kit (Abcam, catalog number ab126556) following the manufacturer's suggested protocol. Briefly, cells were seeded at a density of 1 x 10^4 cells/well in normal growth media. Additionally, a duplicate set of cells to those in the experiment were seeded and did not receive BrdU labeling reagent, while several wells were also left absent of cells to thoroughly account for background. For 24-hour measurements, growth media was supplemented with ACSS2 inhibitor or doxycycline, where applicable, and allowed to incubate for 22 hours at 37°C. BrdU labeling solution was then added for the final 2 hours of the 24-hour incubation before continuing the ELISA protocol. For 48-hour measurements growth media was supplemented with ACSS2 inhibitor and incubated for 24 hours at 37°C before adding the BrdU labeling solution for the final 24 hours of the 48-hour incubation. Following BrdU labeling, cells were fixed for 30 minutes with 1X fixing solution, washed 3 times with 1X wash buffer, and then incubated with detector antibody for 1 hour at room temperature. After incubation with the detector antibody, cells were washed 3 times with 1X wash buffer followed by the addition of 1X Peroxidase Goat Anti-Mouse IgG Conjugate for 30 minutes at room temperature. The ELISA reactions were washed 3 more times with 1X wash buffer and a final time with distilled water. After patting plates dry, TMB Peroxidase substrate was added and reactions incubated for 30 minutes at room temperature in the dark. The reactions were stopped via the addition of stop solution and BrdU incorporation was quantified by reading the plates at 450 nm on a GloMax plate-reader.

Tumor Sphere Formation Assay

786-O cells were seeded onto 12- well ultra-low attachment plates (Corning) at a density of 1 x 10^3 cells per well. Cells were subjected to treatment immediately following plating and fresh growth media with vehicle or the ACSS2 inhibitor was provided every other day. Sphere formation was monitored via microscope daily and after 7 days, images were captured at 4X and 10X magnification on a Keyence BZ-X800 microscope.

Anchorage-Independent Growth Assay

Tumorigenic potential was evaluated using an anchorage-independent growth assay. Briefly, a 2.5% agarose solution was prepared using low-melt agarose and autoclaved dH_2O. This solution was diluted 1:4 in growth media and used to coat 6-well plates which were returned to the 37°C incubator and allowed to set, while cells were trypsinized and counted. For each condition, 1 x 10^5 cells were resuspended in 9 mL of growth media and 1 mL of a 3% agarose solution (0.3% final concentration) and 1 mL (1 x 10^4 cells) of the cell solution was added per well. The top layer was allowed to set for an hour at 37°C at which point growth 2 mL growth media was added to each well. Growth medium and treatments, where applicable, were exchanged every other day and colony formation was monitored via microscope. Images of colonies were captured using EVOS imaging system.

Immunohistochemistry Staining and Imaging

Tumor specimens were harvested immediately following animal euthanasia. Each tumor was washed with cold PBS and then fixed overnight with 10% neutral buffered formalin. The

tumors were then paraffin-embedded, sectioned, and stained by the Translational Pathology

Shared Resource facility at VUMC using standard IHC procedures. Images of stained tumor

sections were obtained using an Olympus light microscope and "name" imaging software.

HIF-2α Protein Stability Assay

To assess the impact of ACSS2 inhibition on HIF-2α protein stability, 786-O cells were

treated with 1 µM of the proteasomal inhibitor MG132 for 1 hour. After the initial incubation with

MG132, growth media was exchanged and supplemented with DMSO or the ACSS2 inhibitor for

24 hours. Upon completion of the treatments, cell lysates were prepared as described above

and analyzed via SDS-PAGE and western blotting. HIF-2α expression was quantified using

ImageJ and normalized to β-Actin.

GC/MS for Cholesterol Quantification

Lipids were extracted using a previously described method [54]. Briefly, the extracts

were filtered, and lipids recovered in the chloroform phase. Individual lipid classes were

separated by thin layer chromatography using Silica Gel 60 A plates developed in petroleum

ether, ethyl ether, acetic acid (80:20:1) and visualized by rhodamine 6G. Phospholipids,

diglycerides, triglycerides and cholesteryl esters were scraped from the plates and methylated

using BF3 /methanol as described previously [55]. The methylated fatty acids were extracted

and analyzed by gas chromatography. Gas chromatographic analyses were carried out on an

Agilent 7890A gas chromatograph equipped with flame ionization detectors, a capillary column

(SP2380, 0.25 mm x 30 m, 0.25 µm film, Supelco, Bellefonte, PA). Helium was used as a

carrier gas. The oven temperature was programmed from 160 °C to 230 °C at 4 °C/min. Fatty

acid methyl esters were identified by comparing the retention times to those of known

standards. Inclusion of lipid standards with odd chain fatty acids permitted quantitation of the

amount of lipid in the sample. Dipentadecanoyl phosphatidylcholine (C15:0), diheptadecanoin

(C17:0), trieicosenoin (C20:1), and cholesteryl eicosenoate (C20:1) were used as standards.

For total cholesterol, internal standard (5-a-cholestane) was added to a portion of the lipid

extract and then saponified at 80 °C in 1 N KOH in 90% methanol for 1 hour. The

nonsaponifiable sterol was extracted into hexane, concentrated under nitrogen, and then

solubilized in hexane to inject onto the gas chromatograph. For unesterified cholesterol, internal

standard is added to a portion of the lipid extract, concentrated under nitrogen and then

solubilized in hexane to inject onto the gas chromatograph. The Agilent 7890A gas

chromatograph was equipped with an HP-50+ column (0.25 mm i.d x 30 m, Agilent) and a flame

ionization detector. The oven temperature was programmed from 260 °C to 280 °C and helium

was used as the carrier gas [56].

Acetate Quantitation Assay

To monitor ACSS2 activity, acetate concentrations in the growth media were measured

over time using an acetate quantitation colorimetric assay (BioVision, catalog number K658)

using the manufacturer's suggested protocol. Briefly, cells were seeded at a density of 2.5 x 10^5

cells/well in a 6-well plate. The following day fresh growth media was added to the cells, as well

as a blank well, supplemented with ACSS2i or doxycycline, where applicable, and incubated for

24 hours at 37°C. Growth media was collected after the 24-hour incubation and immediately

processed for analysis with the colorimetric assay. For each sample, 25 µl of media was

combined with 25 µl of assay buffer and 50 µl of the reaction buffer using the manufacturer

provided reagents and guidelines. Following a 40-minute incubation at room temperature,

acetate concentration measurements were obtained via colorimetric detection at OD450 nm

using a GloMax plate-reader.

Electron Microscopy

Electron microscopy was performed in part through the use of the Vanderbilt Cell Imaging Shared Resource (supported by NIH grants CA68485, DK20593, DK58404, DK59637 and EY08126). All electron microscopy reagents were purchased from Electron Microscopy Sciences. Cell cultures were grown to approximately 80% confluency, subjected to treatment conditions, and fixed with 2.5% glutaraldehyde in 0.1 M cacodylate for 1 hour at room temperature followed by 24 hours at 4°C. After fixation the cells were mechanically lifted from the tissue culture plates and pelleted then sequentially post-fixed with 1% tannic acid, 1% OsO4, and en bloc stained with 1% uranyl acetate. Samples were dehydrated in a graded ethanol series, infiltrated with Quetol 651 based Spurr's resin, and polymerized at 60°C for 48 hours. Ultrathin sections were prepared on a UC7 ultramicrotome (Leica) with a nominal thickness of 70 nm and collected onto 300 mesh nickel grids. Sections were stained with 2% uranyl acetate and lead citrate.

Samples were imaged using a Tecnai T12 operating at 100 kV equipped with an AMT NanoSprint CMOS camera using AMT imaging software. Analysis of the TEM data was performed in FIJI.

Mitochondrial Biogenesis Assay

The synthesis of mitochondria was assessed using a Mitochondrial Biogenesis In-Cell ELISA colorimetric kit (Abcam, catalog number ab110217), which measures the abundance of subunit I of complex IV (mtDNA-encoded) and the 70 kDa subunit of complex II (nuclear-encoded). For each experiment, the manufacturer's suggested procedures were followed. Briefly, 1.5×10^4 cells were seeded into the wells of a 96-well plate and subjected to treatments

24 hours later. Upon completion of the experimental conditions, cells were fixed for 20 minutes

with 4% paraformaldehyde followed by 3 washed with PBS and the plates were then blotted dry.

Subsequently, 100 µl of 0.5% acetic acid was added to each well for 5 minutes to block

endogenous alkaline phosphatase activity. The wells were then washed once again with PBS

followed by the addition of 1X permeabilization buffer for 30 minutes. Following

permeabilization, cells were incubated with 1X blocking solution for 2 hours followed by

overnight incubation with primary antibody at 4°C. After incubation with primary antibody, cells

were washed 3 times with PBS and then incubated with secondary antibody for 1 hour at room

temperature. Secondary antibody was then removed and the cells were washed 4 times with 1X

wash buffer. Expression of the 70 kDa subunit of complex II (SDH-A) and subunit I of complex

IV (COX-I) was then detected and quantified by wavelength detection at 405 nm and 600 nm,

respectively, using a GloMax plate reader.

UALCAN and GEPIA Databases

TCGA and CPTAC expression analyses were performed using the publicly accessible

UALCAN cancer database and GEPIA database [57, 58]. The effect of MUL1 gene expression

on patient survival in the kidney renal clear cell carcinoma dataset (KIRC) was determined by

comparing patient outcome in low vs high expression populations. MUL1 gene expression was

analyzed across individual cancer stages in the KIRC TCGA dataset and expression was

statistically compared to expression in normal adjacent tissue. Mul1 protein expression in

tumors was analyzed in the clear cell RCC dataset and statistically compared to expression in

normal adjacent tissue.

Author's Disclosures

WKR received an unrestricted grant from the VICC and support from the VICC-Incyte alliance

JCR is a founder, scientific advisory board member, and stockholder of Sitryx Therapeutics, a

scientific advisory board member and stockholder of Caribou Biosciences, a member of the

scientific advisory board of Nirogy Therapeutics, has consulted for Merck, Pfizer, and Mitobridge

within the past three years, and has received research support from Incyte Corp., Calithera

Biosciences, and Tempest Therapeutics. KEB has received funding to the institution from BMS-

IASLC-LCFA, consulted for Aravive, and served on the advisory board for Aveo, BMS, Exelexis,

and Seagen. The remaining co-authors have no conflicts to disclose.

Author's Contributions

Z.A.B. performed most of the experimental work, data analysis, and statistical analysis. Z.A.B.,

J.C.R., and W.K.R. participated in study conception and design, data interpretation, and drafting

manuscript. W.A.B. and E.K.A. helped with experimental work. E.S.K. was responsible for

obtaining electron microscopy images. M.W.W., R.A.H., and M.L. processed patient samples

and prepared primary cell cultures for experimental work. K.E.B. provided access to patient

samples. All co-authors assisted with reviewing and editing manuscript.

Acknowledgments

The authors give special thanks to their relevant funding sources. This work was supported by

AACR (WKR), R01 CA21797 (JCR) and institutional research funding from Bristol-Myers

Squibb, Merck, Pfizer, Calithera Biosciences, Peloton, and Incyte. ZAB is supported by the

Integrated Biological Systems Training in Oncology Ruth L. Kirschstein NRSA training grant

(5T32CA119925-12). VUMC Lipid Core which is supported by DK059637 and DK020593.

Electron microscopy was performed in part through the use of the Vanderbilt Cell Imaging

Shared Resource (supported by NIH grants CA68485, DK20593, DK58404, DK59637 and

EY08126). Immunohistochemical staining was performed in part through the use of the

Translational Pathology Shared Resource at VUMC. Authors acknowledge use of an active

BioRender license to generate model.

REFERENCES

1.   Siegel, R.L., et al., *Cancer Statistics, 2021.* CA: A Cancer Journal for Clinicians, 2021. **71**(1): p. 7-
33.
2.   Muglia, V.F. and A. Prando, *Renal cell carcinoma: histological classification and correlation with*
imaging findings. Radiologia brasileira, 2015. **48**(3): p. 166-174.
3.   Gerlinger, M., et al., *Genomic architecture and evolution of clear cell renal cell carcinomas*
defined by multiregion sequencing. Nature genetics, 2014. **46**(3): p. 225-233.
4.   Gnarra, J.R., et al., *Mutations of the VHL tumour suppressor gene in renal carcinoma.* Nature
Genetics, 1994. **7**(1): p. 85-90.
5.   Herman, J.G., et al., *Silencing of the VHL tumor-suppressor gene by DNA methylation in renal*
carcinoma. Proc Natl Acad Sci U S A, 1994. **91**(21): p. 9700-4.
6.   Nickerson, M.L., et al., *Improved identification of von Hippel-Lindau gene alterations in clear cell*
renal tumors. Clin Cancer Res, 2008. **14**(15): p. 4726-34.
7.   Sato, Y., et al., *Integrated molecular analysis of clear-cell renal cell carcinoma.* Nature Genetics,
2013. **45**(8): p. 860-867.
8.   Moore, L.E., et al., *Von Hippel-Lindau (VHL) inactivation in sporadic clear cell renal cancer:*
associations with germline VHL polymorphisms and etiologic risk factors. PLoS Genet, 2011.
7(10): p. e1002312.
9.   Ivan, M., et al., *HIFα targeted for VHL-mediated destruction by proline hydroxylation:*
implications for O2 sensing. Science, 2001. **292**(5516): p. 464-468.
10.   Semenza, G.L. and G.L. Wang, *A nuclear factor induced by hypoxia via de novo protein synthesis*
binds to the human erythropoietin gene enhancer at a site required for transcriptional activation.
Molecular and Cellular Biology, 1992. **12**(12): p. 5447-5454.
11.   Cancer Genome Atlas Research, N., *Comprehensive molecular characterization of clear cell renal*
cell carcinoma. Nature, 2013. **499**(7456): p. 43-49.
12.   Toma, M.I., et al., *Loss of heterozygosity and copy number abnormality in clear cell renal cell*
carcinoma discovered by high-density affymetrix 10K single nucleotide polymorphism mapping
array. Neoplasia (New York, N.Y.), 2008. **10**(7): p. 634-642.
13.   Bacigalupa, Z.A. and W.K. Rathmell, *Beyond glycolysis: Hypoxia signaling as a master regulator*
of alternative metabolic pathways and the implications in clear cell renal cell carcinoma. Cancer
letters, 2020. **489**: p. 19-28.
14.   Brown, W.A., W.K. Rathmell, and Z.A. Bacigalupa, *It's Clear as Day: HIF Signaling is Driving Force*
of the Clear Cell Morphology. KIDNEY, 2021. **19**(1): p. 17.
15.   Du, W., et al., *HIF drives lipid deposition and cancer in ccRCC via repression of fatty acid*
metabolism. Nature Communications, 2017. **8**(1): p. 1769.
16.   Syafruddin, S.E., et al., *A KLF6-driven transcriptional network links lipid homeostasis and tumour*
growth in renal carcinoma. Nature Communications, 2019. **10**(1): p. 1152.
17.   Qiu, B., et al., *HIF2α-Dependent Lipid Storage Promotes Endoplasmic Reticulum Homeostasis in*
Clear-Cell Renal Cell Carcinoma. Cancer Discov, 2015. **5**(6): p. 652-67.
18.   Choueiri, T.K., et al., *Inhibition of hypoxia-inducible factor-2α in renal cell carcinoma with*
belzutifan: a phase 1 trial and biomarker analysis. Nature Medicine, 2021. **27**(5): p. 802-805.

19. Courtney, K.D., et al., *Phase I Dose-Escalation Trial of PT2385, a First-in-Class Hypoxia-Inducible Factor-2α Antagonist in Patients With Previously Treated Advanced Clear Cell Renal Cell Carcinoma.* Journal of Clinical Oncology, 2018. **36**(9): p. 867-874.

20. Wallace, E.M., et al., *A Small-Molecule Antagonist of HIF2α Is Efficacious in Preclinical Models of Renal Cell Carcinoma.* Cancer Research, 2016. **76**(18): p. 5491-5500.

21. Xu, R., et al., *3-[(1S,2S,3R)-2,3-Difluoro-1-hydroxy-7-methylsulfonylindan-4-yl]oxy-5-fluorobenzonitrile (PT2977), a Hypoxia-Inducible Factor 2α (HIF-2α) Inhibitor for the Treatment of Clear Cell Renal Cell Carcinoma.* Journal of Medicinal Chemistry, 2019. **62**(15): p. 6876-6893.

22. Jonasch, E., et al., *Belzutifan for Renal Cell Carcinoma in von Hippel-Lindau Disease.* N Engl J Med, 2021. **385**(22): p. 2036-2046.

23. Zhang, D., et al., *JNK1 mediates degradation HIF-1α by a VHL-independent mechanism that involves the chaperones Hsp90/Hsp70.* Cancer research, 2010. **70**(2): p. 813-823.

24. Cilenti, L., et al., *Mitochondrial MUL1 E3 ubiquitin ligase regulates Hypoxia Inducible Factor (HIF-1α) and metabolic reprogramming by modulating the UBXN7 cofactor protein.* Scientific reports, 2020. **10**(1): p. 1-15.

25. Li, J., et al., *Mitochondrial outer-membrane E3 ligase MUL1 ubiquitinates ULK1 and regulates selenite-induced mitophagy.* Autophagy, 2015. **11**(8): p. 1216-29.

26. Prudent, J., et al., *MAPL SUMOylation of Drp1 stabilizes an ER/mitochondrial platform required for cell death.* Molecular cell, 2015. **59**(6): p. 941-955.

27. Yun, J., et al., *MUL1 acts in parallel to the PINK1/parkin pathway in regulating mitofusin and compensates for loss of PINK1/parkin.* eLife, 2014. **3**: p. e01958-e01958.

28. Courtney, K.D., et al., *HIF-2 Complex Dissociation, Target Inhibition, and Acquired Resistance with PT2385, a First-in-Class HIF-2 Inhibitor, in Patients with Clear Cell Renal Cell Carcinoma.* Clinical Cancer Research, 2020. **26**(4): p. 793-803.

29. Chen, R., et al., *Coordinate regulation of stress signaling and epigenetic events by Acss2 and HIF-2 in cancer cells.* PLoS One, 2017. **12**(12): p. e0190241.

30. Chen, R., et al., *The acetate/ACSS2 switch regulates HIF-2 stress signaling in the tumor cell microenvironment.* PloS one, 2015. **10**(2): p. e0116515.

31. Nagati, J.S., et al., *Mammalian acetate-dependent acetyl CoA synthetase 2 contains multiple protein destabilization and masking elements.* Journal of Biological Chemistry, 2021. **297**(3).

32. Dioum, E.M., et al., *Regulation of hypoxia-inducible factor 2α signaling by the stress-responsive deacetylase sirtuin 1.* Science, 2009. **324**(5932): p. 1289-1293.

33. Schug, Z.T., et al., *Acetyl-CoA synthetase 2 promotes acetate utilization and maintains cancer cell growth under metabolic stress.* Cancer cell, 2015. **27**(1): p. 57-71.

34. Miller, K.D., et al., *Targeting ACSS2 with a Transition-State Mimetic Inhibits Triple-Negative Breast Cancer Growth.* Cancer Research, 2021. **81**(5): p. 1252-1264.

35. Comerford, S.A., et al., *Acetate dependence of tumors.* Cell, 2014. **159**(7): p. 1591-1602.

36. Mashimo, T., et al., *Acetate is a bioenergetic substrate for human glioblastoma and brain metastases.* Cell, 2014. **159**(7): p. 1603-1614.

37. Ciraku, L., et al., *O-GlcNAc transferase regulates glioblastoma acetate metabolism via regulation of CDK5-dependent ACSS2 phosphorylation.* Oncogene, 2022: p. 1-15.

38. Bulusu, V., et al., *Acetate Recapturing by Nuclear Acetyl-CoA Synthetase 2 Prevents Loss of Histone Acetylation during Oxygen and Serum Limitation.* Cell Reports, 2017. **18**(3): p. 647-658.

39. Li, X., et al., *Nucleus-translocated ACSS2 promotes gene transcription for lysosomal biogenesis and autophagy.* Molecular Cell, 2017. **66**(5): p. 684-697. e9.

40. Covello, K.L., et al., *HIF-2alpha regulates Oct-4: effects of hypoxia on stem cell function, embryonic development, and tumor growth.* Genes Dev, 2006. **20**(5): p. 557-70.

41.	Hu, C.J., et al., *Differential regulation of the transcriptional activities of hypoxia-inducible factor 1 alpha (HIF-1alpha) and HIF-2alpha in stem cells.* Mol Cell Biol, 2006. **26**(9): p. 3514-26.

42.	Palikaras, K., E. Lionaki, and N. Tavernarakis, *Balancing mitochondrial biogenesis and mitophagy to maintain energy metabolism homeostasis.* Cell Death & Differentiation, 2015. **22**(9): p. 1399-1401.

43.	Shi, G. and G.A. McQuibban, *The mitochondrial rhomboid protease PARL is regulated by PDK2 to integrate mitochondrial quality control and metabolism.* Cell reports, 2017. **18**(6): p. 1458-1472.

44.	Shao, D., et al., *PGC-1 beta-regulated mitochondrial biogenesis and function in myotubes is mediated by NRF-1 and ERR alpha.* Mitochondrion, 2010. **10**(5): p. 516-27.

45.	Xu, M., et al., *An acetate switch regulates stress erythropoiesis.* Nature medicine, 2014. **20**(9): p. 1018-1026.

46.	Kondo, K., et al., *Inhibition of HIF2α is sufficient to suppress pVHL-defective tumor growth.* PLoS biology, 2003. **1**(3): p. e83.

47.	Kondo, K., et al., *Inhibition of HIF is necessary for tumor suppression by the von Hippel-Lindau protein.* Cancer Cell, 2002. **1**(3): p. 237-246.

48.	Shen, C., et al., *Genetic and functional studies implicate HIF1α as a 14q kidney cancer suppressor gene.* Cancer discovery, 2011. **1**(3): p. 222-235.

49.	Gordan, J.D., et al., *HIF-α effects on c-Myc distinguish two subtypes of sporadic VHL-deficient clear cell renal carcinoma.* Cancer cell, 2008. **14**(6): p. 435-446.

50.	Cho, H. and W.G. Kaelin. *Targeting HIF2 in clear cell renal cell carcinoma*. in *Cold Spring Harbor symposia on quantitative biology*. 2016. Cold Spring Harbor Laboratory Press.

51.	Yuan, Y., et al., *Mitochondrial E3 ubiquitin ligase 1 promotes autophagy flux to suppress the development of clear cell renal cell carcinomas.* Cancer science, 2019. **110**(11): p. 3533-3542.

52.	Wellen, K.E., et al., *ATP-Citrate Lyase Links Cellular Metabolism to Histone Acetylation.* Science, 2009. **324**(5930): p. 1076-1080.

53.	Zhao, S., et al., *ATP-Citrate Lyase Controls a Glucose-to-Acetate Metabolic Switch.* Cell Reports, 2016. **17**(4): p. 1037-1052.

54.	Folch, J., M. Lees, and G.H. Sloane Stanley, *A simple method for the isolation and purification of total lipids from animal tissues.* J biol Chem, 1957. **226**(1): p. 497-509.

55.	Morrison, W.R. and L.M. Smith, *Preparation of fatty acid methyl esters and dimethylacetals from lipids with boron fluoride–methanol.* Journal of lipid research, 1964. **5**(4): p. 600-608.

56.	Rudel, L.L., et al., *Dietary monounsaturated fatty acids promote aortic atherosclerosis in LDL receptor–null, human apoB100–overexpressing transgenic mice.* Arteriosclerosis, thrombosis, and vascular biology, 1998. **18**(11): p. 1818-1827.

57.	Chandrashekar, D.S., et al., *UALCAN: A Portal for Facilitating Tumor Subgroup Gene Expression and Survival Analyses.* Neoplasia, 2017. **19**(8): p. 649-658.

58.	Tang, Z., et al., *GEPIA: a web server for cancer and normal gene expression profiling and interactive analyses.* Nucleic Acids Research, 2017. **45**(W1): p. W98-W102.

 FIGURE LEGENDS

Figure 1. ACSS2 is essential for ccRCC growth and proliferation. **A.** Representative images of

crystal violet staining from a dose-response time-course treatment with DMSO or ACSS2

inhibitor in HKC cells. Quantification of three-independent replicates are provided in the graph

below. Statistical significance was determined using Bonferroni's multiple comparisons test. **B.**

Representative images of crystal violet staining from a dose-response time-course treatment

with DMSO or ACSS2 inhibitor in 786-O cells. Quantification of three-independent replicates are

provided in the graph below. Statistical significance was determined using Bonferroni's multiple

comparisons test. **C.** Box and whisker plots showing the absorbance values detected at OD450

nm of BrdU ELISA assays performed on HKC cells treated for 24 hours (left) or 48 hours (right)

with DMSO, 1 µM, 5 µM, or 10 µM ACSS2 inhibitor (n=4). Statistical significance was

determined using Bonferroni's multiple comparisons test (**, $P < 0.01$; ****, $P < 0.0001$; n.s., not

significant). **D.** Box and whisker plots showing the absorbance values detected at OD450 nm of

BrdU ELISA assays performed on 786-O cells treated for 24 hours (left) or 48 hours (right) with

DMSO, 1 µM, 5 µM, or 10 µM ACSS2 inhibitor (n=4). Statistical significance was determined

using Bonferroni's multiple comparisons test (**, $P < 0.01$; ***, $P < 0.001$; ****, $P < 0.0001$). **E.**

Representative images of 786-O cells transduced with pTRIPZ Control (left), pTRIPZ shACSS2

#1 (center), or pTRIPZ shACSS2 #2 (right) untreated (top) or treated with 2 µg/mL doxycycline

for 48 hours (bottom) stained with crystal violet. Bar graph displaying the quantification of crystal

violet staining from three-independent experiments. Statistical significance was determined

using a one-way ANOVA (**, $P < 0.005$). **F.** Box and whisker plots showing the absorbance

values detected at OD450 nm of BrdU ELISA assays performed on 786-O pTRIPZ Control or

786-O pTRIPZ shACSS2 cells treated with 2 µg/mL doxycycline for 24 hours (n=4). Two-tailed

paired t-test used to assess statistical significance (*, $P<0.05$). **G.** Line plot showing tumor

volume over time in mice inoculated with 786-O pTRIPZ Control (black, n=5) or 786-O pTRIPZ

shACSS2 (red, n=5) cells and fed a doxycycline rodent chow. Statistical significance was

determined using multiple two-tailed t-tests. **H.** Box and whisker plot showing individual data

points of tumor weight for pTRIPZ Control and pTRIPZ shACSS2 tumors. Statistical significance

was determined using an unpaired, two-tailed t-test (***, $P < 0.001$). **I.** Line plot showing tumor

volume over time in mice inoculated with 786-O cells treated with vehicle (black, n=6) or 15

mg/kg ACSS2 inhibitor (red, n=6). Statistical significance was determined using the Holm-Sidak

method of multiple t-tests. **J.** Box and whisker plot showing individual data points of tumor

weight for mice treated with vehicle (black, n=6) or 15 mg/kg ACSS2 inhibitor (red, n=6).

Statistical significance was determined using an unpaired, two-tailed t-test (*, $P < 0.05$). **K.**

Representative images taken at 10X magnification of sections from Control (top) and shACSS2

(bottom) tumors stained with CD31 (left) and Periodic Acid Schiff (PAS, right). **L.** Representative

images taken at 10X magnification of sections from Vehicle (top) and 15 mg/kg ACSS2 inhibitor-

treated (bottom) tumors stained with CD31 (left) and Periodic Acid Schiff (PAS, right).

Figure 2. HIF-2α gene and protein expression is regulated by ACSS2 activity. **A.** Heatmap

showing differential gene expression expressed as fold change (downregulated by $-1.5 \geq$ and

upregulated by $1.5 \leq$, p-Adj ≤ 0.05) between Control (blue) and shACSS2 (red) groups. Gene

expression clustering on the Y-axis depicts genes that are upregulated (purple) and

downregulated (green) in the Control group. **B.** Heatmap showing differential gene expression

following pathway enrichment (Wikipathways) as fold change (downregulated by $-1.5 \geq$ and

upregulated by $1.5 \leq$, p-Adj ≤ 0.05) between Control (blue) and shACSS2 (red) groups. Gene

expression clustering on the Y-axis depicts genes that are upregulated (purple) and

downregulated (green) in the Control group. Bar graph (right) showing transcript counts from

HIF-2α RCC Signaling pathway comparing Control (black) and shACSS2 (red) groups.

Statistical significance determined using unpaired t-tests without correction for multiple

comparisons (**, P < 0.01; ***, P < 0.001; ###, P < 0.0001). **C.** Western blot analysis of HIF-2α

and ACSS2 expression following 24-hour doxycycline induction of 786-O pTRIPZ Control and

pTRIPZ shACSS2 cells. **D.** Western blot analysis assessing expression of HIF-2α, Epo, and

VEGFR2 in 786-O cells treated with DMSO, 1 μM, 5 μM, or 10 μM ACSS2 inhibitor for 24 hours.

E. Western blot analysis showing expression of ACSS2, V5-tag, HIF-2α, VEGFR2, Epo, and

EGLN3 in 786-O cells transduced pLX304 empty vector or pLX304 V5-ACSS2 overexpression

vector. **F.** Box and whisker plot showing the absorbance values detected at OD450 nm of BrdU

ELISA assays performed on 786-O cells transduced pLX304 empty vector or pLX304 V5-

ACSS2 overexpression vector. Statistical significance was determined using an unpaired, two-

tailed t-test (***, P < 0.001). **G.** Representative images taken at day 21 of an anchorage-

independent growth assay performed using 786-O control or V5-ACSS2 cells transduced

express shControl or shHIF2A. Bar graph showing the number of colonies formed for each

condition from three-independent experiments. Statistical significance determined using two-

way ANOVA and Tukey's multiple comparisons test (*, P-value < 0.05; ***, P-value < 0005).

Figure 3. ACSS2 activity mediates HIF-2α stability. **A.** Western blot depicting HIF-2α

expression in 786-O cells in the absence or presence of a 1-hour incubation with 10 μM MG132

prior to being treated with DMSO or 5 μM ACSS2 inhibitor for 24 hours. Bar chart (right)

showing densitometry analysis performed using ImageJ on western blot images from three-

independent experiments. Statistical significance determined using multiple unpaired, two-tailed

t-tests (*, P < 0.05; ***, P < 0.0005; ##, P < 0.01; * = t-test performed vs No pre-treatment

DMSO; # = t-test performed vs No pre-treatment ACSS2i). **B.** Kaplan-Meier plot generated from

the KIRC TCGA dataset assessing patient outcome as it relates to MUL1 expression (MUL1[HIGH]

= black, MUL1[LOW] = red). Statistical significance determined automatically via GEPIA database.

C. Representative images at day 16 of anchorage-independent growth assays performed on

786-O cells transduced to express shControl or shMUL1 and treated with either DMSO, 5 µM

ACSS2 inhibitor, or 10 µM PT-2385. **D.** Western blot images showing expression of the

mitophagy marker LC3 A/B, K48 polyubiquitination, MUL1, and p62 in 786-O cells treated with

DMSO or 5 µM ACSS2 inhibitor for 48 hours. **E.** Western blot analysis showing expression

changes to ACSS2, HIF-2α, and MUL1 in response to induction of shACSS2. **F.** Western blot

images assessing protein interactions from immunoprecipitations of HIF-2α or MUL1 performed

on 786-O cells treated with DMSO or 5 µM ACSS2 inhibitor for 24 hours.

Figure 4. ACSS2 activity supports glucose and cholesterol metabolism. **A.** Heatmap (left)

showing differential gene expression expressed as fold change (downregulated by –1.1 ≥ and

upregulated by 1.1 ≤, p-Adj ≤ 0.05) between 24-hour treatment of 786-O cells with DMSO (blue)

and 5 µM ACSS2 inhibitor (red) groups. Gene expression clustering on the Y-axis depicts genes

that are upregulated (purple) and downregulated (green) in the DMSO group. Heatmap (right)

showing differential gene expression following pathway enrichment (Wikipathways) as fold

change (downregulated by –1.1 ≥ and upregulated by 1.1 ≤, p-Adj ≤ 0.05) between 24-hour

treatment of DMSO (blue) and 5 µM ACSS2 inhibitor (red) groups. Gene expression clustering

on the Y-axis depicts genes that are upregulated (purple) and downregulated (green) in the

DMSO group. **B.** Bar graph showing total cholesterol levels (µg) in 786-O cells treated with

DMSO (black; n=3) or 5 µM ACSS2i (red; n=3) for 24 hours. Statistical significance determined

using an unpaired, two-tailed Student's t-test. **C.** Bar graphs showing relative mRNA abundance

of SLC2A1 (left) and HKII (right) in 786-O cells following 24-hour treatment with DMSO (black)

or 5 µM ACSS2 inhibitor (red). Statistical significance determined using an unpaired, two-tailed

Student's t-test. **D.** Western blot analysis showing expression of HIF-2α, GLUT1, HKII, SREBP-

1, and SREBP-2 in 786-O cells treated with DMSO, 5 µM, or 10 µM ACSS2 inhibitor for 24

hours. **E.** Western blot analysis of ACSS2, HIF-2α, GLUT1, and HKII protein expression in 786-

O cells transduced to express pLX304 empty vector or pLX304 V5-ACSS2 overexpression

vector.

Figure 5. Targeting ACSS2 elicits cancer cell-specific mitochondrial defects. **A.** Representative

images captured via electron microscope of HKC cells treated with DMSO for 72 hours or with 5

μM ACSS2 inhibitor for 24, 48, and 72 hours. Black square inset indicates area of advanced

magnification. **B.** Representative images captured via electron microscope of 786-O cells

treated with DMSO for 72 hours or with 5 μM ACSS2 inhibitor for 24, 48, and 72 hours. Black

square inset indicates area of advanced magnification. **C.** Representative images captured via

electron microscope of 786-O pTRIPZ Control and shACSS2 cells induced with 2 μg/mL

doxycycline for 48 hours. Black square inset indicates area of advanced magnification. **D.**

Representative images captured via electron microscope of 786-O cells transduced to

overexpress ACSS2 (left) followed by transduction with shHIF2A. Black square inset indicates

area of advanced magnification. **E.** Representative images captured via electron microscope of

786-O cells treated with 10 μM PT-2385 for 72 hours. Black square inset indicates area of

advanced magnification.

Figure 6. ACSS2 inhibition selectively impedes cancer cell growth and HIF-2α expression in

ccRCC patient samples. **A.** Representative images of crystal violet staining from a dose-

response time-course treatment with DMSO or ACSS2 inhibitor in epithelial cells isolated from

ccRCC patient's normal adjacent tissue. Quantification of three-independent replicates are

provided in the graph below. Statistical significance was determined using Bonferroni's multiple

comparisons test (P values provided in table below). **B.** Representative images of crystal violet

staining from a dose-response time-course treatment with DMSO or ACSS2 inhibitor in

epithelial cells isolated from ccRCC patient tumor sections. Quantification of three-independent

replicates are provided in the graph below. Statistical significance was determined using

Bonferroni's multiple comparisons test (P values provided in table below). **C.** Representative

western blot images showing expression of HIF-2α and HKII in cells isolated from ccRCC

patient normal adjacent tissue treated with DMSO, 1μM, 5 μM, or 10 μM ACSS2 inhibitor for 24

hours. **D.** Representative western blot images showing expression of HIF-2α and HKII in

epithelial cells isolated from ccRCC patient tumor sections treated with DMSO, 1μM, 5 μM, or

10 μM ACSS2 inhibitor for 24 hours. **E.** Bar graph showing crystal violet quantification of 769-P

cells (three-independent experiments) in the presence or absence of 10 μM PT-2385 treated

with DMSO, 1μM, 5 μM, or 10 μM ACSS2 inhibitor for 24, 48, and 72 hours. Statistical

significance determined using a two-way ANOVA and Tukey's multiple comparisons test.

Representative images of crystal violet staining from a 72-hour dose-response treatment with

DMSO or ACSS2 inhibitor in 769-P cells.

Supplemental Figure 1. A. Representative images of crystal violet staining from a dose-

response time-course treatment with DMSO or ACSS2 inhibitor in A498 cells. Quantification of

three-independent replicates are provided in the graph below. Statistical significance was

determined using Bonferroni's multiple comparisons test. **B.** Box and whisker plots showing the

absorbance values detected at OD450 nm of BrdU ELISA assays performed on A498 cells

treated for 24 hours (top) or 48 hours (bottom) with DMSO, 1 μM, 5 μM, or 10 μM ACSS2

inhibitor (n=4). Statistical significance was determined using Bonferroni's multiple comparisons

test (*, $P < 0.05$; ***, $P < 0.001$; ****, $P < 0.0001$). **C.** Representative images of 786-O tumor

spheres at day 7 of growth in ultra-low attachment plates treated with either DMSO or 5 μM

ACSS2i. Quantification of average sphere area from three-independent replicates are provided

in the bar graph (right). Statistical significance was determined using an unpaired, two-tailed t-

test (***, $P < 0.001$). **D.** Bar graph showing the body weight of shControl + DOX and shACSS2 +

DOX mice at the endpoint of study. **E.** Bar graph showing the body weight of Vehicle- and 15

mg/kg ACSS2i-treated mice at the endpoint of study. **F.** Bar graphs showing quantification of the

percentage of total area staining positive for CD31 (left) or Periodic Acid Schiff (right) in tumor

sections from shControl + DOX or shACSS2 + DOX mice. Statistical significance determined

using an unpaired, two-tailed t-test with Welch's correction (**, P-value < 0.005; ***, P-value <

0.001). **G.** Bar graphs showing quantification of the percentage of total area staining positive for

CD31 (left) or Periodic Acid Schiff (right) in tumor sections from mice treated with Vehicle or 15

mg/kg ACSS2i. Statistical significance determined using an unpaired, two-tailed t-test with

Welch's correction (*, P-value < 0.05).

Supplemental Figure 2. A. Bar graph depicting relative abundance of *Epas1* mRNA expression

determined by qRT-PCR in 786-O cells treated with DMSO or 5 µM ACSS2 inhibitor for 24

hours. Statistical significance determined by unpaired, two-tailed Student's t-test (***, P <

0.001). **B.** Bar graph depicting relative abundance of *Epas1* mRNA expression determined by

qRT-PCR in 786-O cells transduced to express shControl or shHIF2A. Statistical significance

determined by unpaired, two-tailed Student's t-test (***, P < 0.001). **C.** Western blot analysis of

HIF-2α and Actin in the HKC cells treated with DMSO, 1µM, 5 µM, or 10 µM ACSS2i for 24 and

48 hours.**D.** Western blot analysis of HIF-2α, ACSS2, and Actin in the 786-O cells transduced

with empty vector (top) or stably overexpressing V5-ACSS2 (bottom) previously developed in

Figure 2 transduced with shControl or shHIF2A targeting constructs.

Supplemental Figure 3. A. Western blot from replicate experiments depicting HIF-2α

expression in 786-O cells in the absence or presence of a 1-hour incubation with 10 µM MG132

prior to being treated with DMSO or 5 µM ACSS2 inhibitor for 24 hours. **B.** Box and whisker plot

generated using UALCAN database to access KIRC TCGA dataset and assess MUL1 gene

expression spanning the individual stages of disease. Statistical significance determined

automatically via UALCAN database. **C.** Box and whisker plot generated using UALCAN

database to access KIRC CPTAC dataset and assess MUL1 protein expression in normal

kidney and ccRCC samples. Statistical significance determined automatically via UALCAN

database. **D.** Bar graph showing the number of colonies formed for each condition from three-

independent experiments. Statistical significance determined using two-way ANOVA and

Tukey's multiple comparisons test (***, P-value < 0.0005; ****, P-value < 0001). **E.** Western blot

analysis of HIF-2α, ACSS2, MUL1, and Actin in the HKC cells treated with DMSO, 5 µM, or 10

µM ACSS2i for 24 hours.

Supplemental Figure 4. A. Bar graph showing transcript counts for genes involved in the

cholesterol biosynthesis pathway extracted from the NanoString dataset developed from the

experiment in Figure 4A where 786-O cells were treated with DMSO (black) or 5 µM ACSS2i

(red) for 24 hours. Statistical significance determined by performing multiple unpaired, two-tailed

t-tests (*, P-value < 0.05; ***, P-value < 0.0005). **B.** Bar graph showing average glucose

concentrations in growth media in 786-O cells treated with DMSO (black) or 5 µM ACSS2i (red)

at 0, 24, and 48 hour time points. Statistical significance determined using unpaired, two-tailed t-

test (***, P-value < 0.005). **C.** Bar graph showing relative concentrations of acetate in growth

media of 786-O cells treated with DMSO, 1 µM, 5 µM, or 10 µM ACSS2 inhibitor for 24 hours.

Data are presented as Log_2 values of fold change. Statistical significance determined using an

ordinary one-way ANOVA and Bonferroni's multiple comparisons test (**, P-value < 0.01; ***, P-

value < 0.001). **D.** Bar graph showing relative concentrations of acetate in growth media of 786-

O cells transduced to express V5-ACSS2 overexpression vector, or the DOX-inducible

shControl or shACSS2 constructs. Data are presented as Log_2 values of fold change. Statistical

significance determined using an ordinary one-way ANOVA and Bonferroni's multiple

comparisons test (*, P-value < 0.05; **, P-value < 0.01; ****, P-value < 0.0001).

Supplemental Figure 5. A. Western blot analysis of PGC1β, Opa-1, Mitofusin 1, Mitofusin 2,

and Actin using the same membrane from Supplemental Figure 3E. Experiment was performed

in HKC cells treated with DMSO, 5 µM or 10 µM ACSS2 inhibitor for 24 hours. **B.** Western blot

analysis of PGC1α, PGC1β, Opa-1, Mitofusin 1, Mitofusin 2, Drp1, phospho-Drp1 (S616), and

Actin using same membrane as Figure 3D. Experiment performed in 786-O cells treated with

DMSO or 5 µM ACSS2 inhibitor for 24 hours. **C.** Bar graph with individual data points showing

quantification of mitochondrial biogenesis in HKC cells treated with DMSO, 5 µM or 10 µM

ACSS2 inhibitor for 24 hours. Data is shown as the ratio of expression of COX-I to SDH-A

quantified by wavelength detection at 405 nm (SDH-A) and 600 nm (COX-I). Statistical

significance assessed using an ordinary one-way ANOVA and Bonferroni's multiple

comparisons test. **D.** Bar graph with individual data points showing quantification of

mitochondrial biogenesis in 786-O cells treated with DMSO, 5 µM or 10 µM ACSS2 inhibitor for

24 hours. Data is shown as the ratio of expression of COX-I to SDH-A quantified by wavelength

detection at 405 nm (SDH-A) and 600 nm (COX-I). Statistical significance determined using an

ordinary one-way ANOVA and Bonferroni's multiple comparisons test (*, P-value < 0.05; ***, P-

value < 0.001). **E.** Bar graph with individual data points showing quantification of mitochondrial

biogenesis in 786-O pTRIPZ shControl and shACSS2 cells treated with doxycycline for 24

hours. Data is shown as the ratio of expression of COX-I to SDH-A quantified by wavelength

detection at 405 nm (SDH-A) and 600 nm (COX-I). Statistical significance determined using an

unpaired, two-tailed t-test (****, P-value < 0.0001). **F.** Western blot analysis of PGC1α, PGC1β,

Opa-1, Mitofusin 1, Mitofusin 2, and Actin probing the same membrane as used in Figure 3E.

Experiment performed in 786-O pTRIPZ shControl and shACSS2 cells treated with doxycycline

for 24 hours.

Supplemental Figure 6. A. Representative images of crystal violet staining from a dose-

response time-course treatment with DMSO or ACSS2 inhibitor in epithelial cells isolated from

ccRCC patient's normal adjacent tissue. **B.** Representative images of crystal violet staining

from a dose-response time-course treatment with DMSO or ACSS2 inhibitor in epithelial cells

isolated from ccRCC patient tumor sections. **C.** Representative images of crystal violet staining

from a dose-response time-course treatment with DMSO or ACSS2 inhibitor in epithelial cells

isolated from ccRCC patient's normal adjacent tissue. **D.** Representative images of crystal violet

staining from a dose-response time-course treatment with DMSO or ACSS2 inhibitor in

epithelial cells isolated from ccRCC patient tumor sections. **E.** Bar graph showing average fold

change values for gene expression of *HK2* and *EPAS1* extracted from Nanostring experiment

performed in ccRCC patient tumor samples treated with DMSO (n=3) or 5 μM ACSS2i (n=3) for

48 hours. Statistical significance determined by performing unpaired, two-tailed t-tests (***, P-

value < 0.001). **F.** Bar graph showing average fold change values for gene expression of genes

involved in the cholesterol biosynthesis pathway extracted from Nanostring experiment

performed in ccRCC patient tumor samples treated with DMSO (n=3) or 5 μM ACSS2i (n=3) for

48 hours. Statistical significance determined by performing multiple unpaired, two-tailed t-tests

(**, P-value < 0.01; ***, P-value < 0.001; ****, P-value < 0.0001). **G.** Model of mechanism.

Figure 1. ACSS2 is essential for ccRCC growth and proliferation

Supplemental Figure 1

Figure 2. HIF-2α gene and protein expression is regulated by ACSS2 activity.

Supplemental Figure 2

Figure 3. ACSS2 activity mediates HIF-2α stability.

A

786-O

MG132 (1h):	-	-	-	-	-	+	+
5 µM ACSS2i (h):	0	2	6	12	24	0	24

HIF-2α

Actin

786-O

MG132 (1h):	-	-	-	+	+	+
DMSO (24h):	+	-	-	+	-	-
ACSS2i (24 h):	-	+	++	-	+	++

HIF-2α

Actin

B

MUL1 – KIRC Individual Cancer Stages

Comparison	Statistical Significance
Normal-vs-Stage 1	5.4400928E-15
Normal-vs-Stage 2	1.9221002E-10
Normal-vs-Stage 3	<1E-12
Normal-vs-Stage 4	4.4408920E-16

C

Protein expression of MUL1 in ccRCC

Comparison	Statistical Significance
Normal-vs-Primary	3.990813E-04

D

# of colonies

- ■ DMSO
- ■ 5 µM ACSS2i
- ■ 10 µM PT-2385

shControl   shMUL1

****
***

E

HKC

	DMSO	ACSS2i (µM) 5	10

HIF-2α

ACSS2

Mul1

Actin

Supplemental Figure 3

Figure 4. ACSS2 activity supports glucose and cholesterol metabolism.

A

Cholesterol Biosynthesis

Transcript Counts (y-axis: 0, 1000, 2000, 3000, 4000, 5000, 6000, 7000)

*** (SQLE)
* (HMGCS1)
* (HMGCR)
* (MVK)
* (DHCR7)
* (DHCR24)

x-axis: SREBF1, SREBF2, HMGCS1, HMGCR, MVK, FDPS, SQLE, DHCR7, DHCR24

■ DMSO ■ ACSS2i

B

[Glucose] mg/dL (y-axis: 0, 50, 100, 150, 200, 250)

n.s. (24h)
** (48h)

x-axis: 0h, 24h, 48h

■ DMSO ■ ACSS2i

C

Acetate

Relative [Acetate] in Media Log$_2$(Fold Change) (y-axis: -0.1, 0.0, 0.1, 0.2, 0.3)

** (1 μM ACSS2i)
*** (5 μM ACSS2i)
*** (10 μM ACSS2i)

x-axis: DMSO, 1 μM ACSS2i, 5 μM ACSS2i, 10 μM ACSS2i

786-O

D

Acetate

Relative [Acetate] in Media Log$_2$(Fold Change) (y-axis: -3, -2, -1, 0, 1)

* (Control)
** (shACSS2)
* (Control + DOX)
**** (ACSS2 O/E)

x-axis: Growth Media, Control, shACSS2, Control + DOX, shACSS2 + DOX, ACSS2 O/E

786-O

Supplemental Figure 4

Figure 5. Targeting ACSS2 elicits cancer cell-specific mitochondrial defects.

Supplemental Figure 5

Figure 6. ACSS2 inhibition selectively impedes cancer cell growth and HIF-2α expression in ccRCC patient samples.

Supplemental Figure 6