Title: Multisite imaging of neural activity using a genetically encoded calcium

sensor in the honey bee *Apis mellifera*

Short title: Neural recordings using transgenic honey bees

Authors: Julie Carcaud[1*], Marianne Otte[2], Bernd Grünewald[3], Albrecht Haase[4,5,6], Jean-Christophe

Sandoz[1,6], Martin Beye[2,6]

*Corresponding author: Julie Carcaud, Evolution, Genomes, Behavior and Ecology, Université Paris-

Saclay, CNRS, IRD, Gif-sur-Yvette, France. Email: julie.carcaud@egce.cnrs-gif.fr. Phone: +33169823752

Author's addresses:

[1] Evolution, Genomes, Behavior and Ecology, Université Paris-Saclay, CNRS, IRD, Gif-sur-Yvette, France

[2] Evolutionnary Genetics, Heinrich Heine University Düsseldorf, Düsseldorf, Germany

[3]Institut für Bienenkunde, Polytechnische Gesellschaft, FB Biowissenschaften, Goethe-University,

Frankfurt am Main, Germany

[4] Center for Mind/Brain Sciences, University of Trento, Rovereto, Italy

[5] Department of Physics, University of Trento, Trento, Italy

[6]These authors jointly supervised this work

ABSTRACT

Sociality is classified as one of the major transitions in evolution, and the most advanced level of sociality is found in eusocial insects such as the honey bees. This sophisticated social behavior seems to be associated with higher cognitive abilities in these insects, however the study of their neural bases has been limited by the lack of tools that allow measuring neuronal activity simultaneously in different brain regions of the honey bee brain. Here, we developed the first pan-neuronal genetic driver in a Hymenopteran model organism, the honey bee, and expressed the calcium indicator GCaMP6f under the control of the honey bee *synapsin* promoter. We show that GCaMP6f is widely expressed in the honey bee brain, allowing to record neural activity from multiple brain regions simultaneously, including from poorly investigated brain regions such as the mushroom body calyces or the lateral horn. Our recordings show that while odorant quality (chemical structure) and quantity are faithfully encoded in the honeybee antennal lobe, odor coding in the lateral horn departs from this simple physico-chemical coding, in line with the role of this structure in coding the biological value of odorants. These brain recordings represent the first use of a neurogenetic tool for recording neural activity in a eusocial insect.

Keywords: Insect, brain, olfaction, transgenesis, calcium imaging, GCaMP

INTRODUCTION

Sociality is classified as one of the major transitions in evolution, and animals often form social groups because the benefits of grouping (either direct or indirect) outweigh the costs of breeding independently (1, 2). The most advanced level of sociality is found in eusocial insect societies (3, 4), and the insect order Hymenoptera (including ants, bees and wasps) presents the largest number of eusocial species. Among them, honey bees are a classical model for the study of eusocial behavior, as they live in colonies composed of up to 60,000 individuals, consisting of three adult castes (queen, worker, and male). Within the worker caste, honey bees show a clear division of labor with a specialization of roles (5). The success of honey bee colonies lies in the capacity of all members of the society to behave in a well-organized and context-dependent manner, a social behavior mediated in part by olfactory cues such as pheromones used for communication among the colony (6).

Previous research found that this sophisticated social behavior is associated with higher cognitive abilities in these insects, and in this context, honey bees is a mainstream model for studying higher-order insect cognition such as navigation (7), rule learning (8), social learning (9), dance communication(10) but also olfactory perception and learning (11-13). However, whether the evolutionary rise of these sophisticated social behaviors is associated with the formation of specific neuronal pathways and structures has not been sufficiently resolved (14, 15). Progress in understanding these sophisticated behaviors and how social cues are processed and integrated with higher order cognitive abilities have in part been limited due to the lack of tools allowing to measure neuronal activity simultaneously in different regions of the honey bee brain.

For now, honey bee neuroscience remains limited to the use of conventional neuroanatomy (16), pharmacology (17), electrophysiology (18), and imaging tools (13, 19). However, the development of neurogenetic tools during the 20th century has provided unprecedented progress in our understanding of the neural basis of behavior in other model species. Among others, this approach allowed to decipher the brain circuits underlying aggressive behavior (20), courtship (21), or memory

(22) in a limited set of model species, both vertebrates such as mouse or zebrafish and invertebrates.

In insects, the fruit fly *Drosophila melanogaster* has been for many years the leading model for

investigating the neural basis of behavior, from gene expression to neural circuits (23-25). Recently,

neurogenetic approaches have been developed in other Diptera such as mosquitoes (26, 27) and in

some Lepidoptera (28, 29) to understand specific behaviors such as human-host seeking (30, 31) or

insect-plant interactions (32), mainly focusing on their olfactory capabilities. However, genetic

methods are particularly difficult to apply in eusocial insects, since genetic transformation rates are

low, endogenous promotors for neuronal expressions are unknown and the genetically manipulated,

reproductive individuals (the queens) have to be maintained in larger colonies with workers in

containments (33).

Despite the difficulty, the recent use of the genome editing tool CRISPR/Cas9 in the honey bee

allowed to knock out specific genes, such as the sex-determining *dsx* gene (34), the olfactory co-

receptor gene *orco* (35), the gustatory receptor *AmGr3* gene (36) or the *Amyellow-y* gene (37). Even if

these studies evaluated the effect of specific mutations at the neuronal (35) or the behavioral levels

(36), versatile genetic tools allowing to investigate the neural basis of a wide range of higher-order

social behaviors and learning in honey bees are still missing. The advent of genetically-encoded neural

activity sensors, in particular calcium sensors, has represented a major breakthrough in *Drosophila* or

mouse research (38-40), but such a critical tool for circuit dissection is still lacking in Hymenoptera.

To this aim, we developed the first pan-neuronal genetic driver in a Hymenopteran model

organism, the honey bee, and expressed the calcium indicator GCaMP6f under the control of the honey

bee *synapsin* promoter. We characterized its expression pattern in the honey bee brain and evaluated

its potential as a functional tool by recording neural activity upon olfactory stimulation. We show that

GCaMP6f expression allows to record olfactory responses from multiple brain regions simultaneously,

after simply opening the brain capsule. By using a controlled panel of well-characterized odorants, we

show that the recorded signals reveal robust odor coding rules. This new pan-neuronal genetic driver

also permits to record neural activity from poorly recorded regions of the brain such as the lateral horn

and the mushroom body calyces, and allowed to find that olfactory chemical features are less

represented in the lateral horn than in the antennal lobe. This study opens extraordinary new

possibilities for neuroethological research in the honey bee to study the neural basis of advanced social

behaviors and cognitive skills.

RESULTS

This work aimed at expressing the calcium-sensitive protein GCaMP6f in a pan-neuronal manner in the

honey bee brain to record neural activity in this social insect.

Generation of transgenic honey bees

In order to generate a bee with a possibly pan-neuronally expressed calcium sensor, we introduced a

synapsin (syn) promoter GCaMP6f expression cassette (*syn-GCaMP6f*) into the genome using the

piggyBac (PB) transposon system (33). To generate this *syn-GCaMP6f* expression cassette we cloned

the 1 kb promoter region together with the entire 5' untranslated region (5' UTR) of the *synapsin* gene

from the honey bee, which we then fused with the coding sequence of the GCaMP6f sensor protein

(Figure 1 A). We obtained a transgenic *syn-GCaMP6f* queen using previously published procedures and

a hyperactive transposase (33, 41). We instrumentally inseminated the queen and reared from the

offspring 2^{nd} generation queens. Only 1% of those queens were carrying the transgene, which we

identified from transgene amplifications. The 2^{nd} generation *syn-GCaMP6f* queens produced 50% *syn-*

GCaMP6f worker offspring bees, which we used in the neuroanatomy and imaging experiments.

Pattern expression of the GCaMP6f in the brain

We first studied the pattern expression of GCaMP6f in the brain using immunostaining against

GFP (Figure 1B-C), the green fluorescent protein which the GCaMP6f sensor contains. We found clear

and widespread staining throughout the brain, with strong staining in the antennal lobes (ALs), the

optic lobes (OLs), and the mushroom bodies (MBs) suggesting that the GCaMP6f protein is ubiquitously expressed in all major brain structures. Within neurons, we found evidence of staining in somata (Figure 1B left, white arrows) as well as neuronal processes (see for instance neural tracts around the α-lobe; Figure 1B left, white arrows). At the level of the AL (Figure 1C left), lateral and medial clusters of projection neurons and local interneurons somata show strong GCaMP6f immunostaining. In addition, glomeruli seem homogeneously stained: staining in the cortex suggests expression in olfactory sensory neurons (OSNs) terminal projections, whereas staining in the glomerulus core suggests expression in projection neurons (PNs) dendrites and possibly local neurons. Finally, the clearest expression of GCaMP6f was found in somata at the level of the calyces of the MB (Figure 1C right). In these structures, different neuron types are stained, with somata of class I Kenyon Cells in the cup-shaped calyces and class II Kenyon Cell somata lying outside the calyces (see white arrow in Figure 1C right) (42, 43).

We next asked whether the cloned synapsin promoter used for GCaMP6f expression drives the same expression pattern as the one of the *synapsin* gene. To answer this question, we performed a complementary immunostaining targeting the synapsin protein (Figure 1B right, anti-SYNORF1 in red). We found co-localized staining of GCaMP6f and synapsin within AL glomeruli both in the cortex and the core (Figure 1C), demonstrating co-expression within AL neurons. This experiment also shows differences between GCaMP6f and synapsin stainings. The synapsin immunostaining allows to clearly visualize presynaptic zones, for instance in AL glomeruli or microglomeruli in the MB calyces, but is not present in the somata unlike GCaMP6f. This result indicates that even if GCaMP6f is expressed in cells expressing synapsin (in theory all neurons), these 2 proteins appear to be partially differently localized within neurons.

We conclude that GCaMP6f was widely expressed in the honey bee brain, one of the prerequisites for whole-brain functional recordings of neural activity.

Calcium imaging

Neural recordings in the whole brain

GCaMP6f expression in neurons was then used to record neural activity in the honey bee brain (Figure

2). As a proof of concept for the use of these bees in neuroethological research, we decided to focus

here on olfactory information processing, as olfaction is the best understood sensory modality in honey

bees. Right after placing the bee in a recording chamber and opening its head capsule, we first

presented a few standard odorants while recording the whole brain surface accessible under the

microscope objective. Odorant presentations triggered a clear GCaMP fluorescence change from

several brain regions, including the primary olfactory centers (the antennal lobes, ALs) and both higher-

order centers (the lateral horns, LHs and the mushroom bodies, MBs), known for their role in olfactory

processing (Figure 2A). The use of GCaMP-expressing bees allowed us to record neural activity

simultaneously in all these structures (Figure 2B), while the presentation of an unscented air control

did not induce such activity. Response amplitudes were stronger in the AL (Figure 2C left) than in the

LH (Figure 2C middle) or in the MB (Figure 2C right). In all regions, the recorded calcium signals showed

a biphasic response, with a fluorescence increase upon odor presentation followed by a long

undershoot. Such biphasic signals are reminiscent of calcium signals previously recorded using bath-

applied calcium-sensitive dyes (44, 45). This experiment represents the first recording of neural activity

using neurogenetic tools outside of Diptera.

Neural recordings in the antennal lobe (AL)

We then asked if the signals recorded using this genetically-encoded calcium sensor represent

neural signals that are meaningful in terms of sensory coding, focusing first on the primary olfactory

center, the antennal lobe (AL), since olfactory coding was best studied in this structure. To this aim,

we presented 16 aliphatic odorants used in previous imaging studies either after bath-application of a

cell-permeant calcium-sensitive dye (Calcium-green 2-AM, OSN recordings) (45) or after insertion of a

migrating calcium-sensitive dye within a neural tract (Fura-2 dextran, PN recordings) (19). The

response patterns to the 16 aliphatic odors differed systematically according to their functional group

(primary or secondary alcohols, aldehydes, or ketones) or their carbon chain length (from 6 to 9

carbons). As observed with other calcium reporters, the presentation of each odorant induced a

biphasic signal in a different set of AL glomeruli while presentation of the air control did not induce

such response (Figure 3A-B). The averaged AL response amplitudes were significantly different from

the air control for all 16 odors (Figure 3C, RM-ANOVA, odor effect $F_{16,160}$ = 12.49, p = 8e-21, post hoc

Dunnett tests p < 0.0097). Signal amplitudes differed according to the odor (odor effect, p = 8e-21),

and were related to the quantity of molecules in headspace, since odor-evoked responses were highly

correlated with the vapor pressure of the odorants (Figure 3D, R^2 = 0.84, Fisher test $F_{1,14}$ = 73.72, p =

6e-7). Such a correlation was also found in previous studies recording at AL input (OSNs)(45) or output

(PNs)(19). This strong dependence explains why odor-evoked intensities recorded using genetically

encoded GCaMP6f in this study are highly correlated with odor-evoked intensities recorded in OSNs

(Sup. Figure 1A, R^2 = 0.84, $F_{1,14}$ = 74.23, p = 6e-7) or in PNs (Sup. Figure 1B, R^2 = 0.78, $F_{1,14}$ = 49.73, p =

6e-6).

We then evaluated odor coding in the GCaMP bees by comparing similarity relationships

among odorants, by calculating Euclidian distances between response maps obtained for the different

odor pairs. First, we confirmed that within each bee, each odorant evokes a specific activity pattern,

as distances were lower for different presentations of the same odorant than for the presentation of

different odorants (Figure 3E, paired t-test t = 6.15, p = 1.08e-4). This result fits with previous calcium

imaging data both at AL input and output (46, 47). We then performed a hierarchical cluster analysis

using Euclidian distances between odorants. We found a first segregation between odorants with long

chain lengths and odorants with short chain lengths (Figure 3F, C6-C7 vs. C8-C9). Within the shorter

chain length group, a further separation between functional groups appeared based on the oxygen

moiety (C=O for aldehydes and ketones vs. C-OH for alcohols). Within the longer chain length group,

aldehydes were separated from the other odorants. This analysis shows a clear coding depending on

the chemical structure of the odorants, and with remarkably similar results as in previous studies using

non-transgenic approaches (19, 45). Accordingly, inter-odorant distances found in the GCaMP bees

were highly significantly correlated with those measured previously in OSNs using calcium dye bath

application (R^2 = 0.53, Mantel test p = 1.0e-4) or measured in PNs using dye injection (R^2 = 0.44, Mantel

test p = 1.0e-4). The correlation coefficients were not significantly different (R^2 = 0.53 vs. R^2 = 0.44, z =

0.98, p = 0.33, NS), suggesting the possibility that both OSN and PN contribute to the signals recorded

in the AL using genetically encoded GCaMP6f.

Finally, we asked if odorant response maps recorded with the GCaMP bees relate to bees'

actual perception of the odorants. We thus correlated the Euclidian distances measured between all

odor pairs in this study with distances measured in an appetitive conditioning experiment(48). We

found that behavioral distances can be predicted using signals recorded with genetically encoded

GCaMP6f (data not shown; R^2 = 0.34, Mantel test p = 1.0e-4). Thus, odorants evoking similar activity

patterns in the AL are treated as similar by honey bees in their behavior, a further confirmation that

GCaMP6f expression allows to record meaningful neural signals.

Neural recordings in the lateral horn (LH)

We then studied olfactory information processing in higher-order brain centers, first focusing

on the LH. All the aliphatic odorants used in this panel induced calcium signals in the LH (Figure 4A).

Odor-evoked signals systematically followed a biphasic time course, like that recorded in the AL (Figure

4B). In the LH, all but one of the presented odorants induced a response that was significantly higher

than to the air control (RM-ANOVA, odor effect $F_{16,112}$ = 8.091, p = 0.0014, post hoc Dunnett tests, p <

0.04 except p = 0.09 for 2-hexanol). As in the AL, response amplitudes correlated with odorants' vapor

pressures (R^2 = 0.49, $F_{1,14}$ = 13.87, p = 0.0023). However, this correlation appeared to be weaker in the

LH with a near-significant difference between the two correlation coefficients (R^2 = 0.84 in the AL vs.

R^2 = 0.49 in the LH, z = 1.76, p = 0.078). In other words, response intensity in the LH was not a simple

product of the number of volatile molecules in the odor puff, possibly revealing a yet undocumented

gain control mechanism in the LH*.*

We then compared similarity relationships (Euclidian distances among odorants) measured in

the LH and in the AL and found a weak but still significant correlation (R^2 = 0.07, Mantel test p = 0.023).

This suggests a transformation of odor similarity relationships between the two structures. A cluster

analysis performed on Euclidian distances in the LH confirmed this finding. The analysis roughly

segregated alcohols (-OH moiety) on the one side and aldehydes/ketones (=O moiety) on the other.

However, different rules apply here, as the two longest chain molecules nonanal and 2-nonanone

clustered with the former group, while the shortest chain secondary alcohols, 2-hexanol and 2

heptanol clustered with the latter group. Thus, the study of odor coding in the LH using GCaMP bees

suggests less clearly dependence on chemical features than in the AL.

Nevertheless, Euclidian distances measured in the LH correlated significantly with behavioral

distances measures (48) (data not shown; R^2 = 0.12, Mantel test p = 0.0021), demonstrating that

odorants evoking similar activity patterns in the LH are treated as similar by honey bees in their

behavior.

Neural recordings in the mushroom bodies (MB)

Lastly, we turned to the second higher-order center, the mushroom bodies (Figure 5), a

multisensory integration center in the honey bee brain. Calcium signals upon odorant presentation

were observed at the level of the calyx lip. Calcium signals systematically showed a biphasic time

course (Figure 5B) like that recorded in the AL and in the LH, but in all cases with a small first

component and a much stronger second component. In this structure, the five presented odorants

induced significant activity in the MB calyx in comparison to the air control (Figure 5C; RM-ANOVA,

odor effect $F_{5,25}$ = 7.99, p = 0.0033, post hoc Dunnett tests $p < 0.05$, except 2-heptanone with p = 0.07).

However, signal-to-noise ratio was considered too low for further analysis of inter-odor relationships.

DISCUSSION

We developed a line of honey bees with pan-neural expression of a genetically-encoded calcium sensor allowing to record neural activity from honey bee neurons. To our knowledge, this represents the first neurogenetic tool for recording neural activity in a non-dipteran insect and eusocial insect. Neurogenetic tools, especially the use of genetically-encoded calcium indicators (GECIs), have been widely developed in both vertebrates (40, 49) and invertebrates (50) and supported major progress in our understanding of the neural basis of behavior. In insects, such possibility was limited until now to a few Dipteran species (fruit flies and mosquitoes (26)) and was lacking in other insects with different lifestyles. The present development in a social Hymenoptera represents a major progress for the neuroethology of social behavior but also a significant achievement in a species like honey bees in which neurogenetic tools are difficult to establish due to its peculiar reproductive biology and the lack of knowledge of appropriate promoter sequences driving the expressions.

To develop our pan-neuronal calcium indicator, we expressed GCaMP6f under the control of the promoter of the gene coding for synapsin, a vesicle–associated presynaptic protein found throughout the honey bee brain (51). This strategy was successful, as honey bees displayed broad labeling with GCaMP6f in the major brain structures of the nervous system at sufficient amounts for performing calcium imaging. Our neuroanatomical data confirm significant expression throughout the bee brain (optic lobes, antennal lobe, mushroom bodies, all other regions of the protocerebrum), but we noticed substantial cell-to-cell variation in some regions, for instance in the cup of the mushroom body calyx, which contains the somata of type I Kenyon cells (see Figure 1C): there, a strong contrast appeared among neurons, suggesting differences in the level of GCaMP expression among Kenyon cells or even some Kenyon cells lacking GCaMP expression. The GCaMP staining pattern did not correspond to any of the Kenyon cell subtypes previously described based on their differing gene expression profiles (52). In contrast to the synapsin protein, GCaMP6f is located in the entire neuron, since we detected staining in the somata (Figure 1C), the neurites, and the synaptic regions. This widespread expression could represent a great advantage for performing functional recordings in the bee brain.

To assess the power of this new tool for studying brain activity, we focused on the olfactory

system, which is the best-known sensory system in this animal. We compared the results obtained

using the GCaMP6f bees with previous results acquired using conventional dye injections or bath-

application (19, 45). For the first time, we could observe the simultaneous activation of the three main

brain structures involved in insect olfaction, the antennal lobe, the lateral horn, and the mushroom

body calyx. The possibility to record from these structures at the same time is a first and important

step towards the long-term goal to understand brain circuits responsible for bees' sophisticated social

behaviors and cognitive abilities, that may unravel yet undescribed sensory and/or behavior-related

pathways.

To follow this, we first validated our tool using individual structures and we confirmed its

validity for measuring relevant neural activity with regard to bees' behavior. Recordings obtained using

the GCaMP6f-expressing honey bees revealed the same olfactory coding rules in the AL as in previous

studies (19, 53). Thus, odor-induced activity followed the same intensity rule (depending on odorant

vapor pressure – Figure 3D) and inter-odorant neural distance measures were organized according to

odorants' chemical features, chain length, and functional group. Accordingly, calcium imaging

measured with the GCaMP-expressing bees allowed to predict how similarly honey bees perceive these

odorants, since inter-odor relationships among GCaMP response maps correlated with inter-odor

behavioral distances previously measured in an associative conditioning experiment (48). This result,

even if limited to a single sensory modality, is a strong confirmation of the biological pertinence of the

signals measured with this new neurogenetic tool.

In order to demonstrate that the activity can be followed to higher-order brain structures, we

studied the role of the lateral horn in encoding the chemical features of our odorants. The lateral horn

is a poorly studied structure in Hymenoptera, which could only be recorded previously using invasive

dye injection (54). Importantly, while previous recordings only measured from a limited subpopulation

of projection neurons in this structure (the l-ALT), the GCaMP6f-expressing honey bees allowed

recording from the whole structure. The recordings in response to the 16 aliphatic odorants showed

that odor coding in this higher-order center is less clearly influenced by odorant quantity (*i.e.* lower

response amplitude differences among odorants) or by the chemical structure of the odorants than in

the AL. These first results suggest the existence of an additional gain control mechanism in this

structure, fitting with current discussions on its role in *Drosophila* (55, 56) and the possible coding of

hedonic odorant valence rather than chemical features using this new tool.

Calcium signals were also recorded in the mushroom body calyx, and odorants clearly activated

the lip, the olfactory input region of the calyx, as observed previously with a bath-applied dye (57). The

signal amplitude was generally low, which reproduces the known sparseness of the MB odor code (58).

Together with the random arrangement of PN terminals and KC dendritic arbors in the lip, this

prevented the observation of odor-specific response maps at this point. However, advanced 3D

microscopy techniques (59) will benefit from this neurogenetic marker to unravel coding mechanisms

also in this neuropil, as experiments in Drosophila promise (60).

We expect that recording neuronal activity simultaneously and in detail in brain structures

using a genetically expressed Ca reporter in a eusocial insect, the honeybee, as demonstrated here,

will open new possibilities to decipher the neural bases of the sophisticated social behavior and its

associations with higher cognitive abilities. It will also allow the study of other sensory modalities than

the most commonly studied olfaction and vision. As discussed above, this new development enables

to record neural activity in different regions of the brain at the same time, which could become a major

advantage over non-transgenic approaches, as it allows to reveal a whole neural circuit underlying a

specific behavior. Moreover, the study of mechanosensory information processing notably used during

the waggle dance remains limited(61, 62) and should be facilitated using GCaMP-expressing honey

bees. Lastly, transgenic bees could be instrumental for studying neural signals in response to

multisensory inputs, for instance in the mushroom bodies, the higher-order center known for its role

in multimodal sensory integration(63).

The future of insect neuroethology beyond the use of model organisms like *Drosophila* will

depend on the further development of neurogenetic tools in species like honeybees, with their rich

behavioral repertoire involving intricate social behaviors as well as elaborated cognitive skills. First, a

better resolution will have to be achieved using cell-type-specific gene expression to visualize (*i.e.* using

fluorescent proteins such as GFP) or monitor activity (using GECIs as here) from specific neuronal

populations. The same precision shall then be exploited to specifically silence or activate specific

neuron populations, allowing a temporal control of neural function. We hope that the new line of

GCaMP-expressing bees will represent the first step in this direction.

MATERIAL AND METHODS

Piggy-Bac **transgenesis**

The promoter region together with the three exons of the 5' UTR of the honey bee *synapsin* (*syn*) gene

(LOC551737; assembly Amel_4.5) were successively cloned (Sup. Figure 2). This single fragment (1811

bp) was inserted into the PBac plasmid(33) using the *Asc*I/*Nco*I restriction sites. The coding sequence

of *GCaMP6f* was inserted downstream of the promoter region using *Nco*I/*Mss*I restriction sites that

finally resulted in the PBac [*syn-GCaMP6F*] plasmid. Egg injections, larvae and queen rearing were

performed following standard methodology by employing the hyperactive transposase which we have

customized for the honey bee (33, 41). A total of 4231 honey bee eggs not older than 1.5 h were

injected and the hatched larvae were reared to queens. Semen from wildtype drones was used for

instrumental insemination of queens. Inseminated queens were laying fertilized, usually female-

determined eggs that were reared either to queens or worker bees. Transgenic queens were identified

by amplification of the transgene in the offspring(41) using oligonucleotide primers

(CCACACCTCCCCCTGAACCTGAAAC and GAGGTAAGAATAAACATTGTTGGTC) targeting the PBac

sequence.

Anatomical immunostaining

To visualize the expression pattern of the GCaMP6f in the honey bee brain, transgenic workers were caught at emergence and were maintained in an incubator in the dark at 34°C for 2 weeks (*i.e.* the same age as for calcium imaging). The brains of workers were dissected and immediately immersed in cold 1% zinc formaldehyde in PBS (ZnFa 1%(64)) and kept overnight at 4°C. Brains were then washed six times in PBS (10 min each), permeabilized in PBS containing 1% Triton X-100 for 30 min, and pre-incubated 3h in PBS containing 0.3% Triton and 1% BSA (Bovine Serum Albumine in PBS, #37525, ThermoScientific). To stain *GCaMP6f* and *synapsin* (as background staining), the brains were then incubated in PBS containing 0.3% Triton and 0.1% BSA with rabbit polyclonal anti-GFP (Thermofisher, #A-11122, France) at 1:100 dilution and mouse monoclonal anti-SYNORF1 (DHSB, #3C11, US) at 1:100 dilution for 7 days. Brains were then washed 6 times in PBS containing 0.3% Triton and incubated in secondary antibodies directed against rabbit coupled to Alexa 488 (Thermofisher, #A11122, France) and against mouse coupled to Alexa 555 (Thermofisher, #A-21147, France) both diluted at 1:200 for 5 days. Brains were then washed in PBS, dehydrated in an ascending ethanol series (30% to 100%), cleared, and finally mounted in methyl salicylate (M6752, Sigma-Aldrich, France) for observation. Brains were scanned using a laser-scanning confocal microscope (Zeiss LSM 700) with a W Plan-Apochromat 20x/NA 1 objective using sequential excitation wavelengths of 488 nm and 555 nm, observing via two color-filtered channels around 510 nm and 590 nm, respectively.

Honey bee preparation for *in vivo* calcium imaging

Transgenic workers were collected at emergence and were maintained in small cages in the dark at 34°C for 2 weeks. On the day of the experiment, they were individually chilled on ice for 5 min until they stopped moving. Then, they were prepared following the standard preparation used to image the ventral part of the honey bee brain(13, 19, 44, 47). Briefly, the honey bee was fixed in a plastic chamber with its antennae oriented to the front and the proboscis was fixed using beeswax to avoid movement of the brain during the experiment. A pool was built with beeswax and pieces of plastic around the head capsule, and a small window was then cut in the head cuticle. Glands as well

as trachea were removed to expose the brain, and the pool was filled with ringer solution (in mM: NaCl, 130; KCl, 6; MgCl$_2$, 4; CaCl$_2$, 5; sucrose, 160; glucose, 25; Hepes, 10; pH 6.7, 500 mOsmol; all chemicals from Sigma-Aldrich, France), to avoid desiccation of the brain surface. The honey bee was then left 30 min in a moisturized and dark place before the calcium imaging.

Calcium imaging

A T.I.L.L. Photonics imaging system (Martinsried, Germany) was used to perform *in vivo* optical recordings, as described elsewhere(19, 65). An epifluorescence microscope (Olympus BX51WI) was used to record activity in the different regions of the brain with either a 4× dry objective (Olympus, PlanCN; NA 0.10) for whole-brain recordings, a 10× water-immersion objective (Olympus, UMPlanFL; NA 0.3) for AL recordings or a 20x water-immersion objective (Olympus, UMPlanFL; NA 0.5) for LH and MB recordings. GCaMP6f was excited using 475 nm monochromatic light (T.I.L.L. Polychrom IV). The fluorescence signal was separated by a 505 nm dichroic filter and a long-pass 515 nm emission filter and recorded with a 640 × 480 pixels 12-bit monochrome CCD camera (T.I.L.L. Imago) cooled to -12°C with 4 × 4 binning on chip. Each measurement consisted of 100 frames recorded at a rate of 5 Hz (integration time for each frame ~50ms).

Odor stimuli

A constant airstream was directed from a distance of 1 cm to the bee's antennae, and odor stimuli were given at the 15th frame for 1 s. For each odor stimulus (all obtained from Sigma-Aldrich, France), 5 µL of the solution were deposited on a filter paper inserted in a Pasteur pipette. A pipette containing a clean piece of filter paper was used as control stimulus.

For whole-brain recordings, we tested a small set of 3 odorant stimuli known from previous work to trigger strong neural activity: 2-heptanone, octanal, and heptanal. For AL and LH recordings, we tested 16 aliphatic odorants previously used in calcium imaging studies (19, 45) belonging to 4

functional groups (primary and secondary alcohols, aldehydes, and ketones) and having 4 different

carbon chain lengths (6, 7, 8, and 9 carbons). The use of these odorants which are part of floral blends

honey bees encounter while foraging (Knudsen et al. 1993) allowed to compare the results obtained

in this study using GCaMP6f imaging with previous studies using classical calcium-sensitive dyes such

as Fura-2 dextran(19) or Calcium green 2-AM(45). These 16 odorants were used to record neural

activity at the level of the AL and at the level of the LH on two different groups of bees. For recordings

at the level of the MB, a reduced list of odorants which give strong neural activity were used: 1-hexanol,

2-hexanone, 2-heptanone, heptanal, octanal, and the air control.

Each odorant stimulus was presented twice in a pseudo-randomized order, avoiding the

consecutive presentation of stimuli with the same functional group or the same carbon chain length.

Data processing and analyses

All analyses were carried out using custom-made software written in IDL 6.0 (Research

Systems, Boulder, CO). Each odor presentation corresponded to a three-dimensional array consisting

of two spatial dimensions (*x*- and *y*-coordinates) along time (100 frames). First, the relative

fluorescence changes were calculated as $\Delta F/F = (F - F_0)/F_0$ by taking as reference background F_0 the

average of three frames just before the odorant stimulation (frames 9-11). Possible irregularities of

lamp illumination and bleaching were corrected by subtracting the median pixel value of each frame

from every single pixel of the corresponding frame. Finally, the two spatial dimensions were filtered

with a Gaussian filter of window size 7×7 pixels for AL recordings or 3×3 pixels for LH and MB recordings

for noise reduction. A biphasic calcium signal was observed in all recordings. As in previous studies

using bath-applied Calcium Green(45), a high contrast measure for the intensity of the odor-induced

response was obtained by averaging three consecutive frames at the end of the odor presentation

(frames 19-21) and subtracting the average of 3 frames during the second, negative component of the

signal (frames 49-51).

For the quantification of response intensity and similarity relationships, a mask was precisely

drawn in order to exclude regions outside of the imaged structure. A pixel-wise analysis was then

performed, using all the pixels within the mask. Response intensity was calculated by averaging the

intensity of all pixels located within the unmasked area. Similarity relationships between neural activity

patterns $I(x, y)$ were calculated by measuring Euclidian distances

447
$$d(o_1, o_2) = \sqrt{\sum_{x,y} (I_{o_1}(x, y) - I_{o_2}(x, y))^2}$$

for all the 120 odorant pairs o_1, o_2 within each animal.

For all analyses, average values for the two presentations of each odorant were used except

for the comparison of Euclidian distances for the same vs. different odorants (Figure 3E) in which each

single odorant presentation was used.

We used the Euclidian distances between odorants calculated using behavioral experiments

by Guerrieri et al.(48) to compare similarity relationships among odorants at the neural level with data

obtained at the behavioral level. All results are displayed as means over individuals ± SEM.

Statistical analyses

Odor response intensities were compared with ANOVA for repeated measurements, using

odors as within-group factors. A Dunnett post hoc test was applied to compare the intensity of the

response to each stimulus with a common reference, the air control. Wilcoxon matched-pairs tests

were applied to compare Euclidian distances between the same and different odors for AL recordings.

Pearson correlation analyses were performed between response intensity and the logarithm of

odorants' vapor pressure in mmHg (16 aliphatic odors recorded in the AL and in the LH), and between

response intensities recorded in this and previous studies (OSNs(45) or PNs(19)). A Fisher test, giving

a z value, was used to test the significance of the linear regression following Pearson correlation (R

package *multilevel*, *cordif* function). Mantel tests were used to compare matrices containing the

Euclidian distances for the 120 possible odor-pair among 16 odorants, either between different

imaging studies, or between neural and behavioral measures (R package *ade4*, *mantel.rtest* function).

To explore similarity relationships among odorants, hierarchical clustering using Ward's classification

method was used (R package *stats*, *hclust* function). All tests were performed with GraphPadPrism

V7.00 and R (www.r-project.org).

ACKNOWLEDGEMENTS

We thank Robin Parlow, Eva Theilenberg, Marion Müller-Borg and Virginie Larcher for technical

assistance. We thank Sara Bariselli for her support with the egg injection and Amélie Cabirol for fruitful

discussions. We thank the French Research National Agency (ANR to JCS, Project ANR-17-CE20-0003

Bee-O-Choc), the CNRS (JCS) and the University Paris-Saclay (JC and JCS), the DFG (Deutsche

Forschungsgemeinschaft to MB) for funding.

AUTHOR CONTRIBUTIONS

J.C., M.O., A.H., J.C.S., and M.B. conceived the experiments. J.C. and M.O. collected the data. J.C., M.O.,

A.H., J.C.S., and M.B. analysed the data and interpreted the results. J.C., A.H., J.C.S., and M.B. wrote

the manuscript. All authors read and approved the final version of the manuscript.

COMPETING INTEREST STATEMENT

The present research project was conducted in the absence of any commercial relationships that could

be construed as a potential conflict of interest.

	FIGURE LEGENDS

Figure 1: Genetically encoded GCaMP6f and its neural expression

A- Scheme of the honey bee *synapsin* promoter *GCaMP6f* expression cassette. CDS: coding sequence;

UTR: untranslated region; ATG: translation start. **B-** GCaMP6f expression in the honey bee brain

revealed by anti-GFP immunostaining (left, in green). GCaMP6f is widely expressed in the bee brain,

including in somata and neural tracts (white arrows). For comparison with *synapsin* expression, an anti-

SYNORF1 immunostaining (right, in red) was also performed. Scale bar=200 μm **C-** GCaMP6f expression

(anti-GFP in green) and *synapsin* expression (anti-SYNORF1 in red) in the AL (left) and in the mushroom

bodies (right). Remarkable and strong expression is observed in the somata of projection neurons and

local neurons (near the AL) and in some somata of Kenyon cells (in the cup of the calyces). Scale bar=50

μm. AL: Antennal Lobe, MB: Mushroom body, OL: Optic Lobe, Lo: Lobulla, Me: Medulla, vL: vertical

lobe, lc: lateral cluster of projection neurons somata, mc: medial cluster of projection neurons somata,

Li: Lip, BR: Basal Ring, Co: Collar.

Figure 2: Neural activity in the whole brain of the honey bee

A- Hymenopteran olfactory pathway (adapted from (45)). Odorant molecules are detected by olfactory

sensory neurons (OSNs) on the antenna, which send olfactory information to the antennal lobe (AL).

Then, projection neurons (PNs) convey information to higher-order centers, the mushroom body (MB)

and the lateral horn (LH). Lo: lobula, Me: medulla, v: vertical lobe, h: horizontal lobe. **B-** Calcium signals

in the whole brain evoked by 3 different odorants (2-heptanone, octanal, heptanal) and the air control.

Relative fluorescence changes ($\Delta F/F$ [%]) are presented in a false-color code, from dark blue (minimal

response) to red (maximal response). **C-** Time course of odor-evoked responses ($\Delta F/F$ [%]) for one

individual to the presentation of 2-heptanone (in blue), octanal (in pink), and the air control (in grey),

simultaneously recorded in the AL (left), the LH (middle) and the MB (right). The calcium signals show

a biphasic time course, with a fluorescence increase upon odorant presentation (blue bars) followed

by a long-lasting fluorescence undershoot, in all brain regions.

Figure 3: Neural activity recorded in the antennal lobe (AL)

A- GCaMP6f calcium signals in the antennal lobe evoked by a panel of 16 odorants varying

systematically according to their carbon chain length (C6–C9) and their functional group (primary and

secondary alcohols, aldehydes, and ketones). Different odorants induce different glomerular activity

patterns. **B-** Time course of odor-evoked responses ($\Delta F/F$ [%]) recorded in the AL (in white squares

shown in A) to 2-heptanone (in blue), octanal (in pink), and to the air control (in grey). **C-** Amplitude of

calcium responses ($\Delta F/F$ [%]) to the 16 aliphatic odorants and to the air control. All odorants induce

significant activity in comparison to the air control ($n = 11$, ** $p < 0.0097$). **D-** Amplitude of calcium

responses ($\Delta F/F$ [%]) as a function of odorant vapor pressure (in log units). The linear regression shows

a significant correlation ($R^2 = 0.84$, *** $p = 6e\text{-}7$). **E-** Different presentations of the same odorants (2-

octanone and heptanal) show similar glomerular patterns in the AL (left). Dissimilarity measures

(Euclidian distance) between representations of the same or of different odorants. Activity maps are

more similar (shorter distances) when the same odorant is presented (*** $p = 1.1e\text{-}4$), showing, as

expected, a clear odor coding in this structure. **F-** Cluster analysis showing similarity relationships

among odorants (Ward's classification method). Functional groups are shown in different colors:

primary alcohols in blue, secondary alcohols in green, aldehydes in black, and ketones in red. The

analysis shows a first separation between odorants with short and long carbon chain lengths. Odorants

with a short carbon chain are then subdivided into alcohols (primary and secondary, C-OH functional

group) and ketones/aldehydes (C=O functional group).

Figure 4: Neural activity recorded in the lateral horn (LH)

A- Calcium signals in the lateral horn evoked by the same panel of 16 odorants. Different odorants

induce different activity patterns in the LH. **B-** Time course of odor-evoked responses ($\Delta F/F$ [%])

recorded in the LH (in white squares shown in A) to 2-heptanone (in blue), octanal (in pink), and to the

air control (in grey). The calcium signals also show a biphasic response, with a fluorescence increase

upon odor presentation (blue bar) followed by a long undershoot. **C-** Amplitude of calcium responses

($\Delta F/F$ [%]) to the 16 aliphatic odorants and to the air control. All odorants induce a significant activity

in comparison to the air control *(n = 8, * p < 0.04)*. **D-** Cluster analysis showing similarity relationships

among odorants (Ward's classification method).

Figure 5: Neural activity recorded in the mushroom bodies (MB)

A- Calcium signals in the calyces of the mushroom bodies in response to the presentations of 2-

heptanone, heptanal, octanal, and the air control. **B-** Time course of odor-evoked responses ($\Delta F/F$ [%])

recorded in the lip of the MB (white squares shown in A) to 2-heptanone (in blue), octanal (in pink),

and to the air control (in grey). Calcium signals are weaker here than in other structures, with a smaller

first component and a larger second component. **C-** Amplitude of calcium responses ($\Delta F/F$ [%]) to the

5 tested odorants and to the air control. All odorants induce significant activity in comparison to the

air control ($n = 6$, * $p < 0.05$, except 2-heptanone with $p = 0.07$).

Supplementary Figure 1: Correlation between amplitudes of calcium responses recorded in the AL

A- Amplitude of calcium responses ($\Delta F/F$ [%]) recorded in Olfactory Sensory Neurons (OSNs) in[45] as

a function of amplitude of calcium responses ($\Delta F/F$ [%]) recorded in the AL in this study. The linear

regression shows a significant correlation ($R^2 = 0.84$, *** $p = 6e\text{-}7$). **B-** Amplitude of calcium responses

($\Delta F/F$ [%]) recorded in Projection Neurons (PNs) in[19] as a function of amplitude of calcium responses

($\Delta F/F$ [%]) recorded in the AL in this study. The linear regression shows a significant correlation ($R^2 =$

0.78, *** $p = 6e\text{-}6$).

Supplementary Figure 2: Synapsin sequence of the honey bee

The promoter and the 5' UTR sequence from the honey bee synapsin (syn) gene. Different colors

indicate the different exons that were fused.

REFERENCES

1.   Bourke AF (2011) *Principles of social evolution* (Oxford University Press).
2.   Hamilton WD (1964) The genetical evolution of social behaviour. II. *Journal of theoretical*
569   *biology* 7(1):17-52.
570   3.   Hölldobler B & Wilson EO (1990) *The ants* (Harvard University Press) pp 1-732.
571   4.   Wilson EO (2012) *The social conquest of earth* (WW Norton & Company).
572   5.   Oster GF & Wilson EO (1978) Caste and ecology in the social insects. *Monogr Popul Biol* 12:1-
573   352.
574   6.   Paoli M & Galizia GC (2021) Olfactory coding in honeybees. *Cell and Tissue Research*:1-24.
575   7.   Woodgate JL*, et al.* (2021) Harmonic radar tracking reveals that honeybee drones navigate
576   between multiple aerial leks. *iScience* 24(6):102499.
577   8.   Avargues-Weber A, Dyer AG, Combe M, & Giurfa M (2012) Simultaneous mastering of two
578   abstract concepts by the miniature brain of bees. *Proc Natl Acad Sci U S A* 109(19):7481-7486.
579   9.   Chole H*, et al.* (2019) Social Contact Acts as Appetitive Reinforcement and Supports Associative
580   Learning in Honeybees. *Curr Biol* 29(8):1407-1413 e1403.
581   10.   Barron AB & Plath JA (2017) The evolution of honey bee dance communication: a mechanistic
582   perspective. *J Exp Biol* 220(Pt 23):4339-4346.
583   11.   Giurfa M & Sandoz JC (2012) Invertebrate learning and memory: Fifty years of olfactory
584   conditioning of the proboscis extension response in honeybees. *Learn Mem* 19(2):54-66.
585   12.   Wycke MA, Coureaud G, Thomas-Danguin T, & Sandoz JC (2020) Configural perception of a
586   binary olfactory mixture in honey bees, as in humans, rodents and newborn rabbits. *J Exp Biol*
587   223(Pt 21).
588   13.   Paoli M*, et al.* (2018) Neuronal Response Latencies Encode First Odor Identity Information
589   across Subjects. *J Neurosci* 38(43):9240-9251.
590   14.   Farris SM (2016) Insect societies and the social brain. *Curr Opin Insect Sci* 15:1-8.
591   15.   Lihoreau M, Latty T, & Chittka L (2012) An exploration of the social brain hypothesis in insects.
592   *Front Physiol* 3:442.
593   16.   Cabirol A, Brooks R, Groh C, Barron AB, & Devaud JM (2017) Experience during early adulthood
594   shapes the learning capacities and the number of synaptic boutons in the mushroom bodies
595   of honey bees (Apis mellifera). *Learn Mem* 24(10):557-562.
596   17.   Suenami S, Iino S, & Kubo T (2018) Pharmacologic inhibition of phospholipase C in the brain
597   attenuates early memory formation in the honeybee (Apis mellifera L.). *Biol Open* 7(1).
598   18.   Duer A, Paffhausen BH, & Menzel R (2015) High order neural correlates of social behavior in
599   the honeybee brain. *J Neurosci Methods* 254:1-9.
600   19.   Carcaud J, Giurfa M, & Sandoz JC (2018) Differential Processing by Two Olfactory Subsystems
601   in the Honeybee Brain. *Neuroscience* 374:33-48.
602   20.   Sherer LM & Certel SJ (2019) The fight to understand fighting: neurogenetic approaches to the
603   study of aggression in insects. *Current opinion in insect science* 36:18-24.
604   21.   McKelvey EG & Fabre CC (2019) Recent neurogenetic findings in insect courtship behaviour.
605   *Current opinion in insect science* 36:103-110.
606   22.   Ehmann N & Pauls D (2020) Optogenetics: illuminating neuronal circuits of memory formation.
607   *Journal of neurogenetics* 34(1):47-54.
608   23.   Sadanandappa MK, Sathyanarayana SH, Kondo S, & Bosco G (2021) Neuropeptide F signaling
609   regulates parasitoid-specific germline development and egg-laying in Drosophila. *PLoS*
610   *genetics* 17(3):e1009456.
611   24.   Pooryasin A*, et al.* (2021) Unc13A and Unc13B contribute to the decoding of distinct sensory
612   information in Drosophila. *Nature Communications* 12(1):1-14.
613   25.   Grover D, Katsuki T, Li J, Dawkins TJ, & Greenspan RJ (2020) Imaging brain activity during
614   complex social behaviors in Drosophila with Flyception2. *Nature communications* 11(1):1-10.
615   26.   Jové V*, et al.* (2020) Sensory discrimination of blood and floral nectar by aedes aegypti
616   mosquitoes. *Neuron*.

27. Vinauger C*, et al.* (2019) Visual-olfactory integration in the human disease vector mosquito
Aedes aegypti. *Current Biology* 29(15):2509-2516. e2505.
28. Koutroumpa FA*, et al.* (2016) Heritable genome editing with CRISPR/Cas9 induces anosmia in
a crop pest moth. *Sci Rep* 6:29620.
29. Fandino RA*, et al.* (2019) Mutagenesis of odorant coreceptor Orco fully disrupts foraging but
not oviposition behaviors in the hawkmoth Manduca sexta. *Proceedings of the National*
Academy of Sciences 116(31):15677-15685.
30. DeGennaro M*, et al.* (2013) orco mutant mosquitoes lose strong preference for humans and
are not repelled by volatile DEET. *Nature* 498(7455):487-491.
31. McMeniman CJ, Corfas RA, Matthews BJ, Ritchie SA, & Vosshall LB (2014) Multimodal
integration of carbon dioxide and other sensory cues drives mosquito attraction to humans.
Cell 156(5):1060-1071.
32. Zhang ZJ*, et al.* (2019) A determining factor for insect feeding preference in the silkworm,
Bombyx mori. *PLoS Biol* 17(2):e3000162.
33. Schulte C, Theilenberg E, Muller-Borg M, Gempe T, & Beye M (2014) Highly efficient integration
and expression of piggyBac-derived cassettes in the honeybee (Apis mellifera). *Proc Natl Acad*
Sci U S A 111(24):9003-9008.
34. Roth A*, et al.* (2019) A genetic switch for worker nutrition-mediated traits in honeybees. *PLoS*
biology 17(3):e3000171.
35. Chen Z*, et al.* (2021) Neurodevelopmental and transcriptomic effects of CRISPR/Cas9-induced
somatic orco mutation in honey bees. *Journal of Neurogenetics*:1-13.
36. Değirmenci L*, et al.* (2020) CRISPR/Cas 9-mediated mutations as a new tool for studying taste
in honeybees. *Chemical Senses* 45(8):655-666.
37. Nie H-Y*, et al.* (2021) CRISPR/Cas9 mediated knockout of Amyellow-y gene results in
melanization defect of the cuticle in adult Apis mellifera. *Journal of Insect Physiology*
132:104264.
38. Wang JW, Wong AM, Flores J, Vosshall LB, & Axel R (2003) Two-photon calcium imaging reveals
an odor-evoked map of activity in the fly brain. *Cell* 112(2):271-282.
39. Muto A*, et al.* (2011) Genetic visualization with an improved GCaMP calcium indicator reveals
spatiotemporal activation of the spinal motor neurons in zebrafish. *Proc Natl Acad Sci U S A*
108(13):5425-5430.
40. Chen TW*, et al.* (2013) Ultrasensitive fluorescent proteins for imaging neuronal activity. *Nature*
499(7458):295-300.
41. Otte M*, et al.* (2018) Improving genetic transformation rates in honeybees. *Sci Rep* 8(1):16534.
42. Strausfeld NJ (2002) Organization of the honey bee mushroom body: representation of the
calyx within the vertical and gamma lobes. *J Comp Neurol.* 450(1):4-33.
43. Suenami S, Oya S, Kohno H, & Kubo T (2018) Kenyon Cell Subtypes/Populations in the
Honeybee Mushroom Bodies: Possible Function Based on Their Gene Expression Profiles,
Differentiation, Possible Evolution, and Application of Genome Editing. *Front Psychol* 9:1717.
44. Joerges J, Küttner A, Galizia CG, & Menzel R (1997) Representations of odours and odour
mixtures visualized in the honeybee brain. *Nature* 387:285-288.
45. Carcaud J, Hill T, Giurfa M, & Sandoz JC (2012) Differential coding by two olfactory subsystems
in the honeybee brain. *J Neurophysiol* 108(4):1106-1121.
46. Galizia CG, Nägler K, Hölldobler B, & Menzel R (1998) Odour coding is bilaterally symmetrical
in the antennal lobe of honeybees (Apis mellifera   ). *European Journal of Neuroscience*
10:2964-2974.
47. Sachse S & Galizia CG (2002) The role of inhibition for temporal and spatial odor representation
in olfactory output neurons: a calcium imaging study. *Journal of Neurophysiology* 87:1106-
1117.
48. Guerrieri F, Schubert M, Sandoz JC, & Giurfa M (2005) Perceptual and neural olfactory
similarity in honeybees. *PLoS Biology* 3(4):e60.

49. Lin MZ & Schnitzer MJ (2016) Genetically encoded indicators of neuronal activity. *Nat Neurosci*
19(9):1142-1153.
50. Dana H*, et al.* (2019) High-performance calcium sensors for imaging activity in neuronal
populations and microcompartments. *Nat Methods* 16(7):649-657.
51. Groh C, Lu Z, Meinertzhagen IA, & Rössler W (2012) Age-related plasticity in the synaptic
ultrastructure of neurons in the mushroom body calyx of the adult honeybee Apis mellifera. *J
Comp Neurol* 520(15):3509-3527.
52. Suenami S, Oya S, Kohno H, & Kubo T (2018) Kenyon cell subtypes/populations in the honeybee
mushroom bodies: possible function based on their gene expression profiles, differentiation,
possible evolution, and application of genome editing. *Frontiers in psychology* 9:1717.
53. Sachse S, Rappert A, & Galizia CG (1999) The Spatial representation of chemical structures in
the antennal lobe of honeybees: steps towards the olfactory code. *European Journal of
Neuroscience* 11:3970-3982.
54. Roussel E, Carcaud J, Combe M, Giurfa M, & Sandoz JC (2014) Olfactory coding in the honeybee
lateral horn. *Curr Biol* 24(5):561-567.
55. Strutz A*, et al.* (2014) Decoding odor quality and intensity in the Drosophila brain. *eLife*
3:e04147.
56. Dolan MJ*, et al.* (2019) Neurogenetic dissection of the Drosophila lateral horn reveals major
outputs, diverse behavioural functions, and interactions with the mushroom body. *eLife* 8.
57. Faber T & Menzel R (2001) Visualizing mushroom body response to a conditioned odor in
honeybees. *Naturwissenschaften* 88(11):472-476.
58. Szyszka P, Ditzen M, Galkin A, Galizia CG, & Menzel R (2005) Sparsening and temporal
sharpening of olfactory representations in the honeybee mushroom bodies. *J.Neurophysiol.*
94(5):3303-3313.
59. Paoli M & Haase A (2018) In Vivo Two-Photon Imaging of the Olfactory System in Insects.
Methods in molecular biology 1820:179-219.
60. Bilz F, Geurten BRH, Hancock CE, Widmann A, & Fiala A (2020) Visualization of a Distributed
Synaptic Memory Code in the Drosophila Brain. *Neuron* 106(6):963-976 e964.
61. Ai H, Kai K, Kumaraswamy A, Ikeno H, & Wachtler T (2017) Interneurons in the Honeybee
Primary Auditory Center Responding to Waggle Dance-Like Vibration Pulses. *J Neurosci*
37(44):10624-10635.
62. Tiraboschi E, Leonardelli L, Segata G, Rigosi E, & Haase A (2021) Parallel processing of olfactory
and mechanosensory information in the honeybee antennal lobe.
bioRxiv:2021.2005.2011.443598.
63. Groh C & Rössler W (2020) Analysis of synaptic microcircuits in the mushroom bodies of the
honeybee. *Insects* 11(1):43.
64. Ott SR (2008) Confocal microscopy in large insect brains: zinc-formaldehyde fixation improves
synapsin immunostaining and preservation of morphology in whole-mounts. *J Neurosci
Methods* 172(2):220-230.
65. Carcaud J, Giurfa M, & Sandoz JC (2015) Differential combinatorial coding of pheromones in
two olfactory subsystems of the honey bee brain. *J Neurosci* 35(10):4157-4167.

Figure 1

Figure 1: Genetically encoded GCaMP6f and its neural expression

A- Scheme of the honey bee *synapsin* promoter *GCaMP6f* expression cassette. CDS: coding sequence; UTR: untranslated region; ATG: translation start. **B-** GCaMP6f expression in the honey bee brain revealed by anti-GFP immunostaining (left, in green). GCaMP6f is widely expressed in the bee brain, including in somata and neural tracts (white arrows). For comparison with *synapsin* expression, an anti-SYNORF1 immunostaining (right, in red) was also performed. Scale bar=200 µm **C-** GCaMP6f expression (anti-GFP in green) and *synapsin* expression (anti-SYNORF1 in red) in the AL (left) and in the mushroom bodies (right). Remarkable and strong expression is observed in the somata of projection neurons and local neurons (near the AL) and in some somata of Kenyon cells (in the cup of the calyces). Scale bar=50 µm. AL: Antennal Lobe, MB: Mushroom body, OL: Optic Lobe, Lo: Lobulla, Me: Medulla, vL: vertical lobe, lc: lateral cluster of projection neurons somata, mc: medial cluster of projection neurons somata, Li: Lip, BR: Basal Ring, Co: Collar.

Figure 2

Figure 2: Neural activity in the whole brain of the honey bee
A- Hymenopteran olfactory pathway (adapted from [40]). Odorant molecules are detected by olfactory sensory neurons (OSNs) on the antenna, which send olfactory information to the antennal lobe (AL). Then, projection neurons (PNs) convey information to higher-order centers, the mushroom body (MB) and the lateral horn (LH). Lo: lobula, Me: medulla, v: vertical lobe, h: horizontal lobe. **B-** Calcium signals in the whole brain evoked by 3 different odorants (2-heptanone, octanal, heptanal) and the air control. Relative fluorescence changes ($\Delta F/F$ [%]) are presented in a false-color code, from dark blue (minimal response) to red (maximal response). **C-** Time course of odor-evoked responses ($\Delta F/F$ [%]) for one individual to the presentation of 2-heptanone (in blue), octanal (in pink), and the air control (in grey), simultaneously recorded in the AL (left), the LH (middle) and the MB (right). The calcium signals show a biphasic time course, with a fluorescence increase upon odorant presentation (blue bars) followed by a long-lasting fluorescence undershoot, in all brain regions.

Figure 3

Figure 3: Neural activity recorded in the antennal lobe (AL)
A- GCaMP6f calcium signals in the antennal lobe evoked by a panel of 16 odorants varying systematically according to their carbon chain length (C6–C9) and their functional group (primary and secondary alcohols, aldehydes, and ketones). Different odorants induce different glomerular activity patterns. **B-** Time course of odor-evoked responses ($\Delta F/F$ [%]) recorded in the AL (in white squares shown in A) to 2-heptanone (in blue), octanal (in pink), and to the air control (in grey). **C-** Amplitude of calcium responses ($\Delta F/F$ [%]) to the 16 aliphatic odorants and to the air control. All odorants induce significant activity in comparison to the air control ($n = 11$, ** $p < 0.0097$). **D-** Amplitude of calcium responses ($\Delta F/F$ [%]) as a function of odorant vapor pressure (in log units). The linear regression shows a significant correlation ($R^2 = 0.84$, *** $p = 6e{-7}$). **E-** Different presentations of the same odorants (2-octanone and heptanal) show similar glomerular patterns in the AL (left). Dissimilarity measures (Euclidian distance) between representations of the same or of different odorants. Activity maps are more similar (shorter distances) when the same odorant is presented (*** $p = 1.1e{-4}$), showing, as expected, a clear odor coding in this structure. **F-** Cluster analysis showing similarity relationships among odorants (Ward's classification method). Functional groups are shown in different colors: primary alcohols in blue, secondary alcohols in green, aldehydes in black, and ketones in red. The analysis shows a first separation between odorants with short and long carbon chain lengths. Odorants with a short carbon chain are then subdivided into alcohols (primary and secondary, C-OH functional group) and ketones/aldehydes (C=O functional group).

Figure 4

Figure 4: Neural activity recorded in the lateral horn (LH)

A- Calcium signals in the lateral horn evoked by the same panel of 16 odorants. Different odorants induce different activity patterns in the LH. **B-** Time course of odor-evoked responses ($\Delta F/F$ [%]) recorded in the LH (in white squares shown in A) to 2-heptanone (in blue), octanal (in pink), and to the air control (in grey). The calcium signals also show a biphasic response, with a fluorescence increase upon odor presentation (blue bar) followed by a long undershoot. **C-** Amplitude of calcium responses ($\Delta F/F$ [%]) to the 16 aliphatic odorants and to the air control. All odorants induce a significant activity in comparison to the air control ($n = 8$, * $p < 0.04$). **D-** Cluster analysis showing similarity relationships among odorants (Ward's classification method).

Figure 5

Figure 5: Neural activity recorded in the mushroom bodies (MB)
A- Calcium signals in the calyces of the mushroom bodies in response to the presentations of 2-heptanone, heptanal, octanal, and the air control. **B-** Time course of odor-evoked responses ($\Delta F/F$ [%]) recorded in the lip of the MB (white squares shown in A) to 2-heptanone (in blue), octanal (in pink), and to the air control (in grey). Calcium signals are weaker here than in other structures, with a smaller first component and a larger second component. **C-** Amplitude of calcium responses ($\Delta F/F$ [%]) to the 5 tested odorants and to the air control. All odorants induce significant activity in comparison to the air control (n = 6, * p < 0.05, except 2-heptanone with p =0.07).

Sup. Figure 1

Supplementary Figure 1: Correlation between amplitudes of calcium responses recorded in the AL
A- Amplitude of calcium responses ($\Delta F/F$ [%]) recorded in Olfactory Sensory Neurons (OSNs) in[40] as a function of amplitude of calcium responses ($\Delta F/F$ [%]) recorded in the AL in this study. The linear regression shows a significant correlation (R^2 = 0.84, *** p = 6e-7). **B-** Amplitude of calcium responses ($\Delta F/F$ [%]) recorded in Projection Neurons (PNs) in[26] as a function of amplitude of calcium responses ($\Delta F/F$ [%]) recorded in the AL in this study. The linear regression shows a significant correlation (R^2 = 0.78, *** p = 6e-6).

ACGAGAACCGGGAAGCGTAGTGATAATCTTAATGAAAATCCCCCACGTTCGTGCCTCCTTTGATAGTTTTCACC
CCGGGAAAATGTTTGTGACCCACTTTTCCCACAGAAAACGCTCTAATGCCGTTTTACCTGGGGTGCAAAAAGGT
AGAGTGGGGAAAAAAAAAAAAAAAAAGAGGCGGAACCGGAGGAGAGAAGAGAAGAGAGAACCCCGTCTTCT
ACTTGAATACCGTGTCTCGCGCCGTATACGCAAACGTGTACAATATATATAACAATTTTATTGCAAGAATTTCAA
ACAACGCTACGGAGATCAAGGGATCATATAAGAATTTTTATACGTCCTTACAACAGGAAAATTCGTTCCCAAGA
TTCCACGAGTTGTAACGCATGCTGTCTAACCGCACGCATACACACGCACCCCGATACACAATAGAGAAACTCTC
TCTCCTCTCTCTCTCTCCCGAAACCGTTTCAAAAATGTACGATCTCCGTAAGATCGTACAAAAGCTCGGGAAT
AAATGCAACGACGAGGAATCACGAATAACTGGGTCGCTGCATTAATTCTTTATCCAGGCCGACTTATCCGTGG
AGAATGGCGGTCGAGTGGCGAGCAACGAGAAGAATCAACATCATCCATCCGGCACACTCTCTCTATAATTATC
GCGATATCGAAGGAAGGGAAAAAGCGAGGAGTAGAATTTCGAGTTGCAAAGAGATAATATTATATCCCGTTT
CCCTTCCTCCTCCTATTAACTTTCAATCTCATTGCGAACCTCCAGTTTCCTCCAATTTTATATAATATTTTGAAAAG
AGGCGGCTATGCAGAGAATAGAAAAGAAAAAAAAAAGAAAAAGAAGAAATCAGAAGCGCATGCAAACGCGT
TTACCAATGAGCGGGCCGACGTTTCGTCTTCGGTTCGGCCAGCAGGTGGAGCGTGTTTGATGTGAATCGAACA
ACGTCACCCTGTACAAACACCGGGGACCGGGCAGCATATAGGGGCTCGTCCTTTTTTTCCCCGTCGTCGTCGTC
GTCGTTAGCTCGCGAGACT<mark>GGGTGAGGAGGGGGTGATAAGGAGGACGCAAGCGCGGACCTATCGACCAGCT
AACCCTCCGGGTAACACCCTCGCGATCCTCGGCCGAGGCTCACTGCTTCACCGGCAAACTAGTAGGTAGTACA
CGCACGCGACACAG</mark>ATTTCAACTTGGATTGTTTAGTTTCTCCTTCCTCACCTCTCTTATTTCATCCTTTCCTTTCGG
ATTACCTTACGTTCAACTGTTATTCCACAAACTCGGCAAGGGAACGCAACATCTCGTTCGAACGCACATCGTCG
AGCGAAGAAGAGATCACGTCGGCGAGACGACGTCCTTTCCGGTAACCGCCGCGTTTACGAGAAAGGGACAGA
CAGAAGGGGGAGAGCGTAGCGGTGTCCACGCCAACGAAAACACCAACGAAGCGGAATTCTGCGGTTTTCGCG
AGGAGGGCGAGGTCGGTGTTAAACCAGCAAGATTACCGGCCGCGGCGGTTGCGCCAAATACTGTGGTACAGC
TACTTCACCCGGCAACATTTCATTCTCCAGCTTTCGACAGTCCTTCACCTCCAATGTTAACTACCTGAAACGGAG
GTTCTCGTCAGGGGATTTGAGCTCGGAGCTCGACGATGAACCAAGCACAGACTCCGGGCTGGCCTCCATG

Supplementary Figure 2: Synapsin sequence of the honey bee

The promoter and the 5' UTR sequence from the honey bee synapsin (syn) gene. Different colors

indicate the different exons that were fused.