**Title: IL-17A controls CNS autoimmunity by regulating gut microbiota and inducing**

**regulatory T cells**

**Authors:** Shailesh K. Shahi[1], Sudeep Ghimire[1], Samantha N. Jensen[1,2,†], Peter Lehman[1], Nicholas

Borcherding[3], Katherine N. Gibson-Corley[4], Sukirth M. Ganesan[5], Nitin J. Karandikar[1,2], and

Ashutosh K. Mangalam[1,2]*

**Affiliations:**

[1]Department of Pathology, University of Iowa, Iowa City, USA

[2]Graduate Program in Immunology, University of Iowa, Iowa City, USA

[3]Department of Pathology and Immunology, Washington University School of Medicine, Saint

Louis, MO,

[4]Department of Pathology, Microbiology & Immunology; Vanderbilt University Medical Center,

Nashville, TN

[5]Department of Periodontics College of Dentistry, University of Iowa, Iowa City, USA

† Present address- Department of Microbiology and Immunology, Weill Cornell Medicine,

Cornell University, New York, USA

*Corresponding Author

Ashutosh K. Mangalam, Ph.D.
Department of Pathology,
University of Iowa Carver College of Medicine,
25 S. Grand Avenue, 1080A ML
Iowa City, IA  52242
Phone: 1-319-335-8558
Email. ashutosh-mangalam@uiowa.edu
**ORCID  0000-0002-9926-2531**

**Abstract:** A disrupted equilibrium between IL-17A-producing CD4 T-cells (Th17) and $CD4^+CD25^+FoxP3^+$ regulatory T cells (Tregs) play an important role in the pathobiology of Multiple sclerosis (MS). Gut bacteria help in maintaining immune homeostasis by regulating the balance between anti-inflammatory Tregs and pro-inflammatory Th17 cells. Although, both gut bacteria and Tregs can regulate Th17 cells, the impact of IL-17A on gut microbiota and Tregs is unclear. Utilizing HLA-DR3 transgenic mouse model of MS, we show that IL-17A deficiency (HLA-DR3.IL17A$^{-/-}$ mice) expands Treg-inducing gut bacteria such as *Prevotella*, *Parabacteroides,* and *Bacteroides* and consequently Tregs, resulting in a milder disease in an animal model of MS. Notably, IL-17A sufficient DR3 mice develop milder disease on cohousing with IL-17A-deficient mice, highlighting a dominant role for gut microbiota in inducing Treg and reducing disease severity. Further, we observed an enrichment of bacterial-specific Treg promoting short-chain-fatty-acid metabolic pathways and induction of tolerogenic dendritic cells in HLA-DR3.IL17A$^{-/-}$ mice. Thus, our study shows a novel role of IL-17A in immune homeostasis and inflammation through regulation of the gut microbiota-Treg axis which can be used for the development of gut bacteria as therapeutics for MS.

**Significance of our work**

IL-17A a pro-inflammatory cytokine, is linked with pathobiology of multiple inflammatory diseases including multiple sclerosis (MS) and regulated by both gut microbiota and regulatory CD4 T cells (Tregs). However, the importance of IL-17A in the regulation of gut microbiota and Treg is unknown. Here we show that IL-17A can regulate Treg and disease phenotype by modulating gut microbiota and provide a novel mechanism by which immune-mediators such as IL-17A impact the gut microbiota to alter immune cell function and ultimately disease outcomes. Transfer of milder disease phenotype from IL-17A deficient mice to IL-17A sufficient mice on cohousing indicate a dominant role of gut microbiota in disease suppression. Thus, our study lays the foundation for future studies to unravel the interplay between immunological responses and the gut microbiota which will result in the development of microbiota-based therapeutics to treat autoimmune diseases.

**Main text:**

Multiple sclerosis (MS) is a chronic, inflammatory, neurodegenerative disease of the central nervous system (CNS), in which CD4 T helper (Th) cells of Th1 and/or Th17 phenotype have been shown to play an important role in pathobiology of the disease. IL-17A is a signature pro-inflammatory cytokine of the CD4 T helper 17 (Th17) subset that has emerged as being critical to the pathobiology of MS (1, 2). IL-17A plays a crucial role in resolving extracellular bacterial and fungal infections, by recruiting immune cells, specifically neutrophils, to the site of infection and inducing pro-inflammatory mediators from immune cells as well as fibroblasts, epithelial cells, and endothelial cells (3). However, an aberrant IL-17A response can lead to unresolved inflammation and tissue pathology, as observed in multiple autoimmune diseases including MS (4-7). Thus, although IL-17A is considered a major contributor to the pathogenesis of multiple autoimmune diseases, its precise role in the pathobiology of MS is still being explored. Th17 cells have been shown to be regulated by $CD4^+CD25^+FoxP3^+$ regulatory T cells (Treg) and balance between Treg/Th17 is shown to shift toward Th17 phenotype in inflammatory diseases such as MS (8). Additionally, gut bacteria help in maintaining immune homeostasis by regulating the balance between $CD4^+CD25^+FoxP3+$ regulatory T cells (Tregs) and pro-inflammatory Th1/Th17 cells (9, 10). The importance of gut microbiota in regulating development of Treg and Th17 is highlighted by studies performed in germ-free mice where absence or reduced Treg and Th17 cells is evident. However, transplant of single bacterium such as segmented filamentous bacterium can restore Th17 (11, 12) and mixture of chloroform resistant *Clostridium* (13) or *Bacteroides fragilis* can induce/restore Treg population in mice (14). These studies highlight an important role of gut microbiota in regulation of Th17 and Treg population and their role in the pathobiology of MS/EAE. However, whether IL-17A can regulate gut microbiota and/or Treg cells is unknown.

In addition to IL-17A, HLA class II genes, specifically HLA-DR2 and -DR3 alleles, are linked as the strongest genetic factors predisposing individuals to MS (15-22). Evidence suggests MS-associated HLA-DR and –DQ alleles regulate disease outcome and severity by modulating pro- and anti-inflammatory cytokine networks (23). Specifically, myelin antigen processed and presented by susceptible HLA-class II molecules such as HLA-DR2, -DR3, or –DR4, can differentiate into Th1 or Th17 secreting cells which can traffic to the CNS and cause neuroinflammation and demyelination, a characteristic of MS (23-27). Thus, HLA class II

transgenic mice have helped in validating the significance of HLA-class II molecules in the

pathogenesis of MS (24-28). Additionally, we showed that loss of IFNγ was associated with

increased IL-17A levels and severe disease in an HLA-class II transgenic mouse model of

experimental autoimmune encephalomyelitis (EAE) (29). Collectively these data suggest that IL-

17A might be responsible for severe disease in the absence of IFNγ, however it is unclear whether

IL-17A is essential for disease induction or severity or both.

To elucidate the role of IL-17A in the pathogenesis of MS, we generated HLA-DR3 (DR3) class

II transgenic mice lacking IL-17A (DR3.IL17A$^{-/-}$) and assessed the outcome on EAE pathogenesis.

Mice that express HLA-DR3 were chosen over other EAE models as our prior study suggested a

role of IL-17A in disease severity in the absence of IFNγ (30). Unexpectedly, DR3.IL17A$^{-/-}$

transgenic mice were not resistant to EAE, yet developed milder disease compared to IL-17A-

sufficient HLA-DR3 transgenic mice. In the absence of IL-17A, interferon-I related gene

expression was increased, and CD4$^{+}$CD25$^{+}$FoxP3$^{+}$ Treg were expanded, in part due to enhanced

growth of gut bacteria that induce Treg proliferation. The importance of these gut bacteria in

modulating IL-17A-mediated disease was confirmed by fecal transplant/cohousing studies. IL-

17A-sufficient mice that received fecal transplants from, or that were cohoused with, IL-17A-

deficient mice had lower EAE severity relative to IL-17A deficient mice. Bacterial metabolic

pathways related to short-chain fatty acid (SCFA) production were enriched in DR3.IL17A$^{-/-}$

transgenic mice. Also, IL10 production from IL-17A deficient mice was higher for both splenic

and bone marrow derived dendritic cells suggesting that IL-17A deficiency induces an anti-

inflammatory environment. Thus, our study demonstrates that IL-17A regulates Treg levels by

modulating the gut microbiota, which alters immune homeostasis and inflammation to promote

EAE.

**Results**

**IL-17A deficient HLA-class-DR3 transgenic mice develop milder EAE disease**

In the HLA class-II transgenic model of EAE, mice lacking IFNγ develop severe brain-specific

disease characterized by elevated levels of IL-17A (29). Thus, we hypothesized that in the absence

of IL-17A, mice would be resistant to EAE. To investigate the role of IL-17A in EAE, we first

generated HLA-DR3 mice as described in Materials and Methods. All transgenic mice developed

normally with no obvious signs of pathology and analysis of splenocytes confirmed these mice

lacked IL-17A-expressing CD4$^+$ T cells (Fig. S1).

Interestingly, both DR3 and DR3.IL17A$^{-/-}$ transgenic mice develop EAE disease with similar onset

of disease (Fig. 1B). However, DR3.IL17A$^{-/-}$ transgenic mice developed milder disease over time

and had a significantly lower cumulative clinical EAE ($\pm$) as well as peak EAE score ($\pm$) compared

to DR3 mice (Fig. 1B, 1C, 1D). Histological analysis of the brain and spinal cord tissues

demonstrated milder inflammation and demyelination in DR3.IL17A$^{-/-}$ transgenic mice, consistent

with decreased inflammatory cell infiltration in CNS (Fig. 1E, 1F, 1G, 1H). Thus, our data indicate

that IL-17A was not required for induction phase of disease. However, the milder disease that

occured in the absence of IL-17A over time suggests that it may play a role in disease progression.

To test this, we depleted IL-17A in DR3 transgenic mice by treating them with an anti-IL-17A

antibody before induction of EAE (Fig. 2). We treated one cohort of mice with the anti-IL-17A

antibody prior to immunization with PLP$_{91-110}$, on days -19, -14, -11, -8, and -4 (Fig. 2A).

Compared to the control group. DR3 transgenic mice pre-treated with anti-IL-17A still developed

EAE however showed a milder disease course which was similar to EAE oberved in DR3.IL17A$^{-}$

$^{/-}$ transgenic mice (Fig. 2B, 2C). Thus, the development of milder disease in DR3 mice in the

absence of IL-17A (genetic deficiency as well as  IL-17A neutralization)  suggests IL-17A is

redundent for the induction of disease but might be crucial for disease progression.

To determine if the effect of IL-17A on EAE disease was universal or limited to the HLA class II

transgenic mouse model, we assessed the effects of depleting IL-17A in wild-type C57BL/6J (B6)

mice. In this mouse strain, EAE was induced with MOG$_{35-55}$ as described previously (31). B6 mice

treated with anti-IL-17A antibody after the onset of EAE (i.e., at days 5, 8, 11, 14, 17, 20, 23, 26,

29, and 32 post disease induction) showed significantly lower average clinical scores over time

compared to mice treated with PBS or IgG1a (Fig. S2A, 2B). Similar to DR3 transgenic mice, B6

mice that received pre-treatment (i.e., at days -24,-21,-18,-15,-12,-9,-6, and -3) and post-treatment

(days 5, 8, 11, 14, and 17) of anti-IL-17A antibody still developed EAE but exhibited lower disease

severity (Fig. S2C, 2D). These data indicated that the IL-17A is redundant for disease induction

both in PLP-DR3-EAE and MOG-B6-EAE models. to exacerbate EAE progression occurs in both

HLA-DR3 transgenic and wild-type B6 mice, and is not limited to one model of disease.

Even though IL-17A is considered as an important cytokine for the development of EAE (32), our

results suggest that IL-17A is more crucial for disease progression. Further, the development of

EAE disease in the absence of IL-17A, suggests IL-17A is not essnetial for development of EAE

and other inflammatory cytokines might compensate for IL-17A to induce disease. The potential

candidates are IL-17F, GM-CSF and IFNγ as they have been shown to play an important role in

EAE disease.  As IL-17F share 55% homology at the amino acid level with IL-17A, and may

compensate for IL-17A , we analyzed the levels of IL-17F and found that IL-17A deficient mice

have significantly higher levels of the IL-17F compared to IL-17A sufficient DR3 mice (Fig. S3).

However, neutralization of IL-17F had no effect on disease incidence or severity (Fig. S4A, 4B).

Next, we investiagted whether GM-CSF was compensating for EAE induction in IL-17A deficient

mice, as GM-CSF is shown to be a pathogenic cytokine in EAE (33). Although antigen specific

CD4 T cells from DR3.IL17A$^{-/-}$ transgenic mice did not show increased levels of GM-CSF but

neutralization of the GM-CSF in DR3.IL17A$^{-/-}$ resulted in increased average disease score at later

point of the disease (Fig. S4A, S4B). However, the cumulative disease score was similar between

isotype control and anti-GM-CSF treated mice (Fig. S4C). Thus, our data suggested that both GM-

CSF and IL-17F did not compensate for IL-17A in DR3.IL-17A$^{-/-}$ transgenic mice for disease

induction (Fig. S4). To determine the role of IFNγ in disease progression, we generated DR3

transgenic mice in which both IFNγ and IL-17A were deleted (DR3.IL17A$^{-/-}$IFNγ$^{-/-}$). Interestingly,

these mice still developed disease similar to that observed in DR3 transgenic mice (Fig. S5)

suggesting that neither IL-17A nor IFNγ is required for the development of EAE disease. Taken

together, our data indicate that neither IL-17A nor IL-17For GM-CSF, or IFNγ are required for the

development of EAE disease. Thus, our data point toward an inflammatory mediator besides IL-

17A, IL-17F, IFNγ or GM-CSF in the induction of EAE disease.

**IL-17A deficiency enhances proliferation of CD4$^+$CD25$^+$FOXP3$^+$ Treg cells**

The milder course of EAE disease observed in IL-17A-deficient mice suggested the possibility of

an immune-regulatory mechanism at play to dampen the chronic immune response. Studies have

shown that CD4$^+$CD25$^+$Foxp3$^+$ regulatory T cells can regulate autoimmune inflammation in the

body and suppress IL-17A-secreting CD4+ Th17 cells (34, 35). However, few studies have

examined the ability of IL-17A to modulate Treg. Therefore, we analyzed the total number of

CD4$^+$CD25$^+$Foxp3$^+$ regulatory T cells in peripheral blood of DR3 and DR3.IL17A$^{-/-}$ transgenic

mice. We observed that DR3.IL-17A$^{-/-}$ transgenic mice had a significantly higher percentage (Fig.

3A, 3B) and number (Fig. 3C) of CD4$^+$CD25$^+$Foxp3$^+$ regulatory T cells compared to DR3

transgenic mice. To determine whether the increased Treg levels were responsible for milder

disease phenotype in DR3.IL17A$^{-/-}$ transgenic mice, we treated these mice with an anti-CD25

blocking antibody to neutralize the Treg population as shown in Fig 3D (36). CD25 neutralization

in DR3.IL17A$^{-/-}$ transgenic mice resulted in severe disease phenotype similar to DR3 transgenic

mice, indicating that Treg were involved in establishing milder disease phenotype in DR3.IL17A$^-$

$^{/-}$ transgenic mice (Fig. 3E). Notably, CD4$^+$CD25$^+$ Treg that were isolated from the spleen of naïve

DR3.IL17A$^{-/-}$ transgenic mice had significantly higher suppressive ability against antigen specific

CD4$^+$CD25$^-$ effector cells when compared to those of DR3 transgenic mice (Fig. 3F), further

supporting the role of Treg in determining clinical EAE outcomes in DR3.IL17A$^{-/-}$ transgenic

mice. Collectively, our data indicate that deficiency of IL-17A results in a higher frequency and

number of CD4$^+$CD25$^+$ Treg cells with a higher suppressive ability, which likely contributes to a

milder disease phenotype in DR3.IL-17A$^{-/-}$ transgenic mice.

**Loss of IL-17A enriched interferon-I-related genes in the colon of DR3 transgenic mice**

The majority of immune population reside near mucosal surfaces specifically in the gut to protect

large surface area. Trillions of bacteria (microbiota) living in the gut specifically in the colon had

been shown to play an important role in the generation of Treg and Th17 cells. The importance of

gut microbiota in generation of Treg is highlighted by studies performed in germ-free/antibiotic

treated mice, which showed significantly reduced Treg population (37, 38). Interestingly,

colonization of germ-free/antibiotic treated mice with specific commensals can restore Treg

population and function (37-39).

Therefore, to identify whether the genes that might be involved in modulating the disease in IL-

17A deficient transgenic mice, we collected the colonic tissues from both DR3 and DR3.IL17A-/-

transgenic mice before inducing EAE and performed transcriptome analysis (Fig. 4A). We

observed that the colonic tissues from mice of the different genotype (e.g., DR3 and DR3.IL17A$^-$

$^{/-}$) were distinctly separated (Fig. 4B). Upon differential analysis, we identified 123 genes that were

differentially expressed in the colon between the two groups (Fig. 4C, 4D, Table S1). Thirty-three

out of 123 differentially expressed genes were upregulated in DR3.IL17$^{-/-}$ transgenic mice relative

to DR3 transgenic mice. This included genes in the following GO functional categories: *immune*

*function, immune effector process*, and *immune system process* (*Oasl2*, *Oas3*, and *Cfd*), *activation*

*of the immune response, regulation of immune system process*, and *immune response* (*Cfd*),

*signaling* (*Cfd*), and *response to stimulus* (*Oasl2*, *Oas3*, *Cfd*, *Fabp4*, and *Ifit3*). Majority of these

genes belong to type-I interferon category and are shown to be involved in anti-inflammatory

responses (40). The induction of type I interferon genes (e.g., *Oas*-like proteins, *Ifit3*) have been

associated with the stability and function of Treg under stress conditions and promotes induction

of IL-10 expression to enhance their suppressive ability (41, 42). In addition, 90 genes were

downregulated in DR3.IL17$^{-/-}$ transgenic mice relative to DR3 transgenic mice (Fig. 4C, 4D, Table

S1), of which only *Ccl28* was related to immune function and was identified in the GO categories

*immune system process* and *immune response*.  Thus, our observation of enrichment of type-I

interferon genes in the colon of DR3.IL17A$^{-/-}$ transgenic mice suggested that these changes may

contribute to the difference in disease phenotype between DR3 and DR3.IL17A$^{-/-}$ transgenic mice

specifically the ability of type-1 IFN to induce Tregs.

**IL-17A deficiency modulates the composition of the gut microbiome**

The enhancement of Treg and type-I interferon genes in DR3.IL17A$^{-/-}$ transgenic mice relative to

DR3 transgenic mice prompted us to investigate the changes in the composition and function of

the gut microbiota that might impact the immune response. As gut microbiome is known to

modulate the Treg, first, we hypothesized that absence of IL-17A modulated the composition of

the gut microbiome which then can induce Treg population and leading to milder disease

phenotype in DR3.IL17A$^{-/-}$ transgenic mice. Secondly, we hypothesize that the microbiota

enriched in IL-17A deficient mice when transferred to IL-17A sufficient mice should alter the

microbiota of DR3 transgenic mice to enhance Treg in the gut.

To test our first hypothesis i.e., if IL-17A could alter gut microbiota, we collected fecal samples

from both DR3 and DR3.IL17A$^{-/-}$ transgenic mice before cohousing and performed shotgun

metagenomic sequencing to characterize the microbiome in these mice (Fig 5A). The microbial

composition of DR3 transgenic mice clustered distinctly from that of DR3.IL17A$^{-/-}$ transgenic

mice (Fig. S6). Specifically, we observed a significant increase in the average relative abundance

of *Prevotella* sp. MGM1, *Parabacteroides distasonis,* and *Bacteroides sartorii* in DR3.IL17A$^{-/-}$

transgenic mice (Fig. 5B, 5C, 5D, 5E). We did observe that DR3.IL17A$^{-/-}$ transgenic mice had a

significant reduction in the population of *B. vulgatus*, *Lachnospiraceae* bacterium 10_1,

*Lachnospiraceae* bacterium 28_4, *Lachnospiraceae* bacterium COE1, and *Lachnospiraceae*

bacterium A2 (Fig. 5F, 5G, 5H, 5I, 5J). A complete list of bacteria showing trend but not reaching

statistical significance have been provided in Figure S7.

For efficient microbiota transfer we first performed fecal transplantation through oral gavage

followed by cohousing as described previously (43, 44) . We next analyzed the microbiota of

DR3.IL17A$^{-/-}$ and DR3 transgenic mice after fecal transplantation/cohousing and observed that the

microbial composition in DR3 transgenic mice shifted to become more similar to that in

DR3.IL17A$^{-/-}$ transgenic mice (Fig. 5B, Fig. S6). Specifically, *Prevotella* sp. MGM2 (Fig. 5K)

and *E. coli* (Fig. 5L) were significantly increased in DR3 transgenic mice after cohousing with

DR3.IL17A$^{-/-}$ transgenic mice. *Prevotella* species were observed to be significantly higher in

DR3.IL17A$^{-/-}$ transgenic mice compared to DR3 transgenic mice prior to cohousing too and

especially *Prevotella* sp. MGM2 had significantly higher relative abundance in DR3 transgenic

mice after being cohoused with DR3.IL17A$^{-/-}$ transgenic mice. Also, *A. muciniphila*, and *H.*

*typhlonius* were absent in DR3 transgenic mice before they were cohoused but were increased but

not significantly after cohousing with DR3.IL17A$^{-/-}$ transgenic mice (Fig. 5B, Fig. S7). Similarly,

the abundance of *Prevotella* sp. MGM1, *P. distasonis* and *B. sartorii* were increased, although not

to significant levels, in DR3 transgenic mice after being cohoused with DR3.IL17A$^{-/-}$ transgenic

mice (Fig. 5C, 5D, 5E). In contrast, *Lachnospiraceae* bacterial species (Fig. 5G, 5I and 5J) and

*Muribaculaceae* bacterium DSM 10370 (Fig. 5M) were significantly decreased in DR3 transgenic

mice after being cohoused with DR3.IL17A$^{-/-}$ transgenic mice. Thus, the cohousing of DR3

transgenic mice with DR3.IL1-17A$^{-/-}$ transgenic mice transformed microbiota in DR3 mice so that

bacteria that are supportive of Treg were expanded, whereas those that diminished Treg function

were decreased.

**Cohousing DR3 mice with DR3.IL17A$^{-/-}$ transgenic mice transfers a Treg-inducing**

**phenotype**

As deficiency of IL-17A was associated with alterations in the composition of the gut microbiota

and Treg numbers and fecal transplantation plus cohousing resulted in successful colonization by

microbial species from IL-17A deficient mice to IL-17A sufficient mice, we hypothesized that

transferring the gut microbiome from DR3.IL17A$^{-/-}$ transgenic mice to DR3 transgenic mice would

also increase Treg frequency and subsequently reduce EAE severity in recipients. Therefore, we

assessed the levels of CD4$^+$CD25$^+$FoxP3$^+$ Treg cells present in the peripheral blood for both IL-

17A deficient and IL-17A sufficient transgenic mice after fecal transplantation plus cohousing

(Fig. 6A). We observed that CD4$^+$CD25$^+$FoxP3$^+$ T cell populations were significantly increased

in DR3 transgenic mice that received a combination of fecal transplants plus cohousing with

DR3.IL17A$^{-/-}$ transgenic mice (Fig. 6B). In contrast, these Treg populations were similar in

DR3.IL17A$^{-/-}$ transgenic mice that received fecal transplants from DR3 transgenic both before and

after the mice were cohoused (Fig. 6C). These data suggest that microbes transferred from

DR3.IL17A$^{-/-}$ transgenic mice to DR3 transgenic mice, followed by cohousing, were crucial to

enhance the Treg population.

As the transfer of microbiota from DR3.IL17A$^{-/-}$ transgenic mice to DR3 transgenic mice increased

the Treg population (Fig. 6B, 6C), we sought to examine if these changes also affected the outcome

of EAE disease phenotype. DR3 and DR3.IL17A$^{-/-}$ transgenic mice were cohoused for four weeks

and then housed separately while EAE was induced, and the course of disease was assessed. When

housed separately DR3 transgenic mice showed severe disease compared to DR3.IL17A$^{-/-}$

transgenic mice as expected (Fig. 6D, 6E). In contrast, DR3 transgenic mice that were cohoused

with DR3.IL17A$^{-/-}$ transgenic mice showed milder disease compared to DR3 control mice that

were never cohoused (Fig 6D, 6E). However, the disease in DR3.IL-17$^{-/-}$ transgenic mice were

similar pre and post-cohousing with DR3 transgenic mice. Thus, our results confirmed that the

lack of IL-17A alters the composition of the gut microbiota to allow the expansion of bacteria such

as *Prevotella* sp. MGM2, leading to the expansion of the Treg population and ultimately, to milder

EAE disease phenotype.

**Short-chain fatty acid metabolic pathways are transferred from DR3.IL17A$^{-/-}$ transgenic**

**mice to DR3 transgenic mice after cohousing.**

Next to identify the mechanism through which IL-17A influenced gut microbiota can modulate

Treg, we compared bacterial metabolic pathways enriched in DR3.IL-17A$^{-/-}$ mice. We first

assessed short-chain fatty acids (SCFAs), as these have been previously reported to affect the

differentiation of T cells (45) and ameliorate EAE severity (46). The metabolic pathways

associated with pyruvate to propanoate (P108-PWY) and pyruvate fermentation to acetate and

lactate II (PWY-5100) trended higher, although not to significant levels, in DR3.IL17A$^{-/-}$

transgenic mice compared to DR3 transgenic mice before cohousing. P108-PWY was low in

abundance in both DR3 and DR3.IL17A$^{-/-}$ transgenic mice before cohousing but was significantly

higher in both strains after cohousing (Fig. S8A). Likewise, pyruvate fermentation to acetate and

lactate II was lower in DR3 transgenic mice but became significantly enriched after mice were

cohoused with DR3.IL17A$^{-/-}$ transgenic mice (Fig. S8B). The transfer of SCFA pathways from

DR3.IL17A$^{-/-}$ transgenic mice to DR3 transgenic mice, but not vice versa, indicates that the

DR3.IL17A$^{-/-}$ transgenic mice harbor microbes that are responsible for SCFAs production and

these microbes can be transferred to DR3 transgenic mice through co-housing. Thus, our data

suggest the potential role of bacterial SCFAs in inducing Treg to produce milder disease in the

absence of IL-17A.

**323 Dendritic cells in DR3.IL17A$^{-/-}$ transgenic mice produce higher levels of IL-10**

SCFAs such as butyrate and propionate can exhibit strong anti-inflammatory effects by inhibiting

expression of IL-6, IL-12 and IL-23 from dendritic cells (DCs) (47). Butyrate and propionate also

suppress lipopolysaccharide (LPS)-induced DC activation (48). As bacterial metabolic pathway

analysis showed a significant increase in the number of propionate I production pathway, we

hypothesized that higher levels of SCFAs in the gut will alter DCs phenotype and make them

tolerogenic. Therefore, we analyzed the production of the anti-inflammatory cytokine, IL-10, from

DCs in naïve DR3 and DR3.IL17A$^{-/-}$ transgenic mice that were stimulated with LPS. Interestingly,

CD11c$^+$ DCs from naïve DR3.IL17A$^{-/-}$ transgenic mice produced significantly higher levels of IL-

10 compared to DR3 transgenic mice after stimulation with LPS (Fig. 7A). Similarly, bone

marrow-derived DCs from DR3.IL17A$^{-/-}$ transgenic mice produced significantly higher levels of

IL-10 when compared to DR3 transgenic mice (Fig. 7B). Finally, analysis of antigen presentation

capacity showed that bone marrow-derived DCs from DR3.IL17A$^{-/-}$ transgenic mice had a reduced

ability to induce antigen-specific CD4$^+$ T cell proliferation compared to DR3 transgenic mice (Fig.

7C). Thus, our data suggests CD11c+ DC from DR3.IL17A$^{-/-}$ transgenic mice adopted a

tolerogenic phenotype characterized by higher levels of IL-10 with reduced antigen presentation

capacity. Therefore, this phenotype suggests CD11c+ DCs may be involved in limiting disease
severity in DR3.IL17A$^{-/-}$ transgenic mice.

**Discussion**

The ability of IL-17A to influence autoimmune diseases has been well established but the precise
mechanism by which this occurs is poorly understood. In this study, we show that IL-17A is critical
for regulating EAE disease progression through its ability to modulate the gut microbiota and alter
the number and function of Treg. Although not required for disease initiation, our work shows that
IL-17A regulates disease severity by suppressing the Treg population through depletion of Treg-
promoting bacteria. This protective effect of gut microbiota was transferable as cohousing of IL-
17A sufficient DR3 transgenic with DR3.IL17A$^{-/-}$ transgenic mice resulted in gut microbiota
profile, Treg phenotype and milder disease similar to DR3.IL17A$^{-/-}$ transgenic mice. Thus, our data
identify an important pathway through which an immune-mediated cytokine modulates the gut
microbiota to influence the course of inflammatory diseases such as MS.

The development of milder disease in DR3.IL17A$^{-/-}$ transgenic mice suggests induction of
regulatory pathways in the absence of IL-17A. In support of this, we observed an increase in the
CD4$^+$CD25$^+$FoxP3$^+$ Treg population in DR3.IL17A$^{-/-}$ transgenic mice. Further, disease was
restored after Treg depletion using anti-CD25 (PC61) antibody, confirming an important role for
CD4+CD25+ Treg subset in disease regulation. Interestingly, similar observation had been also
reported in GM-CSF deficient mice which are resistant to EAE development but treatment with
anti-CD25 antibody (PC61) resulted in the development of severe disease (49). Thus, these data
suggests that pathogenic cytokine such as IL-17A and GM-CSF might influence EAE disease
progression through downregulation of regulatory cells such as CD4$^+$CD25$^+$ Treg cells.
Development of disease in mice lacking IL-17A or mice receiving CD4$^+$ T cells from *Csf2-/-* (mice
lacking GM-CSF) support our argument that neither IL-17A nor GM-CSF are essential for disease
development in EAE model.

In recent years, gut microbiota has emerged as an important factor in the induction and regulation
of both Treg and Th17 cell subsets (38). However, effect of IL-17A on gut microbiome is poorly
understood and it has been suggested that IL-17A/F can act as modulators of intestinal homeostasis

(50). Therefore, we hypothesized that the absence of IL-17A enables the expansion of certain

species in the gut microbiota that are responsible for the induction of Treg. The presence of distinct

microbiota profiles between DR3 and DR3.IL17A$^{-/-}$ transgenic mice strengthen our hypothesis.

Specifically, we observed an expansion of gut bacteria that are known to promote the Treg

population, including *Prevotella* species, *Parabacteroides distasonis* and *Bacteroides species* (9,

51-53). Of these species, *Prevotella* sps. MGM2 successfully transferred to the IL-17A sufficient

mice from IL-17A deficient mice in high numbers. At the same time, the abundances of none of

the microbial species were affected significantly in IL-17A deficient mice after cohousing with

IL-17A sufficient mice suggesting that the microbial ecosystem of DR3.IL17A$^{-/-}$ transgenic mice

prior to cohousing was at higher stable state compared to DR3 transgenic mice. Also, the microbial

community of DR3 transgenic mice which clustered separately from DR3.IL17A$^{-/-}$ transgenic mice

before fecal transplant and cohousing were clustering with DR3.IL17A$^{-/-}$ transgenic mice after

fecal transplant and cohousing (Fig. S6) indicating that microbial composition of DR3.IL17A$^{-/-}$

transgenic mice were resilient to change compared to DR3 transgenic mice. Ecologically, the

degree of resistance of the microbiota against incoming microbial species is crucial in determining

health and disease (54). The lower stability and the loss of resistance of the gut microbiota in DR3

transgenic mice was ultimately reflected by exacerbated disease severity, which was reduced upon

cohousing with DR3.IL17A$^{-/-}$ transgenic mice, suggesting that the gut microbiota of DR3

transgenic mice was re-configured to restore its protective functions. In particular, cohousing IL-

17A-deficient mice with IL-17A sufficient mice resulted in the transfer of crucial Treg-stimulating

bacteria, such as *Prevotella* species into IL-17A sufficient DR3 transgenic mice. Indeed,

introducing *Prevotella* species can reduce the severity of EAE due to changes in the gut microbiota

that ultimately induce Treg (51, 52). Taken together, our study demonstrates gut microbiota

dependent internal mechanism to maintain intestinal homeostasis and Treg function and

perturbation of the same can promote inflammation and CNS autoimmunity. Suppression of

disease in IL-17A sufficient mice on cohousing with IL-17A deficient mice strongly suggest that

Treg promoting gut microbiota is dominant over disease promoting bacteria and thus have strong

clinical translation potential. However, the microbial community network and the interplay

between the microbes in the community to exacerbate or protect from EAE warrants deeper

exploration.

SCFAs are produced by the gut microbiota through anaerobic fermentation of polysaccharides and

are involved in enhancing IL-10 production and are crucial for maintaining gut homeostasis (55).

Oral administration of a lactate-producing bacterium *P. acidilactici* suppressed EAE by inducing

IL-10-producing Treg (56). In addition, SCFAs maintain the integrity of the gut barrier (57) and

blood-brain barrier (BBB) in the CNS (58, 59). Correspondingly, germ-free mice have reduced

expression of tight junction proteins, such as occludin and claudin, in the BBB, an effect that

increases with age. Recolonization of these mice with diverse microbiota increases the integrity of

both gut and the BBB (60). We have previously shown that mice with EAE have leaky gut and

BBB, however treatment with *Prevotella* restored both gut and blood brain barrier (51). The lower

abundance of these SCFAs pathways in DR3 transgenic mice compared to DR3.IL17A$^{-/-}$

transgenic mice indicate that in the presence of IL-17A, there is a decreased abundance of

microbial populations with the capability to produce SCFAs. However, cohousing of DR3

transgenic mice with DR3.IL17$^{-/-}$ transgenic mice restored microbial SCFA pathways involved in

propionate, acetate, and lactate production. Propionate had been shown to decrease the

differentiation of Th17 cells and enhance the differentiation of Treg (45, 61). Also, the significant

rise in pyruvate fermentation to propionate I (P108-PWY) and pyruvate fermentation to acetate

and lactate II (PWY-5100) in DR3 transgenic mice after being cohoused with DR3.IL17A$^{-/-}$

transgenic mice suggests that the transfer of microbial metabolic pathways can alter immune cell

functions and consequently disease outcome. Thus, absence of IL-17A resulted in the enrichment

of gut bacteria with ability to produce SCFA which then can modulate disease through induction

of Treg.

The majority of Treg that are induced by gut bacteria are found in the colon. Therefore, we

performed gene expression analysis of the colon of IL-17A-sufficient and -deficient DR3

transgenic mice to determine the differentially expressed genes which might be important for

reduced severity of disease phenotype in IL-17A deficient mice. Interestingly, IL-17A-deficient

mice, which exhibit an enhanced Treg population and altered gut microbiota composition, showed

increased expression of type I interferon genes.  Genes belonging to type-I interferon categories

are indicated as important mediators of anti-inflammatory responses (40). Indeed, type I interferon

genes are previously known to promote the suppressive ability of Treg (41, 42). Although IL-17A

deficiency had no effect on the Treg population in a previous study that utilized both IL-17A and

IL-17F (50), we report that the levels of CD4$^+$CD25$^+$FoxP3$^+$ T cell populations were significantly

higher in the DR3.IL17A$^{-/-}$ transgenic mice compared to DR3 transgenic mice. Thus, the induction

of Treg in the colon in the absence of IL-17A is likely critical for regulating inflammation in the

context of EAE.

Next, the induction of Treg in IL-17A deficient mice was concurrent with the increased production

of IL-10 from splenic dendritic cells in naïve, immunized DR3.IL17A$^{-/-}$ transgenic mice. It

suggested that the tolerogenic DCs are induced in the absence of IL-17A and DC-derived IL-10 is

essential to ameliorate EAE severity (62). The reduced antigen presentation capacity of DCs from

DR3.IL17A$^{-/-}$ transgenic mice further support a role of tolerogenic DCs in mice lacking IL-17A.

SCFAs such as propionate and butyrate have been demonstrated as potent inducers of tolerogenic

DCs previously (48). Thus, it is possible that SCFAs produced by the gut microbiota in the current

study could have induced tolerogenic DCs in IL-17A deficient mice, however, details of how the

gut microbiota affect DC proliferation and prompt IL-10 production are only beginning to be

investigated.

Finally, we observed that IL-17A significantly affects disease progression without affecting the

onset of disease. This is consistent with other studies such as Haak et al., who showed that either

loss of IL-17A or overexpression of IL-17A in CNS has a major impact on disease development

(63). It is possible that other Th17 (IL-17F and GM-CSF) or even Th1 cytokine (IFNγ) can

compensate for IL-17A loss for disease development. However, in our study, mice that were

treated with blocking antibodies against IL-17F or GM-CSF still developed disease. IL-17F is a

member of the IL-17 family of cytokines, a number of studies have investigated its role in EAE,

however the majority of studies have concluded that IL-17F is either redundant or don't play a role

in EAE model (63, 64). IL-17F deficient mice treated with anti-IL-17A blocking antibodies were

susceptible to EAE (63). The role of GM-CSF in EAE and MS is still matter of debate, with some

studies suggesting it is essential for EAE disease as GM-CSF deficient mice are resistant to EAE

development (65) whereas others have shown that GM-CSF is not essential for EAE development

(33, 66). Interestingly, Duncker et al., observed that although GM-CSF is required for disease

progression, it was dispensable for disease induction phase (33). Development of severe disease

in DR3 mice lacking both IL-17A and IFNγ ruled out a role of IFNγ in disease induction in the

absence of IL-17A. Thus, our data suggest that IL-17A together with IL-17F, GM-CSF or IFNγ,

is dispensable for disease induction phase as DR3.IL-17A$^{-/-}$ transgenic mice still develop disease,

although a milder course. Future studies will be necessary to determine the mechanisms driving

disease in the absence of IL-17A.

Our results have strong clinical relevance as modulation of the gut microbiota are being considered

as potential therapeutics to treat diseases ranging from inflammatory to neurological diseases. As

MS is an inflammatory disease characterized by gut dysbiosis, combination therapies targeted to

reduce inflammation as well as correct gut dysbiosis might results in better therapeutic outcome

than either one alone. Although IL-17A had been linked with the pathobiology of MS, two anti-

IL17A clinical trials in MS (NCT01874340 and NCT01433250) were terminated early. In that

regard, while anti-IL-17A therapy alone did not prove as effective as current MS drugs,

combination of anti-IL-17A and microbiota modulation might provide better therapeutic efficacy.

Additionally, Treg induction is also being tried as potential treatment option in MS (67).  Our

cohousing experiment showing ability of gut microbiota to colonize, induce Tregs and suppress

disease in susceptible mice suggests that gut microbiota-based induction of host immunity might

have added advantage as it will not only induce Tregs but can correct gut dysbiosis too. Future

studies testing gut commensal with Treg inducing capabilities in combination with anti-IL-17A

are warranted to test whether a combination of anti-inflammatory plus gut commensal modulation

might provide added benefits.

In summary, our study highlights an important pathway through which IL-17A regulates

inflammation, specifically by modulating the gut microbiota to suppress Treg expansion. Most

notably, our data show that the Treg inducing gut microbiota from DR3.IL17A$^{-/-}$ can transfer

disease protection to DR3 transgenic mice to reduce disease severity and progression. In the

absence of IL-17A, the gut microbiota appears stable and resilient to changes in its composition

compared to IL-17A sufficient mice. Thus, our study illuminates a novel mechanism by which

immune-mediators such as cytokines impact the gut microbiota to alter immune cell function and

ultimately disease outcomes and lays the foundation for future studies that unravel the interplay

between immunological responses and the gut microbiota provides long-term implications for the

development of microbiota-based therapeutics to treat autoimmune diseases.

# References

1. T. Korn, E. Bettelli, M. Oukka, V. K. Kuchroo, IL-17 and Th17 Cells. *Annual review of immunology* **27**, 485-517 (2009).

2. J. Milovanovic *et al.*, Interleukin-17 in Chronic Inflammatory Neurological Diseases. *Front Immunol* **11**, 947 (2020).

3. M. M. Curtis, S. S. Way, Interleukin-17 in host defence against bacterial, mycobacterial and fungal pathogens. *Immunology* **126**, 177-185 (2009).

4. W. P. Chong *et al.*, The Cytokine IL-17A Limits Th17 Pathogenicity via a Negative Feedback Loop Driven by Autocrine Induction of IL-24. *Immunity* **53**, 384-397 e385 (2020).

5. P. S. Biswas, IL-17 in Renal Immunity and Autoimmunity. *J Immunol* **201**, 3153-3159 (2018).

6. Y. Iwakura, S. Nakae, S. Saijo, H. Ishigame, The roles of IL-17A in inflammatory immune responses and host defense against pathogens. *Immunological reviews* **226**, 57-79 (2008).

7. M. J. McGeachy, D. J. Cua, S. L. Gaffen, The IL-17 Family of Cytokines in Health and Disease. *Immunity* **50**, 892-906 (2019).

8. B. M. Segal, Stage-specific immune dysregulation in multiple sclerosis. *J Interferon Cytokine Res* **34**, 633-640 (2014).

9. K. M. Nutsch, C. S. Hsieh, T cell tolerance and immunity to commensal bacteria. *Curr Opin Immunol* **24**, 385-391 (2012).

10. C. T. Weaver, R. D. Hatton, Interplay between the TH17 and TReg cell lineages: a (co-)evolutionary perspective. *Nat Rev Immunol* **9**, 883-889 (2009).

11. Y. Yang *et al.*, Focused specificity of intestinal TH17 cells towards commensal bacterial antigens. *Nature* **510**, 152-156 (2014).

12. Ivanov, II *et al.*, Induction of intestinal Th17 cells by segmented filamentous bacteria. *Cell* **139**, 485-498 (2009).

13. K. Atarashi *et al.*, Treg induction by a rationally selected mixture of Clostridia strains from the human microbiota. *Nature* **500**, 232-236 (2013).

14. K. M. Telesford *et al.*, A commensal symbiotic factor derived from Bacteroides fragilis promotes human CD39(+)Foxp3(+) T cells and Treg function. *Gut Microbes* **6**, 234-242 (2015).

15. C. Alvarado-de la Barrera *et al.*, HLA class II genotypes in Mexican Mestizos with familial and nonfamilial multiple sclerosis. *Neurology* **55**, 1897-1900 (2000).

16. L. F. Barcellos *et al.*, HLA-DR2 dose effect on susceptibility to multiple sclerosis and influence on disease course. *American journal of human genetics* **72**, 710-716 (2003).

17. G. C. DeLuca *et al.*, An extremes of outcome strategy provides evidence that multiple sclerosis severity is determined by alleles at the HLA-DRB1 locus. *Proceedings of the National Academy of Sciences of the United States of America* **104**, 20896-20901 (2007).

18. D. A. Dyment *et al.*, Complex interactions among MHC haplotypes in multiple sclerosis: susceptibility and resistance. *Human molecular genetics* **14**, 2019-2026 (2005).

19. J. R. Oksenberg *et al.*, Mapping multiple sclerosis susceptibility to the HLA-DR locus in African Americans. *American journal of human genetics* **74**, 160-167 (2004).

20. O. Olerup, J. Hillert, HLA class II-associated genetic susceptibility in multiple sclerosis: a critical evaluation. *Tissue antigens* **38**, 1-15 (1991).

21. S. V. Ramagopalan *et al.*, Analysis of 45 candidate genes for disease modifying activity in multiple sclerosis. *J Neurol* **255**, 1215-1219 (2008).

22. S. V. Ramagopalan *et al.*, The inheritance of resistance alleles in multiple sclerosis. *PLoS genetics* **3**, 1607-1613 (2007).

23. A. K. Mangalam, V. Taneja, C. S. David, HLA class II molecules influence susceptibility versus protection in inflammatory diseases by determining the cytokine profile. *Journal of immunology* **190**, 513-518 (2013).

24. N. Kaushansky, A. Ben-Nun, DQB1*06:02-Associated Pathogenic Anti-Myelin Autoimmunity in Multiple Sclerosis-Like Disease: Potential Function of DQB1*06:02 as a Disease-Predisposing Allele. *Front Oncol* **4**, 280 (2014).

25. J. Klehmet *et al.*, T cell epitope spreading to myelin oligodendrocyte glycoprotein in HLA-DR4 transgenic mice during experimental autoimmune encephalomyelitis. *Clin Immunol* **111**, 53-60 (2004).

26. J. A. Quandt *et al.*, Myelin basic protein-specific TCR/HLA-DRB5*01:01 transgenic mice support the etiologic role of DRB5*01:01 in multiple sclerosis. *J Immunol* **189**, 2897-2908 (2012).

27. S. A. Nalawade *et al.*, Aire is not essential for regulating neuroinflammatory disease in mice transgenic for human autoimmune-diseases associated MHC class II genes HLA-DR2b and HLA-DR4. *Cellular immunology* **331**, 38-48 (2018).

28. D. Luckey, D. Bastakoty, A. K. Mangalam, Role of HLA class II genes in susceptibility and resistance to multiple sclerosis: studies using HLA transgenic mice. *J Autoimmun* **37**, 122-128 (2011).

29. A. K. Mangalam *et al.*, Absence of IFN-gamma increases brain pathology in experimental autoimmune encephalomyelitis-susceptible DRB1*0301.DQ8 HLA transgenic mice through secretion of proinflammatory cytokine IL-17 and induction of pathogenic monocytes/microglia into the central nervous system. *J Immunol* **193**, 4859-4870 (2014).

30. A. Mangalam *et al.*, HLA-DQ6 (DQB1*0601)-restricted T cells protect against experimental autoimmune encephalomyelitis in HLA-DR3.DQ6 double-transgenic mice by generating anti-inflammatory IFN-gamma. *J Immunol* **180**, 7747-7756 (2008).

31. S. N. Jensen *et al.*, Isoflavone diet ameliorates experimental autoimmune encephalomyelitis through modulation of gut bacteria depleted in patients with multiple sclerosis. *Sci Adv* **7** (2021).

32. Y. Komiyama *et al.*, IL-17 plays an important role in the development of experimental autoimmune encephalomyelitis. *J Immunol* **177**, 566-573 (2006).

33. P. C. Duncker, J. S. Stoolman, A. K. Huber, B. M. Segal, GM-CSF Promotes Chronic Disability in Experimental Autoimmune Encephalomyelitis by Altering the Composition of Central Nervous System-Infiltrating Cells, but Is Dispensable for Disease Induction. *J Immunol* **200**, 966-973 (2018).

34. Z. Li, D. Li, A. Tsun, B. Li, FOXP3+ regulatory T cells and their functional regulation. *Cell Mol Immunol* **12**, 558-565 (2015).

35. S. Hori, T. Nomura, S. Sakaguchi, Control of regulatory T cell development by the transcription factor Foxp3. *Science* **299**, 1057-1061 (2003).

36. A. D. Christensen, S. Skov, P. H. Kvist, C. Haase, Depletion of regulatory T cells in a hapten-induced inflammation model results in prolonged and increased inflammation driven by T cells. *Clin Exp Immunol* **179**, 485-499 (2015).

37. K. Atarashi *et al.*, Induction of colonic regulatory T cells by indigenous Clostridium species. *Science* **331**, 337-341 (2011).

38. J. L. Round, S. K. Mazmanian, Inducible Foxp3+ regulatory T-cell development by a commensal bacterium of the intestinal microbiota. *Proceedings of the National Academy of Sciences of the United States of America* **107**, 12204-12209 (2010).

39. M. B. Geuking *et al.*, Intestinal bacterial colonization induces mutualistic regulatory T cell responses. *Immunity* **34**, 794-806 (2011).

40. E. N. Benveniste, H. Qin, Type I interferons as anti-inflammatory mediators. *Sci STKE* **2007**, pe70 (2007).

41. C. A. Stewart *et al.*, Interferon-dependent IL-10 production by Tregs limits tumor Th17 inflammation. *J Clin Invest* **123**, 4859-4874 (2013).

42.   L. Zhang, S. Yuan, G. Cheng, B. Guo, Type I IFN promotes IL-10 production from T cells to suppress
Th17 cells and Th17-associated autoimmune inflammation. *PLoS One* **6**, e28432 (2011).
43.   T. L. Montgomery *et al.*, Interactions between host genetics and gut microbiota determine
susceptibility to CNS autoimmunity. *Proceedings of the National Academy of Sciences of the*
*United States of America* **117**, 27516-27527 (2020).
44.   V. K. Ridaura *et al.*, Gut microbiota from twins discordant for obesity modulate metabolism in
mice. *Science* **341**, 1241214 (2013).
45.   A. Haghikia *et al.*, Dietary Fatty Acids Directly Impact Central Nervous System Autoimmunity via
the Small Intestine. *Immunity* **43**, 817-829 (2015).
46.   M. Mizuno, D. Noto, N. Kaga, A. Chiba, S. Miyake, The dual role of short fatty acid chains in the
pathogenesis of autoimmune disease models. *PLoS One* **12**, e0173032 (2017).
47.   C. Nastasi *et al.*, Butyrate and propionate inhibit antigen-specific CD8(+) T cell activation by
suppressing IL-12 production by antigen-presenting cells. *Sci Rep* **7**, 14516 (2017).
48.   M. M. M. Kaisar, L. R. Pelgrom, A. J. van der Ham, M. Yazdanbakhsh, B. Everts, Butyrate Conditions
Human Dendritic Cells to Prime Type 1 Regulatory T Cells via both Histone Deacetylase Inhibition
and G Protein-Coupled Receptor 109A Signaling. *Front Immunol* **8**, 1429 (2017).
49.   D. Ghosh, A. D. Curtis, 2nd, D. S. Wilkinson, M. D. Mannie, Depletion of CD4+ CD25+ regulatory T
cells confers susceptibility to experimental autoimmune encephalomyelitis (EAE) in GM-CSF-
deficient Csf2-/- mice. *Journal of leukocyte biology* **100**, 747-760 (2016).
50.   T. Regen *et al.*, IL-17 controls central nervous system autoimmunity through the intestinal
microbiome. *Sci Immunol* **6** (2021).
51.   A. Mangalam *et al.*, Human Gut-Derived Commensal Bacteria Suppress CNS Inflammatory and
Demyelinating Disease. *Cell Rep* **20**, 1269-1277 (2017).
52.   S. K. Shahi *et al.*, Prevotella histicola, A Human Gut Commensal, Is as Potent as COPAXONE(R) in
an Animal Model of Multiple Sclerosis. *Front Immunol* **10**, 462 (2019).
53.   E. Cekanaviciute *et al.*, Gut bacteria from multiple sclerosis patients modulate human T cells and
exacerbate symptoms in mouse models. *Proc Natl Acad Sci U S A* **114**, 10713-10718 (2017).
54.   C. A. Lozupone, J. I. Stombaugh, J. I. Gordon, J. K. Jansson, R. Knight, Diversity, stability and
resilience of the human gut microbiota. *Nature* **489**, 220-230 (2012).
55.   M. Sun *et al.*, Microbiota-derived short-chain fatty acids promote Th1 cell IL-10 production to
maintain intestinal homeostasis. *Nat Commun* **9**, 3555 (2018).
56.   K. Takata *et al.*, The lactic acid bacterium Pediococcus acidilactici suppresses autoimmune
encephalomyelitis by inducing IL-10-producing regulatory T cells. *PLoS One* **6**, e27644 (2011).
57.   M. Priyadarshini, K. U. Kotlo, P. K. Dudeja, B. T. Layden, Role of Short Chain Fatty Acid Receptors
in Intestinal Physiology and Pathophysiology. *Compr Physiol* **8**, 1091-1115 (2018).
58.   L. Hoyles *et al.*, Microbiome-host systems interactions: protective effects of propionate upon the
blood-brain barrier. *Microbiome* **6**, 55 (2018).
59.   Y. P. Silva, A. Bernardi, R. L. Frozza, The Role of Short-Chain Fatty Acids From Gut Microbiota in
Gut-Brain Communication. *Front Endocrinol (Lausanne)* **11**, 25 (2020).
60.   V. Braniste *et al.*, The gut microbiota influences blood-brain barrier permeability in mice. *Sci Transl*
*Med* **6**, 263ra158 (2014).
61.   A. Duscha *et al.*, Propionic Acid Shapes the Multiple Sclerosis Disease Course by an
Immunomodulatory Mechanism. *Cell* **180**, 1067-1080 e1016 (2020).
62.   V. P. Kashi, S. B. Ortega, N. J. Karandikar, Neuroantigen-specific autoregulatory CD8+ T cells inhibit
autoimmune demyelination through modulation of dendritic cell function. *PLoS One* **9**, e105763
(2014).
63.   S. Haak *et al.*, IL-17A and IL-17F do not contribute vitally to autoimmune neuro-inflammation in
mice. *The Journal of clinical investigation* **119**, 61-69 (2009).

64.   F. Wanke *et al.*, Expression of IL-17F is associated with non-pathogenic Th17 cells. *J Mol Med (Berl)*
**96**, 819-829 (2018).
65.   J. L. McQualter *et al.*, Granulocyte macrophage colony-stimulating factor: a new putative
therapeutic target in multiple sclerosis. *The Journal of experimental medicine* **194**, 873-882 (2001).
66.   E. R. Pierson, J. M. Goverman, GM-CSF is not essential for experimental autoimmune
encephalomyelitis but promotes brain-targeted disease. *JCI Insight* **2**, e92362 (2017).
67.   K. Chwojnicki *et al.*, Administration of CD4(+)CD25(high)CD127(-)FoxP3(+) Regulatory T Cells for
Relapsing-Remitting Multiple Sclerosis: A Phase 1 Study. *BioDrugs* **35**, 47-60 (2021).
68.   S. X. Ge, D. Jung, R. Yao, ShinyGO: a graphical gene-set enrichment tool for animals and plants.
*Bioinformatics* **36**, 2628-2629 (2020).

**Acknowledgments:** We thank the Waldschmidt, Legge, Jabbari, and Lieberman laboratories for
helpful discussion. We thank Jian Zhang and John Harty for reading the manuscript and provinding
their valuable inputs. We also thank Comparative Pathology Laboratory at the University of Iowa,
Department of Pathology for their time and efforts.

**Funding:**  We acknowledge funding from following sources-

National Institutes of Health/NIAID 1R01AI137075  (AKM)

Veteran Affairs Merit Award  1I01CX002212 (AKM)

University of Iowa Environmental Health Sciences Research Center, NIEHS/NIH P30 ES005605

(AKM)

Gift from P. Heppelmann and M. Wacek to (AKM)

Carver Trust Pilot Grant (AKM)

National Institute of Health training grant in Immunology T32AI007485 PI- G. Bishop (SNJ)

**Author Contributions:**

Conceptualization: AKM, SKS, NJK

Formal analysis: SKS, SG, NB, KNGC, SMG

Methodology: AKM, SKS, NB, NJK

Investigation: AKM, SKS, SNJ, SG, NB, PL, KNGC

Visualization: AKM, SKS, SNJ, SG, NB, KNGC, SMG

Funding acquisition: AKM

Project administration: AKM, SKS

Supervision: AKM, NJK

Writing – original draft: AKM, SKS, SG,

Writing – review & editing: AKM, SKS, SG, SMG, NJK

**Competing interests:** AKM is inventor of a technology claiming the use of *Prevotella*

*histicola* for the treatment of autoimmune diseases. The patent for the technology is owned by

Mayo Clinic, who has given exclusive license to Evelo Biosciences. AKM received royalties

from Mayo Clinic (paid by Evelo Biosciences). However, no fund or product from the patent

were used in the present study.

The remaining authors declare that the research was conducted in the absence of any commercial

or financial relationships that could be construed as a potential conflict of interest.

**Data and materials availability:** All data needed to evaluate the conclusions in the manuscript

are present in the manuscript and/or the Supplementary Materials. Shotgun metagenomic

sequencing and RNA-seq data had been uploaded under bio-project PRJNA808406. No custom

codes were used for the data analysis.

**Figures:**

**Fig. 1: IL-17A deficiency ameliorates EAE in DR3 transgenic mice** (**A**) Schematic design of disease course in DR3 mice. HLA-DR3 (n= 8) and HLA-DR3.IL17A$^{-/-}$ (n= 8) mice were immunized with PLP$_{91-110}$ /CFA at day 0 and Pertussis toxin at days 0 and 2 to induce EAE. Mice were monitored for EAE disease progression before euthanizing on day 30 post-immunization. (**B**) Clinical EAE scores from mice in A, acquired over time. (**C**) Cumulative EAE scores from mice in A. **D**) Peak EAE score from mice in A (**E-H**) Representative histological images of tissues from the CNS from mice in A. Data are representative of 3 independent experiments with 4-5 mice per group. Bullet points (B) or bars (C, D) represent the mean and error bars represent the standard error of the mean for each experimental group. *p*-value was determined by multiple unpaired t-test (B), and unpaired t-test with Welch correction (C, D).

**Fig. 2: Anti-IL-17A treatment reduces the severity of EAE in HLA-DR3 mice.** (**A**) Schematic design of anti-IL-17A treatment in mice induced with EAE. 8-10-week-old HLA-DR3 mice were treated with an anti-IL-17A monoclonal antibody before EAE induction with PLP$_{91-110}$/CFA. (**B**) Average clinical EAE scores of mice treated as in A. Bullet points represent the mean and bars represent the standard error of the mean of each experimental group. (**C**) Average cumulative EAE scores of mice in A. Bars represent the mean and standard error of mean in each group. Data are from two independent experiments (n ≥4 mice per group). $p$-value determined by multiple unpaired t-test (B) and unpaired t-test with Welch correction (C).

**Fig. 3: IL-17A deficiency in DR3 transgenic mice results in expansion of CD4+CD25+FOXP3+ Treg cells.** (**A**) Representative flow cytometric plots of peripheral blood mononuclear cells (PBMCs) isolated from 8-10-week-old HLA-DR3 and HLA-DR3.IL17A$^{-/-}$ transgenic mice. Plots were gated on live lymphocytes and singlets. (B) Percentage and (C) total number of CD4+CD25+FoxP3+ regulatory T cells from mice shown in A. (**D**) Schematic diagram of experimental workflow. HLA-DR3 and HLA-DR3.IL17A$^{-/-}$ transgenic mice were treated with anti-CD25 to neutralize Treg prior to immunization with PLP$_{91-110}$ /CFA to induce EAE. Arrowheads represent days in which mice received anti-CD25 treatment. (**E**) Average clinical EAE scores of HLA-DR3 and HLA-DR3.IL17A$^{-/-}$ mice that were treated with anti-CD25 or PBS prior to induction of EAE. (**F**) Counts per minute (CPM) of HLA-DR3 CD4 effector cells and antigen presenting cells cultured with CD4+CD25+ Treg from the spleen of naïve HLA-DR3 or HLA-DR3.IL17A$^{-/-}$ transgenic mice. The data represent two experiments performed at different time points. Bars represent the mean and standard error of the mean from two independent experiments. n =4-8 mice per group. *p*-value was determined by unpaired t-test with Welch correction (B, C, F) and two-way ANOVA for EAE clinical scores (E).

Fig. 4: DR3 and DR3.IL17A$^{-/-}$ transgenic mice exhibit distinct gene expression patterns in the colon. Colonic tissue was harvested from DR3 (n=3) or DR3.IL-17A$^{-/-}$ transgenic mice (n=3) and subjected to RNA-seq analysis. (A) Schematic diagram of workflow. B) Multi-dimensional scaling plot showing the separation of HLA-DR3 and HLA-DR3.IL17A$^{-/-}$ colonic samples. Each letter represents an RNA-Seq sample. Sample groups are indicated by different colors, as indicated in the legend. (C) Volcano plot displaying the pattern of gene expression of DR3.IL17A$^{-/-}$ mice relative to DR3 mice. Genes that are significantly differentially expressed, with a Log$_2$ fold change of >1 and an FDR-corrected p-value <0.05, are highlighted in red, blue lines represents the boundary for Log$_2$ fold change and the adjusted p-value cutoff. D) Heatmap depicting all differentially expressed genes between HLA-DR3 and HLA-DR3.IL17A$^{-/-}$ colonic samples. Labelled genes are of immunity-related categories as identified using ShinyGO webserver (68).

**Fig. 5: Taxonomic characterization of the gut microbiome in DR3 and DR3.IL17A$^{-/-}$ mice).**
Fecal samples from DR3 and DR3.IL-17A$^{-/-}$ transgenic mice before cohousing (BCH), or after that one time fecal transferred (orally gavaged) and cohoused (ACH) were obtained and subjected to shotgun metagenomic sequencing. (**A**) Schematic diagram representing the experimental flow. B) Heatmap depicting bacterial species with > 1% average relative abundance in at least one group (n=5). (**C-N**). Statistically significantly abundant bacterial taxa that were statistically different in abundance in at least one of the comparisons (e.g., DR3-BCH to DR3IL17A$^{-/-}$-BCH, DR3-BCH to DR3-ACH, and DR3IL17A$^{-/-}$-BCH to DR3IL17A$^{-/-}$-ACH). Kruskal Wallis test followed by Dunn's *posthoc* analysis was performed to identify significant differences between groups. "*" and "**" represent significance at $p<0.05$ and $p<0.01$ respectively.

**Fig. 6: Treg-inducing bacteria from DR3.IL17A<sup>-/-</sup> mice reduce disease severity when transferred to DR3 mice.** (**A**) Schematic of experimental design showing experimental outline. 8-10-week-old female DR3 transgenic mice were cohoused for four weeks with DR3.IL17A$^{-/-}$ transgenic mice after one-time fecal transferred (orally gavaged) from each other mice strains. Peripheral blood mononuclear cells (PBMCs) were isolated from blood that was collected by retro-orbital bleeding, and regulatory T cells were assessed by flow cytometric analysis. Percentages of CD4$^+$CD25$^+$FoxP3$^+$ T cell populations in (**B**) DR3 transgenic mice before (BCH) and after (ACH) being cohoused with DR3.IL17A$^{-/-}$ transgenic mice and in (**C**) DR3.IL17A$^{-/-}$ transgenic mice before (BCH) and after (ACH) being cohoused with DR3 transgenic mice. (**D**) Average clinical EAE scores of DR3 transgenic mice (non-cohoused control and after cohoused with DR3.IL17A$^{-/-}$ transgenic mice) and DR3.IL17A$^{-/-}$ transgenic mice (non-cohoused control and after cohoused with DR3 transgenic mice) that were immunized with PLP$_{91-110}$/CFA and followed for 30 days. (**E**) Average cumulative EAE scores of mice in C. Data represent the mean and standard error of the mean for each experimental group from two independent experiments: n ≥4 mice per group. *p*-value was determined by unpaired t-test with Welch correction (A, B, D), two-way ANOVA for clinical EAE scores (C).

**Fig. 7**: **Dendritic cells adopt a tolerogenic phenotype in the absence of IL-17A**. (**A**)
Quantification of IL-10 cytokine production in naïve dendritic cells from DR3 and DR3.IL17A[-/-]
transgenic mice following stimulation with LPS. Splenocytes were harvested from naïve 8-10-
week-old DR3 and DR3.IL17A[-/-] transgenic mice and CD11c+ dendritic cells were isolated using
BD™ IMag Particles. Purified CD11c+ cells were activated with LPS for 24 hours and IL-10 was
measured by ELISA. Media without LPS was used as control. Bars represent the mean and
standard error of the mean from each experimental group; n ≥3 mice per group. p-value was
determined by Mann-Whitney test. (**B**) Quantification of IL-10 cytokine production from bone
marrow-derived DCs (BMDCs). Bone marrow was collected from naïve DR3 or DR3.IL17A[-/-]
transgenic mice and cultured in the presence of GM-CSF and IL4 for 4 days. BMDCs were then
cultured with or without LPS for 24 hours and IL-10 was measured by ELISA. (**C**) Antigen-
specific proliferation of CD4+ T cells from DR3 and DR3.IL17A[-/-] transgenic mice. 8-10-week-
old DR3 and DR3.IL17A[-/-] transgenic mice were immunized with PLP$_{91-110}$/CFA. After 7 days,
mice were euthanized, and splenocytes were harvested to isolate CD4[+] T cells using BD™ IMag
Particles. CD4+ T cells and antigen presenting cells were cultured together in presence of PLP$_{91-}$
$_{110}$ for 5 days, and thymidine was added in the last 12 hours of culture. CD4[+] T cells from mice of
either genotype were cultured in the presence of antigen presenting cells. Bars represent the mean
and standard error of the mean from each experimental group after subtracting the baseline medium
values. n =3 mice in each group; *p*-value determined by unpaired t-test with Welch correction (A,
C).

**Supplementary Materials**

**Materials and methods**

**Study design**

The study utilized female HLA-DR3 and HLA-DR3.IL17A$^{-/-}$ transgenic mice to investigate the

contributions of IL-17A to EAE. Disease was induced by immunizing mice with the PLP$_{91-110}$

peptide. The number of sample sizes are indicated in the figure captions. The disease induction

and scoring of the EAE was performed as described below and elsewhere (*1*). No data, including

outliers, were excluded from the analysis. Experiments were performed at least twice with some

exceptions as indicated in the figure legends.

**Mouse procurement, disease induction, and scoring**

HLA-DR3 (DRA1*0101, DRB1*0301) transgenic mice on the B6/129 background have been

characterized previously (Das et al., 2000; Mangalam et al., 2009). These mice lack endogenous

murine major histocompatibility complex (MHC) class II genes (AE$^{-/-}$) and express alleles

DRA1*0101, DRB1*0301, as described previously (Das et al., 2000; Mangalam et al., 2009). The

HLA-DR3.IL17A$^{-/-}$ mice (referred to as DR3.IL17A$^{-/-}$) that were used in this study were generated

by crossing HLA-DR3 (referred to as DR3) and AE$^{-/-}$.IL17A$^{-/-}$ mice and validated using

genotyping. C57BL/6 mice were also utilized in this study and were purchased from the Jackson

Laboratories (Bar Harbor, ME). Female mice DR3.IL17A$^{-/-}$ and C57BL/6 mice (8-10 weeks of

age) were used in this study. Mice were bred and maintained in the University of Iowa animal

facility in accordance with NIH and institutional guidelines. All experiments were approved by

the Institutional Animal Care and Use Committee at the University of Iowa.

To induce EAE in mice, DR3 and DR3.IL17A$^{-/-}$ transgenic mice (8 to 12 weeks old) were

immunized subcutaneously in both flanks with 50 μg of PLP$_{91-110}$ (YTTGAVRQIFG

DYKTTICGK) (GenScript, Piscataway NJ, USA) that was emulsified in CFA containing 200

μg/mouse *Mycobacterium tuberculosis* H37Ra (Becton, Dickinson and Company, Sparks, MD,

USA). 80 ng of pertussis toxin (PTX) (Sigma Chemicals, St. Louis, MO; 100 ng) was given

intraperitoneal at day 0 and 2 post immunization. C57BL/6 mice were immunized subcutaneously

in both flanks with MOG$_{35-55}$ CFA/PTX, as just described for immunization with PLP (*2*). Mice

were observed daily for disease symptoms and EAE was scored with the following scoring system,

as described previously (*1*): normal, 0; loss of tail tone, 1; hind limb weakness, 2; hind limb

paralysis, 3; hind limb paralysis and forelimb paralysis or weakness, 4; and morbidity/death, 5.

**Pathology**

Brains and spinal cords were collected from DR3, and DR3.IL17A$^{-/-}$ transgenic mice at day 30

post immunization. Histological analysis was performed to assess inflammation and

demyelination, as described previously (*3, 4*). Briefly, DR3 and DR3.IL17A$^{-/-}$ transgenic mice

were perfused with approximately 50 ml of 10% neutral buffered formalin via intracardiac

puncture. The calvaria and spinal columns were collected, and immersion fixed in 10 % neutral

buffered formalin (10% BFA) for 24-48 hours followed by decalcification in EDTA for

approximately 24 hours. Decalcified tissues were cut into 2 mm coronal blocks, embedded in

paraffin and routinely processed. The slide were cut at 4-5 mm thickness and stained with

hematoxylin and eosin. HE-stained slides including the brainstem, cortex, corpus callosum,

cerebellum, hippocampus, and striatum were examined and scored by a board-certified veterinary

pathologist as described previously (*3, 4*).

**Flow cytometry**

Peripheral blood mononuclear cells (PBMCs) were isolated from naïve DR3 and DR3.IL17A$^{-/-}$

transgenic mice using Histopaque-1077 (Millipore Sigma, USA) density centrifugation method as

per instructions from the manufacturer (Sigma-Aldrich, St. Louis, USA). Briefly, 200 µl of blood

was collected in an EDTA-containing Eppendorf tube by retroorbital bleeding. Blood was diluted

with 800 ml of phosphate-buffered saline (PBS; Gibco/Lubioscience) and added to 1 ml of

Histopaque-1077, and then centrifuged at 400 x g for 30 minutes at room temperature, without the

brake. Following centrifugation, the buffy coat was collected into 5 mL tubes and washed with

PBS (400 x g for 10 minutes at 4°C). The cells were washed with FACS buffer stained with

antibodies to detect surface expression of CD4 (GK1.5) and CD25 (PC61) (BD Biosciences,

Franklin Lakes, NJ), whereas intracellular expression of FoxP3$^{+}$ was stained using an anti-

Mouse/Rat FoxP3 (FJK-16s) staining kit (eBiosciences, San Diego, CA). Intracellular staining for

IL17, IFNγ, and GM-CSF were performed using the intracellular fixation permeabilization kit and

anti-mouse IL17 (TC11-18H10.1), IFNγ (XMG1.2), and GM-CSF (MP1-22E9) specific

antibodies from eBioscience™. Cells were also stained with antibodies to detect surface

expression of CD45 (30-F11) and CD4 (clone GK1.5) to gate on the leukocyte population.

**T cell proliferation and cytokine assay**

Mice were euthanized 10 days after immunization with $PLP_{91-110}$ and draining lymph nodes were

removed and challenged *in vitro* with $PLP_{91-110,}$ as described previously (Mangalam et al., 2004

Eur. J. Immunol. 34: 280–290). The results are presented as stimulation indices, which are counts

per minute (cpm) of test sample/cpm of the control). For cytokine analysis, supernatants were

collected from culture 48 h after peptide stimulation, and the concentration of cytokines (IL-17F,

and GM-CSF) were measured by sandwich ELISA using pairs of relevant anti-cytokine

monoclonal antibodies (Pharmingen, San Diego, CA).

**Isolation of $CD4^+CD25^+$ Treg, $CD4^+CD25^-$ T effector, and Dendritic cells**

$CD4^+CD25^+$ T cells isolated from the spleens of DR3 and $DR3.IL17A^{-/-}$ transgenic mice using the

EasySep™ Mouse $CD4^+CD25^+$ regulatory T Cell Isolation Kit II (STEMCELL Technologies,

Vancouver, Canada) according to the manufacturer's protocols. $CD4^+CD25^-$ T effector cells were

isolated from the Treg-depleted fraction of splenocytes using Anti-Mouse CD4 Magnetic Particles

– DM (BD Biosciences, CA, USA). DCs were isolated from total splenocytes using CD11c

microbeads according to the manufacturer's protocols (BD Biosciences, CA, USA). The purity of

specific cell populations was analyzed by flow cytometry; all populations used were >90% pure.

***In vivo* depletion of $CD4^+CD25^+$ Treg cells**

DR3 and $DR3.IL17A^{-/-}$ transgenic mice received two intraperitoneal injections (200 µg/injection)

of monoclonal anti-mouse CD25 (IL-2Ra) antibody (clone: PC-61.5.3; BioXcell) in 200 µl of PBS

per dose on days -5 and -3 before induction of EAE (*5*). Controls received two intraperitoneal

injections of 200 µl of PBS. $CD4^+CD25^+FoxP3^+$ T cell depletion was confirmed by flow

cytometric analysis of PBMCs on the day EAE was induced. For IL-17A depletion, DR3

transgenic mice were injected intraperitoneally with 100 ug/mouse anti-IL17A (Clone 17F3;

BioXcell), IgG1 (Clone MOPC-21; BioXcell) or PBS. IL-17A was blocked in C57BL/6 mice by

administering 100 ug/mouse anti-IL-17A, IgG1 or PBS by intraperitoneally injection every three

days following induction of EAE, starting at day 5 through day 32 (e.g., days -24, -21, -18, -15, -

12, -9, -6 and -3 pre EAE induction and post EAE induction (at day 5, 8, 11, 14, and 17). IL-17F

or GM-CSF cytokine inhibition was performed *in vivo* in DR3.IL17A$^{-/-}$ transgenic mice by

intraperitoneally injection of 100µg/mouse anti-IL-17F (clone; MM17F-8F5; BioXcell) or

100µg/mouse of anti-GM-CSF (clone; MP1-22E9; BioXcell) post induction of EAE on days 5,

10, and 14. Control DR3.IL17A$^{-/-}$ transgenic mice received 100µg/mouse IgG1 (Clone MOPC-21;

BioXcell).

**Treg suppression assay**

Spleens from 8-10-week-old DR3 and DR3.IL17A$^{-/-}$ transgenic mice were harvested for CD4$^{+}$ T

cells by BD™ IMag Particles as per manufacturer instruction. T cell-depleted splenocytes were

used as antigen presenting cells after UV irradiation. CD4$^{+}$CD25$^{+}$ T cells were isolated from

splenocytes of DR3 and DR3.IL17A$^{-/-}$ transgenic mice using EasySep™ Mouse CD4$^{+}$CD25$^{+}$

regulatory T cell isolation kit II (STEMCELL Technologies, Vancouver, Canada) according to the

manufacturer's protocols. Treg suppression assays were performed as described previously (*6*).

**Co-housing of DR3 and DR3.IL17A$^{-/-}$ transgenic mice**

For fecal transplantation, DR3.IL17A$^{-/-}$ transgenic mice were recipients of fecal transplants from

DR3 transgenic mice through oral gavage, and conversely, DR3 transgenic mice were recipients

of fecal transplants from DR3.IL17A$^{-/-}$ transgenic mice through oral gavage. Five days after oral

gavage, DR3 transgenic mice were cohoused with DR3.IL17A$^{-/-}$ transgenic mice. Fecal samples

were collected before and after co-housing for shotgun metagenomic sequencing.

**Microbiota DNA isolation, sequencing, and analysis**

Five female mice of each genotype, DR3 and DR3.IL17A$^{-/-}$, were assessed for the composition of

their gut microbiota before and after cohousing, resulting in a total of 20 fecal samples. DNA was

extracted using PowerSoil DNA Isolation Kit (Qiagen) following the manufacturer's protocol.

Each metagenomic DNA was quantified and sequenced using an Illumina MiSeq resulting in more

than 55 Gb of raw paired-end reads from 20 samples after shotgun metagenomic sequencing. The

raw paired-end metagenomes were quality trimmed using Trim Galore version 0.5.0 (*7*), followed

by host read filtering by mapping to the mouse genome (GRCm39) using Bowtie2 (*8*), with default

parameters resulting in 49.43 Gb of high-quality data. The quality trimmed and filtered forward

and reverse reads were taxonomically characterized using MetaPhlAN 3.0 (*9*) and

mpa_v30_CHOCOPhlAN_201901 database with default parameter settings. The obtained relative

percentages of the microbial taxa were compared between the groups using Kruskal-Wallis test

followed by Dunn's multiple t-test in GraphPad Prism unless stated otherwise. The filtered and

trimmed sequences were further assembled using MegaHit (*10*). After assembly, Prokka was used

to predict ORF based protein coding genes and assign functions to the predicted genes (Seemann

T, 2014). BBMap aligner was used to produce quantifiable gene counts in the FPKM format

(fragments per kilobase of gene per million) (https://jgi.doe.gov/data-and- tools/bbtools/).

InterProScan was utilized for assignment for KEGG pathways and GO processes

(https://www.ebi.ac.uk/interpro/search/sequence/).  FPKM values of every protein/gene assigned

to each of the KEGG pathway or GO process were derived from mapping the gene to the FPKM

list of the genes identified in the previous step. Relative abundances of genes and pathways were

used for the downstream statistical analysis. Statistical significance between the groups were

determined using non-parametric Wilcoxon signed rank test, and data visualized using GraphPad

Prism.

**169    In vivo depletion of IL-17A in C57BL/6 mice**

C57BL/6 mice were placed in three groups that received antibody injections prior to EAE

induction (day -24, -21, -18, -15, -12, -9, -6, and -3): the first group of mice received intraperitoneal

injections (200 µg/injection) of monoclonal anti-mouse IL-17A (clone 17F3; BioXCell) in 200µl

PBS per dose; the second group of mice received 200 µg/injection of isotype control (mouse IgG1)

in 200µl PBS per dose; and the third group of mice received 200µl PBS to serve as vehicle control.

Similarly, three groups of C57BL/6 mice were treated with either monoclonal anti-mouse IL-17A

(200 µg/injection), isotype control mouse IgG1 (200 µg/injection), or PBS starting on day 5 post

EAE induction, and continuing every third day until day 32. Mice were observed daily for disease

symptoms and EAE was scored as per the scoring system described (*11*).

**180    Transcriptome sequencing and analysis**

Tissue sections from the colon of both DR3.IL17A$^{-/-}$ and DR3 mice before EAE induction were

collected, chopped into small pieces, stored in RNAlater (Millipore Sigma, MO, USA), and frozen

to -80$^{0}$C until further use. RNA was extracted using the RNeasy Kit (Qiagen) following the

manufacturer's instructions. The quality and quantity of the RNA was evaluated using a NanoDrop

and sequenced at Novogene Co. Ltd, CA. Sequences were quality controlled using default *fastqc*

(*12*). Paired FASTQ files were aligned to the mm10 genome build with the kallisto pseudoaligner

(*13*) using 100 bootstraps. Estimated counts for transcripts were then aggregated into gene-level

quantifications based on gene symbol. An average of 14.79 ±1.3 million transcripts were obtained

for each sample. Further downstream processess were then carried out in R 4.0.3 (*14*) using edgeR

3.32.1 (*15*) to identify differentially expressed genes, as described previously. Differentially

expressed genes were fed into ShinyGO (*16*) to identify changes to genes involved in GO

biological pathways using the mouse reference database.

**Supplementary Figures**

**Fig. S1: HLA-DR3.IL17A$^{-/-}$ mice lack IL-17A-expressing CD4+ T cells in the spleen.** 8-10-

week-old HLA-DR3 and HLA-DR3.IL17A$^{-/-}$ transgenic mice were euthanized and splenocytes

were harvested for analysis. Quantification of the frequency of CD4$^+$IL-17A$^+$ T cells from mice in

A. HLA-DR3: n=3; HLA-DR3.IL17A$^{-/-}$: n = 4. *p*-value was determined using Mann-Whitney test

between the experimental groups. Symbols represent individual mice, bars represent mean±SEM.

**Fig. S2: Anti-IL-17A treatment lowers the severity of MOG$_{35-55}$ induced EAE in C57BL/6J mice. (A-B)** 8-10-week-old B6 mice were immunized with MOG$_{35-55}$/CFA. Five days after immunization, mice received anti-mouse IL-17A monoclonal antibody, isotype control mouse IgG1, or PBS intraperitoneally every three days until the end of the experiment (day 32). (A) Average clinical EAE scores over time. Treatment days are indicated by arrowheads. (B) Average cumulative EAE scores. **(C-D)** 8-10-week-old B6 mice were treated every three days with anti-IL-17A, IgG1, or PBS pre- and post-immunization with MOG$_{35-55}$/CFA. (C) Average clinical EAE scores. Treatment days are indicated by arrowheads. (D) Average cumulative EAE scores. For plots in A, C, each bullet point represents the mean and bars represent the standard error of the mean from each experimental group. *p*-value was determined using two-way ANOVA following Tukey correction for EAE clinical scores (A, C) and unpaired t-test with Welch correction for cumulative EAE score (B, D). For graphs in B, D, each bar represents mean±SEM.

**Fig. S3: Levels of inflammatory cytokines IL-17F in HLA-DR3 and HLA-DR3.IL17A[-/-] mice.**
8-10-week-old mice were immunized with $PLP_{91-110}$ /CFA to induce EAE. After 20 days, mice
were euthanized, spleens were harvested, and splenic cells were cultured in the presence of $PLP_{91-110}$
antigen for five days following stimulation with Phorbol myristate acetate (PMA) and
ionomycin in the presence of Brefeldin A (BFA). The supernatant was collected and levels of
IL17F were measured using ELISA. Cytokine values from DR3 mice were compared to those of
DR3.IL17A[-/-] mice in either media (Med) or after stimulation with $PLP_{91-110}$. *p*-value was
determined using unpaired t-test with Welch correction. Bars represent mean±S.E.M.

Shahi et al., 2022

**Fig. S4: Treatment with anti-GM-CSF or anti-IL-17F monoclonal antibodies increases**
245

**disease severity in HLA-DR3.IL17A$^{-/-}$ mice.** (**A**) Schematic diagram of the experimental
246

workflow. EAE was induced in 8-10-week-old in HLA-DR3.IL17A$^{-/-}$ transgenic mice. At days 5,
247

10, and 14 post-EAE induction, mice received intraperitoneal injections of either: 1) anti-GM-CSF
248

monoclonal antibody, 2) anti-IL-17F monoclonal antibody, or 3) IgGa1 isotype control and clinical
249

EAE scores were monitored for 31 days. (**B**) Average clinical EAE score of mice in each group
250

over time. Bullet points represent the mean and bars represent the standard error of the mean of
251

each experimental group from two independent experiments (n ≥3 mice per group). *p*-value was
252

determined by a two-way ANOVA following Tukey correction for average clinical EAE score
253

(B). (**C**) Average cumulative EAE scores. *p*-value was determined by unpaired t-test with Welch
254

correction (C). For plots in B, C, each symbol represents the mean and bars represent the standard
255

error of the mean from each experimental group.
256

**Fig. S5: HLA-DR3.IL17A$^{-/-}$ mice develops milder EAE in absence of IFNγ .** EAE was induced in 8-10-week-old in HLA-DR3 and HLA-DR3.IL17A$^{-/-}$IFNγ$^{-/-}$ transgenic mice using PLP$_{91-110}$, CFA and PTX. The disease was monitored and scored daily until day 25 of the experiment. Average clinical EAE score of mice in each group over time. Bullet points represent the mean and bars represent the standard error of the mean of each experimental group from two independent experiments (n =4 mice per group). $p$-value was determined by multiple unpaired t-tests with correction using two-stage step up method of Benjamini, Krieger and Yekutieli for average clinical EAE score.

**Fig. S6:** Similarity matrix of microbial compositional communities present in DR3 and DR3.IL17A$^{-/-}$ mice before (BCH) and after (ACH) cohousing. Matrix was computed from the relative abundance matrix using 1-Spearman rank correlation metric using average linkage method and clustered hierarchically using Morpheus webserver (https://software.broadinstitute.org/morpheus).

**Fig. S7:** Composition of the gut bacteria in DR3 and DR3IL17A[-/-] transgenic mice before (BCH)
and after (ACH) mice were cohoused. Each bar represents the average relative abundance (%) of
the bacterium in each group. Kruskal Wallis test followed by Dunn-test was performed to identify
the significance between the groups. 'p-value' from A-N represents the significance for the
Kruskal Wallis test. Those bacteria with significant/non-significant p-value from Kruskal Wallis
test but non-significantly different in at least one of the groups of interest (DR3-BCH to
DR3IL17A[-/-]-BCH, DR3-BCH to DR3-ACH and DR3IL17A[-/-]-BCH to DR3IL17A[-/-]-ACH)
following Dunn test are shown.

**Fig. S8**: Abundance of short chain fatty acid pathways, as determined by shotgun metagenomics analysis in DR3.IL17$^{-/-}$ and DR3 transgenic mice before (BCH) and after (ACH) they were cohoused. Pairwise Wilcoxon test was performed to obtain the statistical significance BCH and ACH for each DR3 and DR3.IL17$^{-/-}$ transgenic mice.

311 **Tables:**

312 **Table S1:** Differentially expressed gene list in the colon of DR3.IL17A$^{-/-}$ compared to DR3

313 transgenic mice calculated using *limma* package within *edgeR* in R. All 109 differentially

314 expressed genes with *LogFC*>1.0 and *adj.p.val*<0.05 are shown.

| SN | Genes | logFC | AveExpr | t | P.Value | adj.P.Val |
|---|---|---|---|---|---|---|
| 1 | Eno1b | -12.154 | 1.443 | -19.576 | 0.000 | 0.000 |
| 2 | Hoxb13 | -11.448 | -1.890 | -19.897 | 0.000 | 0.000 |
| 3 | Hoxd13 | -9.589 | -1.902 | -10.871 | 0.000 | 0.000 |
| 4 | Casp14 | -8.793 | -1.658 | -13.128 | 0.000 | 0.000 |
| 5 | Glo1-ps | -8.601 | -0.491 | -9.588 | 0.000 | 0.000 |
| 6 | 2310079G19Rik | -8.148 | -2.842 | -8.433 | 0.000 | 0.000 |
| 7 | Rpl15-ps6 | -7.886 | 0.116 | -6.560 | 0.000 | 0.002 |
| 8 | Gm13237 | -7.823 | -0.170 | -8.445 | 0.000 | 0.000 |
| 9 | Ly6g | -7.548 | -2.237 | -11.113 | 0.000 | 0.000 |
| 10 | Tgm3 | -6.671 | 3.560 | -31.635 | 0.000 | 0.000 |
| 11 | Vtcn1 | -6.393 | -3.095 | -8.575 | 0.000 | 0.000 |
| 12 | Mettl7a2 | -6.307 | -3.303 | -8.664 | 0.000 | 0.000 |
| 13 | Igkv8-18 | -6.293 | -0.666 | -8.186 | 0.000 | 0.000 |
| 14 | Fxyd4 | -6.059 | -1.910 | -4.809 | 0.000 | 0.025 |
| 15 | 5033404E19Rik | -5.855 | -2.774 | -5.566 | 0.000 | 0.008 |
| 16 | Gm7334 | -5.659 | 1.342 | -5.672 | 0.000 | 0.007 |
| 17 | Sval1 | -5.631 | 3.704 | -19.493 | 0.000 | 0.000 |
| 18 | Evx2 | -5.587 | -2.795 | -6.518 | 0.000 | 0.002 |
| 19 | Gm8540 | -5.532 | -3.496 | -7.667 | 0.000 | 0.000 |
| 20 | Atp12a | -5.477 | 0.033 | -18.752 | 0.000 | 0.000 |
| 21 | Chit1 | -4.989 | -3.144 | -5.998 | 0.000 | 0.004 |
| 22 | Trpv6 | -4.912 | -0.557 | -8.873 | 0.000 | 0.000 |
| 23 | Igkv2-112 | -4.725 | 1.275 | -9.585 | 0.000 | 0.000 |
| 24 | Fut9 | -4.701 | -1.951 | -6.631 | 0.000 | 0.002 |
| 25 | Ighv7-1 | -4.595 | 0.648 | -6.919 | 0.000 | 0.001 |
| 26 | Ninj2 | -4.593 | -1.906 | -5.754 | 0.000 | 0.006 |
| 27 | St8sia5 | -4.551 | -1.819 | -5.220 | 0.000 | 0.013 |
| 28 | Ighv1-61 | -4.532 | 1.365 | -7.272 | 0.000 | 0.001 |
| 29 | Tmprss13 | -4.269 | -0.183 | -8.813 | 0.000 | 0.000 |
| 30 | Gjb5 | -4.261 | -1.541 | -4.778 | 0.000 | 0.025 |
| 31 | Hoxd12 | -4.259 | -3.247 | -4.345 | 0.000 | 0.049 |
| 32 | Vsig1 | -4.255 | -1.212 | -6.516 | 0.000 | 0.002 |
| 33 | Ighv10-3 | -4.253 | 3.640 | -11.518 | 0.000 | 0.000 |

| 34 | B3gnt7 | -4.195 | 5.973 | -34.648 | 0.000 | 0.000 |
|----|----------|--------|--------|---------|-------|-------|
| 35 | Rpl15-ps2 | -3.829 | -0.635 | -4.963 | 0.000 | 0.019 |
| 36 | Itih2 | -3.817 | -1.521 | -4.471 | 0.000 | 0.041 |
| 37 | Mettl7a3 | -3.571 | -1.774 | -4.445 | 0.000 | 0.042 |
| 38 | Pla2g4f | -3.458 | 0.531 | -5.797 | 0.000 | 0.006 |
| 39 | Trpv3 | -3.403 | 3.252 | -12.943 | 0.000 | 0.000 |
| 40 | Thbs4 | -3.334 | 3.057 | -21.508 | 0.000 | 0.000 |
| 41 | Hoxa13 | -3.280 | -1.700 | -8.765 | 0.000 | 0.000 |
| 42 | Itln1 | -3.187 | 6.879 | -10.731 | 0.000 | 0.000 |
| 43 | Sycn | -3.151 | 2.937 | -24.593 | 0.000 | 0.000 |
| 44 | Nxpe4 | -3.057 | 3.610 | -15.040 | 0.000 | 0.000 |
| 45 | Insl5 | -2.941 | -1.314 | -6.488 | 0.000 | 0.002 |
| 46 | Ighv2-5 | -2.897 | 2.075 | -4.897 | 0.000 | 0.021 |
| 47 | Adgrf1 | -2.818 | 0.607 | -9.433 | 0.000 | 0.000 |
| 48 | Igkv3-4 | -2.711 | 3.280 | -8.454 | 0.000 | 0.000 |
| 49 | Igkv4-57 | -2.655 | 4.588 | -12.494 | 0.000 | 0.000 |
| 50 | Slc28a3 | -2.572 | 2.546 | -4.625 | 0.000 | 0.032 |
| 51 | Igkv4-68 | -2.565 | 4.316 | -8.356 | 0.000 | 0.000 |
| 52 | Pla2g2f | -2.494 | 3.879 | -7.097 | 0.000 | 0.001 |
| 53 | Ighv3-3 | -2.458 | 1.456 | -8.997 | 0.000 | 0.000 |
| 54 | Cyp2d12 | -2.407 | 1.294 | -9.324 | 0.000 | 0.000 |
| 55 | Best2 | -2.363 | 0.866 | -10.732 | 0.000 | 0.000 |
| 56 | Gm47283 | -2.290 | 5.085 | -10.117 | 0.000 | 0.000 |
| 57 | Adh1 | -2.274 | 5.559 | -21.681 | 0.000 | 0.000 |
| 58 | Ighv1-39 | -2.271 | 3.886 | -4.873 | 0.000 | 0.022 |
| 59 | Cyp2c68 | -2.248 | 4.529 | -8.006 | 0.000 | 0.000 |
| 60 | B3gnt5 | -2.203 | 3.517 | -6.569 | 0.000 | 0.002 |
| 61 | Ighv5-4 | -2.151 | 2.640 | -5.001 | 0.000 | 0.018 |
| 62 | Ggh | -2.100 | 5.137 | -16.144 | 0.000 | 0.000 |
| 63 | Cyp2d9 | -2.078 | 0.066 | -4.490 | 0.000 | 0.040 |
| 64 | Zfp791 | -2.041 | 2.587 | -4.591 | 0.000 | 0.034 |
| 65 | Slc15a2 | -2.026 | 2.158 | -4.924 | 0.000 | 0.020 |
| 66 | Nox1 | -2.008 | 3.528 | -7.361 | 0.000 | 0.001 |
| 67 | Nt5e | -1.973 | 5.284 | -10.629 | 0.000 | 0.000 |
| 68 | Duoxa2 | -1.952 | 6.303 | -10.058 | 0.000 | 0.000 |
| 69 | Ang4 | -1.951 | 7.834 | -9.720 | 0.000 | 0.000 |
| 70 | Ighv2-9-1 | -1.949 | 3.374 | -4.962 | 0.000 | 0.019 |
| 71 | Slc15a1 | -1.948 | 6.682 | -8.009 | 0.000 | 0.000 |
| 72 | Igkv3-5 | -1.942 | 3.424 | -5.541 | 0.000 | 0.008 |

| 73 | Cldn8 | -1.909 | 4.312 | -9.203 | 0.000 | 0.000 |
|---|---|---|---|---|---|---|
| 74 | Eno3 | -1.762 | 3.750 | -6.374 | 0.000 | 0.002 |
| 75 | Clic6 | -1.762 | 3.833 | -5.762 | 0.000 | 0.006 |
| 76 | Cd177 | -1.749 | 6.807 | -16.983 | 0.000 | 0.000 |
| 77 | Adamts15 | -1.691 | 4.728 | -5.747 | 0.000 | 0.006 |
| 78 | Ccl28 | -1.628 | 7.068 | -12.416 | 0.000 | 0.000 |
| 79 | Igkv9-120 | -1.622 | 4.390 | -4.733 | 0.000 | 0.027 |
| 80 | Hoxa11 | -1.621 | 0.018 | -5.452 | 0.000 | 0.009 |
| 81 | Ceacam10 | -1.593 | 5.986 | -8.932 | 0.000 | 0.000 |
| 82 | Aqp4 | -1.573 | 6.556 | -14.118 | 0.000 | 0.000 |
| 83 | Saa1 | -1.550 | 7.353 | -7.683 | 0.000 | 0.000 |
| 84 | Ano1 | -1.550 | 3.699 | -6.163 | 0.000 | 0.003 |
| 85 | Tcf7l1 | -1.529 | 2.331 | -4.795 | 0.000 | 0.025 |
| 86 | Cela1 | -1.499 | 4.613 | -7.343 | 0.000 | 0.001 |
| 87 | Gm15401 | -1.437 | 3.493 | -4.550 | 0.000 | 0.036 |
| 88 | Prdx6 | -1.317 | 8.165 | -8.030 | 0.000 | 0.000 |
| 89 | Retnlb | -1.259 | 5.150 | -4.445 | 0.000 | 0.042 |
| 90 | Scd2 | -1.133 | 9.600 | -5.131 | 0.000 | 0.015 |
| 91 | Plin4 | 1.287 | 6.266 | 4.615 | 0.000 | 0.033 |
| 92 | Cyp4b1 | 1.379 | 4.923 | 4.760 | 0.000 | 0.026 |
| 93 | Slc13a1 | 1.393 | 7.159 | 5.688 | 0.000 | 0.006 |
| 94 | Hbb-bs | 1.412 | 5.567 | 5.082 | 0.000 | 0.016 |
| 95 | Naaladl1 | 1.451 | 8.878 | 5.901 | 0.000 | 0.005 |
| 96 | Oas2 | 1.473 | 3.640 | 4.420 | 0.000 | 0.044 |
| 97 | Tef | 1.533 | 4.008 | 5.337 | 0.000 | 0.011 |
| 98 | Oasl2 | 1.568 | 5.620 | 8.209 | 0.000 | 0.000 |
| 99 | Fabp4 | 1.636 | 8.267 | 8.666 | 0.000 | 0.000 |
| 100 | Ifit1 | 1.636 | 6.835 | 13.024 | 0.000 | 0.000 |
| 101 | Rpl3 | 1.638 | 7.213 | 9.483 | 0.000 | 0.000 |
| 102 | Thrsp | 1.641 | 5.122 | 4.949 | 0.000 | 0.020 |
| 103 | Ddx60 | 1.649 | 4.067 | 6.229 | 0.000 | 0.003 |
| 104 | Ifit3 | 1.670 | 4.893 | 6.902 | 0.000 | 0.001 |
| 105 | Retn | 1.716 | 4.432 | 5.022 | 0.000 | 0.018 |
| 106 | Sis | 1.727 | 7.589 | 5.489 | 0.000 | 0.009 |
| 107 | Igkv12-98 | 1.757 | 2.715 | 4.414 | 0.000 | 0.044 |
| 108 | Oas3 | 1.774 | 4.742 | 8.831 | 0.000 | 0.000 |
| 109 | Dbp | 1.855 | 4.802 | 9.977 | 0.000 | 0.000 |
| 110 | Rpl15 | 1.893 | 7.014 | 8.315 | 0.000 | 0.000 |
| 111 | Igkv14-100 | 1.910 | 3.744 | 5.128 | 0.000 | 0.015 |

| 112 | Car3 | 2.300 | 7.529 | 24.744 | 0.000 | 0.000 |
|-----|------|-------|-------|--------|-------|-------|
| 113 | Cfd | 2.334 | 7.125 | 24.230 | 0.000 | 0.000 |
| 114 | Adipoq | 2.474 | 4.756 | 11.111 | 0.000 | 0.000 |
| 115 | Plin1 | 2.574 | 4.541 | 7.430 | 0.000 | 0.001 |
| 116 | Lgals2 | 2.688 | 5.834 | 6.063 | 0.000 | 0.004 |
| 117 | Tcaim | 2.984 | 2.326 | 6.378 | 0.000 | 0.002 |
| 118 | Trgc2 | 2.984 | 1.211 | 4.446 | 0.000 | 0.042 |
| 119 | Cyp2e1 | 3.009 | 5.145 | 15.014 | 0.000 | 0.000 |
| 120 | Lyrm7 | 3.114 | 0.716 | 4.632 | 0.000 | 0.032 |
| 121 | Gm16011 | 4.232 | -0.321 | 4.370 | 0.000 | 0.047 |
| 122 | Igkv14-126 | 5.013 | 0.842 | 7.976 | 0.000 | 0.000 |
| 123 | Cylc2 | 8.644 | -1.325 | 9.999 | 0.000 | 0.000 |

## References

1. S. K. Shahi, S. N. Freedman, R. A. Dahl, N. J. Karandikar, A. K. Mangalam, Scoring disease in an animal model of multiple sclerosis using a novel infrared-based automated activity-monitoring system. *Sci Rep* **9**, 19194 (2019).

2. A. F. Tyler, J. P. Mendoza, M. Firan, N. J. Karandikar, CD8(+) T Cells Are Required For Glatiramer Acetate Therapy in Autoimmune Demyelinating Disease. *PLoS One* **8**, e66772 (2013).

3. A. Mangalam *et al.*, HLA-DQ8 (DQB1*0302)-restricted Th17 cells exacerbate experimental autoimmune encephalomyelitis in HLA-DR3-transgenic mice. *J Immunol* **182**, 5131-5139 (2009).

4. A. K. Mangalam *et al.*, Absence of IFN-gamma increases brain pathology in experimental autoimmune encephalomyelitis-susceptible DRB1*0301.DQ8 HLA transgenic mice through secretion of proinflammatory cytokine IL-17 and induction of pathogenic monocytes/microglia into the central nervous system. *J Immunol* **193**, 4859-4870 (2014).

5. L. Goschl *et al.*, A T cell-specific deletion of HDAC1 protects against experimental autoimmune encephalomyelitis. *J Autoimmun* **86**, 51-61 (2018).

6. A. M. Thornton, E. M. Shevach, CD4+CD25+ immunoregulatory T cells suppress polyclonal T cell activation in vitro by inhibiting interleukin 2 production. *J Exp Med* **188**, 287-296 (1998).

7. F. Krueger, Trim Galore!: a wrapper tool around Cutadapt and FastQC to consistently apply quality and adapter trimming to FastQ files.  (2014).

8. B. Langmead, S. L. Salzberg, Fast gapped-read alignment with Bowtie 2. *Nat Methods* **9**, 357-359 (2012).

9. F. Beghini *et al.*, Integrating taxonomic, functional, and strain-level profiling of diverse microbial communities with bioBakery 3. *Elife* **10**,  (2021).

10. D. Li, C. M. Liu, R. Luo, K. Sadakane, T. W. Lam, MEGAHIT: an ultra-fast single-node solution for large and complex metagenomics assembly via succinct de Bruijn graph. *Bioinformatics* **31**, 1674-1676 (2015).

11. A. Mangalam *et al.*, Human Gut-Derived Commensal Bacteria Suppress CNS Inflammatory and Demyelinating Disease. *Cell Rep* **20**, 1269-1277 (2017).

12. S. Andrews. (2010).

13. N. L. Bray, H. Pimentel, P. Melsted, L. Pachter, Near-optimal probabilistic RNA-seq quantification. *Nat Biotechnol* **34**, 525-527 (2016).

14. R Core Team, R: A language and environment for statistical computing *R Foundation for Statistical Computing, Vienna, Austria*,  (2020).

15. M. D. Robinson, D. J. McCarthy, G. K. Smyth, edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics* **26**, 139-140 (2010).

16. S. X. Ge, D. Jung, R. Yao, ShinyGO: a graphical gene-set enrichment tool for animals and plants. *Bioinformatics* **36**, 2628-2629 (2020).