1   **RESEARCH ARTICLE**

2   **Polarly Localized WPR Proteins Interact With PAN Receptors And The**
3   **Actin Cytoskeleton During Maize Stomatal Development**

4   Qiong Nan[1], Si Nian Char[2], Bing Yang[3], Eric J. Bennett[3], Bing Yang[2,4],
5   Michelle R. Facette[1]
6
Orcid IDS:
QN: 0000-0002-9940-3015, SNC: 0000-0002-5759-0764, BY[3]: 0000-0001-6739-2932,
EJB: 0000-0002-1201-331, BY[2,4]: 0000-0002-2293-33844 MRF: 0000-0002-6214-9359

[1]Department of Biology, University of Massachusetts, Amherst, MA, 01003, USA
[2]Division of Plant Sciences, Bond Life Sciences Center, University of Missouri, Columbia, MO
65211, USA
[3]University of California, San Diego, La Jolla, CA 92093, USA
[4]Donald Danforth Plant Science Center, St. Louis, MO 63132, USA

Short title: WPR proteins bind actin and PAN receptors

One-sentence summary: Four related proteins, identified via their physical
interaction with the receptor PAN2, are polarly localized prior to asymmetric
division in stomatal lineage cells, and interact with F-actin.

The author responsible for distribution of materials integral to the findings
presented in this article in accordance with the policy described in the
Instructions for Authors is: Michelle Facette (mfacette@umass.edu)
22
23

ABSTRACT

Polarization of cells prior to asymmetric cell division is crucial for correct cell
divisions, cell fate and tissue patterning. In maize stomatal development,
polarization of subsidiary mother cells prior to asymmetric division is
controlled by the BRK-PAN-ROP pathway. Two catalytically inactive receptor-
like kinases, PAN2 and PAN1, are required for correct division plane
positioning. Proteins in the BRK-PAN-ROP pathway are polarized in
subsidiary mother cells, with the polarization of each protein dependent on the
previous one. As most of the known proteins in this pathway do not physically
interact, possible interactors that might participate in the pathway are yet to be
described. We identified WPR proteins as new players during subsidiary
mother cell polarization. WPRs physically interact with PAN receptors, and
polarly accumulate in subsidiary mother cells. The polarized localization of
WPR proteins depends on PAN2 but not PAN1. CRISPR-Cas9-
induced mutations result in division plane defects in subsidiary mother cells,
and ectopic expression of WPR-RFP results in stomatal defects and
alterations to the actin cytoskeleton. We show certain WPR proteins directly
interact with F-actin through their N-terminus. Our data implicate WPR
proteins as potentially regulating actin filaments, which providing insight into
their molecular function. Together, these results demonstrate that WPR
proteins are important for cell polarization.

INTRODUCTION

Asymmetric cell division, which generates two daughter cells with different fates, is indispensable for cellular differentiation and diversity. In plants, cells do not move, therefore asymmetric cell division is particularly important for cellular patterning (De Smet and Beeckman, 2011; Muroyama and Bergmann, 2019). Certain processes are conserved during asymmetric cell division, although the specific molecular players may vary. A key principle of asymmetric cell division is cell polarization prior to mitosis, which is often marked by the polarization of plasma membrane–associated proteins and nuclear migration to the future division plane (Facette and Smith, 2012; Galatis and Apostolakos, 2004; Yoshida et al., 2019). Cell polarization may be determined by intrinsic cues (pre-existing within the mother cell) or extrinsic cues (originating from outside the mother cell) (Yang 2008; Facette et al., 2019; Lipka et al., 2015).

The asymmetric division of maize subsidiary mother cells (SMCs) has been used as a model to understand cell polarization and asymmetric division in plants. Grasses such as *Zea mays* (maize), *Oryza spp* (rice) and *Brachypodium distachyon* (purple false brome) possess 4-celled stomatal complexes consisting of two guard cells flanked by a pair of subsidiary cells **(Supplemental Figure 1)**. Grass stomatal formation is initiated by an asymmetric division that generates a guard mother cell (GMC). The lateral neighbors of the GMC, called subsidiary mother cells (SMCs), divide asymmetrically to produce subsidiary cells that flank the GMC. It is proposed that SMC divisions are induced by an extrinsic cue from the GMC. Eventually, the GMC divides symmetrically to form a pair of guard cells flanked by the subsidiary cells (Stebbins and Shah, 1960; Facette and Smith, 2012; Raissig et al., 2017**;** Nunes et al., 2020; Gray et al., 2020).

In maize SMCs, several plasma membrane–associated proteins make up the BRK-PAN-ROP pathway, which promotes SMC polarity. BRICK1 (BRK1) is the earliest marker of SMC polarity (Facette et al., 2015). BRICK1 is a member of SCAR/WAVE complex that promotes branched actin networks via activation of the Arp2/3 complex (Frank and Smith 2002; Facette et al., 2015). BRK1, and all subsequent proteins, are polarized within SMCs,

accumulating at the site of GMC contact. BRK1 is required for PANGLOSS2

(PAN2) and PAN1 polarization, which are two leucine-rich repeat receptor-like

proteins with inactive kinase domains. PAN2 is required for the subsequent

polarization of PAN1 (Zhang et al., 2012). Plant Rho family GTPase (ROP) is

a small GTPase that physically interacts with PAN1 and its polarization is

dependent on PAN1 (Humphries et al., 2011). After the polarized

accumulation of these proteins, an actin patch appears at the SMC-GMC

interface and the pre-mitotic SMC nucleus migrates towards the GMC. Within

the pathway, only PAN1 and ROP have been shown to physically interact,

however each protein is required for the next to be polarized. This implies that

there are additional proteins important for the recruitment and/or polarization

of each of the known players. Moreover, the molecular role that each of these

proteins performs during SMC polarization is not known, including how

polarization of these membrane-localized proteins culminates in nuclear

migration and positioning of the division plane.

Since PAN1 and PAN2 are catalytically inactive kinases (Humphries et

al., 2011; Zhang et al., 2012), it is unclear what their molecular function is.

They may act as pseudokinases with important roles in scaffolding; however

other than ROP no other PAN-interacting proteins have yet been

characterized. In some cases, catalytically inactive receptors partner with

active leucine-rich repeat receptor kinases and act as inhibitors of function

(Lee et al., 2012), but to date no active kinases have been identified in the

pathway. PAN1 and PAN2 might perform functions similar to other polarized

plasma membrane proteins and mediate diverse downstream functions that

relate to cellular polarization such as: signal transduction, accumulation of

other cell polarity factors, or eventual nuclear migration (De Smet and

Beeckman, 2011; Ashraf and Facette, 2020). For example, in *A. thaliana*

stomatal development, the polarized proteins BREAKING OF ASYMMETRY

IN THE STOMATAL LINEAGE (BASL) and POLAR LOCALIZATION

DURING ASYMMETRIC DIVISION AND REDISTRIBUTION (POLAR) act as

scaffolding proteins for localized signal transduction (Dong et al., 2009;

Pillitteri et al., 2011; Houbaert et al., 2018). Notably, the *B. distachyon*

orthologue of POLAR was recently discovered to have a distinct localization

pattern during grass stomatal development that is opposite that of PAN1 – it is

excluded from the GMC-SMC contact site (Zhang et al., 2022). Polarized

BASL is required for pre-mitotic microtubule-based nuclear migration and

post-mitotic actin-based nuclear migration (Muroyama et al., 2020). The LRR-

RLK INFLORESCENCE AND ROOT APICES RECEPTOR KINASE (IRK) is

polarly localized in root cells that divide (or have recently divided)

asymmetrically however the molecular pathway downstream of IRK is not

characterized (Campos et al., 2020).

Polarized accumulation of actin filaments is a prominent feature of

maize SMC division, but the functional role of the actin patch that forms late in

SMC polarization is unclear. While SMC nuclear migration in grasses is based

on actin networks (Cho and Wick, 1990), formation of the actin patch and

nuclear migration can be uncoupled (Cartwright et al., 2009; Apostolakos et

al., 2018), implying that the actin patch has a function distinct from promoting

nuclear migration. It is plausible that ROP GTPases are important for

stimulating actin patch formation, however this has not been clearly

demonstrated. In addition to the actin patch, there are other important actin-

related processes during SMC polarization. The early role of BRK proteins,

actin-mediated nuclear migration and actin patch formation imply multiple

functions for actin networks in SMC polarization. SMC actin filaments may

provide force or/and tracks for polarized organelle movement, and/or may

facilitate endo/exocytosis at the polarity site (Hadley et al., 2006; Kimata et al.,

2016; Wu and Bezanilla, 2018).

Actin-dependent movement of chloroplasts has been shown to involve

WEAK CHLOROPLAST MOVEMENT UNDER BLUE LIGHT 1 (WEB1) and

PLASTID MOVEMENT IMPAIRED 2 (PMI2) (Luesse et al., 2006; Kodama et

al., 2010). WEB1, PMI2 and similar WEB1/PMI2-RELATED (WPR) proteins

have a DUF827 domain which is a series of coiled-coils; *A. thaliana* has

fourteen DUF827-containing proteins including the WEB1-like (WEL) clade,

PMI2-LIKE (PMI) clade and the WPRA and WPRB clades (Kodama et al.,

2011). Previous studies in *A. thaliana* suggest proteins with this domain have

diverse functions. WEB1 (Weak Chloroplast Movement under Blue Light 1)

and PMI2 (Plastid Movement Impaired 2) are interacting proteins and are

required for actin-based chloroplast photorelocation (Luesse et al., 2006;

Kodama et al., 2010). TOUCH-REGULATED PHOSPHOPROTEIN1

(TREPH1) belongs to the WPRA protein clade, and is phosphorylated in

response to touch (Wang et al., 2018). TREPH1 is required for touch-induced

growth repression. WEB1 is membrane-associated and PMI2 is cytosolic in

protoplasts, but *in planta* localization for DUF827 proteins has not been

demonstrated. Moreover, while the demonstrated phenotypes of WEB1 and

PMI2 implies an actin-related function, no evidence for physical or regulatory

interactions with actin have been shown for any DUF827 protein thus far.

In this study, we identified members of the DUF827 domain-containing

WPR protein family that participate in maize SMC polarization. WPRs interact

with both PAN2 and PAN1, and WPRs can also form homodimers and

heterodimers. WPRs localize polarly in SMC at sites of guard mother cell

(GMC) contact, and their polarized localization depends on PAN2 but not

PAN1. These findings suggest WPRs are plasma membrane-associated

proteins that are important for pre-mitotic polarity in maize SMCs, and are the

first physical link between PAN1 and PAN2. CRISPR-Cas9-

induced *wprb1;wprb2* double mutants and plants ectopically expressing

WPRB2 have an increased frequency of aberrantly formed subsidiary cells.

WPRB2 overexpression causes a decrease in cellular F-actin. Additionally,

we show that WPRB2 directly interacts with F-actin through its N-terminus. In

total, these results suggest the WPRs are components of the BRK-PAN-ROP

pathway and likely act downstream of PAN2 to regulate actin-associated

processes during cell polarization.

RESULTS

WPRs Exhibit Polarized Localization in Maize SMCs

To better understand the functions of PAN and BRK proteins during

SMC polarization, we identified proteins that physically interact with PAN2-

YFP and PAN1-YFP. Co-immunoprecipitation/mass spectrometry (Co-IP/MS)

of BRK1-CFP, Rab11D-YFP, PIN1-YFP, PDI-YFP and non-transgenic

controls was previously performed using anti-GFP beads (Facette et al.,

2015). In parallel, co-IP/MS of PAN2-YFP and PAN1-YFP was also

performed, but not presented in Facette et al. (2015). As previously published,

a WD-score was calculated to identify high-confidence interactors (Facette et

al., 2015; Sowa et al., 2009). Amongst the PAN2-YFP interactors were four

related proteins containing Domain of Unknown Function 827 (DUF827)

(Supplemental Dataset 1). The DUF827 family includes previously identified

WEB1-PMI2 RELATED (WPR) proteins (Kodama et al., 2010; Gardiner et al.,

2011; Kodama et al., 2011). We identified sixteen WPRs in *A. thaliana* and

seventeen WPRs in maize. A protein tree was inferred, and these

(Supplemental Figure 2) proteins fall into five clades: WPRA, PMI,

WEB/WEL, WPRB and WPRC. The four maize proteins interacting with PAN2

fall into WPRA and WPRB clades; we named these proteins WPRA1,

WPRA2, WPRB1, and WPRB2. WPRs have predicted coiled-coil domains in

their central plant-specific DUF827 region, flanked by uncharacterized N- and

C-terminal regions.

Thus far, all proteins identified in the BRK-PAN-ROP pathway polarly

localize in SMCs prior to division (Cartwright et al., 2009; Humphries et al.,

2011; Zhang et al., 2012; Facette et al., 2015). To determine if WPR proteins

play a role in SMC polarization, we determined if they were similarly polarized.

We created stable transgenic maize lines expressing either CFP-WPRA2 or

RFP-WPRB2. Genomic fragments containing the native promoter, introns and

terminator were used. At early stages of stomatal development prior to SMC

formation, CFP-WPRA2 **(Figure 1A)** and RFP-WPRB2 **(Figure 1B)**

predominantly localize to the cell periphery in all cell types, with low levels of

cytoplasmic fluorescence. RFP-WPRB2 is enriched in cell corners, similar to

BRK1-CFP (Facette et al., 2015). As development proceeds, CFP-WPRA2

and RFP-WPRB2 become enriched in the SMC at the site of GMC contact

and remain polarized after SMC division **(Figure 1)**. Plasmolysis experiments

confirm that CFP-WPRA2 and RFP-WPRB2 are polarized in the SMC, not the

GMC **(Supplemental Figure 3)**. We further validated the localization by

performing immunofluorescence using custom-generated antibodies that

recognize either WRPA1/WPRA2 or WPRB2 (**Supplemental Fig 4**). Similar

to the transgenic lines, we saw polarized localization in SMCs and enrichment

at cell corners. Thus, the localization of WPRA2 and WPRB2 is similar to

known BRK-PAN-ROP pathway components and supports a role in SMC

polarization. Notably, WPR proteins contain no predicted transmembrane

domains or lipidation signals, indicating their association with the membrane

is either peripheral or via interactions with other membrane proteins.

WPRs Interact with PAN1 and PAN2

216   To confirm the WPR-PAN2 interaction, we performed reciprocal co-

IP/MS experiments using three strategies, using either our fluorescent protein

fusion lines or a native antibody. Firstly, we used anti-GFP to

immunoprecipitate WPRA2 from transgenic plants expressing CFP-WPRA2;

non-transgenic wild type plants were used as a negative control. Secondly,

we used anti-RFP to immunoprecipitate WPRB2 from transgenic plants

expressing RFP-WPRB2, again using non-transgenic plants as a negative

control. Thirdly, we used an antibody raised against WPRA1/A2 (see

Methods, and **Supplemental Fig. 4**) to pull down endogenous WPRA from

non-transgenic plant extracts, with reactions omitting this antibody as a

negative control. In each experiment, three biological replicates were

performed. After comparing relative abundances of identified proteins with

corresponding negative controls, we found WPRA family proteins co-

precipitated with WPRB family proteins in all three assays (**Table 1 and**

Supplement Data Set 2). Moreover, we found that both PAN1 and PAN2 co-

precipitated with WPRA1/A2 and PAN2 co-precipitated with RFP-WPRB2. In

addition, enrichment of WPRB3 peptides was detected in CFP-WPRA2 and

WPRA1/A2 immunoprecipitates, suggesting WPRB3 may also associate with

WPRA proteins.

235   To validate these interactions, and test whether WPRs directly interact

with PAN1 and PAN2, yeast two-hybrid assays were conducted using

intracellular portions of PAN1 and PAN2 as bait (Zhang et al., 2012), and

WPRA1, WPRA2, WPRB1, WPRB2, and WPRB3 as prey. In these assays,

no interaction between WPRA1 or WPRA2 and PAN1 or PAN2 was observed.

However, we found that both WPRB1 and WPRB2 interact with PAN1 and

PAN2. WPRB3 interacts with PAN2 but not PAN1 **(Figure 2A and 2B)**. These

data confirm a direct physical interaction between the intracellular regions of

PAN and WPRB proteins. Our co-IP MS data suggest WRPA-WPRB interact,

and previous work has shown that *A. thaliana* WEB1 interacts with and PMI2

and itself (Kodama et al., 2010). We performed yeast 2-hybrid assays to

determine if maize WPR proteins also form homo/heterodimers, and found

several hetero- and homo-dimer combinations **(Figure 2C and 2D)**. This

includes heterodimers between WPRA and WPRB proteins. This potentially

explains why the Co-IP/MS data indicate that PAN proteins interact with both

WPRA and WPRB proteins, yet no direct interaction between WPRA and PAN

proteins was observed. Taken together, the Co-IP/MS combined with Y2H

assays demonstrated an interaction network **(Figure 2F).** We suggest that

WPRB could be mediating a WPRA-WPRB heterodimer interaction with both

PAN1 and PAN2 **(Figure 2G)**.

WPR Proteins Act after PAN2 but before PAN1 in the BRK-PAN-ROP

Pathway

Grass stomatal development occurs sequentially, where cells closest to

the leaf base are developmentally less advanced, and cells more distal from

the leaf base are more developmentally advanced. Therefore, in a single leaf,

all stages of development can be observed **(Supplemental Figure 1A-F).**

SMCs are initially unpolarized. As development proceeds, proteins important

for cell polarization become enriched at the GMC-SMC contact site, in

sequence. Previous studies have demonstrated that BRK-PAN-ROP pathway

components polarize sequentially in SMCs the following order: BRK1, PAN2,

PAN1, followed by formation of a cortical F-actin patch at the GMC contact

site (Cartwright et al., 2009; Zhang et al, 2012; Facette et al., 2015;

Supplemental Figure 1G). To determine the relative timing of WPR protein

polarization, we examined SMCs in plants co-expressing either CFP-WPRA2

or RFP-WPRB2 with BRK1-CFP, PAN2-YFP, PAN1-YFP, or the F-actin

marker FABD2-YFP **(Figure 3).** First, we compared the relative timing of the

two WPR proteins (**Figure 3A**). In plants co-expressing CFP-WPRA2 and

RFP-WPRB2, we observed 276 polarized SMCs, and in all cells CFP-WPRA2

and RFP-WPRB2 polarized at the GMC-SMC interface at the same time. We

never observed polarization of one protein without the other. This is consistent

with WPRA and WPRB proteins acting as a heterodimer. Based on this

observation, we infer that polarization of either WPR protein at the GMC-SMC

interface is likely indicative of the other.

Next, we compared BRK1-CFP, the earliest known marker of polarity,

and RFP-WPRB2. In plants co-expressing BRK1-CFP and RFP-WPRB2, we

frequently saw that BRK1-CFP had become polarized while RFP-WPRB2 had

not yet become polarized (94/232 cells). Arrows in **Figure 3B** indicate cells

where BRK1 is polarized, but WPRB2 is not. Reciprocally, we never saw cells

in which RFP-WPRB2 was polarized and BRK1-CFP was not polarized. This

indicates that BRK1-CFP polarizes prior to RFP-WPRB2, and is consistent

with previous findings that BRK1 is the earliest known marker of polarity

(Figure 3B). When CFP-WPRA2 was co-expressed with either PAN2-YFP

Figure 3C) or PAN1-YFP **(Figure 3D)**, SMCs always showed co-polarization

at the GMC contact site. The relative timing of WPR proteins and PAN1

receptors cannot be resolved from co-localization; our data indicates they

polarizes at the same time. Accumulation of F-actin at the GMC-SMC

interface is the last known marker of polarity. In SMCs co-expressing CFP-

WPRA2 and FABD2-YFP, we could never see polarized accumulation of

FABD2-YFP when CFP-WPRA2 was not polarized. However, we could see

polarized accumulation of CFP-WRPA2 when actin had not yet accumulated

in 67/140 cells, **(Figure 3E)**, suggesting WPRA2 polarizes prior to the actin

patch in developing SMCs. Thus, WPR proteins polarize after BRK1-CFP, but

prior to actin.

Co-expression of WPR proteins and PAN proteins indicate they

polarize at (or very close to) the same time. To investigate the possible

dependence of WPR polarization on the functions of known BRK-ROP-PAN

pathway components, we examined the polarization of CFP-WPRA2 and

RFP-WPRB2 in *brk1,pan1* and *pan2* mutants **(Figure 4)**. Homozygous mutant

plants were compared to heterozygous siblings that were grown in parallel.

Maize transgenics are generated in a hybrid background, and different inbred

lines have a large amount of genotypic variation which may persist after

backcrossing. By comparing mutants to phenotypically wildtype, heterozygous

sibling plants, background genetic variation is accounted for, and any

unexpected silencing of the transgene is also controlled for.

Previously, it was shown that in *brk1* mutants, PAN1 and PAN2 do not

polarize (Facette et al., 2015). Similarly, PAN1 does not polarize in a PAN2

mutant (Zhang et al., 2012). In *brk1* mutants, both CFP-WPRA2 and RFP-

WPRB2 lost their polarized accumulation at the SMC-GMC interface (Figure

4A and 4B). This is consistent with the co-localization data, and indicates that

WPRs act downstream of BRK1. Polarized localization of both CFP-WPRA2

and RFP-WPRB2 at the GMC contact site of SMCs is completely lost

in *pan2* mutants **(Figure 4C and 4D).** In contrast, both WPR proteins remain

polarized in *pan1* mutants **(Figure 4E and 4F)**. Thus, although we observed

no timing difference in the polarized accumulation of PAN2, PAN1 and WPR

proteins, PAN2 acts genetically upstream of WPRs and promotes their

polarized accumulation. However, PAN1, which acts downstream of PAN2, is

not required for WPR polarization.

CRISPR/Cas9-Mediated Knockout of *Wprb1* and *Wprb2* Causes

Subsidiary Cell Defects

To determine whether WPRs are required for asymmetric cell division

during maize stomatal development, we used CRISPR/Cas9 to generate *wpr*

mutants. The coding sequences of *Wpra1* and *Wpra2* are 92% identical,

therefore both genes were targeted using the same two guide RNAs. *Wprb1*

and *Wprb2* are 57% identical, and therefore two different guide RNAs were

used in a single construct to separately target the two genes. Single mutants

were identified for each gene **(Supplemental Figure 5A,B,C).** The edited

lines were backcrossed with B73 and the transgene was segregated away.

Single mutants were then crossed to obtain true-breeding double mutant

lines. Despite PCR and sequence-based genotyping of many hundreds of

plants from several different crosses, we never recovered *wpra1;wpra2*

double homozygous mutants (**Supplemental Figure 5D**). This might reflect

an essential shared function for the two genes in the male gametophyte,

based on the high expression level of *wpra1* and *wpra2* in maize pollen

(**Supplemental Figure 5E**). We were, however, able to generate

wprb1;wprb2 double homozygous mutants by crossing *wprb1/wprb1;wprb2/+*

x *wprb1/+;wprb2/wprb2* plants (which generate double mutants at a higher

frequency than a selfed double heterozygote). No stomatal phenotypes were

observed in *wprb1* or *wprb2* single homozygotes, however

wprb1/wprb1;wprb2/wprb2 double mutants have an increased percentage of

defective subsidiary cells relative to *wprb1/+;wprb2/+, wprb1/b1;wprb2/+ and*

wprb1/+;wprb2/b2 siblings (**Figure 5**). The abnormally shaped subsidiary cells

are phenotypically similar to those found in *brk1, pan1* and *pan2* mutants

(**Supplemental Figure 1H-K**; (Frank et al., 2003; Facette et al., 2015;

Cartwright et al., 2009; Zhang et al., 2012). Stomatal density was unaffected

in *wprb1/wprb1;wprb2/wprb2* mutants (**Supplemental Figure 6**). Since the

aberrant subsidiary cell phenotype of *wprb1;wprb2* double mutants is mild

relative to *pan1* or *pan2* mutants, we wanted to know if *wprb* mutants would

enhance *pan2* phenotypes, similar to what has been observed in *rop2/4/9*

mutants in combination with *pan1* (Humphries et al., 2011). However, we

found that *wprb* mutations do not enhance the *pan2* phenotype

(**Supplemental Figure 7**).We also examined *wprb1;wprb2* mutants for

nuclear polarization and formation of the actin patch, two landmarks of polarity

in SMCs (**Supplemental Figure 8**). No differences in nuclear polarization or

actin patch formation were seen in *wprb1;wprb2* mutants. This may be

because of the relatively mild defect or redundancy within the family. Higher

order mutants, such as with the closely related *wprb3* mutant, could be

informative.

In addition to loss-of-function phenotypes, we examined our native

promoter-fluorescent protein fusion transgenics to determine if mild

overexpression, resulting from co-expression of these FPs in a wild-type

background with endogenous *Wpr* genes, would cause defective stomatal

phenotypes. Plants expressing RFP-WPRB2 or CFP-WPRA2 in B73 indeed

had mild subsidiary cell defects, further supporting that WPR proteins have a

function in subsidiary cell division (**Figure 5 D, E**). These results indicate that

both WPRA and WPRB family members participate in stomatal development,

and suggest that WPRA1 and WPRA2 are important for the viability or

function of male gametophytes.

WPRB Proteins Directly Interact with F-actin

Since WPR proteins contain no discernable domains other than

DUF827, their molecular role is unclear. We hypothesized that WPRs might

interact with F-actin, based on the importance of actin in SMC polarization,

and because *A. thaliana* WEB1 and PMI2 have roles related to actin-based

regulated chloroplast movement (Kodama et al., 2010). Although we did not

see obvious co-localization of WPR proteins with actin filaments in maize,

membrane localization has been observed for other known actin-associated

proteins such as BRK1 and WAL (Facette et al., 2015; Sugiyama et al., 2019).

When overexpressed in tobacco cells, WAL localizes to actin cables (as

opposed to membranes). To determine whether overexpressed WPRs

similarly co-localize with F-actin *in vivo*, we transiently overexpressed GFP-

tagged WPRA2, WPRB1, WPRB2 and WPRB3 proteins in tobacco epidermal

cells. GFP-WPRA2 localized exclusively at the cell periphery/membrane

(Figure 6C). However, GFP-WPRB1 **(Figure 6D)**, GFP-WPRB2 **(Figure 6E)**

and GFP-WPRB3 **(Figure 6F)** were observed in filament-like structures, and

possibly also the plasma membrane. When GFP-WPRB2 is co-expressed

with Lifeact-RFP (actin filament marker), the WPRB2-labeled filamentous

structures extensively overlapped with Lifeact-RFP signals, particularly in the

thick actin cables **(Figure 6L to 6N)**. We treated GFP-WPRB2 expressing

tobacco leaves with Latrunculin B, an inhibitor of actin polymerization. The

GFP-WPRB2 filamentous structures were disrupted, but remained intact in

DMSO control **(Figure 6O and 6P, Supplemental Figure 9)**. Thus, WPRB2

is an F-actin associated protein.

WPR proteins do not show significant sequence similarity to other known

actin-associated proteins. To identify which region of the WPRB2 protein is

responsible for its localization to filamentous structures, various WPRB2

fragments were used for transient tobacco expression assays **(Figure 6A)**.

Surprisingly, the central conserved DUF827 domain is not responsible for

filamentous localization. Instead, we found that the N-terminal 67 amino acids

of WPRB2 is sufficient for filamentous localization in tobacco cells **(Figure**

6J). Longer fragments containing the N-terminal 67 amino acids of WPRB2

were also found in filamentous structures, including GFP-WPRB2ΔC **(Figure**

6G) and GFP-WPRB2N142 **(Figure 6K)**. Any construct lacking the N-

terminal domain does not label actin filaments, such as GFP-WPRB2ΔN

(Figure 6H) and GFP-WPR B2ΔNC **(Figure 6I)**. We obtained similar results

using full-length and truncated CFP-tagged constructs that were fused at C-

terminus (rather than N-terminus) of WPRB2 **(Supplemental Figure 10)**.

Conserved residues in the N-terminus of DUF827 proteins are highlighted in

Supplemental Figure 11. Together, these data confirm that the N-terminal 67

amino acids of WPRB2 are necessary and sufficient for its filamentous

localization.

To ask whether WPRB2 directly interacts with F-actin, we performed *in*

vitro actin co-sedimentation assays. We used the slightly larger WPRB2N240

fragment, as the smaller WPRB2N67 fragment is similar in size to actin and

made interpretation of the results difficult. WPRB2N240 is soluble in the

absence of F-actin, but can be found in the pellet in the presence of F-actin

(Figure 6Q). The amount of WPRB2N240 appearing in the pellet is

concentration dependent. Together, our localization and co-sedimentation

experiments suggest that WPRB2 directly binds to F-actin via its N-terminal

domain.

Physical interactions of WPR proteins with both PAN receptors and the

actin cytoskeleton link together previously identified components of the BRK-

PAN-ROP pathway. Potentially, WPRs may regulate actin polymerization,

stability, or organization during some step of SMC polarization. Future studies

will help resolve the relationship between WPRB proteins and actin.

Intriguingly, during co-localization experiments in stably transformed maize

plants described earlier (**Figure 3**), we noticed that the fluorescence intensity

of the actin marker FABD2-YFP was reduced in RFP-WPRB2 co-expressing

plants compared with RFP negative siblings, in all cell types. **Figure 7** shows

the YFP channel in a plant expressing FABD2-YFP alone (**Figure 7Ai & Aii**),

versus a sibling plant co-expressing both FABD2-YFP and WPRB2-RFP

(**Figure 7Bi & Bii**), using identical acquisition settings. Fluorescence is barely

visible in the co-expressing plants when the images are scaled at a range that

is appropriate for the ABD2-YFP alone plants (Figure 7Ai vs.Bi). Reciprocally,

when the YFP-channel is scaled appropriately for the co-expressing plants,

the image is overexposed (Figure 7Aii vs.Bii).WPRB2-RFP intensity levels

were similar in all sibling plants. We quantified the fluorescence in SMCs, and

found decreased FABD2-YFP fluorescence in at the GMC contact site as well

as at the other site was dramatically reduced, both at the polarized site as well

as at the lateral side of the SMC (**Figure 7H**). The ratio of intensities indicates

the polarized site is more affected (**Figure 7I**). This could be because of a

critical function of WPRB2-RFP in promoting actin accumulation at the

polarized site, rendering it more sensitive. It may be that more of an effect is

seen at the polarized site, however, because more WPRB2-RFP accumulates

there. Western blotting indicates that the overall actin levels are unchanged in

RFP-WPRB2 expressing cells (**Supplemental Figure 12**). This suggests that

the observed decrease in FABD2-YFP fluorescence could have a negative

effects on actin stability, or alters actin in a way such that FABD2-YFP cannot

bind as effectively.

Our previous data indicates that expression of either CFP-WRPA2 or

RFP-WPRB2 results in a low frequency of abnormal subsidiary cells (**Figure

5**), yet only WPRB (and not WPRA) proteins bind F-actin (**Figure 6**). We

quantified whether CFP-WPRA2, like WPRB2, had an effect on FABD2-YFP

intensity (**Supplemental Figure 13**). No similar decrease in FABD2-YFP

intensity was observed in cells expressing CFP-WRPA2, consistent with the

observation that only WPRB, and not WPRA, proteins bind F-actin.

Since actin is important in SMCs in early polarization, prior to WPR

polarization (e.g., BRK proteins) and in late steps after WPR polarization (e.g.

formation of the actin patch), we wanted to know if polarization of WPRB2 is

maintained or stabilized by actin. We treated maize leaves expressing either

RFP-WPRB2 or FABD2-YFP with Latrunculin B. After 4 hours, 40 µm

Latrunculin B treatment led to efficient depolymerization of actin in SMCs,

including the dense patches of actin at the SMC-GMC interface

(Supplemental Figure 14). However, disruption of actin does not affect the

localization of RFP-WPRB2. This suggests that WPRB2 is not dependent on

actin networks for maintaining its polarization. Together, these results suggest

that WPRB2 interacts with actin filaments in a manner that de-stabilizes them,

at least when present in excess of normal amounts. This offers a potential

explanation for how WPRs may impact actin-dependent processes in

polarizing SMCs.

DISCUSSION

We demonstrated that WPR proteins function in the BRK-PAN-ROP

pathway for subsidiary cell formation during maize stomatal development, and

physically interact with the receptor proteins PAN2 and PAN1. Like other

proteins in the BRK-PAN-ROP pathway, WPRs polarize in SMCs at the site of

GMC contact. The polarized accumulation of WPRs in SMC depends on

PAN2, and genetic evidence demonstrates a function for WPRs in promoting

normal SMC divisions. Like WEB1 and PMI2 in *A. thaliana*, different WPR

subfamily members form heterodimers, suggesting WPR subfamily members

co-operate to fulfill their function in different cellular contexts (Kodama et al.,

2010).

Our analyses indicate that WPR proteins act downstream of PAN2, but

upstream of PAN1, yet physically interact with both PAN1 and PAN2. This is

the first observed physical link between PAN2 and PAN1, and offers a

hypothesis of how PAN2 may promote polarization of PAN1, i.e. indirectly via

WPR recruitment. We predict that in *wpr* mutants, PAN1 will no longer

become polarized; however gene redundancy within the WPR family makes

this a non-trivial question to address. Future experiments with high-order *wpr*

mutants and PAN1-YFP will further elucidate the function of WPRs in the

pathway the pathway. Even though WPRA2 and WPRB2 no longer polarize in

pan2 mutants, they still localize to the plasma membrane, indicating that

membrane localization is independent of PAN2.

Our data indicate that both WPRA and WPRB proteins localize

predominantly to the plasma membrane, and may also have cytoplasmic

localization. Similar to the BRK1, which is important for acting regulation,

WPR proteins also localize to cell corners (Facette et al., 2015). Previous

studies using transient protoplast expression of WEB1 and PMI2 indicate

WEB1 localizes to the plasma membrane, and can recruit PMI2 there

(Kodoma et al., 2010). An independent proteomic study indicated that WEB1

is predominantly localized to the cytoplasm but re-localizes to the membrane

(and its phosphorylation status changes) upon blue light irradiation (Deng et

al., 2014). No DUF827 family members have predicted transmembrane

domains, indicating their membrane association is either peripheral or via

interactions with membrane-localized proteins, or perhaps via membrane-

associated actin networks. Besides WEB1 and PMI2, the other characterized

DUF827 protein is TREPH1 (Wang et al., 2018). TREPH1 and maize WPRA2

are orthologues **(Supplemental Figure 2)**. A specific residue in the C-

terminus of TREPH1 is phosphorylated in response to touch, and the

phenotype of the *treph1* mutant is dependent on the phosphorylation status of

this residue (Wang et al., 2018). However, the same residue in maize WPRA2

is not present. If all WPR proteins are similarly phosphoregulated, this

suggests that other active, as of yet undiscovered kinases may be important

for WPR regulation, since both PAN1 and PAN2 are pseudokinases.

Our analysis of CRISPR-Cas9 induced mutants of WPRs

demonstrated that WPRs have diverse functions in maize. We were unable to

recover *wpra1/wpra1*;*wpra2/wpra2* double mutants. WPRA1 and WPRA2 are

89% similar in protein sequence, and are highly expressed in pollen. CFP-

WPRA2 polarly localizes to the subapex of maize pollen tubes

(Supplemental Figure 5), raising the possibility that WPRA proteins play a

role in pollen function. Additionally, we found *wprb1;wprb2* double mutants

affect subsidiary cell formation, albeit not to the same extent as *pan* mutants.

This may be caused by functional redundancy with other *Wpr* genes, such as

Wprb3.

Our discovery that WPRB proteins directly bind actin is significant as it

physically links them to actin network function. WPR proteins do not share

any sequence homology to known actin binding proteins, however several

DUF827 proteins were predicted, based on structural homology, to potentially

bind the cytoskeleton (Gardiner et al., 2011). Surprisingly, the central

conserved DUF827 domain is not the actin-binding domain. We mapped a 67

amino acid sequence that is necessary and sufficient for actin-binding,

thereby identifying a putative novel actin binding domain. Within this

sequence, there appears to be a motif that is conserved in many, but not all

DUF827 proteins **(Supplemental Figure 8)**. Even though WPRA2 also

contains most of this conserved domain, we did not observe co-localization of

WPRA2 to the actin cytoskeleton. This could be because the transient

expression does not reflect *in vivo* function. Alternatively, small differences

between WPRA2 vs. B2 sequence may account for its lack of actin binding

(e.g, one of a pair of glycines conserved in many WPR proteins is aspartate in

WPRA2). Further refining the region of interaction, and determining if other

DUF827 proteins bind actin, would inform their function.

The significance of WPR interaction with the actin cytoskeleton is

unknown, and is complicated by the fact that there are several actin-related

processes that occur during SMC polarization. The most plausible are: a

function related to the SCAR/WAVE complex that includes BRK1, which

polarizes and functions very early to promote SMC polarization; a role in

localization of polarity determinants (such as vesicle trafficking); a role related

to actin patch formation or function; or a role related to actin-based nuclear

migration. These possibilities are not mutually exclusive. Actin filaments can

facilitate polarized protein localization. For example, polarization of PIN

requires an intact actin cytoskeleton (Kleine et al., 2008). It is plausible that

actin networks are responsible for the polarization of PAN1 and/or PAN2, or

even the WPR proteins themselves. This function would be distinct from that

of the actin patch, which appears long after PAN2, PAN1, or WPR proteins

polarize. When actin patches were disrupted with Latrunculin B, this did not

result in the loss of already polarized RFP-WPRB2 in maize leaves,

suggesting the actin patch does not influence WPR protein localization (and

possibly function). A clue to what WPR proteins could be doing *in vivo* can be

inferred from our observation that RFP-WPRB2 transgenic plants WPRB

displayed less F-actin than controls. This suggests WPRB2 negatively affects

the stability of actin filaments, and might promote F-actin depolymerization.

Alternatively, WPRB proteins could compete with ABD2 for binding sites on

the F-actin polymer, or somehow modify F-actin such that ABD2 does not

bind as effectively. Future *in vitro* and *in vivo* work characterizing the

molecular role of WPR proteins on actin dynamics, including the nucleation,

severing and bundling activities will help determine the function of this gene

family – not only in cell polarization, but also during mechanosensing (i.e.,

TREPH1 function) or organelle movements (i.e. WEB1 and PMI2 function).

METHODS

Plant Materials and Plant Growth Condition

BRK1-CFP, PAN2-YFP, PAN1- YFP, and YFP-ABD2-YFP stable transgenic

lines, and *brk1*, *pan1* and *pan2* mutants were described previously (Cartwright

et al., 2009, Zhang et al., 2012; Facette et al., 2015). Plants used for analysis

were grown for 10d to 14d in a greenhouse maintained between 20°C and

29°C. Plants were grown in Pro-mix Professional soil supplemented with .

Supplemental LED lights (Fluence VYPR series) were used to maintain a 16-

hour day length.

Generation of transgenic maize lines

Transgenic lines expressing CFP-WPRA2 or RFP-WPRB2 were created

using a genomic construct. Primers are listed in **Supplemental Table 2.** An

approximately 3kb sequence upstream of the start codon (promoter and 5'

UTR) was amplified from B73 genomic DNA with primers WPRA2-F &

WPRA2-P2 or WPRB2-F & WPRB2-P2. Genomic DNA, including all exons

and introns 1kb after the stop codon (3' UTR), was amplified from B73

genomic DNA using primers WPRA2-P3 & WPRA2-R, or WPRB2-P3 &

WPRB2-R. CFP or RFP was amplified with primer Tag linker-F and Tag

linker-R as described previously (Mohanty et al., 2009). N-terminal fluorescent

protein fusions were assembled through fusion PCR and cloned into

pDONR221. Gateway LR reactions (Thermo Fisher) were used to recombine

the assembled fusion protein into the binary vector pAM1006 (Monhanty et

al., 2009). The constructions were verified and introduced into Hi-II maize

via *Agrobacterium tumefaciens*–mediated transformation by Wisconsin Crop

Innovation Center plant transformation facility. Primary transformants were

crossed to B73 to produce T1 progeny used for immunoprecipitation

experiments, and T1s were crossed with BRK1-CFP, PAN2-YFP, PAN1- YFP,

and YFP-ABD2-YFP to produce T2 progeny used for the imaging

experiments. Primary transformants were also crossed to *brk1*, *pan1* and

pan2 mutants, and the T1 progeny backcrossed again to mutants, to produce

homozygous *brk1*, *pan1* and *pan2* mutants expressing CFP-WPRA2 and

RFP-WPRB2. In all cases, sibling plants grown simultaneously were used as

controls.

Yeast Two-Hybrid Analysis

The GAL4-based yeast two-hybrid interaction system was used for interaction

assays. The cDNA fragments encoding intracellular portions of PAN1 and

PAN2 were cloned into pASGW-attR plasmid as a bait via Gateway cloning as

described previously (Zhang et al., 2012). cDNA was prepared from the maize

leaf stomatal division zone. Full length cDNAs of *Wpra1, Wpra2, Wprb1,*

Wprb2 and *Wprb3* cDNA were amplified from a maize leaf division zone

cDNA preparation with primers WPRA1topo1-F + WPRA1topo-R (*Wpra1*),

WPRA2topo1-F + WPRA2topo-R (*Wpra2*), WPRB1topo1-F + WPRB1topo-

R(*Wprb1*) and WPRB2topo1-F + WPRB2topo-R (*Wprb2*) (**Supplemental**

Table 1). These fragments were cloned into the entry vector pENTR/D-TOPO

(Invitrogen) and further transferred destination vectors pASGW-attR (bait) and

pACTGW-attR (prey) (Nakayama et al., 2002). Bait and prey plasmids were

co-transformed into AH109 yeast strain. Yeast was grown on medium

selecting for diploid cells (SD -Leu-Trp) or detecting interactions (SD -Leu-

Trp-Ade-His) for two days at 30°C.

Confocal Microscopy

Plant tissues were observed using a custom-built spinning disc

confocal unit (3i) equipped with an inverted fluorescence microscope (IX83-

ZDC, Olympus) CSU-W1 spinning disc with 50 micron pinholes (Yokogawa),

a Mesa Illumination Enhancement 7 Field Flattening unit (3i), an Andor iXon

Life 888 EMCCD camera and a UPLANSAPO ×60 Silicone Oil-immersion

objective (NA = 1.20, Olympus) and 4 laser stack with TTL controller (3i). For

CFP, YFP and RFP imaging of transgenic maize plants, a 445/515/561

dichroic (Chroma) was used. All emission filters are from Semrock. A 445 nm

laser line and 483/45 emission filter (CFP), 514 nm laser and 542/27

emission filter (YFP) or 568 laser with 618/50 emission filter (RFP) was used.

For dual imaging of GFP and RFP (tobacco expression) a 405/488/561/640

dichoric (Chroma) was used, with a 488 nm laser and 525/30 emission filter

(GFP) and/or 568 laser with a 618/50 emission filter. Image processing was

performed using Image Fiji and Adobe Photoshop version 8.0 using only

linear adjustments and preserving hard edges.

Quantification of ABD2-YFP fluorescence intensity was performed

using FIJI. Max projections of 7 slices of 16-bit images taken on the same day

with the same acquisition settings were used. Sibling plants were used as

controls.

Transient expression in *N. benthamiana*

To generate 35S-expressed N-terminal GFP-tagged constructs (and

truncated derivatives) and C-terminal CFP-tagged constructs (and truncated

derivatives) for transient tobacco expression, coding sequence were PCR-

amplified from maize leaf cDNA with the appropriate primers (**Supplementary**

Table 1) and cloned into the pENTR/D-TOPO entry vector (Thermo

Scientific). Clones were recombined with the vectors pSITE-2CA and pSITE-

1NB using LR Clonase Mix II (Thermo Scientific) to form N-terminal GFP or C-

terminal CFP (Romit Chakrabarty, 2006).

For transient expression in *N. benthamiana*, *Agrobacterium* strain

GV3101 harboring different constructs were resuspended in infiltration buffer

(10 mM MES (pH 5.7), 10 mM MgCl2, 50 mg/L acetosyringone), and adjusted

OD600 to 1.0. Equal volumes of cultures containing different constructs were

mixed for co-infiltration. The resulting cultures were infiltrated into leaves of 3-

to 4-week-old N. *benthamiana* plants. Leaf samples were harvested 48 hours

after infiltration.

Generation and Purification of WPRA1/A2 and WPRB2-Specific Antibody

Peptides corresponding to amino acids 267 to 288 of maize WPRA1 and

WPRA2 (LRNDFDPAAYDSLKEKLEQTNS), and amino acids 446-465 of

WPRB2 (HPAPRSRDSQNMDIVGVSKGC) were synthesized, conjugated to

KLH, co-injected into rabbits and used for polyclonal antibody production in

rabbits by Pacific Immunology. And the resulting sera was affinity purified

against the corresponding peptide using columns provided by Pacific

Immunology. Affinity purified antibodies were eluted using Gentle Ag/Ab buffer

(Pierce), desalted against TBS Buffer, and concentrated to 1 mg/ml using a

30K MWCO concentrator (Pierce).

Maize Division Zone Membrane Protein Extraction, Western Blotting,

and Co-IP/MS Analysis

Protein extraction for western blotting of WPR proteins and co-IPs were

performed as described previously (Facette et al., 2015). Briefly, 10-14 day

old plants were used to isolate the basal 0.5-2.5 cm of unexpanded leaves 4-

6, the leaf bases were ground in liquid nitrogen. Membrane fractions of

extracts from these tissues were prepared. For western blotting, ~0.5 grams

(3-5 plants) of tissue was used and for mass spectrometry, 1.5 grams (8-10

plants) were used. per 0.25 g of grounded tissue was mixed with 1ml

extraction buffer (50mM Tris, pH 7.5, 150 mM NaCl, 5mM EGTA, 5mM EDTA

0.3% β-mercaptoethanol, 1% Sigma Plant Protease Inhibitor), the mixture was

homogenized for 10s using an Omnitip homogenizer on ice. All subsequent

steps were conducted at 4°C. The mixture was centrifuged twice at 13000 *xg*.

The supernatant was pelleted at 110,000xg for 45 min. Solubilization buffer

(50 mM Tris pH 7.5, 150 mM NaCl, 1% NP-40, 10% glycerol, 0.05% Sodium

deoxycholate) was then added to the pellet and following sonicated 3x15s and

left rotating 1 hour. For western blotting, extracts from B73 plants and

corresponded mutants were separated via SDS-PAGE and analyzed using

anti-WPRA1/A2 at 1 µg/ml.

For western blots to detect actin in B73 and RFP-WPRB2 transgenic

plants, the basal 0.5-2.5 cm of unexpanded leaves 4-6 from 14 days plant

were ground in liquid nitrogen. Groundtissue was mixed with 500 µL

extraction buffer (50mM Tris, pH 7.5, 150 mM NaCl, 5mM EGTA, 5mM EDTA

0.3% β-mercaptoethanol, 1% Sigma Plant Protease Inhibitor) and

homogenize immediately by Omnitip homogenizer for 10s on ice. Then, the

mixture was centrifuged at 15,000 x g for 10 min at 4 °C.

Transfer supernatant (protein extract) to a new tube and keep on ice for

immediate use. 100 ng extracted protein for each plants were separated via

SDS-PAGE and analyzed using anti-actin (Sigma A0480, MFCD00145889) at
1:1000.

Co-IP/MS experiments were performed using the solubilized
membrane proteins. Immunoprecipitation of PAN2-YFP was performed as in
Facette et al. (2015) using anti-GFP beads (Miltenyi). PAN2-YFP
immunoprecipitations and subsequent mass spectrometry runs were
performed in parallel with B73 (control), BRK1-CFP, PAN1-YFP, PDI-YFP,
PIN1-YFP and RAB11D-YFP using anti-GFP beads (Miltenyi). Note that these
co-immunoprecipitations, except PAN2-YFP and PAN1-YFP, were presented
previously in Facette et al. (2015) A WD-score (Sowa et al., 2009) was
calculated to determine high confidence interactors. Tryptic digestion and
mass spectrometry was performed using a Q-Exactive mass spectrometer
(Thermo Scientific, San Jose, CA) as described (Facette et al., 2015).

Anti-WPRA1/A2 and anti-RFP (Rabbit polyclonal, Invitrogen R10367,
Lot 2127438) were coupled with Dynabeads prepared according to the
Dynabeads kit instructions (Thermo Fisher). The coupled WPRA1/A2
antibody-beads complex was added to the membrane protein extracts from
B73 plants; uncoupled beads were used as a negative control. Anti-RFP
coupled beads were added to the membrane protein extracts from RFP-
WPRB2 transgenic plants, and B73 plants as a negative control. All these
samples were incubated rotating at room temperature for 30 min. Then the
Dynabeads-CO-IP complex was washed according to Dynabeads kit
instructions. After washing, the Dynabeads-CO-IP complex was digested with
trypsin and run with the mass spectrometer.

For Co-IP using anti-GFP to pull down WPRA2, anti-GFP beads
(Miltenyi) were added to the extracts from CFP-WPRA2 transgenic plants and
B73 plants (negative control), then the extracts were rotated for an additional
30 minutes. Miltenyi μ columns were equilibrated using a solubilization buffer
as described by instruction. The extracts were applied to the columns, the
columns were washed and eluted as Miltenyi μ columns kit instruction, the
eluted samples were digested with trypsin and run with the mass
spectrometer.

For all WPR immunoprecipitations, peptides were separated by

reverse-phase chromatography using nano-flow EASY-nLC 1000 UHPLC

coupled to Orbitrap Fusion mass spectrometer (Thermo Scientific) with

PepMap RSLCnano column (75 µm ID, 15 cm). Peptides were eluted over a

90 minute 5-35% ACN gradient at 300 nL/min. Survey scans were measured

in the Orbitrap analyzer at 60,000 resolution. Data-dependent MS/MS data

were collected in the linear ion trap using a 2-second cycle time with a full MS

mass range from 400-1800 m/z. Peptides (charge state 2-6) were fragmented

using higher energy collisional dissociation using a normalized collision

energy setting of 27. A dynamic exclusion time of 5 seconds was used and

the peptide match setting was enabled. RAW files were analyzed in

Proteome Discoverer 2.4 (Thermo Scientific) using the SEQUEST search

algorithm using version 4 of the maize genome (B73 RefGen_v4). The search

parameters used are as follows: 10 ppm precursor ion tolerance and 0.4 Da

fragment ion tolerance; up to two missed cleavages were allowed; dynamic

modifications of methionine oxidation and N-terminal acetylation. Peptide

matches were filtered to a protein false discovery rate of 5% using Percolator

algorithm. Peptides were assembled into proteins using maximum parsimony

and only unique and razor peptides were retained for subsequent analysis.

Genomic editing of maize WPRs by CRISPR–Cas9

A dual gRNA maize CRISPR/Cas9 system was used to generate *Wpr*

mutants (Char et al., 2017). Transgenic Hi-II maize plants were generated at

the Iowa State Plant Transformation facility, expressing Cas9 from

Streptococcus pyogenes under the maize UBI1 promotor and two guide RNAs

under the rice U6.1 and U6.2 promotor, respectively. To create *wpra1* and

wpra2 mutants, two tandem gRNAs, gRNA1: GAAACATCTTTGGATATG and

gRNA2: GTGCAAGCACACGAAGAAG were used to target identical

sequences within *Wpra1* and *Wpra*2. To create *wprb1* and *wprb2* mutants,

guide RNAs targeting *Wprb1* using gRNA1: GCTACATGTGATCTGGCTG or

Wprb2 using gRNA2: GCCTCCGTCGAGTCGCTG were used. Genomic

edits were verified by Sanger sequencing of the target regions. Plants

harboring edits were crossed to B73. Edited plants that no longer contain the
transgene were confirmed and used for further analyses.

Phalloidin staining

Phalloidin staining in maize leaves was performed as previously
described (Cartwright et al., 2009, Nan et al., 2019). The basal 0.5-2.5 cm of
leaf 4 from *wprb1/+;wprb2/+* and *wprb1/b1;wprb2/b2* mutants were fixed and
stained with Alexa fluor 488-phalloidin (Thermo Fisher). Nuclei and cell walls
were stained using 10 µg mL-1 propidium iodide (Thermo Fisher). Samples
were mounted in ProLong Gold Antifade (Thermo Fisher).

Immunofluorescent detection of WPR proteins

Immunolocalization of WPRA2 and WPRB2 were performed in leaf
tissue excised from the basal 1 to 3 cm of unexpanded leaves of 2-week-old
plants as described previously (Cartwright et al., 2009, Nan et al., 2019) using
affinity-purified anti-WPRA1/A2 or WPRB2 at 2 µg/ml. Secondary Alexa Flour
488-conjugated anti-rabbit (Invitrogen) were used at a dilution of 1:500. Nuclei
were stained with 10 µg/mL propidium iodide (Sigma-Aldrich) prior to
mounting in in ProLong Gold Antifade (Thermo Fisher).

Bacterial Expression of WPR proteins

The DNA sequence of WPRB2N (1-240), which was used in the actin
co-sedimentation assay, was amplified by PCR using the primers listed in
Supplemental Table 2 and cloned into pGEX4T-1. Fragments of WPRA1 (63-
503) WPRA2 (63-570) WPRB1 (73-421) and WPRB2 (78-429), which were
used to confirm the specificity of the WPRA1/WPRA2 antibody, were also
amplified by PCR using the primers listed in Supplemental Table 2 and cloned
into pGEX4T-1 (Novagen) using BamHI/EcoRI sites to N-terminal glutathione
S-transferase (GST) tag fusions. These constructions were transformed into
Escherichia coli BL21 host strains (Novagen). GST fused proteins were
induced with 0.5 mM isopropyl β-d-1-thiogalactopyranoside for 20 hours at
28°C. Affinity purified using glutathione-Sepharose 4B column (GE
Healthcare) was carried out as described (Harper and Speicher, 2011).

F-Actin Co-sedimentation Assay

WPRB2N240 was dialyzed overnight against buffer A3 (10 mM Tris-

HCl, 0.2 mM $CaCl_2$, 0.2 mM ATP, and 0.5 mM DTT, PH=7.0). Prior to use,

the protein was further purified by centrifugation at 100,000g for 1 hour at 4°C

and only soluble protein was used for the assay. A high-speed co-

sedimentation assay was performed as described previously (Xiang et al.,

2007), Briefly, 5 μM and 10 μM WPRB2N240 alone or mixed with 4 μM

preformed F-actin and incubated in 200 μL volume of buffer A3 for 1 hour at

25°C. The samples were centrifuged at 100,000g for 1 hour at 4°C. GST

protein was used as a negative control. The supernatants and pellets were

separated and subjected to SDS-PAGE and visualized by Coomassie Blue

staining.

Latrunculin B treatment

To transiently depolymerize actin microfilaments in maize leaves, leaf 4 of

two-week-old plants were excised and the basal 1 to 3 cm part was immersed

in 40 μM Lat B (20 mM stock in DMSO) for 4 hours at room temperature. To

examine the LatB effect on the localization of GFP-WPRB2 in tobacco leaves,

the GFP expressed tobacco leaves were cut into 0.5 cm×0.5 cm and covered

with 40 μM LatB for 2 hours at room temperature.

Protein tree construction

Amino acid sequences of maize WPR homologs were identified using

pBLAST against maize genomes. The protein tree was constructed in

Molecular Evolutionary Genetics Analysis 5 (MEGA 5) software (Tamura et

al., 2011) with the following parameters: multiple amino acids sequence

alignment with ClustalW, phylogenetic construction with the maximum

likelihood method, and bootstrap tests of 1,000 replicates. WPRs sequences

used for the phylogenetic analysis are provided in **Supplementary File 1**.

Statistical Tests

T-tests and one-way ANOVA test were performed in JMP (SAS). Chi-squared

test was performed using an online calculator released by GraphPad Prism

(https://www.graphpad.com/quickcalcs/chisquared1/) with expected values

calculated by hand.

Accession Numbers

Gene and protein sequences can be found at MaizeGDB www.maizegdb.org .

Accession numbers for Version 4.0/Version 5.0 of the B73 genome are:

Wpra1: Zm00001d023629/Zm00001eb408590; *Wpra2*: Zm00001d041088/

Zm00001eb133280; *Wprb1:* Zm00001d0475516/Zm00001eb395070; *Wprb2:*

Zm00001d007164/Zm00001eb111490; *Wprb3:* and Zm00001d010610/

Zm00001eb351980; *Brk1:* no v4.0/ Zm00001eb259430; *Pan2:*

Zm00001d007862/Zm00001eb117610; *Pan1*: Zm00001d031437/

Zm00001eb034900.

ACKNOWLEDGEMENTS

We thank Steve Eyles at the UMass Institute of Applied Life Sciences Mass
Spectrometry Facility, RRID:SCR_019063. We thank Chris Phillips, Bob
Skalbite, Neal Woodward, and other greenhouse and farm staff at the
University of Massachusetts Amherst for essential help. This work was
supported by a grant from NSF (IOS-1754665) to MRF and NIH (DP2-
GM119132) to EJB.

AUTHOR CONTRIBUTIONS

M.F. and Q.N. designed and performed the research, analyzed the data and
wrote the manuscript. B.Y. (UCSD) and E.J.B. performed the mass
spectrometry for the initial set of co-IPs. S.N.C. and B.Y.(MO) designed the
WPRs CRISPR sgRNAs and prepared the constructs for maize
transformation.

REFERENCES

Apostolakos, B.G.P. (2004). The role of the cytoskeleton in the morphogenesis and function of stomatal complexes. New Phytol. **161,** 613–639.

Apostolakos, P., P. Livanos, E. Giannoutsou, E. Panteris, and Galatis B. (2018) The intracellular and intercellular cross-talk during subsidiary cell formation in Zea mays: Existing and novel components orchestrating cell polarization and asymmetric division. Annals of Botany **122**, 679-696.

Ashraf, M.A., and Facette, M. (2020). Plant Biology: BASL Gives the Plant Nucleus a Sense of Direction. Curr. Biol. **30,** R1375-R1377.

Campos, R., Goff, J., Rodriguez-Furlan, C., and Van Norman, J.M. (2020). The Arabidopsis Receptor Kinase IRK Is Polarized and Represses Specific Cell Divisions in Roots. Dev. Cell **52,** 183-195.

Cartwright, H.N., Humphries, J.A., and Smith, L.G. (2009). PAN1: a receptor-like protein that promotes polarization of an asymmetric cell division in maize. Science **323,** 649-651.

Char, S. N., Neelakandan, A. K., Nahampun, H., Frame, B., Main, M., Spalding, M. H., Becraft, P. W., Meyers, B. C., Walbot, V., Wang, K., & Yang, B. (2017). An Agrobacterium-delivered CRISPR/Cas9 system for high-frequency targeted mutagenesis in maize. Plant Biotechnology Journal *15*, 257–268.

Deng, Z., Oses-Prieto, J.A., Kutschera, U., Tseng, T.S., Hao, L., Burlingame, A.L., Wang, Z.Y., and Briggs, W.R. (2014). Blue light-induced   proteomic changes in etiolated Arabidopsis seedlings. J. Proteome Res. **13,** 2524-2533.

De Smet I., and Beeckman T. (2011). Asymmetric cell division in land plants and algae: the driving force for differentiation. Nat. Rev. Mol. Cell Biol. **12,** 177– 188.

Dong, J., MacAlister, C.A., and Bergmann, D.C. (2009). BASL controls asymmetric cell division in Arabidopsis. Cell **137,** 1320-1330.

Facette, M.R., and Smith, L.G. (2012). Division polarity in developing stomata. Curr. Plant Biol. **15,** 585-592.

Facette M.R., Park Y., Sutimantanapi D., Luo A., Cartwright H., Yang, B., Bennett E.J., Sylvester A.W., and Smith L.G. (2015). The SCAR/WAVE complex polarizes PAN receptors and promotes division asymmetry in maize. Nat. Plants **1,** 14024.

Facette, M.R., Rasmussen, C.G., and Van Norman, J.M. (2019). A plane choice: coordinating timing and orientation of cell division during plant development. Curr. Plant Biol. **47,** 47-55.

Frank, M. J. and Smith, L. G. (2002). A small, novel protein highly conserved in plants and animals promotes the polarized growth and division of maize leaf epidermal cells. Curr. Boil. **12,** 849-853.

Galatis B., and Apostolakos P. (2004). The role of the cytoskeleton in the morphogenesis and function of stomatal complexes. New Phytol. **161,** 613–639.

Gardiner, J., Overall, R., and Marc, J. (2011). Putative Arabidopsis homologues of metazoan coiled-coil cytoskeletal proteins. Cell Biol. Int. **35,** 767-774.

Gray, A., Liu, L., and Facette, M. (2020). Flanking Support: How Subsidiary Cells Contribute to Stomatal Form and Function. Front. Plant Sci. **11,** 881.

Gu, Y., Fu, Y., Dowd, P., Li, S., Vernoud, V., Gilroy, S., and Yang, Z. (2005). A Rho family GTPase controls actin dynamics and tip growth via two counteracting downstream pathways in pollen tubes. J. Cell Biol. **169,** 127-138.

Hadley, R., Hable, W.E., and Kropf, D.L. (2006). Polarization of the endomembrane system is an early event in fucoid zygote development. BMC Plant Boil. **6,** 5.

Harper S, Speicher DW. (2011) Purification of proteins fused to glutathione S-transferase. Methods Mol Biol. **681,** 259–80.

Houbaert, A., Zhang, C., Tiwari, M., Wang, K., de Marcos Serrano, A., Savatin, D.V., Urs, M.J., Zhiponova, M.K., Gudesblat, G.E., Vanhoutte, I., Eeckhout, D., Boeren, S., Karimi, M., Betti, C., Jacobs, T., Fenoll, C., Mena, M., de Vries, S., De Jaeger, G., and Russinova, E. (2018). POLAR-guided signalling complex assembly and localization drive asymmetric cell division. Nature **563,** 574-578.

Humphries, J.A., Vejlupkova, Z., Luo, A., Meeley, R.B., Sylvester, A.W., Fowler, J.E., and Smith, L.G. (2011). ROP GTPases act with the receptor- like protein PAN1 to polarize asymmetric cell division in maize. Plant cell **23,** 2273-2284.

Kennard J. L., Cleary A. L. (1997). Pre-Mitotic Nuclear Migration in Subsidiary Mother Cells of Occurs in G1 of the Cell Cycle and Requires F-Actin. Cell Motil. Cytoskelet. **36,** 55-67.

Kleine-Vehn, J., Langowski, L., Wisniewska, J., Dhonukshe, P., Brewer, P.B., and Friml, J. (2008). Cellular and molecular requirements for polar PIN targeting and transcytosis in plants. Mol. plant **1,** 1056-1066.

Kimata, Y., Higaki, T., Kawashima, T., Kurihara, D., Sato, Y., Yamada, T., Hasezawa, S., Berger, F., Higashiyama, T., and Ueda, M. (2016). Cytoskeleton dynamics control the first asymmetric cell division in Arabidopsis zygote. Proc Natl Acad Sci USA **113,** 14157-14162.

Kodama, Y., Suetsugu, N., and Wada, M. (2011). Novel protein-protein interaction family proteins involved in chloroplast movement response. Plant Signal. Behav. **6,** 483-490.

Kodama, Y., Suetsugu, N., Kong, S.G., and Wada, M. (2010). Two interacting coiled-coil proteins, WEB1 and PMI2, maintain the chloroplast photorelocation movement velocity in Arabidopsis. Proc. Natl. Acad. Sci. USA **107,** 19591-19596.

Larkin, M.A., Blackshields, G., Brown, N.P., Chenna, R., McGettigan, P.A.,   McWilliam, H., Valentin F., Wallace, I.M., Wilm A., Lopez, R., Thompson, J.D., and Gibson, T.J. (2007) Clustal W and Clustal X version 2.0. Bioinformatics, **23,** 2947–2948.

Lau, O.S., and Bergmann, D.C. (2012). Stomatal development: a plant's perspective on cell polarity, cell fate transitions and intercellular communication. Development **139,** 3683-3692.

Lee JS, Kuroha T, Hnilova M, Khatayevich D, Kanaoka MM, McAbee JM, Sarikaya M, Tamerler C, Torii KU. (2012). Direct interaction of

951    ligand–receptor pairs specifying stomatal patterning. *Genes Dev* **26,**
952     126–136.
953 **Lipka, E., Herrmann, A., and Mueller, S.** (2015). Mechanisms of plant cell
954     division. Wiley interdisciplinary reviews. Dev. Biol. **4,** 391-405.
955 **Luesse, D.R., DeBlasio, S.L., and Hangarter, R.P.** (2006). Plastid
956     movement impaired 2, a new gene involved in normal blue-light-
957     induced chloroplast movements in Arabidopsis. Plant Physiol. **141,**
958 1328-1337.
959 **Mohanty, A., Luo, A., DeBlasio, S., Ling, X., Yang, Y., Tuthill, D.E.,**
960     **Williams, K.E., Hill, D., Zadrozny, T., Chan, A., Sylvester, A.W.,**
961     **and Jackson, D.** (2009). Advancing cell biology and functional
962     genomics in maize using fluorescent protein-tagged lines. Plant
963     Physiol. **149,** 601-605.
964 **Muroyama, A., Gong, Y., and Bergmann, D.C.** (2020). Opposing, Polarity-
965     Driven Nuclear Migrations Underpin Asymmetric Divisions to Pattern
966     Arabidopsis Stomata. Curr. Biol. **30,** 4467-4475
967 **Nakayama M., Kikuno R., Ohara O.** (2002). Protein-protein interactions
968     between large proteins: Two-hybrid screening using a functionally
969     classified library composed of long cDNAs. Genome Res. **12,** 1773–
970     1784.
971 **Nan Q. Mendoza, J. and Facette M.** (2019). Double Labeling of Microtubules
972     and Actin Filaments in Maize Leaf Division Zone. BIO-PROTOCOL **9:**
973     e3262-e3262.
974
975 **Nunes, T.D.G., D. Zhang, and M.T. Raissig.** (2020). Form, development and
976     function of grass stomata. Plant J. **101,** 780-799
977 **Pillitteri, L.J., Peterson, K.M., Horst, R.J., and Torii, K.U.** (2011). Molecular
978     profiling of stomatal meristemoids reveals new component of
979     asymmetric cell division and commonalities among stem cell
980     populations in Arabidopsis. Plant Cell **23,** 3260 – 3275.
981 **Pillitteri, L.J., and Torii, K.U. (2012).** Mechanisms of stomatal development.
982     Annu. Rev. Plant Biol. **63,** 591 – 614.
983 **Romit Chakrabarty, R.B., Sang-Min Chung, Mark Farman, Vitaly**
984     **Citovsky, Saskia A. Hogenhout, Tzvi Tzfira, and Michael**
985     **Goodin**   (2006). pSITE Vectors for Stable Integration or Transient
986     Expression of Autofluorescent Protein Fusions in Plants: Probing
987     Nicotiana benthamiana–Virus Interactions Plant Physiol. **20,** 740–750.
988 **Raissig MT, Matos JL, Anleu Gil MX, Kornfeld A, Bettadapur A, Abrash**
989     **E, Allison HR, Badgley G, Vogel JP, Berry JA, Bergmann DC.**
990     (2017). Mobile MUTE specifies subsidiary cells to build physiologically
991     improved grass stomata. Science **355,** 1215–1218.
992 **Shao, W., and Dong, J.** (2016). Polarity in plant asymmetric cell division:
993     Division orientation and cell fate differentiation. Dev. Biol. **419,** 121-
994     131.
995 **Sowa, M. E., Bennett, E. J., Gygi, S. P., & Harper, J. W. (2009).** Defining
996     the human deubiquitinating enzyme interaction landscape. Cell, **138,**
997     389-403.
998 **Sugiyama, Y., Nagashima, Y., Wakazaki, M., Sato, M., Toyooka, K.,**
999     **Fukuda, H., and Oda, Y.** (2019). A Rho-actin signaling pathway

shapes cell wall boundaries in Arabidopsis xylem vessels. Nat. Commun. **10,** 468.

Sutimantanapi, D., Pater, D., and Smith, L.G. (2014). Divergent roles for maize PAN1 and PAN2 receptor-like proteins in cytokinesis and cell morphogenesis. Plant Physiol. **164,** 1905-1917.

Stebbins, G.L., and Shah, S.S. (1960). Developmental studies of celldifferentiation in the epidermis of monocotyledons. II. Cytological features of stomatal development in the Gramineae. Dev. Biol. **2,** 477 – 500.

Tamura K. Peterson D. Peterson N. Stecher G. Nei M. Kumar S. (2011). MEGA5: Molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. Mol. Biol. Evol. **28,** 2731–2739.

Walley, J.W., Sartor, R.C., Shen, Z., Schmitz, R.J., Wu, K.J., Urich, M.A., Nery, J.R., Smith, L.G., Schnable, J.C., Ecker, J.R., and Briggs, S.P. (2016). Integration of omic networks in a developmental atlas of maize. Science. **353,** 814-818

Wang, K., Yang, Z., Qing, D., Ren, F., Liu, S., Zheng, Q., Liu, J., Zhang, W., Dai, C., Wu, M., Chehab, E.W., Braam, J., and Li, N. (2018). Quantitative and functional posttranslational modification proteomics reveals that TREPH1 plays a role in plant touch-delayed bolting. Proc. Natl. Acad. Sci. USA **115,** E10265-E10274.

Wu, S.Z., and Bezanilla, M. (2018). Actin and microtubule cross talk mediates persistent polarized growth. J. Cell Biol. **217,** 3531-3544.

Xiang, Y., Huang, X., Wang, T., Zhang, Y., Liu, Q., Hussey, P.J., and Ren, H. (2007). ACTIN BINDING PROTEIN 29 from Lilium pollen plays an important role in dynamic actin remodeling. Plant cell **19,** 1930-1946.

Yang, Z. (2008). Cell polarity signaling in Arabidopsis. Annu. Rev. Cell Dev. Biol. **24,** 551-575.

Yoshida, S., van der Schuren, A., van Dop, M., van Galen, L., Saiga, S., Adibi, M., Moller, B., Ten Hove, C.A., Marhavy, P., Smith, R., Friml, J., and Weijers, D. (2019). A SOSEKI-based coordinate system interprets global polarity cues in Arabidopsis. Nat. Plants **5,** 160-166.

Zhang, X., Facette, M., Humphries, J.A., Shen, Z., Park, Y., Sutimantanapi, D., Sylvester, A.W., Briggs, S.P., and Smith, L.G. (2012). Identification of PAN2 by quantitative proteomics as a leucine-rich repeat-receptor-like kinase acting upstream of PAN1 to polarize cell division in maize. Plant Cell **24,** 4577-4589

Zhang, D., Abrash, E.B., Nunes, T.D., Hidalgo, I., Gil, M.X.A., Jesenofsky, B., Lindner, H., Bergmann, D.C. and Raissig, M.T., (2022). Opposite polarity programs regulate asymmetric subsidiary cell divisions in grasses. bioRxiv. https://doi.org/10.1101/2022.04.24.489281

List of Figures and Tables.

Figure 1. CFP-WPRA2 and RFP-WPRB2 polarize in developing subsidiary mother cells. The stomatal division zone of the leaf epidermis of transgenic maize plants expressing fluorescent fusion proteins was dissected and analyzed. Three different developmental stages were selected according to GMC width and SMC division status to observe the localization of CFP-WPRA2 (A) and RFP-WPRB2 (B). Arrows point to sites of CFP-WPRA2 or RFP-WPRB2 accumulation in SMCs where they contact the adjacent GMC. Asterisks mark a GMC within each stomatal row. Scale bar = 5 µm

Figure 2. Yeast 2-hybrid analysis of WPR and PAN protein interactions. Interactions were assessed using the GAL4-based yeast two-hybrid system. Yeast was grown at different dilutions on non-selective (-L-T) and selective (-L-T-A-H) media. Soluble intracellular regions of PAN1 and PAN2 were used. **(A)** and **(B)**, interactions of PAN1 and PAN2 with WPRA1, WPRA2, WPRB1, WPRB2, and WPRB3. **(C)** Heterodimer formation was assessed between WPRA1, WPRA2, WPRB1, WPRB2, and WPRB3. **(D)** Homodimer formation was assessed between WPRA1, WPRA2, WPRB1, WPRB2, and WPRB3. **(E)** Negative controls using empty bait plasmid pASGW-attR or prey plasmid pACTGW-attR. **(F)** Network diagram showing the observed interactions between WPRs and PAN1/PAN2. **(G)** A model of protein complex formation between PAN and WPR proteins.

Figure 3. WPR proteins polarize in SMCs after BRK1 and before actin. The stomatal division zone of leaf 4 from plants co-expressing **(A)** CFP-WPRA2 & RFP-WPRB2; **(B)** BRK1-CFP & RFP-WPRB2; **(C)** CFP-WPRA2 & PAN1-YFP; **(D)** CFP-WPRA2 & PAN2-YFP; or **(E)** CFP-WPRA2 & FABD2-YFP were analyzed for enrichment at the GMC-SMC interface. Fluorescent proteins were scored as polarized when fluorescence was brighter at the SMC-GMC interface than at the cell periphery distal to the GMC. Within each panel, the same GMCs imaged in the different channels (and the subsequent merged panel) are numbered. Arrows point to SMCs where one fluorescent protein is polarized, but not the other. Counts of individual SMCs that had only one fluorescent protein, or both, are listed at right of the images. All images are at the same scale and are Z-projections of 7 confocal slices. Scale bar in the merged panel of E is 10 µm.

Figure 4. **Polarization of WPR proteins depends on BRK1 and PAN2, but not PAN1.** Localization of CFP-WPRA2 and RFP-WPRB2 in *brk1*, *pan2* and *pan1* mutants and non-mutant siblings was analyzed in developing leaf 4. CFP-WPRA2 **(A)** and RFP-WPRB2 **(B)** in developing stomata of *brk1* mutants

and non-mutant (heterozygous) siblings. CFP-WPRA2 **(C)** and RFP-WPRB2
(D) in developing stomata of *pan2* mutants and non-mutant (heterozygous)
siblings. CFP-WPRA2 **(E)** and RFP-WPRB2 **(F)** in developing stomata of
pan1 mutants and non-mutant (heterozygous) siblings. Asterisks mark GMC
rows. Arrows point to SMCs without polarized localization of CFP-WPRA2 or
RFP-WPRB2. 3-6 plants of each genotype were used for data collection.
Scale bar, 5 μm.

1092

Figure 5. CRISPR-Cas9 induced *wprb1*;*wprb2* double mutants and CFP-
WPRA2 and RFP-WPRB2 expressing lines have subsidiary cell defects.
(A-F) Representative image of the third leaf epidermis of *wprb1/+; wprb2/+*
and *wprb1/wprb1;wprb2/wprb2* mutants **(A and D)**, CFP-WPRA2 transgenic
plants and non-transgenic siblings **(B and E)** and RFP-WPRB2 transgenic
plants and non-transgenic siblings **(C and F).** Examples of normal subsidiary
cells in A, B and D are false-colored blue, and abnormal subsidiary cells in D,
E and F are false-colored yellow and marked with black arrows. (G)
Quantification of abnormal subsidiary cells in *wprb1/+; wprb2/+* (n = 27
plants), *wprb1/b1;wprb2/+* (n = 14 plants), *wprb1/+;wprb2/wprb2* (n = 22
plants) and *wprb1/wprb1;wprb2/wprb2* (n=18 plants). For each plant 100-200
subsidiary cells were examined. p-value from Student's T-test comparing
each genotypes, ***$P \leq 0.001$. (H) and (I) Quantification of abnormal
subsidiary cells in CFP-WPRA2 or RFP-WPRB2 expressing plants and non-
transgenic siblings grown in parallel. (n = 11-12 plants and 120-200 cells per
genotype). Student's T-test were performed, ***$p \leq 0.001$.

1109

Figure 6. WPRB interacts with F-actin. (A) Schematic depicting WPRB2
and truncated versions. **(B-F)** Confocal images of transiently expressed full-
length GFP fusion proteins in tobacco. Transient expression of GFP only **(B)**,
GFP-WPRA2 **(C)**, GFP-WPRB1 **(D)**, GFP-WPRB2 **(E)** or GFP-WPRB3 **(F)**.
(G-K) Transient expression of truncated WPRB2 proteins fused to GFP in
tobacco leaves. Co-expression of GFP-WPRB2 **(L)** with Lifeact-RFP labeled
actin microfilaments **(M)**; merged image **(N)** of the same cell. Tobacco leaves
expressing GFP-WPRB2 were treated with DMSO **(negative control, O)** or 40
μM Latrunculin B **(P)** for 2 hours. All images at the same scale, Scale bar in
(P) = 10 μm. **(Q)** High-speed co-sedimentation of GST-tagged WPRB2N240
with F-actin. After centrifugation at 100,000*g*, the proteins in the supernatant
(S) and pellet (P) were resolved by SDS-PAGE and visualized with
Coomassie Blue staining. GST protein was used as a negative control.

1123

Figure 7. Fluorescence intensity of ABD2-YFP decreases in RFP-WPRB2 expressing plants.

Plants expressing RFP-WPRB2 and FABD2-YFP were crossed, and the progeny independently segregated the two markers. (Ai & Aii) Plant only expressing FABD2-YFP. The identical image is shown in Ai and Aii. In Ai, the 16-bit image was scaled from 600 to 12000 prior to converting to 8 bit. Aii was scaled 500 to 2000. (B) Plant only expressing RFP-WPRB2. (C-E) Plant co-expressing FABD2-YFP (C, green in E) with RFP-WPRB2 (D, magenta in E). Ci and Cii show the identical image of the ABD2-YFP channel, where Ci is scaled the same as Ai and Cii is scaled the same as Aii, for comparative purposes. Images in B and D are scaled identically. Scale bar in E = 10 microns. (F) Quantification of fluorescence intensity of SMC lateral cell side (site A) and SMC-GMC interface (site B) in FABD2-YFP only plants (n = 5 plants, 292 cells) and RFP-WPRB2 and FABD2-YFP co-expressing plants (n = 3 plants, 164 cells). One sided t-tests indicate lower values in co-expressing plants (p < 0.0001). (G) Ratio of intensities measured in panel F. Two-sided test indicates a significantly lower ratio in the co-expressing cells (p<0.0001).

Table 1. **WPR proteins interact with other WPR family members and PAN receptors.** WPR-interacting proteins identified by Co-IP/MS. For a complete list of all proteins, see Supplementary Data Set 1. Three different experiments were performed. 1: Anti-GFP was used to pull down WPRA2 and interacting proteins from extracts of plants expressing CFP-WPRA2, and from non-transgenic plants as a negative control. 2: Anti-WPRA1/A2 was used to pull down WPRA1/A2 and interacting proteins from B73 extracts; negative controls used the same extracts but omitted the anti-WPRA1/A2 antibody. 3: Anti-RFP was used to pull down WPRB2 and interactors from extracts of plants expressing RFP-WPRB2, and from non-transgenic plants as a negative control. B73 vs4 gene codes are used to identify proteins, and the number of unique peptides (# peptides) is listed. The baits WPRA2, WPRA1/A2, and WPRB2 are highlighted in bold.

Supplemental Data

Supplemental Figure 1. Progression of maize stomatal development in leaves.

Supplemental Figure 2. Protein tree of the WPR family in plants.

Supplemental Figure 3. Plasmolysis assays of CFP-WPRA2 and RFP-WPRB2 transgenic plants.

Supplemental Figure 4. Immunolocalization using validated WPRA1/2 and WPRB2 antibodies and their validation

Supplemental Figure 5. CRISPR-Cas9 induced *wpra1 and wpra2* mutants might have transmission defects.

Supplemental Figure 6. CRISPR-Cas9 induced *wprb1;wprb2* double
mutants have no effect on stomatal density.

Supplemental Figure 7. Mutations in *wprb1* and *wprb2* do not enhance
***pan2/pan2* phenotype.**

Supplemental Figure 8. CRISPR-Cas9 induced *wprb1;wprb2* double
mutants have no effect on polarized actin accumulation or nuclear
polarization during maize SMC development.

Supplemental Figure 9 Latrunculin B-treated tobacco leaf cells
transiently expressing GFP-WPRB2 and Lifeact-RFP.

Supplemental Figure 10. Transient expression of WPRB2-CFP, and its
truncated proteins in tobacco leaves.

Supplemental Figure 11. Alignment of the actin-binding domain of the
WPR protein family.

Supplemental Figure 12. RFP-WPRB2 expression has no effect on the
total amount of actin.

Supplemental Figure 13. Co-expression of CFP-WPRA2 with ABD2-YFP
does not decrease FABD2-YFP intensity

Supplemental Figure 14. F-actin disruption does not affect the polarized
localization of RFP-WPRB2 in SMCs.

Supplemental Table1.   Primer sequences used in this study.

Supplemental Data Set 1. Co-IP/MS results using an anti-GFP antibody
and membrane extracts from PAN2-YFP, PAN1-YFP, BRK1-CFP, PDI-
YFP, PIN1-YFP, Rab11D-YFP and B73

Supplemental Data Set 2. Co-IP/MS results using WPRA1/2, WPRA2-
CFP, WPRB2-RFP as bait (and corresponding controls)

Supplemental File1. Amino acid sequences of WPRs used for the
phylogenetic analysis in Supplemental Figure 2.

	Gene ID	Gene name	Rep1 # Unique Peptides	Rep2 # Unique Peptides	Rep3 # Unique Peptides	Control 1 # Unique Peptides	Control 2 # Unique Peptides	Control 3 # Unique Peptides
Anti-GFP	Zm00001d041088	**WPRA2 (bait)**	28	42	2	2	0	0
	Zm00001d007164	WPRB2	5	3	0	0	0	0
	Zm00001d010610	WPRB3	15	13	0	0	0	1
	Zm00001d029420	WEB subfamily protein	5	7	0	0	0	0
Anti-WPRA1/A2	Zm00001d023629	**WPRA1 (bait)**	24	19	26	0	1	2
	Zm00001d041088	**WPRA2 (bait)**	26	19	27	0	0	0
	Zm00001d047516	WPRB1	5	4	6	0	0	0
	Zm00001d007164	WPRB2	19	21	21	0	0	0
	Zm00001d010610	WPRB3	24	21	24	0	1	0
	Zm00001d026139	WPRC subfamily protein	2	1	0	0	0	0
	Zm00001d000443	WPRC subfamily protein	2	2	0	0	0	1
	Zm00001d048523	WEB subfamily protein	2	2	2	0	0	0
	Zm00001d007862	PAN2	4	5	4	0	0	2
	Zm00001d031437	PAN1	2	5	2	0	0	0
Anti-RFP	Zm00001d007164	**WPRB2 (bait)**	18	20	17	0	1	0
	Zm00001d023629	WPRA1	18	16	16	0	0	0
	Zm00001d041088	WPRA2	15	21	20	0	2	0
	Zm00001d010610	WPRB3	1	2	1	0	0	0
	Zm00001d007826	PAN2	3	0	1	0	0	0

1193
Table 1. **WPR proteins interact with other WPR family members and**
PAN receptors. WPR-interacting proteins identified by Co-IP/MS. For a
complete list of all proteins, see Supplementary Data Set 1. Three different
experiments were performed. 1: Anti-GFP was used to pull down WPRA2 and
interacting proteins from extracts of plants expressing CFP-WPRA2, and from
non-transgenic plants as a negative control. 2: Anti-WPRA1/A2 was used to
pull down WPRA1/A2 and interacting proteins from B73 extracts; negative
controls used the same extracts but omitted the anti-WPRA1/A2 antibody. 3:

Anti-RFP was used to pull down WPRB2 and interactors from extracts of
plants expressing RFP-WPRB2, and from non-transgenic plants as a negative
control. B73 vs4 gene codes are used to identify proteins, and the number of
unique peptides (# peptides) is listed. The baits WPRA2, WPRA1/A2, and
WPRB2 are highlighted in bold.

Supplemental Table 1. Primer sequences used in this study.

Primer name	Sequence (5'-3')	Description
WPRA2-F	GGGGACA AGTTTGTACAAAAAAGCAGGCTGC ATAACCATTTTATGCTTCAAGC	Primers for *Wpra2* promoter and 5' UTR amplification
WPRA2-R	GGGG AC CAC TTT GTA CAA GAA AGC TGG GTA GAAGTTGCAATTACACCTCTTTGT	
WPRA2-P2	GCTCCACCTCCACCTCCCATTTATTTAAATATTTTACGC	Primers for *Wpra2* coding region and 3' UTR amplification
WPRA2-P3	GCTGCTGCGGCCGCTGGGGCCATGAAGGCAAATATGGGCAC	
WPRB2-F	GGGGACAAGTTTGAAAAAAGCAGGCTGCGAGATTCG AGACAACATACATG	Primers for *Wprb2* promoter and 5' UTR amplification
WPRB2-R	GGGGACCACTTTGTACAAGAAAGCTGGGTATAAAACT CTCACACGAGAGATAC	
WPRB2-P2	GCTCCACCTCCACCTCCCATGGGGACTCGTGCCCCTCT	Primers for *Wprb2* coding region and 3' UTR amplification
WPRB2-P3	GCTGCTGCGGCCGCTGGGGCCATGAACGACGCCGGCGA GGCCG	
Tag linker-F	ATGGGAGGTGGAGGTGGAGCT	Primers for CFP or RFP amplification
Tag linker-R	GGCCCCAGCGGCCGCAGCAGC	
WPRB2 topo-F	CACCATGAACGACGCCGGCGAGG	Primers for WPRB2N67 amplification
WPRB2N topo (1-67)-R	CTTTGGGCTCGGGGGGCAGCG	
WPRB2N topo (1-142)-R	AGCGGCAGTGGGCAGCTCGGCTTG	WPRB2N142 R primer
WPRB2F -GW	GGGGACAAGTTTGTACAAAAAAGCAGGCTGCGC ATGAACGACGCCGGCGAGG	Primers for WPRB2 without stop code amplification
WPRB2R -GW(no stop code)	GGGGACCACTTTGTACAAGAAAGCTGGGTA ATGCGATCCCTTTTCCTTCTTC	
WPRB2 △N F-GW	GGGGACAAGTTTGTACAAAAAAGCAGGCTGC GCCGCTGAAGAGGTTGCAGAAG	Primers for WPRB2△C, WPRB2△N,WPRB2△NC amplification
WPRB2 △C R-GW	GGGGACCACTTTGTACAAGAAAGCTGGGTA CCTGTGTCCAGTCGTGGATCG	
WPRA1 topo-F	CACCATGAAGGCAAATATGGGC	Primers for WPRA1 CDS amplification
WPRA1 topo-R	TCACACAGATTTTTCCCCAGGAAG	
WPRA2 topo-F	CACCATGGGGGTGGGGAAGAGCA	Primers for WPRA2 CDS

WPRA2 topo-R	TTACACAGATTTTTCCCCAGGAAG	amplification
WPRB1 topo-F	CACCATGAGCTCCGAAGCAGAA	Primers for WPRB1 CDS amplification
WPRB1 topo-R	TTAAAAGGATTCTCTCTTCTT	
WPRB2 topo-F	CACCATGAACGACGCCGGCGAG	Primers for WPRB2 CDS amplification
WPRB2 topo-R	CTAATGCGATCCCTTTTC	
WPRB3 topo-F	CACC ATGGCGGAAGTGGCGGCGCGGC	Primers for WPRB3 CDS amplification
WPRB3 topo-R	GCTCAAGCCACCACGAGTGCAGGAG	
WPRA1 BamHI 63F	CGCGGATCC TTAGCCAAGGAGACAGA	Primers of WPRA1(63-503) amplification and GST-WPRA1 expression
WPRA1 EcorI 503R	CCGGAATTC CAACTTCTCTGACTC	
WPRA2 BamHI 63F	CGCGGATCC TTAGCCAAGGAGACAGA	Primers of WPRA1(63-570) amplification and GST-WPRA2 expression
WPRA2 EcorI 570R	CCGGAATTC CTTCTCGCGCCATCT	
WPRB1 BamHI 73F	CGCGGATCC GCCGAGGAGCAGGCT	Primers of WPRB1(73-421) amplification and GST-WPRB1 expression
WPRB1 EcorI 421R	CCGGAATTC TTACCCGCTCTCAGATTT	
WPRB2 BamHI 78F	CGCGGATCC GAGGAGCAGACCGCG	Primers of WPRA1(78-429) amplification and GST-WPRB2 expression
WPRB2 EcorI 429R	CCGGAATTC GGATCGACATCTGCG	
WPRB2 BamHI F	CGCGGATCCATGAACGACGCCGGCGAG	Primers of WPRB2N240 amplification and GST-WPRB2N240 expression
WPRB2 N240 EcorI R	CCGGAATTCTTACTTCTGTGCAGTCCGGTTCAGCTC	

1208

WPRA2-CFP

WPRB2-RFP

Early   Middle   Late stage

Figure 1. CFP-WPRA2 and RFP-WPRB2 polarize in developing subsidiary mother cells. The stomatal division zone of the leaf epidermis of transgenic maize plants expressing fluorescent fusion proteins was dissected and analyzed. Three different developmental stages were selected according to GMC width and SMC division status to observe the localization of CFP-WPRA2 (A) and RFP-WPRB2 (B). Arrows point to sites of CFP-WPRA2 or RFP-WPRB2 accumulation in SMCs where they contact the adjacent GMC. Asterisks mark a GMC within each stomatal row. Scale bar = 5 μm

Figure 2. Yeast 2-hybrid analysis of WPR and PAN protein interactions. Interactions were assessed using the GAL4-based yeast two-hybrid system. Yeast was grown at different dilutions on non-selective (-L-T) and selective (-L-T-A-H) media. Soluble intracellular regions of PAN1 and PAN2 were used. (A) and (B), interactions of PAN1 and PAN2 with WPRA1, WPRA2, WPRB1, WPRB2, and WPRB3. (C) Heterodimer formation was assessed between WPRA1, WPRA2, WPRB1, WPRB2, and WPRB3. (D) Homodimer formation was assessed between WPRA1, WPRA2, WPRB1, WPRB2, and WPRB3. (E) Negative controls using empty bait plasmid pASGW-attR or prey plasmid pACTGW-attR. (F) Network diagram showing the observed interactions between WPRs and PAN1/PAN2. (G) A model of protein complex formation between PAN and WPR proteins.

Figure 3. WPR proteins polarize in SMCs after BRK1 and before actin. The stomatal division zone of leaf 4 from plants co-expressing (A) CFP-WPRA2 & RFP-WPRB2; (B) BRK1-CFP & RFP-WPRB2; (C) CFP-WPRA2 & PAN1-YFP; (D) CFP-WPRA2 & PAN2-YFP; or (E) CFP-WPRA2 & FABD2-YFP were analyzed for enrichment at the GMC-SMC interface. Fluorescent proteins were scored as polarized when fluorescence was brighter at the SMC-GMC interface than at the cell periphery distal to the GMC. Within each panel, the identical GMCs imaged in the different channels (and the subsequent merged panel) are numbered. Arrows point to SMCs where one fluorescent protein is polarized, but not the other. Counts of individual SMCs that had only one fluorescent protein, or both, are listed at right of the images. All images are at the same scale and are Z-projections of 7 confocal slices. Scale bar in the merged panel of E is 10 μm.

Figure 4. Polarization of WPR proteins depends on BRK1 and PAN2, but not PAN1. Localization of CFP-WPRA2 and RFP-WPRB2 in *brk1, pan2* and *pan1* mutants and non-mutant siblings was analyzed in developing leaf 4. CFP-WPRA2 (A) and RFP-WPRB2 (B) in developing stomata of *brk1* mutants and non-mutant (heterozygous) siblings. CFP-WPRA2 (C) and RFP-WPRB2 (D) in developing stomata of *pan2* mutants and non-mutant (heterozygous) siblings. CFP-WPRA2 (E) and RFP-WPRB2 (F) in developing stomata of *pan1* mutants and non-mutant (heterozygous) siblings. Asterisks mark GMC rows. Arrows point to SMCs without polarized localization of CFP-WPRA2 or RFP-WPRB2. 3-6 plants of each genotype were used for data collection. Scale bar, 5 μm.

Figure 5. CRISPR-Cas9 induced *wprb1;wprb2* **double mutants and CFP-WPRA2 and RFP-WPRB2 expressing lines have subsidiary cell defects.** (A-F) Third leaf epidermis of *wprb1/+; wprb2/+* and *wprb1/wprb1;wprb2/wprb2* siblings (A and D), CFP-WPRA2 transgenic plants and non-transgenic siblings (B and E) and RFP-WPRB2 transgenic plants and non-transgenic siblings (C and F). Examples of normal subsidiary cells in A, B and D are false-colored blue, and abnormal subsidiary cells in D, E and F are false-colored yellow and marked with black arrows. (G) Quantification of abnormal subsidiary cells in *wprb1/+; wprb2/+* (n = 27 plants), *wprb1/b1;wprb2/+* (n = 14 plants), *wprb1/+;wprb2/wprb2* (n = 22 plants) and *wprb1/wprb1;wprb2/wprb2* (n=18 plants). For each plant 100-200 subsidiary cells were examined. p-value from Student's T-test comparing each genotypes, ***P≤ 0.001. (H) and (I) Quantification of abnormal subsidiary cells in CFP-WPRA2 or RFP-WPRB2 expressing plants and non-transgenic siblings grown in parallel. (n = 11-12 plants and 120-200 cells per genotype). Student's T-test were performed, ***p≤ 0.001.

Figure 6. WPRB interacts with F-actin. (A) Schematic depicting WPRB2 and truncated versions. (B-F) Confocal images of transiently expressed full-length GFP fusion proteins in tobacco. Transient expression of GFP only (B), GFP-WPRA2 (C), GFP-WPRB1 (D), GFP-WPRB2 (E) or GFP-WPRB3 (F). (G-K) Transient expression of truncated WPRB2 proteins fused to GFP in tobacco leaves. Co-expression of GFP-WPRB2 (L) with Lifeact-RFP labeled actin microfilaments (M); merged image (N) of the same cell. Tobacco leaves expressing GFP-WPRB2 were treated with DMSO (negative control, O) or 40 μM Latrunculin B (P) for 2 hours. All images at the same scale, Scale bar in (P) = 10 μm. (Q) High-speed co-sedimentation of GST-tagged WPRB2N240 with F-actin. After centrifugation at 100,000g, the proteins in the supernatant (S) and pellet (P) were resolved by SDS-PAGE and visualized with Coomassie Blue staining. GST protein was used as a negative control.

Figure 7. Fluorescence intensity of ABD2-YFP decreases in RFP-WPRB2 expressing plants.
Plants expressing RFP-WPRB2 and FABD2-YFP were crossed, and the progeny independently segregated the two markers. (Ai & Aii) Plant only expressing FABD2-YFP. The identical image is shown in Ai and Aii. In Ai, the 16-bit image was scaled from 600 to 12000 prior to converting to 8 bit. Aii was scaled 500 to 2000. (B) Plant only expressing RFP-WPRB2. (C-E) Plant co-expressing FABD2-YFP (C, green in E) with RFP-WPRB2 (D, magenta in E). Ci and Cii show the identical image of the ABD2-YFP channel, where Ci is scaled the same as Ai and Cii is scaled the same as Aii, for comparative purposes. Images in B and D are scaled identically. (F) Quantification of fluorescence intensity of SMC lateral cell side (site A) and SMC-GMC interface (site B) in FABD2-YFP only plants (n = 5 plants, 292 cells) and RFP-WPRB2 and FABD2-YFP co-expressing plants (n = 3 plants, 164 cells). One sided t-tests indicate lower values in co-expressing plants (p < 0.0001). (G) Ratio of intensities measured in panel F. Two-sided test indicates a significantly lower ratio in the co-expressing cells (p<0.0001).

Supplemental Figure 1. Progression of maize stomatal development in leaves. Unexpanded maize leaves mature in a gradient with the developmentally youngest cells towards the base and the developmentally oldest cells towards the tip. **(A)** Stomatal complex formation is initiated by a transverse asymmetric division in a new stomatal row. **(B)** Pavement cells flanking the GMC acquire a SMC fate. At this stage, the nuclei are not yet polarized, but proteins (not shown) in the SMC may be polarized towards the GMC. **(C)** SMC nuclei polarize towards the GMC; preprophase band forms. **(D)** The SMCs divide, with each division forming **(E)** a subsidiary cell and a pavement cell. The GMC undergoes an oriented, symmetric division symmetrically to form (**F**), the guard cell pair. Each stomatal complex is made of two subsidiary and two guard cells. Nuclei/DNA are shown in magenta; microtubule structures are shown in green. **(G)** Ordered sequence of protein polarization, which occurs at the time shown in panel B. (**H-K**) Mutants with polarity defects result in abnormal subsidiary cells exhibit a range of abnormal morphologies. Examples of a single stomatal complex from wild type, *brk1, pan2,* and *pan1* mutants, where abnormal subsidiaries are false-colored yellow. Supports Figures 1 and 5.

Supplemental Figure 2. Protein tree of the WPR family in plants. Protein coding sequences of WPR family members were identified from Phytozome 13 (https://phytozome-next.jgi.doe.gov/) from *A. thaliana* (dicot), *Amborella trichopoda* (dicot), *Solanum lycopersicum* (tomato, dicot), *B. distachyon* (monocot) and *Zea mays* (maize, monocot), and a protein tree was inferred. The proteins fall into 5 subfamilies, including the previously identified WEB/WEL, WPRB, WPRA and PMI clades – we also identified an additional WPRC clade. Previously named proteins are listed, and asterisks mark proteins characterized in this study. Protein sequences were aligned using Clustal Omega and the tree was inferred using MEGA 11. Numbers at nodes represent the percentage values given by 1000 bootstrap analysis samples. At: *Arabidopsis thaliana*, Zm: *Zea mays*. Bd: *Brachypodium distachyon*, Solyc: *Solanum lycopersicum*, scaffold: *Amborella trichopoda*. Supports Figure 1.

Supplemental Figure 3. **Plasmolysis assays of CFP-WPRA2 and RFP-WPRB2 transgenic plants.**
Plants expressing either CFP-WPRA2 or RFP-WPRB2 were treated with 0.8M mannitol to induce
plasmolysis, to determine if WPR proteins were primarily localized to SMCS or GMCs. Similar results were
found with both fusions, with representative images shown. Although plasmolysis results in significant loss
of overall signal, fluorescence was still visible in some cells, especially in SMCs. (A) CFP-WPRA2
expressing cells. Arrows point to bright fluorescent patches in two cells that are SMCS. Based on the
curvature of the membrane, away from GMCs (marked by asterisks), the polarized fluorescence is
predominantly in SMCs. Residual signal could also be seen in GMCs, although typically was much
stronger in SMCs. (B) RFP-WRPB2 expressing cells. In these more mature cells, the SMCs have likely
already divided, and polarized fluorescence remains in subsidiary cells. Arrow points to a cell where
fluorescence is clearly seen in the subsidiary cell, but not the adjacent GMC. (C) and (D) are cartoon
schematics of (A) and (B) respectively, with color-coding of cell types. Scale bar = 10 μm. Supports Figure
1.

Supplemental Figure 4. **Immunolocalization using WPRA1/2 and WPRB2 antibodies and their validation (a)** Immunostaining with anti-WPRA1/A2 (green) antibodies and propidium iodide-staining (magenta) of developing leaf 4. No anti-WPRA1/A2 was used as a negative control. Arrows indicate the localization of WPRA1/A2 at the interface of subsidiary cells and guard cells. **(b and c)** Immunostaining with anti-WPRB2 (green) antibodies and propidium iodide (magenta). CRISPR-Cas9 induced *wprb2* mutants were used as a negative control. Arrows in (b) indicate the localization of WPRB2 at the interface of subsidiary cells and guard cells, in (c) indicate the localization of WPRB2 at pavement cell corner. **(d-f)** Confirmation of anti-WPRA1/A2 **(d and f)** and anti-WPRB2 **(e)** specificity using Western blotting. **(d)** Western blot of proteins extracted from B73 and CFP-WPRA2 transgenic plants probed with affinity-purified anti-WPRA1/A2. A band corresponding to the predicted size of endogenous WPRA1/A2 was recognized in B73. The same band was recognized in CFP-WPRA2 expressing plants; as was a larger band specific to the transgenic plants, which corresponds to the predicted size of CFP-WPRA2. In the right panel, Coomassie blue (CCB) staining of the gel is shown as a loading control. **(e)** Proteins extracted from *wprb2* mutants and B73 were used for western blotting assays. A band around 75 kD which consistent with the mass of WPRB2 protein was recognized in B73 but not in two *wprb2* mutants, Coomassie blue (CCB) staining is shown as a loading control. **(f)** Western blot of bacterially-expressed and purified GST-tagged WPRA1, WPRA2, WPRB1, WPRB2 proteins probed with affinity-purified WPRA1/A2 antibody demonstrates the specificity of this antibody for WPRA proteins. The Coomassie blue–stained gel (CCB) in the lower panel shows the purified proteins of WPRA1, WPRA2, WPRB1, WPRB2. Supports Figure 1, Figure 2 and Supplemental Dataset 2.

Parents	Progeny	Expected	Observed Ratio	χ2 P Value
wpra1/+; wpra2/+ selfing	*a1_;a2_* ፧ *a1_;a2/a2* ፧ *a1/a1;a2_* ፧ *a1/a1;a2/a2*	9:3:3:1	78:15:13:0	P < 0.01
wpra1/+; wpra2/wpra2 selfing	*+/+;a2/a2* ፧ *a1/+;a2/a2* ፧ *a1/a1;a2/a2*	1:2:1	38:48:0	P < 0.01

Supplemental Figure 5. CRISPR-Cas9 induced *wpra1 and wpra2* mutants have transmission defects. Mutated regions are labeled with black squares. **(A)** DNA sequences of Cas9-edited *Wpra1*. *wpra1* has a two nucleotide "GA" deletion at the gRNA2-targeted site. **(B)** and **(C)** DNA sequences of Cas9-edited *Wpra2,* Two mutations were identified: a "T" deletion was found at the gRNA1-targeted site (B), and an 11bp (GCACACGAAGA) deletion was found at gRNA2-targeted site (arrow, C). **(D)** Genotype analysis was performed in the progeny of *wpra1/+; wpra2/+* and *wpra1/+; wpra2/wpra2* mutants. No *wpra1/wpra1; wpra2/wpra2* double mutants were recovered; observed ratios are significantly different from expected ratios. **(E)** RNA-seq expression level in maize growth stages (Data from MaizeGDB, Walley et al., 2016). Stage 21 is B73_Mature Pollen (expression level = 54.4). **(F)** CFP-WPRA2 localization in maize pollen tube. Maize pollen stably expressing CFP-WPRA2 were germinated in liquid culture for 1 hour at room temperature observed using spinning disk confocal and images were collected at × 60 objective (NA = 1.20, Olympus). The arrows point to subapex region localization of CFP-WPRA2. Scale bar in (F) = 10 μm. Supports Figure 5.

Supplemental Figure 6. CRISPR-Cas9 induced *wprb1;wprb2* double mutants have no effect on stomatal density. (A) Representative image of the third leaf of epidermis of *wprb1/+;wprb2/+* and *wprb1/b1;wprb2/b2* mutants. Scale bar, 0.1 mm. **(B)** Cas9-edited *Wprb1* and *Wprb2* genes. Mutated regions are labeled in black square; wprb1 has a "G" insertion and wprb2 has a "C" deletion in the gRNA targeted region. These mutations cause frameshifts and premature termination. **(C)** Quantification of stomatal density in *wprb1/+; wprb2/+* (n = 27 plants), *wprb1/b1;wprb2/+* (n = 14 plants), *wprb1/+;wprb2/wprb2* (n = 22 plants) and *wprb1/wprb1;wprb2/wprb2* (n=18 plants). Student's T-tests were performed among these genotypes, no significant difference in these mutants. Supports Figure 5.

Supplemental Figure 7. Mutations in *wprb1* and *wprb2* do not enhance *pan2/pan2* phenotype.
Stomatal phenotypes of progeny of *pan2/pan2; wprb1/+; wprb2/+* plants were performed using the fully expanded third leaf. The percentage of abnormal subsidiary cell (SC) was quantified in different genotypes, *pan2/pan2; wprb1/+; wprb2/+* (n = 113 plants), *pan2/pan2; wprb1/wprb1;wprb2_*(n = 38 plants), *pan2/pan2;wprb1_; wprb2/wprb2* (n= 41 plants) *and triple mutants pan2/pan2;wprb1/wprb1; wprb2/wprb2*(n= 11 plants). For each plant 100-200 subsidiary cells were genotyped. ANOVA and pair-wise student's T-tests were performed among these genotypes with no significant differences. Supports Figure 5.

Supplemental Figure 8. CRISPR-Cas9 induced *wprb1;wprb2* double mutants have no effect on polarized actin accumulation or nuclear polarization during maize SMC development. The stomatal division zone of developing leaf 4 from *wprb1;wprb2* double mutants, and heterozygous siblings was stained using Alexafluor488-phalloidin (green) and propidium iodide (magenta). SMCs adjacent to GMCs that were greater than than 6 µm in width were used for the quantification, at this stage the majority of SMCs in wild-type had both actin patch and polarized nucleus at SMC-GMC interface. Cells were assayed for the presence of an actin patch, and whether the nucleus was polarized (touching the SMC-GMC interface). (A) Representative image of Alexafluor488-phalloidin stained actin (green) and propidium iodided stained DNA (magenta) in leaf epidermis from *wprb1/+;wprb2/+* and *wprb1/b1;wprb2/b2* plants. Bar = 5 µm. (B) Quantification of SMC classifications in *wprb1/+;wprb2/+* (n = 3 plants, 221 cells) and *wprb1/b1;wprb2/b2* (n = 3 plants, 241 cells) sibling plants. Supports Figure 5.

Supplemental Figure 9 Latrunculin B-treated tobacco leaf cells transiently expressing GFP-WPRB2 and Lifeact-RFP. Tobacco leaves co-expressing GFP-WPRB2 and Lifeact-RFP were treated with DMSO **(**negative control, **A-C)** or 40 μM Latrunculin B **(D-F)** for 2 hours. Supports Figure 6.

Supplement Figure 10. **Transient expression of WPRB2-CFP, and its truncated proteins.**
Localization of WPRB2 in tobacco was confirmed using a CFP (rather than GFP) tag, and a C-terminal
(rather than N-terminal) fusion. See Figure 6A for cartoon of protein truncations tested. (A) Soluble CFP.
(B) Full length WPRB2-CFP. (C) C-terminal truncation WPRB2ΔC-CFP (D) N-terminal truncation
WPRB2ΔN-CFP localizes predominantly to the nucleus. (E) Central DUF827 domain only WPRB2ΔNC.
Filaments structures appears in WPRB2-CFP and WPRB2△C-CFP expressing tobacco cells, but not in
WPRB2ΔN-CFP and WPRB2ΔNC-CFP expressing tobacco cells. Supports Figure 6.

Supplemental Figure 11. Alignment of the actin-binding domain of the WPR protein family. The N terminal amino acids sequence WPRs from *Arabidopsis* and maize were aligned using ClustalX 2.0.5 (Larkin et al., 2007) with the default settings. Supports Figure 6.

Supplemental Figure 12. RFP-WPRB2 expression has no effect on the total amount of actin.
Western blot of total proteins extracted from the stomatal division zone of two independent plants of B73, RFP-WPRB2 transgenic plants, RFP negative siblings. Extracts were probed with an anti-actin antibody. Ponceau S staining is shown as a loading control.

FABD2-YFP alone   **CFP-WPRA2**   **FABD2-YFP**   **Merged**

Co-expression

Supplemental Figure 13. Co-expression of CFP-WPRA2 with ABD2-YFP does not decrease FABD2-YFP intensity. Plants expressing CFP-WPRA2 and FABD2-YFP were crossed, and the progeny independently segregated the two markers. **(A)** Representative image of plant only expressing FABD2-YFP. **(B-D)** Plant co-expressing FABD2-YFP (yellow in D) **(C)** with CFP-WPRA2 (cyan in E) and the merged image **(D),** Scale bar in (D) = 5 μm. **(E, F)** Quantification of fluorescence intensity and intensity ratio of SMC lateral cell side (site A) and SMC-GMC interface (site B) in FABD2-YFP only plants (n = 3 plants, 254 cells) and CFP-WPRA2 + FABD2-YFP co-expressing plants (n = 5 plants, 379 cells). Images are 7-slice max projections. Dots of the same color are from the same plant. Supports Figure 7.

Supplemental Figure 14. **F-actin disruption does not affect the polarized localization of RFP-WPRB2 in SMCs.**(A) FABD2-YFP expressing maize leaves were treated with DMSO (negative control) and 40 µM Latrunculin B for 4 hours. **(B)** RFP-WPRB2 expressed maize leaves were treated with DMSO (negative control) and 40 µM Latrunculin B for 4 hours. Asterisks mark GMCs.