**Communication**

**cMYC protein interactions and chromatin association in NUT carcinoma**

Barry M. Zee[1,2#], Artyom A. Alekseyenko[1,2#], Anne E. Smolko[1,2], Mitzi I. Kuroda[1,2]*

*Corresponding author

#Equal contribution

[1]Div. of Genetics, Dept. of Medicine, Brigham and Women's Hospital, Boston, MA.

[2]Dept. of Genetics, Harvard Medical School, Boston, MA.

**Simple Summary:**

A longstanding goal in biology is to understand how protein interactions influence and reflect

cellular disease states. MYC is a critical regulator of cellular proliferation that is mis-regulated in

many cancers, including those with NUTM1-rearrangements featured in this issue. NUT

carcinoma cells are dependent on MYC expression, which blocks differentiation. Using a small

molecule inhibitor to induce differentiation of a NUTM1-rearranged patient cell line, we sought

to understand how MYC protein interactions are affected by cellular status. When cells

differentiate, we found MYC levels were globally diminished, yet the identities of its protein

partners and genomic binding sites remained constant. During both growth and differentiation,

MYC protein interacted with subunits of the NuA4 lysine acetyltransferase complex, similar to its

associations in non-NUTM1 cell lines. Taken together, our results suggest that lysine

acetyltransferase activities participate in multiple oncogenic steps in NUT carcinoma.

**Abstract:**

The oncogene c*MYC* (HGNC:7553) is a critical genomic target of the BRD4-NUT (B4N) protein

that defines many of the NUTM1-rearrangement cancer subtypes in NUT carcinoma (NC). B4N

interacts with the EP300 lysine acetyltransferase (KAT3B) to create hyperacetylated

"megadomains" that activate downstream genes such as *cMYC*. However, it is less well understood

how mis-regulation of cMYC in turn affects its protein interactions and target genes. The goals of

this study are to define the protein and genomic interactions of cMYC in NC, and to test whether

those change in response to B4N inactivation. We used CRISPR-Cas9 mediated knock-in of a

BioTAP affinity tag to analyze cMYC protein expressed from its normal chromosomal context.

This allowed us to adapt a crosslinking purification method termed BioTAP-XL to preserve cMYC

integrity and chromatin association for genomic and mass spectrometry-based proteomic analyses.

We found that cMYC interacts primarily with the NuA4 KAT5 lysine acetyltransferase complex

and that the interactions were maintained despite a decrease in cMYC levels after JQ1 treatment.

We conclude that a cascade of aberrant acetyltransferase activities, first via EP300 recruitment and

then by KAT5 interaction with MYC, drive NC cell proliferation and blockade to differentiation.

**Keywords:** NUT carcinoma; cMYC; NuA4 acetyltransferase; KAT5; BioTAP-XL

**Graphical Abstract:**

Promote Proliferation
Block Differentiation

**1. Introduction**:

NUT-rearrangement cancers are a collection of aggressive and poorly-differentiated squamous cell carcinomas whose unique molecular hallmark is somatic expression of *NUTM1* (HGNC:29919)[1]. Chromosomal rearrangements result in the otherwise germline-restricted *NUTM1* gene to be fused downstream of a somatically expressed gene, most commonly *BRD4* (HGNC:13575) to generate the fusion protein BRD4-NUT (B4N). At least two NC fusion proteins: B4N and ZNF532-NUT (a fusion product with *ZNF532* (HGNC:30940)) have been shown to regulate expression of the proto-oncogene *cMYC* (hereafter referred to as *MYC*). This is thought to occur through formation of unusual hyperacetylated chromatin regions termed "megadomains" that can span several megabases [2, 3]. Except for this common genomic region, the hundreds of other B4N and ZNF532-NUT megadomains generally map to distinct regions. Treatment of NC patient cells in culture with a small molecule, JQ1, that inhibits interaction of the BRD4 bromodomains with acetylated lysines, leads to loss of megadomains, down-regulation of *MYC,* and differentiation of B4N cells [2, 4]. The unique importance of *MYC* was revealed in studies where induced *MYC* transgene expression reversed the differentiation induction caused by B4N knockdown [5]. Furthermore, non-B4N-dependent pathways such as *KLF4* (HGNC:6348) expression help maintain *MYC* levels in cells resistant to JQ1 [6]. Thus, understanding the mechanism of action of MYC in NC cells should provide valuable insights into this therapeutically intractable disease.

Much of what is understood about MYC stems from non-NC cells [7-10]. MYC regulates the expression of critical enzymes involved in growth and metabolism, thereby affecting global RNA and protein expression. However, the direct target specificity of MYC remains a subject of debate, in part due to an incomplete picture of the relationship between binding and function. There are

several technical challenges with studying how MYC is assembled with other factors. First, the

MYC protein has a relatively short half-life of approximately 30 minutes, a result of ubiquitin-

dependent and ubiquitin-independent pathways and of the even shorter (10 minute) half-life of

*MYC* RNA itself [11-14]. Second, expression of the *MYC* gene depends on many long-range distal

regulatory elements [15]. These challenges necessitate the ability to inactivate multiple cellular

degradation machinery components and the ability to preserve the complex transcriptional

regulatory circuits of the *MYC* locus. Here, we applied our previously published technique,

BioTAP-XL [3, 16], to characterize the MYC protein and genomic interactions in NC and non-NC

cells.  We found that MYC interacts with subunits of the NuA4 acetytransferase complex in both

cell types [17], and that MYC interactions in NC cells are largely maintained when JQ1 treatment

reduces overall MYC levels, suggesting that protein complex assembly is independent of

differentiation status. These results support models in which chromatin acetylation, especially of

NuA4, plays a central role in the biological function of MYC [18, 19].

## 2. Materials and Methods:

### 2.1 Generation of N-BioTAP-cMYC NC797 cells using CRISPR/Cas9

To generate N-BioTAP and N-ProteinA (PrA) cMYC fusions in NC797 cells [20], the following DNA fragments: pAVV-MCS-NheI-5'cMyc_BamHI-ATG-loxP-Blasti-P2A-loxP-PrA-Bio-3'cMyc and pAAV-5'-cMYC-ATG-LoxP-Blasti-P2A-loxP-PrA-3'-cMYC respectively, were synthesized as gBlock gene fragments (Integrated DNA Technologies, **Datafile S1**) and introduced into the NheI/BamHI restriction sites of pAAV-MCS2 by Gibson assembly (New England Biolabs). N-BioTAP and N-PrA cassettes targeted the first ATG of the *MYC* gene. The pAAV-nEFCas9 (Addgene, plasmid 87115) was used to express Cas9, and the pAAV-tagBFP U6-gRNA expression vector was used to express a gRNA targeting the start codon of cMYC: gRNA-cMYC-start (**Table 1**). Recombinant AAV2 was packaged in HEK293T cells using pHelper and pRC2-mi342 plasmids (Takara, Catalog #632608). Three days after transfection, cells were harvested and AAV2 was isolated using AAVpro Extraction Solution (Takara, Catalog #6235). Two million NC797 cells were infected with 30ul of each adeno-associated virus (AAV2). Following 1 week of culture post-AAV infection, cells were selected with 7.5ug/ml Blasticidin (Invitrogen**).** Surviving cells were re-plated and single-cell clones were isolated and expanded, and genotyping PCR and sequencing were performed to check for proper integration of the BioTAP or PrA cassette in the c-MYC locus. We recovered both heterozygous and homozygous clones. Homozygosity of the cassette insertion was determined by absence of the wildtype PCR fragment (1033bp) and presence of the larger PCR fragment (2083bp) with the following PCR primers: Genom-Myc(1S) and Genom-Myc(1A) (**Table 1**). Next, we validated the presence of the correctly sized tagged MYC protein and absence of wild type MYC by Western blot probed with a c-MYC antibody (Rabbit mAb, Cell Signalling Technology, Cat# 5605). We selected homozygous clone

109    number 2 (for N-BioTAP-cMYC) and clone number 4 (for N-PrA-cMYC) for Cre-dependent

110    cassette excision experiments.

112    **Table 1. Sequences of oligonucleotides used in paper.**

| Name | Sequence (5′ to 3′) |
|---|---|
| gRNA-cMYC-start | GACGTTGAGGGGCATCGTCGC |
| Genom-Myc(1S) | GAGTGGGAACAGCCGCAG |
| Genom_Myc(1A) | CAGGTACAAGCTGGAGGTGG |

Next, the recombinant pAAV-CRE plasmid was packaged in HEK293T cells using pHelper and

pRC2-mi342 plasmids (Takara, Catalog #632608). Cells were harvested three days after

transfection and AAV2 was isolated using AAVpro Extraction Solution (Takara, Catalog #6235).

Two million 797 cells with homozygous insertions (clones 2 or 4) were infected with 30ul of each

adeno-associated CRE virus (AAV2). Following one week of culture post-AAV infection, cells

were re-plated, single-cell clones were isolated and expanded, and genotyping PCR and

sequencing were performed to check for absence of the drug resistance cassette in the c-MYC

locus. We selected clone number 2-7 (for N-BioTAP-cMYC) and clone number 4-6 (for N-PrA-

cMYC) to proceed with LC-MS and ChIP-seq experiments.

*2.2 Generation of inducible N and C-BioTAP-cMYC expressing HEK293T cells*

We used lentiviral transduction to generate stable HEK293T cell lines [2] carrying cMYC

constructs with the BioTAP tag at either the N or the C terminus to induce transcription of these

clones from the CMV/2xtetO promoter for genetic complementation experiments. To make N- and

C-terminal BioTAP constructs, we used the Gateway recombination system to introduce the

following Myc ORF clone HsCD00039771 in pDONR221from the CCSB Human ORFeome

Collection into pHAGE-TRE-DEST-NBioTAP (Addgene #53568) and pHAGE-TRE-DEST-

CBioTAP lentiviral vectors (Addgene #53569). Stable 293T-Trex-cell lines containing each

transgene were generated by lentiviral transduction in the presence of 8μg/mL polybrene, followed

by selection with 2μg/mL puromycin dihydrochloride (Sigma #P8833) as described previously [21].

To induce transcription of the cDNA clones from the CMV/2xtetO promoter we added

doxycycline (1μg/mL) to the medium for 48 hours.

*2.3 Proteomic analysis of cMYC interactions using BioTAP-XL*

All cell lines were cultured as monolayers in DMEM (Invitrogen) supplemented with 1x Penicillin

Streptomycin (Hyclone, South Logan, UT), 1x Glutamax (Gibco), and 10% (v/v) Fetal bovine

serum (FBS) (Hyclone). For JQ1 treatment, 500uM of JQ1 (gift from Dr. C.A. French) was added

to NC797 cells for various durations before harvesting for Western Blot analysis. The 12 hour

timepoint was ultimately chosen for BioTAP-XL experiments.

Our primary experiments were performed with 2-7 N-BioTAP-cMYC (clone 2-7). (2.0 × 10^9)

grown as monolayer cultures in 150-mm dishes. 100 plates total were used for BioTAP-XL

purification. The main steps of the BioTAP-XL procedure were as previously described [21]. For

single-step BioTAP-XL, proteins enriched by IgG-agarose beads were eluted and processed with

Agencourt AMPure-XP magnetic beads (Beckman Coulter) as previously described [22]. In brief,

fifty micrograms of eluted proteins were acidified and diluted with acetonitrile prior to binding by

magnetic beads. SDS was removed with ethanol and acetonitrile washes, and proteins were

digested with overnight trypsin incubation, yielding peptides for desalting and LC-MS analysis.

LC-MS acquisition and analyses were performed as previously described [3].

*2.3 Genomic analysis of cMYC binding*

ChIP-seq analysis was performed in parallel with proteomic analyses as previously described [16].

Raw fastq files were aligned to the human genome (hg38) using bowtie2.3.4.3 with default

settings [23]. Duplicate reads were then removed with samtools1.14 [24]. Peak calling was

performed on DMSO treated samples with HOMER4.9 using the findPeaks command [25].

HOMER annotatePeaks.pl was used to assign peaks to genomic regions. Deeptools3.0.2 was

used to generate bigwig files, enrichment profiles, and heatmaps [26]. Bigwig files were

normalized to 1x genome coverage. For heatmaps and enrichment profiles, MYC signal was

normalized to its corresponding input and enrichment was shown as $\log_2$(IP/input). Differential

peak analysis was performed with the DiffBind package in RStudio using default settings:

(http://bioconductor.org/packages/release/bioc/vignettes/DiffBind/inst/doc/DiffBind.pdf).

Datasets can be obtained from the NCBI GEO database (accession# GSE201033).

**3. Results:**

**3.1 Background on Affinity Approach**

To address the poor stability of MYC during conventional biochemical extraction and the risk of

losing key interactions during DNA removal, we applied a crosslinking approach termed BioTAP-

XL for MYC complex purification in NC797 cells, an NC line, and in HEK293 cells, a non-NC

line for comparison. The BioTAP affinity tag itself is composed of the Z domain from

*Staphylococcal* protein A and a biotinylation target domain from *Propionibacterium*

transcarboxylase. After generating a cell line that expresses the BioTAP-tagged bait, the BioTAP-

XL workflow involves three steps: 1) formaldehyde to inactivate cellular enzymes and to crosslink

proximal protein and nucleic acid interactions, 2) sonication to solubilize the crosslinked

chromatin pellet, and 3) enrichment of the tagged bait using tandem ProteinA and streptavidin

pulldowns combined with stringent washes to reduce nonspecific interactions. This approach has

successfully characterized the proteomic and genomic interactions of B4N, wild-type BRD4,

ZNF532-NUT, wild-type ZNF532, and other proteins [3].

**3.2 Proteomic profiling**

We next characterized the protein interactions of N-BioTAP-MYC in NC797 cells. Using a

CRISPR-Cas9 mediated knock-in approach, we successfully inserted the BioTAP affinity tag into

both copies of the endogenous *cMYC* gene in NC797 cells (see Methods, **Figure 1**). A critical

concern with interpreting affinity tag results is whether the tagged protein preserves the functions

of the non-tagged protein. In NC models such as NC797 cells, B4N regulates transcription of the

*MYC* gene, and NC797 cells are dependent on functional MYC expression [1, 2, 4, 5]. That we could

create viable homozygous knock-in NC797 cell lines expressing only BioTAP- or ProteinA-tagged

MYC strongly suggests that neither epitope tag interferes with normal MYC function, thus

enabling our biochemical and genomic interrogation of MYC using the BioTAP-XL procedure.

**Figure 1.** Western blot of NC797 cells carrying homozygous knock-in alleles of BioTAP-tagged

MYC probed with anti-MYC (D84C12 Rabbit mAb, Cell Signalling Technologies). The untagged

size of MYC in unedited NC797 cells is ~57 Kd, (lane 1). The size of N-BioTAP-MYC is ~80

Kd, seen in Clones 16 and 2 (lanes 2 and 3). Clone 2 was used for subsequent experiments.

We performed BioTAP-XL to enrich for MYC containing protein complexes and used liquid

chromatography-mass spectrometry (LC-MS) to sequence the peptides resulting from trypsin

digestion of the enriched complexes. MYC protein interactions were identified as the proteins with

the greatest fold enrichment in total peptides identified in the pulldown compared to the input

sample (**Datafile S2**). As expected, we recovered both MYC and its partner MAX as among the

top twenty-three most enriched proteins (corresponding to approximately the top one to two

percent). Beyond MYC and MAX, eleven of the most enriched proteins are related to chromatin

function. In particular, eight are components of the NuA4 histone acetyltransferase complex [17, 27]

(**Figure 2**). The enriched components include DMAP1 (HGNC:18291), EP400 (HGNC:11958),

EPC1 (HGNC:19876), EPC2 (HGNC:24543), KAT5 (HGNC:5275), MBTD1 (HGNC:19866),

TRRAP (HGNC:12347), and YEATS4 (HGNC:24859).

A

B

| Name | MolWt | Input | JQ1 input | MYC BioTAP | | MYC PrA Vehicle | | MYC PrA JQ1 | | No Bait IgG |
|---|---|---|---|---|---|---|---|---|---|---|
| | | | | Rep #1 | Rep #2 | Rep #1 | Rep #2 | Rep #1 | Rep #2 | |
| MYC | 50.53 | 0 | 0 | 3 | 3 | 7 | 8 | 3 | 6 | 0 |
| MAX | 18.26 | 0 | 0 | 3 | 4 | 7 | 9 | 6 | 7 | 0 |
| DMAP1 | 52.96 | 2 | 2 | 9 | 9 | 9 | 8 | 5 | 6 | 0 |
| EP400 | 343.28 | 8 | 7 | 58 | 66 | 53 | 46 | 35 | 29 | 0 |
| EPC1 | 93.41 | 0 | 0 | 7 | 10 | 4 | 7 | 2 | 4 | 0 |
| EPC2 | 91.04 | 0 | 0 | 9 | 8 | 10 | 8 | 6 | 7 | 0 |
| ING3 | 46.71 | 0 | 0 | 7 | 6 | 6 | 7 | 5 | 4 | 0 |
| KAT5 | 61.76 | 1 | 0 | 10 | 8 | 4 | 6 | 1 | 3 | 0 |
| MBTD1 | 70.5 | 0 | 0 | 5 | 5 | 3 | 2 | 2 | 3 | 0 |
| TRRAP | 437.32 | 9 | 8 | 106 | 95 | 87 | 85 | 52 | 79 | 0 |
| VPS72 | 41.82 | 0 | 1 | 4 | 5 | 4 | 4 | 2 | 3 | 0 |
| YEATS4 | 26.48 | 1 | 0 | 6 | 8 | 2 | 3 | 2 | 2 | 0 |
| ZNHIT6 | 53.88 | 0 | 0 | 3 | 5 | 1 | 1 | 0 | 1 | 0 |

**Figure 2.** Characterization of MYC protein interactions in NC797 cells.

(a) Scatterplots depicting enrichment of proteins from N-terminally tagged MYC using BioTAP-XL workflow in NC797 cells compared to non-enriched input. Each gray dot represents an individually identified protein, where its position is a measure of enrichment efficiency based on the number of total peptides identified in pulldown compared to input and normalized to the molecular weight of the protein from two technical replicates. The dashed gray lines denote the 98th percentile of enrichment. The right plot is a zoomed in region of the most enriched proteins in the left plot from replicate experiments. (b) Number of total peptides mapping to each protein from a given pulldown experiment.

To test whether the NuA4 interaction would change upon B4N inactivation and the resulting

cellular differentiation, we performed affinity enrichment of MYC after cells were treated with the

bromodomain inhibitor JQ1. We knew from prior experience that the amount of MYC bait protein

expression after inactivation of B4N would be significantly decreased, which might be insufficient

for a full tandem affinity purification. Therefore we modified our approach to utilize a single rather

than dual tag, utilizing NC797 clones in which the endogenous gene had only the ProteinA portion

of the affinity tag (PrA-MYC). Consistent with B4N directly regulating the endogenous *MYC*

locus, treatment of NC797 cells with JQ1 reduced levels of PrA-MYC in a time-dependent manner

(**Figure 3**). Interestingly, despite choosing a 12 hour timepoint that corresponded to a five-fold

reduction in PrA-MYC levels, we found the NuA4 complex interactions with MYC were largely

preserved after affinity enrichment (**Figure 2B**). To test whether recovery of NuA4 components is

due to non-specific capture of the proteins by the IgG agarose beads, we performed a parallel

pulldown with NC797 cells expressing no tagged bait. We observed no recovery of peptides from

MYC or NuA4 components (**Figure 2B**).

**Figure 3**. Effect of JQ1 treatment on MYC protein expression. Levels of MYC protein in N-PrA-cMyc-NC797 cells were assayed by Western blot, following different durations of 500uM JQ1 treatment, with approximate fold change in MYC levels provided below the plot compared to no treatment control.

Finally, to test whether the NuA4 complex interaction was NC-cell dependent, we performed the

BioTAP-XL experiment in the non-NC cell line HEK293-TREx. Similar to NC797 cells, we found

enrichment of NuA4 components with an inducibly expressed MYC-BioTAP bait in HEK293-

TREx cells (**Datafile S3**). Collectively, our results suggest that in NC797 patient cells the most

significant protein interactions of MYC besides MAX are members of the NuA4 acetyltransferase

complex. These interactions are maintained during differentiation induced by JQ1 treatment.

**3.3 Genomic profiling**

In parallel with the proteomic analyses, we mapped the genomic occupancy of PrA-MYC in

NC797 cells following JQ1 or DMSO treatment, to determine if its set of target genes changed as

its levels decreased. Differential binding analysis resulted in 2,238 regions with significantly

altered MYC binding (FDR < 0.05) (**Figure 4A,B**). Consistent with the decrease in MYC levels,

a majority of MYC-bound regions showed decreased occupancy (n=2,234). Significantly

downregulated regions were located at promoters (25%, n=555/2,234), introns (40%,

n=900/2,234), and intergenic regions (28%, n=623/2,234) (**Figure 4C**). However, while MYC

occupancy was diminished, the identities of MYC binding sites were not globally changed by

JQ1 treatment (**Figure 4B,D**). Our comparison of proliferating and differentiating NC cells

suggests that MYC-dependent changes in cell state generally occur without alteration of key

MYC protein interactions or MYC binding sites.

**Figure 4.** Effect of JQ1 on MYC genomic binding.

(a.) MA plot of MYC differential binding after JQ1 treatment. Each point represents a 400bp MYC-bound regions. Blue points indicate significantly altered regions (FDR < 0.05). The x-axis depicts the $\log_{10}$ of the average read concentration across all samples. The y-axis indicates the fold change after JQ1 treatment. (b.) Enrichment profile plotting the average signal of significantly decreased MYC-bound regions in DMSO and JQ1 treated samples. One representative replicate is shown. (c.) Pie chart depicting the percentage of significantly decreased MYC-bound regions mapped to the specified genomic regions. (d.) Heatmap of MYC enrichment at significantly decreased MYC-bound regions in DMSO and JQ1 treated samples. One representative replicate is shown.

**4. Discussion:**

To characterize MYC protein partners in NC797 cells, we applied a crosslinking affinity pulldown method. We took this approach because, historically, MYC complexes have been difficult to purify. Our data suggest significant interactions between MYC and NuA4 complex components. The canonical NuA4 complex is comprised of the catalytic subunit KAT5, also known as TIP60, and at least four non-catalytic subunits: EPC1, TRAAP, BAF53, DMAP1. Our results are consistent with the links between MYC and histone acetyltransferase activity that were discovered in some of the earliest studies of MYC function [18, 28-32], and more recent proteomic analyses using targeted biotinylation [27] or BioID [33] to identify protein interactors in mouse and human cell lines, respectively. Our datasets and methods should be useful in further study of MYC and downstream targets specifically in NC cells. Furthermore, demonstrations that MYC is dependent on acetyltransferases for its function suggest potential therapeutic options to what is generally considered an undruggable oncoprotein [19, 34]. If specific inhibitors of KAT5 can be developed, they might synergize with JQ1 in the potential future treatment of NC.

**5. Conclusion:**

MYC is a central player in a majority of cancers, yet understanding its regulation, targeting, and mode of action are important unsolved problems. When we initiated this study, we knew that the B4N oncogenic fusion attracted the p300 acetyltransferase to drive oncogenesis, with the Myc gene as a critical target [1]. Since MYC function in NC cells is reversible by the JQ1 small molecule BET bromodomain inhibitor, we saw this as an opportunity to compare MYC interactions in the same cells under oncogenic and differentiating conditions. While we did not find a difference in protein interactions or gene targeting in the distinct growth states, we confirmed a close association

299 between MYC and the NuA4 acetyltransferase in NC cells. We propose that future analysis of

300 MYC transcriptional regulation in NUT carcinoma, and likely other cancers, should prioritize how

301 MYC recruits KAT complexes to modify the chromatin of its many gene targets.

**Supplementary Materials:**

304 The following supporting information can be downloaded at: www.mdpi.com/xxx/s1:

305 Datafile S1: DNA sequences used to create NC797 cells with N-terminal fusions of the BioTAP-

306 or Protein A-epitope tags to the endogenous cMYC coding region

307 Datafile S2: Total peptide counts of NC797 proteomic results

308 Datafile S3: Total peptide counts of HEK293T proteomic results

**Author Contributions:**

311 Conceptualization, AAA and MIK; methodology, validation, formal analysis, investigation, BMZ

312 and AAA; resources, data curation, BMZ and AES; writing – original draft preparation BMZ and

313 AAA; writing – review and editing, BMZ and MIK; visualization, BMZ, AAA, AES; supervision,

314 project administration, funding acquisition, MIK. All authors have read and agreed to the

315 published version of the manuscript.

**Funding:**

318 This research was funded by the National Institutes of Health, grant number R35 GM126944 to

319 MIK. AES is supported by an American Cancer Society postdoctoral fellowship 721690.

**Data Availability Statement**

Proteomic data sets can be found at Harvard Dataverse (https://doi.org/10.7910/DVN/IFKRBT).

Genomic data sets can be found at NCBI GEO database (accession# GSE201033).

**Acknowledgments**

We thank Dr. C.A. French for introducing us to NC research, and Dr. J.L. Makofske for initial work on this project. We thank Ross Tomaino (Taplin Mass Spectrometry Facility, Harvard Medical School) for expert help with analyzing the proteomics samples. We thank Drs. T. Martin and S.J. Elledge for the pAAV-tagBFP U6-gRNA expression vector. Finally, we thank H.J. Kang for helpful assistance with data submission.

**Conflicts of Interest**

The authors declare no conflict of interest.

**References**

1. French, C. A., NUT Carcinoma: Clinicopathologic features, pathogenesis, and treatment. *Pathol Int* **2018,** *68* (11), 583-595.
2. Alekseyenko, A. A.; Walsh, E. M.; Wang, X.; Grayson, A. R.; Hsi, P. T.; Kharchenko, P. V.; Kuroda, M. I.; French, C. A., The oncogenic BRD4-NUT chromatin regulator drives aberrant transcription within large topological domains. *Genes Dev* **2015,** *29* (14), 1507-23.
3. Alekseyenko, A. A.; Walsh, E. M.; Zee, B. M.; Pakozdi, T.; Hsi, P.; Lemieux, M. E.; Dal Cin, P.; Ince, T. A.; Kharchenko, P. V.; Kuroda, M. I.; French, C. A., Ectopic protein interactions within BRD4-chromatin complexes drive oncogenic megadomain formation in NUT midline carcinoma. *Proc Natl Acad Sci U S A* **2017,** *114* (21), E4184-E4192.
4. Filippakopoulos, P.; Qi, J.; Picaud, S.; Shen, Y.; Smith, W. B.; Fedorov, O.; Morse, E. M.; Keates, T.; Hickman, T. T.; Felletar, I.; Philpott, M.; Munro, S.; McKeown, M. R.; Wang, Y.; Christie, A. L.; West, N.; Cameron, M. J.; Schwartz, B.; Heightman, T. D.; La Thangue, N.; French, C. A.; Wiest, O.; Kung, A. L.; Knapp, S.; Bradner, J. E., Selective inhibition of BET bromodomains. *Nature* **2010,** *468* (7327), 1067-73.
5. Grayson, A. R.; Walsh, E. M.; Cameron, M. J.; Godec, J.; Ashworth, T.; Ambrose, J. M.; Aserlind, A. B.; Wang, H.; Evan, G.; Kluk, M. J.; Bradner, J. E.; Aster, J. C.; French, C.

A., MYC, a downstream target of BRD-NUT, is necessary and sufficient for the blockade of differentiation in NUT midline carcinoma. *Oncogene* **2014,** *33* (13), 1736-1742.

6. Liao, S.; Maertens, O.; Cichowski, K.; Elledge, S. J., Genetic modifiers of the BRD4-NUT dependency of NUT midline carcinoma uncovers a synergism between BETis and CDK4/6is. *Genes Dev* **2018,** *32* (17-18), 1188-1200.

7. Meyer, N.; Penn, L. Z., Reflecting on 25 years with MYC. *Nature reviews. Cancer* **2008,** *8* (12), 976-90.

8. Dang, C. V., MYC on the path to cancer. *Cell* **2012,** *149* (1), 22-35.

9. Kress, T. R.; Sabo, A.; Amati, B., MYC: connecting selective transcriptional control to global RNA production. *Nature reviews. Cancer* **2015,** *15* (10), 593-607.

10. Muhar, M.; Ebert, A.; Neumann, T.; Umkehrer, C.; Jude, J.; Wieshofer, C.; Rescheneder, P.; Lipp, J. J.; Herzog, V. A.; Reichholf, B.; Cisneros, D. A.; Hoffmann, T.; Schlapansky, M. F.; Bhat, P.; von Haeseler, A.; Kocher, T.; Obenauf, A. C.; Popow, J.; Ameres, S. L.; Zuber, J., SLAM-seq defines direct gene-regulatory functions of the BRD4-MYC axis. *Science* **2018,** *360* (6390), 800-805.

11. Hann, S. R.; Eisenman, R. N., Proteins encoded by the human c-myc oncogene: differential expression in neoplastic cells. *Mol Cell Biol* **1984,** *4* (11), 2486-97.

12. Hickman, K. A.; Hariharan, S.; De Melo, J.; Ylanko, J.; Lustig, L. C.; Penn, L. Z.; Andrews, D. W., Image-Based Analysis of Protein Stability. *Cytometry A* **2020,** *97* (4), 363-377.

13. Murai, N.; Murakami, Y.; Tajima, A.; Matsufuji, S., Novel ubiquitin-independent nucleolar c-Myc degradation pathway mediated by antizyme 2. *Sci Rep* **2018,** *8* (1), 3005.

14. Dani, C.; Blanchard, J. M.; Piechaczyk, M.; El Sabouty, S.; Marty, L.; Jeanteur, P., Extreme instability of myc mRNA in normal and transformed human cells. *Proc Natl Acad Sci U S A* **1984,** *81* (22), 7046-50.

15. Cole, M. D., MYC association with cancer risk and a new model of MYC-mediated repression. *Cold Spring Harb Perspect Med* **2014,** *4* (7), a014316.

16. Alekseyenko, A. A.; McElroy, K. A.; Kang, H.; Zee, B. M.; Kharchenko, P. V.; Kuroda, M. I., BioTAP-XL: Cross-linking/Tandem Affinity Purification to Study DNA Targets, RNA, and Protein Components of Chromatin-Associated Complexes. *Curr Protoc Mol Biol* **2015,** *109*, 21 30 1-21 30 32.

17. Doyon, Y.; Cote, J., The highly conserved and multifunctional NuA4 HAT complex. *Curr Opin Genet Dev* **2004,** *14* (2), 147-54.

18. Frank, S. R.; Parisi, T.; Taubert, S.; Fernandez, P.; Fuchs, M.; Chan, H. M.; Livingston, D. M.; Amati, B., MYC recruits the TIP60 histone acetyltransferase complex to chromatin. *EMBO Rep* **2003,** *4* (6), 575-80.

19. Numata, A.; Kwok, H. S.; Zhou, Q. L.; Li, J.; Tirado-Magallanes, R.; Angarica, V. E.; Hannah, R.; Park, J.; Wang, C. Q.; Krishnan, V.; Rajagopalan, D.; Zhang, Y.; Zhou, S.; Welner, R. S.; Osato, M.; Jha, S.; Bohlander, S. K.; Gottgens, B.; Yang, H.; Benoukraf, T.; Lough, J. W.; Bararia, D.; Tenen, D. G., Lysine acetyltransferase Tip60 is required for hematopoietic stem cell maintenance. *Blood* **2020,** *136* (15), 1735-1747.

20. Toretsky, J. A.; Jenson, J.; Sun, C. C.; Eskenazi, A. E.; Campbell, A.; Hunger, S. P.; Caires, A.; Frantz, C.; Hill, J. L.; Stamberg, J., Translocation (11;15;19): a highly specific chromosome rearrangement associated with poorly differentiated thymic carcinoma in young patients. *Am J Clin Oncol* **2003,** *26* (3), 300-6.

21. Alekseyenko, A. A.; Gorchakov, A. A.; Kharchenko, P. V.; Kuroda, M. I., Reciprocal
interactions of human C10orf12 and C17orf96 with PRC2 revealed by BioTAP-XL cross-
linking and affinity purification. *Proc Natl Acad Sci U S A* **2014,** *111* (7), 2488-93.
22. Zee, B. M.; Alekseyenko, A. A.; McElroy, K. A.; Kuroda, M. I., Streamlined discovery of
cross-linked chromatin complexes and associated histone modifications by mass spectrometry.
*Proc Natl Acad Sci U S A* **2016,** *113* (7), 1784-9.
23. Langmead, B.; Salzberg, S. L., Fast gapped-read alignment with Bowtie 2. *Nat Methods* **2012,**
*9* (4), 357-9.
24. Danecek, P.; Bonfield, J. K.; Liddle, J.; Marshall, J.; Ohan, V.; Pollard, M. O.; Whitwham,
A.; Keane, T.; McCarthy, S. A.; Davies, R. M.; Li, H., Twelve years of SAMtools and
BCFtools. *Gigascience* **2021,** *10* (2).
25. Heinz, S.; Benner, C.; Spann, N.; Bertolino, E.; Lin, Y. C.; Laslo, P.; Cheng, J. X.; Murre,
C.; Singh, H.; Glass, C. K., Simple combinations of lineage-determining transcription factors
prime cis-regulatory elements required for macrophage and B cell identities. *Mol Cell* **2010,**
*38* (4), 576-89.
26. Ramirez, F.; Ryan, D. P.; Gruning, B.; Bhardwaj, V.; Kilpert, F.; Richter, A. S.; Heyne, S.;
Dundar, F.; Manke, T., deepTools2: a next generation web server for deep-sequencing data
analysis. *Nucleic Acids Res* **2016,** *44* (W1), W160-5.
27. Kim, J.; Woo, A. J.; Chu, J.; Snow, J. W.; Fujiwara, Y.; Kim, C. G.; Cantor, A. B.; Orkin,
S. H., A Myc network accounts for similarities between embryonic stem and cancer cell
transcription programs. *Cell* **2010,** *143* (2), 313-24.
28. McMahon, S. B.; Wood, M. A.; Cole, M. D., The essential cofactor TRRAP recruits the
histone acetyltransferase hGCN5 to c-Myc. *Mol Cell Biol* **2000,** *20* (2), 556-62.
29. Bouchard, C.; Dittrich, O.; Kiermaier, A.; Dohmann, K.; Menkel, A.; Eilers, M.; Luscher,
B., Regulation of cyclin D2 gene expression by the Myc/Max/Mad network: Myc-dependent
TRRAP recruitment and histone acetylation at the cyclin D2 promoter. *Genes Dev* **2001,** *15*
(16), 2042-7.
30. Frank, S. R.; Schroeder, M.; Fernandez, P.; Taubert, S.; Amati, B., Binding of c-Myc to
chromatin mediates mitogen-induced acetylation of histone H4 and gene activation. *Genes Dev*
**2001,** *15* (16), 2069-82.
31. Fernandez, P. C.; Frank, S. R.; Wang, L.; Schroeder, M.; Liu, S.; Greene, J.; Cocito, A.;
Amati, B., Genomic targets of the human c-Myc protein. *Genes Dev* **2003,** *17* (9), 1115-29.
32. Knoepfler, P. S.; Zhang, X. Y.; Cheng, P. F.; Gafken, P. R.; McMahon, S. B.; Eisenman, R.
N., Myc influences global chromatin structure. *Embo J* **2006,** *25* (12), 2723-34.
33. Dingar, D.; Kalkat, M.; Chan, P. K.; Srikumar, T.; Bailey, S. D.; Tu, W. B.; Coyaud, E.;
Ponzielli, R.; Kolyar, M.; Jurisica, I.; Huang, A.; Lupien, M.; Penn, L. Z.; Raught, B., BioID
identifies novel c-MYC interacting partners in cultured cells and xenograft tumors. *J*
*Proteomics* **2015,** *118*, 95-111.
34. Mustachio, L. M.; Roszik, J.; Farria, A.; Dent, S. Y. R., Targeting the SAGA and ATAC
Transcriptional Coactivator Complexes in MYC-Driven Cancers. *Cancer Res* **2020,** *80* (10),
1905-1911.
438