A method to enrich polypeptidyl-tRNAs to capture snapshots of translation in the cell

Ayako Yamakawa[1], Tatsuya Niwa[1,2,*], Yuhei Chadani[2,*] and Hideki Taguchi[1,2,*]

[1] School of Life Science and Technology, Tokyo Institute of Technology, S2-19, Nagatsuta 4259, Midori-ku, Yokohama, 226-8503, Japan.

[2] Cell Biology Center, Institute of Innovative Research, Tokyo Institute of Technology, S2-19, Nagatsuta 4259, Midori-ku, Yokohama, 226-8503, Japan.

*Correspondence to: tniwa@bio.titech.ac.jp, chadani.y.aa@m.titech.ac.jp, taguchi@bio.titech.ac.jp

Abstract

Life depends on proteins, which all exist in nascent states when the growing polypeptide chain is covalently attached to a tRNA within the ribosome. Although the nascent chains; *i.e.,* polypeptidyl-tRNAs (pep-tRNAs), are considered as merely transient intermediates during protein synthesis, recent advances have revealed that they are directly involved in a variety of cell functions, such as gene expression control. An increasing appreciation for fine-tuning at translational levels demands a general method to handle the pep-tRNAs on a large scale. Here, we developed a method to enrich the pep-tRNAs by organic solvent extraction and silica column separation, and then identify their polypeptide moieties by mass spectrometry. As a proof-of-concept experiment using *Escherichia coli*, we identified ~500 peptides derived from the pep-tRNAs, which were markedly biased towards the N-terminal regions, reflecting that the method captured the intermediate pep-tRNA population during translation. Furthermore, the method identified a subset of pep-tRNAs that were resistant to puromycin treatment, suggesting that these pep-tRNAs accumulated due to nascent-chain dependent translational pausing. Although the current specificity and comprehensiveness of the pep-tRNA enrichment have room for improvement, this new method will complement conventional methods for profiling nascent chains.

Significance Statement

In the central dogma of biology, RNA, an informational molecule, and protein, a functional molecule, are usually regarded as two completely independent molecular species. However, they are combined into a single species called peptidyl-tRNA (pep-tRNA) during the translation process in the ribosome. Despite the importance of pep-tRNAs as precursors of all proteins in the cell, a general method to analyze pep-tRNAs on a large scale was lacking. Taking advantage of the properties of pep-tRNAs as RNA and protein, we developed a method termed *pe*ptidyl-*t*RNA *e*nrichment using *o*rganic extraction and *s*ilica adsorption (PETEOS). PETEOS not only provides a unique approach to examine the nascent state of proteins but also may be effective in exploring peptides that cause translational pauses.

Keywords: translation, nascent polypeptide chain, polypeptidyl-tRNA, proteomics, ribosome, mass spectrometry

Introduction

Every protein in the cell is generated through translation. The formation of the proteome involves the regulation of various steps in gene expression. In addition to the well-known importance of gene regulation at transcriptional levels, gene expression control at translational levels has been increasingly appreciated (1, 2).

The translational control is diverse, flexible, and rapid, depending on cellular contexts or stimuli such as a variety of stresses. In addition to the control at the translation initiation step, which is a major rate-determining step in translation, even the translation elongation could be regulated (3, 4). One of the fine-tuning events is elongation rate control, as exemplified by elongation pausing, which can affect the folding, function, and quality control of proteins (3–7). Besides the pausing induced by mRNA secondary structures or tRNA shortages (4, 5, 7, 8), both of which could be essentially predictable, nascent polypeptides within the ribosome could pause the elongation (9–12). For example, the translation product of the gene *secM* in *Escherichia coli* induces translation arrest (13, 14). The ribosomal arrest peptide of SecM, still tethered to its tRNA as a polypeptidyl-tRNA (pep-tRNA), interacts with the inner wall of the ribosomal exit tunnel (3, 14, 15). The translation arrest by SecM is directly involved in the feedback regulation of its downstream gene, *secA* (13, 14). Other genes identified in bacteria and eukaryotes also cause translation arrest (16–19). In addition to the ad hoc identification, biochemical analyses of over 1,000 *E. coli* genes revealed that a majority of them indeed undergo single or multiple pauses in translation, indicating the widespread occurrence of translational pausing (20). Notably, more than half of the pauses are resistant to puromycin (20), suggesting that the ribosomal function has been impaired, as shown for the arrest by SecM (21). However, so far, a comprehensive picture of amino acid sequence motifs that induce pausing remains obscure. Collectively, there is an increasing demand for methods to globally investigate the translation process in the cell.

So far, two main methodologies based on completely different concepts are used to survey the translation process in cells (22). One is ribosome profiling, a deep sequencing technique for ribosome-protected mRNA fragments generated by RNase digestion (23). Ribosome profiling is a very powerful genome-wide analysis with nucleotide resolution, and has been established as a method to monitor the translation status in cells (22–27). However, this method, which relies on a huge amount of "ribosome footprint" sequencing data, is inherently indirect in detecting the nascent proteins or pep-tRNAs, and might cause some controversy about how to collect and interpret the data (*e.g.*, 28). In addition, the method is still time-consuming and costly.

The other methodology is based on proteomics with liquid chromatography-tandem mass

spectrometry (LC-MS/MS). Several "translatome" methods have been developed to distinguish nascent proteins from the pre-existing protein population. The proteomics-based approaches rely on the labeling of nascent polypeptide chains, using noncanonical amino acids such as azide homoalanine (*e.g.,* BONCAT (29), QuaNCAT (30)) and puromycin analogs (*e.g.,* PUNCH-P (31, 32), OPP-ID (33)). Since the vast majority of nascent proteins labeled by the noncanonical amino acids have already been released from the ribosome, these proteins are newly but already "synthesized" mature proteins, and not nascent proteins during translation. The puromycin-related methods rely on the principle that puromycin or its analogs terminate translation by entering the A-site of the ribosome, and are then covalently incorporated into the growing end of the nascent peptide moiety in pep-tRNAs. However, certain nascent peptides are resistant to puromycin (16, 18, 21). In addition, all of these methods that rely on labeling usually require an hour-long timescale or more to accumulate the labeled products (34). In summary, these analytical methods for the nascent proteome do not directly deal with pep-tRNAs and do not capture the moments of the translation elongation status in the cell.

In view of these points, a direct methodology to handle pep-tRNAs is required to complement the prevailing methods for the translatome analyses. Although the point is not always appreciated, it is obvious that all proteins are born via pep-tRNAs, which possess the properties of both RNA and protein. In other words, pep-tRNA is an essential translation intermediate that lies between RNA and protein in the central dogma of biology. Despite their importance in life, pep-tRNAs are low-abundant. Assuming that the approximate numbers of proteins and ribosomes in *E. coli* are ~3 million and ~70,000, respectively (35), the maximum amount of pep-tRNAs is at most only ~2% of total proteins, even if all ribosomes participate in translation.

These "in-between" properties of both RNA and protein and the low abundance make pep-tRNAs challenging to handle on the "pep-tRNA-ome" scale. Ito *et al.* developed an ingenious method to collectively detect cellular pep-tRNAs, the "nascentome," using a combination of pulse-labeling and two-dimensional electrophoresis (36). The "nascentome" analysis can detect the whole pep-tRNAs but cannot identify which protein is part of each pep-tRNA. Therefore, we developed a non-labeling method to enrich and identify pep-tRNAs in cells. Briefly, pep-tRNAs in cells are enriched by organic solvent extraction, followed by isolation on a silica column. The polypeptide moieties in the enriched pep-tRNAs are cleaved under high pH and high temperature conditions. The resultant polypeptides are digested by proteases such as trypsin and then identified using conventional LC-MS/MS-based shotgun proteomics. This method, termed *pe*ptidyl-*t*RNA *e*nrichment using *o*rganic extraction and *s*ilica adsorption (PETEOS), is not only complementary to prevailing translation measurement methods, but also

103   has, in principle, several advantages such as rapid capturing of translation status, non-labeling
104   feature, and compatibility with puromycin treatment to detect elongation pausing candidates.
105
Results
Polypeptidyl-tRNA enrichment and identification using a proteomics approach.
The chemical properties of pep-tRNAs with various polypeptide moieties are diverse. For
example, when the peptide moiety is hydrophilic or relatively short, the hydrophilic property
of the RNA is dominant, but when the polypeptide portion is more hydrophobic or longer, the
more protein-like property is manifested. As far as we know, there is no established method to
purify whole pep-tRNAs from cells. Our strategy to enrich pep-tRNAs takes advantage of their
two distinctive properties as RNA and polypeptide. The method, which we call PETEOS, is
composed of four steps (Fig. 1). Step 1: Enrichment of pep-tRNAs and proteins from cells.
Harvested cells are directly mixed into a phenol-containing reagent (TriPURE) to separate most
RNAs (aqueous phase) from pep-tRNAs and proteins (organic phase). Note that there is no
need for labeling times in this method. After the chloroform addition, the pep-tRNAs and
proteins in the organic phase are precipitated by trichloroacetic acid (TCA)/acetone, under
conditions where the ester bonds between polypeptide and tRNA in the pep-tRNAs are retained.
Step 2: Isolation of the pep-tRNAs. The RNA is isolated by using silica to adsorb the nucleic
acid portions of pep-tRNAs, under highly concentrated chaotropic salt conditions (37, 38). Step
3: Cleavage of peptide moieties from peptidyl-tRNAs. The ester bond between peptide and
tRNA is cleaved by high pH and heat treatment. Step 4: Identification of the peptides after
trypsin treatment, by LC-MS/MS-based shotgun proteomics.
125
Proof-of-concept analysis using *Escherichia coli* cell lysates.
The validity of the PETEOS method was evaluated at each step. An LC-MS/MS analysis
of the TCA/acetone precipitate in Step 1 revealed that ~79% of the identified proteins in the
precipitate overlapped with proteins identified by a conventional method (39) to extract proteins
from cells (Fig. S1), confirming that the protein fraction in the organic phase seems to lack
significant bias. Next, when we analyzed the eluate at Step 2 by SDS-PAGE under neutral pH
conditions (neutral-PAGE), where the ester bonds between peptide and tRNA in the pep-tRNAs
remain intact, smeared silver-stained bands were observed (Fig. 2A). Since an RNase treatment
of the mixture at Step 1 before the adsorption to the silica column drastically reduced the
smeared bands (Fig. 2A), they are derived from pep-tRNA species.
After the peptide moieties from the pep-tRNAs were obtained by the alkaline and heat
treatment (Step 3), the LC-MS/MS analysis (Step 4) identified more than 500 proteins

Figure 1. PETEOS workflow to enrich and identify polypeptidyl-tRNAs (pep-tRNAs) in cells
Step 1: Enrichment of pep-tRNAs and proteins. Log phase *E. coli* cells ($OD_{660} \approx 0.5$) cultured at 37 °C are extracted by phenol/chloroform to separate the hydrophilic RNA molecules into the aqueous phase. Relatively hydrophobic proteins and pep-tRNAs in the organic phase are precipitated by TCA/acetone. **Step 2**: Isolation of the pep-tRNAs by silica column chromatography in the presence of concentrated chaotropic salts. **Step 3**: Hydrolysis of the ester bond between peptide and tRNA in pep-tRNA under high pH and high temperature conditions. **Step 4**: Identification of the peptide moieties of the pep-tRNAs by conventional shotgun proteomics after trypsin digestion.

(Fig. 2B). The number of identified proteins was reduced to ~50 proteins in the RNase-treated fraction after Step 1 (Fig. 2B), indicating that ~90% of the identified proteins would be derived from pep-tRNAs. In addition, the intensity distribution of the MS/MS-identified peptides was significantly higher in the proteins obtained without the RNase treatment (Fig. 2C). Collectively, the PETEOS method can enrich and identify the cellular pep-tRNAs.

Next, we validated the potential of the PETEOS method to identify peptides paused at a specific position within the ribosome. A particular length of peptides in a pep-tRNA mixture would accumulate if the C-terminal end of the pausing peptide is tethered to a tRNA at the pausing site. Accordingly, we overexpressed a well-known pausing peptide, SecM. The translation of the gene *secM* is arrested at the 165th glycine, where tRNA^{Gly} is covalently attached to the elongating end of the nascent peptide (SecM$_{1-165}$-tRNA^{Gly}). As expected, the MS/MS analysis identified the SecM-derived pausing peptide ($_{150}$FSTPVWISQAQGIRAG$_{165}$) from the SecM-overexpressing cells (Fig. 2D), indicating that the method could identify paused peptides at an amino acid sequence resolution. We also overexpressed the known pausing peptide, TnaC ($_1$MNILHICVTS KWFNIDNKIV DHRP$_{24}$-stop codon), which induces ribosome pausing at the C-terminus. The MS/MS analysis identified a TnaC peptide ($_{12}$WFNIDNK$_{18}$), which would be derived from the trypsinized product of the TnaC pausing

peptide. Note that the MS analysis also detected a TnaC-derived peptide in the aqueous phase
at Step 1, indicating that the separation of the TnaC-tRNA is incomplete with the
phenol/chloroform extraction. The insufficient separation is probably due to the hydrophilic
nature of the TnaC-tRNA, since its polypeptide moiety is relatively short (24 amino acids).
Indeed, a silver-stained gel revealed the aqueous layer-biased accumulation of pep-tRNAs in
the TnaC-overexpressing cells (Fig. S2).

A previous nascentome study showed that Tris-base in the buffer stimulates the ester bond
cleavage of pep-tRNAs (36). A nucleophilic attack of tris(hydroxymethyl)aminomethane (Tris)
on the carboxyl group of the C-terminal amino acid, which is ester-bonded to tRNA in the pep-
tRNA, produces a Tris-adduct during the pep-tRNA hydrolysis (Fig. S3). Therefore, we
speculated that some peptides derived from pep-tRNAs could become conjugated with Tris. A
peptide search assuming the Tris-addition at the C-terminus revealed ~10 Tris-conjugated
peptides, including a SecM-derived paused peptide (Fig. 2D, Table S1). Since the conjugation
of the Tris-base confirmed the handling of intact pep-tRNAs until they are exposed to high pH
and high temperature, the production of these Tris-replaced fragments is one of the hallmarks
of the PETEOS method.

Figure 2. Enrichment and identification of pep-tRNAs in *E. coli*
A. The eluate from the silica column (Step 2) was analyzed by neutral pH SDS-PAGE, and detected by silver staining. The eluate treated with RNase after Step 1 was also analyzed. Pep-tRNA molecules are schematically indicated (tRNA moiety, orange; peptide moiety, blue). **B**. Number of proteins identified as pep-tRNA molecules in the method. The peptides detected more than once in the MS/MS analysis were counted. Among the 48 proteins identified in the RNase-treated fraction, 46 overlapped with the proteins derived from pep-tRNAs. **C**. Histogram of summed peptide intensity per protein ($\log_{10}$ scale) obtained by the MS1 peaks for proteins from the silica column eluate or the RNase-treated fraction. **D**. MS2 spectra of the C-terminal peptides of SecM from *E. coli* cells overexpressing SecM. (*Left*) $_{150}$FSTPVWISQAQGIRAG$_{165}$. (*Right*) The C-terminal peptide modified by Tris-base, $_{150}$FSTPVWISQAQGIRAG$_{165}$-Tris.

The polypeptidyl-tRNA population reflects the translation intermediates.

The MS-identified peptides derived from the pep-tRNAs in *E. coli* confirmed that the peptides are intermediate species during the translation elongation. Since nascent peptides elongate from the N- to C-terminus, we expected that the tryptic peptides derived from pep-tRNAs would be enriched with the N-terminus as compared to those derived from pre-existing total cellular proteins (Fig. 3A). A meta-analysis demonstrated that the tryptic peptides from the pep-tRNAs were highly biased toward the N-terminal regions (Fig. 3B). In contrast, a meta-analysis of the total proteins showed equally distributed tryptic peptides (Fig. 3B), as expected since the total proteins are basically composed of the full-length polypeptides after translation.

In addition, a substantial population of pep-tRNAs (228 out of 515) was not identified in the cells treated with puromycin (Fig. 3C), indicating that the puromycin released a subset of the pep-tRNAs from the ribosome by replacing the tRNA moiety with puromycin.

Finally, the amount of Tris-adducted peptides was significantly increased in the pep-tRNA-derived peptides, as compared to that in the total proteins (Fig. S4), confirming that the tRNAs had been tethered to the C-terminus of the peptide moieties before the heat and high pH treatments. Taken together, PETEOS allowed us to capture a global snapshot of nascent polypeptide chains during translation.

Puromycin-resistant pep-tRNAs accumulate in the cells.

We next applied the PETEOS method to explore novel pausing peptide candidates, since in principle, the method can enrich potential SecM-like pausing peptides after the puromycin treatment, which is impossible by puromycin-labeling methods such as PUNCH-P.

Using an *E. coli* strain lacking the AcrAB-TolC multidrug efflux pump, to prevent the efflux of puromycin (40), we identified 400 proteins derived from the pep-tRNAs after the puromycin treatment (Fig. 3C), whereas the number of identified proteins was significantly

Figure 3. Indication that the peptides derived from pep-tRNAs are intermediates during translation elongation.

A. Illustration of pep-tRNAs and pre-existing proteins digested by trypsin. Trypsin digestion of pep-tRNAs, which elongate from the N- to C-terminus, should accumulate more peptides containing the N-terminal portion. In contrast, for pre-existing proteins, which are basically full-length polypeptides, the amounts of trypsin-digested peptides should be equal for all peptides. Red arrowheads indicate the trypsin cleavage sites. The relative positions of the N-terminus and C-terminus are defined as 0 and 1, respectively. **B**. Meta-analysis of peptides identified by LC-MS/MS for pep-tRNAs and total proteins. **C**. Venn diagram representing the identified pep-tRNAs obtained from cells, with or without puromycin treatment.

reduced to ~18 when RNase was added to the mixture after the phenol/chloroform extraction

(Step 1) (Fig. S5). About 70% of the 400 puromycin-resistant pep-tRNAs (287 out of 400)

overlapped with those derived from the puromycin-untreated sample (Fig. 3C), implying that

the puromycin-resistant pausing events might be prevalent.

The list of detected peptides included two known pausing peptides, SecM

($_{150}$FSTPVWISQAQGIR$_{163}$) and TnaC ($_{19}$IVDHRP$_{24}$), only under the puromycin treatment

even though the *E. coli* strain used in this experiment did not overexpress these ORFs (Table

S2). In addition, a portion of the TnaC-derived pausing peptide was detected as a Tris-adduct

($_{19}$IVDHRP$_{24}$-Tris), a hallmark of the peptidyl-tRNA. Taken together, the puromycin treatment

could be effective in identifying puromycin-resistant pausing peptides.

Indeed, several peptides that accumulated during translation could be candidates for

previously unknown pausing peptides (Table S2). One is the *raiA*-derived peptide,

$_{44}$EPQGFVADATINTPNGVL$_{61}$, along with its Tris-adduct ($_{44}$EPQGFVADATINTPNGVL$_{61}$-

Tris) (Fig. 4, Table S3). In fact, the C-terminal 61st leucine is consistent with a peak position in

the ribosome profiling data (Fig. 4), suggesting that the RaiA peptide could be derived from

translation pausing.

We also found another candidate, IvbL-LLPTAPSAAVVVVR, which seems to be

consistent with the ribosome profiling data (Table S2, Fig. S6). The translation of consecutive

valine residues near the C-terminus of IvbL induces pausing, due to the depletion of the valyl-

tRNA (41). Since the pausing due to such tRNA depletion would be abolished by the puromycin

addition, the accumulation of puromycin-resistant IvbL-tRNA implies an alternative pausing

mechanism, such as an association of the nascent peptide with the exit tunnel. In addition to

these candidates, we also detected other pausing candidates for the translation products, such

as *rof* and *udp* (Table S2).

Discussion

We developed a method for the direct isolation and analysis of pep-tRNAs in cells. Pep-

tRNAs are "in-between" molecules with both RNA and protein properties. The method

developed here exploited the "in-between" property to enrich pep-tRNAs from cellular RNAs

(mRNAs, tRNA, and ribosomal RNAs) and proteins that have already completed translation.

The organic solvent extraction and silica column purification with high concentrations of

guanidium salt were very effective for isolating pep-tRNAs. Considering several results,

including the RNase treatment, the meta-analysis showing N-terminally-biased peptides

derived from pep-tRNAs, and the Tris-conjugated C-terminal ends, we conclude that the

method, termed PETEOS, indeed enriches and identifies cellular pep-tRNAs.

Figure 4. Example of an ORF that includes pep-tRNA-derived peptides from puromycin-treated cells.

Representation of the *raiA* ORF. Blue regions schematically indicate the MS/MS-identified peptides, including a peptide adducted with Tris base. The peptides at the bottom represent the frequencies of ribosome footprints around the *raiA* ORF, obtained from a ribosome profiling database (GWIPS-viz browser). The amino acid sequences of the peptides with C-termini that matched with a prominent peak in the ribosome footprints are shown. Arrowheads indicate the trypsin cleavage sites.

A unique feature of the PETEOS method is the direct handling and analysis of pep-tRNAs,

providing a complementary approach to other established translatome analyses. Additionally,

the PETEOS method has several technical advantages. The time required to halt the cellular

processes before the pep-tRNA enrichment is very rapid, since an organic reagent is simply

added to the cells. Such quick halting enables the method to capture snapshots of the translation

status in the cell. In addition, the PETEOS procedure is relatively inexpensive and

straightforward as compared to ribosome profiling.

We assumed that the pep-tRNAs enriched by the PETEOS method are the genuine

intermediates during translation elongation. However, in addition to pep-tRNAs accommodated

within the ribosome as translation intermediates, some pep-tRNAs might be the products that

dropped off from the ribosome for some reasons. Although peptidyl-tRNA hydrolase cleaves

the ester bond of pep-tRNAs released from the ribosome, some pep-tRNAs might remain. The

current protocol cannot discriminate the "drop-off" pep-tRNAs from the pep-tRNAs-

accommodating ribosomes (42, 43). The isolation of the translating ribosomes by sucrose

density gradient centrifugation could enrich the pep-tRNAs committed to further elongation.

The PETEOS method has several limitations and points to improve. First, the inherently

low abundance of pep-tRNAs in the cell makes it difficult to analyze them using LC-MS/MS.

Second, the acidic phenol/chloroform extraction, which concentrates proteins and pep-tRNAs

while removing RNAs, is not sufficient. Some of the pep-tRNAs move into the aqueous phase,

probably due to their hydrophilic nature. For example, the distribution of TnaC-tRNA in the

organic solvent extraction differs between our results (Fig. S2C) and those of an earlier study,

where TnaC-tRNA was partitioned in the phenol phase (44). Although the protocols are

different, such as when the organic solvent is added, subtle differences in conditions and methods may affect the distribution significantly, leading to the incomplete separation of pep-tRNAs. Such incomplete separation would lead to unidentified peptidyl-tRNAs, causing some biases in the population and a lack of quantitative assessment. Finally, identifying pep-tRNA-derived peptides, including the C-terminal end of the growing polypeptide, is inherently difficult. The possible reasons for this are as follows. (i) Unless the translation is paused, the C-terminus of pep-tRNA moves continuously, making it difficult to accumulate detectable amounts of the same peptides for MS/MS. (ii) The peptide fragments closer to the N-terminus are, in principle, more abundant (Fig. 3). These drawbacks are inherent but could be overcome by improving the enrichment efficiency of C-terminal peptides, which are generally considered problematic even in conventional LC-MS/MS-based proteomics.

Despite its incompleteness, PETEOS has potential in that it is complementary to established translatome methods and enables analyses that cannot be achieved by other translatome methods. In addition, although *E. coli* lysates were used as a proof-of-concept experiment, the pep-tRNA analysis by PETEOS could be applicable to any organism.

Materials and Methods
E. coli strains and plasmids.

The *E. coli* DH5α strain was used for cloning, and the *E. coli* BW25113 {genotype: *Δ(araD-araB)567*, *ΔlacZ4787*(::rrnB-3), λ⁻, *rph-1*, *Δ(rhaD-rhaB)568*, *hsdR514*} was used as the standard strain. Phage P1-mediated transduction was used to introduce the Δ*tolC* mutation from Keio collection JW5503 (45). The plasmids used in this study are the pCA24N vector, pCA24N-*secM* (from the ASKA library) (46) and pYA004(pCA24N-*tnaC*). To construct pYA004, a DNA fragment amplified by PCR from a template *E. coli* using primers AY0003 and AY0004 and a DNA fragment amplified from pCA24N using primers PM0066 and PT0123 were assembled by Gibson assembly (47). The primers used in this study are as follows.
AY0003(AAAGAGGAGAAATTAACTATGAATATCTTACATATATGTGTG),
AY0004(TTAGCGGCCGCATAGGCCGATCCCTGTGAATATTACATC),
PM0066(GGCCTATGCGGCCGCTAAGGGTCGACCTGC),
PT0123(CATAGTTAATTTCTCCTCTTTAATGAATTC).

Bacterial growth.

E. coli cells cultured in LB medium overnight were inoculated in fresh LB medium, and grown at 37°C with shaking at 180 rpm, unless otherwise noted. Puromycin was added to a

final concentration of 10 µg/mL and the cells were incubated for 10 min at 37°C where indicated.

E. coli cells harboring pCA24N or its derivatives were grown overnight in LB medium containing 20 µg/ml of chloramphenicol, and were then inoculated into fresh LB medium containing chloramphenicol. They were grown at 37 °C with shaking at 180 rpm until the OD_{660} reached ~0.5. The expression of SecM or TnaC was then induced by adding 100 μM isopropyl β-D-1-thiogalactopyranoside (IPTG) for 30 min.

SDS-PAGE under neutral pH conditions and silver staining.

Samples were diluted in 2x SDS sample buffer (final 62.5 mM Tris-HCl, pH 6.8, 2% SDS, 10% glycerol, 5% β-mercaptoethanol, a trace of bromophenol blue), as described by Laemmli. SDS sample buffer was treated with RNA Secure (Ambion) according to the manufacturer's specifications. The samples were separated by the 11% WIDE Range SDS-PAGE system (Nacalai Tesque), as described previously (20, 48).

Silver staining of the gels was performed using a PlusOne Silver Staining Kit, Protein (GE Healthcare), according to the manufacturer's specifications in protocol "E," except that glutardialdehyde (25% w/v) was omitted from the *sensitizing solution* in the kit.

Phenol extraction and TCA/acetone precipitation to fractionate polypeptides and polypeptidyl-tRNAs from nucleic acids.

E. coli cells grown in 200 ml of LB medium until the mid-log phase were mixed with an approximately 200 mL volume of crushed ice, and collected by centrifugation at 4 °C for 10 min. After supernatant removal, the cell pellet was suspended in 20 mL of TriPURE isolation reagent (Roche) and incubated for 5 min at room temperature. The lysate was then mixed with 4 ml of chloroform, vortexed for 20 sec, incubated for 5 min at room temperature, and centrifuged at 12,000 ×g for 5 min at 4 °C. The colorless to pale yellow transparent upper layer (aqueous phase, 10-12 mL) was removed, leaving an organic phenol phase and a solid middle layer. After the addition of an equal volume (12-15 mL from a 200 mL culture) of ice-cooled 20% TCA/acetone containing 5 mM dithiothreitol (DTT), the mixture was vortexed for 20 sec and incubated for at least 3 hrs at -20 °C. The sample was centrifuged at 15,600 ×g for 5 min at 4 °C, and the supernatant was carefully discarded. The precipitate was vortexed with 10 mL of acetone and centrifuged at 15,600 ×g for 5 min at 4 °C, and then the supernatant was discarded by aspiration. This washing procedure was repeated twice.

Peptidyl-tRNA purification using a silica column.

The precipitate obtained from the organic phenol phase was resolved in a 1 ~ 3 ml volume

of phase-transfer surfactants (PTS) buffer {100 mM Tris-HCl, pH 6.8, 24 mM sodium deoxycholate, 24 mM sodium N-lauroyl sarcosinate (39)} by vortexing at 37 °C for 15 min. After centrifugation, the collected supernatant was divided into two equal portions, and one was treated with 40 µg/ml of RNase A (Promega) at 37 °C for 1 hr.

As the next step, the peptidyl-tRNAs in the PTS buffer were isolated by using a High Pure miRNA Isolation Kit (Roche) according to the manufacturer's instructions, except that 1/2 volume of the *binding enhancer* was used (48). About 750 µL of PTS lysate could be processed by one silica column in the kit. The silica-bound RNA was eluted with 100 µl of PTS buffer, and this purification was repeated twice.

LC-MS/MS sample preparation.

The purified RNA sample in the previous section (100 µl) was mixed with 10 µl of 2 M Tris base (pH= ~11) and incubated at 80 °C for 20 min to hydrolyze the ester bond of the peptidyl-tRNA. After the addition of 90 µl deionized water, the sample was reduced by the addition of 10 mM DTT at room temperature for 30 min, and then alkylated with 50 mM iodoacetamide in the dark at room temperature for 20 min. The mixture was diluted 5-fold with 50 mM ammonium bicarbonate. For the limited digestion of denatured polypeptides into peptide fragments, 0.5 µg of Lys-C (Fujifilm Wako) was added, and the solution was incubated at room temperature for at least 3 hrs. Subsequently, 0.5 µg of Trypsin Gold (Promega) was added and incubated at 37 °C overnight, whereas for the SecM analysis, 0.5 µg of Lys-C was used. After the digestion, an equal volume of ethyl acetate and 0.5 % trifluoroacetic acid (final concentration) was added. The mixture was shaken vigorously for 2 min and centrifuged at 15,700 ×g for 2 min. The upper ethyl acetate layer was removed, and the solvent was removed by a centrifugal evaporator. The residual pellet was dissolved in 0.1 % TFA and 2 % acetonitrile and desalted, as follows. The solution was applied to an SDB-XC StageTip made by an Empore disk (3M, U.S.A.), equilibrated with 0.1 % TFA and 2 % acetonitrile, washed with 0.1 % TFA and 2 % acetonitrile, and eluted with 0.1 % TFA and 80 % acetonitrile. The solvent was removed by a centrifugal evaporator, and the residue was dissolved in 0.1 % TFA and 2 % acetonitrile. This solution was centrifuged at 20,000 ×g for 5 min, and the supernatant was collected and subjected to the LC-MS/MS measurement.

LC-MS/MS measurement.

The LC-MS/MS measurements were conducted with a Q-Exactive tandem mass spectrometer and an Easy-nLC 1000 nano HPLC system (Thermo Fischer Scientific). The trap column used for the nano HPLC was a 2 cm × 75 µm capillary column packed with 3 µm C18-silica particles

(Thermo Fisher Scientific), and the separation column was a 12.5 cm × 75 μm capillary column
packed with 3 μm C18-silica particles (Nikkyo Technos, Japan). The flow rate for the nano
HPLC was 300 nL/min. The separation was conducted using a $10-40\%$ linear acetonitrile
gradient over 70 min, in the presence of 0.1% formic acid. The nanoLC-MS/MS data were
acquired in the data-dependent acquisition (DDA) mode controlled by the Xcalibur 4.0 program
(Thermo Fisher Scientific). The DDA settings were as follows: the resolution was 70,000 for a
full MS scan and 17,500 for an MS2 scan; the AGC target was 3.0E6 for a full MS scan and
5.0E5 for an MS2 scan; the maximum IT was 60 ms for both the full MS and MS2 scans; the
full MS scan range was $310-1,500$ m/z; and the top 10 signals in each full MS scan were
selected for the MS2 scan. The DDA measurement was performed three times for each sample,
and two or three biological replicates were measured in each condition.
374

Data analysis.
The data analysis of the DDA measurements was performed with the Proteome Discoverer 2.4
program and the SEQUEST HT search engine (Thermo Fischer Scientific). The dataset of the
amino acid sequences of *E. coli* genes was obtained from the UniProt database (UP000000625,
downloaded on 181219). Only the peptides with a < 1 % false discovery rate (by the Percolator
algorithm) detected more than twice in the three measurements were selected. All processes
and statistical analyses were conducted by the R software (version 4.4.1 with in-house scripts).
The ribosome distributions on the mRNAs in the ribosome profiling were obtained using the
aggregation mode in the GWIPS-viz browser, a public web tool to visualize the ribosome
distribution (49, 50).
385

386 **Data Availability Statement.**
387 Data in this manuscript have been uploaded to the Mendeley Dataset public repository
388 (https://doi.org/10.17632/xzycy8m6tb.1).
389

390 **Author contributions.**
391 AY and YC performed experiments; AY and TN analyzed the LC-MS/MS data; AY, YC, TN,
392 and HT conceived the study, designed experiments, and analyzed the results; YC, TN, and HT
393 supervised the entire project; AY, YC, TN, and HT wrote the manuscript.
394

395 **Acknowledgments.**
396 We thank Eri Uemura for technical support; the Bio-support Center at Tokyo Tech for DNA
397 sequencing; and the Cell Biology Center Research Core Facility at Tokyo Tech for the Q-

Exactive mass spectrometry measurements. This work was supported by MEXT Grants-in-Aid for Scientific Research (Grant Numbers 17K15073 to TN, 17K15062, 19K16038 to YC, JP26116002, JP18H03984, JP20H05925 to HT) and a grant from the Ohsumi Frontier Science Foundation to YC.

Conflicts of Interest.

The authors declare no competing interest.

References.

1. C. Vogel, E. M. Marcotte, Insights into the regulation of protein abundance from proteomic and transcriptomic analyses. *Nature Reviews Genetics* **13**, 227–232 (2012).

2. S. Pechmann, F. Willmund, J. Frydman, The Ribosome as a Hub for Protein Quality Control. *Molecular Cell* **49**, 411–421 (2013).

3. K. Ito, S. Chiba, Arrest peptides: Cis-acting modulators of translation. *Annual Review of Biochemistry* **82**, 171–202 (2013).

4. K. C. Stein, J. Frydman, The stop-and-go traffic regulating protein biogenesis: How translation kinetics controls proteostasis. *Journal of Biological Chemistry* **294**, 2076–2084 (2019).

5. E. Samatova, J. Daberger, M. Liutkute, M. v Rodnina, Translational Control by Ribosome Pausing in Bacteria: How a Non-uniform Pace of Translation Affects Protein Production and Folding. *Frontiers in Microbiology* **11**, 619430 (2021).

6. D. N. Wilson, S. Arenz, R. Beckmann, Translation regulation via nascent polypeptide-mediated ribosome stalling. *Current Opinion in Structural Biology* **37**, 123–133 (2016).

7. J. Choi, *et al.*, How Messenger RNA and Nascent Chain Sequences Regulate Translation Elongation. *Annu Rev Biochem* **87**, 421–449 (2018).

8. J. L. Chaney, P. L. Clark, Roles for Synonymous Codon Usage in Protein Biogenesis. *Annual Review of Biophysics* **44**, 143–166 (2015).

9. T. Sunohara, T. Abo, T. Inada, H. Aiba, The C-terminal amino acid sequence of nascent peptide is a major determinant of SsrA tagging at all three stop codons. *RNA* **8**, 1416–1427 (2002).

10. M. A. Sørensen, C. G. Kurland, S. Pedersen, Codon usage determines translation rate in Escherichia coli. *Journal of Molecular Biology* **207**, 365–377 (1989).

11. C. S. Hayes, B. Bose, R. T. Sauer, Proline residues at the C terminus of nascent chains induce SsrA tagging during translation termination. *Journal of Biological Chemistry* **277**, 33825–33832 (2002).

12. I. Rychlík, J. ČErná, S. Chládek, P. Pulkrábek, J. ŽEmlička, Substrate Specificity of
Ribosomal Peptidyl Transferase: The Effect of the Nature of the Amino Acid Side Chain
on the Acceptor Activity of 2'(3')-O-Aminoacyladenosines. *European Journal of*
Biochemistry **16**, 136–142 (1970).

13. H. Nakatogawa, K. Ito, Secretion monitor, secM, undergoes self-translation arrest in the
cytosol. *Molecular Cell* **7**, 185–192 (2001).

14. H. Nakatogawa, K. Ito, The ribosomal exit tunnel functions as a discriminating gate. *Cell*
108, 629–36 (2002).

15. J. Zhang, *et al.*, Mechanisms of ribosome stalling by SecM at multiple elongation steps.
Elife **4**, e09684 D (2015).

16. F. Gong, K. Ito, Y. Nakamura, C. Yanofsky, The mechanism of tryptophan induction of
tryptophanase operon expression: Tryptophan inhibits release factor-mediated
cleavage of TnaC-peptidyl-tRNAPro. *Proc Natl Acad Sci U S A* **98**, 8997–9001 (2001).

17. N. Vazquez-Laslop, C. Thum, A. S. Mankin, Molecular Mechanism of Drug-Dependent
Ribosome Stalling. *Molecular Cell* **30**, 190–202 (2008).

18. H. Onouchi, *et al.*, Nascent peptide-mediated translation elongation arrest coupled with
mRNA degradation in the CGS1 gene of Arabidopsis. *Genes and Development* **19**, 1799–
1810 (2005).

19. K. Yanagitani, Y. Kimata, H. Kadokura, K. Kohno, Translational pausing ensures
membrane targeting and cytoplasmic splicing of XBP1u mRNA. *Science* **331**, 586–589
(2011).

20. Y. Chadani, T. Niwa, S. Chiba, H. Taguchi, K. Ito, Integrated in vivo and in vitro nascent
chain profiling reveals widespread translational pausing. *Proc Natl Acad Sci U S A* **113**,
E829–E838 (2016).

21. H. Muto, H. Nakatogawa, K. Ito, Genetically Encoded but Nonpolypeptide Prolyl-tRNA
Functions in the A Site for SecM-Mediated Ribosomal Stall. *Molecular Cell* **22**, 545–552
(2006).

22. S. Iwasaki, N. T. Ingolia, The Growing Toolbox for Protein Synthesis Studies. *Trends*
Biochem Sci **42**, 612–624 (2017).

23. N. T. Ingolia, S. Ghaemmaghami, J. R. S. Newman, J. S. Weissman, Genome-wide analysis
in vivo of translation with nucleotide resolution using ribosome profiling. *Science (1979)*
324, 218–223 (2009).

24. N. T. Ingolia, G. A. Brar, S. Rouskin, A. M. McGeachy, J. S. Weissman, The ribosome
profiling strategy for monitoring translation in vivo by deep sequencing of ribosome-
protected mRNA fragments. *Nature Protocols* **7**, 1534–1550 (2012).

25. G. A. Brar, J. S. Weissman, Ribosome profiling reveals the what, when, where and how
of protein synthesis. *Nature Reviews Molecular Cell Biology* **16**, 651–664 (2015).

26. N. T. Ingolia, J. A. Hussmann, J. S. Weissman, Ribosome profiling: Global views of
translation. *Cold Spring Harbor Perspectives in Biology* **11**, a032698 (2019).

27. A. R. Buskirk, R. Green, Ribosome pausing, arrest and rescue in bacteria and eukaryotes.
Philosophical Transactions of the Royal Society B: Biological Sciences **372**, 20160183
(2017).

28. F. Mohammad, C. J. Woolstenhulme, R. Green, A. R. Buskirk, Clarifying the Translational
Pausing Landscape in Bacteria by Ribosome Profiling. *Cell Reports* **14**, 686–694 (2016).

29. D. C. Dieterich, A. J. Link, J. Graumann, D. A. Tirrell, E. M. Schuman, Selective
identification of newly synthesized proteins in mammalian cells using bioorthogonal
noncanonical amino acid tagging (BONCAT). *Proc Natl Acad Sci U S A* **103**, 9482–9487
(2006).

30. A. J. M. Howden, *et al.*, QuaNCAT: Quantitating proteome dynamics in primary cells.
Nature Methods **10**, 343–346 (2013).

31. R. Aviner, T. Geiger, O. Elroy-Stein, Genome-wide identification and quantification of
protein synthesis in cultured cells and whole tissues by puromycin-associated nascent
chain proteomics (PUNCH-P). *Nature Protocols* **9**, 751–760 (2014).

32. R. Aviner, T. Geiger, O. Elroy-Stein, Novel proteomic approach (PUNCH-P) reveals cell
cycle-specific fluctuations in mRNA translation. *Genes and Development* **27**, 1834–1844
(2013).

33. C. M. Forester, *et al.*, Revealing nascent proteomics in signaling pathways and cell
differentiation. *Proc Natl Acad Sci U S A* **115**, 2353–2358 (2018).

34. M. Dermit, M. Dodel, F. K. Mardakheh, Methods for monitoring and measurement of
protein translation in time and space. *Molecular BioSystems* **13**, 2477–2488 (2017).

35. R. Milo, P. Jorgensen, U. Moran, G. Weber, M. Springer, BioNumbers The database of
key numbers in molecular and cell biology. *Nucleic Acids Research* **38**, D750-3 (2010).

36. K. Ito, *et al.*, Nascentome analysis uncovers futile protein synthesis in Escherichia coli.
PLoS ONE **6**, e28413 (2011).

37. R. Boom, *et al.*, Rapid and simple method for purification of nucleic acids. *Journal of
Clinical Microbiology* **28**, 495–503 (1990).

38. B. Vogelstein, D. Gillespie, Preparative and analytical purification of DNA from agarose.
Proc Natl Acad Sci U S A **76**, 615–619 (1979).

39. T. Masuda, M. Tomita, Y. Ishihama, Phase transfer surfactant-aided trypsin digestion for
membrane proteome analysis. *Journal of Proteome Research* **7**, 731–740 (2008).

40. H. Okusu, D. Ma, H. Nikaido, AcrAB efflux pump plays a major role in the antibiotic
resistance phenotype of Escherichia coli multiple-antibiotic-resistance (Mar) mutants. *J*
Bacteriol **178**, 306–308 (1996).

41. P. Friden, T. Newman, M. Freundlich, Nucleotide sequence of the ilvB promoter-
regulatory region: A biosynthetic operon controlled by attenuation and cyclic AMP. *Proc*
Natl Acad Sci U S A **79**, 6156–6160 (1982).

42. S. Vivanco-Domínguez, *et al.*, Protein Synthesis Factors (RF1, RF2, RF3, RRF, and tmRNA)
and Peptidyl-tRNA Hydrolase Rescue Stalled Ribosomes at Sense Codons. *Journal of*
Molecular Biology **417**, 425–439 (2012).

43. V. Heurgué-Hamard, *et al.*, Ribosome release factor RF4 and termination factor RF3 are
involved in dissociation of peptidyl-tRNA from the ribosome. *The EMBO Journal* **17**,
808–816 (1998).

44. M. Gong, F. Gong, C. Yanofsky, Overexpression of tnaC of Escherichia coli inhibits growth
by depleting tRNA2Pro availability. *Journal of Bacteriology* **188**, 1892–1898 (2006).

45. T. Baba, *et al.*, Construction of Escherichia coli K-12 in-frame, single-gene knockout
mutants: The Keio collection. *Molecular Systems Biology* **2**, 0008 (2006).

46. M. Kitagawa, *et al.*, Complete set of ORF clones of *Escherichia coli* ASKA library (A
complete set of E. coli K-12 ORF archive): unique resources for biological research. *DNA*
Research **12**, 291–299 (2005).

47. D. G. Gibson, Enzymatic assembly of overlapping DNA fragments. *Methods Enzymol* **498**,
349–361 (2011).

48. Y. Chadani, *et al.*, Intrinsic Ribosome Destabilization Underlies Translation and Provides
an Organism with a Strategy of Environmental Sensing. *Molecular Cell* **68**, 528-539.e5
(2017).

49. A. M. Michel, S. J. Kiniry, P. B. F. O'Connor, J. P. Mullan, P. v. Baranov, GWIPS-viz: 2018
update. *Nucleic Acids Research* **46**, D823–D830 (2018).

50. A. M. Michel, *et al.*, GWIPS-viz: Development of a ribo-seq genome browser. *Nucleic*
Acids Research **42**, D859–D864 (2014).