A revisit to universal single-copy genes in bacterial genomes

Saidi Wang[1, *], Minerva Ventolero[2, *], Haiyan Hu[1,3, ¶], Xiaoman Li[2, ¶]

[1]Department of Computer Science, University of Central Florida, Orlando, Florida, United States of America,

[2]Burnett School of Biomedical Science, College of Medicine, University of Central Florida, Orlando, Florida, United States of America.

[3]Genomics and Bioinformatics Cluster, University of Central Florida, Orlando, Florida, United States of America.

[*]These authors contributed equally to this work.

[¶]Corresponding Authors:

Email: xiaoman@mail.ucf.edu, haihu@cs.ucf.edu

Abstract

Universal single-copy genes (USCGs) are widely used for species classification and taxonomic profiling. Despite many studies on USCGs, our understanding of USCGs in bacterial genomes might be out of date, especially how different the USCGs are in different studies, how well a set of USCGs can distinguish two bacterial species, whether USCGs can separate different strains of a bacterial species, to name a few. To fill the void, we studied USCGs in the most updated bacterial RefSeq genomes. We showed that different USCG sets are quite different

while coming from highly similar functional categories. We also found that although USCGs

occur once in almost all bacterial genomes, each USCG does occur multiple times in certain

genomes. We demonstrated that USCGs are reliable markers to distinguish different species

while they cannot distinguish different strains of most bacterial species. Our study shed new light

on the usage and limitations of USCGs, which will facilitate their applications in evolutionary,

phylogenomic, and metagenomic studies.

Introduction

Universal single-copy genes (USCGs) are marker genes that occur once and only once in

almost every genome [1]. Because of this property of ubiquitous existence and uniqueness in

each genome, USCGs are widely used to study the evolution and classification of species [2-4].

In the past two decades, with the enormous amount of metagenomic data generated, USCGs are

also routinely employed for taxonomic profiling of microbial species and completeness

evaluation of metagenome-assembled genomes in shotgun metagenomic studies [5-9].

A widely used marker gene is the 16S rRNA gene, which is thought to be single-copied

while shown otherwise and is thus not a perfect USCG [10]. 16S rRNA gene has been used as

the gold standard in amplicon sequencing and demonstrated its power for taxonomic profiling in

various metagenomic studies [4, 11-14]. It is the most sequenced gene, with its sequences often

the only ones we know about a species. Despite its indisputable value and popularity, the 16S

rRNA gene may have multiple copies in an unknown genome [10]. Moreover, its sequences

might be too conserved to distinguish certain species and/or correctly measure the species

divergence [4]. It is thus natural to consider other USCGs in evolution and metagenomics.

Previous studies have identified USCGs other than the 16S rRNA gene [1-4, 9, 15-19]. For instance, Ciccarelli et al. obtained 31 USCGs to reconstruct the tree of life across all three domains [1]. Later, Creevey et al. inferred 40 USCGs from the same set of 191 complete genomes [16], which were then used for taxonomic profiling in metagenomics by the tools mOTUs and mOTUs2 [5, 6]. Wu and Eisen developed the AMPHORA pipeline for automatic phylogenomic analysis of microbial species based on 31 USCGs [4]. These 31 USCGs were combined with 104 archaeal USCGs for phylogenomic studies [19]. Later, the same group developed the phyEco gene sets that comprised 40 USCGs for "all bacteria and archaea" and 114 USCGs for "all bacteria" [3]. Alneberg et al. analyzed all clusters of orthologous groups in 525 genomes and identified 36 USCGs that occurred in >97% of the 525 genomes with a frequency <1.03 per genome [15]. Lan et al. discovered 73 USCGs in >90% of 1897 genomes to classify prokaryotic species [2]. Parks et al. extracted 120 USCGs present in >90% of genomes and single-copied in >95% of their present genomes [9]. These sets of USCGs are inferred from different groups of sequenced genomes with different purposes. They are thus different, although they do share a fraction of USCGs.

With multiple USCGs, in addition to studying evolution, classifying and taxonomic profiling of species, several studies attempted to investigate the strain diversity [7, 20-23]. For instance, Quince et al. developed tools to de novo reconstruct bacterial strain genomes from shotgun metagenomic reads with the aforementioned 36 USCGs [7, 23]. The StrainPhlAn tool infers bacterial strains directly from shotgun reads based on ~200 clade-specific marker genes [22]. Nayfach et al. proposed a pipeline for strain profiling in metagenomic datasets with 15 USCGs [24]. Such studies of bacterial strains are important for understanding drug resistance, microbial diversity, and the cure of various complex diseases [25-30].

Despite many studies on USCGs, our understanding of USCGs is somewhat outdated. For instance, how much agreement is there between the USCG sets from different studies? For a set of USCGs, how universal are these USCGs in the ever-increasing number of sequenced genomes? When defining the above sets of USCGs, to our knowledge, only the first two sets of USCGs were selected by requiring their single-copy occurrence in all 191 genomes. None of the USCG sets is tested on the latest set of RefSeq genomes to show their universalism. Moreover, it is also not clear whether certain USCGs can tell two species apart better than others. In addition, it is unknown whether a set of USCGs such as the 40 USCGs from Creevey et al. [16] contain enough variations to distinguish different strains of a bacterial species.

To address these questions, especially the last one, in this study, we compared different sets of USCGs. We systematically studied how similar a USCG is in different species and strains with the latest set of complete RefSeq genomes. We found that almost every USCG occurs multiple times in certain genomes, while more than 99.4% of USCGs are single-copied in a given genome. We also observed that USCGs together are good for separating species, while they cannot distinguish strains from the same bacterial species in general. Our study provides a more updated picture of USCGs and their potential applications in evolutionary and metagenomic studies.

Results

USCG sets are different in gene content while similar in gene function

We compared USCG sets from seven studies [1-4, 9, 15, 16] (Material and Methods,

Supplementary Table S1). They were (1). 31 USCGs from Ciccarelli et al.; (2). 31 USCGs from

Wu and Eisen; (3). 40 USCGs from Creevey et al.; (4). 40 USCGs from Wu et al. for "all"

bacteria and archaea; (5). 36 USCGs from Alneberg et al.; (6). 73 USCGs from Lan et al.; and

(7). 120 USCGs from Parks et al. These USCG sets require different universalism across all

genomes and uniqueness in individual genomes. For instance, USCGs in the last two sets

occurred in only >90% of the genomes, while that was at least >97% in the third to the fifth sets.

We found that even the USCG sets from the same research group could be very different

(Figure 1A). For instance, only 18 (77%) of the 31 USCGs from Wu and Eisen were shared with

the 40 USCGs from Wu et al. Both sets were inferred by the same research group, with the latter

inferred from a much larger number of genomes [3, 4]. Their difference thus implied the

significant effect of the genomes used to infer these USCGs On the other hand, the USCG sets

from the same research group could be highly consistent as well. For instance, the USCG set

from Creevey et al. was a superset of the USCG set from Ciccarelli et al., as the same research

group inferred them with the same genomes and a slightly different strategy. However, the

additional nine USCGs from Creevey et al. also demonstrated the binding effect of such different

strategies on defining USCGs.

Figure1. USCG sets and functional categories. (A) Overlap of the seven USCG sets. (B) The

percentage of the functional categories of the USCGs in the seven sets.

The USCG sets from different research groups differed more (Supplementary Table S2).

For instance, only 14 (45%) of the 31 USCGs from Ciccarelli et al. were shared with those from

Wu and Eisen. Again, the difference in these two sets corroborated the effect of different

strategies and different genomes. When we considered the two sets refined later by the

corresponding research groups, the third and fourth sets, 28 (70%) of the 40 USCGs in these two

sets were shared. Although the USCGs in these two sets were still drastically different, they

shared much more USCGs than their earlier versions, likely because the number of genomes was

large enough to choose a more representative set of genomes by Wu et al. Compared with the

sets from Alneberg et al., Lan et al., and Parks et al., the USCGs from Creevey et al. was at least

comparable with those from Wu et al., if not better. Overall, the USCG set from Creevey et al. is

likely as reliable as any other set, if not more reliable, because of its stricter requirement of

universalism and uniqueness, the similar evolutionary trajectory of these USCGs, and likely

more representative genomes (consistent results with other sets that used much more genomes).

Despite the difference in the number and members of USCGs, the function of the USCGs

in each of the above sets was quite consistent (Figure 1B) [31-35]. By checking the functional

annotations at https://www.ncbi.nlm.nih.gov/research/cog, we found that the vast majority of

USCGs in every set are annotated with the functional category J (Translation, ribosomal

structure, and biogenesis). The remaining USCGs were annotated with other translation and

metabolism related categories, such as L (Replication, recombination, and repair), H (Coenzyme

transport and metabolism), F (Nucleotide transport and metabolism), O (Posttranslational

modification, protein turnover, chaperones), etc. Each category other than J was annotated with

much fewer USCGs, from one to a handful of USCGs, compared with several dozen for J.

Overall, the USCG sets are enriched with functions related to translation and metabolism, no

matter which USCG set is concerned about. For instance, among the 40 USCGs from Wu et al.,

35 are involved in the translation process, while the remaining five are related to the cellular

metabolic process [3].

USCGs are indeed universal across species and unique in individual genomes

To our knowledge, the number of genomes used to infer the above USCG sets was no more than 2000 except those from Parks et al. With more than 25,000 complete bacterial genomes at National Center for Biotechnology Information (NCBI), it was unclear whether the USCGs were still universal and unique. We thus studied the occurrence of the USCG sets in the latest set of complete RefSeq genomes (Material and Methods). We focused on the sets from Creevey et al., Wu et al., and Alneberg et al. below because they are widely used in metagenomics and more updated than their previous versions (the first and third set). Moreover, they have more stringent criteria for universalism and uniqueness than the sets from Lan et al. and Parks et al. [2, 9]. In the following, we presented our study on the above three sets.

We found that the 40 USCGs from Creevey et al. were universally distributed in almost each of the 25271 RefSeq genomes (Table 1, Supplementary Table S3, Material and Methods). For every USCG, it occurred in at least 97.7% of the 25271 complete RefSeq genomes. For genomes without a copy of a USCG, we applied BLAST with an arbitrary cutoff of 1E-15 to search for this USCG. BLAST could identify a copy with this cutoff in most of these missed genomes. A tiny fraction of the genomes still did not have a USCG copy, likely due to the cutoff, the quality of the genomes, and the imperfectness of the gene retrieval methods used. Despite these limitations, each USCG occurred in at least 99.7% of the RefSeq genomes. Of note, the number of USCGs occurring in each genome also showed the universal distribution of these USCGs. The mean number of USCGs occurring in each RefSeq genome was 39.6. In other

words, almost each of the 40 USCGs occurred in every genome, indicating the universalism of

the USCGs from Creevey et al.

We also observed that each of the 40 USCGS from Creevey et al. indeed had only one

copy in almost every genome (Table 1). For the 25271 RefSeq genomes, the mean and median

percentage of the genomes with only one copy of a USCG were 98.5% and 99.0%, respectively.

For individual USCGs, 24 (60%) of the 40 USCGs occurred more than once in <0.2% of the

RefSeq genomes, while several other USCGs occurred more than once in >2.0% of the RefSeq

genomes. Interestingly, these USCGs that occurred more than once in >2% of the RefSeq

genomes were often not identified in other USCG sets. For instance, for the top ten USCGs

occurring more than once, only one of them was also identified in the set from Wu et al.,

suggesting that these USCGs might have occurred multiple times in other studies and were thus

filtered by those studies. Overall, the average number of USCGs with only one copy in a genome

was 39.4. This average number for the RefSeq genomes suggests that almost all USCGs have

only one copy in almost all complete bacterial genomes.

We further studied the universalism and uniqueness of the USCG sets from Wu et al. and

Alneberg et al. (Supplementary Table S3). Compared with the set from Creevey et al., the other

two sets had a similar median (Figure 2). For instance, the median of the universalism was

99.3% for all three sets, and the corresponding median of the uniqueness was 99.0%, 99.1%, and

99.1%. However, there were more individual USCGs in the other two sets that were not so

universal and unique as those from Creevey et al. It is thus evident that the above 40 USCGs

from Creevey et al. might be a better USCG set.

Figure 2. The universalism and uniqueness of the three USCG sets. Comparison of universalism and uniqueness in complete RefSeq genomes.

USCGs well separate different species, but not strains of most species

With the USCGs indeed universal and unique in bacterial genomes, we investigated how well they together could distinguish two bacterial species in the same genus and whether they together are enough to set two bacterial strains of the same bacterial species apart. To address these questions, we studied all 440 genera with at least two sequenced species genomes and all 747 bacterial species with at least two sequenced strain genomes in the 25271 complete RefSeq genomes (Material and Methods). We found that at least the 40 USCGs from Creevey et al. together were enough to distinguish almost all pairs of species in the same genus. However, they could not separate pairs of strains from most species (Table 2).

With the Creevey et al. USCG set, we studied the similarity of 40 pairs of corresponding USCGs in every two species from the same genus for all 440 genera with at least two complete genomes (Supplementary Table S4). We measured the similarity by the percentage of identity (PID) in the alignment of the 40 pairs of USCGs from a pair of species because the PID determines how similar the shotgun metagenomics reads from these USCGs are. The PID had a mean and median of 90.2% and 90.6%. However, about 2.3% of species pairs had a PID larger than 99.5%, which was the limit that the current computational tools could distinguish two genomes [7, 36-40]. Given that the total nucleotide length of the 40 USCGs is about 33,852 base pairs, the 99.5% PID means there are at least 170 varied positions in the 40 USCGs required to

separate the two species. Therefore, the 40 USCGs could tell most pairs of species apart, which

may leave about 2.3% of the pairs of species indistinguishable.

With the same USCG set, we also studied the similarity of 40 pairs of corresponding

USCGs in every two strains from the same species for all 747 species with at least two complete

genomes (Table 2, Supplementary Table S5). The PID in a pair of strains had a mean of 99.8%

and a median of 99.9%. About 91.2% of the PID scores between pairs of strains from the same

species were already larger than 99.5% (Table 2). In other words, the 40 USCGs alone may have

difficulty in separating more than 91.2% of pairs of bacterial strains.

We also studied how well the USCG set from Wu et al. and from Alneberg et al.

separated species and strains (Table 2). Compared with the USCG set from Creevey et al., these

two sets had a slightly lower performance, especially in terms of separating strains of the same

species. Many more pairs of species and strains had close to 100% PID based on the Wu et al. set

compared with the other two sets. Overall, the USCG set from Creevey et al. is likely to perform

better in distinguishing species and strains.

Discussion

We systematically studied the occurrence of USCGs in the most updated list of

sequenced bacterial genomes. We showed that USCGs were universally distributed and uniquely

present in almost every bacterial species. The 40 USCGs from Creevey et al. together could

distinguish different species from the same genus and different strains from the same species

well. Barely was there one individual USCG that could do so alone.

There are many other sets of USCGs not compared here. Wrighton et al. used 30 USCGs

to measure the completeness of the reconstructed 49 bacterial genomes [18]. Haroon et al.

employed the Amphora2 marker genes, including the 31 USCGs for bacteria from Wu and Eisen

and 104 archaeal genes, to measure the completeness of the assembled genomes [17]. Rinke et

al. estimated the completeness of the assembled genomes based on 139□bacterial and

162□archaeal conserved marker genes [41]. Wu et al. reported 114 USCGs for "all" bacteria in

addition to the 40 USCGs compared here [3]. We did not compare these sets of USCGs because

they were more for other purposes and not as universal and unique as those compared in this

study.

We found that at least three of the seven sets of USCGs were universal and unique. Other

sets of USCGs, except the one from Ciccarelli et al., were unlikely to be as universal and unique

as the three sets because their selection was not as strict. For instance, Lan et al. and Parks et al.

required USCGs to occur in only more than 90% of the genomes instead of all genomes [2, 9].

The Ciccarelli et al. set was already included in the set from Creevey et al.

For the complete RefSeq genomes, we demonstrated that USCGs could distinguish

different species but not strains. This was based on the fact that the PID of 99.5% is the limit of

current tools to separate similar sequences. When the strain similarity is higher, say more than

99.5%, we are left with only a few dozen variable loci to distinguish different strains. In this

case, with the available USCG sets, it may be challenging to distinguish them, even if possible.

Note that all aforementioned USCG sets are not created to distinguish bacterial species and

strains in microbiomes. In the future, one may hope to generate new USCG sets for this purpose

by developing novel methods and integrating different sources of information [42-48].

Alternatively, one may consider overlapping the assembled contigs or metagenome-assembled-

genomes with the USCGs so that more polymorphic sites are available to distinguish strains [7,

23].

Material and Methods

Seven sets of USCGS

246   We studied seven sets of USCGs: the 31 set from Ciccarelli et al. [1], the 31 set from Wu

247 and Eisen [4], the 40 set from Creevey et al. [16], the 40 USCGs from Wu et al. [3], the 36

248 USCGs from Alneberg et al. [15], the 73 USCGs from Lan et al. [2], and the 120 USCGs from

249 Parks et al. [9]. We focused on these sets for the separation of similar strains, not for the

250 completeness of the assembled genomes in metagenomics.

The complete RefSeq genomes

252   We studied the complete bacterial Refseq genomes at NCBI. We retrieved the RefSeq

253 genomes from the NCBI website (https://www.ncbi.nlm.nih.gov/genome/browse#!/prokaryotes/)

254 by selecting the 'Bacteria' Kingdom and 'Complete' Assembly level. There were 25271

255 genomes with downloadable reference sequences.

Identification of USCGs in a genome

To identify the corresponding sequences of a USCG in a given genome, we used the tool

fetchMG [6]. This tool was used previously to identify and quantify the microbial species in

assembled contigs from shotgun metagenomic reads. We used the following command to fetch

the USCG sequence in a genome: ./fetchMG.pl -m extraction -p proteins.fasta -o output. Because

fetchMG might be imperfect and might miss certain USCG sequences in a genome, we also

applied BLAST with the following command to retrieve USCG sequences that were not

identified in a genome: psiblast -in_msa marker_aligned.fasta -db db -evalue 1E-15 -

num_alignments num_keep. The E-value cutoff used was 1E-15 as previously [3, 4, 19]. We

considered the fetched sequences were the corresponding USCG sequences in the genomes. Note

that neither fetchMG nor BLAST is perfect for identifying copies of USCGs in a genome, as

they sacrifice the accuracy for the sake of speed. However, the identified copies of USCGs are

likely to be reliable. We obtained the functional categories of each USCG from

https://www.ncbi.nlm.nih.gov/research/cog.

The USCG similarity measurement

We aligned the extracted USCG protein sequences for every USCG with the tool MAFFT

[49]. In other words, we did multiple alignments of the USCG sequences extracted from each

genome under consideration for each USCG. In order to assess the similarity of the USCG

sequences, we calculated the score based on the blosum62 matrix, the matched number, the

mismatch number, the indel number, and the PID. We chose the PID to evaluate the similarity of

each pair of USCG sequences because it directly relates to the shotgun metagenomic reads

mapped to different USCG sequences. To measure the PID, we obtained the two corresponding

sequences in the alignment and removed the loci with an indel versus an indel. We then

calculated the PID as the ratio of the number of matched positions to the number of all remaining

aligned positions.

Acknowledgment

282   This work was supported by the National Science Foundation (1661414, 2015838, 2120907).

Author Contributions

H.H. and X.L. conceived the idea. S.W. and M.V. implemented the idea and generated results.

S.W., M.V., X.L. and H.H. analyzed the results and wrote the manuscript. All authors reviewed

the manuscript.

Conflict of interest

We declare that there is no conflict of interest regarding the publication of this article.

Data availability

The NCBI genomes are from https://www.ncbi.nlm.nih.gov/genome/microbes/. The seven sets of

USCGs are from the corresponding publications [1-4, 9, 15, 16] and are listed in Table S1.

References

1. Ciccarelli, F.D., et al., *Toward automatic reconstruction of a highly resolved tree of life.* Science, 2006. **311**(5765): p. 1283-7.
2. Lan, Y., G. Rosen, and R. Hershberg, *Marker genes that are less conserved in their sequences are useful for predicting genome-wide similarity levels between closely related prokaryotic strains.* Microbiome, 2016. **4**(1): p. 18.
3. Wu, D., G. Jospin, and J.A. Eisen, *Systematic identification of gene families for use as "markers" for phylogenetic and phylogeny-driven ecological studies of bacteria and archaea and their major subgroups.* PLoS One, 2013. **8**(10): p. e77033.
4. Wu, M. and J.A. Eisen, *A simple, fast, and accurate method of phylogenomic inference.* Genome Biol, 2008. **9**(10): p. R151.
5. Sunagawa, S., et al., *Metagenomic species profiling using universal phylogenetic marker genes.* Nat Methods, 2013. **10**(12): p. 1196-9.
6. Milanese, A., et al., *Microbial abundance, activity and population genomic profiling with mOTUs2.* Nat Commun, 2019. **10**(1): p. 1014.
7. Quince, C., et al., *DESMAN: a new tool for de novo extraction of strains from metagenomes.* Genome Biol, 2017. **18**(1): p. 181.
8. Ventolero, M.F., et al., *Computational analyses of bacterial strains from shotgun reads.* Brief Bioinform, 2022. **23**(2).

9.   Parks, D.H., et al., *Recovery of nearly 8,000 metagenome-assembled genomes substantially*
expands the tree of life. Nature Microbiology, 2017. **2**(11): p. 1533-1542.
10.   Vetrovsky, T. and P. Baldrian, *The variability of the 16S rRNA gene in bacterial genomes and its*
consequences for bacterial community analyses. PLoS One, 2013. **8**(2): p. e57923.
11.   Wang, Y., H. Hu, and X. Li, *MBMC: An Effective Markov Chain Approach for Binning*
Metagenomic Reads from Environmental Shotgun Sequencing Projects. OMICS, 2016. **20**(8): p.
470-9.
12.   Eisen, J.A., *Environmental shotgun sequencing: its potential and challenges for studying the*
hidden world of microbes. PLoS Biol, 2007. **5**(3): p. e82.
13.   Brooks, J.P., et al., *The truth about metagenomics: quantifying and counteracting bias in 16S*
rRNA studies. BMC Microbiol, 2015. **15**: p. 66.
14.   Wang, Y. and P.Y. Qian, *Conservative fragments in bacterial 16S rRNA genes and primer design*
for 16S ribosomal DNA amplicons in metagenomic studies. PLoS One, 2009. **4**(10): p. e7401.
15.   Alneberg, J., et al., *Binning metagenomic contigs by coverage and composition.* Nat Methods,
2014. **11**(11): p. 1144-6.
16.   Creevey, C.J., et al., *Universally distributed single-copy genes indicate a constant rate of*
horizontal transfer. PLoS One, 2011. **6**(8): p. e22099.
17.   Haroon, M.F., et al., *Anaerobic oxidation of methane coupled to nitrate reduction in a novel*
archaeal lineage. Nature, 2013. **500**(7464): p. 567-70.
18.   Wrighton, K.C., et al., *Fermentation, hydrogen, and sulfur metabolism in multiple uncultivated*
bacterial phyla. Science, 2012. **337**(6102): p. 1661-5.
19.   Wu, M. and A.J. Scott, *Phylogenomic analysis of bacterial and archaeal sequences with*
AMPHORA2. Bioinformatics, 2012. **28**(7): p. 1033-4.
20.   Segata, N., et al., *Metagenomic microbial community profiling using unique clade-specific*
marker genes. Nature Methods, 2012. **9**(8): p. 811-+.
21.   Truong, D.T., et al., *MetaPhlAn2 for enhanced metagenomic taxonomic profiling (vol 12, pg 902,*
2015). Nature Methods, 2016. **13**(1): p. 101-101.
22.   Truong, D.T., et al., *Microbial strain-level population structure and genetic diversity from*
metagenomes. Genome Research, 2017. **27**(4): p. 626-638.
23.   Quince, C., et al., *STRONG: metagenomics strain resolution on assembly graphs.* Genome Biol,
2021. **22**(1): p. 214.
24.   Nayfach, S., et al., *An integrated metagenomics pipeline for strain profiling reveals novel*
patterns of bacterial transmission and biogeography. Genome Research, 2016. **26**(11): p. 1612-
1625.
25.   Forbes, N.S., *Engineering the perfect (bacterial) cancer therapy.* Nat Rev Cancer, 2010. **10**(11):
p. 785-94.
26.   Hartstra, A.V., et al., *Insights into the role of the microbiome in obesity and type 2 diabetes.*
Diabetes Care, 2015. **38**(1): p. 159-65.
27.   Jiang, C., et al., *The Gut Microbiota and Alzheimer's Disease.* J Alzheimers Dis, 2017. **58**(1): p.
1-15.
28.   Ott, S.J., et al., *Detection of diverse bacterial signatures in atherosclerotic lesions of patients*
with coronary heart disease. Circulation, 2006. **113**(7): p. 929-37.
29.   Wang, Y., et al., *Prognostic cancer gene signatures share common regulatory motifs.* Sci Rep,
2017. **7**(1): p. 4750.
30.   Zaky, A., et al., *The Role of the Gut Microbiome in Diabetes and Obesity-Related Kidney*
Disease. Int J Mol Sci, 2021. **22**(17).
31.   Ding, J., et al., *ChIPModule: systematic discovery of transcription factors and their cofactors*
from ChIP-seq data. Pac Symp Biocomput, 2013: p. 320-31.
32.   Harris, M.A., et al., *The Gene Ontology (GO) database and informatics resource.* Nucleic Acids
Res, 2004. **32**(Database issue): p. D258-61.

33.   Tatusov, R.L., et al., *The COG database: an updated version includes eukaryotes.* BMC
363   Bioinformatics, 2003. **4**: p. 41.
364   34.   Young, M.D., et al., *Gene ontology analysis for RNA-seq: accounting for selection bias.* Genome
365   Biol, 2010. **11**(2): p. R14.
366   35.   Zhao, C., X. Li, and H. Hu, *PETModule: a motif module based approach for enhancer target
367   gene prediction.* Sci Rep, 2016. **6**: p. 30043.
368   36.   Li, X., H. Hu, and X. Li, *mixtureS: a novel tool for bacterial strain reconstruction from reads.*
369   Bioinformatics, 2020.
370   37.   Li, X., et al., *BHap: a novel approach for bacterial haplotype reconstruction.* Bioinformatics,
371   2019. **35**(22): p. 4624-4631.
372   38.   Pulido-Tamayo, S., et al., *Frequency-based haplotype reconstruction from deep sequencing data
373   of bacterial populations.* Nucleic Acids Res, 2015. **43**(16): p. e105.
374   39.   Smillie, C.S., et al., *Strain Tracking Reveals the Determinants of Bacterial Engraftment in the
375   Human Gut Following Fecal Microbiota Transplantation.* Cell Host & Microbe, 2018. **23**(2): p.
376   229-+.
377   40.   Cleary, B., et al., *Detection of low-abundance bacterial strains in metagenomic datasets by
378   eigengenome partitioning.* Nat Biotechnol, 2015. **33**(10): p. 1053-60.
379   41.   Rinke, C., et al., *Insights into the phylogeny and coding potential of microbial dark matter.*
380   Nature, 2013. **499**(7459): p. 431-7.
381   42.   Chen, I.A., et al., *IMG/M v.5.0: an integrated data management and comparative analysis system
382   for microbial genomes and microbiomes.* Nucleic Acids Res, 2019. **47**(D1): p. D666-D677.
383   43.   Federhen, S., *The NCBI Taxonomy database.* Nucleic Acids Res, 2012. **40**(Database issue): p.
384   D136-43.
385   44.   Langr, J. and V. Bok, *GANs in action : deep learning with generative adversarial networks.*
386   2019, Shelter Island, New York,: Manning Publications. xxiii, 214 pages.
387   45.   Li, X., et al., *Integrative analyses shed new light on human ribosomal protein gene regulation.*
388   Sci Rep, 2016. **6**: p. 28619.
389   46.   Shi, J.Q. and T. Choi, *Gaussian process regression analysis for functional data.* 2011, Boca
390   Raton, FL: CRC Press. xix, 196 p.
391   47.   Talukder, A., et al., *EPIP: a novel approach for condition-specific enhancer-promoter interaction
392   prediction.* Bioinformatics, 2019. **35**(20): p. 3877-3883.
393   48.   Tatusov, R.L., E.V. Koonin, and D.J. Lipman, *A genomic perspective on protein families.*
394   Science, 1997. **278**(5338): p. 631-7.
395   49.   Katoh, K., et al., *MAFFT: a novel method for rapid multiple sequence alignment based on fast
396   Fourier transform.* Nucleic Acids Res, 2002. **30**(14): p. 3059-66.

Figure 1. USCG sets and functional categories. (A) Overlap of the seven USCG sets. (B) The

percentage of the functional categories of the USCGs in the seven sets.

Figure 2. The universalism and uniqueness of the three USCG sets.

Table 1. Universalism and uniqueness of USCGs.

	mean			Standard deviation			minimum			median			maximum		
	C	W	A	C	W	A	C	W	A	C	W	A	C	W	A
fetchMG	99.0	98.6	99.0	0.5	2.7	0.8	97.7	83.9	95.9	99.3	99.3	99.3	99.6	99.7	99.6
BLAST	96.1	88.9	94.2	3.0	7.0	4.5	87.1	58.9	76.7	96.7	90.9	96.0	99.7	96.8	99.0
universal	100	99.8	99.9	3.4	9.4	5.3	99.7	93.4	99.0	100	99.9	100	100	100	100
unique	98.5	97.3	98.2	1.0	3.9	2.2	95.4	82.6	89.0	99.0	99.1	99.1	99.5	99.6	99.5

The subcolumns with the name C, W, and A refer to the corresponding percentage for the

USCGs from Creevey et al., Wu et al., and Alneberg et al., respectively.

440 **Table 2**. **The cumulative distribution of the PID of species pairs and strain pairs**.

PID cutoff		<99.5%	<99.6%	<99.7%	<99.8%	<99.9%	<=1
Species	Creevey et al.	97.2	97.7	98.1	98.6	99	100
	Wu et al.	97.2	97.7	98.2	98.5	98.8	100
	Alneberg et al.	97.3	97.8	98.2	98.5	98.8	100
Strains	Creevey et al.	7.6	8.8	10.6	15.5	25.4	100
	Wu et al.	5.2	6.1	7.7	20.4	43.1	100
	Alneberg et al.	5.9	6.7	8.5	14.9	36.9	100

441 The PID cutoff is provided in the first row. The percentage of pairs with their PID smaller than

442 the cutoff is provided in the corresponding entry.

443