# Altered basal ganglia output during self-restraint.

Bon-Mi Gu[1*] and Joshua D Berke[2]

[1]Department of Neurology and Neurological Sciences, Stanford University, Stanford, CA 94305, USA.

[2]Department of Neurology, Department of Psychiatry and Behavioral Sciences, Neuroscience Graduate Program, Kavli Institute for Fundamental Neuroscience, Weill Institute for Neurosciences, University of California, San Francisco.

*Correspondence: bonmigu@stanford.edu

## Abstract

Suppressing actions is essential for flexible behavior. Multiple neural circuits involved in behavioral inhibition converge upon a key basal ganglia output nucleus, the substantia nigra pars reticulata (SNr). To examine how changes in basal ganglia output contribute to self-restraint, we recorded SNr neurons during a proactive behavioral inhibition task. Rats responded to *Go!* cues with rapid leftward or rightward movements, but also prepared to cancel one of these movement directions as a *Stop!* cue might occur. This action restraint – visible as direction-selective slowing of reaction times – altered both rates and patterns of SNr spiking. Overall firing rate was elevated before the *Go!* cue, and this effect was driven by a subpopulation of direction-selective SNr neurons. In neural state space, this corresponded to a shift away from the restrained movement. Furthermore, SNr activity showed an increase in spike variability during proactive inhibition, similar to that reported with movement slowing in Parkinson's Disease. Increased spike variability corresponded to variable state-space trajectories, slowing reaction times via reduced preparation to move. These findings open new perspectives on basal ganglia mechanisms for movement preparation and cognitive control, in both healthy states and brain disorders.

Keywords: behavioral inhibition, substantia nigra pars reticulata, population dynamics, spike variability, Parkinson's disease.

**Introduction**

Fluid, efficient behavior often involves simply triggering well-learned behaviors. However, flexibility requires that such behaviors can be suppressed, should circumstances change. This capacity for behavioral inhibition is considered central to cognitive control (Bari and Robbins, 2013), and is compromised in a range of neurological and psychiatric disorders (Chambers *et al.,* 2009).

Behavioral inhibition can be 'reactive' – for example, promptly responding to an unexpected *Stop!* cue by cancelling upcoming actions. Reactive inhibition has been shown to involve fast cue responses in frontal cortex and basal ganglia pathways (Jahanshahi *et al.,* 2015; Wager *et al.,* 2005), including from the subthalamic nucleus (STN) to SNr (Schmidt *et al.* 2013). This rapid response to stimuli can transiently, and broadly, retard action initiation, providing time for a second set of basal ganglia mechanisms to cancel actions (Mallet *et al.* 2016; Schmidt and Berke 2018).

By contrast, 'proactive' inhibition refers to a state of preparation, in which particular actions are restrained (Cai *et al.,* 2011, Claffey *et al.,* 2010). Proactive inhibition has been argued to be especially important for human life (Aron 2011), and is behaviorally apparent as longer reaction times (RTs) selectively for the restrained action. The underlying mechanisms are not well understood, but have been proposed (Aron 2011) to involve the pathway from striatum "indirectly" to SNr, via globus pallidus pars externa (GPe). In a prior study (Gu *et al.* 2020) we therefore recorded from GPe neurons as rats engaged proactive inhibition towards a specific action. The state of being prepared to stop did not involve an overall net change in GPe firing rate, but rather a shift at the level of neural population dynamics away from action initiation, and towards the alternative action. One objective of the present work was to determine whether a corresponding preparatory change in population dynamics is visible "downstream" in SNr, thereby altering basal ganglia output.

Basal ganglia output neurons are thought to affect behavior not just via their firing rates, but also via their firing patterns and synchrony (Rubin *et al.,* 2012). In particular, Parkinson's disease (PD) is associated with an increase in firing variability and synchronous bursting, often without rate changes (Lobb, 2014; Willard *et al.,* 2019). As PD is characterized by slowed movement initiation (Low *et al.*, 2002), we assessed whether related physiological changes are present when movements are slowed as the result of proactive inhibition.

**Results.**

**Reaction times are selectively slowed with proactive inhibition**.

We used a selective proactive inhibition task (Gu *et al.,* 2020), a variant of our extensively-characterized rat stop-signal task (Leventhal *et al.,* 2012; Schmidt *et al.*, 2013; Mallet *et al.,* 2016) (Fig. 1A). Rats start a trial by nose-poking an illuminated start-port. To proceed, they need to maintain this position for a variable delay (500-1250ms, uniform distribution) until a *Go!* cue is presented (1 kHz or 4 kHz tone, indicating a leftward or rightward movement respectively). If, the movement is initiated rapidly after the *Go!* cue (RT limit < 800ms) and completed correctly and promptly (movement time limit, MT < 500ms), rats are rewarded with a sugar pellet dispensed at a separate food hopper. On a subset of trials, the *Go!* cue is followed by a *Stop*! cue (white noise burst; delay after *Go!* cue onset = 100 - 250ms). This indicates that the rat should not initiate a movement, and instead hold their nose in the Center port for at least 800ms (after *Go!* cue onset) to trigger reward delivery.

To probe selective proactive inhibition, the three possible start ports were associated with different *Stop!* cue probabilities (counterbalanced across rats; Gu *et al.*, 2020, Fig. 1B, Table 1). These were: no possibility of *Stop!* cue ("No-Stop"); 50% probability that a left *Go!* cue will be followed by the *Stop!* cue ("Maybe-Stop-left"); and 50% probability that a right *Go!* cue will be followed by the *Stop!* cue ("Maybe-Stop-right"). We obtained SNr microelectrode recordings from rats (n=10) that had successfully learned this proactive task, as indicated by

proactive control significantly slowing RTs for movements contraversive to the implant side. For example, if electrodes were placed in the right SNr, contraversive proactive inhibition would mean longer RTs for Maybe-Stop-left trials, compared to No-Stop trials. In the same rats we compared SNr activity in sessions in which this proactive effect was significant (n=60), to sessions in which it was not (n=25; Fig 1C).

**Selective proactive inhibition recruits specific SNr subpopulations.**

We examined the task-related activity of individual SNr neurons (n=446; mean firing rate=38Hz, locations are shown in Supp. Fig. 1) recorded during the sessions with significant proactive inhibition. We first compared overall cell activity between Maybe-Stop-Contra and No-Stop trials. We focused on the epoch just before the *Go!* cue, as we presume that this time is critical for being "prepared-to-stop". Average firing rates were significantly higher in the Maybe-Stop-Contra condition (Fig. 2A, top), and this difference was generated by a significant fraction of SNr neurons (Fig. 2A, bottom). Example cells show significant differences before the *Go!* cue between conditions (Fig. 2B). This elevated SNr firing with action restraint was not observed in the sessions without significant behavioral evidence for proactive inhibition (Fig. 2A, right; 173 neurons, mean FR=44Hz).

A common categorization of SNr neurons distinguishes those that increase, versus decrease, firing rate in conjunction with behavioral events (e.g. Bryden *et al.*, 2011; Fan *et al.,* 2012; Gulley *et al.*, 2002; Sato and Hikosaka, 2002). Decreases in firing shortly before movement onset are thought to enable movements, by disinhibiting downstream structures including the superior colliculus (Hikosaka and Wurtz, 1983b). If "decrease-type" neurons are receiving more excitation during proactive inhibition, this might delay their decrease to the level needed to release movements, resulting in longer reaction times. We therefore hypothesized that proactive inhibition is associated with elevated firing specifically of decrease-type neurons.

We categorized cells as increase-type or decrease-type based on their change in firing rate during the 200ms preceding movement onset (Fig. 2C). Increase-type were more numerous, as previously reported (Bryden *et al.*, 2011; Joshua *et al.*, 2009). Contrary to our hypothesis, both increase-type and decrease-type neurons contributed to elevated SNr activity with action restraint (Fig. 2D, left). There was no relationship between the extent of elevated firing in Maybe-Stop trials, and the firing rate change before movement onset (Fig. 2D, right).

We then examined whether proactive inhibition effects were related to neurons' response selectivity, operationally defined as the firing rate difference between Contra and Ipsi actions just before Center Out (Fig. 2E, inset). We found a significant relationship: neurons more active just before Ipsi compared to just before Contra movements ("Ipsi > Contra") showed elevated firing when Contra actions might need to be cancelled (Fig. 2F,G). No such relationship was found for neurons with the opposite selectivity ("Contra > Ipsi"; Fig. 2F,G).

We next considered the interaction between direction selectivity and increases vs. decreases in firing, in proactive inhibition. A cell could be classified as "Ipsi > Contra" because it preferentially increases firing with Ipsi movements, or because it preferentially decreases firing with Contra movements. We found that both subtypes had elevated firing before the *Go!* cue on Maybe-Stop-Contra trials (Supp. Fig. 2). Therefore, the proactive effect was not simply a matter of SNr cells that pause with Contra movements starting from a higher baseline rate, though this may contribute.

**Restraining one action biases population dynamics towards the alternative action.**

The elevated average firing rate of Ipsi > Contra cells suggests a preparatory bias towards Ipsi action, at times when Contra actions might need to be cancelled. To examine this further we turned to a state-space analysis, as we previously used with GPe data (Gu et al., 2020). We extracted principal components from the average firing rates of each neuron during Contra and Ipsi movements (Supp. Fig. 3) and used these to visualize neural population

trajectories (Fig. 3A, top). We defined "Initiation" and "Selection" axes as the common, and distinct, aspects of these trajectories respectively, during the 200ms before action initiation. In particular, the Initiation Axis is a line drawn between the average state space positions at 200ms and 0ms relative to Center Out (disregarding the direction of movement). The Selection Axis connects the midpoints of the Contra and Ipsi trajectories (averaging across the same time epoch).

When we projected Maybe-Stop-Contra data into this space, we found that overall SNr population activity showed a clear shift before the *Go!* cue (Fig. 3A, bottom). This shift was especially pronounced on the Selection Axis, with a highly significant bias towards Ipsi (Fig. 3B). There was also a significant bias on the Initiation Axis (towards movement). These two biases were associated with distinct cell subtypes (Fig. 3C, E). Ipsi > Contra cells showed a strong Selection Axis bias, without a significant Initiation Axis bias (Fig. 3D), and the converse was seen for Contra > Ipsi cells (Fig. 3F). This finding provides further evidence that Ipsi > Contra cells generate an Ipsi bias during selective proactive inhibition.

Moreover, the state of preparation affects the subsequent outcome of the trial. The neural trajectory showed that trajectories during wrong choice trials were further biased toward Ipsi action and towards initiation at the time of *Go!* cue (Supp. Fig. 3D), consistent with our prior GPe results (Gu *et al.,* 2020). Trials in which rats failed to initiate actions (limited hold violations) did not show any statistically significant bias at the *Go!* cue time, although the trajectory after the *Go!* cue showed movement away from action initiation.

**Less regular firing with proactive inhibition.**

As movement slowing has been associated with changes in basal ganglia firing patterns in Parkinson's Disease (Sharott *et al.*, 2014; Tai 2022), we examined whether changes in SNr firing patterns accompany slower movement initiation during proactive inhibition. For each neuron we calculated the coefficient of variation (CV) of inter-spike intervals, and the proportion

of spikes within bursts (Fig. 4A; using the Poisson surprise method; Legendy and Salcman 1985). To obtain sufficient numbers of inter-spike-intervals, both analyses used longer (3s) epochs before the *Go!* cue.

Proactive slowing was associated with altered spiking variability, in several different ways. Neurons recorded in sessions with significant proactive inhibition behavior (n=446) showed more irregular and bursty firing compared to neurons (n=173) in sessions that did not (Fig. 4B). The degree of irregularity was correlated with the mean session-wide reaction times (Supp. Fig. 4A). Increased irregularity and bursting were also seen during Maybe-Stop-Contra trials compared to No-Stop trials in the sessions with significant proactive inhibition effect (Fig. 4C, left), but not in the sessions without it (Fig. 4C, right). Moreover, the degree of increased irregularity between conditions was correlated with the magnitude of the proactive effect on reaction times (Supp. Fig. 4B).

**Proactive inhibition increases the variability of neural trajectories.**

How could increased spiking variability contribute to the slowing of RTs? At the population level, spike variability would correspond to more erratic state-space trajectories before the *Go!* cue (Fig. 5A). This would result in a more variable state at the (unpredictable) time of *Go!* cue onset. If effective movement preparation involves positioning neural activity within a "optimal subspace", as previously proposed (Churchland *et al.* 2006), this more variable state would in turn result in longer and more variable RTs.

To assess this idea we quantified trajectory fluctuations on individual trials (Fig. 5B-D). We included sessions (n=27) in which more than 5 neurons were recorded simultaneously, and examined trajectories in the 3s window before the *Go!* cue (for comparison to our CV measure). In the same way as CV measures variability over time of an individual neuron (one dimension) compared to its mean rate, we can measure the activity variability of a population (*n* dimensions) by comparing the Euclidean distance of each time point to the mean position (Fig. 5C).

As expected, there was a strong relationship between CV of individual neurons and their corresponding population trajectory variability (Fig. 5E). Furthermore, trajectory variability was significantly increased by proactive inhibition (Figs. 5F). Trajectory variability increased during the hold period (examining the 0.8s epoch just before the *Go!* cue, excluding trials with hold duration <0.8s, Wilcoxon signed rank test, $p$=0.005), and was also observed even beforehand (0.8s before Nose In, Wilcoxon signed rank test, $p$=0.006). State space position at *Go!* cue onset was indeed more variable with proactive inhibition (Fig. 5G), and this variability was correlated with RT (Fig. 5H).

**Proactive modulation of firing rate, and variability, are dissociable**.

Changes in preparation to move or stop can be evoked by explicit cues, as in our task design, but also by the subject's ongoing experience – notably, the events on the previous trial (Bissett and Logan, 2011; Pouget *et al.* 2011). We therefore examined how such experience affects the proactive influence over SNr firing described above. The behavioral effect of slower RT on Maybe-Stop trials was present regardless of previous trial type (Fig. 6A, top row), as was the SNr state-space bias toward Ipsi (Fig. 6A, third row). By contrast, the increases in SNr spiking CV were selectively observed following Stop-fail trials (Fig. 6A, bottom). This suggests that the increased CV occurs when rats are especially concerned to avoid hasty responses, having just failed to sufficiently restrain behavior on the previous trial.

**Discussion**

Taking advantage of a sophisticated behavioral task, we found that that action restraint is reflected in altered basal ganglia output, in two distinct respects. As rats wait for a *Go!* cue, if it may be followed by a *Stop!* cue, a subpopulation of movement-selective SNr neurons shows elevated firing. This change is associated with a shift in population activity away from the specific restrained action and towards the alternative. At the same time, increased variability of

SNr firing results in a more variable state at the time of the *Go!* cue. This variability is associated with slowed RTs, plausibly because effective movement preparation involves achieving a more constrained range of network states. Moreover, these firing rate and variability modulations were independently controlled by different trial outcomes, suggesting multiple mechanisms involved in selective proactive inhibition.

**Basal ganglia dynamics and the nature of restraint**

Beyond simple sensory or motor correlates, prior studies have found modulation of SNr firing rates by "internal" factors such as task context (Hikosaka and Wurtz 1983a; Lintz and Felsen 2016). Basal ganglia activity is especially affected by changes in reward expectation (Lauwereyns *et al*. 2002) including in SNr (Sato and Hikosaka 2002; Yasuda and Hikosaka 2017), producing faster RTs towards more rewarding movements. In our proactive task the Maybe-Stop direction receives rewards at a lower rate (due to failures-to-Stop), so asymmetrical reward expectation might be at least partially responsible for the neural and choice bias towards Ipsi movements on Maybe-Stop-Contra trials. However, the slower Contra RTs on Maybe-Stop-Contra trials were not accompanied by faster Ipsi RTs, suggesting that proactive inhibition of Contra movements did not consist simply of greater preparation of Ipsi movements.

The SNr state-space bias during proactive inhibition may arise from the "indirect pathway" projection from GPe, which we found to have a similar neural bias towards Ipsi movements in a prior study (Gu *et al*., 2020). However, proactive inhibition produced no overall change in GPe firing rates, and we were not able to identify a distinct subpopulation of modulated GPe neurons. The reasons for this GPe : SNr difference are not clear. Recent modeling has demonstrated that the influence of the GABAergic GPe inputs over SNr neurons can be far more complex (Simmons *et al*. 2020) than shown in classic rate models of basal ganglia function, for example switching from inhibitory to excitatory in an activity-dependent manner (Phillips *et al.* 2020). Alternatively, the shift in SNr Ipsi > Contra neuron firing with

proactive inhibition may reflect other basal ganglia inputs, particularly the direct pathway input from striatum, or hyperdirect from the subthalamic nucleus (Schmidt *et al.* 2013).

**Behavioral control and neural variability**

Our study also leaves unresolved the origins of the increased SNr spike variability we observed with proactive inhibition. Like GPe and STN neurons, SNr neurons are intrinsic pacemakers that spontaneously fire regularly even if their inputs are blocked (Zhou and Lee 2011). These inputs – which include the aforementioned direct and indirect pathways, and also extensive local collaterals within SNr (Brown *et al.*, 2014; Mailly *et al.,* 2003) – thus alter SNr spiking by either accelerating or delaying the occurrence of the next spike. Changes in spike time variability presumably reflect either changes in local network properties, e.g. due to neuromodulation (Delaville *et al.* 2012) or changes in the statistics of extrinsic inputs.

In the cortex, decreases in spike variability have been previously linked to various processes including stimulus onset (Churchland *et al.*, 2010), attention (Cohen and Maunsell, 2009), and movement preparation (Churchland *et al.,* 2006), and there is some evidence for similar phenomena in the striatum (Berke 2011). To our knowledge, there are no prior observations of specific task cues producing *increases* in variability. Our finding that variability increases with proactive inhibition – in particular following Stop-fail trials – suggests that variability may be actively elevated as part of a behavioral strategy. This would fit with proposals that neural variability can confer behavioral advantages, such as increased flexibility (Waschke *et al.* 2021). Alternatively, increased neural variability with proactive inhibition may simply reflect the *absence* of a reduction in variability that accompanies movement preparation. In this way, preparation to stop would consist, at least in part, of less preparation to go. Either way, our results demonstrate that cognitive control strategies can operate through deliberate shifts in neural variability.

**From proactive inhibition to Parkinson's disease**

      Connecting single-neuron measures of variability to state-space concepts of movement preparation offers an intriguing perspective on PD, which is characterized by both elevated neural variability (Dorval *et al.*, 2008; Lobb, 2014; Willard *et al.,* 2019) and slowed movements. Irregular firing produces more erratic trajectories; when the *Go!* cue arrives, the network is at a greater average distance from an "optimal subspace" for initiating movements (Churchland *et al.* 2006). This increased variability could help explain why average RTs and movement times are slowed in PD, yet the fastest movements are still preserved (Mazzoni *et al.* 2007). An important goal for future studies is to better understanding how shifts in neural variability occur in the basal ganglia, and whether they originate from the same mechanisms during both proactive inhibition and Parkinson's Disease.

**Acknowledgements.** We thank Charles Wilson and members of the Berke Lab for valuable feedback. This work was supported by UCSF, NIH (R01MH101697, R01NS123516), and CHDI.
**Author contributions.** B.G., Conceptualization, Data curation, Software, Formal analysis, Investigation, Visualization, Methodology, Writing – original draft; J.D.B., Conceptualization, Resources, Supervision, Funding acquisition, Writing - original draft, Writing - review and editing
**Declaration of interests.** The authors have no competing interests.

**Figure legends**

**Figure 1. Selective proactive inhibition**. (A) Left, operant box configuration, with dashed red lines indicating photobeams for nose detection; right, event sequence for Go and Stop trials. RT, reaction time; MT, movement time; SSD, stop-signal delay; Reward, delivery of a sugar pellet to the food port. (B) Trial start location indicates stop probabilities. In this example configuration with left SNr recording, illumination of the middle hole indicates that *Go!* cues instructing rightward movement may be followed by a *Stop!* cue, but *Go!* cues instructing leftward

movements will not ("Maybe-Stop-Contra"). (C) Overall, the sessions in which SNr units were successfully recorded showed strong proactive inhibition effect (n=85, Wilcoxon signed rank tests, *p*=8.2e-15). Among them, the individual sessions are considered to show proactive inhibition (red) if the reaction time difference between Maybe-Stop-Contra and No-Stop trials is statistically significant (one-tail Wilcoxon rank sum test, *p*<0.05). (D) The proactive sessions show RT slowing selectively for the Maybe-Stop direction (Wilcoxon signed rank tests on median RTs of Maybe-Stop-Contra versus No-Stop: Contra cues: $z = 6.8$, $p = 1.1e\text{-}11$; Ipsi cues: $z = \text{-}1.2$, $p = 0.22$). Additionally, on Maybe-Stop-Contra trials rats were more likely to fail to respond quickly enough (RT limit errors; Wilcoxon signed rank tests, $z = 6.4$, $p = 1.5e\text{-}10$) and to select the wrong choice (i.e. not matching the *Go!* cue; Wilcoxon signed rank tests, $z = 6.5$, $p = 1.1e\text{-}10$). Error bars, S.E.M. across n = 60 sessions. RT limit error = nose remained in Center port for >800ms after *Go!* cue onset; MT limit error = movement time between Center Out and Side port entry >500ms.

**Figure 2. Elevated firing rates of specific SNr subpopulations with selective proactive inhibition.** (A) Top left: before the Go! cue SNr firing is elevated in Maybe-Stop-Contra compared to No-Stop conditions. This occurs selectively in sessions with behavioral evidence of proactive inhibition (left). Each neuron's firing rate is *Z*-scored and averaged (over all trials in which a *Go!* cue was presented, regardless of *Stop!* cues) Shaded band, +-S.E.M across n=446 neurons. Thicker lines indicate significant differences between conditions (*p*<0.05; *t*-test at each time point). Bottom left: fraction of SNr neurons whose firing rate significantly differs between conditions, across time (*p*<0.05; *t*-test in each 50ms bin). Higher firing with Maybe-Stop-Contra, No-Stop conditions are shown as positive (orange) or negative (blue), respectively. Horizontal grey lines indicate thresholds for a significant proportion of neurons (binomial test, *p*<0.05 without or with multiple-comparisons correction, thin and thick grey lines respectively). Light and dark color-filled bars are those for which the threshold was crossed without or with multiple-

comparisons correction. Right, sessions without significant behavioral evidence of proactive

inhibition do not show this firing rate difference between conditions (same format as left panels;

n=173 neurons). (B) Two individual example neurons demonstrating the proactive elevation of

firing rate before the *Go!* cue. (C) Neurons were categorized as decrease-type or increase-type,

based on an "Initiation Score". We defined the "Initiation Score" for each neuron as the change

in (*Z*-scored) firing rate in the 0.2s before Center Out (inset shows example neuron).  Plots

show average firing of each subpopulation (top; +-S.E.M) and individual cells sorted by Initiation

Score (bottom). (D) No relation between Initiation Score and proactive inhibition (assessed as

the difference between Maybe-Stop-Contra and No-Stop trials, in the 200ms before *Go!* cue).

Bar graph (inset) shows that on average, both increase- and decrease- type neurons modestly

increase firing with proactive inhibition (Wilcoxon signed rank tests). (E) We defined the

"Selection Score" for each neuron as the integral of the difference in (*Z*-scored) firing rate

between Contra and Ipsi actions during the 0.2s epoch before Center Out. Remainder of panel

is as C, but for Selection Score. (F) Significant negative correlation between Selection Score

and proactive inhibition. Bar graph (inset) shows that Ipsi>Contra neurons preferentially

increase activity on Maybe-Stop-Contra trials. (G) Same result as F, using the format of panel A

to illustrate time course.

**Figure 3**. **Biased SNr population dynamics during proactive inhibition.** (A). Top, Overall

SNr state-space trajectories before and during Contra (blue) and Ipsi movements (green), in the

state space of the first 3 principal components (PCs). Trajectories show +-250ms around

detected movement onset (Center Out, larger circles), with each small circle separated by 4ms.

'Initiation Axis' joins the positions (black asterisks) 200ms before and at action initiation

(averaging Contra and Ipsi actions). 'Selection Axis' joins the means of each trajectory in the

same epoch (colored asterisks). Bottom, comparing Maybe-Stop-Contra (orange) and No-Stop

(blue) trials (+-100ms around Go! cue) in the same state space as A. Overall population state is

visibly biased toward Ipsi along the Selection Axis. (B) Permutation tests of bias along each axis (average during -200ms to 0 relative to *Go!* cue), using all 10 PCs. Red bars, observed results; Grey, distributions of surrogate data from 10000 random shuffles of trial type labels. Dark grey indicates 5% of distributions at each tail. (C,D). As A-B, but for Ipsi > Contra cells only. The proactive bias towards Ipsi along the Selection Axis is more clearly visible. (E, F). As A-B, but for Contra > Ipsi cells. These do not show a proactive bias on the Selection Axis, but on the Initiation Axis instead.

**Figure 4**. **SNr firing is more irregular and bursty with proactive inhibition.** (A) Two example neurons, showing spike rasters (left; aligned on *Go!* cues) and autocorrelograms (right; for the 3s preceding *Go!* cues). CV: coefficient of variation of inter-spike intervals. (B) Elevated CV and burst ratio of neurons recorded in sessions with behavioral evidence of proactive inhibition, compared to sessions without (Wilcoxon rank sum tests). This effect was also seen at the level of individual rats (Supp. Fig4C). (C) Within-session comparison of Maybe-Stop-Contra and No-Stop trials shows increased CV and burst ratio when proactive inhibition is engaged (left). This effect is not present on sessions without proactive slowing of reaction times (right). Wilcoxon signed rank tests.

**Figure 5. Altered variability of state trajectories with proactive inhibition.** (A) Conceptual illustration for the relationship between trajectory variability and RTs. Larger fluctuations in trajectory will tend to result in a position further away from the "optimal subspace" when the *Go!* cue arrives. (B) Individual trial trajectories (10 trials each for Contra and Ipsi movements) for one example session (n=46 neurons). (C) Trajectory variability was defined as the mean of the Euclidean distances at each time point to the mean position over the trajectory (in the full neural state space, without PCA). (D) Example trajectories for Maybe-Stop-Contra (orange) and No-Stop (blue) trials, before the *Go!* cue. (E) Correlations between CV and trajectory variability for

each session. (F) Trajectory variability increased on Maybe-Stop-Contra, compared to No-Stop trials. (G) State space position at *Go!* cue (-200ms to 0) was more variable for Maybe-Stop-Contra, compared to No-Stop trials. (H) Variability of state space position at Go! cue (-200ms to 0) is positively correlated with RT. Trials with less than 100ms reaction times were excluded because they would have already initiated the movement trajectory. For (E)-(H), sessions with more than 5 neurons (# of session = 27) were used for analysis (Wilcoxon signed rank test on each session values).

**Figure 6. Feedback effect by previous trials and relation to trial outcomes.** Reaction times, firing rates, selection axis and CVs of maybe-stop-contra trials plotted by the previous trial types to show if feedback controls proactive inhibition effects. Values in No-stop condition were used as baseline. Ipsi > contra cells are used for firing rates and selection axis values (during 200ms before *Go!* cue) and all cells are used for CV calculation. **$p <0.05$ with Bonferroni; *$p<0.05$ without multiple comparison correction (Wilcoxon signed rank test).

**Tables**

**Table 1. Information on individual rats.**

| RAT | Port Location (port1 port2 port3) | # of sessions (with/without proactive inhibition effect) | # of cells (from sessions with/without proactive inhibition effect) | Sessions with contralateral proactive inhibition effect | | |
|---|---|---|---|---|---|---|
| | | | | Contralateral RTs (Maybe-Stop-Contra/No-Stop conditions, ms) | Ipsilateral RTs (Maybe-Stop-Contra/No-Stop conditions, ms) | Stop success rates (%) |
| 1015 | 0% L50% R50% | 7/1 | 184/20 | 563/363 | 428/415 | 55.3 |
| 1019 | L50% 0% R50% | 5/2 | 6/4 | 369/275 | 271/321 | 47.8 |
| 1042 | L50% 0% R50% | 2/0 | 3/0 | 399/293 | 277/377 | 81.1 |

| 1043 | L50% R50% 0% | 8/1 | 30/5 | 474/353 | 384/387 | 70.5 |
|------|--------------|-----|-------|---------|---------|------|
| 1063 | L50% R50% 0% | 8/4 | 21/15 | 347/226 | 266/191 | 26.2 |
| 1064 | L50% 0% R50% | 6/4 | 23/26 | 446/338 | 303/309 | 57.3 |
| 1098 | 0% L50% R50% | 5/2 | 5/2 | 456/330 | 324/327 | 42.2 |
| 1202 | 0% R50% L50% | 10/1 | 114/12 | 494/328 | 359/356 | 56.3 |
| 1296 | R50% 0% L50% | 7/2 | 53/11 | 435/330 | 344/345 | 61.5 |
| 1328 | R50% L50% 0% | 2/8 | 7/78 | 370/298 | 275/368 | 51.8 |

**Methods**

**Animals.** All animal experiments were approved by the University of California, San Francisco Committee for the Use and Care of Animals. Adult male Long-Evans rats were housed on a 12 hr/12 hr reverse light-dark cycle, with training and testing performed during the dark phase.

**Behavior.** The rat proactive stop task has been previously described in detail (Gu *et al.,* 2020). Briefly, rats were trained in an operant chamber (Med Associates, Fairfax VT) which had five nose-poke holes on one wall, a food dispenser on the opposite wall, and a speaker located above the food port. Each trial starts with one of the three starting ports illuminated to indicate the *Stop!* cue probabilities ('Light On'), and the same start port was repeated for 10-15 trials. Inter-trial intervals were randomly selected between 5 and 7s.

**Electrophysiology.** We recorded SNr data from ten rats (all animals in which we successfully recorded SNr neurons during contraversive proactive inhibition). Each rat was implanted with 15 or 30 tetrodes in bundles bilaterally targeting SNr (Supp. Fig1). Wide-band (0.1–9000 Hz) electrophysiological data were recorded with a sampling rate of 30000/s using an Intan RHD2000 recording system (Intan Technologies). All signals were initially referenced to a skull screw (tip-flattened) on the midline 1 mm posterior to lambda. For spike detection we re-

referenced to an electrode common average, and wavelet-filtered (Wiltschko *et al.*, 2008) before thresholding. For spike sorting we performed automatic clustering units using MountainSort (Chung *et al.*, 2017) followed by manual curation of clusters. Approximately every 2-3 sessions, screws were turned to lower tetrodes by 100-160 μm; to avoid duplicate neurons, we did not include units from the same tetrode from multiple sessions unless the tetrodes had been moved between those sessions. We further excluded a small number of neurons on the same tetrode that appeared to be potential duplicates based on waveforms and firing properties (e.g. firing rates, CV, and behavioral correlations), even though the tetrode had been moved. After recording was complete, we anesthetized rats and made small marker lesions by applying 5-10 μA current for 20 s on one or two wires of each tetrode. After perfusing the rats, tissue sections (at 40 μm) were stained with cresyl violet or with CD11b antibody and compared to the nearest atlas section (Paxinos, 2006).

**Data analysis.** *Bursting.* Spike bursts were detected using the Poisson surprise method (Legendy and Salcman 1985) with a surprise threshold of 5, and the burst ratio was calculated as the number of spikes fired in bursts divided by the number of all spikes in each unit. The CV of inter-spike intervals and burst ratio were calculated in each trial using a 3s time window before the *Go!* cue, and averaged across all trials (for sessions with/without proactive inhibition comparison) or across each trial type (for conditional differences).

*PCA analysis* was done largely as previously described (Gu *et al.*, 2020). First, firing rates were smoothed using a Gaussian kernel (30ms SD) and normalized (Z-scored). Then, the normalized average time series for contra and ipsi actions (Supp. Fig. 2A) were used for PCA. To match the analysis to the Initiation and Selection scores (Figure 2), we defined the Initiation Axis by connecting state space points 200ms before and at action initiation (averaging Contra and Ipsi actions), and the Selection Axis by connecting the mean of Contra trajectories to the mean of Ipsi trajectories (again using the epoch from -200 to 0ms relative to action initiation).

*Variability of neural trajectories:* To examine how the increased variability of individual neurons affects population dynamics, we calculated the temporal variability of neural state-space trajectories. We included data from 3s time windows before the *Go!* cue, in each session (n=27) with at least 5 recorded units and a significant proactive inhibition effect. Trajectory variability V corresponds to the mean Euclidean distance between each time point of the trajectory and the center of the trajectory, i.e. how much the trajectory wanders around. For this analysis, we did not apply PCA to reduce dimensionality. Thus, for example, if the normalized FR of population neurons at time (t) is defined as a vector $\vec{R_t}$ , and the center of trajectory (mean FR across t=1…T, where T = 1500 for 3s time window with 500Hz sampling rate) is defined as a vector $\vec{R_c}$ , Variability (V) of a neural trajectory is defined as:

$$V = \frac{1}{T} \sum_{t=1}^{T} || \vec{R_t} - \vec{R_c} ||$$

**References**

Aron AR. 2011. From reactive to proactive and selective control: developing a richer model for stopping inappropriate responses. Biological Psychiatry 69:e55–e68. DOI: https://doi.org/10.1016/j.biopsych.2010.07.024, PMID: 20932513

Bari A, Robbins TW. Inhibition and impulsivity: behavioral and neural basis of response control. Prog Neurobiol. 2013 Sep;108:44-79. doi: 10.1016/j.pneurobio.2013.06.005. Epub 2013 Jul 13. PMID: 23856628.

Berke JD. Functional properties of striatal fast-spiking interneurons. Front Syst Neurosci. 2011 Jun 20;5:45. doi: 10.3389/fnsys.2011.00045. PMID: 21743805; PMCID: PMC3121016.

Bissett PG, Logan GD. Balancing cognitive demands: control adjustments in the stop-signal paradigm. J Exp Psychol Learn Mem Cogn. 2011 Mar;37(2):392-404. doi: 10.1037/a0021800. PMID: 21171806; PMCID: PMC3064521.

Brown J, Pan WX, Dudman JT. The inhibitory microcircuit of the substantia nigra provides feedback gain control of the basal ganglia output. Elife. 2014 May 21;3:e02397. doi: 10.7554/eLife.02397. PMID: 24849626; PMCID: PMC4067753.

Bryden DW, Johnson EE, Diao X, Roesch MR. Impact of expected value on neural activity in rat substantia nigra pars reticulata. Eur J Neurosci. 2011 Jun;33(12):2308-17. doi: 10.1111/j.1460-9568.2011.07705.x. Epub 2011 Jun 6. PMID: 21645133; PMCID: PMC3334837.

Cai W, Oldenkamp CL, Aron AR. 2011. A proactive mechanism for selective suppression of response tendencies. Journal of Neuroscience 31:5965–5969. DOI: https://doi.org/10.1523/JNEUROSCI.6292-10.2011, PMID: 21508221

Chambers CD, Garavan H, Bellgrove MA. Insights into the neural basis of response inhibition from cognitive and clinical neuroscience. Neurosci Biobehav Rev. 2009 May;33(5):631-46. doi: 10.1016/j.neubiorev.2008.08.016. Epub 2008 Sep 11. PMID: 18835296.

Chung JE, Magland JF, Barnett AH, Tolosa VM, Tooker AC, Lee KY, Shah KG, Felix SH, Frank LM, Greengard LF. A Fully Automated Approach to Spike Sorting. Neuron. 2017 Sep 13;95(6):1381-1394.e6. doi: 10.1016/j.neuron.2017.08.030. PMID: 28910621; PMCID: PMC5743236.

Churchland MM, Byron MY, Ryu SI, Santhanam G, Shenoy KV. Neural variability in premotor cortex provides a signature of motor preparation. Journal of Neuroscience. 2006 Apr 5;26(14):3697-712.

Churchland MM, Yu BM, Cunningham JP, Sugrue LP, Cohen MR, Corrado GS, Newsome WT, Clark AM, Hosseini P, Scott BB, Bradley DC, Smith MA, Kohn A, Movshon JA, Armstrong KM, Moore T, Chang SW, Snyder LH, Lisberger SG, Priebe NJ, Finn IM, Ferster D, Ryu SI, Santhanam G, Sahani M, Shenoy KV. Stimulus onset quenches neural variability: a widespread cortical phenomenon. Nat Neurosci. 2010 Mar;13(3):369-78. doi: 10.1038/nn.2501. Epub 2010 Feb 21. PMID: 20173745; PMCID: PMC2828350.

Claffey MP, Sheldon S, Stinear CM, Verbruggen F, Aron AR. Having a goal to stop action is associated with advance control of specific motor representations. Neuropsychologia. 2010 Jan;48(2):541-8. doi: 10.1016/j.neuropsychologia.2009.10.015. Epub 2009 Oct 29. PMID: 19879283; PMCID: PMC2813913.

Cohen MR, Maunsell JH. Attention improves performance primarily by reducing interneuronal correlations. Nat Neurosci. 2009 Dec;12(12):1594-600. doi: 10.1038/nn.2439. Epub 2009 Nov 15. PMID: 19915566; PMCID: PMC2820564.

Delaville C, Navailles S, Benazzouz A. Effects of noradrenaline and serotonin depletions on the neuronal activity of globus pallidus and substantia nigra pars reticulata in experimental parkinsonism. Neuroscience. 2012 Jan 27;202:424-33.

Dorval AD, Russo GS, Hashimoto T, Xu W, Grill WM, Vitek JL. Deep brain stimulation reduces neuronal entropy in the MPTP-primate model of Parkinson's disease. J Neurophysiol. 2008 Nov;100(5):2807-18. doi: 10.1152/jn.90763.2008. Epub 2008 Sep 10. PMID: 18784271; PMCID: PMC2585386.

Fan D, Rossi MA, Yin HH. Mechanisms of action selection and timing in substantia nigra neurons. J Neurosci. 2012 Apr 18;32(16):5534-48. doi: 10.1523/JNEUROSCI.5924-11.2012. PMID: 22514315; PMCID: PMC6703499.

Gu BM, Schmidt R, Berke JD. Globus pallidus dynamics reveal covert strategies for behavioral inhibition. Elife. 2020 Jun 10;9:e57215. doi: 10.7554/eLife.57215. PMID: 32519952; PMCID: PMC7314538.

Gulley JM, Kosobud AE, Rebec GV. Behavior-related modulation of substantia nigra pars reticulata neurons in rats performing a conditioned reinforcement task. Neuroscience. 2002;111(2):337-49. doi: 10.1016/s0306-4522(02)00018-0. PMID: 11983319.

Hikosaka O, Wurtz RH. Visual and oculomotor functions of monkey substantia nigra pars reticulata. III. Memory-contingent visual and saccade responses. Journal of Neurophysiology. 1983a; 49:1268–1284.

Hikosaka O, Wurtz RH. Visual and oculomotor functions of monkey substantia nigra pars reticulata. IV. Relation of substantia nigra to superior colliculus. J Neurophysiol. 1983b May;49(5):1285-301. doi: 10.1152/jn.1983.49.5.1285. PMID: 6306173.

Jahanshahi M, Obeso I, Rothwell JC, Obeso JA. 2015. A fronto-striato-subthalamic-pallidal network for goal-directed and habitual inhibition. Nature Reviews Neuroscience 16:719–732. DOI: https://doi.org/10.1038/nrn4038, PMID: 26530468

Joshua M, Adler A, Rosin B, Vaadia E, Bergman H. Encoding of probabilistic rewarding and aversive events by pallidal and nigral neurons. J Neurophysiol. 2009 Feb;101(2):758-72. doi: 10.1152/jn.90764.2008. Epub 2008 Dec 3. PMID: 19052110.

Lauwereyns J, Watanabe K, Coe B, Hikosaka O. A neural correlate of response bias in monkey caudate nucleus. Nature. 2002 Jul;418(6896):413-7.

Legéndy CR, Salcman M. Bursts and recurrences of bursts in the spike trains of spontaneously active striate cortex neurons. J Neurophysiol. 1985 Apr;53(4):926-39. doi: 10.1152/jn.1985.53.4.926. PMID: 3998798.

Leventhal DK, Gage GJ, Schmidt R, Pettibone JR, Case AC, Berke JD. Basal ganglia beta oscillations accompany cue utilization. Neuron. 2012 Feb 9;73(3):523-36. doi: 10.1016/j.neuron.2011.11.032. PMID: 22325204; PMCID: PMC3463873.

Lintz MJ, Felsen G. Basal ganglia output reflects internally-specified movements. Elife. 2016 Jul 5;5:e13833. doi: 10.7554/eLife.13833. PMID: 27377356; PMCID: PMC4970866.

Lobb C. Abnormal Bursting as a Pathophysiological Mechanism in Parkinson's Disease. Basal Ganglia. 2014 Apr 1;3(4):187-195. doi: 10.1016/j.baga.2013.11.002. PMID: 24729952; PMCID: PMC3979569.

Low KA, Miller J, Vierck E. Response slowing in Parkinson's disease: a psychophysiological analysis of premotor and motor processes. Brain. 2002 Sep;125(Pt 9):1980-94. doi: 10.1093/brain/awf206. PMID: 12183344.

Mailly P, Charpier S, Menetrey A, Deniau JM. Three-dimensional organization of the recurrent axon collateral network of the substantia nigra pars reticulata neurons in the rat. J Neurosci. 2003 Jun 15;23(12):5247-57. doi: 10.1523/JNEUROSCI.23-12-05247.2003. PMID: 12832549; PMCID: PMC6741183.

Mallet N, Schmidt R, Leventhal D, Chen F, Amer N, Boraud T, Berke JD. 2016. Arkypallidal cells send a stop signal to striatum. Neuron 89:308–316. DOI: https://doi.org/10.1016/j.neuron.2015.12.017, PMID: 26777273

Mazzoni P, Hristova A, Krakauer JW. Why don't we move faster? Parkinson's disease, movement vigor, and implicit motivation. J Neurosci. 2007 Jul 4;27(27):7105-16. doi: 10.1523/JNEUROSCI.0264-07.2007. PMID: 17611263; PMCID: PMC6794577.

Paxinos CW. 2006. The Rat Brain in Stereotaxic Coordinates. Academic Press. DOI: https://doi.org/10.1046/j.1469-7580.1997.191203153.x

Phillips RS, Rosner I, Gittis AH, Rubin JE. The effects of chloride dynamics on substantia nigra pars reticulata responses to pallidal and striatal inputs. Elife. 2020 Sep 7;9:e55592.

Pouget P, Logan GD, Palmeri TJ, Boucher L, Paré M, Schall JD. Neural basis of adaptive response time adjustment during saccade countermanding. J Neurosci. 2011 Aug 31;31(35):12604-12. doi: 10.1523/JNEUROSCI.1868-11.2011. PMID: 21880921; PMCID: PMC3173043.

Rubin JE, McIntyre CC, Turner RS, Wichmann T. Basal ganglia activity patterns in parkinsonism and computational modeling of their downstream effects. Eur J Neurosci. 2012 Jul;36(2):2213-28. doi: 10.1111/j.1460-9568.2012.08108.x. PMID: 22805066; PMCID: PMC3400124.

Sato M, Hikosaka O. Role of primate substantia nigra pars reticulata in reward-oriented saccadic eye movement. J Neurosci. 2002 Mar 15;22(6):2363-73. doi: 10.1523/JNEUROSCI.22-06-02363.2002. PMID: 11896175; PMCID: PMC6758246.

Schmidt R, Leventhal DK, Mallet N, Chen F, Berke JD. 2013. Canceling actions involves a race between basal ganglia pathways. Nature Neuroscience 16:1118–1124. DOI: https://doi.org/10.1038/nn.3456, PMID: 23852117

Schmidt R, Berke JD. 2017. A Pause-then-Cancel model of stopping: evidence from basal ganglia neurophysiology. Philosophical Transactions of the Royal Society B: Biological Sciences 372:20160202. DOI: https://doi.org/10.1098/rstb.2016.0202

Sharott A, Gulberti A, Zittel S, Tudor Jones AA, Fickel U, Münchau A, Köppen JA, Gerloff C, Westphal M, Buhmann C, Hamel W, Engel AK, Moll CK. Activity parameters of subthalamic nucleus neurons selectively predict motor symptom severity in Parkinson's disease. J Neurosci. 2014 Apr 30;34(18):6273-85. doi: 10.1523/JNEUROSCI.1803-13.2014. PMID: 24790198; PMCID: PMC4004813.

Simmons DV, Higgs MH, Lebby S, Wilson CJ. Indirect pathway control of firing rate and pattern in the substantia nigra pars reticulata. Journal of Neurophysiology. 2020 Feb 1;123(2):800-14.

Tai CH. Subthalamic burst firing: A pathophysiological target in Parkinson's disease. Neurosci Biobehav Rev. 2022 Jan;132:410-419. doi: 10.1016/j.neubiorev.2021.11.044. Epub 2021 Nov 29. PMID: 34856222.

Wager TD, Sylvester CY, Lacey SC, Nee DE, Franklin M, Jonides J. Common and unique components of response inhibition revealed by fMRI. Neuroimage. 2005 Aug 15;27(2):323-40. doi: 10.1016/j.neuroimage.2005.01.054. PMID: 16019232.

Waschke L, Kloosterman NA, Obleser J, Garrett DD. Behavior needs neural variability. Neuron. 2021 Mar 3;109(5):751-66.

Willard AM, Isett BR, Whalen TC, Mastro KJ, Ki CS, Mao X, Gittis AH. State transitions in the substantia nigra reticulata predict the onset of motor deficits in models of progressive dopamine depletion in mice. Elife. 2019 Mar 6;8:e42746. doi: 10.7554/eLife.42746. PMID: 30839276; PMCID: PMC6402832.

Wiltschko AB, Gage GJ, Berke JD. Wavelet filtering before spike detection preserves waveform shape and enhances single-unit discrimination. J Neurosci Methods. 2008 Aug 15;173(1):34-40. doi: 10.1016/j.jneumeth.2008.05.016. Epub 2008 May 28. PMID: 18597853; PMCID: PMC2602872.

Yasuda M, Hikosaka O. To wait or not to wait—separate mechanisms in the oculomotor circuit of basal ganglia. Frontiers in Neuroanatomy. 2017 Apr 11;11:35.

Zhou FM, Lee CR. Intrinsic and integrative properties of substantia nigra pars reticulata neurons. Neuroscience. 2011 Dec 15;198:69-94. doi: 10.1016/j.neuroscience.2011.07.061. Epub 2011 Aug 2. PMID: 21839148; PMCID: PMC3221915.

**Supplemental information**

**Supplementary Figure 1.** Estimated locations of recorded cells, within coronal atlas sections (Paxinos & Watson 2006).

**Supplementary Figure 2.** Ipsi>Contra type cells are further divided into increase (A) and decrease (B) type cells. Both group of cells show significantly increase firing rates before *Go!* cues. Same formats as in the figure 2.

**Supplementary Figure 3.** (A) PCA was performed using 500 ms epoch around Center Out for contra and ipsi movements (averaged, normalized firing rates were concatenated). (B) Variance explained by each of the first 10 PCs. (C) The first 10 principal components. (D) Population dynamics of different trial outcomes (200ms around Go! cue) show different neural trajectory patterns. Permutation test shows positioning differences at *Go!* cue compared to correct contra go trials. The trials with wrong choice are biased toward ipsi action initiation at the time of *Go!* cue. Formats of permutation histograms are same as in figure 3B.

**Supplementary Figure 4.** (A) Sessions with high CV (averaged across units) shows longer reaction times (mean reaction times of all trials). Each data point shows the value of each session. (B) Sessions with bigger increase of CV during proactive inhibition shows bigger proactive inhibition effects. (C) Sessions with proactive inhibition effect show bigger CV compared to the sessions without proactive inhibition effect in individual rats (one-tailed Wilcoxon signed rank test, $p<0.05$).

**Supplementary Figure 5**. Both Contra>Ipsi and Ipsi>Contra type of cells show increased CV and burst ratios with selective proactive inhibition, however only Contra>Ipsi cells shows a statistically significant effect (Wilcoxon signed rank test). Same format as in the figure 4C.

**Supplementary Figure 6.** Individual rat data shows firing rates (during 200ms before *Go!* cue) and CV (3s before *Go!* cue) differences between two conditions.

## Figure 1

# *Figure 2*

# *Figure 3*

## *Figure 4*

## Figure 5

**A**
Positioning at Go cue
Trajectory before Go cue
Optimal subspace
Long RT
Short RT

**B** Individual trials in a single session
3s Before Go cue
Contra
Ipsi
Center out
Go cue

**C**
Neuron 3
$t_2$
$d(t_2)$
Center of trajectory
$t_1$
$d(t_1)$
Neuron 1
Neuron 2

**D** Temporal variations of example trajectories
*Maybe-Stop-Contra example*
CV: 1.53  Temporal variation: 4.02
*No-Stop-example*
CV: 1.22  Temporal variation: 3.19
Go cue
3s before Go cue

**E**
*p = 5.6X10⁻⁰⁶*
Correlation over trials
[CV, temporal variations]

**F**
*p = 0.0004*
ΔTemporal variations of trajectory
[Maybe-Stop-Contra - No-Stop]

**G**
*p = 0.027*
ΔTrial variations before Go cue
[Maybe-Stop-Contra - No-Stop]

**H**
*p = 0.005*
Correlation over trials
[Contra Go RT, Euclidean distance]

# Figure 6

**A** **Maybe-Stop-Contra** compared to **No-Stop**

**Previous trial :** Success    Fail      Contra   Ipsi

Stop          Go