Title: Growth phase estimation for abundant bacterial populations sampled longitudinally from

human stool metagenomes.

Authors: Joe J. Lim[1], Christian Diener[2], Sean M. Gibbons[2-4,*]

Affiliations: [1] Department of Environmental & Occupational Health Sciences, University of
Washington, Seattle, WA 98105; [2] Institute for Systems Biology, Seattle, WA 98109; [3]
Department of Bioengineering, University of Washington, Seattle, WA 98105; [4] eScience
Institute, University of Washington, Seattle, WA 98105

[*] corresponding author: sgibbons@isbscience.org

ABSTRACT

Longitudinal sampling of the stool has yielded important insights into the ecological dynamics of
the human gut microbiome. However, due to practical limitations, the most densely sampled
time series from the human gut are collected at a frequency of about once per day, while the
population doubling times for gut commensals are on the order of minutes-to-hours. Despite
this, much of the prior work on human gut microbiome time series modeling has, implicitly or
explicitly, assumed that day-to-day fluctuations in taxon abundances are related to population
growth or death rates, which is likely not the case. Here, we propose an alternative model of the
human gut as a continuous flow ecosystem at a dynamical steady state, where population
dynamics occur internally and the bacterial population sizes measured in stool represent an
endpoint of these internal dynamics. We formalize this idea as stochastic logistic growth of a
population held at a constant dilution rate. We show how this model provides a path toward
estimating the growth phases of gut bacterial populations *in situ*. We assess our model
predictions against densely-sampled human stool metagenomic time series data. Consistent
with our model, donors with slower defecation rates tended to harbor a larger proportion of taxa
in later growth phases, while faster defecation rates were associated with more taxa in earlier
growth phases. We discuss how these growth phase estimates may be used to better inform
metabolic modeling in flow-through ecosystems, like animal guts or industrial bioreactors.

INTRODUCTION

The human gut microbiome is an anaerobic bioreactor, ecologically distinct to each individual, that transforms dietary and host substrates into bioactive molecules important to host health [1–3]. Disruptions to the ecological composition of the gut have been shown to mediate the progression of various complex diseases [4–8]. Furthermore, the ecological dynamics of the gut appear to be relevant to both health and disease states [9, 10]. However, the biological interpretation of densely-sampled adult human fecal microbiome time series is fraught.

Various dynamical models have been applied to gut microbial abundance data collected from adult human donors [11–15]. These models often assume, either explicitly or implicitly, that day-to-day changes in abundance are proportional to population growth and/or death [16]. However, the underlying data often do not match this assumption [11, 16–20]. The gut is a flow-through ecosystem and commensal gut bacteria must grow fast enough to avoid dilution-to-extinction. As such, gut bacterial doubling times tend to be fast, ranging from minutes-to-hours [21–23]. However, stool sampling frequency is usually limited to, at most, about once per day. Consequently, rapid internal population dynamics likely cannot be directly estimated from the day-to-day measurements obtained from stool [16].

In the absence of major perturbations, is it possible to extract meaningful information about commensal population dynamics from adult human gut microbiome time series, despite the fundamental limitations in sampling timescales mentioned above? One work around to inferring growth rates of bacterial populations *in situ* is to leverage metagenome-inferred replication rates [22]. Briefly, instantaneous replication rates can be estimated for abundant bacterial populations in metagenomic samples by taking advantage of the fact that fast-growing taxa show an asymmetry in reads mapping to different genomic loci, with higher read depth near the origin of replication and a lower depth near the terminus due to the initiation of multiple

replication forks [21–23]. However, even when replication rates and population abundances can

both be estimated from the same metagenomic samples, it is unclear how these measurements

are related to the *in situ* growth phase of a population (e.g. lag, log, or stationary phases). As

such, biological interpretations regarding population size and replication rate fluctuations in flow-

through ecosystems like the human gut, where internal dynamics are much faster than sampling

rates, remain challenging.

Early experiments by Jaques Monod [24] identified distinct growth phases for bacterial

populations in culture, which can be captured by the stochastic logistic growth equation (sLGE)

[25]. The sLGE has been shown to be a good fit for bacterial population growth *in vitro* and in

real-world, steady-state ecosystems [26–31]. We used the sLGE to study statistical

relationships between population sizes and growth rates across the various phases of growth

(i.e., lag, acceleration, log, deceleration, and stationary phases) to see if we could extract *in situ*

growth phase information from longitudinal data from a steady-state, flow-through ecosystem

sampled at a consistent frequency. Overall, the sLGE model yields statistical relationships that

can be leveraged to identify the *in situ* growth phase of a bacterial population periodically

sampled from a continuous-flow ecosystem, like the human gut.

To assess our model predictions, we calculated population abundance and

growth/replication rate trajectories from more than a dozen organisms in four densely sampled

human gut metagenomic time series [32]. On average, gut commensal growth rates and

population sizes were positively correlated within each of the stool donor time series, which

suggests that most abundant taxa in the gut are growing exponentially. We were able to identify

signatures of specific growth phases in 20-40% of the abundant bacterial populations in the guts

of these four individuals. We describe how our growth phase inference approach can serve to

inform more accurate mechanistic modeling of flow-through ecosystems (e.g., community-scale

metabolic models, which usually assume exponential growth), which could have broad

implications for the gut microbiome and host health [8, 33, 34], flow-through agricultural systems

[35, 36], climate change [35, 37, 38], and industrial bioreactor production processes [39, 40].

RESULTS

Framing the gut as an anaerobic flow-through bioreactor

The mammalian gut can be understood as an anaerobic bioreactor with a continuous input (i.e.,

dietary and host substrates) and output (i.e., stool) [41], and microbial taxa must grow fast

enough to avoid dilution to extinction (Fig. 1A). Stool sampling captures the endpoint of internal

gut bacterial population dynamics. For example, in our cartoon figure we see that Taxon 1 starts

growing higher up in the colon and is in stationary phase by the time a stool sample is collected,

while Taxon 3 starts growing lower in the colon and is still growing exponentially at the point of

stool sampling (Fig. 1A). Overall, the daily abundances of Taxa 1-3 represent the average (μ)

steady-state population size, plus or minus some amount of biological and technical noise, at

the time of stool sampling (Fig. 1A). To investigate improved methods for interpreting the

dynamics of human gut microbial time series, we downloaded shotgun metagenomic time series

data from the BIO-ML cohort (i.e., health-screened stool donors who provided fecal-transplant

material to the stool bank OpenBiome) [32]. The BIO-ML cohort contained 74 donors, 70 of

which had 1-3 time points collected [32]. To filter for dense longitudinal data, we selected a

subset of donors with more than 50 time points. Four donors (i.e. donors ae, am, an, and ao)

met this criterion, with 3-5 fecal samples per week for >50 days (Fig. 1B).

Characterizing the relationships between gut commensal population size and growth rate

using metagenomic time series data

We first investigated the statistical properties of day-to-day fluctuations in gut bacterial

population sizes, estimated from metagenomic time series. Specifically, we looked at the

associations between population abundance estimates (t_n) and the changes in abundance

estimates (i.e., deltas) between time points ($t_{n+1} - t_n$). Naïvely, if most bacterial populations in

the stool were growing exponentially, we would expect that population abundances and growth

rates would be positively correlated. However, prior work has indicated an overall negative

correlation between abundances and changes in abundances in stool 16S rRNA gene amplicon

sequencing data generated from densely sampled human stool time series [15]. Indeed, we

found that abundant bacterial populations in the stool of the four BIO-ML donors maintained

stable average abundances over time (μ), with day-to-day fluctuations above and below this

average, as pictured in the example of *Bacteroides cellulosilyticus* in donor am (Fig. 2A-B). This

kind of pattern fits a regression-to-the-mean model, which one would expect when randomly

sampling from a stationary distribution (Fig. 2B). This kind of regression-to-the-mean process

will give rise to a negative correlation between population abundances and changes in

abundance between time points, consistent with what has been observed previously [15]. We

observed that the deltas ($t_{n+1} - t_n$) for the same gut taxon (*Bacteroides uniformis*) measured

across each donor time series, when plotted against their respective normalized abundances

(t_n), showed the expected negative association (Fig. 2C). Furthermore, similar negative

associations were observed across all taxa analyzed, across all donors (Fig. 2D). Overall, these

results support our assertion that stool samples provide steady-state population abundance

estimates for gut commensal bacteria, which are representative of the endpoint of internal

dynamics.

Next, we looked at the statistical associations between calculated peak-to-trough ratios

(i.e., PTRs; a proxy for growth-rate) [22] of abundant bacterial populations from each

metagenomic sample and their respective metagenomic population abundance estimates. If the

deltas, presented above, were truly proportional to growth and/or death rates, we would expect

that the statistical relationships between deltas and population size would be similar to those

between PTRs and population size. However, unlike the regression-to-the-mean signature

identified for the deltas, we found variable statistical relationships between PTR and centered

log-ratio (CLR) transformed population abundances for the same taxon (*Bacteroides ovatus*)

across the four donors (Fig. 3A). Similarly, we saw a wide range of positive, negative, and null

associations between PTRs and CLR abundances across all measured taxa within each donor

(Fig. 3B). These results are inconsistent with a regression-to-the-mean signal, and suggest a

more complex relationship between growth rate and population size [42–44]. Finally, we

calculated temporally-averaged PTRs and population sizes for each abundant taxon within each

of the donors. Overall, there was a significantly positive (linear regression, *p-values* = 0.0318,

0.125, 0.155, 0.031 for donors ae, am, an, and ao, respectively; combined *p-value* using

Fisher's method = 0.005), albeit noisy, association between average PTR and average CLR

abundance across all four donors (Fig. 3C), indicating that taxa with higher average population

sizes tend to have higher average growth rates. This result is consistent with what we would

expect to observe in exponentially-growing populations.

Stochastic logistic growth equation provides insights into growth phases

In order to better understand and interpret the varying relationships we observe between PTRs

and CLR abundances, we turn to modeling. The basic properties of growth curves of microbial

taxa can be captured using the logistic growth equation (Fig. 4). This model is defined such that

the change in abundance for each taxon i (dx_i/dt) is captured by the current abundance at time t,

$x_i(t)$, multiplied by the maximal growth rate, r, and the carrying capacity (k) term $(1-x_i(t)/k)$ [45]. In

this model, population size over time shows a sigmoidal curve, with the abundance

asymptotically approaching k (Fig. 4A, top panel). The derivative of this curve with respect to

time yields the change in growth rate over time, which peaks during log-phase growth (Fig. 4A,

middle panel). The second derivative of abundance with respect to time, which is the

instantaneous change in growth with respect to time and is often referred to as the acceleration

rate, shows a peak during the acceleration phase and a trough during the deceleration phase

(Fig. 4A, bottom panel). Based on this second-derivative curve, we show the expected

relationships between growth rate and abundance as you move across the logistic growth

curve, along the time axis (Fig. 4B). These expected relationships provide a potential path

forward for inferring the *in situ* growth phase of a bacterial population sampled at a consistent

frequency from a flow-through ecosystem.

The logistic growth model is a deterministic equation. However, the abundances of

commensal bacterial populations in the gut fluctuate due to myriad factors including interspecies

competition, resource fluctuations, and stool residence time [46]. In order to approximate these

fluctuations in our modeling, we introduced a stochastic term to the logistic growth model (Fig.

5A). Herein, σ denotes the noise magnitude and $\omega(t)$ represents a white noise term [27]. Five

growth phases (i.e., lag, acceleration, log, deceleration, and stationary phases) were defined

using the half-maximum and half-minimum, respectively, of the second derivative LGE curve

(Fig. S1A). We grouped these phases into three major categories: lag-acceleration phase, log

phase, and deceleration-stationary phase. We simulated 100 iterations of the stochastic logistic

growth equation (sLGE) for each of a range of parameterizations (see Methods), which

recapitulated the expected statistical relationships between growth rates and abundances for

populations consistently sampled within our three major growth phase categories (Fig. 5A-C).

For example, the Pearson's R values between growth rates and abundances were significantly

positive in lag-acceleration phase and significantly negative in deceleration-stationary phase

(Fig. 5B). Log phase growth was more variable, but showed little-to-no significant association

between growth rates and abundances. These results were reproduced across a wide range of

parameter space and were robust to varying the noise term (Fig. S1B). Overall, the relationships

between growth rate and abundance across growth phases were highly consistent with our

expectations (Fig. 5C).

Inferring in situ growth phases for abundant gut commensal populations sampled in

metagenomic time series

Based on these sLGE results, we assigned putative *in situ* growth phases to abundant gut

bacterial populations from the four BIO-ML gut metagenomic time series. Specifically, we

suggest that significantly positive associations (linear regression, adjusted *p*-value < 0.05, with a

positive beta-coefficient) between PTRs and CLR abundances indicate early-phase exponential

growth (i.e. acceleration phase; we can likely exclude lag phase due to the fact that we would

be unlikely to detect taxa with very low levels of biomass), significantly negative associations

(linear regression, adjusted *p*-value < 0.05, with negative beta-coefficient) indicate deceleration

or stationary phases, and the absence of a significant association could indicate either log-

phase growth or a false negative (i.e., not powered enough to detect a positive or negative

association with the number of time points sampled). *Bacteroides cellulosilyticus*, *Bacteroides*

ovatus 1, and *Megaspaera eldenii* showed significantly positive PTR-abundance associations

within donor ae (Figs. 6A and S2). *Bacteroides ovatus* 1 and *Parabacteroides distasonis*

showed positive PTR-abundance associations, while *Alistipes finegoldii*, *Bacteroides uniformis*,

and *Bacteroides xylanisolvens* showed negative associations in donor am (Figs. 6A and S3).

Alistipes shahii, *Bacteroides intestinalis*, *Bacteroides thetaiotaomicron*, *Bacteroides uniformis*,

Bacteroides xylanisolvens, and *Odoribacter splanchnicus* showed significantly negative PTR-

abundance associations in donor an (Fig. 6A and S4). Finally, *Favonifractor plautii* showed a

positive PTR-abundance association and *Bacteroides fragilis*, *Bacteroides ovatus* 1,

Bacteroides uniformis, and *Bacteroides xylanisolvens* showed negative associations in donor ao

(Fig. 6A and S5).

We observed a slight difference in the number of significantly positive and negative PTR-

abundance associations between donors ae/am, and an/ao, with donors an and ao tending to

have a larger proportion of negative associations when compared to donors ae and am.

Interestingly, donors an and ao had a lower average defecation frequency (≤ 1 per day) than

donors ae and am (> 1 per day). Concordantly, based on our flow-through model of the gut

ecosystem (Fig. 1A), we would expect that bacterial populations would be pushed towards

earlier growth phases at faster flow rates (Fig. 6B). Over half of the taxa with PTR and

abundance time series data did not show significant associations (Fig. 6A and Fig. S2-5). This

suggests that either these taxa are in the log growth phase or we were not powered enough to

detect significant positive or negative associations for these taxa given the effect sizes and the

number of samples. We suggest that many of these taxa may well be in log phase, due to the

significant association observed between average PTRs and average CLR abundances across

donors (Fig. 3C). Overall, our approach provides a new path toward identifying the *in situ* growth

phase of microbial populations in flow-through ecosystems.

DISCUSSION

Many prior studies assumed, either implicitly or explicitly, that the growth and death rates of gut

bacterial populations were proportional to day-to-day changes in abundances, as measured

from human stool samples. However, we outline how this assumption is likely invalid due to the

fact that human gut bacterial population growth/death processes inside the intestinal tract are

known to be faster (minutes-to-hours) than our sampling timescales (days). In support of this

assertion, we show how the statistical relationships between changes in abundance ($t_{n+1} - t_n$)

and abundances (t_n), estimated from stool metagenomic time series, indicate a regression-to-

the-mean effect that one would expect when sampling from a stationary distribution (Figs. 1-2).

Thus, as prior work has indicated [15], bacterial taxa in the gut have stable average population

sizes, which likely represent steady-state endpoints of internal dynamics (Figs. 1-2). Despite the

fundamental mismatch between gut bacterial population dynamics and sampling timescales, we

attempt to identify statistical signatures within these daily-sampled human gut time series that

might provide accurate insights into *in situ* population dynamics.

While changes in abundance between time points (i.e., deltas) do not appear to be

related to population growth, PTRs enable direct estimates of *in situ* growth rates from

metagenomic samples [22, 23, 47–49]. Unlike the relationships between deltas and

abundances, which were always negative (Fig. 2C-D), the relationships between PTRs and

abundances were quite variable (Fig. 3A-3B). While regression-to-the-mean is a plausible

mechanism for the consistent negative delta-abundance relationships (Fig. 2), the underlying

processes driving variable PTR-abundance relationships appear to be more nuanced (Fig. 3).

We turned to the sLGE to explore relationships between growth rate and abundance

across different phases of growth (Fig. 4). The sLGE recapitulated the key relationships

observed in the metagenomic time series between PTRs and abundances and provided

predictions for *in situ* growth phases (Figs. 5-6). Consistent with our sLGE predictions, we found

that individuals with higher defecation rates tended to be enriched for taxa in earlier growth

phases (Fig. 6). Thus, our results reveal a promising approach to inferring *in situ* growth phases

for abundant organisms detected in human gut metagenomic time series. A major limitation of

this approach is our lack of knowledge about the effect size distributions for these relationships

in real-world data and an understanding of the statistical power needed for detecting these

associations from metagenomic time series. The absence of a PTR-abundance relationship

could represent a false negative or it could indicate log-phase growth. Future *in vitro*

experimental work (e.g., in chemostats) should focus on better quantifying these PTR-

abundance relationships across parameter space to build a more quantitative understanding of

these phenomena (e.g., through varying maximal growth rates, the carrying capacities, the flow

rates, and volumes).

We observed that the average PTR and average abundance of a given taxon over time

were positively correlated, which is consistent with exponentially-growing populations (Fig. 3C).

This result is especially relevant to the metabolic modeling community. Ecological interactions

within free-living and host-associated microbial communities are largely governed by exchanges

of small-molecule metabolites [50, 51]. Genome-scale metabolic modeling and flux-balance

analysis (FBA) has been effective mechanistic tools for simulating these metabolic exchanges,

especially in controlled bioreactor systems [52]. The objective function used to find a unique

solution to bacterial FBA models is often biomass maximization, which assumes that these

organisms are growing exponentially at steady state. Exponential growth is a valid assumption

for organisms in acceleration or log growth phases, but this assumption is violated for

organisms in deceleration or stationary phases. Prior work has demonstrated that biomass

composition can change depending on the growth phase of a population, which ideally would be

taken into account to more accurately model metabolic fluxes within the system [53–55].

Overall, our work suggests that most organisms in the human gut are amenable to FBA, and our

growth phase estimation approach allows for the identification of populations that may not fit

classical FBA assumptions.

In conclusion, we provide a new path forward for the biological interpretation of

metagenomic time series data generated from adult human stool samples. We hope that *in situ*

growth phase estimation will be applied more broadly to other kinds of flow-through

environments to improve our understanding of internal dynamics in these systems and provide

improved constraints for mechanistic modeling of microbial communities.

METHODS

Shotgun metagenomics data processing and analysis

Longitudinal shotgun metagenomics sequencing data from healthy human stool samples was

downloaded from NCBI BioProject accession PRJNA544527, and the associated metadata was

downloaded from the associated article [32]. Raw FASTQ files were filtered and trimmed using

FASTP [56], removing the first 5 nucleotides of the read 5' end to avoid leftover primer and

adapter sequencing not removed during demultiplexing and an adaptive sliding window filter on

the 3' end of the read with a required minimum quality score of 20. Reads containing ambiguous

base calls, having a mean quality score less than 20, or with a length smaller than 50nt after

trimming were removed from the analysis. Taxonomic assignment on the read level was

performed with Kraken2 using the Kraken2 default database [57]. Abundances on the kingdom,

phylum, genus, and species ranks were then obtained using Bracken [58]. Trimmed and filtered

reads were then aligned to 2,935 representative bacterial reference genomes taken from the

IGG database (version 1.01) using Bowtie2 [59, 60]. Coverage profiles and log2 estimates of

peak-to-trough ratios were estimated using COPTR v1.1.2 on the species level within each

sample [61]. PTR estimates were then merged with Bracken abundance estimates, retaining

only those species identified by both methods (Kraken2 and Bowtie2 alignment to IGGdb).

The processed data containing the raw reads and $\log_2$ peak-to-trough ratios ($\log_2$PTRs)

were read into R version 4.1.3 for analysis ([62]). All plots were generated using ggplot2 [63],

unless indicated otherwise. Donor time series were selected by only retaining individuals with

over 50 metagenomic time points, resulting in four time series (i.e., donors ae, am, an, and ao).

Distinct *Bacteroides ovatus* strains across all four donors contained duplicated taxon names

with unique taxonomic identifiers, and were renamed to "*Bacteroides ovatus_1*" and

"*Bacteroides ovatus_2.*" Raw read counts for a given taxon within a sample were centered log-

ratio (CLR) transformed [64]. Taxa that had matched $\log_2$PTR information available across more

than 5 time points within an individual, with time differences between samples less than three

days, were used in subsequent analyses. Changes in normalized abundance were calculated

as $Abundance\ changes(delta) = x(t+1) - x(t)$, where $\Delta t < 3\ days$. To assess the regression-

to-the-mean effect, CLR-normalized abundances were plotted against deltas for each taxon,

and the regression coefficients, aggregating all microbial taxa, were plotted as boxplots

(showing median and interquartile range), summarized by donor.

For each donor, to estimate the growth phase of each individual taxon, we used linear

regression of CLR-normalized abundances vs. $\log_2$PTRs, followed by a Benjamini-Hochberg p-

value correction to control for the false discovery rate (FDR) in base R. FDR-adjusted p-values

< 0.05 were considered significant. Taxa with significantly positive or negative associations

were considered to be in lag-acceleration or deceleration-stationary phase, respectively. Those

with no correlation were not assigned a growth phase, as this result could either be a false

negative or indicative of log-phase growth. Linear regression was also used to test whether or

not average CLR-normalized abundances and average $\log_2$PTRs were significantly associated

within each donor, and p-values from individual tests were combined using Fisher's method

[65].

Stochastic logistic growth model simulation

The stochastic logistic growth equation (SLE) was implemented as: $\frac{dx_i}{dt} = rx_i(t)\left(1 - \frac{x_i(t)}{K}\right) +$

$\sigma x_i(t)\omega(t)$, where *t* is time, *r* is the growth rate, x_i is the abundance of taxon *i*, *K* is the carrying

capacity, σ is the noise magnitude term, and $\omega(t)$ is the noise distribution term. Using the R

package sde [66], taxonomic growth was simulated with $x_{i,0} = 1$, $t_0 = 1$ to $t_{final} = 100$, for 100

iterations. The other parameters were varied as described in the results and below. To

investigate the impact of noise on sLGE trajectories, noise levels were set from 0.001 to 1, with

r and *K* ranging from 1 to 3 and 10 to 1000, respectively. To investigate the statistical

relationships between deltas and abundances across growth phases and across model

parameterizations, Pearson's R coefficients and p-values were calculated for each of the three

growth phase categories. The growth phases for each model parameterization were defined

using the non-stochastic logistic growth equation (LGE): $\frac{dx_i}{dt} = rx_i(t)\left(1 - \frac{x_i(t)}{K}\right)$, the solution for

which can be written as $x_i = \frac{x_{i,0}Ke^{rt}}{(K-x_{i,0})+x_{i,0}e^{rt}}$.

The x_i values for each simulated time point from solving the LGE were used to calculate

the first derivative (i.e., the growth rate), which is exactly equal to the LGE. The second

derivative (i.e., growth acceleration), $\frac{d^2x_i}{dt^2} = K^2 x_i \left(1 - \frac{x_i}{K}\right)\left(1 - \left(\frac{2x_i}{K}\right)\right)$, was calculated using

solved x_i values. Growth phases from the SLM were defined using the second derivative curves.

First, the intersections of the acceleration curve and the half-max, a_1 and a_2, and the half-

min, a_3 and a_4, were calculated (Fig. S1). The corresponding simulated time points of a_j,

denoted as s_j, where *j* = 1 - 4, were then used to define growth phases as follows: lag

phase: $t < s_1$; acceleration phase: $s_1 < t < s_2$; log phase: $s_2 < t < s_3$; deceleration phase: $s_3 <$

$t < s_4$; and stationary phase: $t > s_4$. Here, lag and acceleration, and deceleration and stationary

phases were combined, as these phases display similar delta-abundance relationships along

the logistic growth curve. Conceptual diagrams were created using BioRender.

Data and code availability

Nextflow pipelines implementing the processing of metagenomic shotgun sequencing data from

raw reads to taxonomic abundance matrices and PTR estimates can be found at

https://github.com/Gibbons-Lab/pipelines/ (metagenomics pipelines). R scripts and code used to analyze the data, run the sLGE simulations, and produce the figures in the manuscript have been deposited at https://github.com/Gibbons-Lab/human-microbiome-time-series-growth-phase-estimation.

Acknowledgements

We would like to thank Shijie Zhao for suggesting that we investigate PTR-abundance relationships in the BIO-ML data set. We would also like to thank Nitin Baliga, Amy Willis, Julia Cui, and the members of the Gibbons Lab for helpful discussions of this work. SMG and CD were supported by a Washington Research Foundation Distinguished Investigator Award and by startup funds from the Institute for Systems Biology. JL was supported by the Environmental Pathology/Toxicology training grant (ES007032).

References

1. El Aidy S, Hooiveld G, Tremaroli V, Bäckhed F, Kleerebezem M. The gut microbiota and mucosal homeostasis: colonized at birth or at adulthood, does it matter? *Gut Microbes* 2013; **4**: 118–124.

2. Martin AM, Sun EW, Rogers GB, Keating DJ. The Influence of the Gut Microbiome on Host Metabolism Through the Regulation of Gut Hormone Release. *Front Physiol* 2019; **10**: 428.

3. Zhang P, Meng X, Li D, Calderone R, Mao D, Sui B. Commensal Homeostasis of Gut Microbiota-Host for the Impact of Obesity. *Front Physiol* 2017; **8**: 1122.

4. Stefan KL, Kim MV, Iwasaki A, Kasper DL. Commensal Microbiota Modulation of Natural Resistance to Virus Infection. *Cell* 2020; **183**: 1312–1324.e10.

5. Fischbach MA. Microbiome: Focus on Causation and Mechanism. *Cell* 2018; **174**: 785–790.

6.   Zmora N, Zilberman-Schapira G, Suez J, Mor U, Dori-Bachash M, Bashiardes S, et al.

Personalized Gut Mucosal Colonization Resistance to Empiric Probiotics Is Associated with

Unique Host and Microbiome Features. *Cell* 2018; **174**: 1388–1405.e21.

7.   Suez J, Zmora N, Zilberman-Schapira G, Mor U, Dori-Bachash M, Bashiardes S, et al.

Post-Antibiotic Gut Mucosal Microbiome Reconstitution Is Impaired by Probiotics and

Improved by Autologous FMT. *Cell* 2018; **174**: 1406–1423.e16.

8.   Bullman S, Pedamallu CS, Sicinska E, Clancy TE, Zhang X, Cai D, et al. Analysis of

Fusobacterium persistence and antibiotic response in colorectal cancer. *Science* 2017;

358: 1443–1448.

9.   Halfvarson J, Brislawn CJ, Lamendella R, Vázquez-Baeza Y, Walters WA, Bramer LM, et

al. Dynamics of the human gut microbiome in inflammatory bowel disease. *Nat Microbiol*

2017; **2**: 17004.

10. Cuna A, Morowitz MJ, Ahmed I, Umar S, Sampath V. Dynamics of the preterm gut

microbiome in health and disease. *Am J Physiol Gastrointest Liver Physiol* 2021; **320**:

G411–G419.

11. Fisher CK, Mehta P. Identifying keystone species in the human gut microbiome from

metagenomic timeseries using sparse linear regression. *PLoS One* 2014; **9**: e102451.

12. Bashan A, Gibson TE, Friedman J, Carey VJ, Weiss ST, Hohmann EL, et al. Universality of

human microbial dynamics. *Nature* 2016; **534**: 259–262.

13. Stein RR, Bucci V, Toussaint NC, Buffie CG, Rätsch G, Pamer EG, et al. Ecological

modeling from time-series inference: insight into dynamics and stability of intestinal

microbiota. *PLoS Comput Biol* 2013; **9**: e1003388.

14. Faust K, Lahti L, Gonze D, de Vos WM, Raes J. Metagenomics meets time series analysis:

unraveling microbial community dynamics. *Curr Opin Microbiol* 2015; **25**: 56–66.

15. Gibbons SM, Kearney SM, Smillie CS, Alm EJ. Two dynamic regimes in the human gut

microbiome. *PLoS Comput Biol* 2017; **13**: e1005364.

16. Carr A, Diener C, Baliga NS, Gibbons SM. Use and abuse of correlation analyses in

microbial ecology. *ISME J* 2019; **13**: 2647–2655.

17. Momeni B, Xie L, Shou W. Lotka-Volterra pairwise modeling fails to capture diverse

pairwise microbial interactions. *Elife* 2017; **6**.

18. Berry D, Widder S. Deciphering microbial interactions and detecting keystone species with

co-occurrence networks. *Front Microbiol* 2014; **5**: 219.

19. Freilich MA, Wieters E, Broitman BR, Marquet PA, Navarrete SA. Species co-occurrence

networks: Can they reveal trophic and non-trophic interactions in ecological communities?

Ecology 2018; **99**: 690–699.

20. Coenen AR, Weitz JS. Limitations of Correlation-Based Inference in Complex Virus-

Microbe Communities. *mSystems* 2018; **3**.

21. Korem T, Zeevi D, Suez J, Weinberger A, Avnit-Sagi T, Pompan-Lotan M, et al. Growth

dynamics of gut microbiota in health and disease inferred from single metagenomic

samples. *Science* 2015; **349**: 1101–1106.

22. Brown CT, Olm MR, Thomas BC, Banfield JF. Measurement of bacterial replication rates in

microbial communities. *Nat Biotechnol* 2016; **34**: 1256–1263.

23. Gibson B, Wilson DJ, Feil E, Eyre-Walker A. The distribution of bacterial doubling times in

the wild. *Proc Biol Sci* 2018; **285**.

24. Monod J. THE GROWTH OF BACTERIAL CULTURES. *Annu Rev Microbiol* 1949; **3**: 371–

394.

25. Logistic stochastic growth models and applications. *Handbook of the Logistic Distribution*.

1991. CRC Press, pp 419–419.

26. Descheemaeker L, de Buyl S. Stochastic logistic models reproduce experimental time

series of microbial communities. *Elife* 2020; **9**.

27. Grilli J. Macroecological laws describe variation and diversity in microbial communities. *Nat

Commun* 2020; **11**: 4743.

28. Lobry JR, Flandrois JP, Carret G, Pave A. Monod's bacterial growth model revisited. *Bull*
Math Biol 1992; **54**: 117–122.

29. Fujikawa H, Kai A, Morozumi S. A new logistic model for bacterial growth. *Shokuhin*
Eiseigaku Zasshi 2003; **44**: 155–160.

30. Baranyi J, McClure PJ, Sutherland JP, Roberts TA. Modeling bacterial growth responses. *J*
Ind Microbiol 1993; **12**: 190–194.

31. Ho P-Y, Good BH, Huang KC. Competition for fluctuating resources reproduces statistics of
species abundance over time across wide-ranging microbiotas. *Elife* 2022; **11**.

32. Poyet M, Groussin M, Gibbons SM, Avila-Pacheco J, Jiang X, Kearney SM, et al. A library
of human gut bacterial isolates paired with longitudinal multiomics data enables
mechanistic microbiome research. *Nat Med* 2019; **25**: 1442–1452.

33. Kadosh E, Snir-Alkalay I, Venkatachalam A, May S, Lasry A, Elyada E, et al. The gut
microbiome switches mutant p53 from tumour-suppressive to oncogenic. *Nature* 2020; **586**:
133–138.

34. Kolodziejczyk AA, Zheng D, Shibolet O, Elinav E. The role of the microbiome in NAFLD and
NASH. *EMBO Mol Med* 2019; **11**.

35. Day JA, Diener C, Otwell AE, Tams KE, Bebout B, Detweiler AM, et al. Lettuce (Lactuca
sativa) productivity influenced by microbial inocula under nitrogen-limited conditions in
aquaponics. *PLoS One* 2021; **16**: e0247534.

36. Levy R, Borenstein E. Metabolic modeling of species interaction in the human microbiome
elucidates community-level assembly rules. *Proc Natl Acad Sci U S A* 2013; **110**: 12804–
12809.

37. Otwell AE, Carr AV, Majumder ELW, Ruiz MK, Wilpiszeski RL, Hoang LT, et al. Sulfur
Metabolites Play Key System-Level Roles in Modulating Denitrification. *mSystems* 2021; **6**.

38. Wallace RJ, Snelling TJ, McCartney CA, Tapio I, Strozzi F. Application of meta-omics
techniques to understand greenhouse gas emissions originating from ruminal metabolism.

Genet Sel Evol 2017; **49**: 9.

39. Roy P, Dutta A, Deen B. Greenhouse gas emissions and production cost of ethanol

produced from biosyngas fermentation process. *Bioresour Technol* 2015; **192**: 185–191.

40. Zhong C. Industrial-Scale Production and Applications of Bacterial Cellulose. *Front Bioeng*

Biotechnol 2020; **8**: 605374.

41. Guzman-Rodriguez M, McDonald JAK, Hyde R, Allen-Vercoe E, Claud EC, Sheth PM, et

al. Using bioreactors to study the effects of drugs on the human microbiota. *Methods* 2018;

149: 31–41.

42. Blätke M-A, Bräutigam A. Evolution of C4 photosynthesis predicted by constraint-based

modelling. *Elife* 2019; **8**.

43. Atolia E, Cesar S, Arjes HA, Rajendram M, Shi H, Knapp BD, et al. Environmental and

Physiological Factors Affecting High-Throughput Measurements of Bacterial Growth. *MBio*

2020; **11**.

44. Nguyen J, Lara-Gutiérrez J, Stocker R. Environmental fluctuations and their effects on

microbial communities, populations and individuals. *FEMS Microbiol Rev* 2021; **45**.

45. Tsoularis A, Wallace J. Analysis of logistic growth models. *Math Biosci* 2002; **179**: 21–55.

46. O'Sullivan O, Cronin O, Clarke SF, Murphy EF, Molloy MG, Shanahan F, et al. Exercise

and the microbiota. *Gut Microbes* 2015; **6**: 131–136.

47. Long AM, Hou S, Ignacio-Espinoza JC, Fuhrman JA. Benchmarking microbial growth rate

predictions from metagenomes. *ISME J* 2021; **15**: 183–195.

48. Szafrańska AK, Junker V, Steglich M, Nübel U. Rapid cell division of Staphylococcus

aureus during colonization of the human nose. *BMC Genomics* 2019; **20**: 229.

49. Gao Y, Li H. Quantifying and comparing bacterial growth dynamics in multiple

metagenomic samples. *Nat Methods* 2018; **15**: 1041–1044.

50. Braga RM, Dourado MN, Araújo WL. Microbial interactions: ecology in a molecular

perspective. *Braz J Microbiol* 2016; **47 Suppl 1**: 86–98.

51.   Zelezniak A, Andrejev S, Ponomarova O, Mende DR, Bork P, Patil KR. Metabolic

478   dependencies drive species co-occurrence in diverse microbial communities. *Proc Natl*

479   *Acad Sci U S A* 2015; **112**: 6449–6454.

480   52.   Orth JD, Thiele I, Palsson BØ. What is flux balance analysis? *Nat Biotechnol* 2010; **28**:

481   245–248.

482   53.   Pramanik J, Keasling JD. Stoichiometric model of Escherichia coli metabolism:

483   incorporation of growth-rate dependent biomass composition and mechanistic energy

484   requirements. *Biotechnol Bioeng* 1997; **56**: 398–421.

485   54.   Dikicioglu D, Kırdar B, Oliver SG. Biomass composition: the 'elephant in the room' of

486   metabolic modelling. *Metabolomics* 2015; **11**: 1690–1701.

487   55.   Schulz C, Kumelj T, Karlsen E, Almaas E. Genome-scale metabolic modelling when

488   changes in environmental conditions affect biomass composition. *PLoS Comput Biol* 2021;

489   **17**: e1008528.

490   56.   Chen S, Zhou Y, Chen Y, Gu J. fastp: an ultra-fast all-in-one FASTQ preprocessor.

491   *Bioinformatics* 2018; **34**: i884–i890.

492   57.   Wood DE, Lu J, Langmead B. Improved metagenomic analysis with Kraken 2. *Genome Biol*

493   2019; **20**: 257.

494   58.   Lu J, Breitwieser FP, Thielen P, Salzberg SL. Bracken: estimating species abundance in

495   metagenomics data. *PeerJ Comput Sci* 2017; **3**: e104.

496   59.   Nayfach S, Shi ZJ, Seshadri R, Pollard KS, Kyrpides NC. New insights from uncultivated

497   genomes of the global human gut microbiome. *Nature* 2019; **568**: 505–510.

498   60.   Langmead B, Wilks C, Antonescu V, Charles R. Scaling read aligners to hundreds of

499   threads on general-purpose processors. *Bioinformatics* 2019; **35**: 421–432.

500   61.   Joseph TA, Chlenski P, Litman A, Korem T, Pe'er I. Accurate and robust inference of

501   microbial growth dynamics from metagenomic sequencing reveals personalized growth

502   rates. *Genome Res* 2022; **32**: 558–568.

62. Team RC. RA language and environment for statistical computing. R Foundation for

Statistical Computing Vienna Austria URL (2022).

63. Wickham H. ggplot2: Elegant Graphics for Data Analysis. 2016. Springer.

64. Lin H, Peddada SD. Analysis of microbial compositions: a review of normalization and

differential abundance analysis. *NPJ Biofilms Microbiomes* 2020; **6**: 60.

65. Fisher SRA. Statistical Methods for Research Workers. 1925. Oliver and Boyd.

66. Iacus SM. SDE : simulation and inference for stochastic differential equations. 2007.

Figure 1. Conceptual figure showing two flow-through microbial ecosystems: a
bioreactor and a human gut. **A.** Both bioreactors and guts are continuous flow-through
systems. Prior to reaching the measured abundances in stool, taxa grow in the large intestine
with varying growth rates, carrying capacities, and steady-state population sizes, which may be
in different growth phases at the time of measurement. For example, see dynamics for Taxa 1-
3. Daily stool collections show variation in abundances, but this variation likely does not reflect
internal growth dynamics in the gut. **B.** Healthy BIO-ML stool donors (subject IDs: ae, am, an,
and ao) with samples collected 3-5 days per week for a total of >50 time points. Red indicates
presence of shotgun metagenomic sequencing data and gray represents absence of
metagenomic data from consecutive daily time points.

Figure 2. Regression-to-the-mean effect in human microbial time series data. A. Yellow
line represents the mean abundance (μ) of *Bacteroides cellulosilyticus* over time in donor am.
Time points t_1 and t_3 indicate fluctuations below and above the mean abundance, and t_2 and t_4
show the return to the mean abundance. **B.** Distribution of time series delta values (e.g., t_2-t_1)
for *Bacteroides cellulosilyticus* in donor am, which is approximately normally distributed. **C.**
Deltas vs. abundances for *Bacteroides uniformis* time series from donors ae, am, an, and ao.
D. Boxplots (showing median and interquartile range) of linear regression coefficients for deltas
vs. abundances across all taxa time series in all four donors. Red line indicates a regression
coefficient of 0.

Figure 3. Variable relationships between PTRs and CLR-normalized abundances across human gut microbial time series. A. Log$_2$(PTR) and CLR-normalized abundance relationships for donors ae, am, an, and ao. Orange and blue lines show significantly positive and negative linear regression coefficients (linear regression, FDR adjusted *p*-value < 0.05), respectively. Gray lines indicate no statistically significant association. **B.** Boxplots (showing median and interquartile range) of linear regression coefficient combined for all filtered taxa for each donor. **C.** Mean log$_2$(PTR) and mean CLR-normalized abundance for all abundant taxa in each donor (*p-values* for regressions run within each donor were combined using Fisher's method; combined *p-value* = 0.005).

A Logistic growth of microbiota

$$\frac{dx_i}{dt} = r\,x_i(t)\left(1 - \frac{x_i(t)}{k}\right)$$

x

dx/dt

d²x/dt²

Time

B

Lag-Acceleration

Log

Deceleration-stationary

dx/dt

dx/dt

dx/dt

x

x

x

560   **Figure 4. Logistic growth model. A.** The logistic growth curve models abundance (x) with
561   respect to time (top panel). The first derivative of the logistic growth curve models the growth
562   rate with respect to time (middle panel). The second derivative of the logistic growth curve
563   models growth rate acceleration with respect to time (bottom panel). **B.** Expected relationships
564   between abundance and growth rate at different locations along the logistic growth curve.

Figure 5. Distinguishing growth phases using the stochastic logistic growth model. A.
Stochastic logistic growth curves with growth rate (r) = 1.2, carrying capacity (k) = 100, and
noise level (n) = 0.1 across 100 iterations. Major growth phase groups in orange (lag-
acceleration), gray (log), and blue (deceleration-stationary). **B.** Pearson's R values between
abundances and growth rates in each of our three growth phase windows across variable model
parameterizations (r = 1-3, k = 10-1000) and a fixed noise level ($\square$ = 0.1). **C.** Scatter plots
showing relationships between abundances and deltas across the three growth phase regions
defined in panel A.

Figure 6. *in vivo* growth phase estimation. A. We find variable relationships between PTRs and population abundances across taxa in each of the four donors, consistent with the growth phase patterns observed in sLGE simulations. Donors with higher defecation rates tended to have a larger fraction of taxa with positive PTR-abundance associations and fewer with negative associations, indicating lag-acceleration and deceleration-stationary phases, respectively. **B.** We suggest that higher defecation rates (i.e., higher dilution rates) push bacterial populations towards earlier growth phases, which is consistent with our results in panel A. **C.** Growth phase estimates can be leveraged to identify taxa that are more-or-less amenable to metabolic modeling techniques, like Flux Balance Analysis, which assumes exponential growth.

Figure S1. Definition of major growth phases using the stochastic logistic growth model.
A. The half-maximum of the peak and half-minimum of the trough of the second derivative of
abundance were used to define growth phases across model parameterizations. **B.** Pearson's R
values between abundances and growth rates in the three growth phase categories obtained
from combined sLGE simulation results across a range of growth rates (r = 1-3), carrying
capacities (k = 10-1000), and noise levels (n = 0.001-1).

599
Figure S2. Relationships between abundance and log2(PTR) for abundant taxa in donor ae. Abundant taxa with relatively dense longitudinal PTR and abundance data (at least 5 matched data points; time differences between adjacent samples less than three days) were selected for analysis. Gray trend lines show no significant correlations, orange trend lines indicate significant positive correlations, and blue trend lines represent significant negative correlations (linear regression, BH-FDR < 0.05).

Figure S3. Relationships between abundance and log2(PTR) for abundant taxa in donor am. Abundant taxa with relatively dense longitudinal PTR and abundance data (at least 5 matched data points; time differences between adjacent samples less than three days) were selected for analysis. Gray trend lines show no significant correlations, orange trend lines indicate significant positive correlations, and blue trend lines represent significant negative correlations (linear regression, BH-FDR < 0.05).

613
Figure S4. Relationships between abundance and log2(PTR) for abundant taxa in donor an. Abundant taxa with relatively dense longitudinal PTR and abundance data (at least 5 matched data points; time differences between adjacent samples less than three days) were selected for analysis. Gray trend lines show no significant correlations, orange trend lines indicate significant positive correlations, and blue trend lines represent significant negative correlations (linear regression, BH-FDR < 0.05).

Figure S5. Relationships between abundance and log2(PTR) for individual taxon in donor ao. Abundant taxa with relatively dense longitudinal PTR and abundance data (at least 5 matched data points; time differences between adjacent samples less than three days) were selected for analysis. Gray trend lines show no significant correlations, orange trend lines indicate significant positive correlations, and blue trend lines represent significant negative correlations (linear regression, BH-FDR < 0.05).