1 **Three-Dimensional Mitochondria Reconstructions of Murine Cardiac Muscle Changes in**
2 **Size Across Aging**
3
Zer Vue[1*], Kit Neikirk[1*], Larry Vang[1*], Edgar Garza-Lopez[2], Trace A. Christensen[3], Jianqiang
Shao[4], Jacob Lam[2], Heather K. Beasley[1], Andrea G. Marshall[1], Amber Crabtree[1], Josephs B Jr.[1],
Benjamin Rodriguez[1], Benjamin Kirk[2], Serif Bacevac[2], Taylor Barongan[1], Bryanna Shao[1],
Dominique C. Stephens[1,5], Kinuthia Kabugi[1], Ho-Jin Koh[6], Alice Koh[1], Chantell S. Evans[7],
Brittany Taylor[8], Anilkumar K. Reddy[9], Tyne Miller-Flemming[10], Ky'Era V. Actkins[10], Elma
Zaganjor[1], Nastaran Daneshgar[11], Sandra A. Murray[12], Bret C. Mobley[13], Steven Damo[5], Jennifer
A. Gaddy[13,14,15], Blake Riggs[16], Celestine Wanjalla[17], Annet Kirabo[17], Melanie McReynolds[18],
Jose A. Gomez[19], Mark A. Phillips[20], Vernat Exil[21,22#], Dao-Fu Dai[23#], Antentor Hinton, Jr[1#]
12
13
14
Affiliations:
1 Department of Molecular Physiology and Biophysics, Vanderbilt University, Nashville, TN, 37232, USA
2 Department of Internal Medicine, University of Iowa, Iowa City, IA, 52242, USA
3 Microscopy and Cell Analysis Core Facility, Mayo Clinic, Rochester, MN, 55905, USA
4 Central Microscopy Research Facility, University of Iowa, Iowa City, IA, 52242, USA
5 Department of Life and Physical Sciences, Fisk University, Nashville, TN, 37208, USA
6 Department of Biological Sciences, Tennessee State University, Nashville, TN 37209, USA
7 Department of Cell Biology, Duke University School of Medicine, Durham, NC, 27709, USA
8 J. Crayton Pruitt Family Department of Biomedical Engineering, University of Florida, Gainesville, FL, 32612, USA
9 Department of Medicine, Baylor College of Medicine, Houston, TX, 77002, USA
10 Division of Genetic Medicine, Department of Medicine, Vanderbilt University Medical Center, Nashville, TN, USA.
11 Department of Pathology, Carver College of Medicine, University of Iowa, Iowa City, IA, 52243, USA
12 Department of Cell Biology, School of Medicine, University of Pittsburgh, Pittsburgh, PA, 15261, USA
13 Department of Pathology, Microbiology and Immunology, Vanderbilt University Medical Center, Nashville, TN, 37233 USA
14 Department of Medicine, Vanderbilt University Medical Center, Nashville, TN 37232, USA
15 Tennessee Valley Healthcare Systems, U.S. Department of Veterans Affairs, Nashville, TN 37212, USA
16 Department of Biology at San Francisco State University, San Francisco, CA 94133, USA.
17 Department of Medicine / Clinical Pharmacology Division. Vanderbilt University Medical Center, Nashville, TN 37232
18 Department of Biochemistry and Molecular Biology, Pennsylvania State University, State College, PA 16801
19 Department of Medicine, Vanderbilt University Medical Center, Nashville, TN, 37233, USA
20 Department of Integrative Biology, Oregon State University, Corvallis, OR, 97332, USA
21 Department of Pediatrics, Div. of Cardiology, St. Louis University School of Medicine, St. Louis, MO, 63105, USA
22 Department of Pediatrics, Carver College of Medicine, University of Iowa, Iowa City, IA 52243, USA
23 Department of Pathology, University of John Hopkins, Baltimore, MD, USA.
39
40
Corresponding Author:
Antentor O. Hinton, Jr
Department of Molecular Physiology and Biophysics
Vanderbilt School of Medicine Basic Sciences
The Vanderbilt Diabetes Research and Training Center
750 Robinson Research Building

2200 Pierce Ave
Nashville, TN 37232-0615
319-383-3095 (cell)
antentor.o.hinton.jr@Vanderbilt.Edu

51
52
53

ABSTRACT: With sparse treatment options, cardiac disease remains a significant cause of
death among humans. As a person ages, mitochondria break down and the heart becomes less
efficient. Heart failure is linked to many mitochondria-associated processes, including
endoplasmic reticulum stress, mitochondrial bioenergetics, insulin signaling, autophagy, and
oxidative stress. The roles of key mitochondrial complexes that dictate the ultrastructure, such as
the mitochondrial contact site and cristae organizing system (MICOS), in aging cardiac muscle
are poorly understood. To better understand the cause of age-related alteration in mitochondrial
structure in cardiac muscle, we used transmission electron microscopy (TEM) and serial block
facing-scanning electron microscopy (SBF-SEM) to quantitatively analyze the 3D networks in
cardiac muscle samples of male mice at aging intervals of 3 months, 1 year, and 2 years. Here,
we present the loss of cristae morphology, the inner folds of the mitochondria, across age. In
conjunction with this, the 3D volume of mitochondria decreased. These findings mimicked
observed phenotypes in murine cardiac fibroblasts with CRISPR/Cas9 knockout of *Mitofilin,*
Chchd3, Chchd6 (some members of the MICOS complex), and *Opa1*, which showed poorer
oxidative consumption rate and mitochondria with decreased mitochondrial length and volume.
In combination, these data show the need to explore if loss of the MICOS complex in the heart
may be involved in age-associated mitochondrial and cristae structural changes.

71
72

73   **Keywords**: MICOS, aging, mitochondria, 3D morphometry; mitochondrial disease;
74   mitochondrion; reconstruction; reticulum; serial block-face SEM; cardiac muscle.

75
76
INTRODUCTION:
The mitochondrion is an organelle that critically carries out oxidative phosphorylation in
the cell and is implicated in many diseases, including Alzheimer's disease, diabetes, and heart
failure (1–3). Given the dependence on mitochondria to meet energetic demands, mitochondrial
dysfunction and deterioration are associated with cell death, including autophagy and apoptosis
(4–6). In addition to producing adenosine triphosphate (ATP), mitochondria also regulate cell
signaling during various cellular processes, including calcium homeostasis, apoptosis, and
immunity (5, 7–9). Interestingly, research also suggests that mitochondria play a significant role
in aging (10, 11). Although the functions of mitochondria are well understood, the relationship
between mitochondrial dysfunction and age-associated diseases requires further investigation.
Mitochondrial morphology is associated with mitochondrial bioenergetic capacity.
Mitochondrial dynamics, mainly fusion and fission, provide a means for mitochondria to readily
adapt to the specific energy demands of the cell. Fusion and fission are regulated by optic
atrophy 1 (OPA1) and dynamin-related protein-1 (DRP1), respectively (4, 12–14). While the loss

of DRP1 results in an abundance of overly elongated and fused mitochondria (5), the loss of
OPA1 gives rise to fragmented mitochondria (12, 15). Mitochondria may go through several
forms in coordinated cycles of fusion and fission. If OPA1 and mitofusions become upregulated,
fusion rates increase producing long, tubular mitochondria with more complex network
formation (16). Under normal physiological conditions, tubular mitochondria function well.
However, under conditions of stress, mitochondria have been shown to alter their formation into
donut, blob, or fragmented shapes, the latter of which is caused by a high fission-fusion ratio
(16–18). Importantly, the evaluation of mitochondrial structure is vital as deviations from typical
tubular mitochondrial shape indicate lower ATP production and increased susceptibility to
autophagosome degradation (19). Moreover, the loss of several mitochondrial genes has been
implicated in cristae structure alterations (13, 20).
Cristae, the folds of the mitochondria inner membrane, house electron transfer chain
complexes responsible for establishing the proton motive force, which ultimately produces ATP
(21). The morphological arrangement of cristae ultimately determines the energetic capacity of
mitochondria and has been implicated in playing a role in other functions including apoptosis
and homeostasis (21). The mitochondrial contact site and cristae organizing system (MICOS)
maintains the morphology of cristae (22). Similar to OPA1, the MICOS complex governs the
dynamics of cristae structure (20, 22–24). The MICOS complex is also involved in the
maintenance of cristae junctions, which are sub-compartments for metabolites (25). Given the
significance of cristae junctions, cristae may also be involved in other processes such as
regulating calcium homeostasis and molecular signaling. Considering the significance of
mitochondrial and cristae morphology in normal function, dysfunction of the MICOS complex
could negatively impact health (24). However, the role of the MICOS complex in mitochondrial
health during cardiac aging has yet to be defined. Given the understanding that mitochondria are
implicated in the aging process (11), we hypothesized that changes in the MICOS complex,
which is critical to maintaining cristae morphology (26, 27), may lead to age-associated changes
in mitochondria.
Specifically, we sought to understand the role of the MICOS complex in cardiac muscle
mitochondria as it may provide novel insights into maintaining normal mitochondrial health in
the heart. This study combines an *in vivo* and *in vitro* approach that looks at cardiac muscle in
mice, cardiac fibroblasts, and human induced pluripotent stem cell-derived cardiomyocytes
(iPSC-CMs). Previously, mitochondrial-focused aging studies have focused on cardiomyocytes
(28); however, these studies overlooked the MICOS complex in relation to the aging of the
cardiac fibroblasts. Cardiac health is linked to insulin resistance and diabetes (1, 29). Heart
failure is also associated with other mitochondrial processes (6, 30–32) such as endoplasmic
reticulum stress, mitochondrial bioenergetics, insulin signaling, autophagy, and oxidative stress
(33). Mitochondria have specialized types and structures associated with their roles in glycolysis
and oxidative metabolism in cardiac fibroblasts (17, 34–36). While young mitochondria show
extensive specialization, it is unclear whether this is also true for aged cardiac murine
mitochondria, and there is little information quantitively analyzing how aging affects cristae and
3D mitochondrial structure across aging in cardiac tissue. In cardiac muscle, research has shown
that across aging, mitochondria in between myofibrils, called intermyofibrillar mitochondria,
have the most significant decrease in mitochondrial oxidative respiration function (37).
However, the full extent of causes of these changes in the oxidative phosphorylation of
intermyofibrillar mitochondria have yet to be elucidated. Given that heart failure risk increases

with aging (38), it is vital to consider the role of mitochondrial dysfunction in the loss of heart
function (33).
To better understand the relationship between cardiac mitochondrial structure and the
aging process, we used transmission electron microscopy (TEM) and fluorescence for 2D
micrographs to observe mitochondria and cristae. Different muscle types have distinct
mitochondrial function (17, 34). However, it is unclear how these unique functioning differences
change over age. To further understand how aging affected mitochondrial morphology, we
looked at cristae morphology at three-time points; 3-month, 1-year, and 2-year, which represent
"young," "middle-aged," and "old," mice, respectively. We expanded this to include serial
block-face scanning electron microscopy (SBF-SEM) with manual contour tracing to reconstruct
3D mitochondrial morphology and structure. We then quantitatively analyzed the 3D networks in
mouse cardiac muscle samples at different age intervals. Notably, 3D reconstruction is an
important avenue as the 3D morphology of mitochondria has been linked to their functional
capacity (17). Using SBF-SEM, we were able to look at single organelles to compare the shape,
morphology, count, complexity, and branching across different ages. Furthermore, we compared
the aging effects to the loss of the MICOS complex in relation to mitochondrial morphology
alterations. Specifically, we used CRISPR/Cas9 on cardiac fibroblasts to knockout (KO) three
genes of the MICOS complex: *Chchd3* (Mic19), *Chchd6* (Mic25)*, and *Mitofilin* (Mic60), as well
as *Opa1*, as a positive control*, to see how the loss of the MICOS complex affected mitochondrial
size, morphology, and oxygen consumption rate. Finally, we sought to understand the role of the
MICOS complex, specifically *Chchd6*, in modulating the reactive oxygen species response in
iPSC-CMs.
158
159
160 **RESULTS:**
161
Ultrastructural Changes in Mitochondria and Cristae with Age
It is commonly understood that heart function declines with age in human and murine
models. However, these changes and their association to mitochondrial morphological changes
are unclear. To investigate the cardiac function of aged mice cohorts, prior to TEM analysis, we
collected and analyzed echometric data. We observed changes with age that included increased
cardiac output, stroke volume, and LV thickness, without significant changes in normalized heart
mass, ejection fraction, or heart rate (Figure S1).
As expected, mice increased in body weight across aging (Fig S2A). Specifically, the fat
mass increased significantly faster than lean mass, especially in the 2-year sample (Fig S2B-C).
In accordance with this, heart mass also significantly increased with age (Fig S2D). After a 6-
hour fast, a glucose tolerance test (GTT) showed that aged mice had higher plasma glucose
levels post-GTT, indicating impaired glucose tolerance (Fig S2E-F). These results are consistent
with aging mice having larger fat depots and abnormal glucose status in comparison to younger
mice. Critically, on average, even with decreased glucose tolerance, aged mice did not reach
blood glucose levels that are indicative of diabetes. Therefore, this cohort served as a strong
representation of age-independent from pathology in this study. To further understand how
mitochondrial affected these dynamics, TEM was utilized to understand aging mitochondrial and
cristae morphology.
We examined mitochondria and cristae morphology in cardiac muscle at 3-month, one-
year, and 2-year-old male mice. Different cell types can be studied in cardiac muscle (36) and

182 cardiac muscle can give information about potential causes of loss of optimal heart function with
183 age (38). TEM is a powerful tool for studying cristae in mitochondria as it gives very high-
184 quality micrographs (39). Young male mice (n=3, per age group) showed very clear
185 mitochondria with electron-dense membranes, whereas the aged mice had fewer mitochondria
186 and cristae (Figure 1A-A''). The number of mitochondria more than quadrupled between the 3-
187 month to 1-year aged samples before slightly leveling out between 1-year to 2-year (Figure 1B).
188 Although mitochondrial abundance increased, the average mitochondrial area was significantly
189 smaller in samples older than 3-month; the decrease is smaller between the 1-year and 2-year
190 samples (Figure 1C). This shows a compensating effect as mitochondrial area per fiber area does
191 not show significant changes across the total aging process (Figure 1D). The mitochondria also
192 became slightly more circular (Figure 1E). To measure ultrastructural details of mitochondria,
193 cristae numbers in each mitochondrion were evaluated and consistently showed that the number
194 of cristae decreased with age (Figure 1F). Finally, the cristae score was used to assess the cristae
195 quantity and morphology as it is often used to holistically evaluate cristae (39, 40). A cristae
196 score of 0 represents no clearly defined cristae, 1 represents that greater than 50% of
197 mitochondrial area is devoid of cristae, a cristae score of 2 represents that less than 75% of the
198 mitochondria area has cristae, and the maximum cristae score of 4 represents overall typically
199 defined cristae with normal architecture. There was a large drop in the quality of cristae in
200 cardiac muscle with age. The young 3-month sample had mostly regular and slightly irregular
201 cristae, with cristae scores of about 3.3. The aged samples both had cristae scores around 2,
202 which suggests there are many areas lacking cristae or having irregular cristae (Figure 1G). Thus,
203 we found that cristae count and quality both decreased across the aging process. These important
204 findings that indicate a loss of cristae folds with aging caused us to wonder if cristae were
205 differentially affected in a female model (Figure 1H-H'').
206  Previous research has indicated that male mice have impaired mitochondrial function in
207 cardiac muscle in comparison to female mice, in both healthy and cardiac pathological mice (41).
208 To see mitochondrial and cristae structure changes across aging in a sex-dependent manner, we
209 first looked at mitochondrial quantity which increased after a year of aging, while mitochondrial
210 area was inversely proportional (Figure 1I-K). Like the male model, the circularity index of
211 mitochondria increased with age (Figure 1L). Importantly, the female model recapitulated
212 findings of significant loss of cristae across the aging process in murine cardiac tissue.
213 Specifically, the number of cristae decreased (Figure 1M) consistently across aging, while the
214 cristae score also significantly decreased past the 3-month age point (Figure 1N). Although in the
215 female model, the cristae score improved slightly in comparing 2-year mice to 1-year mice, in
216 both cohorts the average cristae score remained below 2, representing that 25% or more of
217 mitochondrial area lacked typical cristae. Based on these findings, in tandem, we propose that
218 mitochondrial fission, resulting in decreased mitochondrial area, is increased while the quality of
219 cristae is decreased with age in a non-sex-dependent manner (Figure 1O). However, it should be
220 noted that TEM can be limited in analyzing mitochondrial changes beyond those of cristae
221 structure changes across the aging process. Given these changes did not present as sex-
222 dependent, to properly understand how potential novel phenotypes arose in cardiac tissue,
223 alterations in mitochondrial morphology were analyzed with 3D reconstruction in a male mouse
224 model.
225
226 **Aging Changes Mitochondrial Size in Cardiac Muscle: 3D Reconstruction Analysis**

Based on our observations of the lack of cristae folding in aging cardiac muscle, we
employed 3D techniques to image cardiac muscle biopsies from young (3-month-old), mature (1-
year-old), and aged (2-year-old) mice with SBF-SEM. With resolutions of 10 µm for the x- and
y- planes and 50 µm for the z-plane, SBF-SEM enables 3D reconstruction of mitochondria with
an accurate spatial resolution that cannot be seen in 2D. Given TEM did not reveal a sex-specific
change, we focused on a male model. Specifically, we examined the morphological changes in
the intermyofibrillar region as mitochondrial frequency has been demonstrated to increase in this
region (42). Thus, there is a need to determine if such changes in frequency are associated with
other mitochondrial-related aging changes, including mitochondrial orientation, network, and
nanotunnel alterations. To elucidate the changes in intermyofibrillar mitochondria in relation to
aging, we surveyed approximately 250 intermyofibrillar mitochondria from three male mice
(n=3) (Figure 2A) sampled at each age time point with SBF-SEM 3D reconstruction methods. At
a 10 µm by 10 µm image stack resolution, ~50 50-µm ortho slides (Figure 2B) were manually
traced at the transverse intervals (Figure 2C). This allowed for 3D reconstructions of
mitochondria to be created, as observed in the flowchart of Figure 2 (Figure 2D). In totality,
across the 3 mice, approximately 750 mitochondria for each age cohort were surveyed. All
measurements used 3D metrics to understand the volumetric dynamics of mitochondria (Figure
2H).
In Figure 2, we show representative images of the cardiac muscle at each aging point
(Figure 2F-H). The overlay of the 3D reconstruction (Figure 2F'-H'), and the isolated 3D
reconstruction (Figure 2F''-H''), allow for the mitochondrial structure to be viewed better
(Video 1-2). Each color represents an independent mitochondrion. The perimeter, 3D area (or
surface area), and volume decreased between 1-year to 2-year, which was a greater change than
between 3-month and 1-year (Figure 2I-K). Importantly, this shows that while TEM was able to
capture some dynamics of mitochondria, it neglected the alteration that is far greater in mice
moving from adulthood to seniority (1-year to 2-year) than from adolescence to adulthood (3-
month to 1-year). For these quantifications, three individual mice were sampled at each age
(Figure 2L-N). When comparing the mitochondrial quantifications from each mouse, they
showed overall little inter-group heterogeneity but persistent intra-individual variability.
With these observations, we sought to determine if mitochondrial networks changed in
response to aging. To further elucidate age-related changes, we looked at cardiac muscle from
both transverse (Figure 3A-C) and longitudinal views (Figure 3A'-C'). The mitochondrial
branching index (MBI) was utilized to better understand mitochondrial branching (Figure 3D).
The MBI measures network complexity by examining the ratio between transverse and
longitudinal mitochondria (36) and is similar to aspect ratio which measures the ratio between
major and minor axis (43). With this technique, we demonstrated that mitochondrial branching
decreased between 3-month and the 1-year (Figure 3E). Similarly, within an age cohort, there is
no significant variation between mice, but there was a large amount of heterogeneity in
mitochondrial samples in each mouse (Figure 3F). To further characterize the mitochondrial
types in each age cohort, we used mito-otyping, a method similar to karyotyping, to organize
mitochondria based on their volumes to better visualize the overall mitochondrial diversity
(Figure 3G). This allows for comparison of the mitochondria across ages at each volume. We
measured sphericity to understand how the surface area changed during the aging process
(Figure 3H). Sphericity generally showed minimal changes (Figure 3I) while there was
homogeneity across all 9 mice measured (Figure 3J). For all of these metrics, some outliers were
observed, which were omitted in the presentation for ease of view, but they were included in all

statistical analysis (Figure S3). Critically, together, this approach revealed that there were few
significant changes in morphology with only branching showing reductions. In combination, the
aged cardiac muscle mitochondrial morphology resembled healthy mitochondria with a reduced
size that lacks a phenotype or fragmentation (Figure 3). Since that the largest findings are loss in
cristae structure and mitochondria size across aging, we sought to understand the underlying
cause of these observed changes. Given the principal role of the MICOS complex in cristae
formation and maintenance (27, 44), we investigated its role in cristae and mitochondrial
remodeling across the aging process in cardiac muscle.
281
Aging Changes in MICOS Complex in Fibroblasts: 3D Reconstruction Analysis
Although it is established that the MICOS complex is critical for mitochondrial dynamics
(22, 26), it is unclear how aging affects the MICOS complex. Studies have shown that *Opa1*,
which is epistatic to the MICOS complex and physically interacts with components of the
MICOS complex (23), decreases with age (45). With *Opa1* as a positive control, we sought to
determine if the MICOS complex mRNA expression also decreased in cardiac muscle with age.
As previously suggested (45), *Opa1* mRNA decreased by over 50% between 3-months and 2-
years (Figure 4A). *Mitofilin* also decreased by 50% (Figure 4B). Likewise, *Chchd3* and *Chchd6*
also progressively decreased with age but not as much as the decline of other transcripts (Figure
4C-D). While *Opa1* interacts with the MICOS complex, it is not required for the formation of
cristae junctions at which the MICOS complex forms, nor does *Opa1* loss negatively affect
MICOS components (46). This suggests that the loss of the MICOS complex across aging occurs
in an *Opa1*-independent manner.
To further understand the role of mitochondrial dynamics upon the loss of these MICOS
genes, we utilized CRISPR/Cas9 to make a knockout (KO) model in fibroblasts, which was
validated by qPCR (Figure 5A-D). We have previously validated DRP1, MFN2, and OPA1
CRISPR-mediated gene knockout translates to protein level changes as shown by transmission
electron microscopy (39, 47). Coupled with the current use of qPCR, gene knockout translates to
reduced transcript and protein expression. Specifically, we measured approximately 1250
mitochondria across 10 cells. Since the loss of *Opa1* triggers changes in morphology (12, 15, 20,
39), we used it as a positive control for morphological changes. We marked mitochondria with
MitoTracker Red and verified the successful deletion of *Opa1, Mitofilin, Chchd3,* and *Chchd6*
(Figure 5E). Then z-stacks were reconstructed using Bitplane Imaris to quantify the number of
cardiac fibroblasts with loss of MICOS expression (Figure 5E). Deletions of *Chchd3, Mitofilin,*
Chchd6, or *Opa1* led to significant decreases in mitochondria length (Figure 5F) and volume
(Figure 5G). We further sought to determine if the mitochondria were more tubular, which
represents normal states, or fragmented, which represents stressed states (Figure 5H). Notably,
Chchd3-KO cells had nearly completely fragmented mitochondria, like those in *Opa1*-KO.
While the WT had mostly tubular mitochondria, the deletion of any of these critical
mitochondrial proteins led to a much higher proportion of fragmented mitochondria. We further
used live-confocal imaging to observe the dynamics of mitochondrial changes in real-time. The
WT mitochondria (Video 3) were normal and potentially undergoing fusion. However, the cells
with deletion of *Opa1* (Video 4), *Chchd3* (Video 5), or *Mitofilin* (Video 6) showed fragmented
mitochondria. These videos aid in showing the real-time mitochondrial dynamics that may be
contributing to changes in the 3D structure observed.
To understand how the loss of the MICOS complex affects mitochondrial activity, we
measured oxygen consumption rates (OCR) with an XF24 extracellular flux bioanalyzer upon

knockout of *Opa1* and the MICOS genes in cardiac fibroblasts (Figure 5I). The OCR
measurements encompassed four stages: basal respiration, ATP-linked respiration, maximal
respiratory capacity, and reserve capacity. Basal respiration reflects the cellular OCR under
normal, non-stressed conditions, while ATP-linked respiration represents the OCR directly
related to ATP synthesis, which can be measured following the application of oligomycin, an
inhibitor of ATP synthase. Maximal respiratory capacity refers to the maximum OCR a cell can
achieve under stress or high-energy demand and is measured following the application of
carbonyl cyanide-p-trifluoromethoxyphenylhydrazone (FCCP), an uncoupler of mitochondrial
oxidative phosphorylation. Reserve capacity represents the difference between maximal
respiratory capacity and basal respiration, indicating the cell's ability to respond to increased
energy demand and is measured following the application of Antimycin A/rotenone, inhibitors of
the electron transport chain. The *Opa1*-KO, *Chchd3*-KO, *Chchd6*-KO, and *Mitofilin*-KO cells
showed decreased basal OCR, ATP-linked OCR, and maximum OCR compared with the control
(Figure 5J-M), indicating a general loss of OCR at baseline and during high demand.
These results suggest that the depletion of the MICOS complex can lead to the impairment of the
electron transfer chain, mitochondrial respiration, and bioenergetics, providing valuable insights
for the development of targeted therapies for diseases associated with mitochondrial dysfunction.
Of relevance, while *Opa1*-KO had a dramatic decrease in the reserve capacity of OCR, *Mitofilin*-
KO cells showed no alteration in reserve capacity. These data show decreased oxygen
consumption rate at basal consumption, decreased ATP-linked respiration following oligomycin
application, decreased maximal respiratory capacity following FCCP application, and decreased
reserve capacity, as measured following AntimycinA/rotenone treatment.
Loss of OPA1 does not yield changes in mtDNA (14), while alterations in the MICOS
complex can affect mtDNA content (48, 49). Given these differential changes in mtDNA
content, as shown previously, total protein is the most appropriate way to normalize to
mitochondrial content (14, 50). Accordingly, cells were plated at a density of 20×10^{3}
consistently with normalized protein, which together will take into account changes from the
mitochondrial mass. Together this shows that changes in oxidative consumption occur
independently from changes in mitochondrial mass. Overall, depletion of these critical
mitochondrial proteins led to a general loss of oxygen consumption rate both at baseline and
during high demand. These findings suggest that depletion of the MICOS complex can lead to
the impairment of the electron transfer chain along with mitochondrial respiration and
bioenergetics.
352
Loss of *Chchd6* Results in Changes in Mitochondrial Size, Shape, and Fluorescence of
Reactive Oxygen Species in Induced Pluripotent Stem Cell-Derived Cardiomyocytes
To understand the impact of the MICOS complex beyond murine cardiac fibroblast
models, we knocked down *Chchd6*, a component of the MICOS complex, in induced pluripotent
stem cell-derived cardiomyocytes (iPCS-CMs). To verify changes in mitochondrial size and
morphology, we analyzed iPCS-CM through TEM (Figure 6A-B). We found that *Chchd6*-
deficient iPCS-CM had reduced mitochondrial area and perimeter while mitochondrial
circularity index increased (Figure 6C-F). We then moved to fluorescence live-cell imaging and
overlaid tetramethylrhodamine, ethyl ester (TMRE) red dye, a label of active mitochondria and
their membrane potential, dichlorodihydrofluorescein diacetate (DCFDA) green dye to assay for
overall reactive oxygen species, and Hoechst blue dye to label for DNA in control iPCS-CMs
(Figure 6G-G''') and *Chchd6*-depleted IPCS-CMs (Figure 6H-H'''). Analysis showed that while

mitochondrial membrane potential did not significantly change (Figure 6I), general reactive
oxygen species increased (Figure 6J). Following verifying the cardiomyocyte identity of our
iPSC-CM using troponin T staining (Figure 6K), to further validate this response and to look at
alterations in levels of superoxides (51) upon loss of the MICOS complex, we examined Mitosox
live-staining (Figure 6L-L'), which shows a significant increase of Mitosox fluorescence in the
Chchd6-depleted, suggesting that the loss of this subunit of the MICOS complex results in
increased in mitochondrial superoxides.
We also sought to understand the functional impact of loss of Chchd6 using the
Vanderbilt BioVU bank (Figure S4A). This is a powerful biobank of over 270,000 collected by
the Vanderbilt University Medical Center (52). This database includes a mixture of individuals
of European ancestry and African American ancestry (Figure S4B). Genetically regulated gene
expression (GREX) of CHCHD6 was calculated across cardiac tissues in individuals of
European ancestry. The relationship between CHCHD6 GREX and the "Heart failure with
reduced EF [Systolic or combined heart failure]" phenotype was evaluated across cases and
controls within BioVU. The associations between CHCHD6 GREX and heart failure was
nominally significant in 2 of the 3 tissues tested in individuals of European ancestry and 1 of the
3 tissues tested in individuals of African ancestry ($p < 0.05$). None of the associations met the
across-tissue Bonferroni-corrected p-value (0.0167). Cases were required to have two instances
of the "Heart failure with reduced EF [Systolic or combined heart failure]" phecode, controls had
no reports of "Heart failure with reduced EF [Systolic or combined heart failure]" in their
records. Individuals with similar codes were excluded from the analysis (Congestive heart
failure; non-hypertensive, Congestive heart failure (CHF) NOS, Heart failure NOS, Heart failure
with preserved EF [Diastolic heart failure], Ill-defined descriptions and complications of heart
disease, Heart transplant/surgery, Abnormal function study of cardiovascular system, Symptoms
involving cardiovascular system, Cardiac complications, not elsewhere classified). Covariates
included sex, age, median age of medical record, genotype batch, and genetic ancestry (principle
components 1-10). Together, this suggests that the MICOS complex may be linked to genetic
factors of heart failure, and plays a role in modulating membrane potential and oxidative stress in
iPCS-CM.
394
395 **DISCUSSION:**
396
Heart failure may be intrinsically linked to mitochondria (2, 33, 53), and understanding
the interplay of cristae dynamics, MICOS complex component expression, and mitochondrial
dynamics during aging may aid in the development of future therapies to restore energetic
production. Here, we demonstrated abnormal mitochondrial structure and impaired function
related to the loss of the MICOS complex. To our knowledge, this study is the first to show that
abnormal MICOS complex alters mitochondrial morphology in cardiac tissue assessed by 3D-
EM. Previous studies looking at mitochondria in aged cardiomyocytes did not find significant
changes in mitochondrial number, however, there was no comprehensive quantitative analysis of
the mitochondrial size and morphology (54). With the innovative 3D-EM technology, we
reported decreased mitochondrial volume and altered morphology with age. Importantly, our
study combined TEM and SBF-SEM to create novel 3D reconstructions of aging cardiac muscle
cells. Although the high resolution of TEM allows us to measure cristae (39), it only creates 2D
images that do not provide accurate spatial resolutions. Therefore, the 3D reconstructions of
mitochondria by SBF-SEM allow us to better understand mitochondrial morphology. Future

studies are needed that use focused ion beam-scanning electron microscopy (FIB-SEM) and thus,
allow for better resolution of the details of smaller subcellular objects, such as cristae (36, 55). In
the past, FIB-SEM was successfully performed in mouse cardiac muscle to characterize
mitochondria, but not sub-organellar structures, such as cristae (54, 56). However, mixed-
microscope approaches may still be required to achieve the resolution and sample size to analyze
changes in mitochondria associated with aging and organ, particularly muscle, behavior.
417   Previous studies have quantified mitochondria with SBF-SEM in human and mouse
muscles, yet the aging heart remains understudied with 3D techniques(36). Given that cardiac
muscle is responsible for proper heart contractions, there remains a gap in our understanding of
the changes of cardiac mitochondrial 3D structure in aging. When only examining TEM, our
results show that much of the age-dependent changes in mitochondrial architecture occurs only
from 3-months to 1-year, which is equivalent from adolescence to adult. However, when
switching to 3D reconstruction, we observe that mitochondrial size and morphology
progressively decrease across the aging process, including from 1-year to 2-years, which better
emulates geriatric aging. Interestingly, however, mitochondrial networking does not appear as
affected by the aging process past 1 year. In tandem, our 3D reconstruction elucidates how
cardiac muscle mitochondria change across aging in mice, highlighting the importance of the
MICOS complex in aging and its effects on cristae morphology and mitochondrial dynamics.
Future studies are needed that examine mtDNA and other modulators of cristae and
mitochondrial dynamics (57) as they may provide a further understanding of the MICOS
complex in relation to mitochondrial structure.
432   When viewing mitochondrial 3D structure across aging, we looked for the presence of
nanotunnels. Previous studies suggest that the formation of nanotunnels occurs during
mitochondrial stress. Nanotunnels are mitochondrial structures that allow for the transport of
materials and inter-mitochondrial communication (35). Nanotunnels have also been found in
human skeletal muscles and in diseased states (35). We did not find evidence of nanotunnel
formation in aged heart. This warrants future studies to investigate the factors leading to
nanotunnel formation and whether this mechanism also exists in cardiac muscle (58).
439   We observed a decline in MICOS components gene expression with age (Figure 7).
Additionally, with 3D reconstructions, we observed that the loss of *Opa1, Mitofilin, Chchd3,* and
Chchd6 in cardiac fibroblasts may be correlated with high mitochondrial dysfunction. This was
marked by smaller mitochondria, altered morphology, and reduced OCR. Given the higher
percentage of fragmented mitochondria and reduced mitochondrial volume of *Opa1*- KO,
Chchd3-KO, *Chchd6*-KO, and *Mitofilin*-KO. The exact cause of this fragmentation is unclear,
although past studies have linked cristae dysfunction and mitochondrial fragmentation,
potentially through mtDNA mutations (25). However, further study is needed to determine if the
loss of the MICOS complex is causing mitochondrial stress-related fragmentation, modulation of
fission-fusion dynamics, or acts on mitochondrial dynamics in an altogether different way.
449   It was predicted that these phenotypes would also arise in aging as expression of *Opa1*,
Mitofilin, *Chchd3*, and *Chchd6* is reduced across aging (Figure 4). In cardiac tissue, we observed
that the mitochondrial number increased with age, while the area decreased (Figure 1B-D). It is
possible that fission rates may be increasing over time, due to pathway changes such as *Opa1*
loss or *Drp1* upregulation (5, 20). Furthermore, the loss of cristae morphology supports the
occurrence of age-associated-decline of the MICOS complex. When looking at 3D
reconstruction, we principally noticed alterations in mitochondrial volume across aging which
parallel findings from the loss of MICOS complex. In combination, these findings suggest that

the loss of MICOS complex genes may be associated with a loss of overall mitochondrial
volume. However, in comparing morphological changes in aged samples with MICOS complex
KO fibroblasts, we noticed a differential change in morphology. Mitochondrial morphology in
the heart did not change as drastically as anticipated in response to aging. In general, the
sphericity of mitochondria did not undergo significant changes (Figure 3I).
Changes in mitochondrial shape are implicated in heart failure (33), which makes the
specific study of cardiac muscle across aging essential for the development of new therapeutics.
This study showed less networking, as measured through MBI, in cardiac mitochondria across
aging than other studies have suggested (Figure 3). It is possible that some cardiac metabolic
pathways are more resilient to aging changes, supporting the retention of mitochondrial
morphology. Interestingly, a recent study showed that spermidine treatment restores
mitochondrial morphology in the aged hearts (54), potentially indicating that spermidine
differentially is regulated specifically in cardiac tissue. However, future studies are warranted to
examine age-dependent mitochondrial changes in other tissues, to better understand tissue-
specific mitochondrial changes that occur during aging.
Looking at iPCS-CMs, the loss of the MICOS complex led to increased ROS (Figure 6),
consistent with previous studies in other models showing that loss of the MICOS complex may
cause oxidative stress (59). Although the mitochondrial free radical theory of aging has been
proposed for decades (60), the detailed mechanistic interplay between ROS and mitochondrial
dynamics remains incompletely understood. Previous studies suggest that a positive feed loop
within mitochondria promotes more ROS and thus, amplifies the ROS-related damage and
structural decline (61). Consistent with this, loss of *Opa1* has been reported to cause the
accumulation of ROS (62). Taken together, it is possible that during the aging process, the loss
of *Opa1* and the MICOS complex allows for more oxidative stress to occur, resulting in ROS-
induced ROS and the age-related functional decline of mitochondria (63). TMRE we used here
to validate TEM findings that there are no significant changes in active mitochondria, thus
oxidative stress changes can be attributed mainly due to altered generation (64). Interestingly, we
also noticed that when using TMRE as a baseline measurement of mitochondrial membrane
potential, we diverge from the literature in showing there is no change in the membrane
potential. Loss of the MICOS complex, specifically Mic60, results in a decrease in membrane
potential in Drosophila (48). A loss of membrane potential in OPA1 loss has also been seen in a
murine model (14). Yet, TMRE may not be the best monitor of membrane potential as loss of
negative charge affects its retention; thus, future studies may better investigate the MICOS-
dependent loss of membrane potential (64, 66). Using the BioVU BioBank, we also observed
linkages of *Chchd6* to factors of heart failure, with a differential ethnicity association (Figure
S4), suggesting that loss of the MICOS complex and concomitant modulation of mitochondria
structure has important implications in human health which must need to be further elucidated.
In summary, we combined TEM and 3D reconstructions to evaluate mitochondria and
cristae morphology in male murine cardiac muscle. Our findings indicate that the MICOS
complex decreases with age, and loss of the MICOS complex results in mitochondrial
dysfunction and changes in mitochondrial structure. These findings better our understanding of
how mitochondrial structure quantitively changes in cardiac muscle, and aid in understanding a
potential that can be targeted to protect cardiac muscle mitochondria from complete
fragmentation. Future studies are needed to explore this mechanism and continue to elucidate the
link between age-related changes in cardiac cristae structural integrity, the MICOS complex,
mitochondrial dysfunction, and ROS.

503
LIMITATIONS:
While we explored mitochondrial structure of around 750 mitochondria in each age
cohort of male murine myocardium tissue, there are several potential limitations of our study.
While the overall mitochondrial count was high, the quantity of mice in each age cohort is
limited, although we did not note large heterogeneity between the mice for most mitochondrial
quantifications. Notably, while Figure 7 shows a human model, this study was performed in a
male murine model. There are several known critical differences between human and rodent
hearts, which have differential developmental processes although they are generally considered
to have similar anatomy (65). For example, while human hearts are predominantly composed of
beta-Myosin heavy chain (MHC), in murine hearts, alpha-MHC is relatively more abundant
while pathophysiology and aging causes a shift toward beta-MHC (67, 68). Notably, myocardial
performance is partially dependent on the relative quantities of these MHCs, suggesting that they
may modulate mitochondrial structure and function (69). While mitochondrial-related
cardiovascular pathophysiology parallels each other between these models, greater research is
still necessary in potential model-dependent differences. Beyond this, while we show through
TEM that there are no sex-related differences on mitochondria across aging (Figure 1), future
studies may still consider performing 3D reconstruction of female mice. While it is commonly
understood that men are more vulnerable to age-related cardiovascular pathologies, female hearts
have been shown to have increased resilience towards oxidative stress (70), which suggests that
there may be a sex-dependent response to ROS. Finally, while the study sheds light on some of
the molecular mechanisms underlying age-related decline in mitochondrial function, it does not
address potential interventions or treatments to prevent or reverse this decline. Future research
may look at if restoring the MICOS complex can restore these age-related changes. Since we
only show a loss of MICOS transcripts over aging, it remains to be elucidated if increasing
MICOS protein amount may restore mitochondrial structure.
529
530
EXPERIMENTAL PROCEDURES
Animal Care and Maintenance
These protocols are previously described (14), and animals used in this study were cared for
using standard procedures approved by The University of Iowa Animal Care and Use Committee
(IACUC) that follow the National Institute of Health Guide for the Care and Use of Laboratory
Animals recommendations. WT male C57Bl/6J mice were exclusively used in experiments.
They were housed at 22°C on a 12-hour light, 12-hour dark cycle. They had free access to water
and standard chow. Mice were grown to various ages as described in the main text. 5%
isoflurane/95% oxygen was used to anesthetize mice.
540
RNA Extraction and RT-qPCR
RNA was extracted from tissue using TRIzol (Invitrogen) and RNeasy kit (Qiagen Inc). The
subsequent concentration of isolated RNA samples was measured using a NanoDrop 1000
(NanoDrop products, Wilmington, DE, USA) spectrophotometer at an absorbance of 260 nm and
280 nm. Reverse transcription was conducted on isolated RNA (~1 µg) using a High-Capacity
cDNA Reverse Transcription Kit (Applied Biosciences, Carlsbad, CA). SYBR Green (Life
Technologies, Carlsbad, CA) (1) was used for real-time quantitative PCR (qPCR). Three samples
for each qPCR (~50 ng) were placed in a 384-well plate that subsequently underwent qPCR in

the ABI Prism 7900HT instrument (Applied Biosystems) (Table 1)(14). The following qPCR
conditions were used: 1 cycle at 95 °C for 10 min; 40 cycles of 95 °C for 15 s; 59 °C for 15 s, 72
°C for 30 s, and 78 °C for 10 s; 1 cycle at 95 °C for 15 s; 1 cycle of 60 °C for 15 s; and one cycle
of 95 °C for 15 s. The results were normalized to glyceraldehyde-3-phosphate dehydrogenase
(*GAPDH*); data are shown as fold changes.
554
555   Table 1: qPCR Primers Used

Gene	Primers	Sequence
Opa1	Forward	5'-ACCAGGAGACTGTGTCAA-3'
	Reverse	5'-TCTTCAAATAAACGCAGAGGTG-3'
Chchd3	Forward	5'-GAAAAGAATCCAGGCCCTTCCACGCGC-3'
	Reverse	5'-CAGTGCCTAGCACTTGGCACAACCAGGAA-3'
Chchd6	Forward	5'-CTCAGCATGGACCTGGTAGGCACTGGGC-3'
	Reverse	5'-GCCTCAATTCCCACATGGAGAAAGTGGC-3'
Mitofilin	Forward	5'-CCTCCGGCAGTGTTCACCTAGTAACCCCTT-3'
	Reverse	5'-TCGCCCGTCGACCTTCAGCACTGAAAACCTAT-3'

556
557
CRISPR-Cas9 Knockouts
Primary mouse fibroblasts were isolated as previously described (71, 72). Briefly, after
cervical dislocation of 8-12 week mice, via sternotomy the heart was expose and transferred to
15 mL PBS centrifuge tube incubated at 37 °C. Aorta was removed and heart was washed with
10 mL of PBS warmed to 37 °C. Once ventricles were dissected, they minced into small pieces
of 1 mm^3, which were then subjected to enzymatic digestion using a mixture of
Collagenase/Dispase and DNase I in PBS, while the tissues were incubated at 37 °C with 5%
CO_2 atmosphere for 10 min on a rocking platform. After being centrifuged for 5 minutes at 1500
rpm, the tissue digest pellet was resuspended in 1 mL of fibroblast culture medium (M199
medium supplemented with 10% FBS and 2% penicillin-streptomycin). This process was
repeated approximately 9 times until the tissue was completely digested. Individual cell
suspensions were combined pelleted again by running in a centrifuge at 1500 rpm for 7 minutes.
Cells were resuspended in cardiac fibroblast media, and plated on 35 mm collagen-coated soft
hydrogel-bound polystyrene plates. After the media is aspirated, plates with adherent cells were
washed after 150 minutes with fibroblast culture medium. From there, 2 mL of fresh media was
added, and the plate was incubated at 37 °C with 5% CO_2 atmosphere. After 20 hours of
digestion, this washing and replacing of media process was repeated, and cells were grown to
90% confluency.
Once isolated, CRISPR/Cas9 was used to infect cardiac fibroblasts to produce the
following knockouts—control CRISPR/Cas9 (sc-418922), *Chchd6* (Mic25) CRISPR (sc-
425817), *Chchd3* (Mic19) CRISPR (sc-425804), and *Mitofilin* (Mic60) CRISPR (sc-429376)
(Santa Cruz Biotechnology, California, US) (Table 2). For each CRISPR, 2.5% of the CRISPR
was combined with 2.5% RNAiMax (ThermoFisher Scientific; 13778075), and 95% Opti-MEM
(Gibco; 31985070). This mixture was incubated at room temperature for 20 min. The media was
removed from the cells and washing twice with PBS. 200 µL of the CRISPR mixture and 800 µL
of Opti-MEM were added to each sample, then incubated at 37 °C for 4 hrs. An additional 1 mL
of DMEM medium was added before cells were incubated at 37 °C overnight. Fibroblasts were

washed with PBS, and a fresh medium was added. Experiments were performed 3- and 7-days
following knockouts.
587
588 Table 2: Guide RNA and Plasmids Used

Gene Name	Type of Plasmid	CAS Number
Mitofilin	CRISPR/Cas9 KO (m)	sc-429376
Chchd6	CRISPR/Cas9 KO (m)	sc-425817
Chchd3	CRISPR/Cas9 KO (m)	sc-425804
Control	CRISPR/Cas9 KO (m)	sc-418922

589
590
Processing of Mouse Left Ventricles for Serial Block-Face Scanning Electron Microscope (SBF-
SEM)
SBF-SEM was performed as previously described (15). Male mice were euthanized using 5%
isoflurane. The heart was excised and incubated in 2% glutaraldehyde with 100 mM phosphate
buffer for 30 min. Left ventricles were dissected, cut into 1-mm^3 cubes, and then incubated in
2.5% glutaraldehyde, 1% paraformaldehyde, and 120 mM sodium cacodylate solution for 1 hr.
597
From there, the tissue was three times washed with 100 mM cacodylate buffer at room
temperature. The tissue was immersed in 3% potassium ferrocyanide and 2% osmium tetroxide
for 1 hr at 4 °C, then washed three times with deionized water and incubated in 0.1%
thiocarbohydrazide and 2% filtered osmium tetroxide 30 min. Finally, the tissue was washed
three times with deionized water, before transferring to 1% uranyl acetate and left overnight at 4
°C. The next day, the samples were washed with deionized water, then incubated at 0.6% lead
aspartate solution for 30 min at 60 °C. From there, the samples were dehydrated using an acetone
gradient (20%, 50%, 70%, 90%, 95%, and 100% acetone for five mins each). Cardiac tissues
were impregnated in Epoxy Taab 812 hard resin, then moved to new resin; polymerization
occurred at 60°C for 36–48 hrs. Blocks of resin were sectioned, cut to 0.5 mm × 0.5 mm, and
glued to aluminum pins. These pins were transferred to the FEI/Thermo Scientific Volumescope
2 SEM. 300–400 ultrathin (0.09 µm) serial sections from each block were collected for
conventional TEM. All sections were collected onto formvar-coated slot grids (Pella, Redding
CA), stained, and imaged.
612
Measurement of OCR Using Seahorse
Oxygen consumption rate was measured for *Opa1*, *Cchchd3*, *Chchd6*, or *Mitofilin* KD
fibroblasts using an XF24 bioanalyzer (Seahorse Bioscience: North Billerica, MA, USA), as
previously described (14, 73). Fibroblasts were plated at a density of 20×10^3 per well and
differentiated for 3 days. For KO models, CRISPR/Cas9 for the relevant KO was added as per
the protocol above prior to plating. The media was changed to XF☐DMEM (supplemented with
1 g/L D-Glucose, 0.11 g/L sodium pyruvate, and 4 mM L-Glutamine) and cells were incubated
without CO_2 for 60 minutes. oligomycin (1 µg/ml), carbonyl cyanide 4-
(trifluoromethoxy)phenylhydrazone (FCCP; 1 µM), rotenone (1 µM), and antimycin A (10 µM
(Figure 5I) was performed in that order while remaining in the XF-DMEM media. After
measurement, cells were lysed accordingly to prior protocols (73), using 20 µl of 10 mM Tris
with 0.1% Triton X-100 added at pH 7.4, and media replaced with 480 µl of Bradford reagent.
Total protein concentration was then measured at absorbance at 595 nm and used for

normalization. For each sample, three independent experiments were performed for each
condition with representative data from the replicates being shown.
628
Quantification of TEM Micrographs and Parameters Using ImageJ
The National Institutes of Health (NIH) *ImageJ* software was used to quantify TEM images, as
previously described (39). Specifically, to ensure no bias in TEM quantification, one team
member was responsible for conducting the experiment and fixing the cells and tissue, while
another individual was responsible for processing and acquiring images using the electron
microscope in a blinded and randomized manner. Each fiber of interest was divided into four
equal quadrants, and two quadrants were randomly selected for measurements. Three additional
blinded members were responsible for quantifying the anonymized samples. By averaging their
collective findings, the potential for individual subjective bias was reduced. Moreover, those in
charge of quantification were given randomized images at both whole-cell and higher
magnification levels to further minimize bias.
640
Segmentation and Quantification of 3D SBF-SEM Images Using Amira
SBF-SEM images were manually segmented in Amira to perform 3D reconstruction, as
previously described (15). For each 3D reconstruction, (300–400 slices) were obtained and
transferred to Amira. By hand, an individual blinded to samples, traced structural features
manually on sequential slices of micrograph blocks. For each of the 3D reconstructions of
cardiac muscle in mice, 50-100 serial sections were chosen at approximately equal z-direction
intervals, stacked, aligned, and visualized using Amira to make videos and quantify volumetric
structures. Algorithms for measurements were entered manually for those not already in the
system (algorithms used are displayed in the main text; Figure 3D; Figure 3H). A total of 750
mitochondria from three mice were collected for each quantification.
651
652
Confocal mCherry-Mito-7 Labeling
To label the mitochondria of cardiac fibroblasts, the mCherry-Mito-7 plasmid was transfected
into the cells using a transfection reagent according to the manufacturer's instructions (74).
Briefly, the plasmid and the transfection reagent were diluted in Opti-MEM medium separately,
mixed gently, and incubated for 20 min at room temperature. The mixture was added to the
culture medium of the cells, and the cells were incubated for 24-48 h to allow expression of the
mCherry-Mito-7 protein. The localization of the mCherry-Mito-7 protein in cardiac fibroblasts
was visualized using a Leica SP8 Confocal Microscope. The cells were washed with PBS, fixed
with 4% paraformaldehyde for 10 min, and mounted with DAPI-containing mounting medium.
The fluorescent signals were observed using appropriate filters and recorded using a digital
camera.
664
Immunostaining protocol
For immunostaining, the hiPS-CMs were plated on glass-bottom dishes and were fixed in 4%
paraformaldehyde for 15 minutes, permeabilized with 0.1% Triton-X-100 for 8 minutes and
blocked with normal donkey serum for one hour, followed by incubation with primary antibody,
Cardiac Troponin-T (Invitrogen, Catalog # MA5-12960) overnight. They were then probed with
Donkey anti mouse AlexaFluor 555 secondary antibody (Thermo) and counterstained with

DAPI. Images were acquired using a Leica SP8 confocal microscope followed by quantification
using ImageJ software.

Segmentation and Quantification of 3D Fibroblasts Using Confocal Microscopy and Imaris
For the 3D reconstruction of fibroblasts, a minimum of 10 cells were chosen, and approximately
20 mitochondria from each cell were segmented for a total of about 200 mitochondria. 3D
structures were quantified as previously described (15) using the Imaris software (Bitplane),
which automatically measured many parameters. All individuals measuring the parameters were
blind to the experimental conditions but familiar in identifying diverse mitochondrial
morphology. For each of the 3D reconstructions of cardiac fibroblasts, 50-100 serial sections
were chosen at approximately equal z-direction intervals, stacked, aligned, and visualized.

Maintenance of induced pluripotent stem cell-derived cardiomyocytes
Cultured on Matrigel☐coated plates (Corning, Life Sciences), iPSC was fed 75% SFM and 25%
mTeSR$^+$. Cardiomyocyte differentiation was performed according to prior procedures (75), using
STEMdiff™ Cardiomyocyte Differentiation and Maintenance kits. Typically, CMs began
beating at day 8, and further enriched in day 10 of differentiation with lactic acid
(sodium l☐lactate, Sigma–Aldrich, USA). To aid CM maturation, triiodothyronine (T3) was
added across Days 20-40.

Live-Cell Staining

Live cell staining procedures for iPS☐CMs were performed according to prior protocols (75).
Once plated on glass☐bottom dishes, iPS-CMs media were replaced with 30 minutes with
culture medium (37°C) supplemented with DCFDA (5 mM) and TMRE (25 nM). Media was
replaced with new media supplemented with Hoechst 33342 for counterstaining. A Leica SP8
confocal microscope was used for image acquisition and ImageJ software was used for the
analysis of fluorescence intensity.

MitoSox
A stock solution of MitoSOXTM Red CMXRos-M7512 at a concentration of 1mM was prepared
by adding 1 vial of MitoSOXTM Red (50µg) to 127µl of DMSO, which was diluted to a
working solution of MitoSOXTM Red CMXRos at a concentration of 2.5µM by adding 10 µl of
the stock solution to 10 ml of media. Cells were first aspirated of their media and then incubated
with the working solution of MitoSOXTM Red at 37°C for 35 minutes. Following rinsing with
warm media, cells were fixed with 16% paraformaldehyde at 37°C for 15 minutes. Following
fixation, the cells were rinsed and imaged at 579 nm excitation and 599 nm emission
wavelengths. Quantification of relative intensity normalized to standardized image area.

Echocardiographic Parameters Collection
Animals were studied in accordance with protocols approved by the Institutional Animal Care
and Use Committees of the Carver College of Medicine of the University of Iowa. Data from 9-
10 animals at each age cohort were averaged to study in-vivo cardiac structure and function per
previous procedures (76, 77). After anesthetization in an induction chamber using 2.5%
isoflurane, mice were placed on a heated electrocardiography platform for heart rate monitoring
during the imaging procedure and maintained at 37°C. Mice had a nose cone administering 1%
isoflurane while in the left lateral decubitus position during imaging. A 13-MHz probe (Vivid V

echocardiograph; GE Healthcare, Tampa, FL) was used to take standard B-mode M-mode
images, along the short axis position of the papillary muscles.
717
Data Analysis
All SBF-SEM and TEM data were obtained from at least three independent experiments and are
presented as the mean across these experiments. In the presentation, black bars represent the
standard error of mean, and dots represent individual data points shown. For all data with only
two groups, an unpaired t-test was used. If more than two groups were compared, one-way
ANOVA was performed, and significance was assessed using Tukey *post hoc* tests for multiple
comparisons. For both analyses, GraphPad Prism software package was used (La Jolla, CA,
USA). A minimum threshold of $p < 0.05$ indicated a significant difference. Higher degrees of
statistical significance (**, ***, ****) were defined as $p < 0.01$, $p < 0.001$, and $p < 0.0001$,
respectively. For genetically-regulated gene expression statistical analysis was performed as
previously described (78). Briefly, transcriptome-wide association studies using gene, splicing,
and proteome were compared with Bonferroni correction (0.05/#genes tested) for multiple
testing correction. For all mouse models (Figure 2-3), each dot represents an independent mouse
(n=3) but the statistical analysis shown represents the 750 mitochondria quantified.
732

733 **ACKNOWLEDGEMENTS**
734 We would like to thank Mariya Sweetwyne for her advice as an aging expert.
735
FUNDING
This work was supported by National Institute of Health (NIH) NIDDK T-32, number entitled
Multidisciplinary Training in Molecular Endocrinology to Z.V.; This work was supported by
National Institute of Health (NIH) NIDDK T-32, number DK007563 entitled Multidisciplinary
Training in Molecular Endocrinology to A.C.; Integrated Training in Engineering and Diabetes",
Grant Number T32 DK101003; Burroughs Wellcome Fund Postdoctoral Enrichment Program
#1022355 to D.S.; The UNCF/ Bristol-Myers Squibb (UNCF/BMS)- E.E. Just Postgraduate
Fellowship in Life sciences Fellowship and Burroughs Wellcome Fund/ PDEP #1022376 to
H.K.B.; NSF grant MCB #2011577I and NIH T32 5T32GM133353 to S.A.M.; I01 BX005352
from the Dept. of Veterans Affairs Office of Research and National Institutes of Health
R01HD090061 to J.A.G., NSF EES2112556, NSF EES1817282, NSF MCB1955975, and CZI
Science Diversity Leadership grant number 2022-253614 from the Chan Zuckerberg Initiative
DAF, an advised fund of Silicon Valley Community Foundation to S.D.; The UNCF/Bristol-
Myers Squibb E.E. Just Faculty Fund, Career Award at the Scientific Interface (CASI Award)
from Burroughs Welcome Fund (BWF) ID # d, BWF Ad-hoc Award, NIH Small Research Pilot
Subaward to 5R25HL106365-12 from the National Institutes of Health PRIDE Program,
DK020593, Vanderbilt Diabetes and Research Training Center for DRTC Alzheimer's Disease
Pilot & Feasibility Program. CZI Science Diversity Leadership grant number 2022- 253529 from
the Chan Zuckerberg Initiative DAF, an advised fund of Silicon Valley Community Foundation
to A.H.J.; Its contents are solely the responsibility of the authors and do not necessarily represent
the official view of the NIH. The funders had no role in study design, data collection and
analysis, decision to publish, or preparation of the manuscript.
758
759 **CONFLICT OF INTEREST**
760 The authors declare that they have no conflict of interest.

References:

1. **Boudina S**, **Sena S**, **Theobald H**, **Sheng X**, **Wright JJ**, **Hu XX**, **Aziz S**, **Johnson JI**, **Bugger H**, **Zaha VG**, **Abel ED**. Mitochondrial energetics in the heart in obesity-related diabetes: direct evidence for increased uncoupled respiration and activation of uncoupling proteins. *Diabetes* 56: 2457–2466, 2007. doi: 10.2337/db07-0481.

2. **Brown DA**, **Perry JB**, **Allen ME**, **Sabbah HN**, **Stauffer BL**, **Shaikh SR**, **Cleland JGF**, **Colucci WS**, **Butler J**, **Voors AA**, **Anker SD**, **Pitt B**, **Pieske B**, **Filippatos G**, **Greene SJ**, **Gheorghiade M**. Mitochondrial function as a therapeutic target in heart failure. *Nat Rev Cardiol* 14: 238–250, 2017. doi: 10.1038/nrcardio.2016.203.

3. **Tyumentsev MA**, **Stefanova NA**, **Kiseleva EV**, **Kolosova NG**. Mitochondria with Morphology Characteristic for Alzheimer's Disease Patients Are Found in the Brain of OXYS Rats. *Biochemistry Moscow* 83: 1083–1088, 2018. doi: 10.1134/S0006297918090109.

4. **Alaimo A**, **Gorojod RM**, **Beauquis J**, **Muñoz MJ**, **Saravia F**, **Kotler ML**. Deregulation of Mitochondria-Shaping Proteins Opa-1 and Drp-1 in Manganese-Induced Apoptosis. *PLOS ONE* 9: e91848, 2014. doi: 10.1371/journal.pone.0091848.

5. **Lee Y**, **Jeong S-Y**, **Karbowski M**, **Smith CL**, **Youle RJ**. Roles of the mammalian mitochondrial fission and fusion mediators Fis1, Drp1, and Opa1 in apoptosis. *Molecular biology of the cell* 15: 5001–5011, 2004.

6. **Zuo W**, **Zhang S**, **Xia C-Y**, **Guo X-F**, **He W-B**, **Chen N-H**. Mitochondria autophagy is induced after hypoxic/ischemic stress in a Drp1 dependent manner: the role of inhibition of Drp1 in ischemic brain damage. *Neuropharmacology* 86: 103–115, 2014.

7. **Gunter TE**, **Yule DI**, **Gunter KK**, **Eliseev RA**, **Salter JD**. Calcium and mitochondria. *FEBS letters* 567: 96–102, 2004.

8. **Mehta MM**, **Weinberg SE**, **Chandel NS**. Mitochondrial control of immunity: beyond ATP. *Nat Rev Immunol* 17: 608–620, 2017. doi: 10.1038/nri.2017.66.

9. **Nicholls DG**. Mitochondria and calcium signaling. *Cell calcium* 38: 311–317, 2005.

10. **Barja G**. The mitochondrial free radical theory of aging. *Progress in molecular biology and translational science* 127: 1–27, 2014.

11. **Figueiredo PA**, **Mota MP**, **Appell HJ**, **Duarte JA**. The role of mitochondria in aging of skeletal muscle. *Biogerontology* 9: 67–84, 2008. doi: 10.1007/s10522-007-9121-7.

12. **Cipolat S**, **de Brito OM**, **Dal Zilio B**, **Scorrano L**. OPA1 requires mitofusin 1 to promote mitochondrial fusion. *Proceedings of the National Academy of Sciences* 101: 15927–15932, 2004. doi: 10.1073/pnas.0407043101.

13. **Otera H**, **Miyata N**, **Kuge O**, **Mihara K**. Drp1-dependent mitochondrial fission via MiD49/51 is essential for apoptotic cristae remodeling. *Journal of Cell Biology* 212: 531–544, 2016.

14. **Pereira RO**, **Tadinada SM**, **Zasadny FM**, **Oliveira KJ**, **Pires KMP**, **Olvera A**, **Jeffers J**, **Souvenir R**, **Mcglauflin R**, **Seei A**. OPA 1 deficiency promotes secretion of FGF 21 from muscle that prevents obesity and insulin resistance. *The EMBO journal* 36: 2126–2145, 2017.

15. **Garza-Lopez E**, **Vue Z**, **Katti P**, **Neikirk K**, **Biete M**, **Lam J**, **Beasley HK**, **Marshall AG**, **Rodman TA**, **Christensen TA**, **Salisbury JL**, **Vang L**, **Mungai M**, **AshShareef S**, **Murray SA**, **Shao J**, **Streeter J**, **Glancy B**, **Pereira RO**, **Abel ED**, **Hinton A**. Protocols for Generating Surfaces and Measuring 3D Organelle Morphology Using Amira. *Cells* 11: 65, 2022. doi: 10.3390/cells11010065.

16. **Wang C**, **Du W**, **Su QP**, **Zhu M**, **Feng P**, **Li Y**, **Zhou Y**, **Mi N**, **Zhu Y**, **Jiang D**, **Zhang S**, **Zhang Z**, **Sun Y**, **Yu L**. Dynamic tubulation of mitochondria drives mitochondrial network formation. *Cell Res* 25: 1108–1120, 2015. doi: 10.1038/cr.2015.89.

17. **Glancy B**, **Kim Y**, **Katti P**, **Willingham TB**. The Functional Impact of Mitochondrial Structure Across Subcellular Scales [Online]. *Frontiers in Physiology* 11, 2020. https://www.frontiersin.org/articles/10.3389/fphys.2020.541040 [16 Dec. 2022].

18. **Ahmad T**, **Aggarwal K**, **Pattnaik B**, **Mukherjee S**, **Sethi T**, **Tiwari B**, **Kumar M**, **Micheal A**, **Mabalirajan U**, **Ghosh B**, **Roy S**, **Agrawal A**. Computational classification of mitochondrial shapes reflects stress and redox state. *Cell death & disease* 4: e461, 2013. doi: 10.1038/cddis.2012.213.

19. **Rambold AS**, **Kostelecky B**, **Elia N**, **Lippincott-Schwartz J**. Tubular network formation protects mitochondria from autophagosomal degradation during nutrient starvation. *Proc Natl Acad Sci U S A* 108: 10190–10195, 2011. doi: 10.1073/pnas.1107402108.

20. **Frezza C**, **Cipolat S**, **De Brito OM**, **Micaroni M**, **Beznoussenko GV**, **Rudka T**, **Bartoli D**, **Polishuck RS**, **Danial NN**, **De Strooper B**. OPA1 controls apoptotic cristae remodeling independently from mitochondrial fusion. *Cell* 126: 177–189, 2006.

21. **Cogliati S**, **Enriquez JA**, **Scorrano L**. Mitochondrial cristae: where beauty meets functionality. *Trends in biochemical sciences* 41: 261–273, 2016.

22. **Kozjak-Pavlovic V**. The MICOS complex of human mitochondria. *Cell and tissue research* 367: 83–93, 2017.

23. **Glytsou C**, **Calvo E**, **Cogliati S**, **Mehrotra A**, **Anastasia I**, **Rigoni G**, **Raimondi A**, **Shintani N**, **Loureiro M**, **Vazquez J**. Optic atrophy 1 is epistatic to the core MICOS component MIC60 in mitochondrial cristae shape control. *Cell reports* 17: 3024–3034, 2016.

24. **Eramo MJ**, **Lisnyak V**, **Formosa LE**, **Ryan MT**. The 'mitochondrial contact site and
cristae organising system' (MICOS) in health and human disease. *The Journal of*
Biochemistry 167: 243–255, 2020. doi: 10.1093/jb/mvz111.

25. **Kondadi AK**, **Anand R**, **Hänsch S**, **Urbach J**, **Zobel T**, **Wolf DM**, **Segawa M**, **Liesa M**,
Shirihai OS, **Weidtkamp☐Peters S**, **Reichert AS**. Cristae undergo continuous cycles of
membrane remodelling in a MICOS☐dependent manner. *EMBO Rep* 21: e49776, 2020.
doi: 10.15252/embr.201949776.

26. **Friedman JR**, **Mourier A**, **Yamada J**, **McCaffery JM**, **Nunnari J**. MICOS coordinates
with respiratory complexes and lipids to establish mitochondrial inner membrane
architecture. *Elife* 4: e07739, 2015.

27. **Stephan T**, **Brüser C**, **Deckers M**, **Steyer AM**, **Balzarotti F**, **Barbot M**, **Behr TS**, **Heim**
G, **Hübner W**, **Ilgen P**. MICOS assembly controls mitochondrial inner membrane
remodeling and crista junction redistribution to mediate cristae formation. *The EMBO*
journal 39: e104105, 2020.

28. **Zhang J**, **He Z**, **Fedorova J**, **Logan C**, **Bates L**, **Davitt K**, **Le V**, **Murphy J**, **Li M**, **Wang**
M, **Lakatta EG**, **Ren D**, **Li J**. Alterations in mitochondrial dynamics with age-related
Sirtuin1/Sirtuin3 deficiency impair cardiomyocyte contractility. *Aging Cell* 20: e13419,
2021. doi: 10.1111/acel.13419.

29. **Parra V**, **Verdejo HE**, **Iglewski M**, **del Campo A**, **Troncoso R**, **Jones D**, **Zhu Y**,
Kuzmicic J, **Pennanen C**, **Lopez☐Crisosto C**, **Jaña F**, **Ferreira J**, **Noguera E**, **Chiong**
M, **Bernlohr DA**, **Klip A**, **Hill JA**, **Rothermel BA**, **Abel ED**, **Zorzano A**, **Lavandero S**.
Insulin Stimulates Mitochondrial Fusion and Function in Cardiomyocytes via the Akt-
mTOR-NFκB-Opa-1 Signaling Pathway. *Diabetes* 63: 75–88, 2013. doi: 10.2337/db13-
0340.

30. **Kim J**, **Wei Y**, **Sowers JR**. Role of mitochondrial dysfunction in insulin resistance.
Circulation research 102: 401–414, 2008.

31. **Friederich M**, **Hansell P**, **Palm F**. Diabetes, oxidative stress, nitric oxide and mitochondria
function. *Current diabetes reviews* 5: 120–144, 2009.

32. **Ngoh GA**, **Papanicolaou KN**, **Walsh K**. Loss of mitofusin 2 promotes endoplasmic
reticulum stress. *J Biol Chem* 287: 20321–20332, 2012. doi: 10.1074/jbc.M112.359174.

33. **Kenny HC**, **Abel ED**. Heart Failure in Type 2 Diabetes Mellitus. *Circ Res* 124: 121–141,
2019. doi: 10.1161/CIRCRESAHA.118.311371.

34. **Willingham TB**, **Ajayi PT**, **Glancy B**. Subcellular Specialization of Mitochondrial Form
and Function in Skeletal Muscle Cells. *Front Cell Dev Biol* 9: 757305, 2021. doi:
10.3389/fcell.2021.757305.

35. **Vincent AE**, **Turnbull DM**, **Eisner V**, **Hajnóczky G**, **Picard M**. Mitochondrial
Nanotunnels. *Trends in Cell Biology* 27: 787–799, 2017. doi: 10.1016/j.tcb.2017.08.009.

36. **Vincent AE**, **White K**, **Davey T**, **Philips J**, **Ogden RT**, **Lawless C**, **Warren C**, **Hall MG**,
Ng YS, **Falkous G**, **Holden T**, **Deehan D**, **Taylor RW**, **Turnbull DM**, **Picard M**.
Quantitative 3D Mapping of the Human Skeletal Muscle Mitochondrial Network. *Cell*
Reports 26: 996-1009.e4, 2019. doi: 10.1016/j.celrep.2019.01.010.

37. **Lesnefsky EJ**, **Chen Q**, **Hoppel CL**. Mitochondrial Metabolism in Aging Heart. *Circ Res*
118: 1593–1611, 2016. doi: 10.1161/CIRCRESAHA.116.307505.

38. **Chaudhary KR**, **El-Sikhry H**, **Seubert JM**. Mitochondria and the aging heart. *J Geriatr*
Cardiol 8: 159–167, 2011. doi: 10.3724/SP.J.1263.2011.00159.

39. **Lam J**, **Katti P**, **Biete M**, **Mungai M**, **AshShareef S**, **Neikirk K**, **Garza Lopez E**, **Vue Z**,
Christensen TA, **Beasley HK**, **Rodman TA**, **Murray SA**, **Salisbury JL**, **Glancy B**, **Shao**
J, **Pereira RO**, **Abel ED**, **Hinton A**. A Universal Approach to Analyzing Transmission
Electron Microscopy with ImageJ. *Cells* 10: 2177, 2021. doi: 10.3390/cells10092177.

40. **Eisner V**, **Picard M**, **Hajnóczky G**. Mitochondrial dynamics in adaptive and maladaptive
cellular stress responses. *Nat Cell Biol* 20: 755–765, 2018. doi: 10.1038/s41556-018-0133-
0.

41. **John C**, **Grune J**, **Ott C**, **Nowotny K**, **Deubel S**, **Kühne A**, **Schubert C**, **Kintscher U**,
Regitz-Zagrosek V, **Grune T**. Sex Differences in Cardiac Mitochondria in the New
Zealand Obese Mouse [Online]. *Frontiers in Endocrinology* 9, 2018.
https://www.frontiersin.org/articles/10.3389/fendo.2018.00732 [26 Jan. 2023].

42. **Vendelin M**, **Béraud N**, **Guerrero K**, **Andrienko T**, **Kuznetsov AV**, **Olivares J**, **Kay L**,
Saks VA. Mitochondrial regular arrangement in muscle cells: a "crystal-like" pattern.
American Journal of Physiology-Cell Physiology 288: C757–C767, 2005. doi:
10.1152/ajpcell.00281.2004.

43. **Picard M**, **White K**, **Turnbull DM**. Mitochondrial morphology, topology, and membrane
interactions in skeletal muscle: a quantitative three-dimensional electron microscopy study.
J Appl Physiol (1985) 114: 161–171, 2013. doi: 10.1152/japplphysiol.01096.2012.

44. **Hu C**, **Shu L**, **Huang X**, **Yu J**, **Li L**, **Gong L**, **Yang M**, **Wu Z**, **Gao Z**, **Zhao Y**, **Chen L**,
Song Z. OPA1 and MICOS Regulate mitochondrial crista dynamics and formation. *Cell*
Death Dis 11: 1–17, 2020. doi: 10.1038/s41419-020-03152-y.

45. **Tezze C**, **Romanello V**, **Desbats MA**, **Fadini GP**, **Albiero M**, **Favaro G**, **Ciciliot S**,
Soriano ME, **Morbidoni V**, **Cerqua C**, **Loefler S**, **Kern H**, **Franceschi C**, **Salvioli S**,
Conte M, **Blaauw B**, **Zampieri S**, **Salviati L**, **Scorrano L**, **Sandri M**. Age-Associated
Loss of OPA1 in Muscle Impacts Muscle Mass, Metabolic Homeostasis, Systemic
Inflammation, and Epithelial Senescence. *Cell Metab* 25: 1374-1389.e6, 2017. doi:
10.1016/j.cmet.2017.04.021.

46. **Barrera M**, **Koob S**, **Dikov D**, **Vogel F**, **Reichert AS**. OPA1 functionally interacts with
MIC60 but is dispensable for crista junction formation. *FEBS Letters* 590: 3309–3322,
2016. doi: 10.1002/1873-3468.12384.

47. **Hinton A**, **Katti P**, **Christensen TA**, **Mungai M**, **Shao J**, **Zhang L**, **Trushin S**, **Alghanem A**, **Jaspersen A**, **Geroux RE**, **Neikirk K**, **Biete M**, **Lopez EG**, **Shao B**, **Vue Z**, **Vang L**, **Beasley HK**, **Marshall AG**, **Stephens D**, **Damo S**, **Ponce J**, **Bleck CKE**, **Hicsasmaz I**, **Murray SA**, **Edmonds RAC**, **Dajles A**, **Koo YD**, **Bacevac S**, **Salisbury JL**, **Pereira RO**, **Glancy B**, **Trushina E**, **Abel ED**. A Comprehensive Approach to Sample Preparation for Electron Microscopy and the Assessment of Mitochondrial Morphology in Tissue and Cultured Cells. .

48. **Wang L**, **Hsu T**, **Lin H**, **Fu C**. Drosophila MICOS knockdown impairs mitochondrial structure and function and promotes mitophagy in muscle tissue. *Biology Open* 9: bio054262, 2020. doi: 10.1242/bio.054262.

49. **Qin J**, **Guo Y**, **Xue B**, **Shi P**, **Chen Y**, **Su QP**, **Hao H**, **Zhao S**, **Wu C**, **Yu L**, **Li D**, **Sun Y**. ER-mitochondria contacts promote mtDNA nucleoids active transportation via mitochondrial dynamic tubulation. *Nat Commun* 11: 4471, 2020. doi: 10.1038/s41467-020-18202-4.

50. **Pereira RO**, **Marti A**, **Olvera AC**, **Tadinada SM**, **Bjorkman SH**, **Weatherford ET**, **Morgan DA**, **Westphal M**, **Patel PH**, **Kirby AK**, **Hewezi R**, **Bùi Trân W**, **García-Peña LM**, **Souvenir RA**, **Mittal M**, **Adams CM**, **Rahmouni K**, **Potthoff MJ**, **Abel ED**. OPA1 deletion in brown adipose tissue improves thermoregulation and systemic metabolism via FGF21. *eLife* 10: e66519, 2021. doi: 10.7554/eLife.66519.

51. **Dikalov SI**, **Harrison DG**. Methods for Detection of Mitochondrial and Cellular Reactive Oxygen Species. *Antioxid Redox Signal* 20: 372–382, 2014. doi: 10.1089/ars.2012.4886.

52. **Davis L**. Psychiatric Genomics, Phenomics, and Ethics Research In A 270,000-Person Biobank (BioVU). *European Neuropsychopharmacology* 29: S739–S740, 2019. doi: 10.1016/j.euroneuro.2017.06.069.

53. **Bayeva M**, **Gheorghiade M**, **Ardehali H**. Mitochondria as a Therapeutic Target in Heart Failure. *Journal of the American College of Cardiology* 61: 599–610, 2013. doi: 10.1016/j.jacc.2012.08.1021.

54. **Messerer J**, **Wrede C**, **Schipke J**, **Brandenberger C**, **Abdellatif M**, **Eisenberg T**, **Madeo F**, **Sedej S**, **Mühlfeld C**. Spermidine supplementation influences mitochondrial number and morphology in the heart of aged mice. *Journal of Anatomy* 242: 91–101, 2023. doi: 10.1111/joa.13618.

55. **Baena V**, **Conrad R**, **Friday P**, **Fitzgerald E**, **Kim T**, **Bernbaum J**, **Berensmann H**, **Harned A**, **Nagashima K**, **Narayan K**. FIB-SEM as a Volume Electron Microscopy Approach to Study Cellular Architectures in SARS-CoV-2 and Other Viral Infections: A Practical Primer for a Virologist. *Viruses* 13: 611, 2021. doi: 10.3390/v13040611.

56. **Glancy B**, **Hartnell LM**, **Malide D**, **Yu Z-X**, **Combs CA**, **Connelly PS**, **Subramaniam S**, **Balaban RS**. Mitochondrial reticulum for cellular energy distribution in muscle. *Nature* 523: 617–620, 2015.

57. **Kondadi AK**, **Anand R**, **Reichert AS**. Functional Interplay between Cristae Biogenesis,
Mitochondrial Dynamics and Mitochondrial DNA Integrity. *Int J Mol Sci* 20: 4311, 2019.
doi: 10.3390/ijms20174311.

58. **Lavorato M**, **Iyer VR**, **Dewight W**, **Cupo RR**, **Debattisti V**, **Gomez L**, **Fuente SD la**,
Zhao Y-T, **Valdivia HH**, **Hajnóczky G**, **Franzini-Armstrong C**. Increased mitochondrial
nanotunneling activity, induced by calcium imbalance, affects intermitochondrial matrix
exchanges. *PNAS* 114: E849–E858, 2017. doi: 10.1073/pnas.1617788113.

59. **Warnsmann V**, **Marschall L-M**, **Meeßen AC**, **Wolters M**, **Schürmanns L**, **Basoglu M**,
Eimer S, **Osiewacz HD**. Disruption of the MICOS complex leads to an aberrant cristae
structure and an unexpected, pronounced lifespan extension in Podospora anserina. *Journal*
of Cellular Biochemistry 123: 1306–1326, 2022. doi: 10.1002/jcb.30278.

60. **Sanz A**, **Stefanatos RK**. The mitochondrial free radical theory of aging: a critical view.
Current aging science 1: 10–21, 2008.

61. **Ježek J**, **Cooper KF**, **Strich R**. Reactive Oxygen Species and Mitochondrial Dynamics: The
Yin and Yang of Mitochondrial Dysfunction and Cancer Progression. *Antioxidants* 7, 2018.
doi: 10.3390/antiox7010013.

62. **Quintana-Cabrera R**, **Manjarrés-Raza I**, **Vicente-Gutiérrez C**, **Corrado M**, **Bolaños JP**,
Scorrano L. Opa1 relies on cristae preservation and ATP synthase to curtail reactive
oxygen species accumulation in mitochondria. *Redox Biol* 41: 101944, 2021. doi:
10.1016/j.redox.2021.101944.

63. **Haas RH**. Mitochondrial Dysfunction in Aging and Diseases of Aging. *Biology* 8: 48, 2019.
doi: 10.3390/biology8020048.

64. **Chazotte B**. Labeling mitochondria with TMRM or TMRE. *Cold Spring Harb Protoc* 2011:
895–897, 2011. doi: 10.1101/pdb.prot5641.

65. **Wessels A**, **Sedmera D**. Developmental anatomy of the heart: a tale of mice and man.
Physiological Genomics 15: 165–176, 2003. doi: 10.1152/physiolgenomics.00033.2003.

66. **Chazotte B**. Labeling Mitochondria with MitoTracker Dyes. *Cold Spring Harb Protoc* 2011:
pdb.prot5648, 2011. doi: 10.1101/pdb.prot5648.

67. **Krenz M**, **Robbins J**. Impact of beta-myosin heavy chain expression on cardiac function
during stress. *Journal of the American College of Cardiology* 44: 2390–2397, 2004. doi:
10.1016/j.jacc.2004.09.044.

68. **Carnes CA**, **Geisbuhler TP**, **Reiser PJ**. Age-dependent changes in contraction and regional
myocardial myosin heavy chain isoform expression in rats. *Journal of Applied Physiology*
97: 446–453, 2004. doi: 10.1152/japplphysiol.00439.2003.

69. **Herron TJ**, **McDonald KS**. Small Amounts of α-Myosin Heavy Chain Isoform Expression
Significantly Increase Power Output of Rat Cardiac Myocyte Fragments. *Circulation*
Research 90: 1150–1152, 2002. doi: 10.1161/01.RES.0000022879.57270.11.

70. **Kander MC**, **Cui Y**, **Liu Z**. Gender difference in oxidative stress: a new look at the
mechanisms for cardiovascular diseases. *Journal of Cellular and Molecular Medicine* 21:
1024–1032, 2017. doi: 10.1111/jcmm.13038.

71. **Gansemer ER**, **McCommis KS**, **Martino M**, **King-McAlpin AQ**, **Potthoff MJ**, **Finck BN**,
Taylor EB, **Rutkowski DT**. NADPH and Glutathione Redox Link TCA Cycle Activity to
Endoplasmic Reticulum Homeostasis. *iScience* 23: 101116, 2020. doi:
10.1016/j.isci.2020.101116.

72. **Sahadevan P**, **Allen BG**. Isolation and culture of adult murine cardiac atrial and ventricular
fibroblasts and myofibroblasts. *Methods* 203: 187–195, 2022. doi:
10.1016/j.ymeth.2021.04.004.

73. **Dranka BP**, **Benavides GA**, **Diers AR**, **Giordano S**, **Zelickson BR**, **Reily C**, **Zou L**,
Chatham JC, **Hill BG**, **Zhang J**, **Landar A**, **Darley-Usmar VM**. Assessing bioenergetic
function in response to oxidative stress by metabolic profiling. *Free Radic Biol Med* 51:
1621–1635, 2011. doi: 10.1016/j.freeradbiomed.2011.08.005.

74. **Olenych SG**, **Claxton NS**, **Ottenberg GK**, **Davidson MW**. The fluorescent protein color
palette. *Curr Protoc Cell Biol* Chapter 21: Unit 21.5, 2007. doi:
10.1002/0471143030.cb2105s36.

75. **Yoon J**, **Daneshgar N**, **Chu Y**, **Chen B**, **Hefti M**, **Vikram A**, **Irani K**, **Song L**, **Brenner C**,
Abel ED, **London B**, **Dai D**. Metabolic rescue ameliorates mitochondrial
encephalo□cardiomyopathy in murine and human iPSC models of Leigh syndrome. *Clin*
Transl Med 12: e954, 2022. doi: 10.1002/ctm2.954.

76. **Tsushima K**, **Bugger H**, **Wende AR**, **Soto J**, **Jenson GA**, **Tor AR**, **McGlauflin R**, **Kenny**
HC, **Zhang Y**, **Souvenir R**, **Hu XX**, **Sloan CL**, **Pereira RO**, **Lira VA**, **Spitzer KW**,
Sharp TL, **Shoghi KI**, **Sparagna GC**, **Rog-Zielinska EA**, **Kohl P**, **Khalimonchuk O**,
Schaffer JE, **Abel ED**. Mitochondrial Reactive Oxygen Species in Lipotoxic Hearts
Induces Post-Translational Modifications of AKAP121, DRP1 and OPA1 That Promote
Mitochondrial Fission. *Circ Res* 122: 58–73, 2018. doi:
10.1161/CIRCRESAHA.117.311307.

77. **Hinton Jr AO**, **Yang Y**, **Quick AP**, **Xu P**, **Reddy CL**, **Yan X**, **Reynolds CL**, **Tong Q**, **Zhu**
L, **Xu J**, **Wehrens XHT**, **Xu Y**, **Reddy AK**. SRC-1 Regulates Blood Pressure and Aortic
Stiffness in Female Mice. *PLOS ONE* 11: e0168644, 2016. doi:
10.1371/journal.pone.0168644.

78. **Pathak GA**, **Singh K**, **Wendt FR**, **Fleming TW**, **Overstreet C**, **Koller D**, **Tylee DS**, **De**
Angelis F, **Mendoza BC**, **Levey DF**, **Koenen K**, **Krystal JH**, **Pietrzak RH**, **O' Donell C**,
Michael Gaziano J, **Falcone G**, **Stein MB**, **Gelernter J**, **Pasaniuc B**, **Mancuso N**, **Davis**
LK, **Polimanti R**. Genetically regulated multi-omics study for symptom clusters of

posttraumatic stress disorder highlights pleiotropy with hematologic and cardio-metabolic
traits. *Mol Psychiatry* 27: 1394–1404, 2022. doi: 10.1038/s41380-022-01488-9.

1021
1022
1023
1024
1025   **Figure Legend:**
1026
1027
Figure 1: Changes in cardiac muscle mitochondria and cristae across aging revealed in TEM and
heart data.
Representative transmission electron micrographs for cardiac muscle at (**A**) 3-months, (**A'**) 1-
year (**A''**) and 2-years in male mice. Blue boxes show cristae magnified to enhance the details of
cristae. Quantification of key mitochondrial characteristics included (**B**) the number of
mitochondria normalized to the area surveyed, (**C**) average mitochondrial area, (**D**) the total
mitochondrial area content per fiber area, and (**E**) the circularity index, which measures
mitochondrial shape. For cristae, (**F**) the number of cristae (**G**) and cristae score, a measurement
of the quality of cristae observed, are shown. Representative transmission electron micrographs
for cardiac muscle at (**H**) 3-months, (**H'**) 1-year (**H''**) and 2-years in female mice. Quantification
of (**I**) the number of mitochondria normalized to the area surveyed, (**J**) average mitochondrial
area, (**K**) mitochondrial area per fiber area, and (**L**) circularity index, (**M**) the number of cristae
(**N**) and cristae score. (**O**) Schematic showing dysfunction of mitochondrial cristae across the
aging process. Each dot represents a single mitochondrion. One-way ANOVA statistical test
performed with post hoc Tukey's Test. Significance values *, **, ***, ****
indicate $p \leq 0.05$, $p \leq 0.01$, $p \leq 0.001$, and $p \leq 0.0001$, respectively.

1044
Figure 2: Changes in cardiac muscle mitochondria morphology across aging revealed in SBF-
SEM. (**A**) Workflow depicting removal of the left ventricle of murine hearts. Following
embedded fixation, SBF-SEM allows for (**B**) ortho slice alignment. (**C**) Manual segmentation of
ortho slices was performed to (**D**) ultimately yield 3D reconstructions of mitochondria. (**E**)
Schematic depicting how metrics are found using pre-installed analyses on Amira 3D Software.
For some graphs, outlying dots were removed for presentation, but all mitochondria values were
considered in statistical analysis. (**F-G**) Representative SBF-SEM orthoslice for cardiac muscle
(n=3 mice). (**F'-G'**) 3D reconstructions of mitochondria in male cardiac tissues of different ages
overlaid on ortho slices. (**F''-G''**) 3D reconstructed and isolated mitochondria for clear
visualization. (**I-N**) 3D reconstructions were then quantified by (**I**) mitochondrial volume in
cardiac muscle of different ages, (**J**) 3D area of the average mitochondria in cardiac muscle, and
(**K**) Perimeter of the average mitochondria in cardiac muscle. Each dot represents the average of
a single mouse. Individual quantifications for (**L**) average volume per mitochondria, (**M**) average
3D area of mitochondria, and (**N**) average perimeter of the mitochondria for each of the three
mice sampled at 3-months, 1-year, and 2-years. One-way ANOVA statistical test performed with
post hoc Tukey's Test. Significance values *, **, ***, ****
indicate $p \leq 0.05$, $p \leq 0.01$, $p \leq 0.001$, and $p \leq 0.0001$, respectively.

1062
Figure 3: Changes in cardiac muscle branching and networking across aging revealed in SBF-
SEM. (**A-C**) 3D reconstruction of individually colored mitochondria from a transverse view for

mouse cardiac muscle of different ages. (**A'-C'**) 3D reconstruction of individually colored
mitochondria from a longitudinal view in cardiac muscle tissues of different ages. (**D**) Schematic
showing how transverse and longitudinal mitochondrial length is used to measure mitochondrial
branching index. (**E**) Mitochondrial branching index was measured to estimate mitochondrial
networks. (**F**) For each of the three male mice sampled at each aging time point, the
mitochondrial branching index is shown. (**G**) Mitochondria 3D reconstructions were further
organized by volume for each of the age cohorts. (**H**) Schematic showing how sphericity is
measured, as a function of volume to surface area ratio. (**I**) To measure shape, sphericity changes
across aging in cardiac muscle was further measured. (**J**) The average sphericity values from
each of the approximately 75 mitochondria surveyed per mice is also shown. Outlying dots were
removed for presentation for some graphs, but all mitochondria values were considered in
statistical analysis. One-way ANOVA statistical test performed with post hoc Tukey's Test.
Significance value **** and ** indicates $p \le 0.0001$ and $p \le 0.01$.
1078
Figure 4: Transcription of *Opa-1* and MICOS genes in aging cardiac muscle.
(**A-D**) qPCR analyzing the gene transcript fold changes of Opa-1 and MICOS across aging. (**A**)
Opa1 transcripts. (**B**) *Mitofilin* transcripts. (**C**) *Chchd3* transcripts. (**D**) *Chchd6* transcripts. One-
way ANOVA statistical test performed with post hoc Tukey's Test. Significance values *, **,
, * indicate $p \le 0.05$, $p \le 0.01$, $p \le 0.001$, and $p \le 0.0001$, respectively. For all
experiments, n=6.
1085
Figure 5: Loss of *Opa1* and MICOS genes result in mitochondrial structure changes and oxygen
consumption rate changes. (**A**) qPCR (n=6) analyzing the gene transcript fold changes of Opa-1
and MICOS in cardiac fibroblasts upon CRISPR/Cas9 knockdown. (**A**) *Opa1* transcripts. (**B**)
Mitofilin transcripts. (**C**) *Chchd3* transcripts. (**D**) *Chchd6* transcripts. (**E**) Confocal fluorescence
(using mCherry-Mito-7) shows changes upon individual knockout (KO) of *Opa1, Mitofilin,*
Chchd3, and *Chchd6* in mouse cardiac muscle. Below this, individual KO of *Opa1, Mitofilin,*
Chchd3, and *Chchd6* was also observed in 3D reconstruction of SBF-SEM. (**F-H**) Quantification
upon KO-state of each MICOS gene and *Opa1* (n=10 cells) was performed in 3D reconstruction.
(**F**) Mitochondria length and (**G**) mitochondria volume across each knockout. (**H**) Additionally,
the relative frequency of mitochondria that presented as fragmented or tubular was altered across
knock outs. (**I**) Seahorse Analyzer was utilized to measure oxygen consumption rate (OCR) in
Opa1, as positive control, along with *Chchd3*, *Chchd6,* and *Mitofilin* KD. (**J**) Basal OCR, which
shows baseline energetic demands, (**K**) ATP-linked, which was measured by ATP-inhibitor
oligomycin, (**L**) maximum OCR, which represents maximum respiration capacity upon
application of FCCP. Finally, the application of rotenone/antimycin A causes only non-
mitochondrial respiration to remain, which allows for measurement of (**M**) spare respiratory
capacity, which signifies the extra amount of ATP which can quickly be generated. For seahorse
n=6 plates for experimental knockouts, while for control n=16. For A-D: Unpaired (independent)
student's parametric statistical t-test performed. For all other statistical tests, One-way ANOVA
statistical test performed with Dunnett's multiple comparisons test.
1106
Figure 6: Chchd6-deficiency results in changes in mitochondrial amounts and reactive oxygen
species in induced pluripotent stem cell-derived cardiomyocytes (iPSC-CMs). (**A**) Transmission
electron microscopy (TEM) representative images of control and (**B**) CHCHD6-deficient
induced pluripotent stem cell-derived cardiomyocytes. TEM analysis of (**C**) area, (**D**) perimeter,

(**E**) circularity index, and (**F**) mitochondrial count (normalized to cell area). Fluorescence live-
cell imaging in control iPSC-CMs used (**G**) tetramethylrhodamine, ethyl ester (TMRE; red) dye
to label active mitochondria, (**G'**) Dichlorodihydrofluorescein diacetate (DCFDA; green) to
assay for overall reactive oxygen species, (**G''**) and Hoechst dye (blue) to label for DNA.
Overlaid fluorescence image is also presented (**G'''**) and fluorescence dye analysis were further
performed in (**H-H'''**) Chchd6-deficient IPSC-SMs. Analysis of fluorescent channels showing
relative fluorescence of (**I**) TMRE and (**J**) DCFDA. Further fluorescence dye with Hoechst for
DNA and Troponin-T (**K**) and MitoSox to measure reactive oxygen species including
superoxides in control (**L**) and (**L'**) Chchd6-deficient IPSC-SMs. (**M**) Analysis of MitoSox
fluorescence. (**C-F**) Each dot represents a single mitochondrion. (**I,J,M**) Each dot represents
quantification from a microscopic field from three independent replicates for each control and
KO groups. Unpaired (independent) student's parametric statistical t-test performed.
Significance value **** and * indicates $p \leq 0.0001$ and $p \leq 0.05$.
1124
1125   **Figure 7:** Graphical representation of loss of MICOS complex across aging that may occur,
1126   affecting the structure of mitochondria. Importantly, here we show this in a murine model, which
1127   may not translate to a human model (See Limitations section), but these findings suggest an
1128   important relation between the MICOS complex and aging which may be reproduced in human
1129   models.
1130
1131   **Figure S1.** Echocardiographic parameters of male and female mice at aged intervals of 3-month,
1132   1-year, and 2-years. Presented as Means±SEM. (**A-I**) (**A**) HR indicates heart rate; (**B**) LVEF, left
1133   ventricle ejection fraction; (**C**) CO, cardiac output; (**D**) SV, stroke volume; (**E**) shows heart mass
1134   and (**F**) heart mass divided by body mass (**G**) LVT, left ventricle thickness; (**H**) ESV, end
1135   systolic volume; (**I**) EDV, end diastolic volume. Variable number of mice surveyed (n=9-10).
1136
1137   **Figure S2.** Changes in (**A**) body weight, (**B**) amount of lean mass, (**C**) amount of fat mass, and
1138   the mass of hearts (**D**) changes in murine mice across their aging. Each dot represents a mouse
1139   sampled. (**E**) Furthermore, glucose tolerance test shows there is altered glucose tolerance. (**F**)
1140   The area under curve of glucose tolerance test comparing the 3 age cohorts. Five mice surveyed
1141   (n=5). One-way ANOVA statistical test performed with post hoc Tukey's Test.
1142
1143   **Figure S3.** Mitochondrial changes in (**A**) volume, (**B**) area 3D, (**C**) perimeter, (**D**) mitochondrial
1144   branching index, (**E**) and sphericity. All data are presented on a logarithmic scale.
1145
Figure S4. (**A**) Genetically-regulated gene expression for CHCHD6 was calculated in BioVU
participants using models built from the GTEx version 8 data (right panel), which contains
genotype data matched to RNA-Seq data from 838 donors across 49 tissues. BioVU participants
with a heart failure diagnosis were identified using ICD9/10 codes mapped to phecodes from
Vanderbilt's de-identified electronic health record database (middle panel). Imputed CHCHD6
expression was calculated and tested for association in heart failure cases versus controls using
logistic regression models (right panel), accounting for genetic ancestry (principle
components/PC 1-10), sex, age, and genotyping batch. (**B**) Demographics of BioVU Patients
with Systolic Heart Failure, with standard deviations of any means in parenthesis (**C**) CHCHD6
GREX Association with Heart Failure Results in Cardiac Tissues. Beta level, standard error
(SE), odds ratio of enrichment (OR), and p-value displayed for each tissue type.

1157
1158
Video 1 – 3D reconstruction of 3-month male murine cardiac tissue
Video 2 -- 3D reconstruction of 2-year male murine cardiac tissue
Video 3 – Confocal imaging of wildtype fibroblasts.
Video 4 – Confocal imaging of *Opa1* KD fibroblasts.
Video 5 – Confocal imaging of *Chchd3* KD fibroblasts.
Video 6 – Confocal imaging of *Mitofilin* KD fibroblasts
1165

Figure 1

Figure 2

Figure 3

Mitochondrial Branching Index = $\dfrac{\text{Transverse Distance}}{\text{Longitudinal Distance}}$

Sphericity = $\dfrac{\pi^{\frac{1}{3}}(6*Volume)^{2/3}}{Surface\ Area}$

Figure 4

Figure 5

Figure 6

Figure 7

Left Ventricle

MICOS Complex

MICOS

OMM

IMS

IMM

12/13

12/13

Cristae Junction

Matrix