# Immune activation state modulates retrieval of infant engrams

Sarah D. Power[1,2], Erika Stewart[1,2], Louisa G. Zielke[1,2,3], Eric Patrick Byrne[1,2], Clara Ortega-de San Luis[1,2], Lydia Lynch[1,6], Tomás J. Ryan[1,2,4,5*].

**Affiliations:**

[1] School of Biochemistry and Immunology, Trinity College Dublin, Dublin, Ireland.

[2] Trinity College Institute for Neuroscience, Trinity College Dublin, Dublin, Ireland.

[3] Faculty of Psychology and Neuroscience, Maastricht University, Netherlands.

[4] Florey Institute of Neuroscience and Mental Health, Melbourne Brain Centre, University of Melbourne, Melbourne, Victoria, Australia.

[5] Child & Brain Development Program, Canadian Institute for Advanced Research (CIFAR), Toronto, Ontario, Canada.

[6] Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA.

*Corresponding author. Email: tomas.ryan@tcd.ie

**Abstract:**
Infantile amnesia is possibly the most ubiquitous form of memory loss in mammals. Despite its widespread relevance, little is known about the biological conditions for infantile amnesia to occur and its effect on the engram cells that encode a memory. We investigated how memories are stored in the brain throughout development by integrating engram labeling technology with mouse models of infantile amnesia. We discovered a sex-specific phenomenon in which male offspring in maternal immune activation models of autism spectrum disorder do not experience infantile amnesia. We rescued the same apparently forgotten infantile memories in mice by optogenetically reactivating dentate gyrus engram cells that were labeled during complex experiences in infant development. Further, we were able to permanently reinstate lost infantile memories by artificially updating the memory engram, demonstrating that infantile amnesia is a reversible process. Our findings suggest that immune activation during development modulates innate, and reversible, forgetting switches that determine whether infantile amnesia will occur.

**Main Text:**

Infantile amnesia is the rapid forgetting of memories formed during early development; and is a largely neglected form of memory loss that seemingly affects 100 % of the human population. Memories of events that occur before the age of three are rarely remembered, and spotty recollection is observed for memories formed from 3 years of age (Rovee-Collier, 1999, Newcombe et al., 2000). Not uniquely a human phenomenon, this form of amnesia has been documented in rodents, which show forgetting of contextual and fear memories formed during the infant period (Campbell and Campbell, 1962, Callaghan and Richardson, 2012, Akers, 2014, Alberini and Travaglia, 2017). Little is known about the basic neurobiology of infantile amnesia and further its effect on the engrams cell ensembles that encode specific memories.

In cases of amnesia, such as Alzheimer's disease, it was previously assumed that memories were permanently lost from the brain. Thanks to the integration of activity-dependent ensemble labeling with optogenetics it is now possible to investigate if a memory engram is still present and/or functional in the brain even in the case of amnesia (Liu et al., 2012). Using this methodology, it has been shown that memory recall can be induced after amnesia by optogenetic of engram cells in the hippocampus and other brain regions, demonstrating that these memories are not only still present in the brain but are recoverable (Ryan et al., 2015, Roy et al., 2016, Perusini et al., 2017, Abdou et al., 2018). This framework provides an opportunity to investigate how development affects the storage and retrieval of early childhood memories.

The environment strongly influences learning and memory (Ryan and Frankland, 2022). During development, there are periods in which the developing brain has heightened sensitivity to environmental influences (Callaghan and Richardson, 2012, Choi et al., 2016, Cowan and Richardson, 2019, Cowan et al., 2019, Ohline and Abraham, 2019, Kalish et al., 2021, Sullivan and Opendak, 2020). Infantile amnesia has been shown to be preventable through pharmacological interventions with GABA agonists (Kim et al., 2006, Kim and Richardson, 2007, Tang et al., 2007) and corticosteroids (Callaghan and Richardson, 2012) or ectopic administration of neurotrophins (Travaglia et al., 2016) but the relevance of these switches to organismal function is unknown. Here we sought to identify naturally occurring variation in infantile amnesia caused by the developmental experience of the animal, and then investigate the effects on engram cell function.

Using a contextual fear conditioning paradigm (Fig. 1A), we trained and tested adult mice at postnatal day 63 (P63) for fear memory recall 1 or 8 days post-training (Fig. 1B). At both time points, the experimental shock group showed significantly more freezing than the no shock control group (Fig. 1C, D). Infant (P17) mice tested for memory recall 1 day following training showed significantly higher levels of freezing compared to the control group (Fig. 1C). While the group tested 8 days after training showed robust infantile amnesia, with similar levels of freezing to the no shock controls (Fig. 1D). Consistent with the literature infant mice demonstrate forgetting as early as one week after training, while adult mice show continuous memory retention (Guskjolen et al., 2018, Akers et al., 2014).

We tested whether altering environmental influences, such as environmental enrichment during infancy and the time of weaning from maternal care, affected the occurrence of infantile amnesia (fig. S1). Since these postnatal interventions had no effect, we next looked at the effect of challenging the embryonic environment. During pregnancy, exposure to pathogens or inflammation during critical periods of brain development (maternal immune activation) has been shown to alter postnatal brain growth and cognitive development (Vlasova et al., 2021).

The viral-mimetic compound Polynosinic-polycytidylic acid (Poly I:C), or the effector cytokine IL-17a, were used to stimulate maternal immune activation (MIA) in pregnant dams at E12.5 in C57BL6/J mice (Fig. 1G). In line with published studies, male offspring from pregnant dams treated with Poly I:C or IL-17a demonstrated repetitive behavior and deficits in social behavior, indicative of autism spectrum disorder (ASD) (fig. S2) (Choi et al., 2016, Shin Yim et al., 2017, Reed et al., 2020). To test if MIA causes any effect on memory retention, we next trained infant male and female offspring from MIA dams using contextual fear conditioning at P17. When tested 1 day after training all cohorts showed similar levels of freezing, irrespective of sex (Fig. 1H, J). However male, but not female offspring from pregnant dams treated with Poly I:C or IL-17a showed retention of infant memory, like that of adult mice, when tested 8 days after training (Fig. 1I, K) (Fig. S2M). This represents a novel phenomenon in which male offspring from MIA dams do not demonstrate infantile amnesia for a contextual fear memory.

**Fig. 1 MIA male offspring do not demonstrate infantile amnesia.**
(**A**) Developmental trajectory of infant mice. (**B**) Behavioral schedule. The black lightning symbol represents foot shocks. Mice are trained using contextual fear conditioning (CFC ) in context A and tested for recall 1 or 8 days later. (**C**-**F**) Memory recall in context A. Adult (P63) C57BLJ mice (n=8) froze significantly more in context A than no shock controls (n=8) both 1 (**C**) and 8 (**D**) days after training. (**E**) Infant (P17) C57BLJ mice (n=9) froze significantly more than no shock controls during recall 1 day after training. (**F**) No significant difference in freezing between groups (n=9) when tested for recall 8 days after training. (**G**) Representative diagram of MIA in mice. The syringe indicates Poly I:C (PIC) injection at E12.5. (**H**-**K**) Memory recall in context A after CFC at P17. (**H**) Male (n=10) and (**J**) female (n=9,10) MIA offspring show similar levels of freezing to controls when tested for recall 1 day after training. (**I**) Male (n=10) but not (**K**) female (n=9) MIA offspring freeze significantly more than controls 8 days after training. Data presented as mean ±SEM. Unpaired t test or one-way ANOVA Bonferroni; *P < 0.05, **P < 0.01, ***P < 0.001.

We characterized and employed an iCre-based engram labeling strategy for whole-brain tagging of engrams cells in infant mice, using a FosTRAP transgenic line expresses iCre from an inducible *c-fos* promoter (Fig. 2A-C) (Guenthner et al., 2013, Allen et al., 2017). Intraperitoneal (IP) injection of the tamoxifen metabolite, 4-hydroxytamoxifen (4-OHT), allows iCre to translocate into the nucleus. Ai32 transgenic mice express channelrhodopsin-2/EYFP (ChR2-EYFP) following exposure to Cre recombinase (Madisen et al., 2010, Madisen et al., 2012). By crossing TRAP mice with Ai32 mice it is possible to produce a stable transgenic line containing both Fos-iCre and ChR2-EYFP. We evaluated the efficiency of the Ai32-FosTRAP engram labeling system in infant (P17) and adult (P42) mice by quantifying the number of ChR2-EYFP+ cells in brain regions associated with context and fear memory including the dentate gyrus (DG), amygdala (AMG), retrosplenial cortex (RSC) and the periaqueductal gray (PAG) at P63 for a home cage, contextual or contextual fear memory (Fig. 2D-F, fig. S3).

We next investigated whether engram cells formed at the time of infant memory encoding are reactivated by exposure to training cues (Fig. 2H). Cellular activation of engram cells was examined by quantifying the number of EYFP$^+$ c-Fos$^+$ cells before (P20) and after (P63) the infantile amnesia period across multiple brain regions (Fig. 2G, fig. S5, S6). Natural recall cues result in above chance EYFP$^+$/c-Fos$^+$ overlap in the DG and AMG at P20 (Fig. 2H, I, fig. S5C, F), an effect that was not seen at P63 (Fig. 2J, K, fig. S6C, F). We next compared engram reactivation after natural recall for a contextual fear memory in male offspring from MIA dams (Fig. 2M). Interestingly, male offspring from MIA dams have an increased number of EYFP$^+$ engram cells in the DG compared to controls (Fig. 2N), indicating a larger engram ensemble also reported for a genetic model of ASD (Li et al., 2020). However, no difference was seen in the relative level of engram reactivation between groups as both demonstrated an above chance level of EYFP$^+$/c-Fos$^+$ overlap in the DG (Fig. 2P). Dendritic spine analysis showed that engram cells in MIA offspring had significantly increased dendritic spine density (Fig. 2R) as well as larger spine head diameter and volume relative to control animals at P25 (Fig. 2S, T). Thus, a combination of larger engram size, and increased dendritic plasticity, may contribute to functional engram expression of infant engrams in MIA offspring.

**Fig. 2. Natural reactivation of engram cells for an infant encoded memory.**
(**A**) Engram labeling method in Ai32-FosTRAP mice. (**B**) Whole-brain engram labeling of an infant engram. Scale bar 100 μm. (**C**) DG engram cells (ChR2-EYFP⁺ green) in Ai32-FosTRAP mice. Scale bar 100 μm. (**D**) Behavioral schedule for engram labeling of a homecage (HC), contextual (Cxt), or contextual fear memory (CFC) in P17 Ai32-FosTRAP mice. The black lightning symbol represents foot shocks. Syringe symbol represents 4-OHT injection 2 h after training. (**E, F**) ChR2-EYFP⁺ cell counts in the DG and AMG. (**G**) DG ChR2-EYFP⁺ (green) and c-Fos⁺ (red) cell counts after recall at P20 and P63. Scale bar 20 μm. (**H, M**) Behavioral schedule. (**I-L**) Engram reactivation in the DG and AMG after recall at (**I, K**) P20 and (**K**, **L**) P63. (**N**) ChR2-EYFP⁺ cell counts in the DG in MIA male offspring. (**O**) c-Fos⁺ cell counts in the DG. (**P**) c-Fos⁺/ChR2-EYFP⁺ cell counts in the DG. (**Q**) Dendritic spines in MIA male offspring. Scale bar 20 μm (**R**) Spine density per 10 μm. (**S**) Average dendritic spine head diameter (μm). (**T**) Average dendritic spine head volume (μm²). Data presented as mean ±SEM. Unpaired t test or one-way ANOVA Bonferroni; *P < 0.05, **P < 0.01, ***P < 0.001.

To investigate the functionality of memories after naturally occurring infantile amnesia, we tested the behavioral effect of optogenetically stimulating infant-labeled DG engram cells in adults (Fig. 3A, B). Engram cells for context A were labeled in Ai32-FosTRAP mice at P17 using CFC (Fig. 3C). Although both cohorts were placed in context A, only the experimental group received foot shocks (S). At P63 both cohorts were placed in context C for a 12 min test session where they received four 3 min epochs of blue light on or off. During the light-off epochs, both groups showed background levels of freezing (Fig. 3D). The experimental group showed significantly higher levels of freezing during the light-on epochs (Fig. 3D, E). This light-induced freezing was not observed for animals that underwent the same behavioral paradigm but did not receive any footshocks at P17. While stimulation of a neutral memory, labeled in infancy, does not result in freezing behavior (fig. S7A-C). Based on these data, 'lost' infantile memories can be acutely recalled by optogenetic stimulation.

Although optogenetic stimulation of a forgotten infant memory in adulthood resulted in a specific behavioral response, it is unclear how the information survives. We investigated the connectivity between engram cells in downstream regions, after infantile amnesia, by histologically assessing engram cell reactivation following optogenetic stimulation of upstream engram cells (Fig. 3F-K) (Redondo et al., 2014, Ryan et al., 2015). The resulting c-Fos+ counts were equivalent in the hippocampus after light activation of a CFC (shock, S) or a contextual (no shock, NS) memory (Fig. 3G). In contrast, c-Fos$^+$ cell counts were significantly higher in the AMG after light activation of an infant CFC memory (Fig. 3J), demonstrating that amygdala activity correlates with the behavioral expression of freezing. Light activation of DG engram cells results in an above chance c-Fos$^+$/EYFP$^+$ overlap in the DG (Fig. 3H, fig. S7D) and crucially in other downstream brain regions including the AMG (Fig. 3K, fig. S7E), RSC (fig. S7H, I) and PAG (fig. S7M, N). These results demonstrate that the connectivity pattern of the engram survives infantile amnesia and persists into adulthood, and optogenetic stimulation of a lost infantile memory is sufficient to reactivate the functional connectivity patterns. Further, encoding of a memory before (P17) or after (P29) the infantile amnesia period results in a similar engram connectivity pattern (fig. S8).

Since optogenetic stimulation results in memory recall under artificial conditions, we sought to permanently reinstate an infant memory by adopting an optogenetic induction procedure to induce plasticity in engram cells (fig. S9A, B) (Ramirez et al., 2013, Ryan et al., 2015). Infant mice can form purely contextual memories that can be updated with shock information (fig. S9C-F). At P63 and in a novel context, we optogenetically stimulated DG engram cells for context A, originally encoded at P17, while simultaneously delivering footshocks to create a false association of an infant context engram with adult shock experience (Fig. 3L). Animals subsequently froze significantly more when exposed to context A even though they were shocked in context C (Fig. 3M). This increased freezing was not due to generalization since the control group did not freeze in context B. There was also an increased level of overlap of c-Fos$^+$/EYFP$^+$ cells in the DG and RSC after exposure to context A, demonstrating that permanent engram reinstatement was also reflected at a cellular level (Fig. 3N-Q, fig. S10C, J). This histological result demonstrates that engram cells active at the time of encoding were reactivated during adult exposure to context A. Together, these findings show that the plasticity that accompanies the targeted updating of an infant engram restores the natural accessibility of that engram to the appropriate perceptual cues, and that context specificity is maintained into adulthood.

**Fig. 3. Artificial updating of infant encoded engram cells permanently reinstates a 'lost' infantile memory.**
(**A**) Behavioral schedule for optogenetic reactivation of infant DG engram cells in adulthood. (**B**) Representative image of optogenetic stimulation of DG engram cells in Ai32-FosTRAP mice. (**C**) Freezing levels of infant mice (n=10) during natural memory recall. Shock (S), no shock (NS). (**D**) Memory recall in context C (engram reactivation) with light-off and light-on epochs. (**E**) Freezing for the two light-off and light-on epochs averaged. (**F**, **I**) ChR2-EYFP+ cell counts in the DG and AMG after optogenetic stimulation at P63. (**G**, **J**) c-Fos+ cell counts in the DG and AMG. (**H**, **K**) c-Fos+/ChR2-EYFP+ cell counts in the DG and AMG. (**L**) Behavioral schedule for artificial updating of an infant engram. (**M**) Freezing levels during recall in context B and context A (n=10). Experimental light group froze significantly more in context A. (**N**, **O**) Histological representation of engram reactivation in the DG and RSC during recall in context A. DAPI+ (blue), ChR2-EYFP+ (green), c-Fos+ (red). Scale bar 20 μm. (**P**, **Q**) DG and RSC c-Fos+/ChR2-EYFP+ cell counts after recall in context A. Data presented as mean ±SEM. Unpaired t test or one-way ANOVA Bonferroni; *P < 0.05, **P < 0.01, ***P < 0.001.

To test the relevance of infantile amnesia to other types of memory, we next looked at this form of forgetting in a novel object recognition behavioral paradigm (Fig. 4A). Infant mice demonstrated rapid forgetting for novel object recognition tasks, following the developmental trend like that of other forms of episodic memory. Both adult and infant cohorts spent more time exploring a novel object relative to a familiar object when tested 1 day after acquisition, demonstrating that a long-term memory was formed (Fig. 4B, fig. S11C). When tested again 8 days later, only the adult cohort spent significantly more time exploring the novel object (Fig. 4C, fig. S11D). We then investigated whether male offspring from MIA dams show infantile amnesia for object memory (Fig. 4D). When tested for object memory retention, male offspring from MIA dams spent significantly more time with the novel object 8 days after acquisition (Fig. 4E-G). Like contextual fear memory, male offspring from Poly I:C or IL-17a injected dams do not show infantile amnesia for an object memory.

We then investigated whether it was possible to rescue a 'lost' infant object memory by optogenetic stimulation of infant-labeled engram cells (Fig. 4H). On the test day, animals that received light stimulation 3 min before the test demonstrated a preference for the novel object (Fig. 4J). While animals that did not receive light stimulation before the test spent the same amount of time exploring both the novel and familiar object (Fig. 4I). These object memory engrams survive infantile amnesia, can be retrieved artificially, and their forgetting can be prevented by MIA.

Spatial and object tasks in rodents are useful for the development of parallel, translational object/context and maze learning assays in human behavior experiments. To expand our analysis of infantile amnesia to an active, maze-based task we probed memory for a spatial location in infant and adult C57BL/6J mice using a cued version of the Barnes maze (Fig. 4K). All cohorts sufficiently learned the escape hole's location during training over five sessions, evidenced by a gradual decrease in the latency, and distance traveled, to reach the escape hole (fig. S12A, B). Unlike adults, infant mice demonstrated memory retention for the location of the escape hole when tested 1, but not 8, days later (Fig. 4L, M, fig. S12C-M). We assessed memory content profiling changes in navigational strategies taken by the animal during the task (Fig. 4Q). Navigational strategies can be categorized as random, procedural, or contextual (Garthe et al., 2009, Garthe et al., 2016, Berdugo-Vega et al., 2020). Infant mice predominantly used contextual search strategies when tested for spatial memory on the Barnes maze 1 day after training (Fig. 4R). However, during the 8-day probe trial infant mice show a switch in navigational strategy, moving to procedural and random strategies (Fig. 4R). This adaptation in navigational search strategies reflects the forgetting of the location of the escape platform. In contrast, adult mice show similar search patterns both 1 and 8 days after training (fig. S12N). Like other forms of memory, we also found that male offspring from MIA models do not show infantile amnesia for a spatial navigation task (Fig. 4N-P, fig. S12O-Q).

By labeling the engram ensembles for the location of the escape platform in the Barnes Maze in infants, we next investigated whether DG optogenetic stimulation of the engram in adulthood results in the location of the escape platform (Fig. 4S). The experimental light group received 3 min light stimulation before being placed on the Barnes maze for a probe test. The experimental light group spent significantly more time in the zone where the escape hole should be when compared to zone 2 and 3 (Fig. 4U). Although there was no significant difference in the total time spent between zone 4 and zone 1, heat map analysis of the mean time spent in each location showed a concentration around the location where the escape hole previously was located during training (Fig. 4U). No difference was seen in the total time spent between zones for the no-light control group (Fig. 4T). Changing the location of the escape hole during

memory encoding did not affect the recall of the location of the escape hole after light stimulation (Fig. 4V-X). Optogenetic stimulation of EYFP expressing neurons in the DG (neurons labeled during infant encoding of the Barnes maze memory) allows for recall of the location of the escape hole in the Barnes maze.

**Fig. 4. Infantile amnesia for context and maze memory.**
**A**, **D**) Behavioral schedule for novel object recognition. (**B, C**) Exploration time of infant C57BLJ mice during novel object recognition test 1 or 8 d after acquisition (n=9). (**E-G**) Exploration time of MIA male offspring (n=9-10). (**H**) Behavioral schedule for DG optogenetic reactivation of an infant encoded engram for an object. (**I, J**) Exploration time of adult mice after optogenetic stimulation (n=9-10). (**K**) Behavioral schedule for the Barnes maze in infant C57BLJ mice. (**L, M**) Time spent in each zone during test (**L**) 1 d or (**M**) 8 d after training. (**N-P**) Time spent in each zone by MIA offspring during test 8 d after training (n=9-11). (**Q**) Example representations of navigational search strategies. (**R**) Infant search strategies during testing 1 and 8 d after training. (**S, V**) Behavioral schedule for DG optogenetic reactivation of an infant encoded engram for the Barnes maze. (**T, U, W, X**) Time spent in each zone during test (n=8-9). Data presented as mean ±SEM. Unpaired t test or one-way ANOVA Bonferroni; *P < 0.05, **P < 0.01, ***P < 0.001.

To test whether the effect of immune activation on infantile amnesia is in fact mediated by the downstream effector cytokine IL-17a, we next injected pregnant *Il17a* knockout (KO) dams with Poly I:C at E12.5 (Fig. 5A). Adult male and female offspring from MIA *Il17a* KO dams did not demonstrate repetitive behavior or deficits in social behavior (fig. S13). Crucially, infant offspring show normal infantile amnesia for a CFC memory at P25 (Fig. 5 B-C). Preventing maternal IL-17a signaling, using a genetic model, occluded the MIA effect on infantile amnesia in the offspring. Therefore, the behavioral effect seen following MIA is specifically dependent on IL-17a signaling. We investigated whether altering the timing of immune activation, from maternal to postnatal, would have the same effect on memory retention (Fig. 5E) (López-Aranda et al., 2021). Postnatal immune activation in C57BLJ mice did not affect infantile amnesia, indicating that there is a critical window during prenatal development where IL-17a and immune activation can affect a mammal's propensity for infantile amnesia (Fig. 5F-H). Induction of an inflammatory response in MIA offspring has been shown to temporarily rescue deficits in sociability in ASD (Reed et al., 2020). Therefore, we investigated whether we could reverse the effect of MIA on infantile amnesia in the offspring through the administration of IL-17a. MIA male offspring were injected with IL-17a three hours before the recall test at P25 (Fig. 5I, J). Regardless of whether injected with IL-17a or PBS, MIA male infant offspring showed normal memory retention for a CFC memory 8 days after training, indicating that the effect of MIA on infantile amnesia cannot be reversed through IL-17a administration.

**Fig. 5. IL-17a is required for MIA effects on infantile amnesia.**
(**A**) Representative diagram of MIA in *Il17a* KO mice. (**B**) Behavioral schedule for CFC in *Il17a* KO male and female mice. (**C**, **D**) Freezing levels of male and female*Il17a* KO infant mice during recall 1 or 8 d after training. (**E**) Behavioral schedule for postnatal immune activation in C57BLJ infant mice. Syringe symbol represents Poly I:C injection at P3, P7, and P14. (**G**, **H**) Freezing levels of infant male and female mice when tested 8 d after training. (**I**, **J**) Behavioral schedule for IL-17a injection 3 h before recall at P25 in MIA C57bLJ offspring. (**K**) Freezing levels of infant male and female mice when tested 8 d after training. Data presented as mean ±SEM. Unpaired t test or two-way ANOVA Bonferroni; *P < 0.05, **P < 0.01, ***P < 0.001.

Together, these findings indicate that infantile amnesia for a wide range of memory types can be modulated by the immunological experiences of the subject during embryonic development. The effects of MIA are sex-specific and preserve the natural retrievability of infant engrams across postnatal development. Our data shows that infant memories are suppressed during development, but that their engrams persist and can be acutely activated in adulthood by direct optogenetic stimulation (Guskjolen et al., 2018). Furthermore, artificially updating an engram restored natural access to the target memory at both a behavioral and cellular level, reversing infantile amnesia. The specific connectivity patterns between engram cells, formed during infancy across distributed brain regions, also remain intact into adulthood and may account for stored information to be retained after infantile amnesia (Guskjolen et al., 2018). Postnatal environment experiences do not seem to affect infantile amnesia, making it plausibly driven by an innate predetermined switch. The brain state that permits infantile amnesia is absent in immunological models of ASD, and these findings are reminiscent of fly and human studies that have shown improved long-term retention of memories in cases of ASD (Dong et al., 2016, Zamoscik et al., 2016). It is conceivable that the MIA offspring brain state may mirror that of precocial mammals, who do not show infantile amnesia (Campbell et al., 1974, Akers, 2014). Infantile amnesia may represent a genetically canalized form of natural forgetting that can be modulated by developmental plasticity during critical periods. Future studies should determine the nature of the switching mechanism that determines when infantile amnesia occurs, its interaction with the immune system, and its reversible effect on engram cell function. Maternal immune activation models may offer new opportunities for translational studies of memory across the lifespan in mice and humans (Shook et al., 2022).

# References

ABDOU, K., SHEHATA, M., CHOKO, K., NISHIZONO, H., MATSUO, M., MURAMATSU, S.-I. & INOKUCHI, K. 2018. Synapse-specific representation of the identity of overlapping memory engrams. *Science,* 360**,** 1227-1231.

AKERS, K. G., MARTINEZ-CANABAL, A., RESTIVO, L., YIU, A. P., CRISTOFARO, A. D., HSIANG, H.-L., WHEELER, A. L., GUSKJOLEN, A., NIIBORI, Y., SHOJI, H., OHIRA, K., RICHARDS, B. A., MIYAKAWA, T., JOSSELYN, S. A. & FRANKLAND, P. W. 2014. Hippocampal Neurogenesis Regulates Forgetting During Adulthood and Infancy. *Science,* 344**,** 598-602.

AKERS, K. G. M.-C., A.; RESTIVO, L.; YIU, A.P; DE CRISTOFARO, A.; HSIANG, L.; WHEELER, A.L.; GUSKJOLEN, A.; NIIBORI, Y.; SHOJI, H.; OHIRA, K.; RICHARDS, B.A.; MIYAKAWA, T.; JOSSELYN, S.A; FRANKLAND, P.W. 2014. Hippocampal Neurogenesis Regulates Forgetting During Adulthood and Infancy. *Science,* 344.

ALBERINI, C. M. & TRAVAGLIA, A. 2017. Infantile Amnesia: A Critical Period of Learning to Learn and Remember. *J Neurosci,* 37**,** 5783-5795.

ALLEN, W. E., DENARDO, L. A., CHEN, M. Z., LIU, C. D., LOH, K. M., FENNO, L. E., RAMAKRISHNAN, C., DEISSEROTH, K. & LUO, L. 2017. Thirst-associated preoptic neurons encode an aversive motivational drive. *Science,* 357**,** 1149-1155.

BERDUGO-VEGA, G., ARIAS-GIL, G., LÓPEZ-FERNÁNDEZ, A., ARTEGIANI, B., WASIELEWSKA, J. M., LEE, C.-C., LIPPERT, M. T., KEMPERMANN, G., TAKAGAKI, K. & CALEGARI, F. 2020. Increasing neurogenesis refines hippocampal activity rejuvenating navigational learning strategies and contextual memory throughout life. *Nature Communications,* 11.

CALLAGHAN, B. L. & RICHARDSON, R. 2012. The effect of adverse rearing environments on persistent memories in young rats: removing the brakes on infant fear memories. *Translational Psychiatry,* 2**,** e138.

CAMPBELL, B. & CAMPBELL, E. H. 1962. Retention and extinction of learned fear in infant and adult rats. *Journal of Comparative and Physiological Psychology,* 55**,** 1-8.

CAMPBELL, B. A., MISANIN, J. R., WHITE, B. C. & LYTLE, L. D. 1974. Species differences in ontogeny of memory: Indirect support for neural maturation as a determinant of forgetting. *Journal of Comparative and Physiological Psychology,* 87**,** 193-202.

CHOI, G. B., YIM, Y. S., WONG, H., KIM, S., KIM, H., KIM, S. V., HOEFFER, C. A., LITTMAN, D. R. & HUH, J. R. 2016. The maternal interleukin-17a pathway in mice promotes autism-like phenotypes in offspring. *Science,* 351**,** 933-939.

COWAN, C. S. M. & RICHARDSON, R. 2019. Early-life stress leads to sex-dependent changes in pubertal timing in rats that are reversed by a probiotic formulation. *Dev Psychobiol,* 61**,** 679-687.

COWAN, C. S. M., STYLIANAKIS, A. A. & RICHARDSON, R. 2019. Early-life stress, microbiota, and brain development: probiotics reverse the effects of maternal separation on neural circuits underpinning fear expression and extinction in infant rats. *Dev Cogn Neurosci,* 37**,** 100627.

DONG, T., HE, J., WANG, S., WANG, L., CHENG, Y. & ZHONG, Y. 2016. Inability to activate Rac1-dependent forgetting contributes to behavioral inflexibility in mutants of multiple autism-risk genes. *Proceedings of the National Academy of Sciences,* 113**,** 7644-7649.

GARTHE, A., BEHR, J. & KEMPERMANN, G. 2009. Adult-Generated Hippocampal Neurons Allow the Flexible Use of Spatially Precise Learning Strategies. *PLoS ONE,* 4**,** e5464.

GARTHE, A., ROEDER, I. & KEMPERMANN, G. 2016. Mice in an enriched environment learn more flexibly because of adult hippocampal neurogenesis. *Hippocampus, 26,* 261-271.

GUENTHNER, C. J., MIYAMICHI, K., YANG, H. H., HELLER, H. C. & LUO, L. 2013. Permanent genetic access to transiently active neurons via TRAP: targeted recombination in active populations. *Neuron, 78,* 773-84.

GUSKJOLEN, A., KENNEY, J. W., DE LA PARRA, J., YEUNG, B. A., JOSSELYN, S. A. & FRANKLAND, P. W. 2018. Recovery of "Lost" Infant Memories in Mice. *Curr Biol, 28,* 2283-2290 e3.

KALISH, B. T., KIM, E., FINANDER, B., DUFFY, E. E., KIM, H., GILMAN, C. K., YIM, Y. S., TONG, L., KAUFMAN, R. J., GRIFFITH, E. C., CHOI, G. B., GREENBERG, M. E. & HUH, J. R. 2021. Maternal immune activation in mice disrupts proteostasis in the fetal brain. *Nature Neuroscience, 24,* 204-213.

KIM, J. H., MCNALLY, G. P. & RICHARDSON, R. 2006. Recovery of fear memories in rats: role of gamma-amino butyric acid (GABA) in infantile amnesia. *Behavioural Neuroscience, 120,* 40-8.

KIM, J. H. & RICHARDSON, R. 2007. Immediate post-reminder injection of gamma-amino butyric acid (GABA) agonist midazolam attenuates reactivation of forgotten fear in the infant rat. *Behavioural Neuroscience, 121,* 1328-32.

LI, J., JIANG, R. Y., ARENDT, K. L., HSU, Y. T., ZHAI, S. R. & CHEN, L. 2020. Defective memory engram reactivation underlies impaired fear memory recall in Fragile X syndrome. *Elife, 9.*

LIU, X., RAMIREZ, S., PANG, P. T., PURYEAR, C. B., GOVINDARAJAN, A., DEISSEROTH, K. & TONEGAWA, S. 2012. Optogenetic stimulation of a hippocampal engram activates fear memory recall. *Nature, 484,* 381-5.

LÓPEZ-ARANDA, M. F., CHATTOPADHYAY, I., BOXX, G. M., FRALEY, E. R., SILVA, T. K., ZHOU, M., PHAN, M., HERRERA, I., TALOMA, S., MANDANAS, R., BACH, K., GANDAL, M., GESCHWIND, D. H., CHENG, G., RZHETSKY, A., WHITE, S. A. & SILVA, A. J. 2021. Postnatal immune activation causes social deficits in a mouse model of tuberous sclerosis: Role of microglia and clinical implications. *Science Advances, 7.*

MADISEN, L., MAO, T., KOCH, H., ZHUO, J.-M., BERENYI, A., FUJISAWA, S., HSU, Y.-W. A., GARCIA, A. J., GU, X., ZANELLA, S., KIDNEY, J., GU, H., MAO, Y., HOOKS, B. M., BOYDEN, E. S., BUZSÁKI, G., RAMIREZ, J. M., JONES, A. R., SVOBODA, K., HAN, X., TURNER, E. E. & ZENG, H. 2012. A toolbox of Cre-dependent optogenetic transgenic mice for light-induced activation and silencing. *Nature Neuroscience, 15,* 793-802.

MADISEN, L., ZWINGMAN, T. A., SUNKIN, S. M., OH, S. W., ZARIWALA, H. A., GU, H., NG, L. L., PALMITER, R. D., HAWRYLYCZ, M. J., JONES, A. R., LEIN, E. S. & ZENG, H. 2010. A robust and high-throughput Cre reporting and characterization system for the whole mouse brain. *Nature Neuroscience, 13,* 133-140.

NEWCOMBE, N. S., DRUMMEY, A. B., FOX, N. A., LIE, E. & OTTINGER-ALBERTS, W. 2000. Remembering Early Childhood. *Current Directions in Psychological Science, 9,* 55-58.

OHLINE, S. M. & ABRAHAM, W. C. 2019. Environmental enrichment effects on synaptic and cellular physiology of hippocampal neurons. *Neuropharmacology, 145,* 3-12.

PERUSINI, J. N., CAJIGAS, S. A., COHENSEDGH, O., LIM, S. C., PAVLOVA, I. P., DONALDSON, Z. R. & DENNY, C. A. 2017. Optogenetic stimulation of dentate gyrus engrams restores memory in Alzheimer's disease mice. *Hippocampus.*

RAMIREZ, S., LIU, X., LIN, P., SUH, J., PIGNATELLI, M., REDONDO, R., RYAN, T. J. & TONEGAWA, S. 2013. Creating a false memory in the hippocampus. *Science,* 341**,** 216-219.

REDONDO, R. L., KIM, J., ARONS, A. L., RAMIREZ, S., LIU, X. & TONEGAWA, S. 2014. Bidirectional switch of the valence associated with a hippocampal contextual memory engram. *Nature,* 513**,** 426-30.

REED, M. D., YIM, Y. S., WIMMER, R. D., KIM, H., RYU, C., WELCH, G. M., ANDINA, M., KING, H. O., WAISMAN, A., HALASSA, M. M., HUH, J. R. & CHOI, G. B. 2020. IL-17a promotes sociability in mouse models of neurodevelopmental disorders. *Nature,* 577**,** 249-253.

ROVEE-COLLIER, C. 1999. The development of infant memory. *Curr Dir Psychol Science***,** 80–85.

ROY, D. S., ARONS, A., MITCHELL, T. I., PIGNATELLI, M., RYAN, T. J. & TONEGAWA, S. 2016. Memory retrieval by activating engram cells in mouse models of early Alzheimer's disease. *Nature,* 531**,** 508-12.

RYAN, T. J. & FRANKLAND, P. W. 2022. Forgetting as a form of adaptive engram cell plasticity. *Nature Reviews Neuroscience,* 23**,** 173-186.

RYAN, T. J., ROY, D. S. & PIGNATELLI, M. 2015. Engram cells retain memory under retrograde amnesia. *Science,* 348**,** 1007-1013.

SHIN YIM, Y., PARK, A., BERRIOS, J., LAFOURCADE, M., PASCUAL, L. M., SOARES, N., YEON KIM, J., KIM, S., KIM, H., WAISMAN, A., LITTMAN, D. R., WICKERSHAM, I. R., HARNETT, M. T., HUH, J. R. & CHOI, G. B. 2017. Reversing behavioural abnormalities in mice exposed to maternal inflammation. *Nature,* 549**,** 482-487.

SHOOK, L. L., SULLIVAN, E. L., LO, J. O., PERLIS, R. H. & EDLOW, A. G. 2022. COVID-19 in pregnancy: implications for fetal brain development. *Trends in Molecular Medicine,* 28**,** 319-330.

SULLIVAN, R. M. & OPENDAK, M. 2020. Defining immediate effects of sensitive periods on infant neurobehavioral function. *Current Opinion in Behavioral Sciences,* 36**,** 106-114.

TANG, H. H., MCNALLY, G. P. & RICHARDSON, R. 2007. The effects of FG7142 on two types of forgetting in 18-day-old rats. *Behav Neurosci,* 121**,** 1421-5.

TRAVAGLIA, A., BISAZ, R., SWEET, E. S., BLITZER, R. D. & ALBERINI, C. M. 2016. Infantile amnesia reflects a developmental critical period for hippocampal learning. *Nature Neuroscience,* 19**,** 1225-1233.

VLASOVA, R. M., IOSIF, A.-M., RYAN, A. M., FUNK, L. H., MURAI, T., CHEN, S., LESH, T. A., ROWLAND, D. J., BENNETT, J., HOGREFE, C. E., MADDOCK, R. J., GANDAL, M. J., GESCHWIND, D. H., SCHUMANN, C. M., VAN DE WATER, J., MCALLISTER, A. K., CARTER, C. S., STYNER, M. A., AMARAL, D. G. & BAUMAN, M. D. 2021. Maternal Immune Activation during Pregnancy Alters Postnatal Brain Growth and Cognitive Development in Nonhuman Primate Offspring. *The Journal of Neuroscience,* 41**,** 9971-9987.

ZAMOSCIK, V., MIER, D., SCHMIDT, S. N. & KIRSCH, P. 2016. Early Memories of Individuals on the Autism Spectrum Assessed Using Online Self-Reports. *Front Psychiatry,* 7**,** 79.

**Acknowledgments:**

We thank R. Cusack and M. Ramaswami for useful discussions; J. O'Leary, M. Pezzoli, L. Marks, and other Ryan Lab members for collegial support and scientific input.

**Funding:**

European Research Council Starting Grant (TJR)

Science Foundation Ireland Future Research Leaders Award (TJR)

Jacobs Foundation Research Fellowship (TJR)

Lister Institute of Preventive Medicine Award (TJR)

Brain & Behavior Research Foundation Young Investigator Grant (TJR)

Boehringer Ingelheim Fonds PhD Fellowship (SDP)

**Author contributions:**

Conceptualization: SDP, TJR

Methodology: SDP, COS, LL, TJR

Investigation: SDP, ES, LZ, COS, EPB

Funding acquisition: TJR

Project administration: SDP, TJR

Supervision: TJR

Writing – original draft: SDP, TJR

Writing – review & editing: SDP, ES, LZ, COS, LL, TJR

**Competing interests:** Authors declare that they have no competing interests.

**Data and materials availability:** All data are available in the main text or the supplementary materials.

**Supplementary Materials**

Materials and Methods

Figs. S1 to S13

**Materials and Methods**

Subjects

The Ai32-FosTRAP mice were generated by crossing Fos-iCre (*28*, *29*) mice with Ai32(RCL-ChR2(H134R)/EYFP) (*30*, *31*) and selecting offspring carrying both the CreETR and EYFP transgenes. All mice were socially housed in numbers 2-5 littermates, on a 12 h light/dark cycle with access to food and water *ad libitum*. The day of birth was designated postnatal day 0 (P0). Infant (P17) mice were housed with parents until the time of weaning when they were group-housed by the same sex. All mice were 7 weeks old before undergoing surgery. Post-operation mice were allowed to recover for a minimum of 2 weeks in their homecage before experimentation. All procedures relating to mouse care and treatment conformed with the health produces regulatory authority (HPRA) Ireland guidelines.

ChR2-EYFP expression in Ai32-FosTRAP mice

To label memory engram cells with this system, mice were injected intraperitoneally with 4-OHT (50mg/kg) 2 h post the learning event. The Fos-iCre line expresses the inducible c-Fos promoter. Injection of 4-OHT activates iCre recombinase. Activated iCre recombinase translocates to the nucleus and acts on two lox sites removing the stop codon that otherwise prevents expression of the ChR2/EYFP transgene. ChR2/EYFP transgene expression is driven by the pCAG promoter in iCre-expressing tissue 72 h after 4-OHT injection. For activity-dependent expression of ChR2-EYFP in Ai32-FosTRAP mice, Kainic acid (5mg/kg) was injected intraperitoneally followed by 4-OHT injection 2 h post kainic acid injection.

Stereotactic optical fiber implant

Mice were anesthetized using 500 mg kg$^{-1}$ avertin. Each animal underwent bilateral craniotomies using a 0.5 mm diameter drill bit at -2.0 mm anteroposterior (AP), ± 1.3 mm mediolateral (ML) for DG injections. A bilateral patch cord optical fiber implant (200 μm core diameter) was lowered above the injection site (-1.9 mm DV). A layer of adhesive cement was applied to secure the optical fiber implant to the skull. Dental cement was applied to secure a protective cap to the implant and close the surgical site. Each animal received meloxicam analgesic (0.075 mL/5g) via subcutaneous injection and remained on a heating pad until fully recovered from anesthesia. All mice were allowed to recover for two weeks before any subsequent experiments. All fiber sites were verified histologically.

Immunohistochemistry

Mice were dispatched by overdose with 50 μl sodium-pentobarbital and perfused transcardially with phosphate-buffered saline (PBS), followed by 4 % paraformaldehyde (PFA) in PBS. Extracted brains were kept in 4 % PFA at 4 °C overnight and stored in PBS. 50 μm coronal slices were cut using a vibratome and collected in PBS. Slices were washed in PBS-Triton X-100 (PBS-T) 0.2% followed by a 1 h blocking in PBS-T with 10% normal goat serum at room temperature before being incubated with the primary antibody at 4 °C overnight. Slices were washed in PBS-T 0.1 % followed by a 1 h incubation with the secondary antibody before undergoing another round of washing using PBS-T 0.1%. Vectashield DAPI was used to mount the slices onto super frost slides before visualization using an Olympus BX51 upright microscope. For cell counting experiments and dendritic spine analysis, an additional incubation with DAPI in PBS (1:1000) was carried out to label nuclei, and images were acquired using Leica SP8 gated stimulated emission depletion (STED) nanoscope. All images were taken at 40 X magnification.

## Cell counting

To measure the extent to which populations of active cells overlap between the exposure to the same or different contexts, the number of EYFP$^+$ and c-Fos$^+$ immunoreactive neurons in the DG, AMG, RSC, and PAG were counted. All animals were sacrificed 45 min post-assay or optical stimulation for immunohistochemical analyses. Four coronal slices were taken from the dorsal hippocampus of each animal (N=4, n=4). For optogenetic stimulation experiments, only slices with accurate implant location sites were used for counting. Sections were imaged on a Leica SP8 gated stimulated emission depletion (STED) nanoscope at a magnification of 40X. The area of interest was manually identified and the area of each region of interest was calculated using Fiji. To calculate the total number of DAPI cells in the DG, the average diameter of a sample of EYFP positive cells in the DG was taken from each animal and the area of the cell was calculated. The total number of cells was estimated as the area of the DG divided by the area of the cell. For the AMG, PAG, and RSC, the number of DAPI cells in 3 randomly selected regions of interest were counted and used along with the total area of the region to determine the total number of DAPI cells. The total number of c-Fos positive and EYFP positive cells were manually identified and counted for each region using Adobe Photoshop CC 2018. To calculate the percentage of cells expressing EYFP in each region, the total number of EYFP-positive cells was divided by the total number of DAPI-positive cells for each area. To calculate the percentage of cells expressing c-Fos in each region, the total number of c-Fos positive cells was divided by the total number of DAPI-positive cells for each area. To quantify engram reactivation, the number of EYFP and c-Fos positive cells were quantified as a percentage of total EYFP or DAPI positive cells.

## Dendritic spine analysis

The dendrites of DG engram cells in Ai32-FosTRAP mice were imaged using the Leica SP8 gated STED scope. Z-stack images (10 μm) of 50 μm coronal brain sections were taken using Leica Application Suite X (LasX) software (line average 2, zoom factor 1.7-1.9) at 40X. Imaris software (Oxford Instruments, Imaris v9.5) was used to carry out dendritic spine analysis. Dendrites were traced (10 μm) using a semi-automated neurofilament tracer tool and dendritic spines were individually highlighted and manually traced with the software. Each fragment represents a different ChR2-EYFP$^+$ cell. Parameters were generated automatically using Imaris software.

## Behavioral assays

All behavioral experiments were conducted during the light cycle of the day (07:00 am to 7:00 pm). All behavioral subjects were individually habituated to handling by the investigator for 3 min on 3 consecutive days. Handling took place in the breeding room. Immediately before each handling session mice were transported in separate cages to and from the vicinity of the experimental room to habituate them to the journey.

## Contextual fear conditioning

Three distinct contexts were used for contextual fear conditioning (CFC). Context A was a 31 x 24 x 21 cm Med Associates chamber with a removable grid floor (bars 3.2 mm diameter, spaced 7.9 mm apart), opaque triangular ceiling, and scented with 0.25 % benzaldehyde. Context B chambers were 29 x 25 x 22 cm Coulbourne Instruments chambers with Perspex white floors, bright white lighting, and scented with 1 % acetic acid. All conditioning sessions were conducted in Context A. Each session was 330 s in duration with three 0.75 mA shocks of 2 s duration delivered at 150 s, 210 s, and 270 s. Mice were removed from the conditioning chamber 60 s following the final footshock and returned to their homecage. All recall or context

generalization tests were 180 s in duration. Testing conditions were identical to training conditioning except that no shocks were presented. Up to 4 mice were run simultaneously in the 4 identical chambers. Floors of chambers were cleaned with Trigene before and between runs. Mice were transported to and from the experimental room in separate Perspex cages. All experimental groups were counter-balanced for chambers within contexts.

### Optogenetic engram reactivation

Optogenetic engram reactivation was carried out in Context C, a 31 x 24 x 21 cm Med Associates chamber with a white Perspex floor and curved insert, a dim light level and scented with 0.25 % acetophenone. The optical fiber implant was connected to a 45 nm laser diode fiber light source (Doric LDFLS 450/080). Habituation sessions lasted for 12 min. During testing, the 12 min session was divided into four 3 min epochs split into two light-off and two light-on epochs. During the light on epochs, blue light stimulation (20 Hz) with a pulse width of 15 ms was delivered through the optical fiber implant for the entire 3 min duration. At the end of the 12 min session, the mouse was immediately detached from the laser and returned to its homecage.

### Contextual pre-exposure

For context pre-exposure mice were exposed to Context A (31 x 24 x 21 cm Med Associates chamber with a removable grid floor) for 10 min. Mice were removed from the chamber and placed back in their homecage. The following day, mice were placed back in Context A where an immediate foot shock (1mA, 2s) was delivered. Mice were removed from the chamber 1 min after the shock and placed back in their homecage. Up to 4 mice were run simultaneously in the 4 identical chambers except for during the update when one mouse was run at a time. All recall or context specificity tests were 180 s in duration. Floors of chambers were cleaned with Trigene before and between runs.

### Artificial updating

Artificial updating of a contextual memory was carried out in 29 x 25 x 22 cm Coulbourne Instruments chambers with removable grid floors, bright white lighting, and scented with 1 % acetic acid. The optical fiber implant was connected to a 45 nm laser diode fiber light source (Doric LDFLS 450/080). The 7 min session consisted of a 2 min acclimatization period followed by 5 min of blue light stimulation (20Hz) with a pulse width of 15 ms delivered through the optical fiber implant for the entire 5 min duration. Three shocks, 0.75 mA of 2 s duration, were delivered at minutes 4, 5, and 6 of the session. At the end of the 7 min session, the mouse was immediately detached from the laser and returned to its homecage.

### Novel object recognition

Each subject was allowed to habituate to the rectangular testing arena for 10 min on the day before object acquisition. During object acquisition, two identical objects were positioned on adjacent walls of the arena. Mice were allowed to explore both objects freely for a 10 min period before being placed back in their homecage. On testing, one familiar object was replaced with a novel object and each subject was introduced for a 5 min exploration period. Object exploration was defined as the time when the subject's nose came within a 2 cm radius of the object. In between each trial, the testing arena and objects were cleaned with Trigene. For optogenetic reactivation experiments, animals received 3 min of blue light stimulation directly before being placed into the testing arena.

### Barnes maze

Training consisted of five trials with a 10 min limit per trial. At the start of each trial, the mouse was placed in a black PVC start chamber located in the center of the maze apparatus. After 15 secs, the start chamber was removed, and the latency and distance traveled to enter the escape tunnel were recorded. Subjects were placed back in their homecage and allowed 40 min in between each trial. The position of the escape tunnel remained in a fixed location relative to spatial cues in the room. For each probe trial, the escape tunnel was removed from the apparatus. Each mouse was allowed 5 min to freely explore the four quadrants of the maze before being removed and placed back in its homecage. The latency and distance traveled before reaching the location of the removed escape tunnel, along with the time spent in each quadrant were recorded. The apparatus and escape tunnel were cleaned with Trigene after each trial. For optogenetic reactivation experiments, animals received 3 min of blue light stimulation directly before being placed into the testing arena.

Maternal immune activation
Mice were mated overnight for one night. Females were weighed and checked for seminal plugs, noted as E0.5. Pregnant dams were injected SC with a single dose of Poly(I: C) (Sigma Aldrich) at 20mg/kg, rmlIL-17A (Immunotools) at 50ug/kg, or control PBS on E12.5. For rmlIL-17A injections at P25, mice were injected SC with a single dose of rmlIL-17A (Immunotools) at 50ug/kg or control PBS 3 h before the recall test.

Postnatal immune activation
Mice were injected SC with a single dose of Poly(I: C) (Sigma Aldrich) at 20mg/kg or control PBS on P3, P7, and P14.

Three-chamber social interaction
Mice were habituated to the chamber with two empty holders for 10 min. The next day, mice were placed in the middle of the chamber and allowed to explore the three chambers for a period of 5 min. During this testing period, a social object (novel mouse) was contained in one holder in one chamber and an inanimate object (lego blocks) was contained in a holder in the other chamber. Object exploration and distance traveled were tracked using ANY-maze video tracking software.

Repetition assay
A large homecage was filled 5 cm deep with wood chipping bedding and lightly tamped down to make an even flat surface. A consistent pattern of 20 identical glass marbles (15mm diameter) was evenly placed (4 cm apart) on the surface of the wood chip bedding. Mice were left alone in the testing arena for 30 min. A picture was taken before and after the test for analysis. A marble was considered buried if 2/3rds of the depth of the marble was buried.

Genotyping
Genomic DNA was extracted from the ear punches of each mouse and sent to Transnetyx for genotyping.

Analysis and statistics
All experiments were analyzed blind to the experimental group. All videos were randomized before manual scoring. Behavioral performance was recorded by a digital video camera. Videos were manually scored individually, and investigators were blind to the experimental condition and test day during all manual scoring. Data analysis and statistics were conducted using GraphPad Prism 6.00 (GraphPad software). Unpaired Student's t-tests were used for independent group comparisons. Paired Student's t-tests were used to assess differences within

groups. ANOVA followed by a Bonferroni post hoc test was used to determine conditions that were significant from each other where appropriate. Outliers were detected using Grubbs' test where $p < 0.05$. Results are displayed as mean with SEM and deemed significant when $p < 0.05$.

**Fig. S1.**

**Delayed weaning or environmental enrichment does not affect infantile amnesia.** (**A**) Behavioral schedule for CFC in infant (P17) C57BLJ mice subjected to normal (P21) or delayed (P24) weaning. The lightening symbol represents footshocks. (**B**-**E**) Freezing levels during recall tests (n=10) at Test 1 (P18), CxtB (P19), Test 3 (P25), and Test 4 (P32). (**F**) Behavioral schedule for CFC in infant C57BLJ mice. (**G**-**J**) Freezing levels during recall tests (n=10).

**Fig. S2.**

**MIA male offspring demonstrate social interaction deficits and repetitive behavior.** (**A**) Behavioral paradigm for social interaction task in adult (P63) C57BLJ mice. (**B**, **C**) Total distance (m) traveled during social interaction test (n=10). (**D**-**I**) Time (s) spent in each chamber. (**F**, **H**) Male MIA offspring spent significantly more time with the object than the novel mouse ($p < 0.05$, $p < 0.01$). (**J**) Representative diagram of behavioral paradigm for marble burying task. (**K**-**M**) Number of marbles buried during the task (n=10). (**K**) Male MIA offspring buried more marbles than controls ($p < 0.05$).

**Fig. S3.**

**Adult Ai32-FosTRAP mice show activity-dependent increases in ChR2-EYFP labelling.**
(**A**) Behavioral schedule. The syringe symbol represents 4-OHT injection 2 h after exposure.
The black lightning symbol represents foot shocks. (**B**-**E**) Quantification of ChR2-EYFP+ cells
in the (**B**) DG (N=4, n=4), (**C**) AMG (N=4, n=4), (**D**) RSC (N=4, n=4) and (**E**) PAG (N=4,
n=2-4) for an adult (P42) encoded homecage (HC), context (Cxt) and contextual fear (CFC)
memory. Exposure to a context resulted in significantly more ChR2-EYFP+ cells in the (**B**) DG
($p < 0.001$) and (**D**) RSC ($p < 0.001$). Quantification of ChR2-EYFP+ cells after CFC was
significantly higher in the (**B**) DG ($p < 0.05$), (**C**) AMG ($p < 0.001$) and the (**D**) RSC ($p <
0.001$). (**F, G**) ChR2-EYFP cell counts (N=4, n=4) in the RSC and PAG for an infant (P17)
encoded HC, Cxt and CFC memory.

**Fig. S4.**
**Infant mice show memory retention for a fear memory at P20.** (**A**) Behavioral schedule for CFC in infant (P17) C57BLJ mice. The shock symbol represents footshocks. (**B**) Freezing levels during recall test (n=9). Infant mice who received footshocks during training froze significantly (p < 0.001) more than controls. (**C**) Comparison of freezing levels during recall at 1 d (P18) or 3 d (P20) after training.

**Fig. S5.**

**Engram reactivation is increased in the DG, AMG, and RSC after natural recall at P20.**
(**A**-**N**) Histological analysis of cell counts (N=4, n=4) after recall at P20. Percentage of (**A**) ChR2-EYFP+ and (**B**) c-Fos+ cells in the DG. There was a significant increase in c-Fos+ cells in the DG after recall at P20 ($p < 0.01$). (**C**) Percentage of DAPI cells positive for both ChR2-EYFP and c-Fos. (**C**) Engram reactivation as a percentage of DAPI+ was significantly higher in the DG after recall at P20 ($p < 0.01$). Percentage of (**D**) ChR2-EYFP+ and (**E**) c-Fos+ cells in the AMG. There was a significant increase in c-Fos+ cells in the AMG after recall at P20 ($p < 0.001$). (**F**) Percentage of DAPI cells positive for both ChR2-EYFP and c-Fos. (**F**) Engram reactivation as a percentage of DAPI+ in the AMG was significantly higher after recall at P20 ($p < 0.05$). Percentage of (**G**) ChR2-EYFP+ and (**H**) c-Fos+ cells in the RSC. (**V**) There was a significant increase in c-Fos+ cells in the RSC after recall at P63 ($p < 0.05$). Engram reactivation in the RSC as a percentage of (**I**) ChR2-EYFP+ and (**J**) DAPI+. (**I**, **J**). Engram reactivation in the RSC was significantly higher after recall at P20 ($p < 0.01$). Percentage of (**K**) ChR2-EYFP+ and (**L**) c-Fos+ cells in the PAG. Engram reactivation in the RSC as a percentage of ChR2-EYFP+ (**M**) and (**N**) DAPI+.

**Fig. S6.**

**Presentation of recall cues for an infant encoded context does not result in engram reactivation at P63.** (**A**-**N**) Histological analysis of cell counts after recall at P63. Percentage of (**A**, **D**) ChR2-EYFP⁺ and (**B**, **E**) c-Fos⁺ cells in the DG and AMG (N=4, n=4). (**B**) There was a significant increase in c-Fos⁺ cells in the DG after recall at P63 (p < 0.01). (**C**, **F**) Engram reactivation as a percentage of DAPI⁺ in the DG and AMG. Percentage of (**G**, **K**) ChR2-EYFP⁺ and (**H**, **L**) c-Fos⁺ cells in the RSC (N=4, n=4) and PAG (N=3, n=4). (**H**) There was a significant increase in c-Fos⁺ cells in the RSC after recall at P63 (p < 0.05). Engram reactivation in the RSC and PAG as a percentage of (**I**, **M**) ChR2-EYFP⁺ and (**J**, **N**) DAPI⁺.

**Fig. S7.**

**Optogenetic engram reactivation of an infant encoded neutral context does not elicit freezing behavior in adult mice.** (**A**) Behavioral schedule for engram reactivation of a neutral context B in Ai32-FosTRAP mice. The black lightning symbol represents foot shocks. The syringe symbol represents 4-OHT injection 2 h after context exposure. (**B**) Memory recall in context C (engram reactivation) with light-off and light-on epochs (n=9). (**C**) Freezing for the two light-off and light-on epochs averaged. (**D**-**N**) Cell counts (N=4, n=4) after optogenetic reactivation of an infant encoded engram in adult Ai32-FosTRAP mice (Fig. 3A). (**D**, **E**) Percentage of DAPI cells positive for both ChR2-EYFP and c-Fos in the DG and AMG. (**E**) Engram reactivation as a percentage of DAPI$^+$ cells was significantly higher ($p < 0.05$) in the AMG after optogenetic stimulation of an infant encoded fear memory (S). Percentage of (**F**) ChR2-EYFP$^+$ and (**G**) c-Fos$^+$ cells in the RSC. Engram reactivation in the RSC as a percentage of (**H**) ChR2-EYFP$^+$ and (**I**) DAPI$^+$. Percentage of (**J**) ChR2-EYFP$^+$ and (**K**) c-Fos$^+$ cells in the PAG (N=4, n=3-4). Engram reactivation in the PAG as a percentage of (**M**) ChR2-EYFP$^+$ and (**N**) DAPI$^+$.

**Fig. S8.**

**Memory engrams encoded before and after the infantile amnesia period have similar engram connectivity patterns.** (**A**) Behavioral schedule for optogenetic reactivation of DG engram cells, labeled at P29, in adult Ai32-FosTRAP mice. The black lightning symbol represents foot shocks. The syringe symbol represents 4-OHT injection 2 h after training. Percentage of ChR2-EYFP+ cells labelled during training in the (**B**) DG, (**F**) AMG (p < 0.05), (**J**) RSC, and (**N**) PAG (N=4, n=4). Percentage of c-Fos+ cells in the (**C**) DG, (**G**) AMG, (**K**) RSC, and (**O**) PAG after optogenetic stimulation at P63. Engram reactivation as a percentage of (**D**, **H**, **L**, **P**) EYFP+ and (**E**, **I**, **M**, **Q**) DAPI+ cells.

**Fig. S9.**
**Infant mice can form context-specific memories that can be behaviorally updated.** (**A**) Behavioral schedule for artificial updating of an adult engram in Ai32-FosTRAP mice. (**B**) Freezing levels during recall in context B and context A (n=10). Experimental light group froze significantly more in context A (p < 0.001). (**C**) Behavioral schedule for updating paradigm in C57BLJ infant mice. (**D**) Freezing levels (n=10) during recall tests at P19 (Recall 1), P20 (CxtB), and P63 (Recall 2). Infant mice that were pre-exposed (PE) to context A froze significantly (p < 0.05) more during recall 1. (**E**) Behavioral schedule. (**F**) Freezing levels (n=9) during recall tests.

**Fig. S10.**

**Artificial updating of infant engram cells results in an above chance level of engram reactivation in the DG, AMG, RSC, and PAG.** (**A**-**N**) Histological analysis of cell counts (N=4, n=4) after context A recall in adult Ai32-FosTRAP mice. Percentage of (**A**) ChR2-EYFP+ and (**B**) c-Fos+ cells in the DG. (**C**) Percentage of DAPI cells positive for both ChR2-EYFP and c-Fos. The experimental light group showed a significant increase (p < 0.05) in engram reactivation in the DG. Percentage of (**D**) ChR2-EYFP+ and (**E**) c-Fos+ cells in the AMG. Engram reactivation in the AMG as a percentage of (**F**) ChR2-EYFP+ and (**G**) DAPI+. Percentage of (**H**) ChR2-EYFP+ and (**I**) c-Fos+ cells in the RSC. (**J**) Percentage of DAPI cells positive for both ChR2-EYFP and c-Fos. The experimental light group showed a significant increase (p < 0.01) in engram reactivation in the RSC. Percentage of (**K**) ChR2-EYFP+ and (**L**) c-Fos+ cells in the PAG. Engram reactivation in the RSC as a percentage of (**M**) ChR2-EYFP+ and (**N**) DAPI+ cells.

**Fig. S11.**
**Adult mice retain an object memory a week after acquisition.** (**A**) Behavioral schedule for novel object recognition task in adult C57BLJ mice. (**B-D**) Time (s) spent exploring each object (n=9). Adult mice spend significantly more time exploring the novel object when tested (**C**) 1 d (p < 0.05) or (**D**) 8 d (p < 0.001) after acquisition. (**E**) Time (s) spent exploring both objects during infant acquisition (n=9). (**F-H**) Time (s) spent exploring both objects during MIA offspring acquisition (n=9-10). (**I, J**) Time (s) spent exploring both objects during acquisition (n=8-9).

**Fig. S12.**
**Adult mice retain a spatial memory for the location of the escape hole on the Barnes maze task.** (**A**) Time (s) and (**B**) distance (m) taken by infant C57BLJ mice to reach the escape hole during each trial (n=18-20). (**C**) Time (s) and (**D**) distance (m) taken to first reach where the escape hole should be located during probe tests (n=9-10). (**E**) Behavioral schedule for the Barnes maze in adult C57BLJ mice. (**F**) Time (s) and (**G**) distance (m) taken to reach the escape hole during each trial (n=10). (**H-K**) Time spent in each zone during probe test (**H, I**) 1 d (p < 0.001) or (**J, K**) 8 d (p < 0.05, p < 0.01) after training. (**L**) Time (s) and (**M**) distance (m) taken to first reach where the escape hole should be located during probe tests. (**N**) Adult search strategies during testing 1 and 8 d after training (n=9-10). (**O-Q**) Heat map analysis of time spent in each zone by MIA offspring during test 8 d after training (n=9-11).

**Fig. S13.**

***Il17a-/-* MIA offspring do not show social interaction deficits or repetitive behavior.** (**A**) Behavioral paradigm for social interaction task in adult (P63) *Il17a-/-* mice. (**B**, **C**) Total distance (m) traveled during social interaction test (n=9-10). (**D**-**G**) Time (s) spent in each chamber. (**H**) Representative diagram of behavioral paradigm for marble burying task. (**I**, **J**) Number of marbles buried during task (n=10).