# First-in-human evaluation of cutaneous innate and adaptive immunomodulation by mosquito bites

David Guerrero[1,*], Hoa Thi My Vo[1,*], Chanthap Lon[3,*], Jennifer A. Bohl[2], Sreynik Nhik[3], Sophana Chea[3], Somnang Man[3,4], Sokunthea Sreng[3,4], Andrea R. Pacheco[3], Sokna Ly[3], Rathanak Sath[3], Sokchea Lay[1], Dorothee Missé[5], Rekol Huy[4], Rithea Leang[4], Hok Kry[6], Jesus G. Valenzuela[2], Fabiano Oliveira[2,$],Tineke Cantaert[1,$], Jessica E. Manning[2,3,$,#]

[1] Institut Pasteur du Cambodge, Pasteur Network, Phnom Penh, Cambodia

[2] Laboratory of Malaria and Vector Research, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Rockville, Maryland, USA

[3] International Center of Excellence in Research, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Phnom Penh, Cambodia

[4] National Center of Parasitology, Entomology, and Malaria Control, Phnom Penh, Cambodia

[5] MIVEGEC, Univ Montpellier, IRD, CNRS, 34000 Montpellier, France.

[6] Kampong Speu Provincial District, Ministry of Health, Phnom Penh, Cambodia

*co-first author

$ co-last author

# corresponding author: jessica.manning@nih.gov

**Abstract**

Mosquito-borne viruses are a growing global threat. Initial viral inoculation occurs in the skin via the mosquito 'bite', eliciting immune responses that shape the establishment of infection and pathogenesis. We aimed to cutaneous innate and adaptive immune responses to mosquito bites in individuals from endemic areas. In this single-arm, cross-sectional interventional study, we enrolled 30 healthy adult participants aged 18 to 45 years of age from Cambodia between October 2020 and January 2021. We performed 3-mm skin biopsies at baseline as well as 30 minutes, 4 hours, and 48 hours after a controlled feeding by uninfected *Aedes aegypti* mosquitos. The primary endpoints were measurement of changes in early and late innate responses in bitten vs unbitten skin by gene expression profiling, immunophenotyping, and cytokine profiling. Results revealed induction of neutrophil degranulation and recruitment of skin-resident dendritic cells and M2-macrophages in 'bitten' skin. As the immune reaction progressed over time, T cell priming and regulatory pathways were upregulated along with a shift to a Th2-driven response and CD8+ T cell activation. In accordance, participants' bitten skin cells produced less pro-inflammatory cytokines when stimulated by *Ae. aegypti* salivary gland extract. No unexpected adverse events occurred, and one patient was lost to follow-up at Day 14. These results identify key immune genes, cell types, and pathways in the human response to mosquito bites that can be leveraged to develop novel therapeutics and vector-targeted vaccine candidates to arboviral diseases.

**Graphical Abstract**

**Panel shows evolution of the human skin response to *Aedes aegypti* bites over time.**

## Introduction

With one million deaths every year, vector-borne diseases remain a major health problem worldwide[1]. In particular, arboviruses are of particular concern given a global distribution and a burden of nearly half a billion infections[2]. Several clinically significant arboviruses such as yellow fever (YFV), dengue (DENV), chikungunya (CHIKV), and Zika (ZIKV) viruses are all transmitted by *Aedes spp.* mosquitoes. These vectors are abundant in tropical and subtropical areas, but current climate change and rapid urbanization are expanding their transmission potential and increasing the at-risk population[3].

Mosquitoes first acquire an arbovirus via insertion of their proboscis into the dermis of an infected host during a blood meal. The virus then replicates in midgut cells to later disseminate and reach the salivary glands (SG)[4]. During a subsequent blood meal, the infectious mosquito inoculates virus and insect-derived molecules, including salivary components, into the dermis of a new human host. The saliva from hematophagous mosquitoes is a cocktail of pharmacologically active molecules, some of which are anti-hemostatic, angiogenic, and immunomodulatory[5-7]. Over the last couple decades, several animal and *in vitro* studies demonstrated that inoculation of vector saliva, and/or concomitant blood-feeding by an arthropod, can immunomodulate the host antiviral response in the skin and periphery[7-10]. These perturbations result in altered cytokine production profiles [11-13], promotion of recruitment of infection-susceptible cells to the bite site[14,15], stimulation of autophagy activation[5], and an induction of neutrophil infiltration to the local inflammation site[15] amongst other consequences. Mosquito-inoculated DENV or ZIKV infection in macaques led to higher or delayed viremia compared to non-vector or needle inoculation routes[16,17]. With regard to clinical pathogenesis, mice inoculated with both Semliki Forest virus and mosquito saliva had higher loads of virus RNA in the skin, earlier brain dissemination, and more frequent lethal

outcomes than animals inoculated with virus alone[12].  Taken together, arthropod-mediated viral

infection leads to increased disease progression, viremia, and mortality[8,9]. Despite this robust

evidence from animal and *in vitro* studies, there are no translational studies on the human skin

response to mosquito bites, much less those in tropical areas. Individuals are continuously exposed

to mosquito bites, possibly shaping their immune responses to mosquito salivary proteins in

contrast to those intermittently or rarely exposed[18,19]. In order to examine the dermal and epidermal

innate and adaptive immune responses to *Ae. aegypti* saliva, we compared the immune responses

before and after *Ae. aegypti* bites in skin biopsies from 30 healthy Cambodian individuals living

in an area where *Ae. Aegypti* mosquitoes are prevalent. Our study provides information on the key

genes and cell populations involved in the human immune response at different time points after

the 'mosquito bite.'

**Results**

**Controlled *Aedes aegypti* bite challenge in healthy volunteers in Cambodia**

From October 2020 to January 2021, we enrolled 30 healthy Cambodian adults at Kampong Speu

District Referral Hospital in Chbar Mon, Cambodia.  The study population comprised 13 females

and 17 males, all of whom were of Khmer (Cambodian) ethnicity (Supplementary Table 1). The

population was randomly divided in three groups to assess immune responses by different

technologies (Fig. 1). The median age was 32.5 years (IQR 22–35) and the primary occupations

were variable, although nearly half (14/30) were unemployed. In terms of mosquito exposure risk,

most participants reported living in a house (20/30), being middle income (25/30), having one to

two domestic water containers (22/30). Notably, most participants did not use larvicide (28/30) or

insecticide (19/30) in their homes.  Only 10% (3/30) reported prior dengue infection, but all 30

participants were positive for dengue anti-NS1 antibodies. All participants had detectable IgG to

*Ae. aegypti* salivary gland extract (SGE) prior to mosquito feeding, and these levels were not

significantly different by age, gender, or 14 days after the mosquito feeding (Supplementary Figure

S1). One participant did not complete Day 14 follow-up and was considered lost to follow-up.

No unanticipated adverse events occurred with the biopsies or mosquito feedings.

Figure 1. **Graphical Representation of Study Design.** Thirty Cambodian participants were

enrolled in the study. Each patient underwent a mosquito feeding (5 females *Aedes aegypti*) for

ten minutes. Each volunteer underwent a 3mm biopsy from a mosquito bite site at 30 minutes, 4

hours, and 48 hours. Distinct bite sites were biopsied at different time points. Additionally, two

3mm biopsies of normal skin (NSK) were taken from the opposite forearm. Blood was collected

to obtain serum.

**Marked clinical changes occurred in the skin post-mosquito feeding**

Exposure to five *Ae. aegypti* mosquitos resulted in 3 to 10 visible bite sites per volunteer. Upon

clinical observation, participants experienced erythema and swelling with a "bite size" reaction

that peaked at 15 minutes after feeding (Fig. 2a). The mean bite size of 4.9mm at 15 min after

feeding decreased to 4.7 mm at 30 minutes, to 2.7 mm at 4 hours and gradually decreased to 2.1mm

by 48 hours (Supplementary Table 1). Observation of representative histological sections shows

increased recruitment of cells to the dermis and epidermis at 48 hours post-bite (Fig. 2b,c). There

was no difference in bite size reaction by gender or age.

| | Pre- | Post- | | |
|---|---|---|---|---|
| | Contr | 30 | 4 | 48 |

Figure 2. **Clinical observations of skin reaction pre- and post-bite on gross examination and**

**H&E staining. a,** Skin immune responses to mosquito bites (left to right) normal skin (NSK), 30

minutes, 4 hours, and 48 hours after exposure to five *Ae. aegypti* mosquitos. **b,** Representative

cross-sectional of 3mm punch biopsies H&E stained at (left to right) 0 hours, 30 minutes, 4 hours,

and 48 hours after exposure to *Ae. aegypti* bites. Distinct bite sites were biopsied at different time

points. Scale bar is 1 mm. Black arrows indicate area of infiltrate. **c,** Magnification of infiltrate

indicated by black areas in B, scale bar 100μM.

**Changes in gene expression in *Aedes aegypti* bitten versus normal skin (NSK) evolved over**

**48 hours**

We extracted RNA and performed next generation sequencing on 40 skin biopsies from 10 participants (baseline normal skin (NSK) biopsy on the forearm, plus 3 follow-up biopsies each at 30 minutes, 4 hours, and 48 hours post *Ae. aegypti* bites on the contralateral arm). Principal Component (PC) analysis of *Ae. aegypti* bite site skin biopsies resulted in segregation of sample clusters based on the time the biopsy was taken (Fig. 3a), with PC1 and PC2 variances defining differences between NSK and 30 minutes versus 4 hours and 48 hours. While PC3 variance associated to the separation between biopsies taken at 4 hours and 48 hours, samples taken from NSK (pink circles) and 30 minutes after *Ae. aegypti* bites clustered in the same spatial area. At 4 hours post-bite, 6 of 10 samples occupied a distinct region (Fig. 3a, blue ellipse) from NSK, 30 minutes or 48 hours after *Ae. aegypti* bites. Similarly at 48 hours post-bite, 7 of 10 samples segregated into a discrete space (Fig. 3a, orange ellipse) compared to NSK, 30 minutes or 4 hours after *Ae. aegypti* bites.  These data indicate that time post-bite rather than inter-individual differences drive unique patterns in gene expression. Differential analysis of RNA expression with a threshold of two-fold ($-1 < \log2FC > 1$; False discovery rate (FDR) $< 0.05$) over NSK biopsies resulted in 709 upregulated genes post-bite compared to 31 downregulated genes (Fig. 3b, red circles). At 30 minutes, 4 hours, and 48 hours after a mosquito bite, 26, 210 and 473 genes were upregulated over NSK, respectively (Fig. 3c, Supplementary Table 2).  Conversely, one gene at 4 hours and 27 genes at 48 hours were downregulated after *Ae. aegypti* bites (Supplementary Table 2).

Upregulated genes at each time point were subjected to over-representation analysis (ORA) for enriched pathways using the Reactome database with an FDR cut off $< 0.05$. Four pathways related

to cell proliferation, cell growth and nuclear transcription were identified (FDR < 0.05) at 30

minutes after *Ae. aegypti* bites (Supplementary Table 3). At both 4 and 48 hours post-bite, the

immune nature of the host response to *Ae. aegypti* bites was evident via the revealed pathways

(Fig. 3d, Supplementary Table 3). At 4 hours post-bite, pathways such as "neutrophil

degranulation", "Interleukin-4 and Interleukin-13 signaling", "Interleukin-10 signaling",

"Interferon-γ signaling" "Chemokine receptors bind chemokines" and "Immunoregulatory

interactions between a Lymphoid and a non-Lymphoid cell" (Fig. 3d, upper chord) were

significantly overrepresented and maintained until 48 hours post-bite (Fig. 3d, lower chord).

Unique pathways discovered at 4 hours post-bite were related to "degradation of extracellular

matrix" (Fig 3d, upper chord) while "antimicrobial peptides" pathway and "adaptive immune

system processes" were present at 48 hours post-bite including "TCR signaling", "Co-stimulation

by the CD28 family", "PD-1 signaling", "MHC class II antigen presentation" (Fig. 3d, lower

chord). A heatmap with differentially expressed genes (DEG) under an FDR cutoff < 0.05 and

log2FC >1 in at least one timepoint was assembled (Supplementary Figure S2, Supplementary

Table 4), showing the dynamics of DE expression throughout the three timepoints. Fig. 3e shows

DEG driving the common pathways found on Fig. 3d.

Overall, few DEG genes were detected at 30 minutes post-bite compared to NSK biopsies. The

main upregulated gene FOSB (up 18.8x) is one of the transcription factors of the FOS family

proteins that regulate cell proliferation and differentiation (Supplementary Table 2). As the

reaction progresses over time, gene expression increased towards neutrophil recruitment followed

by signaling of both Th1, Th2, and regulatory pathways. The five most regulated genes at 4 hours

compared to NSK are KRT6C (related to keratinization; up 80.8x), CXCL8 (neutrophil-attracting

chemokine IL-8 up 75.0x), TNIP3 (inhibitor of NF-kappa-B activation; up 59.3x), IL-20

(stimulates keratinocyte proliferation and differentiation; up 59.1x) and IL-1B (pro-inflammatory

cytokine; up 45.7x) (Supplementary Table 2). The immune response evolved at 48 hours post-bite

to include adaptive immune response pathways related to T-cell priming. The five top genes at 48

hours are KRT6C (related to keratinization; up 59.6x), DEFB4A (an antimicrobial peptide; up

32.7x), GZMB (Granzyme B, up 21.1x), TCL1A (proliferation of T and B cells; up 20.3x) and

CCL18 (chemokine associated with associated with TH2 responses; up 18.5x).

Figure 3. **Differential expression gene (DEG) and pathway analysis induced by *Aedes aegypti***

**mosquito bites over time.  a,** 3D principal component analysis showing clustering of individual

biopsies taken at 30 minutes (green dots), 4 hours (blue dots), 48 hours (orange dots) post-bite or

normal skin (NSK; pink dots). Blue ellipse indicates clustering at 4 hours, while orange ellipse at

48 hours. **b,** Volcano plot comparing DEG from skin biopsies post-bite with NSK (significant

genes with -1 < log2FC >1 and FDR < 0.05 are shown in red). **c,** Venn diagram summarizing the

relationships between significantly upregulated genes over NSK (log2FC >1 and FDR < 0.05) and

time post-bite. **d,** Chord plot shows significantly upregulated genes at 4 hours and 48 hours post-

bite over NSK (left side of chord plot) and overrepresented pathways (right side of chord plot).

Outer left ring shows genes and color range displays log2 fold change (left, key at bottom) or

Reactome terms (outer right ring). Chords connect gene names with Reactome term pathways. **e,**

Heatmap (scaled to log2FC) displaying a list of DEGs based on the common pathways

overrepresented at 4- and 48-hours post-bite.

**Early innate immune response in the skin is characterized by increased proportions of DCs**

**and M2 macrophages and a decreased proportion of NK cells**

To confirm and extend the gene expression analysis, dissociated cells from NSK and skin biopsies

taken at 30 minutes and 4 hours after exposure to mosquitos, were characterized for innate and

adaptive immune subsets by flow cytometry (Supplementary Figure S3, Supplementary Table 5).

No significant changes in cell populations were detected between NSK and 30 minutes post-bite,

mirroring the transcriptomic analysis of DEG.  At 4 hours post-bite, the proportion of dendritic

cells (DC), and M2 macrophages increased, while the proportion of mast cells, natural killer (NK)

and T-cells decreased, when compared to NSK and 30 minutes post-bite (Fig. 4a). The frequencies

of DC subsets increased at 4 hours post-bite compared to NSK and 30 minutes (Fig. 4 b-e). Within the DC subsets, Langerhans cells (CD45+CD3-CD207+) increased at 4 hours compared to NSK (Fig. 4d). A decreasing, though non-significant trend in the NK cell population frequency was observed over time (Fig. 4f). Regarding the macrophage population, M2-skewed macrophages (CD45+CD14+CD163+) frequency increased after 4 hours post-bite as well as their expression of activation markers (CD16/CD69) compared to NSK and 30 minutes (Fig. 4 h-j). Frequencies of activated M2 macrophages, and frequency of Langerhans cells positively correlated with clinically observed changes in bite size at 30 minutes (Supplementary Figure S4 a, b). In order to evaluate if changes in total leukocytes ($CD45^+$) were affecting the reported frequencies, we analysed the changes of the innate cell population within total live cells. Here, similar results were observed (Supplementary Figure S5). Unfortunately, we did not include specific neutrophil markers to our flow cytometry panels. Alternatively, when we gated on CD45+CD11b+CD14-CD56-CD117- cells, that comprises mostly of neutrophils, we detect a significant "neutrophils" accumulation at 4 hours post-bite compared to NSK (Supplementary Figure S6). The relative proportion of $CD4^+$ and $CD8^+$ T cells did not change over time. Taken together, the results suggest that immune cell infiltration and migration occurs in local skin tissue within 4 hours post-bite and is characterized by increased frequencies of DCs, M2 macrophages and neutrophils, but decreased frequencies of NK cells.

Figure 4. **Early innate immune responses in the skin are characterized by increased proportions of DCs and M2 macrophages, and a decreased proportion of NK cells at 4 hours post-bite**. **a,** Pie charts showing changes in the frequencies of skin immune cells during early innate immune response to mosquito bite at 30 minutes and 4 hours after exposure. Frequencies are reported as percentages of total leukocytes (CD45$^+$) **b,** total frequency of dendritic cells calculated as the sum of Langerhans (CD207$^+$), dermal (CD1c$^+$) and plasmacytoid (CD123$^+$) DCs. **c,d,** Frequency of plasmacytoid dendritic cells and Langerhans cells populations reported as percentage of total CD45$^+$ cells shows significant increase of the later, at 4 hours post mosquito

bites. **e,** Frequency of activated dermal DCs (CD1c$^+$CD69$^+$) from the total dermal DCs population

(CD1$^+$). **f,** Frequency of NK cells (CD56) reported as percentage of total CD45+ cells. **g,**

Frequency of cytotoxic NK cells (CD56+CD16+) as percentage of total NK cells. **h-j**, Frequency

of M2 macrophages and MFI of their expressed activation markers CD16 and CD69. Statistical

analysis were performed with Chi-square test (**a**) and Friedman + Dunn's multiple comparisons

test (**b-j**).  Bars indicate median and interquartile range. N=10

**Changes in the T cell compartment are observable at 48 hours after mosquito 'bite'**

Changes in skin T cell populations were assessed using NSK and biopsies taken at 48 hours post-

bite. We found that the frequency of CD8+CD25+ and CD8+PD-1+ T-cells increased at 48 hours

post-bite, compared to the NSK (Fig. 5a). Frequencies of CD8+ T cells correlated inversely with

the bite size at 48 hours (Supplementary Figure S4c). The frequency of effector memory cells re-

expressing CD45RA (T$_{EMRA}$) CD8 T cells (CD8+CCR7-CD45RA+) decreased at 48 hours (Fig.

5b). In the CD4+ T cell compartment, we observed a decrease in the frequency of Th1/Th17

(CD4+CCR4-CXCR3+) cells at 48h compared to NSK and an increase in the frequency of

Th2/Th17 (CD4+CCR4+CXCR3-) cells (Fig. 5c). The frequency of CD4+ effector memory T

cells (T$_{EM}$) (CD4+CCR7-CD45RA-) increased at 48 hours compared to NSK, while the frequency

of T$_{EMRA}$ cells (CD4+CCR7-CD45RA+) decreased (Fig. 5d). Results were similar when analyzing

the changes of T cell frequencies within all live cells (Supplementary Figure S7).  Interestingly,

no changes were observed in the frequencies of skin-resident CD4 (CD4+CD103+CLA+) or CD8

(CD8+CD103+CLA+) T cells (Supplementary Figure S8). At 48 hours post-bite, these results

show an increase in the frequency of activated CD8 T cells in addition to a shift from Th1/Th17

towards Th2/Th17 polarization of CD4 T cells.

Figure 5. **Changes in the T cell compartment are observable at 48 hours after mosquito exposure. CD8 T cells exhibit a high activation phenotype, while CD4 T effector memory cells shift from a Th1/Th17 phenotype to a Th2/Th17 phenotype. a,** Increased frequency of CD25+ and PD1+ CD8 T cells at 48 hours post mosquito exposure. **b,** Frequencies of CD8 T cells naive (CCR7+CD45RA+), central memory (CCR7+CD45RA-), effector memory (CCR7-CD45RA-) and TEMRA (CCR7-CD54RA+) frequencies. **c,** decreased Th1/Th17 and increased Th2/Th17 effector memory CD4 T cell compartment at 48 hours. **d,** CD4 T cells naive (CCR7+CD45RA+), central memory (CCR7+CD45RA-), effector memory (CCR7-CD45RA-) and TEMRA (CCR7-CD54RA+) frequencies. Statistical analysis of 8 samples dedicated to this time point was performed with Wilcoxon signed-rank test. Bars indicate median and interquartile range.

**"Bitten" human skin cells stimulated by *Aedes aegypti* saliva produce significantly less pro-inflammatory cytokines**

Next, we sought to assess if *Ae. aegypti* saliva impacted cytokine production by skin cells taken 48 hours post-bite. We measured pro- and anti-inflammatory cytokines released into the media after a stimulation assay with and without *Ae. aegypti* SGE. Both NSK and SGE-treated cells showed low secretion of cytokines in the media (Supplementary Figure S9). However, we observed a significant reduction in the levels of Interleukin 2 (IL-2) and Interferon gamma (IFN-γ) in the cells treated with both SGE and PMA/Ionomycin compared to the cells treated with PMA/Ionomycin alone (Fig. 6, Supplementary Figure S9). The levels of other pro-inflammatory cytokines like TNF-α, IL-17A and IL-17F showed no significant differences (Fig. 6). Similarly, no differences in the regulatory cytokine IL-10 was observed (Fig. 6). This data suggests that SGE has a dampening effect on the production of pro-inflammatory cytokines IL-2 and IFN-γ under stimulatory conditions in "bitten" human skin cells.

Figure 6. **Human skin cells obtained from *Ae. aegypti* bitten skin stimulated by *Ae. aegypti* salivary gland extract (SGE) produce significantly less pro-inflammatory cytokines (IL-2 and IFN-γ).** Cytokine production measured in skin cell culture supernatant after treatment with

PMA/Ionomycin in the presence or absence of SGE. Skin dissociated cells were seeded on round

bottom 96-well plates (50.000 cells/well) and treated with SGE (10 µg/mL) or PBS for 24 hours.

Cells were stimulated with PMA (0.1µg/mL) and Ionomycin (1µg/mL) or left unstimulated for the

last 6 hours of the culture.  Statistical analysis was performed with Wilcoxon signed-rank test. N=9

individuals.

**Discussion**

Arboviruses enter the host through the skin when the arthropod injects its saliva and other

components into the dermis while taking a bloodmeal.  Data from mice and macaques suggest that

the unique dual presence of virus and saliva orchestrates an immunological cascade of events that

influence the establishment and course of infection[12,16,17].  To translate these findings to humans,

the current study assessed the dynamics of the cutaneous immune response to arthropod bites from

a clinically relevant vector, the *Ae. aegypti* mosquito, in humans from arboviral-endemic areas

with lifelong exposure to *Aedes* mosquitos. At 4 hours post-bite, we observed upregulation of

genes involved in neutrophil recruitment and increased frequencies of dendritic cells and M2

macrophages. As the immune response progressed over two days, we observed an increased

frequency of activated, PD-1+CD8+ T cells, a polarization towards Th2/Th17 CD4+ T-cells, and

upregulation of IL-10, IL-4, and IL-13 and IFN-γ related pathways. Interestingly, these human

skin cells post-bite produced less pro-inflammatory cytokines when stimulated by Ae. aegypti

SGE.

A mosquito 'bite' results in minor tissue damage by the proboscis, injection of mosquito saliva

into the skin, and introduction of both mosquito and host microbiota into the skin. In this study,

we did not parse out the separate impact of each process on the skin immune response. Alterations

in skin immunity due to injury from the punch biopsy itself were mitigated by ensuring biopsy

sites were at least 10 to 20 mm away from one another and were not taken directly at the site of

the needle administration of subcutaneous lidocaine. The effect of local anesthetic was mitigated

by the case-control design that each person's anesthetized, bitten skin was compared to

anesthetized, unbitten skin, presumably negating any immediate cutaneous immune effects

attributed to lidocaine, for which the literature is sparse.

Mosquito bites cause a local cutaneous reaction with wheal and flare manifestation[20,21].

Concordantly, the bite size reaction peaked after 15 minutes post-bite in this endemic study

population. Few changes were detected in gene expression or influx and activation of neutrophils,

dendritic cells subsets and macrophages at the 30 minutes post-bite time point. Upregulated genes

were related to top-level processes such as signal transduction. We did not design this study to

assess the role of pre-stored immune mediators such as histamine (and others) in the development

of edema and their modulation in the immune response. Indeed, mosquito bites can activate

cutaneous mast cells, which are prominent histamine-storing cells, and induce their degranulation.

The release of histamine and other mediators leads to vasodilation, fluid accumulation at the bite

site, and neutrophil influx, which were all ultimately observed here[20,21].

As the immune reaction progressed, the mosquito 'bite' induced degradation of the extracellular

matrix (ECM), and a strong influx of dendritic cells and macrophages at 4 hours post-bite,

consistent with previous animal observations[22]. We observed several disintegrin and

metalloprotease with thrombospondin type I motifs (ADAMTS) were upregulated, mostly at 4

hours post-bite, that can be associated with ECM degradation. Similarly, granzyme B (GZMB)

upregulated at 48 hours, despite its effects on inducing inflammation it may also be involved in

ECM degradation using as substrate fibronectin, vitronectin, laminin and others[23,24]. Saliva from

blood-feeding arthropods contains anti-hemostatic proteins that prevent coagulation and promote

vasodilation, leading to increased leukocyte trafficking[20,25]. These processes – extracellular matrix

degradation and vasodilation – can facilitate viral infection by disrupting vascular functions, as is

the case of sialokinin, a mosquito salivary protein that hinders endothelial barrier integrity[26]. In

addition, elevated leukocyte trafficking can facilitate viral replication by augmenting the

proportion of susceptible cells to the local inoculation site[12,15,26]. Indeed, as most leukocyte subsets

are permissive to infection by most arboviruses, they play a dual role in the infection process: both

as replication targets/dissemination vehicles and as the first line of defense. Indeed, the increased

presence of activated M2 macrophages and dendritic cells at 4 hours post-bite highlight the cellular

trafficking role initiated by 'the bite' that can ultimately result in viral dissemination. The observed

increase in CD163+CD14+ cells at 4 hours is what we believe to be an initiation of the

differentiation of monocytes and macrophages into M2-skewed macrophages given that full

macrophage polarization does not occur as early as 4 hours [27–29]. Upregulated chemokines such as

CCL2 and CCL7, attracting mainly monocytes, DCs and T cell, may be participating in this

recruitment[30]. M2-polarized macrophages produce anti-inflammatory cytokines, mediate wound

healing processes, and support Th2 responses [31]. Moreover, activation of M2 macrophages

positively correlated with the bite size in our study. Upon stimulation, DCs upregulated CD69,

preventing them from exiting the tissue and improving the likelihood of antigen internalization.

Subsequent CD69 downregulation promotes Sphingosine-1-Phosphate (S1P1) leading to DC

migration to the lymph nodes[32]. The CD69-S1P1 mechanism of cell retention in tissue is also well-

described for resident T cells,[33] whose role in saliva-mediated immunity is not yet known.

*Ae. aegypti* mosquito saliva has a clear impact on neutrophil influx and degranulation. Two recently identified *Ae. aegypti* salivary proteins (NeSt1 and AgBr1) that attract and stimulate neutrophils at the 'bite site,' changing the immune microenvironment and leading to increased early ZIKV replication and enhanced pathogenesis[14,15]. Here, at 4 hours post-bite, chemokines CXCL2, CXCL3, CXCL5, and especially CXCL8, plus the cytokine IL-1B were upregulated returning to levels comparable to NSK at 48 hours post-bite. CXCL2-5 chemokines are responsible for attracting neutrophils and exhibiting angiogenic properties[34]. Pro-inflammatory responses through inflammasome activation and IL-1B production can trigger a sustained neutrophilic response[12]. Microbiota egested during infected sand fly bites contributed to IL-1B production[35], but we were unable to assess the effects attributed to vector microbiota in this study. On the other hand, IL-20 was upregulated at 4 hours post-bite. IL-20 is most active at epithelial sites downregulating IL-1B resulting in deactivation of neutrophil functions such as phagocytosis, granule exocytosis, and migration[36,37]. Increased gene expression for neutrophil recruitment and degranulation at 4 hours post-bite in our Cambodian study population further supports the timing of neutrophil influx seen 3 hours post-bite in bite-experienced C57BL/6 immunocompetent mice[12]. How the timing of neutrophil influx would differ in a bite-naïve population is a question that remains to be answered as none of the participants would be considered naïve. Moreover, studies of 'infected bites' would clarify the role of both 'neutrophil influx' and extracellular matrix degradation in facilitating pathogen establishment[12,38], but currently approved human challenge models deliver DENV via needle inoculation, not via mosquito bite[39].

As the immune response transitioned from innate to adaptive, a strong T-cell signature emerged at 4 and 48 hours emerged after the mosquito 'bite'. CD8+ T cell activation was confirmed by increased expression of CD25, an early activation marker. Skin CD8+ T cells also upregulate PD-

1 expression, an immune-checkpoint receptor that is induced by yet can also inhibit T cell activation. PD-1 primarily inhibits effector function acquisition by T cells, the loss of sensitivity after T cell receptor signaling and modulation of cytotoxicity, and inhibition of TNF-α and IL-2 production along with T-cell expansion[40,41]. Interestingly, highly activated CD8+ cells have been observed in skin from dengue shock syndrome patients[42,43]. Here at 4 and 48 hours post-bite, CD4+ T cells had a skewed phenotype to Th2-like cells and IL-10, IL-13, and IL-4 related pathways were upregulated compared to NSK. CCL18, upregulated at 48 hours post-bite, can be expressed by M2 macrophages and is linked to increased IL-4, IL-10, and IL-13, especially in the presence of histamine[44]. Indeed, initial animal experiments also pointed to a Th2-mediated allergic inflammatory response following mosquito bites or inoculation with SGE[13]. In case of arbovirus infection, virus will replicate locally in the skin, and the local skin immune responses will direct the course of the systemic host immune response. At 48 hours post-bite, arboviruses will also have disseminated to draining lymph nodes[45]; hence , the contribution of the localized Th2 polarization at this time point on disease outcome remains to be determined.

At 48 hours post-bite, other upregulated genes such as CD28, ICOS, and CTLA-4 further suggest a regulatory role of the CD4+ compartment in controlling the local immune responses to *Ae. aegypti* saliva. Th2 polarization is essential to avoid excessive collateral damage from an exacerbated pro-inflammatory response, especially in a situation where the skin immune cells are repeatedly exposed to the same type of insult, as is the case with individuals living in tropical areas of mosquito abundance. Blood DC antigen-2 (CLEC4C), upregulated at 48h compared to NSK, is thought to signal through BDCA-2 to inhibit activation of the NF-κB pathway, inhibit production of type I IFNs, and block TRAIL-mediated cytotoxic activity[46]. This may represent a possible mechanism that differentiates chronically exposed individuals compared to naïve individuals.

To consider other potential mechanisms that immunomodulate the microenvironment, we noted

increased expression at 48 hours post-bite of ficolin-1 (FCN1), a complement lectin pathway

protein that may bind and opsonize mosquito microbiota or salivary components, then activate the

complement pathway[47]. Another C-type lectin receptor LSECtin (CLEC4G, increased at 48 hours

compared to NSK) negatively regulates T cell-mediated immunity and may be also linked to

apoptotic cell clearance by macrophages[48]. Of note, the upregulation of CXCL10 chemokine and

its receptor CXCR3 are linked to their major roles of inflammatory reactions, T cell migration, and

IFN-γ in the saliva-mediated process[49]. Besides chemokines, the top expressed molecule was an

antimicrobial peptide hBD-2 (DEFB4A), produced by neutrophils and keratinocytes. DEFB4A is

a marker of inflammation absent in normal skin and responsible for recruitment of memory T cells,

immature DCs, mast cells and neutrophils[50,51].

In this study, the presence of mosquito saliva inhibited pro-inflammatory cytokine production by

'bitten' human skin cells after *in vitro* re-stimulation with SGE. One consideration is that we used

SGE, rather than mosquito saliva as forced salivation did not provide sufficient material, but our

preparations minimize the cellular components of the salivary glands. The cytokine-dampening

effect may partly result from the interaction between salivary components and key checkpoints on

different signaling pathways. One example includes the salivary protein LTRIN known to interfere

with the Nf-kB pathway by binding to LTβR, thus hindering its dimerization and leading to

decreased production of inflammatory cytokines[11]. On the other hand, skin immune tolerance to

constant mosquito exposure may also explain the reducing effect of SGE on cytokine production

as seen here in our bite-experienced participants[12,19,52].

Our results refine and extend the knowledge obtained from mouse models of mosquito biting. For

example, in mouse models, increased frequencies of dendritic cells, total macrophages and

neutrophils have been detected after mosquito bites [21,53,54]. In our human participants, we also

observe an increase in the frequency of these cells, indicating that the mouse models of mosquito

biting recapitulate the early innate events in human skin. However, we expand these data by

showing that the observed increased frequencies are due to increases in M2-like macrophages,

which are activated, and increases in Langerhans cells at 4 hours post-bite in our human

participants. Additionally, we substantially increased our knowledge on human T cell responses

to mosquito saliva by observing CD8 T cells exhibiting a high activation phenotype, while CD4 T

effector memory cells shift from a Th1/Th17 phenotype to a Th2/Th17 phenotype 48 hours post-

bite. Saliva or salivary proteins from ticks and sand flies have shown ability to inhibit proliferation

of both murine and human cells.[55,56] We here confirm the inhibitory effects of SGE to skin cells

recruited during a mosquito bite. Nevertheless, as previously showed in individuals from an

endemic area of leishmaniasis,[57] the immune responses to saliva of arthropods also triggers a pro-

inflammatory response highlighted by the upregulation of pathways related to interferon genes.

This mix of anti- and pro-inflammatory responses need to be further evaluated in animal models

of mosquito bite exposure as well as clinical field settings.

With the current rise of vector-borne diseases around the world, future clinical development of

vector-targeted strategies for both known and unknow vector-borne diseases will require a

fundamental understanding of how vector saliva modulates the cutaneous immune response of

humans. Translational studies, such as the one presented here, significantly expand our current

knowledge on human skin immunity and the immune response orchestrated by a 'mosquito bite'.

These newly identified genes, cell types, and processes of human cutaneous immunity can be

leveraged in the development of vector-targeted vaccine candidates or novel therapeutics such as

those targeting the reversibility of functional macrophage polarization or chemokine blockades

that may have application in other inflammatory disorders[58]com. Such new pathogen-agnostic

strategies have shown promising results in recent years from mosquito saliva-based vaccinations

to topical applications like imiquimod post-bite that activated skin macrophages in order to limit

pan-viral activity[59–62]. Building on these findings, the next generation of countermeasures against

vector-borne disease will require nuanced and comprehensive understanding of skin immunity to

both pathogen and vector saliva alike.

**Materials and methods**

**Study Enrollment and Procedures**

From October 2020 to January 2021, we enrolled thirty Cambodian adults at Kampong Speu

District Referral Hospital. Eligible participants were healthy adults between the ages of 18 and 45

years old who agreed not to use topical or oral antihistamines or steroids for the duration of the

study. Participants were excluded if they were pregnant or reported a history of severe allergic

reaction to mosquito or other insect bites, use of medications that affect blood clotting, or history

of significant scarring such as keloid formation (full criteria found on

https://clinicaltrials.gov/ct2/show/NCT04350905). All participants provided written informed

consent before enrollment, and the study was conducted in accordance with the provisions of the

Declaration of Helsinki and Good Clinical Practice guidelines.

On Day 0, five starved uninfected *Ae. aegypti* female mosquitos were selected from a mosquito

colony dedicated to human feeding studies in the Malaria and Vector Research Laboratory at the

National Center for Parasitology, Entomology and Malaria Control. Mosquitos fed through the

disposable mesh of a feeding device placed on participants' ventral forearm skin for 10 minutes.

Redness and swelling in millimeters were assessed immediately, 15 minutes, 30 minutes, 4 hours

and 48 hours after feeding. Redness and swelling were measured in millimeters at 15 minutes, 30

minutes, and 4 hours.

A total of five 3-mm skin punch biopsies were performed on each individual: one biopsy from

bitten skin per time point (30 minutes, 4 hours, and about 48 hours after feeding is completed) for

a total of three 'bitten' biopsies, and two from NSK collected at baseline. Biopsy sites were

selected based on the 3 bite sites with the largest diameters at 30 minutes after feeding. We tattooed

the area surrounding those 3 bite sites with a permanent marker and randomly picked one of the

three distinct sites for 30 min, 4 hours and 48 hours biopsies. Less than 1cc of 0. 2% lidocaine

without epinephrine was delivered via 30-G needle each chosen 'bite' at the time of biopsy and 3-

mm biopsy was performed on 'bite' adjacent to needle puncture. For feasibility of skin sample

analysis, each participant was assigned to 1 of 3 technical modality cohorts: 1)

immunohistochemistry, 2) RNASeq, and 3) flow cytometry with ten people in each cohort.

Additionally, blood was collected for serological and cellular analyses at baseline and 14 days

after feeding.

**_Aedes aegypti_ SGE preparation**

Salivary glands from _Ae. aegypti_ were dissected and collected in 200µl of PBS 1x (1pairs/1µl of

PBS 1x) in Low protein binding centrifuge tubes. Homogenization was done in an Ultrasonic

SONIFIER® (BRANSON) for 30 seconds on each side of the tube. The homogenates were then

centrifuged at 25,000 × g at 4 °C for 3 minutes, and the supernatant was collected and quantified

using Pierce™ BCA Protein Assay Kit (ThermoFisher Scientific, Cat# 23225) in an xMark™

Microplate Spectrophotometer (BIO-RAD).

**_Aedes aegypti_ salivary IgG enzyme-linked immunosorbent assays**

*Ae. aegypti* SGE was diluted to a concentration of 2 ug/ul in carbonate-bicarbonate buffer

(Millipore-Sigma) and used to coat 96-well Immulon 4x HB plates (Nunc) overnight at 4°C. Plates

were blocked in a solution of TBS 4 % Bovine serum albumin 0.025 % Tween 20 (Millipore-

Sigma). Each plate included sera (1:200 dilution) from all time points, a negative control (non-

reactive human sera to SGE), an internal control (a pool of positive sera to SGE with OD set to

0.25 at 450nm) and blank wells. The controls were chosen accordingly from a stored bank of

human sera, well-characterized by reactivity to the saliva of various vectors, at the Laboratory of

Malaria and Vector Research. Plate-to-plate normalization was done by multiplication of all OD

values by a correcting factor to achieve the preset internal control OD 0.25 at 450nm. Secondary

antibody [goat anti-human IgG alkaline phosphatase (1:10000 in blocking solution; Millipore-

Sigma). A solution of p-nitrophenylphosphate (Millipore-Sigma) was used as substrate and OD

measured at 450nm in arbitrary enzyme-linked immunosorbent assay (ELISA) units and reported

in arbitrary ELISA units after correction stated above.

**Dengue IgG enzyme-linked immunosorbent assays**

The PanBio® Dengue Indirect IgG ELISA (product code 01PE30) was performed according to

the manufacturer's instructions. Two microtiter plates were supplied, one containing stabilized

dengue-1 through dengue-4 (antigen plate) and the other containing either anti-human IgM or anti-

human IgG bound to separate wells (assay plate). Peroxidase-labeled anti-dengue virus

monoclonal antibody (125 μl/well) was added to the antigen plate to solubilize the antigens and

form antibody-antigen complexes. Concurrently, 100 μl of patient serum was diluted 1:100 in the

provided diluent and added to each well of the assay plate containing either bound anti-human IgM

or bound anti-human IgG, and human IgM or IgG in the patient's serum was captured. The plates

were incubated for 1 h at room temperature (antigen plate) or 37°C (assay plate); then the assay

plate was washed, and 100 μl of antibody-antigen complexes per well was transferred from the

antigen plate to the assay plate. These complexes were captured by dengue virus-specific IgM or

IgG during incubation for 1 h at 37°C. The plate was washed, and bound complexes were

visualized through the addition of 100 μl of tetramethylbenzidine substrate per well. After 10 min,

the reaction was stopped by the addition of 100 μl of 1 M phosphoric acid per well, and the strips

were read at 450 nm with a microtiter plate reader. Positivity was determined by comparison to

the IgM and IgG reference sera provided (cutoff calibrators). A positive sample was defined as

having a sample/calibrator absorbance ratio of ≥1.0, and a negative sample was defined as having

a ratio of <1.0.

**Formalin preservation and H&E staining of skin biopsies**

Biopsies were placed in 7ml prefilled 10% formalin vials (Azer Scientific) and stored at 4°C until

paraffin embedded. Unstained slides were cut from paraffin blocks and deparaffinized in xylene,

then hydrated through graded alcohols up to water, placed in Carazzi's hematoxylin, washed in

tap water, and put in one change 95% ethanol.  Next, they were placed in eosin-phloxine solution

and then run through graded alcohols to xylene.  After xylene, the stained slides were than a cover

slip was mounted using Permount Mounting Media (Fisher Scientific).

**RNA extraction, RNA-seq library preparation, sequencing and analysis.**

Biopsies were stored in DNA/RNA Shield (Zymo research) and kept in 4 °C until processing. Skin

was homogenized using 3mm zirconium beads (Biolink laboratories) and a Magnalyser bead

beater (Roche Diagnostics®). After homogenization total RNA was loaded at InnuPure® C16

touch (AnalitikJena) for magnetic separation of total RNA using an InnuPREP® RNA Kit–IPC16

(AnalitikJena). Quality of extractions was verified on a 4200 TapeStation (Agilent) and all samples

had RIN numbers above seven. RNA sequencing was performed by Novogene. Briefly, mRNA

was enriched by Poly (A) capture and RNA libraries were prepared using the NEBNext Ultra II

Directional RNA Library Prep Kit for Illumina (New England Biolabs) and sequenced on a

NovaSeq 6000 (Illumina) generating paired end reads at 150 bp length. After removing low quality

reads (Trimommatic 0.39)[63], mapping to the human genome (hg38; GCF_000001405.39) and

gene-level quantification were generated by RSEM 1.3.2[64], Bowtie 2-2.4.2[65] and SAMtools

1.11[66]. Normalization and differential gene expression (DGE) analysis were performed using

DESeq2 1.34.0[67], employing the variance stabilizing transformed function (VSN 3.62.0)[68] to the

count matrix, that was designed to account for paired samples enabling expression analyses with

normal skin samples from the same subject as reference. Approximate Posterior Estimation for

GLM (Apeglm 1.16.0) method was used to calculate fold change of DEGs[69]. DEGs were

considered significant if apeglm-estimated shrunken log2fold change was <1 or >1 and with a false

discovery rate <0.05 (Benjamini–Hochberg multiple testing correction method[70]). PlotPCA3D

0.0.1(twbattaglia/btools) was used to generate tri-dimensional principal component analysis plots

on DESeq2. EnhancedVolcano 1.12.0 was used to generate volcano plots[71]. GOplot 1.0.[72] was

used to generate chord plots using data obtained from packages clusterProfiler 4.2.0[73] and

ReactomePA 1.38.0[74]. Pathways with a Benjamini–Hochberg corrected P value <0.05 and were

considered as enriched, selected pathways were mapped on chord plots[70]. Heatmaps were prepared

using package pheatmap 1.0.12.[75] and dendsort 0.3.4[76]. All DEG analyses were performed using

R 4.1.0[77].

**Skin biopsy dissociation**

Biopsies were transported in RPMI media (Sigma-Aldrich) at 4 °C. Whole skin dissociation kit

(Miltenyi®) was used to perform the biopsies dissociation. An enzyme mix consisting of enzyme

D, enzyme A and buffer L was prepared according to the manufacturer's specifications. Single

punch biopsies were incubated in 400 µl of dissociation mix, at 37°C – 5% CO2 overnight.

Following incubation, each sample mix was transferred to a Falcon round-bottom tube with cell-

strainer cap (40-75um). Using a rubber pestle and the cell-strainer cap as a mortar, biopsies where

gently macerated to increase cell recovery yield. 500 µl of cold RMPI was used to rinse caps.

Samples were then centrifuged at 1500 rpm 4°C for 10 minutes and the pellet resuspended in 500

µl of cold RPMI. Dissociated skin cells were counted using a pre-programmed setting in a

Countess™ II FL Automated Cell Counter.

**Flow cytometry**

Dissociated cells were seeded on round bottom 96-well plates and washed with 200 µl of

ThermoFisher's PBS buffer 1x (1500 rpm, 4°C for 5 minutes). Twenty µl of Biolegend Zombie

Aqua fixable viability stain (1:500) was added and incubated for 20 minutes in the dark at 4°C.

Excess viability stain was washed away with 200 µl of PBS/BSA/EDTA buffer (1500 rpm, 4°C

for 5 minutes), and the pellet resuspended and incubated for 10 minutes at 4°C in 4 µl of

Biolegend's FcR blocking antibody (1:10). Cells were then incubated for 30 minutes at 4°C with

surface staining antibody master mix. After washing free unbound antibody with 200 µl of PBS

(1500 rpm, 4°C for 5 minutes), cells were fixed with 100 µl of Biolegend's fixation buffer 1x for

20 minutes at 4°C in the dark. Following fixation, cells were permeabilized and washed with 100

µl of Biolegend perm-wash solution 1x (1500 rpm, 4°C for 5 minutes) and the pellet resuspended

and incubated for 30 minutes at 4°C in intracellular staining antibody master mix. Stained cells

were placed in 200 µl of PBS/BSA/EDTA 1x and visualized on a BD Biosciences FACS Aria II

using FACS Diva software. Analysis was performed using BD Biosciences FlowJo software.

Antibody panels are described in supplemental material (Supplementary Table 5).

**In vitro SGE stimulation and Multiplex cytokine screening**

Skin dissociated cells were seeded on round bottom 96-well plates (50.000 cells/well) and treated with SGE (10 µg/mL) or PBS for 24 hours. Cells were stimulated with PMA (0.1µg/mL) and Ionomycin (1µg/mL) or left unstimulated for the last 6 hours of the culture. The supernatants were collected and stored at -20°C. Cytokines in the cell culture supernatants were quantified using Human Th Cytokine Panel (12-plex) (Cat. 741027, BioLegend) according to the manufacturer's instructions. All samples in the experiment were measured in duplicate. Briefly, undiluted supernatants were incubated with beads coated with capture antibodies specific for IL-2, 4, 5, 6, 9, 10, 13, 17A, 17F, 22, IFN-γ and TNF-α for 2 hours at room temperature on shaker. After incubation, beads were washed and incubated with biotin-labeled detection antibodies for 1 hour, followed by a final incubation with streptavidin-PE for 30 minutes at room temperature on shaker. Beads were washed and re-suspended with washing buffer. Beads were analyzed by flow cytometry using a FACS Canto cytometer. Analysis was performed using the LEGENDplex analysis software v8.0, which distinguishes between the 12 different analytes on basis of bead size and internal dye.

**Statistical analyses**

Flow cytometry and multiplex cytokine screening data were analyzed using the non-parametric-based tests, Wilcoxon signed-rank and Chi-square test. Correlation analysis was performed with the Pearson's correlation coefficient. Bars indicate median and interquartile range, and p-values *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$. Statistical analysis of RNAseq data is described above.

**Acknowledgements**

We would like to thank the study participants in Kampong Speu, Cambodia. We would also like to acknowledge Noemi Kedei and Maria Hernandez for their assistance in histology and staining as well as Jon Fintzi for statistical review of the clinical protocol.

## Funding

This work was funded by the Division of Intramural Research at National Institute of Allergy and Infectious Diseases at the National Institutes of Health. T.C. is a HHMI-Wellcome international research scholar (208710/Z/17/Z). D.M is funded by the French National Research Agency (ANR-17-CE15-00029) . D.G. is supported by a Calmette-Yersin PhD scholarship funded by the Institute Pasteur International Network.

## Competing interests

The authors declare no competing interests.

## Author contributions

Conceptualization: F.O., J.E.M., T.C., J.G.V. Methodology: D.G., H.V, S.N., S.C., S.L., J.A.B., F.O., J.E.M.. Investigation: D.G., H.V, J.A.B, S.N., S.C., S.L., A.P., J.G.V., F.O., J.E.M. Visualization: D.G., F.O., J.A.B. Funding acquisition: F.O., J.E.M. Study enrollment and clinical procedures: C.L, S.L., R.S, S.S., S.N., J.M, H.K, R.L. Project administration: C.L., T.C., F.O., J.E.M., Supervision: T.C., D.M, F.O., J.E.M., C.L, R.L., H.R, H.K.  Writing original draft: D.G., H.V., T.C., F.O., J.E.M., J.A.B. Writing review and editing: T.C., D.M., C.L., F.O., J.A.B., J.E.M.

All authors contributed to the article and approved the submitted version

## Data availability

The Illumina raw reads (fastq files) were loaded at the NCBI Sequence Read Archive with accession numbers SRR18215008 to SRR18215047 and deposited in GenBank under bioproject accession code PRJNA812009. All data generated or analyzed during this study are included in this published article (and its supplementary information files). Clinical trial data is

631    available at https://clinicaltrials.gov/ct2/show/NCT04350905.  Flow cytometry data can be made

632    available upon reasonable request directly to the authors.

**Bibliography**

1.    WHO. A global brief on vector-borne diseases. *www.who.int* (2020).

2.    Pierson, T. C. & Diamond, M. S. The continued threat of emerging flaviviruses. *Nat. Microbiol.* **5**,

796–812 (2020).

3.    Ryan, S. J., Carlson, C. J., Mordecai, E. A. & Johnson, L. R. Global expansion and redistribution of

Aedes-borne virus transmission risk with climate change. *PLoS Negl. Trop. Dis.* **13**, e0007213 (2019).

4.    Godoy, R. S. M. *et al.* Dengue and Zika virus infection patterns vary among Aedes aegypti field

populations from Belo Horizonte, a Brazilian endemic city. *PLoS Negl. Trop. Dis.* **15**, e0009839

(2021).

5.    Sun, P. *et al.* A mosquito salivary protein promotes flavivirus transmission by activation of

autophagy. *Nat. Commun.* **11**, 260 (2020).

6.    Ha, Y.-R. *et al.* Detection of Heparin in the Salivary Gland and Midgut of Aedes togoi. *Korean J.*

*Parasitol.* **52**, 183–188 (2014).

7.    Schneider, B. S. & Higgs, S. The enhancement of arbovirus transmission and disease by mosquito

saliva is associated with modulation of the host immune response. *Trans. R. Soc. Trop. Med. Hyg.*

**102**, 400–408 (2008).

8.    Guerrero, D., Cantaert, T. & Missé, D. Aedes Mosquito Salivary Components and Their Effect on the

Immune Response to Arboviruses. *Front. Cell. Infect. Microbiol.* **10**, 407 (2020).

9.    Manning, J. E. & Cantaert, T. Time to Micromanage the Pathogen-Host-Vector Interface:

Considerations for Vaccine Development. *Vaccines* **7**, 10 (2019).

10. Vogt, M. B. *et al.* Mosquito saliva alone has profound effects on the human immune system. *PLoS*

*Negl. Trop. Dis.* **12**, e0006439 (2018).

11. Jin, L. *et al.* Salivary factor LTRIN from Aedes aegypti facilitates the transmission of Zika virus by

interfering with the lymphotoxin-β receptor. *Nat. Immunol.* **19**, 342–353 (2018).

12. Pingen, M. *et al.* Host Inflammatory Response to Mosquito Bites Enhances the Severity of Arbovirus

Infection. *Immunity* **44**, 1455–1469 (2016).

13. Schneider, B. S., Soong, L., Zeidner, N. S. & Higgs, S. Aedes aegypti salivary gland extracts modulate

anti-viral and TH1/TH2 cytokine responses to sindbis virus infection. *Viral Immunol.* **17**, 565–573

(2004).

14. Hastings, A. K. *et al.* Aedes aegypti NeSt1 Protein Enhances Zika Virus Pathogenesis by Activating

Neutrophils. *J. Virol.* **93**, e00395-19, /jvi/93/13/JVI.00395-19.atom (2019).

15. Uraki, R. *et al.* Aedes aegypti AgBR1 antibodies modulate early Zika virus infection of mice. *Nat.*

*Microbiol.* **4**, 948–955 (2019).

16. McCracken, M. K. *et al.* Route of inoculation and mosquito vector exposure modulate dengue virus

replication kinetics and immune responses in rhesus macaques. *PLoS Negl. Trop. Dis.* **14**, e0008191

(2020).

17. Dudley, D. M. *et al.* Infection via mosquito bite alters Zika virus tissue tropism and replication

kinetics in rhesus macaques. *Nat. Commun.* **8**, 2096 (2017).

18. Trager, W. Acquired Immunity to Ticks. *J. Parasitol.* **25**, 57 (1939).

19. Guilbride, D. L., Guilbride, P. D. L. & Gawlinski, P. Malaria's deadly secret: a skin stage. *Trends*

*Parasitol.* **28**, 142–150 (2012).

20. Conway, M. J. Type I hypersensitivity promotes Aedes aegypti blood feeding. *Sci. Rep.* **11**, 14891

(2021).

21. Demeure, C. E. *et al.* Anopheles Mosquito Bites Activate Cutaneous Mast Cells Leading to a Local

Inflammatory Response and Lymph Node Hyperplasia. *J. Immunol.* **174**, 3932–3940 (2005).

22. Conway, M. J. *et al.* Mosquito Saliva Serine Protease Enhances Dissemination of Dengue Virus into

the Mammalian Host. *J. Virol.* **88**, 164–175 (2014).

23. Turner, C. T., Lim, D. & Granville, D. J. Granzyme B in skin inflammation and disease. *Matrix Biol.* **75–**

**76**, 126–140 (2019).

24. Buzza, M. S. *et al.* Extracellular Matrix Remodeling by Human Granzyme B via Cleavage of

Vitronectin, Fibronectin, and Laminin. *J. Biol. Chem.* **280**, 23549–23558 (2005).

25. Champagne, D. E. Antihemostatic Molecules from Saliva of Blood-Feeding Arthropods. *Pathophysiol.*

*Haemost. Thromb.* **34**, 221–227 (2005).

26. Lefteri, D. A. *et al. Mosquito saliva sialokinin-dependent enhancement of arbovirus infection through*

*endothelial barrier leakage*. http://biorxiv.org/lookup/doi/10.1101/2021.02.19.431961 (2021)

doi:10.1101/2021.02.19.431961.

27. Davies, L. C., Jenkins, S. J., Allen, J. E. & Taylor, P. R. Tissue-resident macrophages. *Nat. Immunol.* **14**,

986–995 (2013).

28. Mantovani, A., Sica, A. & Locati, M. Macrophage Polarization Comes of Age. *Immunity* **23**, 344–346

(2005).

29. Mantovani, A., Biswas, S. K., Galdiero, M. R., Sica, A. & Locati, M. Macrophage plasticity and

polarization in tissue repair and remodelling: Macrophage plasticity and polarization in tissue repair

and remodelling. *J. Pathol.* **229**, 176–185 (2013).

30. Bardina, S. V. *et al.* Differential Roles of Chemokines CCL2 and CCL7 in Monocytosis and Leukocyte

Migration during West Nile Virus Infection. *J. Immunol.* **195**, 4306–4318 (2015).

31. Mills, C. D. Anatomy of a Discovery: M1 and M2 Macrophages. *Front. Immunol.* **6**, (2015).

32. Lamana, A. *et al.* CD69 Modulates Sphingosine-1-Phosphate-Induced Migration of Skin Dendritic

Cells. *J. Invest. Dermatol.* **131**, 1503–1512 (2011).

33. Mackay, L. K. *et al.* Cutting Edge: CD69 Interference with Sphingosine-1-Phosphate Receptor

Function Regulates Peripheral T Cell Retention. *J. Immunol.* **194**, 2059–2063 (2015).

34. Rajarathnam, K., Schnoor, M., Richardson, R. M. & Rajagopal, S. How do chemokines navigate

neutrophils to the target site: Dissecting the structural mechanisms and signaling pathways. *Cell.*

*Signal.* **54**, 69–80 (2019).

35. Dey, R. *et al.* Gut Microbes Egested during Bites of Infected Sand Flies Augment Severity of

Leishmaniasis via Inflammasome-Derived IL-1β. *Cell Host Microbe* **23**, 134-143.e6 (2018).

36. Gough, P., Ganesan, S. & Datta, S. K. IL-20 Signaling in Activated Human Neutrophils Inhibits

Neutrophil Migration and Function. *J. Immunol.* **198**, 4373–4382 (2017).

37. Myles, I. A. *et al.* Signaling via the IL-20 receptor inhibits cutaneous production of IL-1β and IL-17A

to promote infection with methicillin-resistant Staphylococcus aureus. *Nat. Immunol.* **14**, 804–811

(2013).

38. Lefteri, D. A. *et al.* Mosquito saliva enhances virus infection through sialokinin-dependent vascular

leakage. *Proc. Natl. Acad. Sci.* **119**, e2114309119 (2022).

39. Kirkpatrick, B. D. *et al.* The live attenuated dengue vaccine TV003 elicits complete protection against

dengue in a human challenge model. *Sci. Transl. Med.* **8**, (2016).

40. Shimizu, K. *et al.* PD-1 Imposes Qualitative Control of Cellular Transcriptomes in Response to T Cell

Activation. *Mol. Cell* **77**, 937-950.e6 (2020).

41. Wei, F. *et al.* Strength of PD-1 signaling differentially affects T-cell effector functions. *Proc. Natl.*

*Acad. Sci.* **110**, E2480–E2489 (2013).

42. Duyen, H. T. L. *et al.* Skin dendritic cell and T cell activation associated with dengue shock syndrome.

*Sci. Rep.* **7**, 14224 (2017).

43. Rivino, L. *et al.* Virus-specific T lymphocytes home to the skin during natural dengue infection. *Sci.*

*Transl. Med.* **7**, (2015).

44. Mommert, S., Schaper, J. T., Schaper-Gerhardt, K., Gutzmer, R. & Werfel, T. Histamine Increases Th2

Cytokine-Induced CCL18 Expression in Human M2 Macrophages. *Int. J. Mol. Sci.* **22**, 11648 (2021).

45. Styer, L. M. *et al.* Mosquito Saliva Causes Enhancement of West Nile Virus Infection in Mice. *J. Virol.*

**85**, 1517–1527 (2011).

46. Chiffoleau, E. C-Type Lectin-Like Receptors As Emerging Orchestrators of Sterile Inflammation

Represent Potential Therapeutic Targets. *Front. Immunol.* **9**, 227 (2018).

47. Holmskov, U., Thiel, S. & Jensenius, J. C. Collectins and Ficolins: Humoral Lectins of the Innate

Immune Defense. *Annu. Rev. Immunol.* **21**, 547–578 (2003).

48. Tang, L. *et al.* Liver Sinusoidal Endothelial Cell Lectin, LSECtin, Negatively Regulates Hepatic T-Cell

Immune Response. *Gastroenterology* **137**, 1498-1508.e5 (2009).

49. Kuo, P. T. *et al.* The Role of CXCR3 and Its Chemokine Ligands in Skin Disease and Cancer. *Front.*

*Med.* **5**, 271 (2018).

50. Niyonsaba, F., Ogawa, H. & Nagaoka, I. Human beta-defensin-2 functions as a chemotactic agent for

tumour necrosis factor-alpha-treated human neutrophils. *Immunology* **111**, 273–281 (2004).

51. Niyonsaba, F., Iwabuchi, K., Matsuda, H., Ogawa, H. & Nagaoka, I. Epithelial cell-derived human β-

defensin-2 acts as a chemotaxin for mast cells through a pertussis toxin-sensitive and phospholipase

C-dependent pathway. *Int. Immunol.* **14**, 421–426 (2002).

52. Rizzo, C. *et al.* IgG1 and IgG4 Antibody Responses to the Anopheles gambiae Salivary Protein gSG6 in

the Sympatric Ethnic Groups Mossi and Fulani in a Malaria Hyperhendemic Area of Burkina Faso.

*PLoS ONE* **9**, e96130 (2014).

53. McKimmie, C. S., Pingen, M., Bryden, S. R. & Lefteri, D. A. Host inflammatory response to mosquito

bites enhances the severity of arbovirus infection. *ISBT Sci. Ser.* **13**, 76–79 (2018).

54. Henrique, M. O. *et al.* Evaluation of inflammatory skin infiltrate following *Aedes aegypti* bites in

sensitized and non-sensitized mice reveals saliva-dependent and immune-dependent phenotypes.

*Immunology* **158**, 47–59 (2019).

55. Qureshi, A. A. *et al.* Immunomodulatory properties of maxadilan, the vasodilator peptide from sand

fly salivary gland extracts. *Am. J. Trop. Med. Hyg.* **54**, 665–671 (1996).

56. Gillespie, R. D., Dolan, M. C., Piesman, J. & Titus, R. G. Identification of an IL-2 binding protein in the

saliva of the Lyme disease vector tick, Ixodes scapularis. *J. Immunol. Baltim. Md 1950* **166**, 4319–

4326 (2001).

57. Oliveira, F. *et al.* Delayed-Type Hypersensitivity to Sand Fly Saliva in Humans from a Leishmaniasis-

Endemic Area of Mali Is TH1-Mediated and Persists to Midlife. *J. Invest. Dermatol.* **133**, 452–459

(2013).

58. Pasparakis, M., Haase, I. & Nestle, F. O. Mechanisms regulating skin immunity and inflammation.

*Nat. Rev. Immunol.* **14**, 289–301 (2014).

59. Manning, J. E. *et al.* Safety and immunogenicity of a mosquito saliva peptide-based vaccine: a

randomised, placebo-controlled, double-blind, phase 1 trial. *The Lancet* S0140673620310485 (2020)

doi:10.1016/S0140-6736(20)31048-5.

60. Oliveira, F. *et al.* A sand fly salivary protein vaccine shows efficacy against vector-transmitted

cutaneous leishmaniasis in nonhuman primates. *Sci. Transl. Med.* **7**, (2015).

61. Sajid, A. *et al.* mRNA vaccination induces tick resistance and prevents transmission of the Lyme

disease agent. *Sci. Transl. Med.* **13**, eabj9827 (2021).

62. Bryden, S. R. *et al.* Pan-viral protection against arboviruses by activating skin macrophages at the

inoculation site. *Sci. Transl. Med.* **12**, eaax2421 (2020).

63. Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina sequence data.

*Bioinformatics* **30**, 2114–2120 (2014).

64.  Li, B. & Dewey, C. N. RSEM: accurate transcript quantification from RNA-Seq data with or without a

reference genome. *BMC Bioinformatics* **12**, 323 (2011).

65.  Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat. Methods* **9**, 357–359

(2012).

66.  Dobin, A. & Gingeras, T. R. Mapping RNA-seq Reads with STAR. *Curr. Protoc. Bioinforma.* **51**, (2015).

67.  Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-seq

data with DESeq2. *Genome Biol.* **15**, 550 (2014).

68.  Huber, W., von Heydebreck, A., Sultmann, H., Poustka, A. & Vingron, M. Variance stabilization

applied to microarray data calibration and to the quantification of differential expression.

*Bioinformatics* **18**, S96–S104 (2002).

69.  Zhu, A., Ibrahim, J. G. & Love, M. I. Heavy-tailed prior distributions for sequence count data:

removing the noise and preserving large differences. *Bioinformatics* **35**, 2084–2092 (2019).

70.  Benjamini, Y. & Hochberg, Y. Controlling the False Discovery Rate: A Practical and Powerful

Approach to Multiple Testing. *J. R. Stat. Soc. Ser. B Methodol.* **57**, 289–300 (1995).

71.  Blighe, K., Rana, S. & Lewis, M. Enhanced Volcano: publication-ready volcano plots with enhanced

colouring and labeling. (2021).

72.  Walter, W., Sánchez-Cabo, F. & Ricote, M. GOplot: an R package for visually combining expression

data with functional analysis: Fig. 1. *Bioinformatics* **31**, 2912–2914 (2015).

73.  Yu, G., Wang, L.-G., Han, Y. & He, Q.-Y. clusterProfiler: an R Package for Comparing Biological

Themes Among Gene Clusters. *OMICS J. Integr. Biol.* **16**, 284–287 (2012).

74.  Yu, G. & He, Q.-Y. ReactomePA: an R/Bioconductor package for reactome pathway analysis and

visualization. *Mol. Biosyst.* **12**, 477–479 (2016).

75.  Kolde, R. Pretty Heatmaps. R package version.

76. Sakai, R., Winand, R., Verbeiren, T., Moere, A. V. & Aerts, J. dendsort: modular leaf ordering

methods for dendrogram representations in R. *F1000Research* **3**, 177 (2014).

77. R Core Team (2021). *R: A language and environment for statistical computing. R Foundation fo*

*Statistical Computing, Vienna, Austria*. (The R Foundation).

798