1   **First-in-human evaluation of cutaneous innate and adaptive immunomodulation by**

2   **mosquito bites**

3

4   David Guerrero[1,*], Hoa Thi My Vo[1,*], Chanthap Lon[3,*], Jennifer A. Bohl[2], Sreynik Nhik[3], Sophana Chea[3], Somnang Man[3,4], Sokunthea Sreng[3,4], Andrea R. Pacheco[3], Sokna Ly[3], Rathanak Sath[3], Sokchea Lay[1], Dorothee Missé[5], Rekol Huy[4], Rithea Leang[4], Hok Kry[6], Jesus G. Valenzuela[2], Fabiano Oliveira[2,$], Tineke Cantaert[1,$], Jessica E. Manning[2,3,$,#]

5

9   [1] Institut Pasteur du Cambodge, Pasteur Network, Phnom Penh, Cambodia

10   [2] Laboratory of Malaria and Vector Research, National Institute of Allergy and Infectious Diseases,

11   National Institutes of Health, Rockville, Maryland, USA

12   [3] International Center of Excellence in Research, National Institute of Allergy and Infectious Diseases,

13   National Institutes of Health, Phnom Penh, Cambodia

14   [4] National Center of Parasitology, Entomology, and Malaria Control, Phnom Penh, Cambodia

15   [5] MIVEGEC, Univ Montpellier, IRD, CNRS, 34000 Montpellier, France.

16   [6] Kampong Speu Provincial District, Ministry of Health, Phnom Penh, Cambodia

17

18   *co-first author

19 $ co-last author

20   # corresponding author: jessica.manning@nih.gov

21

22 Abstract

Mosquito-borne viruses are a growing global threat. Initial viral inoculation occurs in the skin via

the mosquito 'bite', eliciting immune responses that shape the establishment of infection and

pathogenesis. To understand these phenomena, we assessed the cutaneous innate and adaptive

immune responses via controlled *Aedes aegypti* feedings in humans living in an *Aedes*-endemic

country. Gene expression profiling and immunophenotyping revealed induction of neutrophil

degranulation and recruitment of skin-resident dendritic cells and M2-macrophages. As the

immune reaction progressed over time, T cell priming and regulatory pathways were upregulated

along with a shift to a Th2-driven response and CD8+ T cell activation. In accordance, participants'

bitten skin cells produced less pro-inflammatory cytokines when stimulated by *Ae. aegypti* salivary

gland extract. These results identify key immune genes, cell types, and pathways in the human

response to mosquito bites that can be leveraged to develop novel therapeutics and vector-targeted

vaccine candidates to arboviral diseases.

 Graphical Abstract

Introduction

With one million deaths every year, vector-borne diseases remain a major health problem worldwide[1]. In particular, arboviruses are of particular concern given a global distribution and a burden of nearly half a billion infections[2]. Several clinically significant arboviruses such as yellow fever (YFV), dengue (DENV), chikungunya (CHIKV), and Zika (ZIKV) viruses are all transmitted by *Aedes spp*. mosquitoes. These vectors are abundant in tropical and subtropical areas, but current climate change and rapid urbanization are expanding their transmission potential and increasing the at-risk population[3].

Mosquitoes first acquire an arbovirus via insertion of their proboscis into the dermis of an infected host during a blood meal. The virus then replicates in midgut cells to later disseminate and reach the salivary glands (SG)[4]. During a subsequent blood meal, the infectious mosquito inoculates virus and insect-derived molecules, including salivary components, into the dermis of a new human host. The saliva from hematophagous mosquitoes is a cocktail of pharmacologically active molecules, some of which are anti-hemostatic, angiogenic, and immunomodulatory[5–7]. Over the last couple decades, several animal and *in vitro* studies demonstrated that inoculation of vector saliva, and/or concomitant blood-feeding by an arthropod, can immunomodulate the host antiviral response in the skin and periphery[7–10]. These perturbations result in a dampening of the pro-inflammatory responses[11–13], promotion of recruitment of infection-susceptible cells to the bite site[14,15], stimulation of autophagy activation[5], and an induction of neutrophil infiltration to the local inflammation site[15] amongst other consequences. Mosquito-inoculated DENV or ZIKV infection in macaques led to higher or delayed viremia compared to non-vector or needle inoculation routes[16,17]. With regard to clinical pathogenesis, mice inoculated with both Semliki Forest virus and mosquito saliva had higher loads of virus RNA in the skin, earlier brain dissemination, and

more frequent lethal outcomes than animals inoculated with virus alone[12]. Taken together,

arthropod-mediated viral infection leads to increased disease progression, viremia, and

mortality[8,9]. Despite this robust evidence from animal and *in vitro* studies, there are no translational

studies on the human skin response to mosquito bites, much less those in tropical areas. Individuals

are continuously exposed to mosquito bites, possibly shaping their immune responses to mosquito

salivary proteins in contrast to those intermittently or rarely exposed[18,19]. In order to examine the

dermal and epidermal innate and adaptive immune responses to *Ae. aegypti* saliva, we compared

the immune responses before and after *Ae. aegypti* bites in skin biopsies from 30 healthy

Cambodian individuals living in an area where *Ae. Aegypti* mosquitoes are prevalent. Our study

provides information on the key genes and cell populations involved in the human immune

response at different time points after the 'mosquito bite.'

Results

Controlled *Aedes aegypti* bite challenge in healthy volunteers in Cambodia

From October 2020 to January 2021, we enrolled 30 healthy Cambodian adults at Kampong Speu

District Referral Hospital in Chbar Mon, Cambodia. The study population comprised 13 females

and 17 males, all of whom were of Khmer (Cambodian) ethnicity (Supplementary Table 1). The

population was randomly divided in three groups to assess immune responses by different

technologies (Fig. 1). The median age was 32.5 years (IQR 22–35) and the primary occupations

were variable, although nearly half (14/30) were unemployed. In terms of mosquito exposure risk,

most participants reported living in a house (20/30), being middle income (25/30), having one to

two domestic water containers (22/30). Notably, most participants did not use larvicide (28/30) or

insecticide (19/30) in their homes. Only 10% (3/30) reported prior dengue infection, but all 30

participants were positive for dengue anti-NS1 antibodies. All participants had detectable IgG to

Ae. aegypti salivary gland extract (SGE) prior to mosquito feeding, and these levels were not

significantly different by age, gender, or 14 days after the mosquito feeding (Extended Data Fig.

1). One participant did not complete Day 14 follow-up and was considered lost to follow-up. No

unanticipated adverse events occurred with the biopsies or mosquito feedings.

Figure 1. **Graphical Representation of Study Design.** Thirty Cambodian participants were

enrolled in the study. Each patient underwent a mosquito feeding (5 females *Aedes aegypti*) for

ten minutes. Each volunteer underwent a 3mm biopsy from the mosquito bite site at 30 minutes,

4 hours, and 48 hours. Additionally, two 3mm biopsies of normal skin (NSK) were taken from

the opposite forearm. Blood was collected to obtain serum.

95 **Marked clinical changes occurred in the skin post-mosquito feeding**

96 Exposure to five *Ae. aegypti* mosquitos resulted in 3 to 10 visible bite sites per volunteer. Upon

97 clinical observation, participants experienced erythema and swelling with a "bite size" reaction

98 that peaked at 15 minutes after feeding (Fig. 2a). The mean bite size of 4.9mm at 15 min after

99 feeding decreased to 4.7 mm at 30 minutes, to 2.7 mm at 4 hours and gradually decreased to 2.1mm

100 by 48 hours (Supplementary Table 1). Observation of representative histological sections shows

101 increased recruitment of cells to the dermis and epidermis at 48 hours post-bite (Fig. 2b,c). There

102 was no difference in bite size reaction by gender or age.

103

Figure 2. **Clinical observations of skin reaction pre- and post-bite on gross examination and**

H&E staining. a, Skin immune responses to mosquito bites (left to right) normal skin (NSK), 30

minutes, 4 hours, and 48 hours after exposure to five *Ae. aegypti* mosquitos. **b,** Representative

cross-sectional of 3mm punch biopsies H&E stained at (left to right) 0 hours, 30 minutes, 4 hours,

and 48 hours after exposure to *Ae. aegypti* bites. Scale bar is 1 mm. Black arrows indicate area of

infiltrate. **c,** Magnification of infiltrate indicated by black areas in B, scale bar 100μM.

Changes in gene expression in *Aedes aegypti* bitten versus normal skin (NSK) evolved over

48 hours

We extracted RNA and performed next generation sequencing on 40 skin biopsies from 10

participants (baseline normal skin (NSK) biopsy on the forearm, plus 3 follow-up biopsies each at

30 minutes, 4 hours, and 48 hours post *Ae. aegypti* bites on the contralateral arm). Principal

Component (PC) analysis of *Ae. aegypti* bite site skin biopsies resulted in segregation of sample

clusters based on the time the biopsy was taken (Fig. 3a), with PC1 and PC2 variances defining

differences between NSK and 30 minutes versus 4 hours and 48 hours. While PC3 variance

associated to the separation between biopsies taken at 4 hours and 48 hours, samples taken from

NSK (pink circles) and 30 minutes after *Ae. aegypti* bites clustered in the same spatial area. At 4

hours post-bite, 6 of 10 samples occupied a distinct region (Fig. 3a, blue ellipse) from NSK, 30

minutes or 48 hours after *Ae. aegypti* bites. Similarly at 48 hours post-bite, 7 of 10 samples

segregated into a discrete space (Fig. 3a, orange ellipse) compared to NSK, 30 minutes or 4 hours

after *Ae. aegypti* bites.   These data indicate that time post-bite rather than inter-individual

differences drive unique patterns in gene expression. Differential analysis of RNA expression with

a threshold of two-fold (-1 < log2FC >1; False discovery rate (FDR) < 0.05) over NSK biopsies

resulted in 709 upregulated genes post-bite compared to 31 downregulated genes (Fig. 3b, red

circles). At 30 minutes, 4 hours, and 48 hours after a mosquito bite, 26, 210 and 473 genes were

upregulated over NSK, respectively (Fig. 3c, Supplementary Table 2). Conversely, one gene at 4

hours and 27 genes at 48 hours were downregulated after *Ae. aegypti* bites (Supplementary Table

2).

Upregulated genes at each time point were subjected to over-representation analysis (ORA) for

enriched pathways using the Reactome database with an FDR cut off < 0.05. Four pathways related

to cell proliferation, cell growth and nuclear transcription were identified (FDR < 0.05) at 30

minutes after *Ae. aegypti* bites (Supplementary Table 3). At both 4 and 48 hours post-bite, the

immune nature of the host response to *Ae. aegypti* bites was evident via the revealed pathways

(Fig. 3d, Supplementary Table 3). At 4 hours post-bite, pathways such as "neutrophil

degranulation", "Interleukin-4 and Interleukin-13 signaling", "Interleukin-10 signaling",

"Interferon-γ signaling" "Chemokine receptors bind chemokines" and "Immunoregulatory

interactions between a Lymphoid and a non-Lymphoid cell" (Fig. 3d, upper chord) were

significantly overrepresented and maintained until 48 hours post-bite (Fig. 3d, lower chord).

Unique pathways discovered at 4 hours post-bite were related to "degradation of extracellular

matrix" (Fig 3d, upper chord) while "antimicrobial peptides" pathway and "adaptive immune

system processes" were present at 48 hours post-bite including "TCR signaling", "Co-stimulation

by the CD28 family", "PD-1 signaling", "MHC class II antigen presentation" (Fig. 3d, lower

chord). A heatmap with differentially expressed genes (DEG) under an FDR cutoff < 0.05 and

log2FC >1 in at least one timepoint was assembled (Extended Data Fig. 3, Supplementary Table

4), showing the dynamics of DE expression throughout the three timepoints. Fig. 3e shows DEG

driving the common pathways found on Fig. 3d.

Overall, few DEG genes were detected at 30 minutes post-bite compared to NSK biopsies. The

main upregulated gene FOSB (up 18.8x) is one of the transcription factors of the FOS family

proteins that regulate cell proliferation and differentiation (Supplementary Table 2). As the

reaction progresses over time, gene expression increased towards neutrophil recruitment followed

by signaling of both Th1, Th2, and regulatory pathways. The five most regulated genes at 4 hours

compared to NSK are KRT6C (related to keratinization; up 80.8x), CXCL8 (neutrophil-attracting

chemokine IL-8 up 75.0x), TNIP3 (inhibitor of NF-kappa-B activation; up 59.3x), IL-20

(stimulates keratinocyte proliferation and differentiation; up 59.1x) and IL-1B (pro-inflammatory

cytokine; up 45.7x) (Supplementary Table 2). The immune response evolved at 48 hours post-bite

to include adaptive immune response pathways related to T-cell priming. The five top genes at 48

hours are KRT6C (related to keratinization; up 59.6x), DEFB4A (an antimicrobial peptide; up

32.7x), GZMB (Granzyme B, up 21.1x), TCL1A (proliferation of T and B cells; up 20.3x) and

CCL18 (chemokine associated with associated with TH2 responses; up 18.5x).

Figure 3. **Differential expression gene (DEG) and pathway analysis induced by** *Aedes aegypti*

mosquito bites over time. a, 3D principal component analysis showing clustering of individual

biopsies taken at 30 minutes (green dots), 4 hours (blue dots), 48 hours (orange dots) post-bite or

normal skin (NSK; pink dots). Blue ellipse indicates clustering at 4 hours, while orange ellipse at

48 hours. **b,** Volcano plot comparing DEG from skin biopsies post-bite with NSK (significant

genes with -1 < log2FC >1 and FDR < 0.05 are shown in red). **c,** Venn diagram summarizing the

relationships between significantly upregulated genes over NSK (log2FC >1 and FDR < 0.05) and

time post-bite. **d,** Chord plot shows significantly upregulated genes at 4 hours and 48 hours post-

bite over NSK (left side of chord plot) and overrepresented pathways (right side of chord plot).

Outer left ring shows genes and color range displays log2 fold change (left, key at bottom) or

Reactome terms (outer right ring). Chords connect gene names with Reactome term pathways. **e,**

Heatmap (scaled to log2FC) displaying a list of DEGs based on the common pathways

overrepresented at 4- and 48-hours post-bite.

Early innate immune response in the skin is characterized by increased proportions of DCs

and M2 macrophages and a decreased proportion of NK cells

To confirm and extend the gene expression analysis, dissociated cells from NSK and skin biopsies

taken at 30 minutes and 4 hours after exposure to mosquitos, were characterized for innate and

adaptive immune subsets by flow cytometry (Supplementary Table 5). No significant changes in

cell populations were detected between NSK and 30 minutes post-bite, mirroring the

transcriptomic analysis of DEG. At 4 hours post-bite, the proportion of dendritic cells (DC), and

M2 macrophages increased, while the proportion of mast cells, natural killer (NK) and T-cells

decreased, when compared to NSK and 30 minutes post-bite (Fig. 4a). The frequencies of DC

subsets and macrophages increased at 4 hours post-bite compared to NSK and 30 minutes (Fig. 4

b-h); a decreasing, though non-significant, trend in the NK cell population frequency was observed

over time (Fig. 4i). Within the DC subsets, Langerhans cells (CD45+CD3-CD207+) and activated

dermal DCs (CD45+CD3-CD1c+CD69+) increased at 4 hours compared to NSK while only

activated dermal DCs were increased at 30 minutes. (Fig. 4d, e). Regarding the macrophage

population, M2-skewed macrophages (CD45+CD14+CD163+) frequency increased after 4 hours

post-bite as well as their expression of activation markers (CD16/CD69) compared to NSK and 30

minutes (Fig. 4 f-h). Frequencies of activated M2 macrophages, activated dermal DCs, and

frequency of Langerhans cells positively correlated with clinically observed changes in bite size

at 30 minutes (Extended Data Fig. 2 a-c). The relative proportion of CD4$^+$ and CD8$^+$ T cells did

not change over time. Taken together, the results suggest that immune cell infiltration and

migration occurs in local skin tissue within 4 hours post-bite and is characterized by increased

frequencies of various DCs and M2 macrophages, but decreased frequencies of NK cells.

Figure 4. **Early innate immune responses in the skin are characterized by increased proportions of DCs and M2 macrophages, and a decreased proportion of NK cells at 4 hours post-bite**. **a,** Pie charts showing changes in the frequencies of skin immune cells during early innate immune response to mosquito bite at 30 minutes and 4 hours after exposure. **b-e,** Frequency changes in total DCs and subsets: Plasmacytoid DCs (CD123+), Langerhans cells (CD209+), and activated dermal DCs (CD1c+CD69+) at 4 hours post mosquito bites. **f-h,** Increase in M2 macrophages (CD14+CD163+) frequencies and their activation phenotypes (CD16+/CD69+). **i,** Frequency of NK cells (CD56+) in NSK, 30 minutes and 4 hours. **j,** Frequency of cytotoxic NK

209 cells in NSK, 30 minutes and 4 hours (CD56+CD16+). Statistical analysis were performed with

210 Chi-square test (**a**) and Friedman + Dunn's multiple comparisons test (**b-j**). Bars indicate median

211 and interquartile range.

213   Changes in the T cell compartment are observable at 48 hours after mosquito 'bite'

Changes in skin T cell populations were assessed using NSK and biopsies taken at 48 hours post-

bite. We found that the frequency of CD8+CD25+ and CD8+PD-1+ T-cells increased at 48 hours

post-bite, compared to the NSK (Fig. 5a). Frequencies of CD8+ T cells correlated inversely with

the bite size at 48 hours (Extended Data Fig. 2d). The frequency of effector memory cells re-

expressing CD45RA (T_{EMRA}) CD8 T cells (CD8+CCR7-CD45RA+) decreased at 48 hours (Fig.

5b). We observed a decrease in the frequency of Th1/Th17 (CD4+CCR4-CXCR3+) cells at 48h

compared to NSK and an increase in the frequency of Th2/Th17 (CD4+CCR4+CXCR3-) cells

(Fig. 5c). Within the CD4+ T cell compartment, the frequency of CD4+ effector memory T cells

(T_{EM}) (CD4+CCR7-CD45RA-) increased at 48 hours compared to NSK, while the frequency of

T_{EMRA} cells (CD4+CCR7-CD45RA+) decreased (Fig. 5c). Interestingly, no changes were

observed in the frequencies of skin-resident CD4 (CD4+CD103+CLA+) or CD8

(CD8+CD103+CLA+) T cells (Extended Data Fig. 4). At 48 hours post-bite, these results show

an increase in the frequency of activated/exhausted CD8 T cells in addition to a shift from

Th1/Th17 towards Th2/Th17 polarization of CD4 T cells.

Figure 5. **Changes in the T cell compartment are observable at 48 hours after mosquito exposure. CD8 T cells exhibit a high activation/exhausted phenotype, while CD4 T effector memory cells decrease in their Th1Th17 compartment and increase their Th2Th17 compartment. a,** Increased frequency of CD25+ and PD1+ CD8 T cells at 48 hours post mosquito exposure. **b,** Frequencies of CD8 T cells naive (CCR7+CD45RA+), central memory (CCR7+CD45RA-), effector memory (CCR7-CD45RA-) and TEMRA (CCR7-CD54RA+) frequencies. **c,** decreased Th1/Th17 and increased Th2/Th17 effector memory CD4 T cell compartment at 48 hours. **d,** CD4 T cells naive (CCR7+CD45RA+), central memory (CCR7+CD45RA-), effector memory (CCR7-CD45RA-) and TEMRA (CCR7-CD54RA+) frequencies. Statistical analysis was performed with Wilcoxon signed-rank test. Bars indicate median and interquartile range.

241   **"Bitten" human skin cells stimulated by *Aedes aegypti* saliva produce significantly less pro-**

242   **inflammatory cytokines**

243   Next, we sought to assess if *Ae. aegypti* saliva impacted cytokine production by skin cells taken

244   48 hours post-bite. We measured pro- and anti-inflammatory cytokines released into the media

245   after a stimulation assay with and without *Ae. aegypti* SGE. We observed a significant reduction

246   in the levels of Interleukin 2 (IL-2) and Interferon gamma (IFN-γ) in the cells treated with both

247   SGE and PMA/Ionomycin compared to the cells treated with PMA/Ionomycin only (Fig. 6,

248   Extended Data Fig. 5). The levels of other pro-inflammatory cytokines like TNF-α, IL-17A and

249   IL-17F showed no significant change (Fig. 6). Similarly, no change in the regulatory cytokine IL-

250   10 was observed (Fig. 6). This data suggests that SGE has a dampening effect on the production

251   of pro-inflammatory cytokines IL-2 and IFN-γ under stimulatory conditions in "bitten" human

252   skin cells.

253

254   Figure 6. **Human skin cells obtained from *Ae. aegypti* bitten skin stimulated by *Ae. aegypti***

255   **salivary gland extract (SGE) produce significantly less pro-inflammatory cytokines (IL-2**

256   **and IFN-γ)**. Cytokine production measured in skin cell culture supernatant after treatment with

PMA/Ionomycin in the presence or absence of SGE. Statistical analysis was performed with

Wilcoxon signed-rank test.

Discussion

Arboviruses enter the host through the skin when the arthropod injects its saliva and other

components into the dermis while taking a bloodmeal. Data from mice and macaques suggest that

the unique dual presence of virus and saliva orchestrates an immunological cascade of events that

influence the establishment and course of infection[12,16,17]. To translate these findings to humans,

the current study assessed the dynamics of the cutaneous immune response to arthropod bites from

a clinically relevant vector, the *Ae. aegypti* mosquito, in humans from arboviral-endemic areas

with lifelong exposure to *Aedes* mosquitos. At 4 hours post-bite, we observed upregulation of

genes involved in neutrophil recruitment and increased frequencies of dendritic cells and M2

macrophages. As the immune response progressed over two days, we observed an increased

frequency of activated, $PD-1^+CD8^+$ T cells, a polarization towards Th2/Th17 $CD4^+$ T-cells, and

upregulation of IL-10, IL-4, and IL-13 and IFN-γ related pathways. Interestingly, these human

skin cells post-bite produced less pro-inflammatory cytokines when stimulated by *Ae. aegypti*

SGE.

A mosquito 'bite' results in minor tissue damage by the proboscis, injection of mosquito saliva

into the skin, and introduction of both mosquito and host microbiota into the skin. In this study,

we did not parse out the separate impact of each process on the skin immune response. Alterations

in skin immunity due to injury from the punch biopsy itself were mitigated by ensuring biopsy

sites were at least 10 to 20 mm away from one another and were not taken directly at the site of

the needle administration of subcutaneous lidocaine. The effect of local anesthetic was mitigated

by the case-control design that each person's anesthetized, bitten skin was compared to

anesthetized, unbitten skin, presumably negating any immediate cutaneous immune effects

attributed to lidocaine, for which the literature is sparse.

Mosquito bites cause a local cutaneous reaction with wheal and flare manifestation[20,21].

Concordantly, the bite size reaction peaked after 15 minutes post-bite in this endemic study

population. Few changes were detected in gene expression or influx and activation of innate

immune cells at the 30 minutes post-bite time point. Upregulated genes were related to top-level

processes such as signal transduction. We did not design this study to assess the role of pre-stored

immune mediators such as histamine (and others) in the development of edema and their

modulation in the immune response. Indeed, mosquito bites can activate cutaneous mast cells,

which are prominent histamine-storing cells, and induce their degranulation. The release of

histamine and other mediators leads to vasodilation, fluid accumulation at the bite site, and

neutrophil influx, which were all ultimately observed here[20,21].

As the immune reaction progressed, the mosquito 'bite' induced degradation of the extracellular

matrix (ECM), and a strong influx of leukocytes at 4 hours post-bite, consistent with previous

animal observations[22]. We observed several disintegrin and metalloprotease with thrombospondin

type I motifs (ADAMTS) were upregulated, mostly at 4 hours post-bite, that can be associated

with ECM degradation. Similarly, granzyme B (GZMB) upregulated at 48 hours, despite its effects

on inducing inflammation it may also be involved in ECM degradation using as substrate

fibronectin, vitronectin, laminin and others[23,24]. Saliva from blood-feeding arthropods contains

anti-hemostatic proteins that prevent coagulation and promote vasodilation, leading to increased

leukocyte trafficking[20,25]. These processes – extracellular matrix degradation and vasodilation –

can facilitate viral infection by disrupting vascular functions, as is the case of sialokinin, a

mosquito salivary protein that hinders endothelial barrier integrity[26].   In addition, elevated

leukocyte trafficking can facilitate viral replication by augmenting the proportion of susceptible

cells to the local inoculation site[12,15,26]. Indeed, as most leukocyte subsets are permissive to

infection by most arboviruses, they play a dual role in the infection process: both as replication

targets/dissemination vehicles and as the first line of defense.

The strong influx of activated M2 macrophages, DCs and Langerhans cells at 4 hours post-bite

highlight the cellular trafficking role initiated by 'the bite' that can ultimately result in viral

dissemination. Upregulated chemokines such as CCL2 and CCL7, attracting mainly monocytes,

DCs and T cell, may be participating in this recruitment[27]. M2-polarized macrophages produce

anti-inflammatory cytokines, mediate wound healing processes, and support Th2 responses [28].

Moreover, activation of dermal DCs and M2 macrophages positively correlated with the bite size

in our study. Upon stimulation, DCs upregulated CD69, preventing them from exiting the tissue

and improving the likelihood of antigen internalization. Subsequent CD69 downregulation

promotes Sphingosine-1-Phosphate (S1P1) leading to DC migration to the lymph nodes[29]. The

CD69-S1P1 mechanism of cell retention in tissue is also well-described for resident T cells,[30]

whose role in saliva-mediated immunity is not yet known.

Ae. aegypti mosquito saliva has a clear impact on neutrophil influx and degranulation. Two

recently identified *Ae. aegypti* salivary proteins (NeSt1 and AgBr1) that attract and stimulate

neutrophils at the 'bite site,' changing the immune microenvironment and leading to increased

early ZIKV replication and enhanced pathogenesis[14,15]. Here, at 4 hours post-bite, chemokines

CXCL2, CXCL3, CXCL5, and especially CXCL8, plus the cytokine IL-1B were upregulated

returning to levels comparable to NSK at 48 hours post-bite. CXCL2-5 chemokines are responsible

for attracting neutrophils and exhibiting angiogenic properties[31]. Pro-inflammatory responses

through inflammasome activation and IL-1B production can trigger a sustained neutrophilic

response.[32] Microbiota egested during infected sand fly bites contributed to IL-1B production[33].

On the other hand, IL-20 was upregulated at 4 hours post-bite. IL-20 is most active at epithelial

sites downregulating IL-1B resulting in deactivation of neutrophil functions such as phagocytosis,

granule exocytosis, and migration[34,35]. Increased gene expression for neutrophil recruitment and

degranulation at 4 hours post-bite in our Cambodian study population further supports the timing

of neutrophil influx seen 3 hours post-bite in bite-experienced C57BL/6 immunocompetent mice[12].

How the timing of neutrophil influx would differ in a bite-naïve population is a question that

remains to be answered.

As the immune response transitioned from innate to adaptive, a strong T-cell signature emerged at

4 and 48 hours emerged after the mosquito 'bite'. CD8+ T cell activation was confirmed by

increased expression of CD25, an early activation marker. Skin CD8+ T cells also upregulate PD-

1 expression, an immune-checkpoint receptor that is induced by yet can also inhibit T cell

activation. PD-1 primarily inhibits effector function acquisition by T cells, the loss of sensitivity

after T cell receptor signaling and modulation of cytotoxicity, and inhibition of TNF-α and IL-2

production along with T-cell expansion[36,37]. Interestingly, highly activated CD8+ cells have been

observed in skin from dengue shock syndrome patients[38]. Here at 4 and 48 hours post-bite, CD4+

T cells had a skewed phenotype to Th2-like cells and IL-10, IL-13, and IL-4 related pathways were

upregulated compared to NSK. CCL18, upregulated at 48 hours post-bite, can be expressed by M2

macrophages and is linked to increased IL-4, IL-10, and IL-13, especially in the presence of

histamine[39]. Indeed, initial animal experiments also pointed to a Th2-mediated allergic

inflammatory response following mosquito bites or inoculation with SGE[13].

At 48 hours post-bite, other upregulated genes such as CD28, ICOS, and CTLA-4 suggest a

regulatory role of the CD4+ compartment in controlling the local immune responses to *Ae. aegypti*

saliva. Th2 polarization is essential to avoid excessive collateral damage from an exacerbated pro-

inflammatory response, especially in a situation where the skin immune cells are repeatedly

exposed to the same type of insult, as is the case with individuals living in tropical areas of

mosquito abundance. Blood DC antigen-2 (CLEC4C), upregulated at 48h compared to NSK, is

thought to signal through BDCA-2 to inhibit activation of the NF-κB pathway, inhibit production

of type I IFNs, and block TRAIL-mediated cytotoxic activity[40]. This may represent a possible

mechanism that differentiates chronically exposed individuals compared to naïve individuals.

To consider other potential mechanisms that immunomodulate the microenvironment, we noted

increased expression at 48 hours post-bite of ficolin-1 (FCN1), a complement lectin pathway

protein that may bind and opsonize mosquito microbiota or salivary components, then activate the

complement pathway[41]. Another C-type lectin receptor LSECtin (CLEC4G, increased at 48 hours

compared to NSK) negatively regulates T cell-mediated immunity and may be also linked to

apoptotic cell clearance by macrophages[42]. Of note, the upregulation of CXCL10 chemokine and

its receptor CXCR3 are linked to their major roles of inflammatory reactions, T cell migration, and

IFN-γ in the saliva-mediated process[43]. Besides chemokines, the top expressed molecule was an

antimicrobial peptide hBD-2 (DEFB4A), produced by neutrophils and keratinocytes. DEFB4A is

a marker of inflammation absent in normal skin and responsible for recruitment of memory T cells,

immature DCs, mast cells and neutrophils[44,45].

In this study, the presence of mosquito saliva inhibited pro-inflammatory cytokine production by

'bitten' human skin cells after *in vitro* re-stimulation with saliva. This cytokine-dampening effect

may partly result from the interaction between salivary components and key checkpoints on

different signaling pathways. One example includes the salivary protein LTRIN known to interfere

with the Nf-kB pathway by binding to LTβR, thus hindering its dimerization and leading to

decreased production of inflammatory cytokines[11]. On the other hand, skin immune tolerance to

constant mosquito exposure may also explain the reducing effect of SGE on cytokine production

as seen here in our bite-experienced participants[12,19,46].

With the current rise of vector-borne diseases around the world, future clinical development of

vector-targeted strategies for both known and unknown vector-borne diseases will require a

fundamental understanding of how vector saliva modulates the cutaneous immune response of

humans. Translational studies, such as the one presented here, significantly expand our current

knowledge on human skin immunity and the immune response orchestrated by a 'mosquito bite'.

These newly identified genes, cell types, and processes of human cutaneous immunity can be

leveraged in the development of novel therapeutics and vector-targeted vaccine candidates. Such

new pathogen-agnostic strategies have shown promising results in recent years from vector saliva-

based vaccinations to topical antiviral applications post-bite [47–50]. To build the next generation of

countermeasures against vector-borne disease, nuanced and comprehensive understanding of skin

immunity to both pathogen and vector saliva alike will be required.

388   Materials and methods

389   Study Enrollment and Procedures

From October 2020 to January 2021, we enrolled thirty Cambodian adults at Kampong Speu

District Referral Hospital. Eligible participants were healthy adults between the ages of 18 and 45

years old who agreed not to use topical or oral antihistamines or steroids for the duration of the

study. Participants were excluded if they were pregnant or reported a history of severe allergic

reaction to mosquito or other insect bites, use of medications that affect blood clotting, or history

of significant scarring such as keloid formation (full criteria found on

https://clinicaltrials.gov/ct2/show/NCT04350905).   All participants provided written informed

consent before enrollment, and the study was conducted in accordance with the provisions of the

Declaration of Helsinki and Good Clinical Practice guidelines.

On Day 0, five starved uninfected *Ae. aegypti* female mosquitos were selected from a mosquito

colony dedicated to human feeding studies in the Malaria and Vector Research Laboratory at the

National Center for Parasitology, Entomology and Malaria Control. Mosquitos fed through the

disposable mesh of a feeding device placed on participants' ventral forearm skin for 10 minutes.

Redness and swelling in millimeters were assessed immediately, 15 minutes, 30 minutes, 4 hours

and 48 hours after feeding. Redness and swelling were measured in millimeters at 15 minutes, 30

minutes, and 4 hours.

A total of five 3-mm skin punch biopsies were performed on each individual: one biopsy from

bitten skin per time point (30 minutes, 4 hours, and about 48 hours after feeding is completed) for

a total of three 'bitten' biopsies, and two from NSK collected at baseline. Less than 1cc of 0. 2%

lidocaine without epinephrine was delivered via 30-G needle each chosen 'bite' at the time of

biopsy and 3-mm biopsy was performed on 'bite' adjacent to needle puncture. For feasibility of

skin sample analysis, each participant was assigned to 1 of 3 technical modality cohorts: 1)

immunohistochemistry, 2) RNASeq, and 3) flow cytometry with ten people in each cohort.

Additionally, blood was collected for serological and cellular analyses at baseline and 14 days

after feeding.

***Aedes aegypti* salivary IgG enzyme-linked immunosorbent assays**

416 *Ae. aegypti* SGE (prepared from homogenized, dissected salivary glands) was diluted to a concentration of

417 2 ug/ul in carbonate-bicarbonate buffer (Millipore-Sigma) and used to coat 96-well Immulon 4x HB plates

418 (Nunc) overnight at 4°C. Plates were blocked in a solution of TBS 4 % Bovine serum albumin 0.025 %

419 Tween 20 (Millipore-Sigma). Each plate included sera in duplicate (1:200 dilution in blocking solution)

420 from all time points, a negative control (non-reactive human sera to SGE), an internal control (a pool of

421 positive sera to SGE with optical density (OD) set to 0.25 at 450nm) and blank wells. Secondary antibody

422 [goat anti-human IgG alkaline phosphatase (1:10000 in blocking solution; Millipore-Sigma). A solution of

423 p-nitrophenylphosphate (Millipore-Sigma) was used as substrate and OD measured at 450nm in arbitrary

424 enzyme-linked immunosorbent assay (ELISA) units.

Dengue IgG enzyme-linked immunosorbent assays

426 The PanBio® Dengue Indirect IgG ELISA (product code 01PE30) was performed according to

427 the manufacturer's instructions. Two microtiter plates were supplied, one containing stabilized

428 dengue-1 through dengue-4 (antigen plate) and the other containing either anti-human IgM or anti-

429 human IgG bound to separate wells (assay plate). Peroxidase-labeled anti-dengue virus

430 monoclonal antibody (125 µl/well) was added to the antigen plate to solubilize the antigens and

431 form antibody-antigen complexes. Concurrently, 100 µl of patient serum was diluted 1:100 in the

432 provided diluent and added to each well of the assay plate containing either bound anti-human IgM

433 or bound anti-human IgG, and human IgM or IgG in the patient's serum was captured. The plates

434 were incubated for 1 h at room temperature (antigen plate) or 37°C (assay plate); then the assay

435 plate was washed, and 100 µl of antibody-antigen complexes per well was transferred from the

436 antigen plate to the assay plate. These complexes were captured by dengue virus-specific IgM or

437 IgG during incubation for 1 h at 37°C. The plate was washed, and bound complexes were

438 visualized through the addition of 100 µl of tetramethylbenzidine substrate per well. After 10 min,

439 the reaction was stopped by the addition of 100 µl of 1 M phosphoric acid per well, and the strips

440   were read at 450 nm with a microtiter plate reader. Positivity was determined by comparison to

441   the IgM and IgG reference sera provided (cutoff calibrators). A positive sample was defined as

442   having a sample/calibrator absorbance ratio of ≥1.0, and a negative sample was defined as having

443   a ratio of <1.0.

444   Formalin preservation and H&E staining of skin biopsies

Biopsies were placed in 7ml prefilled 10% formalin vials (Azer Scientific) and stored at 4°C until

paraffin embedded. Unstained slides were cut from paraffin blocks and deparaffinized in xylene,

then hydrated through graded alcohols up to water, placed in Carazzi's hematoxylin, washed in

tap water, and put in one change 95% ethanol. Next, they were placed in eosin-phloxine solution

and then run through graded alcohols to xylene. After xylene, the stained slides were than a cover

slip was mounted using Permount Mounting Media (Fisher Scientific).

451   RNA extraction, RNA-seq library preparation, sequencing and analysis.

Biopsies were stored in DNA/RNA Shield (Zymo research) and kept in 4 °C until processing. Skin

was homogenized using 3mm zirconium beads (Biolink laboratories) and a Magnalyser bead

beater (Roche Diagnostics®). After homogenization total RNA was loaded at InnuPure® C16

touch (AnalitikJena) for magnetic separation of total RNA using an InnuPREP® RNA Kit–IPC16

(AnalitikJena). Quality of extractions was verified on a 4200 TapeStation (Agilent) and all samples

had RIN numbers above seven. RNA sequencing was performed by Novogene. Briefly, mRNA

was enriched by Poly (A) capture and RNA libraries were prepared using the NEBNext Ultra II

Directional RNA Library Prep Kit for Illumina (New England Biolabs) and sequenced on a

NovaSeq 6000 (Illumina) generating paired end reads at 150 bp length. After removing low quality

reads (Trimommatic 0.39)[51], mapping to the human genome (hg38; GCF_000001405.39) and

gene-level quantification were generated by RSEM 1.3.2[52], Bowtie 2-2.4.2[53] and SAMtools

1.11[54]. Normalization and differential gene expression (DGE) analysis were performed using

DESeq2 1.34.0[55], employing the variance stabilizing transformed function (VSN 3.62.0)[56] to the

count matrix, that was designed to account for paired samples enabling expression analyses with

normal skin samples from the same subject as reference. Approximate Posterior Estimation for

GLM (Apeglm 1.16.0) method was used to calculate fold change of DEGs[57]. DEGs were

considered significant if apeglm-estimated shrunken log2fold change was <1 or >1 and with a false

discovery rate <0.05 (Benjamini–Hochberg multiple testing correction method[58]). PlotPCA3D

0.0.1(twbattaglia/btools) was used to generate tri-dimensional principal component analysis plots

on DESeq2. EnhancedVolcano 1.12.0 was used to generate volcano plots[59]. GOplot 1.0.[60] was

used to generate chord plots using data obtained from packages clusterProfiler 4.2.0[61] and

ReactomePA 1.38.0[62]. Pathways with a Benjamini–Hochberg corrected P value <0.05 and were

considered as enriched, selected pathways were mapped on chord plots[58]. Heatmaps were prepared

using package pheatmap 1.0.12.[63] and dendsort 0.3.4[64]. All DEG analyses were performed using

R 4.1.0[65].

Skin biopsy dissociation

Biopsies were transported in RPMI media (Sigma-Aldrich) at 4 °C. Whole skin dissociation kit

(Miltenyi®) was used to perform the biopsies dissociation. An enzyme mix consisting of enzyme

D, enzyme A and buffer L was prepared according to the manufacturer's specifications. Single

punch biopsies were incubated in 400 μl of dissociation mix, at 37°C – 5% CO_2 overnight.

Following incubation, each sample mix was transferred to a Falcon round-bottom tube with cell-

strainer cap (40-75um). Using a rubber pestle and the cell-strainer cap as a mortar, biopsies where

gently macerated to increase cell recovery yield. 500 μl of cold RMPI was used to rinse caps.

Samples were then centrifuged at 1500 rpm 4°C for 10 minutes and the pellet resuspended in 500

µl of cold RPMI. Dissociated skin cells were counted using a pre-programmed setting in a

Countess™ II FL Automated Cell Counter.

Flow cytometry

Dissociated cells were seeded on round bottom 96-well plates and washed with 200 µl of

ThermoFisher's PBS buffer 1x (1500 rpm, 4°C for 5 minutes). Twenty µl of Biolegend Zombie

Aqua fixable viability stain (1:500) was added and incubated for 20 minutes in the dark at 4°C.

Excess viability stain was washed away with 200 µl of PBS/BSA/EDTA buffer (1500 rpm, 4°C

for 5 minutes), and the pellet resuspended and incubated for 10 minutes at 4°C in 4 µl of

Biolegend's FcR blocking antibody (1:10). Cells were then incubated for 30 minutes at 4°C with

surface staining antibody master mix. After washing free unbound antibody with 200 µl of PBS

(1500 rpm, 4°C for 5 minutes), cells were fixed with 100 µl of Biolegend's fixation buffer 1x for

20 minutes at 4°C in the dark. Following fixation, cells were permeabilized and washed with 100

µl of Biolegend perm-wash solution 1x (1500 rpm, 4°C for 5 minutes) and the pellet resuspended

and incubated for 30 minutes at 4°C in intracellular staining antibody master mix. Stained cells

were placed in 200 µl of PBS/BSA/EDTA 1x and visualized on a BD Biosciences FACS Aria II

using FACS Diva software. Analysis was performed using BD Biosciences FlowJo software.

Antibody panels are described in supplemental material (Supplementary Table 5).

Multiplex cytokine screening

The supernatants were collected from 24-hour stimulation cultures of dissociated skin cells and

stored at -20°C. Cytokines in the cell culture supernatants were quantified using Human Th

Cytokine Panel (12-plex) (Cat. 741027, BioLegend) according to the manufacturer's instructions.

All samples in the experiment were measured in duplicate. Briefly, undiluted supernatants were incubated with beads coated with capture antibodies specific for IL-2, 4, 5, 6, 9, 10, 13, 17A, 17F, 22, IFN-γ and TNF-α for 2 hours at room temperature on shaker. After incubation, beads were washed and incubated with biotin-labeled detection antibodies for 1 hour, followed by a final incubation with streptavidin-PE for 30 minutes at room temperature on shaker. Beads were washed and re-suspended with washing buffer. Beads were analyzed by flow cytometry using a FACS Canto cytometer. Analysis was performed using the LEGENDplex analysis software v8.0, which distinguishes between the 12 different analytes on basis of bead size and internal dye.

Statistical analyses

Flow cytometry and multiplex cytokine screening data were analyzed using the non-parametric-based tests, Wilcoxon signed-rank and Chi-square test. Correlation analysis was performed with the Pearson's correlation coefficient. Bars indicate median and interquartile range, and p-values *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$. Statistical analysis of RNAseq data is described above.

Acknowledgements

We would like to thank the study participants in Kampong Speu, Cambodia. We would also like to acknowledge Noemi Kedei and Maria Hernandez for their assistance in histology and staining.

Funding

This work was funded by the Division of Intramural Research at National Institute of Allergy and Infectious Diseases at the National Institutes of Health. T.C. is a HHMI-Wellcome international research scholar (208710/Z/17/Z). D.M is funded by the French National Research Agency (ANR-17-CE15-00029) . D.G. is supported by a Calmette-Yersin PhD scholarship funded by the Institute Pasteur International Network.

Competing interests

The authors declare no competing interests.

Author contributions

Conceptualization: F.O., J.E.M., T.C., J.G.V. Methodology: D.G., H.V, S.N., S.C., J.A.B., F.O.,

J.E.M.. Investigation: D.G., H.V, J.A.B, S.N., S.C., A.P., J.G.V., F.O., J.E.M. Visualization: D.G.,

F.O., J.A.B. Funding acquisition: F.O., J.E.M. Study enrollment and clinical procedures: C.L, S.L.,

R.S, S.S., S.N., J.M, H.K, R.L. Project administration: C.L., T.C., F.O., J.E.M., Supervision: T.C.,

D.M, F.O., J.E.M., C.L, R.L., H.R, H.K. Writing original draft: D.G., H.V., T.C., F.O., J.E.M.,

J.A.B. Writing review and editing: T.C., D.M., C.L., F.O., J.A.B., J.E.M.

All authors contributed to the article and approved the submitted version

Data availability

The Illumina raw reads (fastq files) were loaded at the NCBI Sequence Read Archive with

accession numbers SRR18215008 to SRR18215047 and deposited in GenBank under

bioproject accession code PRJNA812009. All data generated or analyzed during this study are

included in this published article (and its supplementary information files). Clinical trial data is

available at https://clinicaltrials.gov/ct2/show/NCT04350905. Flow cytometry data can be made

available upon reasonable request directly to the authors.

Bibliography

1.   WHO. A global brief on vector-borne diseases. *www.who.int* (2020).

2.   Pierson, T. C. & Diamond, M. S. The continued threat of emerging flaviviruses. *Nat.*

Microbiol. **5**, 796–812 (2020).

3.   Ryan, S. J., Carlson, C. J., Mordecai, E. A. & Johnson, L. R. Global expansion and

redistribution of Aedes-borne virus transmission risk with climate change. *PLoS Negl. Trop.*

Dis. **13**, e0007213 (2019).

4.   Godoy, R. S. M. *et al.* Dengue and Zika virus infection patterns vary among Aedes aegypti

field populations from Belo Horizonte, a Brazilian endemic city. *PLoS Negl. Trop. Dis.* **15**,

e0009839 (2021).

5.   Sun, P. *et al.* A mosquito salivary protein promotes flavivirus transmission by activation of

autophagy. *Nat. Commun.* **11**, 260 (2020).

6.   Ha, Y.-R. *et al.* Detection of Heparin in the Salivary Gland and Midgut of Aedes togoi.

Korean J. Parasitol. **52**, 183–188 (2014).

7.   Schneider, B. S. & Higgs, S. The enhancement of arbovirus transmission and disease by

mosquito saliva is associated with modulation of the host immune response. *Trans. R. Soc.*

Trop. Med. Hyg. **102**, 400–408 (2008).

8.   Guerrero, D., Cantaert, T. & Missé, D. Aedes Mosquito Salivary Components and Their

Effect on the Immune Response to Arboviruses. *Front. Cell. Infect. Microbiol.* **10**, 407

(2020).

9.   Manning, J. E. & Cantaert, T. Time to Micromanage the Pathogen-Host-Vector Interface:

Considerations for Vaccine Development. *Vaccines* **7**, 10 (2019).

10. Vogt, M. B. *et al.* Mosquito saliva alone has profound effects on the human immune system.

PLoS Negl. Trop. Dis. **12**, e0006439 (2018).

11. Jin, L. *et al.* Salivary factor LTRIN from Aedes aegypti facilitates the transmission of Zika

virus by interfering with the lymphotoxin-β receptor. *Nat. Immunol.* **19**, 342–353 (2018).

12. Pingen, M. *et al.* Host Inflammatory Response to Mosquito Bites Enhances the Severity of

Arbovirus Infection. *Immunity* **44**, 1455–1469 (2016).

13. Schneider, B. S., Soong, L., Zeidner, N. S. & Higgs, S. Aedes aegypti salivary gland extracts

modulate anti-viral and TH1/TH2 cytokine responses to sindbis virus infection. *Viral*

Immunol. **17**, 565–573 (2004).

14. Hastings, A. K. *et al.* Aedes aegypti NeSt1 Protein Enhances Zika Virus Pathogenesis by

Activating Neutrophils. *J. Virol.* **93**, e00395-19, /jvi/93/13/JVI.00395-19.atom (2019).

15. Uraki, R. *et al.* Aedes aegypti AgBR1 antibodies modulate early Zika virus infection of

mice. *Nat. Microbiol.* **4**, 948–955 (2019).

16. McCracken, M. K. *et al.* Route of inoculation and mosquito vector exposure modulate

dengue virus replication kinetics and immune responses in rhesus macaques. *PLoS Negl.*

Trop. Dis. **14**, e0008191 (2020).

17. Dudley, D. M. *et al.* Infection via mosquito bite alters Zika virus tissue tropism and

replication kinetics in rhesus macaques. *Nat. Commun.* **8**, 2096 (2017).

18. Trager, W. Acquired Immunity to Ticks. *J. Parasitol.* **25**, 57(1939).

19. Guilbride, D. L., Guilbride, P. D. L. & Gawlinski, P. Malaria's deadly secret: a skin stage.

Trends Parasitol. **28**, 142–150 (2012).

20. Conway, M. J. Type I hypersensitivity promotes Aedes aegypti blood feeding. *Sci. Rep.* **11**,

14891 (2021).

21. Demeure, C. E. *et al.* Anopheles Mosquito Bites Activate Cutaneous Mast Cells Leading to a

Local Inflammatory Response and Lymph Node Hyperplasia. *J. Immunol.* **174**, 3932–3940

(2005).

22. Conway, M. J. *et al.* Mosquito Saliva Serine Protease Enhances Dissemination of Dengue

Virus into the Mammalian Host. *J. Virol.* **88**, 164–175 (2014).

23. Turner, C. T., Lim, D. & Granville, D. J. Granzyme B in skin inflammation and disease.

Matrix Biol. **75–76**, 126–140 (2019).

24. Buzza, M. S. *et al.* Extracellular Matrix Remodeling by Human Granzyme B via Cleavage of

Vitronectin, Fibronectin, and Laminin. *J. Biol. Chem.* **280**, 23549–23558 (2005).

25. Champagne, D. E. Antihemostatic Molecules from Saliva of Blood-Feeding Arthropods.

Pathophysiol. Haemost. Thromb. **34**, 221–227 (2005).

26. Lefteri, D. A. *et al. Mosquito saliva sialokinin-dependent enhancement of arbovirus*

infection through endothelial barrier leakage.

http://biorxiv.org/lookup/doi/10.1101/2021.02.19.431961 (2021)

doi:10.1101/2021.02.19.431961.

27. Bardina, S. V. *et al.* Differential Roles of Chemokines CCL2 and CCL7 in Monocytosis and

Leukocyte Migration during West Nile Virus Infection. *J. Immunol.* **195**, 4306–4318 (2015).

28. Mills, C. D. Anatomy of a Discovery: M1 and M2 Macrophages. *Front. Immunol.* **6**, (2015).

29. Lamana, A. *et al.* CD69 Modulates Sphingosine-1-Phosphate-Induced Migration of Skin

Dendritic Cells. *J. Invest. Dermatol.* **131**, 1503–1512 (2011).

30. Mackay, L. K. *et al.* Cutting Edge: CD69 Interference with Sphingosine-1-Phosphate

Receptor Function Regulates Peripheral T Cell Retention. *J. Immunol.* **194**, 2059–2063

(2015).

31. Rajarathnam, K., Schnoor, M., Richardson, R. M. & Rajagopal, S. How do chemokines

navigate neutrophils to the target site: Dissecting the structural mechanisms and signaling

pathways. *Cell. Signal.* **54**, 69–80 (2019).

32. Pingen, M. *et al.* Host Inflammatory Response to Mosquito Bites Enhances the Severity of

Arbovirus Infection. *Immunity* **44**, 1455–1469 (2016).

33. Dey, R. *et al.* Gut Microbes Egested during Bites of Infected Sand Flies Augment Severity of

Leishmaniasis via Inflammasome-Derived IL-1β. *Cell Host Microbe* **23**, 134-143.e6 (2018).

34. Gough, P., Ganesan, S. & Datta, S. K. IL-20 Signaling in Activated Human Neutrophils

Inhibits Neutrophil Migration and Function. *J. Immunol.* **198**, 4373–4382 (2017).

35. Myles, I. A. *et al.* Signaling via the IL-20 receptor inhibits cutaneous production of IL-1β

and IL-17A to promote infection with methicillin-resistant Staphylococcus aureus. *Nat.*

Immunol. **14**, 804–811 (2013).

36. Shimizu, K. *et al.* PD-1 Imposes Qualitative Control of Cellular Transcriptomes in Response

to T Cell Activation. *Mol. Cell* **77**, 937-950.e6 (2020).

37. Wei, F. *et al.* Strength of PD-1 signaling differentially affects T-cell effector functions. *Proc.*

Natl. Acad. Sci. **110**, E2480–E2489 (2013).

38. Duyen, H. T. L. *et al.* Skin dendritic cell and T cell activation associated with dengue shock

syndrome. *Sci. Rep.* **7**, 14224 (2017).

39. Mommert, S., Schaper, J. T., Schaper-Gerhardt, K., Gutzmer, R. & Werfel, T. Histamine

Increases Th2 Cytokine-Induced CCL18 Expression in Human M2 Macrophages. *Int. J. Mol.*

Sci. **22**, 11648 (2021).

40. Chiffoleau, E. C-Type Lectin-Like Receptors As Emerging Orchestrators of Sterile

Inflammation Represent Potential Therapeutic Targets. *Front. Immunol.* **9**, 227 (2018).

41. Holmskov, U., Thiel, S. & Jensenius, J. C. Collectins and Ficolins: Humoral Lectins of the

Innate Immune Defense. *Annu. Rev. Immunol.* **21**, 547–578 (2003).

42. Tang, L. *et al.* Liver Sinusoidal Endothelial Cell Lectin, LSECtin, NegativelyRegulates

Hepatic T-Cell Immune Response. *Gastroenterology* **137**, 1498-1508.e5 (2009).

43. Kuo, P. T. *et al.* The Role of CXCR3 and Its Chemokine Ligands in Skin Disease and

Cancer. *Front. Med.* **5**, 271 (2018).

44. Niyonsaba, F., Ogawa, H. & Nagaoka, I. Human beta-defensin-2 functions as a chemotactic

agent for tumour necrosis factor-alpha-treated human neutrophils. *Immunology* **111**, 273–281

(2004).

45. Niyonsaba, F., Iwabuchi, K., Matsuda, H., Ogawa, H. & Nagaoka, I. Epithelial cell-derived

human β-defensin-2 acts as a chemotaxin for mast cells through a pertussis toxin-sensitive

and phospholipase C-dependent pathway. *Int. Immunol.* **14**, 421–426 (2002).

46. Rizzo, C. *et al.* IgG1 and IgG4 Antibody Responses to the Anopheles gambiae Salivary

Protein gSG6 in the Sympatric Ethnic Groups Mossi and Fulani in a Malaria Hyperhendemic

Area of Burkina Faso. *PLoS ONE* **9**, e96130 (2014).

47. Manning, J. E. *et al.* Safety and immunogenicity of a mosquito saliva peptide-based vaccine:

a randomised, placebo-controlled, double-blind, phase 1 trial. *The Lancet*

S0140673620310485 (2020) doi:10.1016/S0140-6736(20)31048-5.

48. Oliveira, F. *et al.* A sand fly salivary protein vaccine shows efficacy against vector-

transmitted cutaneous leishmaniasis in nonhuman primates. *Sci. Transl. Med.* **7**, (2015).

49. Sajid, A. *et al.* mRNA vaccination induces tick resistance and prevents transmission of the

Lyme disease agent. *Sci. Transl. Med.* **13**, eabj9827 (2021).

50. Bryden, S. R. *et al.* Pan-viral protection against arboviruses by activating skin macrophages

at the inoculation site. *Sci. Transl. Med.* **12**, eaax2421 (2020).

51. Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina

sequence data. *Bioinformatics* **30**, 2114–2120 (2014).

52. Li, B. & Dewey, C. N. RSEM: accurate transcript quantification from RNA-Seq data with or

without a reference genome. *BMC Bioinformatics* **12**, 323 (2011).

53. Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat. Methods* **9**,

357–359 (2012).

54. Dobin, A. & Gingeras, T. R. Mapping RNA-seq Reads with STAR. *Curr. Protoc.*

Bioinforma. **51**, (2015).

55. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for

RNA-seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).

56. Huber, W., von Heydebreck, A., Sultmann, H., Poustka, A. & Vingron, M. Variance

stabilization applied to microarray data calibration and to the quantification of differential

expression. *Bioinformatics* **18**, S96–S104 (2002).

57. Zhu, A., Ibrahim, J. G. & Love, M. I. Heavy-tailed prior distributions for sequence count

data: removing the noise and preserving large differences. *Bioinformatics* **35**, 2084–2092

(2019).

58. Benjamini, Y. & Hochberg, Y. Controlling the False Discovery Rate: A Practical and

Powerful Approach to Multiple Testing. *J. R. Stat. Soc. Ser. B Methodol.* **57**, 289–300

(1995).

59. Blighe, K., Rana, S. & Lewis, M. Enhanced Volcano: publication-ready volcano plots with

enhanced colouring and labeling. (2021).

60. Walter, W., Sánchez-Cabo, F. & Ricote, M. GOplot: an R package for visually combining

expression data with functional analysis: Fig. 1. *Bioinformatics* **31**, 2912–2914 (2015).

61. Yu, G., Wang, L.-G., Han, Y. & He, Q.-Y. clusterProfiler: an R Package for Comparing

Biological Themes Among Gene Clusters. *OMICS J. Integr. Biol.* **16**, 284–287 (2012).

62. Yu, G. & He, Q.-Y. ReactomePA: an R/Bioconductor package for reactome pathway

analysis and visualization. *Mol. Biosyst.* **12**, 477–479 (2016).

63. Kolde, R. Pretty Heatmaps. R package version.

64. Sakai, R., Winand, R., Verbeiren, T., Moere, A. V. & Aerts, J. dendsort: modular leaf

ordering methods for dendrogram representations in R. *F1000Research* **3**, 177 (2014).

65. R Core Team (2021). *R: A language and environment for statistical computing. R*

Foundation fo Statistical Computing, Vienna, Austria. (The R Foundation).

694