The regulation of circadian rhythm by insulin signaling in *Drosophila*

Sho T Yamaguchi, Jun Tomita, Kazuhiko Kume*

Department of Neuropharmacology, Graduate School of Pharmaceutical Sciences,

Nagoya City University, Nagoya 467-8603, Japan

* Correspondence should be addressed to :

kume.kazuhiko@gmail.com, kume@phar.nagoya-cu.ac.jp

Tanabe 3-1, Mizuho, Nagoya 467-8603 Japan

Key words: circadian rhythm, metabolism, insulin, insulin receptor, Drosophila

Acknowledgments

We thank Dr. Taishi Yoshii, BDSC, and VDRC for fly stocks, and the members of Kume lab for discussions.

Funding

This study was supported by JSPS, Japan: Kazuhiko Kume 18H02481, 21H02529; Jun Tomita 20K06744.

Author Contributions

SY, JT, and KK designed the experiments. SY conducted all experiments and data analysis. SY, JT, and KK wrote the manuscript.

Competing Interests

The authors declare no conflict of interests.

Abstract

Circadian rhythm is well conserved across species and relates to numerous biological functions. Circadian

misalignment impairs metabolic function. Insulin signaling is a key modulator of metabolism in the fruit fly as well

as mammals and its defects cause metabolic disease. Daily diet timing affects both circadian rhythmicities of behavior

and metabolism. However, the relationship between circadian clock and insulin signaling is still elusive. Here, we

report that insulin signaling regulates circadian rhythm in *Drosophila melanogaster*. We found the insulin receptor

substrate mutant, *chico[1]*, showed a shorter free-running circadian period. The knockdown of *insulin receptor* (*InR*),

or another signaling molecule downstream of *InR*, *dp110*, or the expression of a dominant-negative form of InR

resulted in the shortening of the circadian period and diminished its amplitude. The impairment of insulin signaling

both in all neurons and restricted circadian clock neurons altered circadian period length, indicating the insulin

signaling plays a role in the regulation of circadian rhythm in clock cells. Among 3 insulin-like ligands expressed in

the brain, *dilp5* showed the largest effect on circadian phenotype when deleted. These results suggested that insulin

signaling contributes to the robustness of the circadian oscillation and coordinates metabolism and behavior.

Highlights

● Insulin receptor substrate mutant, *chico[1]*, displayed circadian rhythm phenotype.

● Pan-neuronal inhibition of insulin receptor signaling shortened circadian cycle.

● Inhibition of insulin signaling only in clock neurons altered circadian cycle.

● Dilp5 is a major insulin receptor ligand for circadian effects.

Introduction

Intrinsic circadian rhythms exist in many species and are involved in a variety of physiological phenomena (Dunlap, 1999; Green et al., 2008; Roenneberg and Merrow, 2005). In humans, circadian misalignment has been shown to cause changes in gene expression as well as affect metabolic functions such as plasma insulin and glucose (Archer et al., 2014; Scheer et al., 2009). It has been suggested that such changes due to disruption of circadian rhythm increase the risk of obesity and diabetes (Depner et al., 2014; Noh, 2020).

The fruit fly, *Drosophila melanogaster*, has been used as a genetic model organism to reveal genes regulating circadian rhythm and sleep (Bedont et al., 2021; Dubowy and Sehgal, 2017; Toda et al., 2019; Tomita et al., 2017). We pioneered its sleep regulation (Kume et al., 2005; Tomita et al., 2015, 2011; Ueno et al., 2012; Ueno and Kume, 2014). To date, we examined the relationship between sleep and metabolism or feeding behavior and found that c-Jun N terminal kinase, Sik3, and nutrition or taste of the food function to regulate sleep (Funato et al., 2016; Hasegawa et al., 2017; Takahama et al., 2012; Yamazaki et al., 2012).

Metabolism and feeding behavior tightly relate to not only sleep but also circadian rhythm. For example, null mutation of clock genes disrupts glucose homeostasis and lipid metabolism as in mammals (Rudic et al., 2004; Seay and Thummel, 2011; Turek et al., 2005). Furthermore, feeding with a high-fat diet or high sucrose diet affected clock gene expression or daily activity profile (Lee et al., 2021; Nayak and Mishra, 2021). However, the molecular mechanisms of how metabolism affects circadian rhythm are still elusive.

Insulin signaling, a critical regulator of metabolism in mammals, is well conserved in flies (Nässel et al., 2015; Nässel and Broeck, 2016). Flies have eight mammalian insulin homologs, *drosophila insulin-like peptides* (*dilps*),

one *insulin receptor* (*InR*), and one mammalian insulin receptor substrate homolog, *chico*. These molecules have

been shown to regulate not only metabolism (Rajan and Perrimon, 2012; Ugrankar et al., 2015) and development

(Böhni et al., 1999; Brogiolo et al., 2001; Fernandez et al., 1995; Sano et al., 2015) but also behaviors, such as

memory, feeding behavior, courtship, and sleep (Barber et al., 2021; Metaxakis et al., 2014; Naganos et al., 2016;

Tanabe et al., 2017; Watanabe and Sakai, 2016). In addition to these reports, previous research has shown that *Akt*

and *target of rapamycin* (*TOR*), downstream components of insulin signaling, affect circadian rhythm (Zheng and

Sehgal, 2010). Although the result suggested that insulin signaling regulates circadian rhythm, it is not clear whether

InR and *chico* in the central nervous system regulate circadian rhythm. In this study, we aimed to clarify the regulation

of circadian rhythm by neuronal insulin signaling.

Material & Methods

Fly strains and culture conditions

Flies (*Drosophila melanogaster*) were reared with conventional food, which contained yeast, glucose, cornmeal,

wheat germ, and agar, at 24.5°C under a 12-h:12-h light: dark (LD) cycle as described before (Kume et al., 2005).

The following stocks were obtained from the Bloomington Stock Center, Indiana University: *elav*-Gal4 (Stock

number: 458), *tim*-Gal4 (7126), UAS-*InR* DN (8252), *chico^1* (10738), *dilp2* KO (30881), *dilp3* KO (30882), *dilp5*

KO (30884), *dilp2*, *3*, and *5* KO (30889), *y^1 v^1*; *P{CaryP}attP2* (36303), *y^1 v^1*; *P{CaryP}attP40* (36304), UAS-*chico*

RNAi (36665), UAS-*InR* RNAi (51518), and UAS-*dp110* RNAi (61182). *DvPdf*-Gal4 (Bahn et al., 2009) and *Pdf*-

Gal4 was provided by Dr. Taishi Yoshii. *Pdf*-GS was gifted from Dr. Takaomi Sakai. UAS-*dilp5* RNAi (VDRC

49520), UAS-*Dicer-2* (60008), and w^{1118} (60000) which is the genetic background of the UAS-*dilp5* RNAi line were

purchased from Vienna Drosophila RNAi Center (VDRC) (Dietzl et al., 2007). *elav*-Gal4, *tim*-Gal4, *dilp2* KO, *dilp3*

KO, *dilp5* KO, and UAS-*InR* DN were backcrossed to the control strain (w^{1118}) at least five times.

Circadian behavioral analysis

The two- to five-days-old male flies were loaded into a glass tube (length, 65 mm; inside diameter, 3 mm) with

sucrose food (1% agar supplemented with 5% sucrose) at one end and were entrained for at least 3 days under LD

cycle conditions before being transferred to constant dark (DD) conditions. In the experiment with *Pdf*-GS, the two-

to five-days-old male flies were loaded into a glass tube with sucrose food containing vehicle (EtOH, - RU486) or

100 µM RU486 (+ RU486) at one end and transferred to DD conditions. Data were collected in 1 min bin for 10 days

under DD conditions using the *Drosophila* Activity Monitor system (Trikinetics). The period length and amplitude

of circadian rhythm were calculated using locomotor activity data for 9 or 10 days by Fourier transform (FT) with a

custom-made R code. Flies with the maximal value of the power spectrum calculated by the FT less than 0.04 were

considered arrhythmic and were removed from the calculation of the mean of circadian period length. The data from

the experiments of the *InR* DN expression with *elav*-Gal4, *tim*-Gal4, and *DvPdf*-Gal4, or the progenies from crosses

between *dilps* KO and TKO were also used in the published paper (Yamaguchi et al., 2021). However, the analysis

methods are completely different between this study and the published paper.

mRNA analysis with quantitative RT-PCR

mRNA levels were examined by quantitative RT-PCR. The 2- to 5-days-old male flies were collected at

zeitgeber time 6. Fifteen to thirty heads were divided into three and used for total RNA extraction with RNAiso

Plus (TAKARA) according to the manufacturer's instruction. cDNA was synthesized from the total RNA using

ReverTra Ace qPCR Master Mix with gDNA Remover (Toyobo), then used for quantitative RT-PCR using

THUNDERBIRD SYBR qPCR Master Mix (Toyobo). *GAPDH2* expression levels were used as an internal control.

The primers were 5'-AGGAAGGAAAAGCGGAAAAG-3' and 5'-GGGGAGGATAAACGAGGTGT-3' for

GAPDH2, 5'-GCTTCGCTGCATTTCAACTT-3' and 5'-AGTGTCCCAGGCGCTTATTA-3' for *chico*, 5'-

GGACAAGGAGCGAATCAAAC-3' and 5'-TGCAGTTCTCCAGCTGATTG-3' for *InR*, 5'-

CCATCCGGTGCTTACACTTT-3' and 5'-AGATCGCTTTCCAGGTACGA-3' for *dp110*, 5'-

CCAGGAAGGTTGTCATCTCG-3' and 5'-GAAAGCCCGCTCGTAGATAAG-3' for *4E-BP*, 5'-

TGAGTATGGTGTGCGAGGAG-3' and 5'-ACAAACTGCAGGGGATTGAG-3' for *dilp2*, 5'-

AGAGAACTTTGGACCCCGTGAA-3' and 5'-TGAACCGAACTATCACTCAACAGTCT-3' for *dilp3*, 5'-

GAGGCACCTTGGGCCTATTC-3' and 5'-CATGTGGTGAGATTCGGAGCTA-3' for *dilp5*. The primer

sequences for *dilp3* and *dilp5* are referred from a previous report (Broughton et al., 2005). Each mRNA expression

level was normalized to GAPDH2, then normalized to the average of three independent control samples.

Statistical analysis

Data were analyzed as described in the figure legend using Microsoft Excel and the freely available statistical

software package R 3.5.3 (https://www.r-project.org/).

Results

***chico[1]* mutation caused a shorter circadian period.**

To investigate the relationship between insulin signaling and circadian rhythm, we used the genetic mutant

related to insulin signaling. Because homozygous *InR* null mutant is lethal, we used *chico[1]*, *chico* mutant line which

has P element insertion at 80 bp downstream of translation start site and this insertion cause complete loss of chico

protein to examine the effect of InR signal on circadian rhythm (Böhni et al., 1999; Naganos et al., 2012). To remove

the genetic background effect, we outcrossed *chico[1]* for five generations into *w[1118]*, but the homozygous outcrossed

chico[1] mutant was lethal. Therefore, we measured the activity of the heterozygous mutants of outcrossed or original

chico[1] with *w[1118]* (*chico[1]* / + (BC5), *chico[1]* / + (F1)) in addition to the homozygous mutants of original *chico[1]* to

investigate the effect of *chico[1]* mutation. *chico[1]* showed a shorter circadian period length (mean ± SEM; 1395 ± 4.78)

than the other groups (Fig. 1A, B), although the results were not directly comparable due to the difference in genetic

background. Nonetheless, there was little difference in circadian period length between the *chico[1]* / + (BC5) (1421 ±

4.05) and *chico[1]* / + (F1) (1420 ± 2.01), implying the little genetic background effect on circadian period length.

Moreover, *chico[1]* / + (BC5) tended to have a shorter circadian period length than *w[1118]* (+, 1433 ± 4.95). These results

suggested that *chico* relates to the regulation of circadian rhythm and its defect leads to a shorter circadian period.

Insulin signaling in neurons, especially in the clock neurons, regulates circadian rhythm.

To determine whether the neuronal insulin signaling regulates circadian rhythm, we inhibited the insulin

signaling in specific regions using Gal4 / upstream activation sequence (UAS) system (Brand and Perrimon, 1993).

First, we used pan-neuronal Gal4 driver, *elav*-Gal4, and UAS-*chico* RNAi to obtain the flies with pan-neuronal

knockdown of *chico*. The results showed that the circadian period length was significantly shortened in flies with

pan-neuronal knockdown of *chico* (Fig. 2A, Sup. Fig. 1A). Also, we conducted the expression of a dominant-negative

form of *InR* (*InR* DN) and knockdown of *InR* or *dp110* (*Phosphatidylinositol 3-kinase 92E*), one of the downstream

of insulin signaling, to validate the effect of insulin signaling inhibition on circadian rhythm. These manipulations

with *elav*-Gal4 caused shorter circadian period length as well as pan-neuronal knockdown of *chico* (Fig. 2A, Sup.

Fig. 1B-D). In addition, pan-neuronal knockdown of *InR* caused decreased amplitude of circadian rhythm (Fig. 2B).

To confirm the efficiency of RNAi, we quantified the mRNA level of each gene and *4E-BP*, the expression of which

is negatively regulated by insulin signaling (Jünger et al., 2003; Puig et al., 2003). Although *4E-BP* mRNA level was

significantly increased only by the pan-neuronal knockdown of *chico*, the mRNA level of *chico* or *dp110* was

significantly declined in flies with pan-neuronal knockdown of *chico* or *dp110*, respectively (Sup. Fig. 2A, C).

Although we couldn't detect any change of *InR* mRNA level in flies with pan-neuronal knockdown of *InR*, there is a

significant increase of *4E-BP* mRNA level in these flies (Sup. Fig. 2B, D), suggesting the inhibition of insulin

signaling. Taken together, insulin signaling in the brain could regulate circadian rhythm.

Since inhibition of neuronal insulin signaling affected behavioral rhythm, we focused on the clock neurons,

which express clock genes and are essential for the generation of circadian rhythms. (Dubowy and Sehgal, 2017). To

manipulate the insulin signaling in the clock neurons, we employed *tim*-Gal4 which expresses GAL4 under the

promoter of *timeless*, one of the clock genes in *Drosophila* (Myers et al., 1995; Sehgal et al., 1995). Inhibition of

insulin signaling with *tim*-Gal4 shortened circadian period length (Fig. 2C, Sup. Fig. 3A-D). Moreover, the expression

of *InR* DN or knockdown of *InR* declined the amplitudes of the circadian rhythm (Fig. 2D). These results indicated

that insulin signaling in clock neurons regulates circadian rhythm.

Among the clock neurons, ventral and dorsal lateral neurons (LNvs and LNds) are recently demonstrated to be

the two key components of the clock neuron network (Delventhal et al., 2019; Schlichting et al., 2019). These studies

showed that the lack of a clock gene both in LNvs and LNds caused complete loss of behavioral rhythmicity and the

molecular clock either in LNvs or in LNds is sufficient to generate the behavioral circadian rhythm. On the other

hand, LNvs themselves are known to be necessary to synchronize the clock neuron network via the release of a

principal circadian messenger, pigment dispersing factor (PDF) (Renn et al., 1999). To validate whether insulin

signaling both in LNvs and LNds or only in LNvs relates to circadian rhythm regulation, inhibition of insulin

signaling was conducted by *DvPdf*-Gal4 and *Pdf*-Gal4. As a result, knockdown of *chico*, *InR* or *dp110* by *DvPdf*-

Gal4 shortened circadian period length (Fig. 3A, Sup. Fig. 4A-D). Although expression of *InR* DN with *DvPdf*-Gal4

did not significantly change period length, it caused arrhythmic-like behavior after four to five days of DD conditions

(Sup. Fig. 4B). Furthermore, all these manipulations with *DvPdf*-Gal4 decreased the amplitudes of the circadian

rhythm (Fig. 3B). In addition, the knockdown of *InR* or *dp110* with *Pdf*-Gal4 shortened circadian period length (Fig.

3C, Sup. Fig. 5A-D). The *InR* knockdown by *Pdf*-Gal4 weakened the circadian rhythm amplitude (Fig. 3D). On the

other hand, both the knockdown of *chico* and the expression of *InR* DN with *Pdf*-Gal4 caused a longer circadian

period length (Fig. 3C). These results indicated that insulin signaling in LNvs contribute to maintain the proper

behavioral rhythm.

Since insulin signaling relates to development and growth (Naganos et al., 2012; Song et al., 2003), constitutive

inhibition of insulin signaling could cause behavioral changes via some developmental defects. To avoid inhibition

of insulin signaling during development and growth stages, we used a drug-inducible gene expression system, Gene-

Switch (GS) system (McGuire et al., 2004). Knockdown of *chico* and expression of *InR* DN specifically at the adult

stage were achieved by using *Pdf*-GS. Without RU486, flies with UAS-*chico* RNAi or UAS-*InR* DN did not show

significant changes in circadian period length (Fig. 4A, B, Sup. Fig. 6A, B). With RU486, flies with *chico* knockdown

or expression of *InR* DN by *Pdf*-GS showed significantly longer circadian period length (Fig. 4A, B). Since the

amplitudes from flies with UAS-*chico* RNAi or UAS-*InR* DN are different from each control without RU486, there

might be some genetical background effects. These results indicated that insulin signaling in LNvs relates to circadian

rhythm regulation, at least in part, independently of its developmental function.

dilp5 contributes to the regulation of circadian rhythm via insulin signaling

Next, we focused on *dilp2*, *dilp3*, and *dilp5*, which are expressed in the brain, among the eight *dilps* (Nässel et

al., 2015; Nässel and Broeck, 2016). To inhibit the compensational expression of other *dilps* in a knockout mutant of

one *dilp* (Grönke et al., 2010), we crossed every single knockout mutant of *dilp2* (*dilp2* KO), *dilp3* (*dilp3* KO), or

dilp5 (dilp5 KO) with a triple knockout mutant of *dilp2*, *dilp3*, and *dilp5* (TKO) and analyzed the flies of one *dilp*

homozygous KO with two heterozygous KO. Among these mutants, only *dilp5* KO (TKO x *dilp5* KO) showed a

significantly shorter circadian period length than control (TKO crossed with w^{1118}, TKO x w^{1118}) (Fig. 5A, Sup. Fig.

7). As we previously reported, each heterozygous mutant expressed an excess amount of *dilp* other than

systematically knockout one (Yamaguchi et al., 2021). To elucidate the effect of systematically *dilps* knockout on

insulin signaling, the mRNA level of *4E-BP* was quantified. *4E-BP* expression was significantly increased only in

TKO x *dilp5* KO, suggesting that insulin signaling is inhibited in TKO x *dilp5* KO (Fig. 4B). On the other hand, *4E-*

BP mRNA level did not change in the heterozygous mutant of TKO and *dilp2* KO (TKO x *dilp2* KO) or *dilp3* KO

(TKO x *dilp3* KO), despite their mutation in *dilps*. To confirm the function of *dilp5* in the regulation of circadian

rhythm, we knocked down *dilp5* with *elav*-Gal4. Accordingly, flies with pan-neuronal knockdown of *dilp5* showed

a significant reduction of circadian period length without any changes of mRNA level of *dilp2* or *dilp3* (Fig. 5C, Sup.

Fig. 8). These results indicated the regulation of circadian rhythm by *dilp5*.

Discussion

In this study, we investigated the relationship between circadian rhythm and insulin signaling. To clarify the

relationship, we conducted inhibition of insulin signaling in the targeted region. We also analyzed the activity of the

dilps mutants to specify the ligand involved in the regulation of circadian rhythm. Through these experiments, we

showed that *InR* and its downstream molecules in the clock neurons regulate circadian rhythm. In addition, we

discovered that *dilp5* contributes to the regulation of circadian rhythm. These findings support the relationship

between circadian rhythm and metabolism in *Drosophila melanogaster*.

Specifically, we examined the effect of inhibition of insulin signaling on circadian rhythm by the expression of

a dominant-negative form of *InR*, *InR* DN, and by the knockdown of *chico*, *InR*, and *dp110*. All of these manipulations

with *elav*-Gal4 and *tim*-Gal4 drivers shortened circadian period length (Fig. 2A, Fig. 2C), although we couldn't detect

a significant change of *InR* expression in flies with pan-neuronal knockdown of *InR*. It may be because *InR* is

expressed in various tissues, including the neurons, glia, and fat body. Knockdown of the three genes with *DvPdf*-

Gal4 also caused the same phenotype in period length (Fig. 3A). Also, knockdown of *InR* or *dp110* by *Pdf*-Gal4

shortened circadian period length. On the other hand, knockdown of *chico* or expression of *InR* DN by *Pdf*-Gal4

lengthened period length (Fig. 3C). In previous research, flies with silencing of *TOR* in the cells expressing *period*,

one of the core clock genes, showed the altered circadian period length (Kijak and Pyza, 2017). Other paper reported

that activation of *Akt* and *TOR* signaling in LNvs lengthened circadian period length(Zheng and Sehgal, 2010).

Although the detailed distribution of *InR* is poorly understood to date, cell-specific RNA sequencing of each clock

cell detected mRNA expression of the insulin signaling molecules in the clock neurons (Abruzzi et al., 2017; Ma et

al., 2021). These results support the idea that neuronal insulin signaling, especially in the clock neurons, regulates

circadian rhythm.

In our experiment, most of the neuronal insulin signaling inhibition altered circadian period length. Although

only the expression of *InR* DN with *DvPdf*-Gal4 did not cause significant changes in the circadian period length (Fig.

3A), these flies seem to have a shorter circadian period length than control according to the activity in the first 4-

days of DD condition (Sup. Fig. 4B). Also, these flies showed arrhythmic-like behavior after the 4-5 days of DD

condition. Such phenotype might cause a loss of accuracy for quantification of circadian period length. Contrary to

the shortened circadian period length in the knockdown of *InR* or *dp110* by *Pdf*-Gal4, the knockdown of *chico* and

the expression of *InR* DN with *Pdf*-Gal4 lengthened circadian period length (Fig. 3C). There are some possible causes

of these contradictions, such as the genetic background, the strength of insulin signaling inhibition, or the

developmental defects. Although there were some contradictions, we speculated that the insulin signaling in the clock

neurons contribute to maintain the proper behavioral rhythm.

Recently, Schlichting et al. reported that knockout of *period* in all or subset of lateral neurons differently affected

circadian period length. In addition, Schlichting et al. and Delventhal et al. demonstrated that the clock neurons act

not in a hierarchy but as a distributed network to regulate circadian activity (Delventhal et al., 2019; Schlichting et

al., 2019). In these papers, cell-specific knockout of clock genes revealed that the clock in each clock neuron,

especially in LNvs and LNds, collaborate to achieve the close to 24-hr period length. Inhibition of insulin signaling

in the lateral neurons might diminish such communication between the lateral neurons. Although inhibition of insulin

signaling by *elav*-Gal4 and *tim*-Gal4 did not cause severe defects in circadian behavior as much as *DvPdf*-Gal4 did,

it might be due to the difference of Gal4 expression levels in LNvs and LNds. Furthermore, *shaggy* (*sgg*, homolog

of *GSK3*), which is negatively regulated by insulin signaling (Papadopoulou et al., 2004), is known to phosphorylate

clock protein and manipulate the intrinsic speed of the molecular clock (Martinek et al., 2001). Additionally, Zhang

and Sehgal reported that *Akt* and *TOR* modulate the timing of nuclear expression of TIM in the clock neurons (Zheng

and Sehgal, 2010). Although we did not assess whether our manipulation of insulin signaling changes the core clock

gene behavior or neural communication, it is possible that insulin signaling modulated the molecular clock via the

regulation of *sgg* activity as previous research reported. Further experiments are needed to clarify the effect of insulin

signaling on the clock gene expression rhythm or the clock neuron activity.

Since insulin signaling is important for development, the inhibition of insulin signaling with constitutive Gal4

might induce some developmental defects (Böhni et al., 1999; Naganos et al., 2012; Song et al., 2003). However, we

demonstrated that the adult stage-specific insulin signaling inhibition in LNvs by *Pdf*-GS altered behavioral rhythm

(Fig. 4A, B). These results suggested that insulin signaling, at least in LNvs, regulates circadian rhythm in a

development-independent manner.

We demonstrated that *dilp5* alone among three insulin ligands tested showed a significant effect on the circadian

period length when only one of the *dilps* was systemically knocked out or knocked down in neurons. An earlier study

reported that *dilp* single mutation caused compensational expression of other *dilps* and little change of *4E-BP*

expression (Grönke et al., 2010). We previously reported the similar compensational effects in the progenies from

crosses between TKO and single KO mutants (Yamaguchi et al., 2021). Especially, *dilp2* mRNA level in the mutants

of *dilp5* was significantly increased. *dilp2* mRNA increase was also observed in the mutants of *dilp3* which did not

show any changes in circadian period length. In addition, pan-neuronal knockdown of *dilp5* shortened circadian

period length without an increase of other *dilps* mRNA (Fig. 5F, Sup. Fig. 8B). According to these results, we

conjectured that increase of *dilp2* mRNA did not cause the changes of circadian period length. However, it is still

possible that *dilp2* or *dilp3* plays a role in the regulation of circadian rhythm. It should also be noted that *dilps*,

especially *dilp2*, *dilp3*, and *dilp5*, are thought to have a functional redundancy (Nässel et al., 2015). Nevertheless, we

demonstrated that *dilp5* has a significant effect on circadian rhythm.

In conclusion, our results suggest that neuronal insulin signaling contributes to the regulation of robust circadian

rhythm in flies. In mammals, previous studies showed the relationship between insulin signaling and circadian rhythm

(Boden et al., 1996; Marcheva et al., 2010; Sato et al., 2014). Recently, Crosby et al. demonstrated that insulin and

insulin-like growth factor 1 are important for the entrainment of behavioral rhythms to feeding time and have some

activity in the suprachiasmatic nucleus, the master clock of mammals (Crosby et al., 2019). Taken together, our results

shed light on the conserved role of insulin and insulin signaling in the regulation of circadian rhythm from flies to

mammals.

References

Abruzzi, K.C., Zadina, A., Luo, W., Wiyanto, E., Rahman, R., Guo, F., Shafer, O., Rosbash, M., 2017. RNA-seq

analysis of Drosophila clock and non-clock neurons reveals neuron-specific cycling and novel candidate

neuropeptides. PLoS Genetics 13, e1006613. https://doi.org/10.1371/journal.pgen.1006613

Archer, S.N., Laing, E.E., Möller-Levet, C.S., van der Veen, D.R., Bucca, G., Lazar, A.S., Santhi, N., Slak, A., Kabiljo,

R., von Schantz, M., Smith, C.P., Dijk, D.J., 2014. Mistimed sleep disrupts circadian regulation of the human

transcriptome. Proc Natl Acad Sci U S A 111. https://doi.org/10.1073/pnas.1316335111

Bahn, J.H., Lee, G., Park, J.H., 2009. Comparative analysis of Pdf-mediated circadian behaviors between Drosophila

melanogaster and D. virilis. Genetics 181, 965–975. https://doi.org/10.1534/genetics.108.099069

Barber, A.F., Fong, S.Y., Kolesnik, A., Fetchko, M., Sehgal, A., 2021. Drosophila clock cells use multiple

mechanisms to transmit time-of-day signals in the brain. Proc Natl Acad Sci U S A 118, 1–8.

https://doi.org/10.1073/pnas.2019826118

Bedont, J.L., Toda, H., Shi, M., Park, C.H., Quake, C., Stein, C., Kolesnik, A., Sehgal, A., 2021. Short and long

sleeping   mutants   reveal   links   between   sleep   and   macroautophagy.   Elife   10,   1–27.

https://doi.org/10.7554/eLife.64140

Boden, G., Ruiz, J., Urbain, J.L., Chen, X., 1996. Evidence for a circadian rhythm of insulin secretion. American

Journal   of   Physiology-Endocrinology   and   Metabolism   271,   E246–E252.

https://doi.org/10.1152/ajpendo.1996.271.2.E246

Böhni, R., Riesgo-Escovar, J., Oldham, S., Brogiolo, W., Stocker, H., Andruss, B.F., Beckingham, K., Hafen, E.,

1999. Autonomous control of cell and organ size by CHICO, a Drosophila homolog of vertebrate IRS1-4. Cell

97, 865–875. https://doi.org/10.1016/S0092-8674(00)80799-0

Brand, a H., Perrimon, N., 1993. Targeted gene expression as a means of altering cell fates and generating dominant

phenotypes. Development 118, 401–15.

Brogiolo, W., Stocker, H., Ikeya, T., Rintelen, F., Fernandez, R., Hafen, E., 2001. An evolutionarily conserved

function of the drosophila insulin receptor and insulin-like peptides in growth control. Current Biology 11, 213–

221. https://doi.org/10.1016/S0960-9822(01)00068-9

Broughton, S.J., Piper, M.D.W., Ikeya, T., Bass, T.M., Jacobson, J., Driege, Y., Martinez, P., Hafen, E., Withers, D.J.,

Leevers, S.J., Partridge, L., 2005. Longer lifespan, altered metabolism, and stress resistance in Drosophila from

ablation of cells making insulin-like ligands. Proc Natl Acad Sci U S A 102, 3105–3110.

https://doi.org/10.1073/pnas.0405775102

Crosby, P., Hamnett, R., Putker, M., Hoyle, N.P., Reed, M., Karam, C.J., Maywood, E.S., Stangherlin, A., Chesham,

J.E., Hayter, E.A., Rosenbrier-Ribeiro, L., Newham, P., Clevers, H., Bechtold, D.A., O'Neill, J.S., 2019.

Insulin/IGF-1 Drives PERIOD Synthesis to Entrain Circadian Rhythms with Feeding Time. Cell 177, 896-

909.e20. https://doi.org/10.1016/j.cell.2019.02.017

Delventhal, R., O'Connor, R.M., Pantalia, M.M., Ulgherait, M., Kim, H.X., Basturk, M.K., Canman, J.C., Shirasu-

Hiza, M., 2019. Dissection of central clock function in Drosophila through cell-specific CRISPR-mediated

clock gene disruption. Elife 8, 1–14. https://doi.org/10.7554/eLife.48308

Depner, C.M., Stothard, E.R., Wright, K.P., 2014. Metabolic consequences of sleep and circadian disorders. Current

Diabetes Reports 14, 1–15. https://doi.org/10.1007/s11892-014-0507-z

Dietzl, G., Chen, D., Schnorrer, F., Su, K.C., Barinova, Y., Fellner, M., Gasser, B., Kinsey, K., Oppel, S., Scheiblauer,

S., Couto, A., Marra, V., Keleman, K., Dickson, B.J., 2007. A genome-wide transgenic RNAi library for

conditional gene inactivation in Drosophila. Nature 448, 151–156. https://doi.org/10.1038/nature05954

Dubowy, C., Sehgal, A., 2017. Circadian rhythms and sleep in Drosophila melanogaster. Genetics 205, 1373–1397.

https://doi.org/10.1534/genetics.115.185157

Dunlap, J.C., 1999. Molecular bases for circadian clocks. Cell 96, 271–90. https://doi.org/10.1016/s0092-

8674(00)80566-8

Fernandez, R., Tabarini, D., Azpiazu, N., Frasch, M., Schlessinger, J., 1995. The Drosophila insulin receptor

homolog: A gene essential for embryonic development encodes two receptor isoforms with different signaling

potential. EMBO Journal 14, 3373–3384. https://doi.org/10.1002/j.1460-2075.1995.tb07343.x

Funato, H., Miyoshi, C., Fujiyama, T., Kanda, T., Sato, M., Wang, Z., Ma, J., Nakane, S., Tomita, J., Ikkyu, A.,

Kakizaki, M., Hotta-Hirashima, N., Kanno, S., Komiya, H., Asano, F., Honda, T., Kim, S.J., Harano, K.,

Muramoto, H., Yonezawa, T., Mizuno, S., Miyazaki, S., Connor, L., Kumar, V., Miura, I., Suzuki, T., Watanabe,

A., Abe, M., Sugiyama, F., Takahashi, S., Sakimura, K., Hayashi, Y., Liu, Q., Kume, K., Wakana, S., Takahashi,

J.S., Yanagisawa, M., 2016. Forward-genetics analysis of sleep in randomly mutagenized mice. Nature 539,

378–383. https://doi.org/10.1038/nature20142

Green, C.B., Takahashi, J.S., Bass, J., 2008. The Meter of Metabolism. Cell 134, 728–742.

https://doi.org/10.1016/j.cell.2008.08.022

Grönke, S., Clarke, D.-F., Broughton, S., Andrews, T.D., Partridge, L., 2010. Molecular evolution and functional

characterization of Drosophila insulin-like peptides. PLoS Genetics 6, e1000857.

https://doi.org/10.1371/journal.pgen.1000857

Hasegawa, T., Tomita, J., Hashimoto, R., Ueno, T., Kume, S., Kume, K., 2017. Sweetness induces sleep through

gustatory signalling independent of nutritional value in a starved fruit fly. Scientific Reports 7, 14355.

https://doi.org/10.1038/s41598-017-14608-1

Jünger, M.A., Rintelen, F., Stocker, H., Wasserman, J.D., Végh, M., Radimerski, T., Greenberg, M.E., Hafen, E.,

2003. The Drosophila forkhead transcription factor FOXO mediates the reduction in cell number associated

with reduced insulin signaling. Journal of Biology 2, 20. https://doi.org/10.1186/1475-4924-2-20

Kijak, E., Pyza, E., 2017. TOR signaling pathway and autophagy are involved in the regulation of circadian rhythms

in behavior and plasticity of L2 interneurons in the brain of Drosophila melanogaster. PLoS ONE 12, e0171848.

https://doi.org/10.1371/journal.pone.0171848

Kume, K., Kume, S., Park, S.K., Hirsh, J., Jackson, F.R., 2005. Dopamine is a regulator of arousal in the fruit fly.

Journal of Neuroscience 25, 7377–7384. https://doi.org/10.1523/JNEUROSCI.2048-05.2005

Lee, S.H., Cho, E., Yoon, S.E., Kim, Y., Kim, E.Y., 2021. Metabolic control of daily locomotor activity mediated by

tachykinin in Drosophila. Communications Biology 4. https://doi.org/10.1038/s42003-021-02219-6

Ma, D., Przybylski, D., Abruzzi, K.C., Schlichting, M., Li, Q., Long, X., Rosbash, M., 2021. A transcriptomic

taxonomy   of   Drosophila   circadian   neurons   around   the   clock.   Elife   10,   e63056.

https://doi.org/10.7554/eLife.63056

Marcheva, B., Ramsey, K.M., Buhr, E.D., Kobayashi, Y., Su, H., Ko, C.H., Ivanova, G., Omura, C., Mo, S., Vitaterna,

M.H., Lopez, J.P., Philipson, L.H., Bradfield, C.A., Crosby, S.D., Jebailey, L., Wang, X., Takahashi, J.S., Bass,

J., 2010. Disruption of the clock components CLOCK and BMAL1 leads to hypoinsulinaemia and diabetes.

Nature 466, 627–631. https://doi.org/10.1038/nature09253

Martinek, S., Inonog, S., Manoukian, A.S., Young, M.W., 2001. A Role for the Segment Polarity Gene shaggy/GSK-

3 in the Drosophila Circadian Clock. Cell 105, 769–779. https://doi.org/10.1016/S0092-8674(01)00383-X

McGuire, S.E., Mao, Z., Davis, R.L., 2004. Spatiotemporal Gene Expression Targeting with the TARGET and Gene-

Switch Systems in Drosophila. Science's STKE 2004, pl6–pl6. https://doi.org/10.1126/stke.2202004pl6

Metaxakis, A., Tain, L.S., Grönke, S., Hendrich, O., Hinze, Y., Birras, U., Partridge, L., 2014. Lowered Insulin

Signalling Ameliorates Age-Related Sleep Fragmentation in Drosophila. PLoS Biology 12, e1001824.

https://doi.org/10.1371/journal.pbio.1001824

Myers, M.P., Wager-Smith, K., Wesley, C.S., Young, M.W., Sehgal, A., 1995. Positional Cloning and Sequence

Analysis of the Drosophila Clock Gene, timeless. Science (1979) 270, 805–808.

https://doi.org/10.1126/science.270.5237.805

Naganos, S., Horiuchi, J., Saitoe, M., 2012. Mutations in the Drosophila insulin receptor substrate, CHICO, impair

olfactory associative learning. Neuroscience Research 73, 49–55. https://doi.org/10.1016/j.neures.2012.02.001

Naganos, S., Ueno, K., Horiuchi, J., Saitoe, M., 2016. Learning defects in Drosophila growth restricted chico mutants

are caused by attenuated adenylyl cyclase activity. Molecular Brain 9, 37. https://doi.org/10.1186/s13041-016-

0217-3

Nässel, D.R., Broeck, J. vanden, 2016. Insulin/IGF signaling in Drosophila and other insects: Factors that regulate

production, release and post-release action of the insulin-like peptides. Cellular and Molecular Life Sciences

73, 271–290. https://doi.org/10.1007/s00018-015-2063-3

Nässel, D.R., Liu, Y., Luo, J., 2015. Insulin/IGF signaling and its regulation in Drosophila. General and Comparative

Endocrinology 221, 255–266. https://doi.org/10.1016/j.ygcen.2014.11.021

Nayak, N., Mishra, M., 2021. High fat diet induced abnormalities in metabolism, growth, behavior, and circadian

clock in Drosophila melanogaster. Life Sciences 281, 119758. https://doi.org/10.1016/j.lfs.2021.119758

Noh, J., 2020. The effect of circadian and sleep disruptions on obesity risk. Journal of Obesity and Metabolic

Syndrome 27, 78–83. https://doi.org/10.7570/JOMES.2018.27.2.78

Papadopoulou, D., Bianchi, M.W., Bourouis, M., 2004. Functional Studies of Shaggy/Glycogen Synthase Kinase 3

Phosphorylation Sites in Drosophila melanogaster . Molecular and Cellular Biology 24, 4909–4919.

https://doi.org/10.1128/mcb.24.11.4909-4919.2004

Puig, O., Marr, M.T., Ruhf, M.L., Tjian, R., 2003. Control of cell number by Drosophila FOXO: Downstream and

feedback regulation of the insulin receptor pathway. Genes and Development 17, 2006–2020.

https://doi.org/10.1101/gad.1098703

Rajan, A., Perrimon, N., 2012. Drosophila cytokine unpaired 2 regulates physiological homeostasis by remotely

controlling insulin secretion. Cell 151, 123–137. https://doi.org/10.1016/j.cell.2012.08.019

Renn, S.C.P., Park, J.H., Rosbash, M., Hall, J.C., Taghert, P.H., 1999. A pdf Neuropeptide Gene Mutation and

Ablation of PDF Neurons Each Cause Severe Abnormalities of Behavioral Circadian Rhythms in Drosophila.

Cell 99, 791–802. https://doi.org/10.1016/S0092-8674(00)81676-1

Roenneberg, T., Merrow, M., 2005. Circadian clocks — the fall and rise of physiology. Nature Reviews Molecular

Cell Biology 6, 965–971. https://doi.org/10.1038/nrm1766

Rudic, R.D., McNamara, P., Curtis, A.M., Boston, R.C., Panda, S., Hogenesch, J.B., FitzGerald, G.A., 2004. BMAL1

and CLOCK, two essential components of the circadian clock, are involved in glucose homeostasis. PLoS

Biology 2. https://doi.org/10.1371/journal.pbio.0020377

Sano, H., Nakamura, A., Texada, M.J., Truman, J.W., Ishimoto, H., Kamikouchi, A., Nibu, Y., Kume, K., Ida, T.,

Kojima, M., 2015. The Nutrient-Responsive Hormone CCHamide-2 Controls Growth by Regulating Insulin-

like Peptides in the Brain of Drosophila melanogaster. PLoS Genetics 11, e1005209.

https://doi.org/10.1371/journal.pgen.1005209

Sato, M., Murakami, M., Node, K., Matsumura, R., Akashi, M., 2014. The Role of the Endocrine System in Feeding-

Induced Tissue-Specific Circadian Entrainment. Cell Reports 8, 393–401.

https://doi.org/10.1016/j.celrep.2014.06.015

Scheer, F.A.J.L., Hilton, M.F., Mantzoros, C.S., Shea, S.A., 2009. Adverse metabolic and cardiovascular

consequences of circadian misalignment. Proceedings of the National Academy of Sciences 106, 4453–4458.

https://doi.org/10.1073/pnas.0808180106

Schlichting, M., Díaz, M.M., Xin, J., Rosbash, M., 2019. Neuron-specific knockouts indicate the importance of

network communication to drosophila rhythmicity. Elife 8, e48301. https://doi.org/10.7554/eLife.48301

Seay, D.J., Thummel, C.S., 2011. The Circadian Clock, Light, and Cryptochrome Regulate Feeding and Metabolism

in Drosophila. Journal of Biological Rhythms 26, 497–506. https://doi.org/10.1177/0748730411420080

Sehgal, A., Rothenfluh-Hilfiker, A., Hunter-Ensor, M., Chen, Y., Myers, M.P., Young, M.W., 1995. Rhythmic

Expression of *timeless* : A Basis for Promoting Circadian Cycles in *period* Gene Autoregulation. Science (1979)

270, 808–810. https://doi.org/10.1126/science.270.5237.808

Song, J., Wu, L., Chen, Z., Kohanski, R.A., Pick, L., 2003. Axons guided by insulin receptor in Drosophila visual

system. Science 300, 502–5. https://doi.org/10.1126/science.1081203

Takahama, K., Tomita, J., Ueno, T., Yamazaki, M., Kume, S., Kume, K., 2012. Pan-neuronal knockdown of the c-

Jun N-terminal Kinase (JNK) results in a reduction in sleep and longevity in Drosophila. Biochemical and

Biophysical Research Communications 417, 807–811. https://doi.org/10.1016/j.bbrc.2011.12.040

Tanabe, K., Itoh, M., Tonoki, A., 2017. Age-Related Changes in Insulin-like Signaling Lead to Intermediate-Term

Memory Impairment in Drosophila. Cell Reports 18, 1598–1605. https://doi.org/10.1016/j.celrep.2017.01.053

Toda, H., Williams, J.A., Gulledge, M., Sehgal, A., 2019. A sleep-inducing gene, nemuri , links sleep and immune

function in Drosophila. Science (1979) 363, 509–515. https://doi.org/10.1126/science.aat1650

Tomita, J., Ban, G., Kume, K., 2017. Genes and neural circuits for sleep of the fruit fly. Neuroscience Research 118,

82–91. https://doi.org/10.1016/j.neures.2017.04.010

Tomita, J., Mitsuyoshi, M., Ueno, T., Aso, Y., Tanimoto, H., Nakai, Y., Aigaki, T., Kume, S., Kume, K., 2011. Pan-

neuronal knockdown of calcineurin reduces sleep in the fruit fly, Drosophila melanogaster. J Neurosci 31,

13137–13146. https://doi.org/10.1523/JNEUROSCI.5860-10.2011

Tomita, J., Ueno, T., Mitsuyoshi, M., Kume, S., Kume, K., 2015. The NMDA Receptor Promotes Sleep in the Fruit

Fly, Drosophila melanogaster. PLoS ONE 10, e0128101. https://doi.org/10.1371/journal.pone.0128101

Turek, F.W., Joshu, C., Kohsaka, A., Lin, E., Ivanova, G., McDearmon, E., Laposky, A., Losee-Olson, S., Easton, A.,

Jensen, D.R., Eckel, R.H., Takahashi, J.S., Bass, J., 2005. Obesity and metabolic syndrome in circadian Clock

mutant nice. Science (1979) 308, 1043–1045. https://doi.org/10.1126/science.1108750

Ueno, T., Kume, K., 2014. Functional characterization of dopamine transporter in vivo using Drosophila

melanogaster behavioral assays. Frontiers in Behavioral Neuroscience 8, 1–11.

https://doi.org/10.3389/fnbeh.2014.00303

Ueno, T., Tomita, J., Tanimoto, H., Endo, K., Ito, K., Kume, S., Kume, K., 2012. Identification of a dopamine pathway

that regulates sleep and arousal in Drosophila. Nature Neuroscience 15, 1516–1523.

421 https://doi.org/10.1038/nn.3238

422 Ugrankar, R., Berglund, E., Akdemir, F., Tran, C., Kim, M.S., Noh, J., Schneider, R., Ebert, B., Graff, J.M., 2015.

423 Drosophila glucome screening identifies Ck1alpha as a regulator of mammalian glucose metabolism. Nature

424 Communications 6, 7102. https://doi.org/10.1038/ncomms8102

425 Watanabe, K., Sakai, T., 2016. Knockout mutations of insulin-like peptide genes enhance sexual receptivity in

426 Drosophila virgin females. Genes and Genetic Systems 90, 237–241. https://doi.org/10.1266/ggs.15-00025

427 Yamaguchi, S.T., Tomita, J., Kume, K., 2021. Insulin signaling in clock neurons regulates sleep in Drosophila.

428 Biochemical and Biophysical Research Communications 591, 44–49.

429 https://doi.org/10.1016/j.bbrc.2021.12.100

430 Yamazaki, M., Tomita, J., Takahama, K., Ueno, T., Mitsuyoshi, M., Sakamoto, E., Kume, S., Kume, K., 2012. High

431 calorie diet augments age-associats sleep impairment in Drosophila. Biochemical and Biophysical Research

432 Communications 417, 812–816. https://doi.org/10.1016/j.bbrc.2011.12.041

433 Zheng, X., Sehgal, A., 2010. AKT and TOR Signaling Set the Pace of the Circadian Pacemaker. Current Biology 20,

434 1203–1208. https://doi.org/10.1016/j.cub.2010.05.027

Figure Legends

Figure 1. *chico¹* mutation caused a shorter circadian period length.

(A) Average activity from wild type (+, n = 14), backcrossed heterozygous (*chico¹* / + (BC5), n = 15), original

heterozygous (*chico¹* / + (F1), n = 31), and original homozygous (*chico¹*, n = 30) *chico* mutant flies. Each fly activity

was recorded for 10 days in constant dark (DD) conditions. Activity counts over 7 days in DD condition are shown.

Subjective day and night are depicted by the gray and black bars, respectively. (B) Circadian period lengths for +,

chico¹ / + (BC5), *chico¹* / + (F1), and *chico¹* flies. Circadian period lengths were calculated using locomotor activity

data for 10 days by FT. (C) Amplitudes of circadian rhythm for +, *chico¹* / + (BC5), *chico¹* / + (F1), and *chico¹*.

Amplitudes were calculated using locomotor activity data for 10 days by FT. *** p < 0.001; *t*-test.

Figure 2. Inhibition of insulin signaling in clock neurons resulted in shorter circadian period length.

(A, B) Circadian period lengths (A) and amplitudes of circadian rhythm (B), which were calculated using locomotor

activity data for 10 days by FT, for each genotype denoted at bottom of the panels. *w¹¹¹⁸*, the genetical background

of UAS-*InR* DN, was used as a control for UAS-*InR* DN. *y¹ v¹*; *P{CaryP}attP2* was used as ctrl for UAS-*chico* RNAi.

y¹ v¹; *P{CaryP}attP40* was used as ctrl for UAS-*InR* RNAi and UAS-*dp110* RNAi. Since each experiment was

conducted independently, the control group for UAS-*InR* RNAi and UAS-*dp110* RNAi showed different circadian

period lengths and amplitudes, although they are the same genotype. The numbers of flies of each bar were 31, 31,

29, 32, 16, 16, 16, and 11 from the left. (C, D) Circadian period lengths (C) and amplitudes of circadian rhythm (D),

which were calculated using locomotor activity data for 10 days by FT, for each genotype denoted at bottom of the

panels. The numbers of flies of each bar were 32, 27, 32, 32, 32, 32, 32, and 32 from the left. * $p < 0.05$, *** $p <$

0.001; *t*-test.

Figure 3. Insulin signaling in the lateral neurons regulates circadian rhythm.

(A, B) Circadian period lengths (A) and amplitudes of circadian rhythm (B), which were calculated using locomotor

activity data for 10 days by FT, for each genotype denoted at bottom of the panels. w^{1118}, the genetical background

of UAS-*InR* DN, was used as a control for UAS-*InR* DN. y^1 v^1; *P{CaryP}attP2* was used as ctrl for UAS-*chico* RNAi.

y^1 v^1; *P{CaryP}attP40* was used as ctrl for UAS-*InR* RNAi and UAS-*dp110* RNAi. Since each experiment was

conducted independently, the control group for UAS-*InR* RNAi and UAS-*dp110* RNAi showed different circadian

period lengths and amplitudes, although they are the same genotype. The numbers of flies of each bar were 32, 31,

31, 30, 32, 32, 32, and 32 from the left. (C, D) Circadian period lengths (C) and amplitudes of circadian rhythm (D),

which were calculated using locomotor activity data for 10 days by FT, for each genotype denoted at bottom of the

panels. The numbers of flies of each bar were 32, 32, 32, 31, 32, 32, 29, and 28 from the left. Error bars indicate

SEM. *** $p < 0.001$; *t*-test.

Figure 4. Insulin signaling in LNvs regulates circadian rhythm in a development-independent manner.

(A) Circadian period lengths and amplitudes of circadian rhythm, which were calculated using locomotor activity

data for 10 days by FT, for control (*Pdf*-GS x ctrl, white bar) and flies with *chico* knockdown by *Pdf*-GS (*Pdf*-GS x

UAS-*chico* RNAi, black bar) without RU486 (- RU486) or with RU486 (+ RU486). $y^1 v^1$; $P\{CaryP\}attP2$ was used

as ctrl for UAS-*chico* RNAi. The numbers of flies of each bar were 24, 28, 18, and 20 from the left. Error bars indica

te SEM. (B) Circadian period lengths and amplitudes of circadian rhythm, which were calculated using locomotor

activity data for 10 days by FT, for control (*Pdf*-GS x w^{1118}, white bar) and flies with *InR* DN expression by *Pdf*-GS

(*Pdf*-GS x UAS-*InR* DN, black bar) without RU486 (- RU486) or with RU486 (+ RU486). The numbers of flies of

each bar were 26, 25, 27, and 26 from the left. Error bars indicate SEM. w^{1118}, the genetical background of UAS-*InR*

DN, was used as a control for UAS-*InR* DN. n.s. $p \geq 0.05$, * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$; Tukey Kramer

method.

Figure 5. *dilp5* relates to the regulation of circadian rhythm by insulin signaling.

(A) Circadian period lengths and amplitudes of circadian rhythm, which were calculated using locomotor activity

data for 10 days by FT, for each genotype denoted at bottom of the panels: TKO x w^{1118} (n = 15), TKO x *dilp2* KO

(n = 16), TKO x *dilp3* KO (n = 16), TKO x *dilp5* KO (n = 13). (B) Relative expression of *4E-BP* for each genotype

denoted at bottom of each panel. (C) Circadian period lengths and amplitudes of circadian rhythm, which were

calculated using locomotor activity data for 10 days by FT, for *elav*-Gal4; UAS-*Dicer-2* x ctrl (n = 14) and *elav*-

Gal4; UAS-*Dicer-2* x UAS-*dilp5* RNAi (n = 11). w^{1118}(VDRC stock number: 60000) was used as ctrl for UAS-*dilp5*

RNAi. Error bars indicate SEM. * $p < 0.05$, ** $p < 0.01$; vs. TKO x w^{1118}; Tukey Kramer method (A, B). ** $p < 0.01$;

t-test (C)

Figure 1

Figure 3