Ultrasensitivity of Microtubule severing rate on the concentration of free Tubulin

Author list:

Chloe E. Shiff[1], Jane Kondev[1], Lishibanya Mohapatra[2,*]

[1]Department of Physics, Brandeis University, Waltham, MA 02454, USA

[2]School of Physics and Astronomy, College of Science, Rochester Institute of Technology, Rochester, NY 14623, USA

* = Corresponding Author, Lead contact.

Corresponding author email: lxmsps@rit.edu

Abstract

Cytoskeletal structures aid in cell polarization, motility, and intracellular transport.
Their functions are predicated on the rapid turnover of cytoskeleton proteins, which
is achieved by the coordinated effort of multiple regulatory proteins whose
dynamics are not well understood. In-vitro experiments have shown that free
tubulin can repair nanoscale damages of microtubules created by severing
proteins. Based on this observation, we propose a model for microtubule severing
as a competition between the processes of damage spreading and tubulin-induced
repair. Using theory and simulations, we demonstrate that this model is in
quantitative agreement with in vitro experiments. We predict the existence of a
critical tubulin concentration above which severing becomes rare but fast, and
hypersensitive to the concentration of free tubulin. Further we show that this
hypersensitivity leads to a dramatic increase in the dynamic range of steady-state
microtubule lengths, when lengths are controlled by severing. Our work
demonstrates how synergy between tubulin and severing proteins can lead to
novel dynamical properties of microtubules.

17
18
19 **Main Text**
20
21 **Introduction**
22
Cells contain a cytoskeleton that aids in various cellular processes such as
maintaining cell polarization, motility, and transport. The cytoskeleton is highly
dynamic – it is composed of various structures that are constantly built and
dismantled to assist in various functions. For example, the single-cell alga
Chlamydomonas uses microtubule-based flagella for motility (Marshall et al., 2005;

Song and Dentler, 2001). Microtubule-based fibers in the mitotic spindle shorten
for chromosome separation during mitosis and meiosis (Bennabi et al., 2016;
Dumont and Mitchison, 2009; Hyman and Karsenti, 1996).

Such dynamic changes in cytoskeletal structures require a rapid turnover which
involves an exchange of their molecular building blocks, with the same molecules
that are freely diffusing in the cytoplasm. Remarkably, the time scales associated
with the dynamics of proteins filaments in vivo are quite different from those
measured in vitro. While actin networks within cells are disassembled on the order
of seconds, the process takes a few minutes in vitro (Kuhn and Pollard, 2005;
Miyoshi and Watanabe, 2013; Pollard, 1986; Watanabe and Mitchison, 2002).
Similarly, the polymerization rate of tubulin in vivo is about five to tenfold higher
than in vitro at a similar concentration of free tubulin (Desai and Mitchison, 1997).
Further, microtubules are known to switch between phases of slow growth and
rapid shrinking by a process known as dynamic instability, which has been studied
extensively in vitro (Desai and Mitchison, 1997). The rates of switching from
shrinking to growth and vice versa (called rates of rescue and catastrophe,
respectively) are sensitive to, and are believed to be modulated by several
microtubule-associated proteins in vivo (Desai et al., 1999; Gardner et al., 2013;
Geisterfer et al., 2020; Goshima et al., 2005; Tournebize et al., 2000). Turnover
rates can also be different at different stages of the cell cycle – for example,
microtubule turnover rates measured in Xenopus extracts range from 3 min to 20s
during interphase and mitosis, respectively(Hyman and Karsenti, 1996). While
much is known about the biochemical processes that facilitate the turnover of these
structures in isolation, it is still not clear how different cytoskeleton-associated
proteins work together to promote filament turnover in cells.
53
Recently, several studies have addressed this question by combining multiple
factors with cytoskeletal filaments in vitro and have reported novel synergistic
results that are not seen when studying each component individually. Shekhar et
al. report that the cofilin-induced depolymerization rate of actin filaments increases
200-fold in the presence of cyclase associated protein/Srv-2(Shekhar et al., 2019).
Similar studies have been done using microtubules with a recent study finding that
collective effects of microtubule-associated proteins like MCAK and XMAP215 can
lead to microtubule treadmilling(Arpağ et al., 2020). Inspired by yet another set of
recent experiments, we consider the interplay between the free pool of tubulin
dimers and microtubule severing proteins, which fragment protein filaments(Vemu
et al., 2018).
65
Free tubulin is known to affect microtubule dynamics by modulating rates of
nucleation and polymerization (Geisterfer et al., 2020; King and Petry, 2020; Ohi
et al., 2021; Woodruff et al., 2017). Notably, using FRET, it was shown that a local
release of tubulin promoted microtubule extensions into lamellipodia,
demonstrating a feedback between filament bound and free tubulin pools(Van
Geel et al., 2020). However, there are also reports of free tubulin affecting
microtubule disassembly via its interactions with proteins such as kinesin-13
(Desai et al., 1999; Walczak et al., 2013), kinesin-8 (Arellano-Santoyo et al., 2017;

Varga et al., 2009) and severing proteins (Kuo et al., 2019b; Vemu et al., 2018).
Here we theoretically describe the effect of free tubulin on microtubule severing
via a recently discovered process of repair of damages to the microtubule lattice
by the severing proteins(Vemu et al., 2018).
78
Severing proteins can play a key role in the reorganization of the cytoskeleton by
breaking down existing structures and making more monomers available for use
in new structures (Kuo and Howard, 2021). Recent studies have found that
severing proteins can seed new filament growth by releasing filaments from
nucleation sites and allowing their transport within the cell (Sharp and Ross, 2012).
Katanin and spastin are well characterized examples of severing proteins for
microtubule-based structures, while cofilin and Srv-2 sever actin-based structures
(Balcer et al., 2003, p. 2; Chaudhry et al., 2013; Díaz-Valencia et al., 2011;
Johnston et al., 2015, p. 2; Kuo et al., 2019b). Katanin is thought to play two
functions in mitosis and meiosis – to amplify the number of microtubules in the
spindle and to uncap microtubule plus ends from the kinetochore and enable
depolymerizing kinesins to target microtubules (Sharp and Ross, 2012). Inhibition
of katanin results in abnormally long mitotic spindle length in *Xenopus* extracts
(Loughlin et al., 2011). Severing proteins play an important role in neurons as well
(Karabay et al., 2004). It has been proposed that katanin stimulates axon growth
by releasing microtubules from centrosomes and severing them into shorter
segments that can be transported along longer microtubules down the axon to
seed new microtubule growth (Ahmad et al., 1999). Mutations in *Drosophila*
spastin reduce dendritic arborization (branching) (Jinushi-Nakao et al., 2007),
suggesting that spastin seeds microtubule growth within newly forming regions of
the dendrite. Additionally, by using in vitro systems and theory, many studies have
also analyzed the effect of severing proteins on the length of cytoskeletal filaments
(Elam et al., 2013; Kuo et al., 2019b, 2019a; Mohapatra et al., 2016; Roland et al.,
2008; Tindemans and Mulder, 2010).
103
Thus far, most studies have focused on the effect of severing proteins in isolation,
but recently experiments have examined the interplay of these proteins in
presence of free tubulin. Electron microscopy images of microtubules have
revealed the mechanisms by which severing proteins induce a break in the
microtubule. One study reported that once severing proteins land on a microtubule,
they create nanoscale damages(Vemu et al., 2018). This study found that not all
of the damage sites proceeded to a severing event, as free tubulin could repair the
damage in some cases. Furthermore, a few in-vitro studies have reported that
severing proteins enhance the rate of rescue in microtubules undergoing dynamic
instability (Kuo et al., 2019b, 2019a; Kuo and Howard, 2021; Vemu et al., 2018).
At this point, it is unknown whether severing proteins have the same effect in vivo,
where they may interact with other microtubule-associated proteins.
116
Inspired by these observations (Vemu et al., 2018), we propose a model for
microtubule severing that incorporates the newly discovered repair process by free
tubulin. Using theory and simulations, we show that this new model quantitatively

agrees with in vitro experiments and makes predictions for new in vitro
experiments. Significantly, we predict the existence of a critical tubulin
concentration above which severing becomes rare but fast. This model also has
several implications for the dynamics of microtubules in vivo – we show that length
control via severing is far more sensitive to changes in tubulin concentrations,
resulting in a dramatically expanded dynamic range of filament lengths. We also
report the probability distribution of lengths and the mean length and its coefficient
of variation as a function of the model parameters, which can be used to test the
model. In summary, our work describes how the concerted action of tubulin and
severing proteins produces novel dynamical properties of microtubules that are not
seen when these proteins are studied independently.
131
132 **Results**
133
1. Model of microtubule severing with repair

Inspired by recent experiments(Vemu et al., 2018), we consider that a microtubule
severing event begins when a severing protein lands on a filament, binds to the
side, and creates a damage spot (Fig. 1A). The fate of the damage created by a
severing protein is determined by a competition between the removal and addition
of tubulin at the damage site, eventually leading to severing or repair. We assume
that monomers are added to the damage site at a rate proportional to the
concentration of free tubulin $k_T[T]$ (s^{-1}) and are removed from the damage site at
a rate k_r (s^{-1}), with each step either decreasing or increasing the size of damage.
If the size of the damage reaches a size N_D removed tubulins, the filament is
severed. On the other hand, if the size reaches zero, then the damage is repaired.
In this way, the size of the damage x measured in monomers (i.e., tubulin dimers)
is modeled as a one-dimensional random walk with absorbing boundaries at $x = 0$
and $x = N_D$. (Fig. 1B)

Figure 1. Microtubule severing with repair. (A) Severing proteins (in blue) create
small damages on microtubule lattice which either get repaired by free tubulin or
lose enough monomers to sever the microtubule. (B) One-dimensional random
walk model for studying the competition between removal of tubulin monomers at
a rate k_r and the addition of monomers at a rate proportional to the free tubulin
concentration, $k_T[T]$.(C) We use published data (Vemu et al., 2018) for the
severing times as a function of tubulin concentration to estimate parameters of the
severing model, which are listed in Table 1. (D) The same parameters are used to
compute the standard deviation (inset) and probability distribution of severing time
using stochastic simulations. Each distribution was computed from 400,000
trajectories. Error bars on experimental data obtained by bootstrapping method
(Section 4 in Materials and Methods). The error bars on the standard deviation
from simulations show SEM calculated from 750 values, with each value computed
from 20,000 trajectories. Comparison with published experimental data (Fig. A2 ,
from (Vemu et al., 2018), in blue) is shown.

In this model of severing with repair, a damage site can either be repaired or
progress to a severing event. Using our one-dimensional random walk model, we
find an analytical expression for the probability of severing (described in section 2
and SI), and the mean time it would take to sever a filament as a function of model
parameters (k_T, k_r, and N_D). Assuming an initial damage size of one monomer,
and defining the ratio of the tubulin addition and removal rate as a dimensionless

tubulin concentration $\tilde{T} = \frac{k_T[T]}{k_r}$, we predict that the time to sever a filament has the following form ($\tilde{T} \neq 1$)

$$T_{sev}(k_T,\ k_r, N_D) = \left(\frac{\frac{1}{k_r}}{\tilde{T}-1}\right)\left(N_D \frac{(\tilde{T}^{N_D+1})}{(\tilde{T}^{N_D}-1)} - \frac{\tilde{T}+1}{\tilde{T}-1}\right),\ (1)$$

In the case, $\tilde{T} = 1$, $T_{sev}(k_T,\ k_r, N_D) = \frac{1}{6k_r}(N_D^2 - 1)$.

Considering $N_D \gg 1$, equation 1 has two limits. When the rate of tubulin removal is more than tubulin addition, i.e., $\tilde{T} \ll 1$, $T_{sev}(k_T,\ k_r, N_D) \approx \frac{N_D}{k_r}(1 + \tilde{T})$, the severing time is linear in tubulin concentration. When $\tilde{T} \gg 1$, $T_{sev}(k_T,\ k_r, N_D) \approx \left(\frac{N_D}{k_r}\frac{1}{\tilde{T}}\right)$, the severing time decreases as tubulin concentration increases. These limits indicate that there is a critical concentration of tubulin, $[T]_c = \frac{k_r}{k_T}$, which delineates different functional dependencies of the severing time on the concentration of free tubulin. We also note that the time scale of the severing time is given by a parameter $\frac{N_D}{k_r}$, which is the mean time for completing N_D tubulin removal steps, each lasting $\frac{1}{k_r}$, on average. Equation 1 was checked against stochastic simulation of severing with repair (described in Section 5(B) of Materials and Methods).

Next, we use published data to estimate a set of parameters for the model of severing with repair. We consider the measured probability to sever and severing time as a function of tubulin concentration published in Fig. 3D and 3H in (Vemu et al., 2018), and fit our model parameters (k_T, k_r, and N_D) using method of least squares, described in Section 4 of Materials and Methods; see Figure 1C. The sen tivity of our results on the chosen parameters is shown in Fig. A1. Using the estimated parameters, listed in Table 1, we conduct stochastic simulations of the model (see section 5(A) in Materials and Methods and Figure A1 in the Appendix) to compute the distribution of severing times as a function of tubulin concentration and compare it to published data. Remarkably, just as in the data (Vemu et al., 2018), we find that the distribution of severing times is asymmetric around the maximum and skewed towards large times; see Figure 1D. A quantitative comparison between the measured standard deviation of the severing time and the predictions from our model is shown in the inset of Figure 1D.

The published experiment (Vemu et al., 2018) suggests that severing time is monotonic in the tubulin concentration up to $5\,\mu M$. Interestingly, our model (equation 1) predicts a non-monotonic relationship between severing time and the concentration of tubulin. We find, as reported in the study, that severing time increases as the concentration of tubulin is increased but only until a certain critical concentration of tubulin. Above this concentration, as shown in equation 1, we predict that the severing time should decrease (See Fig. 2A). As we describe next, at high concentrations of tubulin, we expect that the severing events are rare (but rapid) and almost all damages are likely to quickly be repaired by the free tubulin.

208 Note that the non-monotonicity in the severing time is a prediction of our model
209 that can be used to experimentally test the proposed mechanism of severing with
210 repair.

2. Theoretical predictions of severing with repair

212 Our model makes several other predictions. The probability of severing is the
213 probability that the one-dimensional random walker reaches the $x = N_D$ boundary
214 before the $x = 0$ boundary:

$$P_{sev}([T]) = \frac{\tilde{T} - 1}{\tilde{T}^{N_D} - 1}, \qquad (2)$$

where we have chosen the initial damage size of 1, while N_D is the damage size
that produces a severing event, and we expect $N_D \gg 1$ (see Table 1). Again, this

expression has two limits (shown in dotted black lines in Fig. 2): when $\tilde{T} < 1$, (
$[T] < [T]_c$), $P_{sev}([T]) \approx 1 - \tilde{T}$, whereas when $[T] > [T]_c$ ($\tilde{T} > 1$), $P_{sev}([T]) \approx$
$\left(\frac{1}{\tilde{T}}\right)^{N_D - 1}$. Thus, above the critical concentration, the probability of severing is very
small, and damage events that lead to a severing event, are very unlikely. We
confirmed our analytical expression for the mean severing time and the probability
of severing as a function of tubulin concentration by stochastic simulations (See
Fig. A2).

Figure 2. Theoretical prediction for model for microtubule severing with
repair (A)The model predicts a non-monotonic relationship between severing time
and free tubulin concentration and the existence of a critical concentration of
tubulin, $[T]_c$, beyond which severing time decreases. (B) Probability of severing
decreases as the tubulin concentration is increased (Inset: plot on a log scale).
The dashed lines in both plots show the asymptotic behavior at large $[T] \gg [T]_c$
and small $[T] \ll [T]_c$ free tubulin concentrations. Parameters used to plot these
graphs are listed in Table 1.
233

3. Length control in the presence of repair

The probability that a severing protein which is diffusing in solution lands on a filament is proportional to its length (Berg, 1993). Therefore, the process of severing is a length-dependent disassembly mechanism wherein longer filaments are more likely to be severed. This results in negative feedback of filament length on a growing filament and leads to control of length of cytoskeletal filaments. A number of theoretical and experimental studies (Kuo et al., 2019b, 2019a; Loughlin et al., 2011; Mohapatra et al., 2016; Roland et al., 2008; Tindemans and Mulder, 2010) have explored this aspect of severing. Given the recent observation of tubulin-induced repair (Vemu et al., 2018), we next study the implications of this repair process on microtubule length control by severing. We consider a situation where a microtubule filament undergoes growth with a rate proportional to the free tubulin concentration and severing, where severing competes with tubulin-induced repair, as described above. We make a simplifying assumption, namely we ignore the time it takes for the severing event to complete once the initial damage is created. We also assume that microtubules are stabilized and are not undergoing dynamic instability. Later in the Discussion and in the Appendix, we examine these assumptions and show that they do not qualitatively change our results about the mean steady-state microtubule lengths (see Sections A1 and A2 in Appendix, and Figs. A3 and A4), while allowing us to obtain analytic results for the steady state microtubule length. We compare our results of the severing with repair model to the severing model without repair, as considered previously in (Kuo et al., 2019a; Mohapatra et al., 2016). Results of this comparison are shown in Fig. 3.

Figure 3. Effect of repair on filament length control via severing. (A) Schematic of the two models considered. Severing without repair: A filament grows with a rate $k[T]$ and is severed with a rate $k_d[S]$. Severing with repair: Filaments grow with a rate $k[T]$ and either get severed with a rate $k_d[S]P_{sev}$ (per monomer

of the filament) or repaired with a rate $k_d[S](1 - P_{sev})$ where P_{sev} (probability of severing) is calculated using equation 2. If severing occurs, the end portion of the filament (right part) is removed. (B) Dynamics of filament length (average of 3 length trajectories shown for clarity) for the two models obtained using stochastic simulations. Dynamic range of the steady state filament lengths as [T] is tuned is larger with repair than without. (C) Mean steady state length as a function of tubulin concentration (Log-Log plot inset). Dots are from simulations; lines are from analytic calculations for the model with repair (orange) and without (purple). The dashed lines represent the show the asymptotic behavior at large $[T] \gg [T]_c$ and small $[T] \ll [T]_c$ (D) Distribution of steady state lengths and coefficient of variation (inset) as a function of tubulin concentration, each computed from 10^6 trajectories. Parameters in the steady-state simulations are listed in Table 1.

Using the computational scheme described in the Methods (See section 5(B)), we conduct stochastic simulations of the filament length as a function of time and analyze the process with and without the repair of severing protein-induced damage (Fig. 3A). In our simulations, individual filaments grow at a rate proportional to the tubulin concentration, k [T], where k is the second order rate constant for the tubulin binding to the end of microtubules and has the units of $(\mu M.s)^{-1}$. Severing proteins land on the side of the filament with a rate proportional to the concentration of severing proteins $[S]$ and to the length of the filament, l. For the simulations without the repair process, severing at any position on the filament is $k_d[S]$. In the simulation with the repair process, we take $k_d[S]l$ to represent the rate at which damage sites are created along the length of the microtubule. Assuming this is the rate limiting process in severing, the rate of severing can be written as $k_d[S]lP_{sev}$ where P_{sev} is the probability to sever (equation 2).

For the two cases (severing without and severing with repair), we generate trajectories of length over time. In both cases, we notice that, after a steep growth phase, the filament length fluctuates around a steady-state length, but the values of range of lengths obtained for each case are different. We generate trajectories at three different tubulin concentrations and find that the mean lengths achieved by the filaments (defined as the dynamic range of filament lengths) are larger when the process of repair is included (Fig. 3B).

We also analyzed this process using theory (described in Section 6 in Materials and Methods) and calculated the mean filament length at steady state as $L_s^{without\ repair} = \sqrt{\dfrac{\pi k[T]}{2\,k_d[S]}}$, and $L_s^{with\ repair} = \sqrt{\dfrac{\pi k[T]}{2\,k_d[S]P_{sev}([T])}}$. These expressions were also confirmed using simulations. As shown in Fig. 3C, while the steady state length $L_s \sim \sqrt{[T]}$ without repair, we observe two regimes with tubulin-induced repair present– when $[T] < [T]_c$, $L_s \sim \sqrt{[T]}$ and when $[T] > [T]_c$, $L_s \sim [T]^{N_D/2}$, suggesting a very high sensitivity to tubulin concentration, above the critical concentration (See Section 6 in Materials and Methods). Note that our results for severing without repair ($L_s \sim \sqrt{[T]/[S]}$) are similar to the predictions for the "no catastrophe model" in a recently published paper which studied the effect of the severing protein spastin on microtubule length (Kuo et al., 2019a). In contrast, we find that including the possibility of repair of severing events in our model predicts a new regime of microtubule growth that was not described previously.

We derive analytically (described in Section 6 in Materials and Methods), the probability distribution of the steady state lengths and compare our simulations with the theory in Fig. 3D. We also calculate the coefficient of variation (CV = standard deviation/mean, commonly referred to as the noise) of the steady-state lengths, with and without repair. Significantly, we calculate that CV doesn't change with [T] (See Fig. 3D, inset and Section 6 in Materials and Methods) even though the mean lengths are very different for the two models for [T] above $[T]_c$. This

321 serves as a theoretical prediction for severing controlling length and can be used
322 to discern this mechanism from other control mechanisms(Mohapatra et al., 2016).
323

324 **Discussion**
325

Dynamics of protein filaments in cells (microtubules and actin filaments) are very
different from what is observed in vitro (Zwetsloot et al., 2018). For example, actin
networks in cells disassemble on the order of seconds whereas in vitro actin
filaments disassemble over minutes (Miyoshi and Watanabe, 2013; Watanabe and
Mitchison, 2002). These observations suggest that the dynamics of filament
assembly are affected by proteins that associate with filaments in vivo and exert
control over assembly and/or disassembly (Kuo and Howard, 2021; Sharp and
Ross, 2012; Zanic et al., 2013). Inspired by recent experiments on microtubules
(Vemu et al., 2018), we consider one such control system and demonstrate that it
leads to dynamics that are distinct from those described for microtubules in
isolation.
337
338

PARAMETERS	VALUE USED IN PAPER	REFERENCES
CONCENTRATION OF TUBULIN, [T]	0-5 µM	(Vemu et al., 2018)
RATE OF REPLENISHMENT, K_T	0.11 µM^{-1}s^{-1}	Estimated
RATE OF REMOVAL, K_R	0.54 s^{-1}	Estimated
MINIMUM SIZE OF DAMAGE	1 monomer	Assumed
MAXIMUM SIZE OF DAMAGE N_D	25	Estimated
RATE OF GROWTH, K	3.4 µM^{-1}s^{-1}	(Mickolajczyk et al., 2019)
CONCENTRATION OF SEVERING PROTEINS, [S]	50 nM	(Vemu et al., 2018)
SEVERING PROTEIN LANDING RATE, K_D	0.002 µM^{-1}s^{-1}nm^{-1}	(Kuo et al., 2019a)

340 **Table 1.** Parameters used in the models for the model of severing and length
341 control with tubulin-induced repair. Estimated values were attained by fitting the
342 theoretical expression for the mean severing time to experimental data in ref
343 (Vemu et al., 2018); see Figure 1C.
344
345

We model the severing of microtubules as a competition between the process of removal of monomers from the damage site created by a severing protein and the repair of the damage site by binding of free monomers. We use a one-dimensional random walk model to describe this competition and compute quantities that can be measured in experiments, such as the probability of severing and the time taken to successfully sever a microtubule as a function of the free tubulin concentration. We quantitatively compare our theory to a number of experimental observations(Vemu et al., 2018) pertaining to the effect of severing proteins on microtubules. Notably, we predict a distribution of severing times (Fig. 1D) skewed towards large severing times, which was observed in experimental results(Vemu et al., 2018). Additionally, using our model, we predict that the time to sever has a non-monotonic dependence on tubulin concentration and reaches a maximum at a critical tubulin concentration. Next, we studied the implication of this repair process on the length of a growing filament and found that it leads to a very sensitive dependence of steady state length as a function of free tubulin concentration.

Signatures of severing as a length-control mechanism

Often in cells, several mechanisms can be at play to control the size of a sub-cellular structure, making it challenging to distinguish between them. Key results of this paper can serve as quantitative signatures of the severing mechanism.

A key feature of length-control mechanisms is that they lead to a peaked distribution of steady-state filament lengths. Several theoretical and experimental studies have demonstrated that severing can control the length of filaments (Kuo et al., 2019a; Mohapatra et al., 2016; Roland et al., 2008). Longer microtubules have more binding sites for severing proteins, and therefore have a greater probability of being severed than shorter ones. This leads to negative feedback on the length of a growing microtubule which eventually reaches a steady-state length once the disassembly rate balances the rate of assembly. The resulting distribution of steady-state lengths, in addition to being peaked, has a skewness (see Fig. 3D) which makes it distinct from other control mechanisms studied thus far (Mohapatra et al., 2016). This skewness, predicted in theoretical studies (Kuo et al., 2019a; Mohapatra et al., 2016; Tindemans and Mulder, 2010), was observed in recent experimental studies (Kuo et al., 2019a; Vemu et al., 2018). Additionally, we also find that the coefficient of variation (CV) of steady state microtubule lengths is independent of the concentration of tubulin and has a constant value equal to 0.523 (See Fig. 3D inset and Section 6 in Materials and Methods). This result relies on an assumption that the severing process is uniform along the filament, as was recently reported for spastin (Kuo et al., 2019a). Factors like filament aging, as has been documented in the context of actin filaments, arising due to the ATP hydrolysis (Roland et al., 2008), could invalidate this assumption.

Experimental predictions

1. Probability of severing as a function of tubulin concentration

We have shown that a one-dimensional random walk model quantitatively accounts for recent experiments on microtubule severing. Using this model, we find the probability of severing decreases as the concentration of tubulin increases (See Fig. 2B). This prediction of the model can be used to further test it. Many studies have used a "severing assay" to study the effect of severing proteins on microtubules (Belonogov et al., 2019; Vemu et al., 2018; Ziółkowska and Roll-Mecak, 2013). In these experiments, severing proteins are flown into a reaction chamber with pre-formed microtubules and quantities like the number of successful severing events per unit length of the microtubule (the severing rate) can thus be measured. Note that the maximal rate of severing is achieved when there is no free tubulin to repair the initial damage caused by a severing protein. Therefore, the ratio of the severing rate in the presence of free tubulin to the severing rate with no free tubulin is the probability of severing. At zero concentration of free tubulin, we expect that all severing initiation events lead to successful severing, but that the proportion would decrease as free tubulin can lead to repair, which is captured by equation 2, and Fig. 2B.

2. Non-monotonic relationship of severing time and free tubulin concentration

Importantly, our model predicts a critical tubulin concentration $[T]_c = k_r/k_T$, where k_r is the first order rate of removal and k_T is the second order rate of addition of tubulin from the damage site. Above this critical concentration, the probability of severing drops rapidly, causing the severing time to peak at $[T]_c$; see Fig. 2A. By using our estimated parameters, we predict that the mean severing time should increase for $0 \leq [T] \leq 5\mu M$, a prediction which agrees with results from experiments described in a recent study(Vemu et al., 2018). However, at tubulin concentrations above those which were experimentally investigated in (Vemu et al., 2018), $[T] > 5\,\mu M$, we find that severing time decreases. This prediction can be used to test the proposed model using the set-up described in (Vemu et al., 2018).

Figure 4: Testable predictions of the severing with repair model of length control
(A) Mean filament length and (B) coefficient of variation at steady state as function of

Tubulin concentration at different severing protein concentrations. Dynamic range is larger at lower concentrations of severing proteins, and the coefficient of variation is independent of tubulin and severing protein concentrations. Parameters used in the simulation are listed in Table 1.

3. Ultra-sensitivity of microtubule length control.

Severing of filaments, which are also growing by the addition of monomers, leads to length control. We observe that the tubulin repair process leads to an ultra-sensitivity in the lengths of microtubules as a function of tubulin concentration (Fig. 3C).

In Fig. 4, we show results of calculations of the change in steady-state length (L_S) and the coefficient of variation as a function of two experimentally tunable knobs – tubulin and severing protein concentrations. We find a sharp increase in L_S with $[T]$ above the critical concentration $[T]_c = \frac{k_r}{k_T}$, while the coefficient of variation remains the same across all concentrations of severing proteins. These results can be tested by creating an in vitro system (Díaz-Valencia et al., 2011; Kuo et al., 2019b; Vemu et al., 2018) in which growing microtubules are exposed to severing proteins. We also compute the probability distribution of steady-state lengths as a function of tubulin concentrations which can be used as a more stringent test of the model (see Fig. 3D).

Examining the assumptions of the model:

In order to make the model of severing with repair analytically tractable, we made a few simplifying assumptions. Below we examine how robust our conclusions are with respect to these assumptions:

Effect of finite severing time on microtubule dynamics
In our calculations of microtubule length-control (Figs. 3 and 4), we ignore the time it takes for the severing event to complete once the initial damage is created. This is equivalent to assuming that the time between subsequent damage creation events is much bigger than time to sever. This is a reasonable assumption in the limits of low and high tubulin concentrations, but not a valid one for intermediate tubulin concentrations (Fig A3, section A1 in Appendix).

Using the rates in Table 1, for small tubulin concentrations $[T] \approx 0.01 \mu M$, typical microtubules lengths are small, about 50 nm (using equation $L_S \sim \sqrt{\frac{\pi k[T]}{2k_d[S]}}$ at small $[T]$ approximation) and $P_{sev}[T]$ is close to 1, which means practically all damage events lead to severing. Time taken for a severing protein to bind to a microtubule and induce damage is $\approx \frac{1}{k_d[S]L}$ = 200s for a microtubule of length around 50 nm. In comparison, the severing time after initial damage $\approx \frac{N_D}{k_r}$ = 50 secs. Thus, at low

tubulin concentrations, severing time is shorter than the severing protein binding
time, making it unlikely that multiple damage sites are created. At high tubulin
concentrations$[T] \gtrsim 5 \ \mu M$, the microtubules are $\gtrsim 1$ micron in length, and the time
for a severing protein to bind is around 10 secs. At these tubulin concentrations,
the probability of severing is small ($\lesssim 0.025$) and at least 40 binding events are
required for one severing event. Waiting time for 40 binding events is about 400
seconds, which is greater than the severing time (200s or less) (Fig. A3 D, inset).
Thus, our simplified model that ignores the possibility of multiple damage sites, in
addition to being analytically tractable, provides a good approximation to study the
effect of severing on microtubule length control in the limits of low and high tubulin
concentrations.
478
At intermediate tubulin concentrations, however, time between severing events is
comparable to severing time (Fig A3 D, inset), making it possible that several
damage events are happening at once. This can lead to correlations between
severing events which our calculations ignore. In order to study the implications of
this scenario, we performed stochastic simulations wherein we tracked the
locations of multiple damage sites on a microtubule lattice. Each damage site has
a corresponding severing time, drawn from a distribution of severing times
obtained from the stochastic simulation of the damage spreading simulations
(described in Section 5(A) in Materials and Methods). We observe that results of
large dynamic range in steady-state length from this simulation are similar to what
was seen with the simple model (Figs. 3C, A4). Additionally, the noise as a function
of free tubulin concentration from the full simulation was identical to the one
obtained from the simple model at low and high tubulin concentration. However, at
intermediate values of tubulin concentrations, we notice a peak in the CV vs tubulin
concentration plot (Fig. A3 D), which was in contrast to the constant value seen
with the simple model (Fig. 3D) but to be expected due to correlations in severing
events.
496
497 *Dynamic instability*
498
Microtubules are known to switch between phases of slow growth and rapid
shrinking by a process known as dynamic instability. The rates of switching from
shrinking to growth and vice versa are called rates of rescue and catastrophe,
respectively. In our simulations leading to Fig. 3 and 4, we assumed that the
microtubule are stabilized, and are not undergoing dynamic instability. In order to
check the robustness of our results, we included dynamic instability in our
simulations (see section A2 for simulation protocol). We found that dynamic
instability did not qualitatively change the observation of large dynamic range of
filament lengths. Coincidentally, a previous theoretical study analyzed the effect of
the severing protein spastin on dynamic microtubules, and reported that their
model predictions were similar to a model which ignored the dynamic instability
("no catastrophe" model) (Kuo et al., 2019a).
511

With stabilized microtubules, we had calculated that the coefficient of variation (CV) for the steady state length fluctuations for the models of severing without repair and severing with repair was constant with respect to tubulin concentration. In contrast, when dynamic instability was included, we observed a non-monotonic relationship in CV for both models (see Fig. A4 in section A2). In our simulations, microtubule filaments grow at a rate proportional to tubulin concentration. Since severing depends on microtubule length, which is typically small at low concentrations of tubulin, dynamic instability dominates over severing in both models. In this regime, the length distributions are exponential and resulting in CV being close to 1(Dogterom and Leibler, 1993; Kuo et al., 2019b). However, we observe different behaviors for both models (severing without repair and severing with repair) at higher tubulin concentration. In this regime, filaments grow longer and severing dominates over dynamic instability, as the number of subunits lost in a catastrophe event is small compared to the number of subunits lost due to severing. So, as expected, in the model of severing without repair, CV converges towards 0.523 at large concentration of tubulin. In severing with repair, we observe that CV shows a non-monotonic behavior with tubulin concentration, where it first decreases and then increases, and the dip in CV depends on the rescue rates chosen. Additionally, in-vitro studies have reported an increase in rescue rates in presence of severing proteins. Several hypotheses have been proposed to explain this, such as, the incorporation of new tubulin creating a "GTP-island", which promotes rescue (Vemu et al., 2018) and the accumulation of severing proteins on the tips of shrinking microtubules, where they slow shrinkage and promote rescue (Kuo et al., 2019b). At this point, it is unknown whether severing proteins have the same effect in vivo, where they may interact with other microtubule associated proteins.

In summary, our calculations show that tubulin and severing proteins can cause qualitatively new dynamics of microtubules. Using simulations and theory, our study describes the role of a free pool of building blocks on the process of repair of damage created by severing proteins and its implications for the control of length of microtubules. Here we show the existence of a critical tubulin concentration above which severing becomes strongly dependent on $[T]$, leading to a rare but fast severing. We find that microtubule length control via severing with tubulin-induced repair is ultra-sensitive to changes in the concentrations of tubulin leading to a dramatically increased dynamic range of filament lengths at steady state. This mechanism may have biological significance in processes like mitotic spindle size scaling during development where relatively small changes in microtubule growth rates have been correlated with large changes in mitotic spindle size(Rieckhoff et al., 2020). Our study shows how the antagonistic action of assembly and disassembly factors can produce novel dynamical properties of microtubules.

Material and Methods

1. Calculation of probability of severing

Severing protein lands and creates a damage on a 1- dimensional grid of size N_D. The fate of the damage is determined by a competition between the removal and addition of tubulin at the damage site, eventually leading to severing or repair as shown in Fig. A2 (A) in the Appendix. We assume that monomers are added to the damage site at a rate proportional to the concentration of free tubulin $k_T[T]$ (s^{-1}) and are removed from the damage site at a rate k_r (s^{-1}), with each step either increasing or decreasing the size of damage. If the size of the damage reaches a size N_D, the filament is severed. On the other hand, if the size reaches zero, then the damage is repaired. Thus, the size of the damage x measured in monomers (i.e. tubulin dimers) is modeled as a one-dimensional random walk with absorbing boundaries at $x = 0$ and $x = N_D$.

Probability of severing $P_{sev}(x)$ is the probability that 1-D random walker, starting at position $x(0 \leq x \leq N_D)$ reaches the $x = N_D$ boundary before repair occurs, i.e., before $x = 0$ is reached. In this case, the probability of severing when the damage size is 0, i.e., $P_{sev}(x = 0) = 0$ and probability of severing when the damage size is N_D i.e., $P_{Sev}(x = N_D) = 1$. These two conditions will serve as our two boundary conditions.

If $0 < x < N_D$, then, we can write the following recursion equation

$$P_{sev}(x) = (1 - P_{rm})P_{sev}(x - 1) + P_{rm}P_{sev}(x + 1) \ (1)$$

where $P_{rm} = k_r/(k_r + k_T[T])$ is the probability of damage spot increasing in size (monomers being removed from the site) and $1 - P_{rm}$ is the probability of damage spot decreasing in size (monomers being added at the site).

We use the method of characteristic equations to solve the recursion equation. Substituting, $P_{sev}(x - 1) = 1$, $P_{sev}(x) = s$ and $P_{sev}(x + 1) = s^2$, we get

$$P_{rm}s^2 - s + (1 - P_{rm}) = 0 \ (2)$$

whose two solutions are $s_{1,2} = 1, \frac{1-P_{rm}}{P_{rm}}$. Solutions to (1) are $P_{sev}(x) = A(s_1)^x + B(s_2)^x$. Substituting, we get $P_{sev}(x) = A\left(\frac{1-P_{rm}}{P_{rm}}\right)^x + B$. Using boundary conditions, $P_{sev}(0) = 0$, $P_{sev}(N_D) = 1$, we get $B = -A$ and $A = \frac{1}{\left(\frac{1-P_{rm}}{P_{rm}}\right)^{N_D} - 1}$. We define a dimensionless tubulin concentration $\tilde{T} = \frac{1-P_{rm}}{P_{rm}} = \frac{k_T[T]}{k_r} = \frac{[T]}{[T]_c}$ which is the ratio of the tubulin concentration and a critical concentration $[T]_c$, which is the tubulin concentration where the rate of repair is equal to rate of severing, equal to $k_r/k_{T,}$. We find that $P_{sev}(x, [T]) = \frac{\tilde{T}^x - 1}{\tilde{T}^{N_D} - 1}$, which is a function of Tubulin concentration. We note that when $\tilde{T} << 1$, $P_{sev}(x, [T]) \approx 1 - \tilde{T}^x$ and when $\tilde{T} >> 1$, $P_{sev}(x, [T]) \approx$

$\left(\frac{1}{\tilde{T}}\right)^{N_D-x}$. Also, $P_{sev}(x,[T]) \le \frac{\tilde{T}^x}{\tilde{T}^{N_D}} = \left(\frac{k_r}{k_T[T]}\right)^{N_D-x}$, hence as the tubulin concentration

increases, $P_{sev}(x)$ decreases (Fig. 2 and A2).

If we assume an initial damage spot of a size $x = 1$ as we do in the manuscript,

then $P_{sev}(x = 1, [T]) = \frac{\tilde{T}-1}{\tilde{T}^{N_D}-1}$ which is approximately equal to $\left(\frac{1}{\tilde{T}}\right)^{N_D-1}$ when $\tilde{T} \gg$

1 and $P_{sev}(x,[T]) \approx 1 - \tilde{T}$ when $\tilde{T} \ll 1$. In other words, above the critical tubulin

concentration, the probability of severing becomes a very sensitive function of the

tubulin concentration.

2. Calculation for time to either sever or repair

Let $T^*_{end}(x)$ be the average time (in steps) it takes to for the filament to either get

severed or repaired, starting with a damage of size x. On the first step, the damage

spot can either get replenished with monomers with a probability $1 - P_{rm}$ or

increase in size by having monomers removed with a probability P_{rm} in time Δt i.e.

$$T^*_{end}(x) = \Delta t + T^*_{end}(x - \Delta x)(1 - P_{rm}) + T^*_{end}(x + \Delta x)P_{rm}$$

There are two possible boundary conditions, $T^*_{end}(0) = T^*_{end}(N_D) = 0$. Using $\Delta x =$

$\Delta t = 1$, we get a recursion relation similar to the one before, i.e. $T^*_{end}(x) =$

$T^*_{end}(x - 1)(1 - P_{rm}) + T^*_{end}(x + 1)P_{rm} + 1$, which is an inhomogeneous recursion

relation.

We first find the solution to the homogenous equation:

$$T^*_{end}(x - 1)(1 - P_{rm}) - T^*_{end}(x) + T^*_{end}(x + 1)P_{rm} = 0$$

This is similar to the one we solved in Section 1 and following the same method,

we can show that the solution is $T^{*homogenous}_{end}(x) = A\left(\frac{1-P_{rm}}{P_{rm}}\right)^x + B$ and the solution

to inhomogeneous equation we have to guess (since the inhomogeneous is a

constant) $T^{*inhomogenous}_{end}(x) = Cx + D$

Substituting it in the equation for $T^*_{end}(x)$ we get $C = \frac{x}{1-2P_{rm}}$ and hence the complete

solution is $T^*_{end}(x) = T^{*homogenous}_{end}(x) + T^{*inhomogenous}_{end}(x)$ which is $T^*_{end}(x) =$

$A\left(\frac{1-P_{rm}}{P_{rm}}\right)^x + B + \frac{x}{1-2P_{rm}}$

Using the boundary conditions, and assuming $P_{rm} \ne 1/2$, we find $T^*_{end}(x) =$

$\frac{x}{1-2P_{rm}} - \frac{N_D}{1-2P_{rm}} \frac{\left(\frac{1-P_{rm}}{P_{rm}}\right)^x - 1}{\left(\frac{1-P_{rm}}{P_{rm}}\right)^{N_D} - 1}$. When $P_{rm} = 1/2$, $T^*_{end}(x) = x\,(N_D - x)$.

Note that $T^*_{end}(x)$ is the time (in steps) taken to reach either end (filament gets repaired or severed). We get time taken by a damage spot to be repaired or severed, i.e. $T_{end}(x)$ by multiplying time in steps by $\Delta t_{avg} = \frac{1}{k_T[T]+k_r}$,

$$T_{end}(x,[T]) = \left(\frac{1/k_r}{\tilde{T}+1}\right)\left(\frac{\tilde{T}+1}{\tilde{T}-1}\right)\left(x - N_D\left(\frac{\tilde{T}^x-1}{\tilde{T}^{N_D}-1}\right)\right).\ (3)$$

where $P_{rm} = \frac{k_r}{k_r+k_T[T]}$ and $\tilde{T} = \frac{1-P_{rm}}{P_{rm}} = \frac{k_T[T]}{k_r}$. Equation 3 for $(x = 1)$ is plotted in Fig A1 (C).

3. Calculation of severing time

In this section, we derive an expression for time it takes to sever (equation 1 in main text) a filament. In the one-dimensional random walk model for damage spreading, there are two ways to reach the end – either get severed or get repaired. Hence, the time to reach either end is given by the following equation,

$$T_{end}(x) = P_{sev}(x)T_{sev}(x) + \left(1 - P_{sev}(x)\right)T_{sev}(N_D - x)\ (4)$$

where $T_{end}(x)$ is the time to reach either end if the initial size of the damage is x, $T_{sev}(x)$ is the time to sever and $P_{sev}(x)$ is the probability to sever when starting at x. Using the expression for $T_{end}(x)$ which we computed in Section 2 (Materials and Methods), we can extract $T_{sev}(x)$.

Replacing x with $N_D - x$ in equation 4 we get, $T_{end}(N_D - x) = P_{sev}(N_D - x)T_{sev}(N_D - x) + \left(1 - P_{sev}(N_D - x)\right)T_{sev}(x)$. Using $P_{sev}(x) = \frac{\tilde{T}^x-1}{\tilde{T}^{N_D}-1}$ and $P_{sev}(N_D - x) = \frac{\tilde{T}^{N_D-x}-1}{\tilde{T}^{N_D}-1}$, we get $1 - P_{sev}(x) = \frac{\tilde{T}^{N_D}-\tilde{T}^x}{(\tilde{T}^{N_D}-1)} = \tilde{T}^x P_{sev}(N_D - x)$ and $1 - P_{sev}(N_D - x) = \frac{\tilde{T}^{N_D}-1-\tilde{T}^{N_D-x}+1}{\tilde{T}^{N_D}-1} = \frac{\tilde{T}^{N_D-x}(\tilde{T}^x-1)}{\tilde{T}^{N_D}-1} = \tilde{T}^{N_D-x} P_{sev}(x)$.

From section 2 in Material and methods, $T_{end}(x) = \left(\frac{1/k_r}{\tilde{T}+1}\right)\left(\frac{\tilde{T}+1}{\tilde{T}-1}\right)\left(x - N_D\left(\frac{\tilde{T}^x-1}{\tilde{T}^{N_D}-1}\right)\right) = \left(\frac{1}{k_T[T]+k_r}\right)\left(\frac{\tilde{T}+1}{\tilde{T}-1}\right)\left(x - N_D P_{sev}(x)\right)$.

Substituting the expression for $1 - P_{sev}(x)$ and $1 - P_{sev}(N_D - x)$ in the equations for $T_{end}(x)$ and $T_{end}(N_D - x)$, we get

$$T_{sev}(x) = \frac{T_{end}(x) - \tilde{T}^x T_{end}(N_D - x)}{P_{sev}(x)\left(1 - \tilde{T}^{N_D}\right)}\ (5)$$

After substituting expression for $T_{end}(x)$ and $T_{end}(N_D - x)$ in (3), we get

$$T_{sev}(x) = \left(\frac{1}{k_T[T]+k_r}\right)\left(\frac{\tilde{T}+1}{\tilde{T}-1}\right)\left(\frac{x - N_D P_{sev}(x) - \tilde{T}^x(N_D(\tilde{T}^{N_D-x}P_{sev}(x)) - x)}{P_{sev}(x)\left(1 - \tilde{T}^{N_D}\right)}\right)$$

648 Finally, after substituting the expression for $P_{sev}(x)$ and moving terms around, we
649 get

$$T_{sev}(x) = \left(\frac{1/k_r}{\tilde{T}+1}\right)\left(\frac{\tilde{T}+1}{\tilde{T}-1}\right)\left(N_D\frac{\left(1+\tilde{T}^{N_D}\right)}{\left(\tilde{T}^{N_D}-1\right)} - \frac{x\left(\tilde{T}^x+1\right)}{\left(\tilde{T}^x-1\right)}\right) \quad (6)$$

For an initial damage size $x = 1, T_{sev}(k_T, k_r, N_D) = \left(\frac{\frac{1}{k_r}}{\tilde{T}-1}\right)\left(N_D\frac{\left(\tilde{T}^{N_D}+1\right)}{\left(\tilde{T}^{N_D}-1\right)} - \frac{\tilde{T}+1}{\tilde{T}-1}\right)$,
which is the equation we used to plot the mean severing time in Fig. 1D and A2(C)
inset (see Appendix).

$T_{sev}(k_T, k_r, N_D)$ has two limits: (1) When $\tilde{T} \ll 1$, and $N_D \gg 1$, $T_{sev}(k_T, k_r, N_D) =$
$-\left(\frac{\frac{1}{k_r}}{1-\tilde{T}}\right)\left(-N_D + \frac{\tilde{T}+1}{1-\tilde{T}}\right) \sim \frac{N_D}{k_r}\left(\frac{1}{1-\tilde{T}}\right) \sim \frac{N_D}{k_r}\left(1+\tilde{T}\right)$; the severing time is linear in tubulin
concentration. (2) When $\tilde{T} \gg 1$, and $N_D \gg 1$, $T_{sev}(k_T, k_r, N_D) = \left(\frac{1}{k_r\tilde{T}}\right)(N_D)$; the
severing time decreases as tubulin concentration increases. These limiting
behaviors imply that the severing time has a maximum as a function of tubulin
concentration, which is obtained at a concentration that is of the order critical
tubulin concentration $[T]_c = \frac{k_r}{k_T}$.

661
662 For the diffusive limit, when the rate of damage spreading and repair are balanced
663 i.e. when $\tilde{T} = 1$, $T_{sev}(k_T, k_r, x, N_D) = \frac{1}{6k_r}(N_D^2 - x^2)$.

664
4. Estimation of parameters

In this section, we compare our theoretical predictions with experimental results
from ref. (Vemu et al., 2018), and extract values for the parameters of our model.
In particular we focus on Fig. 3D and 3H in ref (Vemu et al., 2018), where the
authors plot probability to sever and time taken to sever for different concentrations
of tubulin.

Our model for severing with repair has four parameters - k_T, k_r, x and N_D. In
equation 6, we find an expression for the mean severing time as a function of the
tubulin concentration. We assume an initial damage of size $x = 1$, and constrain
the space of possible parameter values using experimental data in ref (Vemu et
al., 2018) . In equation 2, we predict that the probability of severing decreases as
N_D increases (Fig. A1). We use the observation of severing events at $[T] = 5\ \mu M$
in ref (Vemu et al., 2018) to constrain N_D (upper bound = 40) by assuming that
probability of severing is at least 10^{-3}. We obtain a rough lower bound for $N_D \approx 20$
from the TEM images in Fig. 1 in ref. (Vemu et al., 2018) showing patches of tubulin
removed from microtubules which are still intact. Next, we use the method of least
squares on this constrained parameter range to find sets of parameters that fit the

published data Fig. 3H in ref (Vemu et al., 2018) which reports the mean and the standard deviation of severing times as a function of tubulin concentration (Fig. A2). Our estimated values are: $k_T = 0.11\ \mu M^{-1}s^{-1}$, $k_r = 0.54\ s^{-1}$ and $N_D = 25$. Using these estimated values, we plot the severing time and the probability of severing in Fig. 2A and 2B respectively.

Sensitivity analysis of parameters: As seen in Fig. A1, probability of severing increases with a larger initial damage size x but severing time is unaffected. Probability of severing is unaffected by N_D but time to sever increases with a larger N_D. As is captured by the analytical formula in equation 2 in main text, the critical concentration of tubulin $[T_c]$ where the time of severing starts decreasing, is unaffected by x and N_D.

Comparison of experiment and theory via bootstrapping: In Figure 1 in the main text, we show a comparison of experimental data from (Vemu et al., 2018) with a theoretical curve for mean severing time and simulation curve for standard deviation of severing time using our fitted parameters. Since ref(Vemu et al., 2018) reported mean severing time and standard deviation of severing time, we used a bootstrapping scheme using data resampling to compute the error on the mean (error bars in Fig 1C), and error on the standard deviation (error bars in Fig 1D(inset)). The error bars on the mean severing time and standard deviation of severing times were calculated from 750 bootstrapping samples.

5. Stochastic simulation schemes

(A) Model for severing with repair

A severing event begins when a severing protein lands on a filament, binds to the side, and creates a damage of size x. The fate (eventual severing or repair) of the damage created by a severing protein is determined by a competition between the removal and addition of tubulin at the damage site. We use the Gillespie algorithm to simulate this process - monomers are added to the damage site at a rate proportional to the concentration of free tubulin $k_T[T]\ (s^{-1})$ and are removed from the damage site at a rate k_r (s⁻¹), with each step either increasing or decreasing the size of damage. If the size of the damage reaches a size N_D, the filament is severed. On the other hand, if the size reaches zero, then the damage is repaired. By performing this simulation multiple times (typically about 13,000 severing events), we obtain, as a function of tubulin concentration, the probability of severing (Fig. 2), the mean severing time (Fig. 1C) and the probability distribution of severing times (Fig. 1D).

(B) Length trajectory with repair-severing

Simulation of filament growth is performed using a Gillespie algorithm. At each step of the simulation, either the filament gains one subunit, with probability

proportional to the concentration of tubulin ($k\,[T]$), or it gains a new severing site
when a severing protein lands on the filament at position between 1 and $l-1$, with
probability k_r, where l is the length of the filament. In the absence of competition
between severing and repair, a severing event occurs immediately upon creating.
However, if competition is considered, each time a severing site is created, a
severing event occurs at this site with a probability $P_{sev}(x)$ in equation 1.

(C) Filament growth with finite severing time
In order to incorporate a finite severing time into our simulation, we first run our
damage spreading simulation (A) to create a distribution of severing times. Then,
we adjust the filament growth simulation in (B) by tracking the location of each
damage site (where severing protein binds) on our microtubule. The severing
occurs with a probability given by equation (2). For every severing event, we assign
a severing time by sampling it from our distribution of severing times. Once this
time has surpassed in the simulation, the microtubule is severed at this damage
site. We keep track of one part of the severed microtubule. The other part, and any
damage sites on that part, are removed.

6. Calculation of probability distribution of filament lengths, due to
severing with and without repair

In ref(Mohapatra et al., 2016), we considered the case of a uniform rate of severing
along a filament; in other words, severing takes place anywhere along the filament
with equal probability. A filament consisting of l subunits can be broken into two
smaller filaments at any of the $(l-1)$ positions with an equal rate s, for any choice
of severing location; the total rate of severing at any location is then, $s\,(l-1)$. In
addition to the severing rate, there is r, the rate at which subunits are added to a
filament. Using these two rates we find an expression for the probability distribution
of lengths at steady state (reproduced from ref(Mohapatra et al., 2016)), $P(l) =$
$\dfrac{l\,s\,r^{l-1}}{(r+s)(r+2s)(r+3s)\dots(r+ls)}$, which can be further simplified to

755
$$P(l) = \frac{l\,s\,r^{l-1}}{r^l\left(1+\frac{s}{r}\right)\left(1+\frac{2s}{r}\right)\dots\left(1+\frac{ls}{r}\right)} = \frac{ls}{r}\,e^{-\frac{s}{r}\sum_{j=1}^{l}j} = \frac{ls}{r}\,e^{-\frac{s}{r}\frac{l(l+1)}{2}} \quad (7)$$

Here we use the approximation that $e^{js/r} = 1 + \frac{js}{r}$ which is valid for $js \ll r$. For $l >$
> 1 , $P(l) = \frac{ls}{r}e^{-\frac{s}{r}\frac{l^2}{2}}$ and we use this expression to plot the distribution of lengths
without repair with rates $r = k[T]$ and $s = k_d[S]$, and with repair with the rates $r =$
$k[T]$ and $s = k_d[S]P_{sev}([T])$ in Fig. 3D.

Additionally, we can use this approximation to compute the mean $< l >$ and
variance of the filament length distribution, $< l^2 > - < l >$. In general, moments

are defined as, $<l^n> = \frac{\int l^n P(l)dl}{\int P(l)dl}$. For the mean, $n = 1$, and hence $<l> =$

$\frac{\int l^2 \frac{S}{r} e^{-\frac{S}{r}\frac{l^2}{2}} dl}{\int \frac{ls}{r} e^{-\frac{S}{r}\frac{l^2}{2}} dl}$.

Using the properties of gaussian integrals, $\int_0^\infty e^{-\alpha \frac{l^2}{2}} dl = \frac{1}{2}\sqrt{\frac{2\pi}{\alpha}}$, we can show

$\int_0^\infty l e^{-\alpha \frac{l^2}{2}} dl = \frac{1}{\alpha}$, and $\int_0^\infty l^3 e^{-\alpha \frac{l^2}{2}} dl = \frac{2}{\alpha^2}$. Substituting these results, we get $<l>$

$= \frac{\int \alpha l^2 e^{-\alpha \frac{l^2}{2}} dl}{\int \alpha l e^{-\alpha \frac{l^2}{2}} dl} = \sqrt{\frac{\pi}{2\alpha}}$ and $<l^2> = \frac{\int l^3 e^{-\alpha \frac{l^2}{2}} dl}{\int l e^{-\alpha \frac{l^2}{2}} dl} = \frac{2}{\alpha}$.

The variance, $<l^2> - <l>$, is $\frac{2}{\alpha} - \frac{\pi}{2\alpha}$ and standard deviation, $\sigma = \sqrt{\frac{1}{\alpha}\left(2 - \frac{\pi}{2}\right)}$.

We define "Noise" or the coefficient of variation as $\frac{\sigma}{<l>} = \sqrt{\frac{2-\frac{\pi}{2}}{\frac{\pi}{2}}} = \sqrt{\frac{4}{\pi} - 1} = 0.523$,

which is a prediction for the uniform severing mechanism. We use this relationship

to plot Fig. 3D, inset. Additionally, substituting $\alpha = \frac{k_d[S]}{k[T]}$ in to the expression for the

mean and variance of filament lengths at steady state, we get, $L_s^{simple\ severing} = <$

$l> = \sqrt{\frac{\pi k[T]}{2\,k_d[S]}}$, and substituting $\alpha = \frac{k_d[S]P_{sev}([T])}{k[T]}$, we find $L_s^{with\ repair} =$

$\sqrt{\frac{\pi k[T]}{2\,k_d[S]P_{sev}([T])}}$, which is used in Fig. 3C.

Note that $L_s^{with\ repair}$ has two regimes: When $[T] < [T]_c$, $P_{sev}([T]) \approx 1 - \tilde{T}$,

$L_s \sim \sqrt{[T]}$. When $[T] > [T]_c$, $P_{sev}([T]) \approx \left(\frac{1}{\tilde{T}}\right)^{N_D - 1}$ and, $L_s \approx [T]^{N_D/2}$.

 Appendix

791 **Fig A1. Estimation of parameters**. (A) and (B) Density plots for the estimation of
792 k_r, k_T and N_D. Loss function calculates the sum of squares of differences between
793 mean severing time from equation (6) with indicated parameters and experimental
794 data for [T]=0,2,5 µM. Probability of severing and Mean severing time vs
795 concentration of Tubulin as a function of (C) x and (D) N_D.

Fig A2. Comparison of theory and simulation for severing with repair. (A) Schematic of damage spreading in a damage site, corresponding to Figure 1B in the main text. Damage spreads along the microtubule lattice, i.e., tubulin dimers are removed, at a rate k_r, and damage is repaired, i.e. tubulin dimers are incorporated back into the microtubule lattice a rate $k_T[T]$. (B) Probability of severing and (C) time to repair or sever and mean severing time (inset) as a function of tubulin concentration. All plots were computed using 3000 damage spreading simulations, and the SEM error (in blue) was calculated by repeating these 2000 times. Parameters used are listed in Table 1.

A1: Finite severing time

If the severing protein binding time is comparable to severing time, it is possible that, after one severing event, there are other damage sites on the filament which will lead to a severing event shortly thereafter. This will produce correlations in subsequent severing events which we ignore when deriving our analytic formulas for P_{sev} and T_{sev}. In order to account for this situation, we performed a simulation where we allowed multiple damage sites on a microtubule lattice at one time, see Section 5(C) in Materials and Methods. We observed that our results of large dynamic range in steady-state length from the full simulation are similar to what was seen with the simple model where we assumed a negligible severing time (Fig. A3). The coefficient of variation (CV) of filament lengths at steady state calculated in the full simulation was identical to the one obtained from the simple model at low and high tubulin concentration. However, at intermediate values of tubulin concentrations, we observed a peak in the CV as seen in Fig. A3(C), which

was in contrast to the constant value seen with the simple model (Fig. 3D). We
observed that the peak in CV depends on the concentration of severing protein as
shown in Fig. A3(D). As shown in inset, the variation from the typical constant value
occurs when the time between binding events which can lead to severing
(computed by $\frac{1}{k_d[S]\,L_s\,P_{sev}}$ where L_s is the length of microtubule) is less than the time
it takes to sever.

Fig A3. Effect of finite severing time on filament dynamics. (A) Length
trajectory of a microtubule (average of three for clarity) which takes into account
the finite time it takes to sever a filament, at different concentrations of tubulin. (B)
Probability distribution of steady state lengths, computed using 10^6 trajectories.
(C) Mean steady state length and coefficient of variation (CV) (inset) with and
without repair (with and without finite severing time), computed using 10^6
trajectories. (D) Adding a finite time of severing does not change the results
qualitatively for mean length, but there is a peak in the CV that depends on the
severing protein concentration; the CV was computed using 3×10^6 trajectories.
(Inset) The CV differs from its constant value at tubulin concentrations at which the
time between productive events that lead to severing ("effective bind time") is less

than the severing time. Effective bind time is given by $\frac{1}{k_d[S]L_s P_{sev}}$. Parameters used
in all simulations are listed in Table 1.

A2: Dynamic instability

The simulation that includes dynamic instability follows the same algorithm as that
detailed in Section 5(B) in Materials and Methods, with inclusion of an additional
boolean parameter for whether the microtubule is in a state of rescue or
catastrophe. In the rescue state, the microtubule grows (gain subunits) at a rate
proportional to the tubulin concentration, $c_T[T]$. During catastrophe, the
microtubule loses subunits at a rate c_s. The microtubule can undergo severing in
either state. At each timestep, the microtubule can a) grow or shrink depending on
whether it is in catastrophe or rescue, b) have a severing protein land and either
sever the filament or not depending on the outcome of the competition between
severing and repair, or c) switch from catastrophe to rescue or rescue to
catastrophe. We use the parameters listed in Table 1 and A1 for our stochastic
simulations and are in the regime where the growth/shrinking rates are faster than
the rescue/catastrophe rates. As before, we find that the dynamic range of the
mean length of microtubules is greater when the repair process is included (Fig.
A4). However, coefficient of variation shows a non-monotonic relationship with
tubulin concentration.

A3: Conversion between monomer number and filament length

Our simulations are based in subunits. We use the following conversion to convert
from subunits to length: (4/13) (length in # of subunits) = (length in nm).
Microtubules are hollow cylinders and made up of 13 protofilaments built up from
tubulin dimers, each 4 nm in length. A cross section of a microtubule is a rind of
13 dimers that is 4 nm wide.

Fig A4. **Effect of Dynamic Instability**. (A) Length trajectory (average of 3) with
repair and (B) without repair. (C) Mean steady state length, computed using 10^6
trajectories and (D) coefficient of variation without and with repair with two rescue
rates, computed using 10^6 trajectories. Parameters used are listed in Table A1.

872
873

876
877

Table A1. Parameters use for studying Length control in presence of repair with dynamic instability.

Parameters	Value used in paper	References
Catastrophe rate	.004 s^{-1}	(Margolin et al., 2012)
Rescue rate	.02 s^{-1}	(Margolin et al., 2012)
Shrinkage rate in catastrophe	152 dimers/s	(Margolin et al., 2012)

Acknowledgments

We wish to thank Stephanie C. Weber and Shane G. McInally for careful reading of the manuscript, and members of Mohapatra group and Kondev group for stimulating discussions. This work was supported by the National Science Foundation through grants DMR-1610737, MRSEC DMR 2011846 (J.K., C.S.), the Simons and the HHMI foundations (J.K., L.M. and C.S) and Rochester Institute of Technology start-up funds (L.M.). We acknowledge computational support from the Brandeis HPCC which is partially supported by the NSF through DMR-MRSEC 2011846 and OAC-1920147.

Competing interests

Authors declare no competing interests.

References

Ahmad FJ, Yu W, McNally FJ, Baas PW. 1999. An Essential Role for Katanin in Severing Microtubules in the Neuron. *Journal of Cell Biology* **145**:305–315. doi:10.1083/jcb.145.2.305

Arellano-Santoyo H, Geyer EA, Stokasimov E, Chen G-Y, Su X, Hancock W, Rice LM, Pellman D. 2017. A Tubulin Binding Switch Underlies Kip3/Kinesin-8 Depolymerase Activity. *Dev Cell* **42**:37-51.e8. doi:10.1016/j.devcel.2017.06.011

Arpağ G, Lawrence EJ, Farmer VJ, Hall SL, Zanic M. 2020. Collective effects of XMAP215, EB1, CLASP2, and MCAK lead to robust microtubule treadmilling. *Proc Natl Acad Sci U S A* **117**:12847–12855. doi:10.1073/pnas.2003191117

Balcer HI, Goodman AL, Rodal AA, Smith E, Kugler J, Heuser JE, Goode BL. 2003. Coordinated regulation of actin filament turnover by a high-molecular-weight Srv2/CAP complex, cofilin, profilin, and Aip1. *Curr Biol* **13**:2159–2169.

Belonogov L, Bailey ME, Tyler MA, Kazemi A, Ross JL. 2019. Katanin catalyzes microtubule depolymerization independently of tubulin C-terminal tails. *Cytoskeleton (Hoboken)* **76**:254–268. doi:10.1002/cm.21522

912   Bennabi I, Terret M-E, Verlhac M-H. 2016. Meiotic spindle assembly and chromosome
913   segregation in oocytes. *Journal of Cell Biology* **215**:611–619. doi:10.1083/jcb.201607062
914   Berg HC. 1993. Random Walks in Biology. Princeton University Press.
915   Chaudhry F, Breitsprecher D, Little K, Sharov G, Sokolova O, Goode BL. 2013. Srv2/cyclase-
916   associated protein forms hexameric shurikens that directly catalyze actin filament
917   severing by cofilin. *Mol Biol Cell* **24**:31–41. doi:10.1091/mbc.E12-08-0589
918   Desai A, Mitchison TJ. 1997. Microtubule polymerization dynamics. *Annu Rev Cell Dev Biol*
919   **13**:83–117. doi:10.1146/annurev.cellbio.13.1.83
920   Desai A, Verma S, Mitchison TJ, Walczak CE. 1999. Kin I Kinesins Are Microtubule-Destabilizing
921   Enzymes. *Cell* **96**:69–78. doi:10.1016/S0092-8674(00)80960-5
922   Díaz-Valencia JD, Morelli MM, Bailey M, Zhang D, Sharp DJ, Ross JL. 2011. Drosophila Katanin-60
923   Depolymerizes and Severs at Microtubule Defects. *Biophysical Journal* **100**:2440–2449.
924   doi:10.1016/j.bpj.2011.03.062
925   Dogterom M, Leibler S. 1993. Physical aspects of the growth and regulation of microtubule
926   structures. *Phys Rev Lett* **70**:1347–1350. doi:10.1103/PhysRevLett.70.1347
927   Dumont S, Mitchison TJ. 2009. Force and length in the mitotic spindle. *Curr Biol* **19**:R749-761.
928   doi:10.1016/j.cub.2009.07.028
929   Elam WA, Kang H, De la Cruz EM. 2013. Biophysics of actin filament severing by cofilin. *FEBS Lett*
930   **587**:1215–1219. doi:10.1016/j.febslet.2013.01.062
931   Gardner MK, Zanic M, Howard J. 2013. Microtubule Catastrophe and Rescue. *Curr Opin Cell Biol*
932   **25**:14–22. doi:10.1016/j.ceb.2012.09.006
933   Geisterfer ZM, Zhu DY, Mitchison TJ, Oakey J, Gatlin JC. 2020. Microtubule Growth Rates Are
934   Sensitive to Global and Local Changes in Microtubule Plus-End Density. *Curr Biol*
935   **30**:3016-3023.e3. doi:10.1016/j.cub.2020.05.056
936   Goshima G, Wollman R, Stuurman N, Scholey JM, Vale RD. 2005. Length control of the
937   metaphase spindle. *Curr Biol* **15**:1979–1988. doi:10.1016/j.cub.2005.09.054
938   Hyman AA, Karsenti E. 1996. Morphogenetic Properties of Microtubules and Mitotic Spindle
939   Assembly. *Cell* **84**:401–410. doi:10.1016/S0092-8674(00)81285-4
940   Jinushi-Nakao S, Arvind R, Amikura R, Kinameri E, Liu AW, Moore AW. 2007. Knot/Collier and Cut
941   Control Different Aspects of Dendrite Cytoskeleton and Synergize to Define Final Arbor
942   Shape. *Neuron* **56**:963–978. doi:10.1016/j.neuron.2007.10.031
943   Johnston AB, Collins A, Goode BL. 2015. High-speed depolymerization at actin filament ends
944   jointly catalysed by Twinfilin and Srv2/CAP. *Nat Cell Biol* **17**:1504–1511.
945   doi:10.1038/ncb3252
946   Karabay A, Yu W, Solowska JM, Baird DH, Baas PW. 2004. Axonal growth is sensitive to the levels
947   of katanin, a protein that severs microtubules. *J Neurosci* **24**:5778–5788.
948   doi:10.1523/JNEUROSCI.1382-04.2004
949   King MR, Petry S. 2020. Phase separation of TPX2 enhances and spatially coordinates
950   microtubule nucleation. *Nat Commun* **11**:270. doi:10.1038/s41467-019-14087-0
951   Kuhn JR, Pollard TD. 2005. Real-time measurements of actin filament polymerization by total
952   internal reflection fluorescence microscopy. *Biophys J* **88**:1387–1402.
953   doi:10.1529/biophysj.104.047399
954   Kuo Y-W, Howard J. 2021. Cutting, Amplifying, and Aligning Microtubules with Severing
955   Enzymes. *Trends Cell Biol* **31**:50–61. doi:10.1016/j.tcb.2020.10.004
956   Kuo Y-W, Trottier O, Howard J. 2019a. Predicted Effects of Severing Enzymes on the Length
957   Distribution and Total Mass of Microtubules. *Biophysical Journal* **117**:2066–2078.
958   doi:10.1016/j.bpj.2019.10.027

Kuo Y-W, Trottier O, Mahamdeh M, Howard J. 2019b. Spastin is a dual-function enzyme that
severs microtubules and promotes their regrowth to increase the number and mass of
microtubules. *Proc Natl Acad Sci U S A* **116**:5533–5541. doi:10.1073/pnas.1818824116
Loughlin R, Wilbur JD, McNally FJ, Nédélec FJ, Heald R. 2011. Katanin Contributes to Interspecies
Spindle Length Scaling in Xenopus. *Cell* **147**:1397–1407. doi:10.1016/j.cell.2011.11.014
Margolin G, Gregoretti IV, Cickovski TM, Li C, Shi W, Alber MS, Goodson HV. 2012. The
mechanisms of microtubule catastrophe and rescue: implications from analysis of a
dimer-scale computational model. *MBoC* **23**:642–656. doi:10.1091/mbc.e11-08-0688
Marshall WF, Qin H, Rodrigo Brenni M, Rosenbaum JL. 2005. Flagellar length control system:
testing a simple model based on intraflagellar transport and turnover. *Mol Biol Cell*
16:270–278. doi:10.1091/mbc.E04-07-0586
Mickolajczyk KJ, Geyer EA, Kim T, Rice LM, Hancock WO. 2019. Direct observation of individual
tubulin dimers binding to growing microtubules. *Proceedings of the National Academy*
of Sciences **116**:7314–7322. doi:10.1073/pnas.1815823116
Miyoshi T, Watanabe N. 2013. Can filament treadmilling alone account for the F-actin turnover
in lamellipodia? *Cytoskeleton (Hoboken)* **70**:179–190. doi:10.1002/cm.21098
Mohapatra L, Goode BL, Jelenkovic P, Phillips R, Kondev J. 2016. Design Principles of Length
Control of Cytoskeletal Structures. *Annu Rev Biophys*. doi:10.1146/annurev-biophys-
070915-094206
Ohi R, Strothman C, Zanic M. 2021. Impact of the 'tubulin economy' on the formation and
function of the microtubule cytoskeleton. *Current Opinion in Cell Biology* **68**:81–89.
doi:10.1016/j.ceb.2020.09.005
Pollard TD. 1986. Rate constants for the reactions of ATP- and ADP-actin with the ends of actin
filaments. *J Cell Biol* **103**:2747–2754.
Rieckhoff EM, Berndt F, Elsner M, Golfier S, Decker F, Ishihara K, Brugués J. 2020. Spindle Scaling
Is Governed by Cell Boundary Regulation of Microtubule Nucleation. *Curr Biol* **30**:4973-
4983.e10. doi:10.1016/j.cub.2020.10.093
Roland J, Berro J, Michelot A, Blanchoin L, Martiel J-L. 2008. Stochastic severing of actin
filaments by actin depolymerizing factor/cofilin controls the emergence of a steady
dynamical regime. *Biophys J* **94**:2082–2094. doi:10.1529/biophysj.107.121988
Sharp DJ, Ross JL. 2012. Microtubule-severing enzymes at the cutting edge. *J Cell Sci* **125**:2561–
2569. doi:10.1242/jcs.101139
Shekhar S, Chung J, Kondev J, Gelles J, Goode BL. 2019. Synergy between Cyclase-associated
protein and Cofilin accelerates actin filament depolymerization by two orders of
magnitude. *Nat Commun* **10**. doi:10.1038/s41467-019-13268-1
Song L, Dentler WL. 2001. Flagellar Protein Dynamics in Chlamydomonas. *Journal of Biological*
Chemistry **276**:29754–29763. doi:10.1074/jbc.M103184200
Tindemans SH, Mulder BM. 2010. Microtubule length distributions in the presence of protein-
induced severing. *Phys Rev E* **81**:031910. doi:10.1103/PhysRevE.81.031910
Tournebize R, Popov A, Kinoshita K, Ashford AJ, Rybina S, Pozniakovsky A, Mayer TU, Walczak
CE, Karsenti E, Hyman AA. 2000. Control of microtubule dynamics by the antagonistic
activities of XMAP215 and XKCM1 in Xenopus egg extracts. *Nat Cell Biol* **2**:13–19.
doi:10.1038/71330
Van Geel O, Cheung S, Gadella TWJ. 2020. Combining optogenetics with sensitive FRET imaging
to monitor local microtubule manipulations. *Scientific Reports* **10**:6034.
doi:10.1038/s41598-020-62874-3

Varga V, Leduc C, Bormuth V, Diez S, Howard J. 2009. Kinesin-8 motors act cooperatively to
mediate length-dependent microtubule depolymerization. *Cell* **138**:1174–1183.
doi:10.1016/j.cell.2009.07.032
Vemu A, Szczesna E, Zehr EA, Spector JO, Grigorieff N, Deaconescu AM, Roll-Mecak A. 2018.
Severing enzymes amplify microtubule arrays through lattice GTP-tubulin incorporation.
Science **361**. doi:10.1126/science.aau1504
Walczak CE, Gayek S, Ohi R. 2013. Microtubule-Depolymerizing Kinesins. *Annu Rev Cell Dev Biol*
29:417–441. doi:10.1146/annurev-cellbio-101512-122345
Watanabe N, Mitchison TJ. 2002. Single-molecule speckle analysis of actin filament turnover in
lamellipodia. *Science* **295**:1083–1086. doi:10.1126/science.1067470
Woodruff JB, Ferreira Gomes B, Widlund PO, Mahamid J, Honigmann A, Hyman AA. 2017. The
Centrosome Is a Selective Condensate that Nucleates Microtubules by Concentrating
Tubulin. *Cell* **169**:1066-1077.e10. doi:10.1016/j.cell.2017.05.028
Zanic M, Widlund PO, Hyman AA, Howard J. 2013. Synergy between XMAP215 and EB1 increases
microtubule growth rates to physiological levels. *Nat Cell Biol* **15**:688–693.
doi:10.1038/ncb2744
Ziółkowska NE, Roll-Mecak A. 2013. In vitro microtubule severing assays. *Methods Mol Biol*
1046:323–334. doi:10.1007/978-1-62703-538-5_19
Zwetsloot AJ, Tut G, Straube A. 2018. Measuring microtubule dynamics. *Essays Biochem* **62**:725–
735. doi:10.1042/EBC20180035
1025