Prediction of allosterically acting mutations using MD simulations and Rosetta

Nicholas G. M. Wells[1], Colin A. Smith[1,*]

[1]Department of Chemistry, Wesleyan University

[*]Corresponding Author:
Wesleyan University
52 Lawn Ave
+1 860 685-3274
colin.smith@wesleyan.edu

Running Title: Prediction of allosterically acting mutations using MD simulations and Rosetta

Manuscript Pages: 38
Supplementary Figures: 5

Abstract

Computationally modeling how mutations affect protein binding not only helps uncover the biophysics of protein interfaces, but also enables the redesign and optimization of protein-protein interactions. Traditional high-throughput methods for estimating binding free energies are currently limited to mutations directly at the interface due to difficulties in accurately modeling how more distant allosterically acting mutations propagate their effects through the protein structure. However, the modeling and design of such allosteric mutations is of substantial interest as it allows for greater control and flexibility in protein design applications. We have developed a method that combines high-throughput Rosetta-based side-chain optimization with conformational sampling using classical molecular dynamics simulations, finding significant improvements in our ability to accurately predict allosterically acting mutational perturbations to protein binding. Our approach uses an analytical framework grounded in rigorous alchemical free energy calculations while enabling exploration of a vastly larger sequence space. When comparing to experimental data, we find that our method can predict allosteric perturbations with a level of accuracy similar to that of traditional methods in predicting the effects of non-allosteric interface mutations. This work represents a new and generalizable approach to optimize protein free energy landscapes for desired biological functions.

Author Summary

Protein-protein interactions are vital to almost all biological processes, and therefore the ability to accurately and efficiently predict how mutations alter protein binding has far-reaching applications in protein analysis and design. Current

approaches to predict such mutational free energy changes are limited to mutations

directly at the interaction interface. Much research has underlined the prevalence of

allosteric protein regulation in biological processes, indicating the importance of

understanding and predicting the effects of mutations which act over long distances. In

this work we develop a novel method based on theoretically rigorous alchemical free

energy calculations and the Rosetta macromolecular modeling suite which can predict

the effects of allosteric mutations with levels of accuracy rivaling state of the art

interface-specific methods. We hope that our method will serve as a novel framework

for high-throughput allosteric mutational analysis and therefore benefit future protein

design efforts.

Introduction

Protein interactions are central to many vital biological processes such as antibody binding, signal transduction, gene expression, and enzyme regulation. Consequently, changes in the propensity of proteins to associate or dissociate, for example due to mutation or post-translational modification, can be very disruptive to biological function and are therefore often linked to disease(1-3). The ability to understand and control the energetics of protein interactions therefore has far-reaching consequences for the study of disease and for the development of novel protein-based therapeutics.

A number strategies have been developed to quantitatively predict the effects of mutations on protein interactions based on either empirical(4-6) or statistically-based(7, 8) scoring approaches, where energy functions or observations from known protein structures, respectively, are used to describe the physical interactions between protein residues and estimate free energy changes upon mutation. The relatively low computational cost of these scoring methods makes them well suited to large-scale analyses and protein design, albeit at the expense of accuracy(9). Modest improvements have been made to score-based approaches by integrating information from MD simulations to account for solvent effects and variations in protein conformation(10, 11). Additionally, approaches which utilize the more physics-based scoring capabilities of the Rosetta macromolecular modeling suite have recently shown a promisingly high degree of accuracy in predicting mutational binding free energy changes(12-14), finding notable success in design applications(15, 16). However, despite this progress, current physics-based scoring methods are almost exclusively

limited to predicting the energetic effects of mutations at the protein interaction interface and fail to accurately predict the effects of allosterically active mutations.

Other approaches to calculating mutational free energy changes involve applying theoretically rigorous statistical mechanics and conformational sampling from MD simulations to calculate, rather than approximate through heuristics, mutational free energy changes(17-19). Free Energy Perturbation (FEP) is a method to determine the free energy difference between two states by simulating the system in one state while simultaneously evaluating the energies of the system in another state(19, 20). FEP is based on the Zwanzig equation(20), which states that the free energy difference between two states is equal to the exponential average of the energy differences between the two states:

$$\Delta F(A \rightarrow B) = -\beta^{-1} \ln \langle \exp\left(-\frac{E_B - E_A}{k_B T}\right) \rangle_A$$

Here, $\Delta F(A \rightarrow B)$ is the free energy difference between states A and B, β is equal to the inverse of the product of temperature (T) and Boltzmann's constant (k_B), E_A and E_B are the energies of conformations in state A and B respectively, and angular brackets denote the average over conformations taken from an equilibrium ensemble of state A.

While this theorem is numerically exact with an infinite sample of energy differences and only requires simulation of one equilibrium ensemble, it is highly reliant on sufficient sampling of the equilibrium system to achieve acceptable levels of accuracy and convergence(21, 22). It also often requires long and computationally expensive MD simulations simulated at many intermediate states when applied to typical biomolecular systems.

Methods like the Bennet acceptance ratio (BAR) and thermodynamic integration (TI) are non-equilibrium alternatives to FEP which involve calculating the work of a transition in the "forward" as well as the "reverse" directions, i.e. $w_{A \to B}$ and $w_{B \to A}$(23). Such methods have shown high degrees of accuracy in predicting the effects of mutations, both interface-adjacent and allosteric, on biophysical processes such as protein association(24, 25). However, while these "bidirectional" approaches significantly improve the accuracy and convergence of free energy estimates compared to their "unidirectional" counterparts(26), they further increase the need for computationally expensive MD simulations by requiring separate equilibrium simulations for each state of interest to allow calculation of a set of "reverse" work values. In addition to substantial equilibrium sampling, these methods also often rely on computationally expensive simulations to generate non-equilibrium work values(17, 24, 26), making them less suitable for large-scale applications.

Recent studies have emphasized the importance of allosteric regulation in protein interactions throughout biology(27-29), underscoring the power of mutations which can propagate their effects over long distances. The ability to predict the effects of such mutations would significantly improve the current scope of protein design and allow for a greater understanding of disease processes which feature allosterically regulated protein interactions. To the best of our knowledge there does not currently exist an accurate high-throughput method to predict the effects of allosteric mutations on protein-protein interactions. In the present study, we develop the Molecular Dynamics Rosetta-Free Energy Perturbation (MDR-FEP) method which combines the extensive conformational sampling from classical MD simulations with the high-

throughput nature of Rosetta repacking and the theoretical framework of alchemical FEP, allowing for the accurate prediction of allosterically active mutational free energy changes with a high enough throughput for large-scale mutational analyses and forward protein design.

Results and Discussion

Protocol Rationale and Overview

The effect of mutations directly at the protein-protein interface can often be modeled using algorithms that keep the backbone fixed and allow side chain flexibility(14). However, the farther a mutation is away from the protein-protein interface, the more likely it is to affect binding through backbone perturbations like shifts in secondary structure or protein loops. Prediction of those backbone shifts can be done with two different strategies. An "induced-fit" approach involves first making the mutation and then using structural refinement algorithms to predict how the mutation propagates changes to the rest of the structure and alters the binding free energy. Alternatively, a "conformational selection" approach involves first obtaining an ensemble of backbone structures with the wild-type sequence and then later determining whether given mutations will sufficiently alter the energetics of that backbone ensemble to affect the binding free energy.

For the purposes of this work, we assume that sampling of backbone conformations is significantly more computationally costly than sampling of side chain conformations. This is largely supported by the high performance of algorithms like the Rosetta packer(30), which can very efficiently sample both rotamer and sequence space, and the considerably more complex task of backbone sampling. Given this

dramatic difference in performance between the sampling of backbone and side chain conformations, a "conformational selection" algorithm that creates an ensemble of backbone conformations once and then samples sequence space many times is likely to be much more efficient than an "induced-fit" approach. While a "conformational selection" algorithm may perform poorly on mutations that require larger shifts of the backbone ensemble than would be captured in an "induced-fit" approach, there likely remains a large number of mutations that represent low hanging fruit amenable to conformational selection.

We hypothesized that equilibrium molecular dynamics simulations of a WT protein are an effective means of generating a backbone ensemble for a "conformational selection" approach. To be effective, the MD simulations should transiently sample backbone conformations for which the energy with a given mutant sidechain is lower than WT, providing information about that mutation's relative ΔG in the free or bound state. Because backbone conformations which are more stable with a mutant sidechain could have higher energies than with the WT sidechain present, these conformations may be infrequently sampled in WT simulations. Such infrequent sampling potentially leads to a relatively small contribution of these low-energy conformations to the overall ensemble, resulting in negligible shifts in the mean or median energies for the WT and mutant sequences. To detect such shifts at the extremes of the energy distributions, we tested whether the use of analytical methods rooted in statistical mechanics enables better accounting for the contribution of these low-energy conformations to the overall ΔG value. In the present work, the Rosetta packer is used to side-chain optimize and score the energy of each frame in an

equilibrium simulation with both WT and mutant sequences. These energies are then evaluated using exponential averaging in the Zwanzig equation (above) to produce a ΔG value.

The β parameter of that function, which describes the inverse of the equilibrated system's theoretical temperature multiplied by the Boltzmann constant, dictates the extent to which low-energy conformations contribute to ΔG: At low values of β, frames with a high ΔE dominate while those with a low ΔE contribute very little. Conversely, an increase in β causes the contribution of low-energy conformations to increase (Figure S1). In the proposed method, β is treated as a parameter to be optimized in order to find the ideal level of contribution of these low-energy frames to ΔG. This optimization is warranted because energies output by Rosetta are in somewhat abstract Rosetta Energy Units (REU). Furthermore, optimization with the Rosetta packer is a distinctly non-equilibrium process for which an appropriate temperature is not known *a priori*.

For a protein dimer of chains A and B, WT MD simulations are run with both dimeric and monomeric chains (Figure 1A, Step 1). Individual structural snapshots are extracted every 1 ns from each trajectory. For each frame, all residues within 10 Å of the mutation of interest are repacked and scored with both the WT and mutant sidechain present, generating a distribution of ΔE values for both the monomer and dimer states (Figure 1A, Step 2). The set of ΔE values generated for each mutation in the monomer and dimer states is converted into a distribution using kernel density estimation (Figure 1B-C). In order to account for the possibility of outliers, a cutoff is then applied to this density function where all values which fall below the cutoff are set equal to zero. The Zwanzig equation is then applied to each distribution to generate a

ΔG value for the monomer and dimer, allowing for the calculation of $\Delta\Delta G_{Dimerization}$ for the mutation of interest.

Parameter Optimization and Performance Evaluation

We first performed a grid search over three separate mutation datasets in order to determine optimal β and cutoff parameters for the proposed method (Figure 2). Correlation with experimental data was evaluated with β ranging from 0 to 0.1 and cutoff ranging between 0 and 1%. We first evaluated our method against an experimental surface plasmon resonance alanine scan(31) of urokinase-urokinase receptor dimerization for both interior allosteric and interface-adjacent (non-allosteric) mutations (Figure 2A-B). Interestingly, we found that the method's performance in predicting the effects of both allosteric and non-allosteric mutations in the urokinase-urokinase receptor interaction seemed to depend on the cutoff parameter, with correlations appearing to decrease with cutoffs larger than 0.05% and 0.03%, respectively (Figure 2A/B). We interpret these results to indicate that the accurate prediction of the ΔG of these mutations relies more on the sampling of low-energy frames, perhaps due to the increased difficulty of simulating sterically larger mutations to alanine. Conversely, the correlation for each of these sets of mutations seemed to depend less on the β parameter (Figure 2A/B).

We then evaluated the MDR-FEP method using an experimental ITC study(32) of the barnase-barstar dimerization process which included only non-allosteric (interface-adjacent) mutations (Figure 2C), finding that our method was able to reproduce experimental data with reasonable accuracy independent of the cutoff parameter while using a low β value. Correlation with the experimental ITC data did however decrease sharply at higher β values (Figure 2C). The fact that correlation did not decrease as

more low-energy frames were removed by the increasing cutoff parameter suggests

that the accurate prediction of ΔG for these mutations does not rely on the sampling of

low-energy frames. Although the 1BRS and 3BT1 simulations sampled approximately

the same number of low-ΔG conformations (Figure S2), we believe that mutations in the

1BRS system are overall easier to predict by Rosetta repacking alone and therefore rely

less on the sampling of frames which approximate mutation-accommodating backbone

conformations.

We then compared the performance of the proposed method to that of Flex

ddG(12), an existing Rosetta-based $\Delta\Delta G_{Dimerization}$ calculation method. Over a set of 65

allosteric mutations to alanine (Figure 2D), our method achieved an overall correlation

of 0.33 while Flex ddG showed a correlation of approximately 0. We noticed that a

number of mutations were falsely predicted to have ΔΔG values between -1 and 1 REU

by the MDR-FEP method. Interestingly, correlation improved significantly (0.33 to 0.51)

with the exclusion of these "negligible predicted ΔΔG" mutations whereas Flex ddG did

not show any improvement when applying this same filter. Over a set of 49 mutations to

alanine at the urokinase-urokinase receptor interface (Figure 2E), the proposed method

achieved an overall correlation of 0.21 compared to Flex ddG's 0.41.

We observed that the MDR-FEP method was predicting very negative ΔG values

for the D254A mutation in the monomer, even at low β values. Analysis of this mutation

revealed that less than 90% of the contribution to the exponential averaging fell within

the main distribution, while for all other mutations this was not the case. This analysis

indicates an inappropriately large contribution from low-energy conformations for this

particular mutation (Figure S3) Exclusion of this potential outlier resulted in a correlation

of 0.50 with experiment. The correlation for the proposed method improved to 0.58 with

the exclusion of negligible predicted ΔΔG mutations and the outlier, while the correlation

of Flex ddG did not show such an improvement. Finally, for non-allosteric mutations at

the barnase-barstar interface, both the MDR-FEP method and Flex ddG achieved an

overall correlation to experiment of approximately 0.5 over a set of 28 mutations (Figure

2F). Again, the MDR-FEP method showed an improved correlation with experiment (0.5

to 0.61) with the exclusion of low predicted-value mutations, while Flex ddG did not.

After observing that the optimal β value for all three datasets was approximately

0.002, we sought to determine an optimal consensus cutoff parameter that would

maximize correlation independent of the dataset. We compared the correlation of each

dataset using a β value of 0.002 against cutoff values ranging from 0 to 1% (Figure 2G),

finding that all three datasets demonstrated acceptable correlations with no cutoff

applied. Based on these results, we propose an optimal consensus parameter set of β =

0.002 and cutoff = 0%.

Predictive Performance Depends on Mutation Location

After observing good performance with both allosteric and non-allosteric

mutations, we evaluated the performance of our method based on the location of the

allosteric mutations included in the present study (Figure 3). We hypothesized that ΔG

values for mutations farther from the 3BT1 interface would be more difficult to correctly

predict due to the greater distances over which these mutations exert their influence.

Indeed, we observed that mutations which were correctly predicted to be destabilizing

or stabilizing fell closest to the 3BT1 interface (Figure 3A-C, green and blue). Likewise,

mutations incorrectly predicted to be destabilizing or stabilizing tended to fall farther

from the interface (Figure 3A-C, red and magenta). Interestingly, we observed no clear

trend to explain the performance of negligible predicted ΔΔG mutations based on distance to the interface. Additionally, we evaluated the performance of the MDR-FEP method at various distance cutoffs, where only mutations that fall closer to the interface than the cutoff are included in the calculation (Figure 3D). We found that correlation indeed decreased with the inclusion of more distant mutations.

We hypothesized that the exclusion of negligible predicted ΔΔG mutations improves correlation by removing "false negative" mutations, or mutations which are falsely predicted to have near-zero ΔΔG values. These false negatives likely arise due to a higher energy difference between the WT and mutant proteins, resulting in a smaller likelihood of sampling the low-energy frames needed to recapitulate experimental data. Indeed, we observed that negligible predicted ΔΔG mutations tend to sample fewer low-energy frames than their high predicted-value counterparts (Figure S4). Additionally, we found that 13 of the 36 negligible predicted ΔΔG allosteric mutations feature amino acids with hydroxyl groups (Figure 4A). Mutations which remove or introduce hydrogen bonds to the protein structure may be more difficult for the WT equilibrium simulations to sample, perhaps helping to explain this trend.

MDR-FEP Relies on WT Equilibrium Simulations Sampling MUT-like Conformations

We initially predicted that the MDR-FEP method would detect mutational ΔG values by transiently sampling conformations which are more stable with mutant sidechains, and hence have more "low-RMSD frames" similar to conformations produced by mutant equilibrium simulations. We tested this hypothesis by evaluating the performance of the MDR-FEP method with differing numbers of low-RMSD frames removed, predicting that correlation to experiment would decrease as more low-RMSD

frames are excluded. We indeed observed a strong relationship between correlation

and the removal of low-RMSD frames, with overall correlation to experiment dropping

from ~0.55 to ~0.30 with approximately 30% of the lowest-RMSD frames excluded

(Figure 4A).

Likewise, we expected that for mutations whose predicted values disagreed with

experiment, the WT equilibrium simulations would feature higher RMSDs to the MUT

equilibrium simulations' average conformations. We found that the WT equilibrium

simulation was more likely to have higher RMSDs to the average conformation of

mutations which were falsely predicted to be stabilizing to the 3BT1 dimerization

process (Figure 4B). However, we did not observe this trend in the monomer

simulations, perhaps indicating that both monomer and dimer simulations must sample

low-energy frames to accurately predict ΔG (Figure S5).

Finally, we hypothesized that the WT equilibrium simulations would be more

likely to sample low-energy frames with longer simulation times, thereby increasing

correlation with experiment. We observed a strong relationship between correlation and

total simulation time used (Figure 4C), as well as a direct relationship between total

simulation time and total number of frames with sub-Angstrom RMSDs to mean

structures from mutant equilibrium simulations (Figure 4D). These results indicate that

the MDR-FEP method's ability to predict experimental values depends on the WT

equilibrium simulation's transient sampling of low-energy mutant-like frames, and that

this sampling depends on the total simulation time used. Based on these results, we

believe that the performance of the MDR-FEP method on negligible predicted ΔΔG

mutations, which suffer from insufficient sampling of low-energy frames, may improve

with longer equilibrium simulation times.

The MDR-FEP method selects conformations which are sterically favorable with the mutant but not WT sidechains.

We hypothesized that the MDR-FEP method would be most sensitive to

conformations which favor the mutant through a steric rather than electrostatic effect. In

order to investigate this conformational selection mechanism, we analyzed the

individual Rosetta score terms of sample mutations which were correctly identified to be

destabilizing and stabilizing by the MDR-FEP method (Figure 5). We first analyzed the

H143A mutation (Figure 5A-C), which is correctly predicted by the MDR-FEP method to

allosterically destabilize 3BT1 dimerization ($\Delta\Delta G_{calc}$ = 4.8 REU, $\Delta\Delta G_{exp}$ = 2.8 kJ mol^{-1}).

We found that excluding frames with ΔE < -500 REU resulted in a predicted $\Delta\Delta G$ not in

accordance with experiment (0.7 REU). Analysis of the individual score terms of low-

energy frames revealed high Lennard-Jones repulsion ("fa_rep") between residues with

the WT but not mutant sidechain, suggesting that mutating His143 to alanine in these

particular conformations can relieve much more steric strain in monomeric state,

resulting in destabilization of the dimer. A high correlation (R > 0.99) between the ΔE

and Δfa_rep of individual frames in both the H143A monomer and dimer was observed.

Analysis of the W129A mutation, which was correctly predicted to stabilize

dimerization, revealed similar trends (Figure 5D-F). When frames with ΔE < -500 REU

were included, the MDR-FEP method correctly predicts the W129A mutation to stabilize

dimerization ($\Delta\Delta G_{calc}$ = -2.9 REU, ΔG_{exp} = -1.8 kJ mol^{-1}). Removal of these frames

resulted in an incorrect $\Delta\Delta G$ prediction of 1.7 REU. Score term analysis of these frames

again revealed high steric overlap with the WT but not mutant sidechains present,

indicating that mutation to alanine stabilizes the dimer by relieving more steric strain in the dimer than the monomer. The same strong relationship (R > 0.99) between Δfa_rep and overall ΔE was observed.

For the H143A mutation, we found that there were more frames with a calculated ΔE < -500 REU in the monomer than in the dimer. This higher number of low-energy frames contributes to a lower ΔG value in the monomer than in the dimer, resulting in an overall positive ΔΔG. Similarly, there were more low-energy frames in the dimer for W129A than in the monomer, resulting in a negative ΔΔG. Based on this analysis, we believe that the MDR-FEP method is able to correctly predict the ΔΔG values of these mutations by appropriately reproducing the relative number of mutation-favoring conformations in the monomer and dimer.

Conclusions

Prediction of the effects of allosterically active mutations remains an important goal for advancing the field of protein design and analysis. However, current methods which can accurately predict these long-range effects are limited by prohibitively large computational cost and are therefore not useful for large-scale analysis or protein design. Recent advancements using the Rosetta macromolecular modeling suite have allowed for accurate large-scale prediction of mutational ΔΔGs, however these methods are limited to interface mutations and cannot accurately detect allosteric effects. In the current study, we propose a method to combine the theoretical framework and extensive conformational sampling of alchemical free energy calculations with the high-throughput capabilities of Rosetta. In this approach, conformations from MD simulations are repacked and scored with Rosetta, and the effects of mutation-accommodating

conformations are accounted for using theory based in statistical mechanics to allow for the accurate prediction of allosteric effects using only simulations of the WT system.

We evaluated the MDR-FEP method over three sets of mutations from two separate systems with accurate experimental data available. We found that the method is able to predict the effects of interface mutations with levels of accuracy comparable to current methods. Importantly, the MDR-FEP method was also able to predict the effects of allosterically active mutations with similarly high levels of accuracy. We believe that this method represents a new approach to high-throughput analysis of allosterically active mutations.

Materials and Methods

Molecular Dynamics Simulations

Molecular dynamics simulations were generated *in silico* using the reference PDB structure 3BT1 (chains A and U) for the urokinase-urokinase receptor dimer, or 1BRS for the barnase-barstar dimer. Missing residues in 3BT1 were added using RosettaRemodel(33). All mutant 3BT1 structures were generated using the PyMOL software package(34). For all simulations, each PDB structure was placed in a dodecahedral box with 1.5 nm between the protein and box walls and solvated using the TIP3P water model(35). Joung Na^+ and Cl^- ions were added to the simulation box at physiological concentrations of 150 mM(36). Each system was allowed to energy-minimize for up to 10,000 steps using the Steepest-Descent algorithm in GROMACS. Energy-minimized structures were then equilibrated with 20 ps of NVT, and then 20 ps of NPT, simulation using 1000 kJ mol^{-1} nm^{-2} all-atom position restraints. Each system was then equilibrated with three consecutive 20 ps NPT simulations using all-atom

position restraints of 500, 250, and 125 kJ mol^{-1} nm^{-2}, respectively. Finally, two successive 20 ps NPT equilibrations were run with force constants of 125 kJ mol^{-1} nm^{-2}, first on all backbone atoms and finally on C$_\alpha$ atoms only.

Equilibrium simulations were run at constant temperature and pressure. The C-rescale barostat was used to hold pressure at 1 Bar, and the V-rescale thermostat used to hold temperature at 300 K. All system preparations were carried out using GROMACS functions and in-house scripts. Production simulation trajectories were generated using GROMACS 2021[37] with a 2 fs timestep, the Verlet neighbor-searching cutoff-scheme, and Particle-Mesh Ewald (PME) for van der Waals and electrostatic interactions. The AMBER 99SB*-ILDN[38, 39] force field was used to generate production trajectories, constraining all bonds using the LINCS algorithm[40].

Protocol Implementation and Rosetta Repacking

For each system, three independent 500 ns simulations of both the monomer and dimer were run. Conformations were extracted every 1 ns from these trajectories. Each of these conformational frames was then repacked using the Rosetta 2020 fixed-backbone repacker, allowing for extra sub-rotamers for χ_1 and χ_2 angles. The "multi-cool-annealer" repacking option was used, where multiple "cooling" cycles are run and low temperature rotamer substitutions are then run from the 10 best network states generated during the cooling stage. Additionally, sidechain dihedral angle minimization was used on the lowest energy structure from the packer. For each frame, the score (E) from the lowest energy conformation of 50 repacking iterations was recorded with both the WT and mutant sidechains present. The following flags were used: −ex1 −ex2 −multi_cool_annealer 10 −minimize_sidechains −ndruns 50. Only residues which contained at least 1 atom within 10 Å of the mutation of interest in the crystal structure

were repacked. For each mutation, the same set of residues was repacked in the monomer and dimer forms of the protein.

Following repacking and scoring, ΔE values for each frame were determined using the following equation:

$$\Delta E = E_{\text{MUT}} - E_{\text{WT}}$$

Where ΔE represents the difference in score between the frame with a WT and MUT sidechain present. The set of ΔE values generated for each mutation in the monomer and dimer states were converted into a distribution using kernel density estimation with a bandwidth manually set to 0.1, and a cutoff parameter was applied such that all values below the cutoff are set equal to zero. ΔG values were then generated from each distribution of ΔE values using the following equation:

$$\Delta G = -\frac{1}{\beta} \ln \left(\sum_{i=1}^{N} y_i e^{-\beta x_i} \Bigg/ \sum_{i=1}^{N} y_i \right)$$

Where x and y are the vectors describing the X and Y values of the kernel density function, respectively. Finally, $\Delta\Delta G$ values were generated using the following equation:

$$\Delta\Delta G_{\text{MUT,Dimerization}} = \Delta G_{\text{MUT,Dimer}} - \Delta G_{\text{MUT,Monomer}}$$

All experimental data in Figure 2 and Figure 3 were obtained from the SKEMPI 2.0 database(41) with mutations involving glycine removed, as were the locations of all mutations relative to the protein interaction interface ("interior" and "interface"). Flex ddG values shown in Figure 2 were calculated using the recommended parameters: `nstruct = 35, max_minimization_iter = 5000, abs_score_convergence_thresh = 1.0,` and `number_backrub_trials = 35000.`

Data Analysis and Visualization

The solid lines in the data distributions shown in Figure 4B, Figure S4, and Figure S5 represent the mean value over three independent MD simulations. The shrouds in these distributions represent the standard error of these means, as calculated by

$$SE = \frac{\sigma}{\sqrt{N}}$$

where σ is the standard deviation *N* is the number of independent trials. All protein images were generated using PyMol(34).

Acknowledgements

This work was supported by NSF MCB2144396 (CAS). The authors would also like to thank Wesleyan University for computer time supported by the NSF under grant number CNS-0619508 and CNS-0959856, and the Extreme Science and Engineering Discovery Environment (XSEDE), which is supported by National Science Foundation grant number ACI-1548562 (under the allocation ID MCB190110), for resources on the SDSC Expanse and PSC Bridges compute clusters used to produce this work(42). To the best of our knowledge, no authors have any conflict of interest, financial or otherwise.

Figures

Figure 1. Schematic of the MDR-FEP method.

(A) A schematic overview of the proposed method is shown, with the first step involving Gromacs molecular dynamics simulations, the second step using Rosetta repacking simulations, and the third step involving analysis of the resulting energies. **(B)** Box plots containing all ΔE values for the L113A monomer and dimer are shown, where each frame outside the interquartile range represented by a dot. **(C)** Images of the primary ΔE distributions for the L113A monomer and dimer are shown. ΔG values, calculated using β = 0.002 and cutoff = 0.005%, are shown as dashed lines using all frames (solid colors) and using only frames with ΔE > -500 REU (opaque colors). The final ΔΔG value is the difference between the blue and green dashed lines, represented by a solid arrow for the calculation including all frames and by an opaque arrow for the calculation using only frames with ΔE > -500 REU.

Figure 2. Parameter optimization by grid search.

(A-C) Grid search heat maps showing the correlation of MDR-FEP predicted data to experiment for (A) 3BT1 allosteric, (B) 3BT1 interface, and (C) 1BRS interface sets of mutations. Correlations were calculated after excluding negligible predicted ΔΔG mutations for all datasets. Green circles indicate the set of parameters producing the

highest correlation. **(D-F)** The performance of the MDR-FEP method is compared to Flex ddG(12), a leading Rosetta-based ΔΔG calculation method, for each set of mutations in (A-C) shown above. **(G)** The correlation of each set of mutations in (A-C) is plotted against the data cutoff using $\beta = 0.002$. Correlations were calculated after removal of negligible predicted ΔΔG mutations for all datasets, and after the removal of the outlier indicated by the gray circle in (E) for 3BT1 interface mutations.

Figure 3. Calculated ΔΔG$_{Dimerization}$ values show good agreement with experimental data for allosterically active mutations in the 3BT1 interior.

(A) ΔΔG$_{Dimerization}$ values calculated with the optimal parameter set (β = 0.002, cutoff= 0.05%) are plotted against experimentally determined values for 3BT1 mutations which are defined as "interior" by the SKEMPI database(41), meaning that these residues are located away from the protein-protein interface and have a relative solvent-accessible surface area of less than 25%. Lines of best fit are shown for the comparison of all mutations (gray) and for the set which excludes negligible predicted ΔΔG mutations (| MDR-FEP ΔΔG | ≤ 1 REU, orange). Dots are labeled by the identity of the residue being mutated to alanine (WT alanines are mutated to serine). **(B)** Mutations shown in

Panel A are depicted as spheres on 3BT1 chain U (gray backbone) along with the binding partner chain A (yellow backbone). Values which were correctly identified as stabilizing and destabilizing are shown in blue and green, respectively. Values which are incorrectly predicted to be stabilizing and destabilizing are shown in red and magenta, respectively. Negligible predicted $\Delta\Delta G$ mutations are shown in gray. **(C)** Distributions of the minimum distance between a mutation's C_β and the closest C_β (or C_α in the case of glycine) of the opposite chain for each set of mutations in (A). Colored based on (A-B). **(D)** MDR-FEP correlation with experiment, calculated using the same mutations and parameter set as in (A), is plotted against a distance cutoff where mutations with a minimum distance to the opposite chain according to (C) greater than the cutoff are excluded from the calculation. Colored according to (A).

Figure 4. The MDR-FEP method performs well by sampling mutant-like conformations.

(A) The correlation of predicted values to experimental data for all non-charge-changing interior allosteric 3BT1 mutations is compared to the number of frames with the lowest RMSDs to mutant simulations excluded from the calculation of each ΔG value. The optimal parameter set of β = 0.002 and cutoff = 0.05% is used. Correlations are calculated after removing negligible predicted ΔΔG mutations. **(B)** For each mutation

shown in Figure 3A, the backbone RMSD of each frame of the WT equilibrium 3BT1 dimer simulation to the average structure of that mutation's equilibrium simulation is calculated. An average RMSD is then calculated for each mutation, and the distribution of these average values is shown for each set of mutations colored according to Figure 3A. **(C)** The correlation of predicted values to experiment is compared to total simulation time using the same set of mutations, parameters, and correlation calculation scheme as Figure 3A. The total simulation time is split evenly among three independent simulations, for example three independent 400 ns simulations are run for a total simulation time of 1200 ns. **(D)** The relationship between total simulation time and total number of frames with an RMSD to any mutant's average structure of less than 1 Å is shown.

Figure 5. The MDR-FEP method selects conformations which sterically favor the mutant.

(A-B, D-E) Plots comparing the overall ΔE and Δfa_rep (the Rosetta energy term which describes Lennard-Jones overlap) are shown for sample (A-B) destabilizing and (D-E) stabilizing mutations using the optimal parameter set of β = 0.002 and cutoff = 0.05%. Each frame is represented by a dot. **(C/F)** Protein images of the WT frames indicated by green circles in the (A)/(E) (above) are shown in green. Sample structures from frames with ΔE closer to 0, indicated by the blue circles in (A)/(E), are shown in blue for comparison.

References

1.	Stites, W. E. 1997. Protein−Protein Interactions: Interface Structure, Binding Thermodynamics, and Mutational Analysis. Chemical Reviews 97(5):1233-1250.

2.	Sugiki, T., T. Fujiwara, and C. Kojima. 2014. Latest approaches for efficient protein production in drug discovery. Expert Opinion on Drug Discovery 9(10):1189-1204.

3.	Vidal, M., Michael E. Cusick, and A.-L. Barabási. 2011. Interactome Networks and Human Disease. Cell 144(6):986-998.

4.	Guerois, R., J. E. Nielsen, and L. Serrano. 2002. Predicting Changes in the Stability of Proteins and Protein Complexes: A Study of More Than 1000 Mutations. Journal of Molecular Biology 320(2):369-387.

5.	Tuncbag, N., A. Gursoy, and O. Keskin. 2009. Identification of computational hot spots in protein interfaces: combining solvent accessibility and inter-residue potentials improves the accuracy. Bioinformatics 25(12):1513-1520.

6.	Krï¿½ger, D. M., and H. Gohlke. 2010. DrugScorePPI webserver: fast and accurate in silico alanine scanning for scoring protein–protein interactions. Nucleic Acids Research 38(suppl_2):W480-W486.

7.	Dehouck, Y., J. M. Kwasigroch, M. Rooman, and D. Gilis. 2013. BeAtMuSiC: prediction of changes in protein–protein binding affinity on mutations. Nucleic Acids Research 41(W1):W333-W339.

8.	Moal, I. H., and J. Fernandez-Recio. 2013. Intermolecular Contact Potentials for Protein–Protein Interactions Extracted from Binding Free Energy Changes upon Mutation. Journal of Chemical Theory and Computation 9(8):3715-3727.

9.	Kastritis, P. L., and A. M. J. J. Bonvin. 2010. Are Scoring Functions in Protein−Protein Docking Ready To Predict Interactomes? Clues from a Novel Binding Affinity Benchmark. Journal of Proteome Research 9(5):2216-2225.

10.	Yan, Y., M. Yang, C. G. Ji, and J. Z. H. Zhang. 2017. Interaction Entropy for Computational Alanine Scanning. Journal of Chemical Information and Modeling 57(5):1112-1122.

11.	Moreira, I. S., P. A. Fernandes, and M. J. Ramos. 2007. Computational alanine scanning mutagenesis—An improved methodological approach. Journal of Computational Chemistry 28(3):644-654. https://doi.org/10.1002/jcc.20566.

12. Barlow, K. A., S. Ó Conchúir, S. Thompson, P. Suresh, J. E. Lucas, M. Heinonen, and T. Kortemme. 2018. Flex ddG: Rosetta Ensemble-Based Estimation of Changes in Protein–Protein Binding Affinity upon Mutation. The Journal of Physical Chemistry B 122(21):5389-5399.

13. Kim, D. E., D. Chivian, and D. Baker. 2004. Protein structure prediction and analysis using the Robetta server. Nucleic acids research 32(Web Server issue):W526-W531.

14. Kortemme, T., and D. Baker. 2002. A simple physical model for binding energy hot spots in protein–protein complexes. Proceedings of the National Academy of Sciences 99(22):14116-14121.

15. Havranek, B., K. K. Chan, A. Wu, E. Procko, and S. M. Islam. 2021. Computationally Designed ACE2 Decoy Receptor Binds SARS-CoV-2 Spike (S) Protein with Tight Nanomolar Affinity. Journal of Chemical Information and Modeling 61(9):4656-4669.

16. Manrique-Suárez, V., L. Macaya, M. A. Contreras, N. Parra, R. Maura, A. González, J. R. Toledo, and O. Sánchez. 2021. Design and characterization of a novel dimeric blood–brain barrier penetrating TNFα inhibitor. Proteins: Structure, Function, and Bioinformatics 89(11):1508-1521. https://doi.org/10.1002/prot.26173.

17. Gapsys, V., S. Michielssens, D. Seeliger, and B. L. de Groot. 2015. pmx: Automated protein structure and topology generation for alchemical perturbations. J Comput Chem 36(5):348-354.

18. Gapsys, V., and B. L. de Groot. 2017. Alchemical Free Energy Calculations for Nucleotide Mutations in Protein–DNA Complexes. Journal of Chemical Theory and Computation 13(12):6275-6289.

19. Shirts, M. R., and V. S. Pande. 2005. Comparison of efficiency and bias of free energies computed by exponential averaging, the Bennett acceptance ratio, and thermodynamic integration. The Journal of Chemical Physics 122(14):144107.

20. Zwanzig, R. W. 1954. High-Temperature Equation of State by a Perturbation Method. I. Nonpolar Gases. The Journal of Chemical Physics 22(8):1420-1426.

21. Wood, R. H., W. C. F. Muhlbauer, and P. T. Thompson. 1991. Systematic errors in free energy perturbation calculations due to a finite sample of configuration space: sample-size hysteresis. The Journal of Physical Chemistry 95(17):6670-6675.

22. Zuckerman, D. M., and T. B. Woolf. 2004. Systematic Finite-Sampling Inaccuracy in Free Energy Differences and Other Nonlinear Quantities. Journal of Statistical Physics 114(5):1303-1323.

23. Lu, N., J. K. Singh, and D. A. Kofke. 2003. Appropriate methods to combine forward and reverse free-energy perturbation averages. The Journal of Chemical Physics 118(7):2977-2984.

24. Wells, N. G. M., G. A. Tillinghast, A. L. O'Neil, and C. A. Smith. 2021. Free energy calculations of ALS-causing SOD1 mutants reveal common perturbations to stability and dynamics along the maturation pathway. Protein Science 30(9):1804-1817. https://doi.org/10.1002/pro.4132.

25. Patel, D., J. S. Patel, and F. M. Ytreberg. 2021. Implementing and Assessing an Alchemical Method for Calculating Protein–Protein Binding Free Energy. Journal of Chemical Theory and Computation 17(4):2457-2464.

26. Gapsys, V., A. Yildirim, M. Aldeghi, Y. Khalak, D. van der Spoel, and B. L. de Groot. 2021. Accurate absolute free energies for ligand–protein binding based on non-equilibrium approaches. Communications Chemistry 4(1):61.

27. Fenton, A. W., and G. D. Reinhart. 2002. Isolation of a single activating allosteric interaction in phosphofructokinase from Escherichia coli. Biochemistry 41(45):13410-13416.

28. Alontaga, A. Y., and A. W. Fenton. 2011. Effector analogues detect varied allosteric roles for conserved protein-effector interactions in pyruvate kinase isozymes. Biochemistry 50(11):1934-1939.

29. Tang, Q., and A. W. Fenton. 2017. Whole-protein alanine-scanning mutagenesis of allostery: A large percentage of a protein can contribute to mechanism. Hum Mutat 38(9):1132-1143.

30. Kuhlman, B., G. Dantas, G. C. Ireton, G. Varani, B. L. Stoddard, and D. Baker. 2003. Design of a novel globular protein fold with atomic-level accuracy. Science 302(5649):1364-1368.

31. Bdeir, K., A. Kuo, A. Mazar, B. S. Sachais, W. Xiao, S. Gawlak, S. Harris, A. A. Higazi, and D. B. Cines. 2000. A region in domain II of the urokinase receptor required for urokinase binding. J Biol Chem 275(37):28532-28538.

32. Schreiber, G., and A. R. Fersht. 1995. Energetics of protein-protein interactions: analysis of the barnase-barstar interface by single mutations and double mutant cycles. J Mol Biol 248(2):478-486.

33. Huang, P.-S., Y.-E. A. Ban, F. Richter, I. Andre, R. Vernon, W. R. Schief, and D. Baker. 2011. RosettaRemodel: A Generalized Framework for Flexible Backbone Protein Design. PLoS One 6(8):e24109.

34. The PyMOL Molecular Graphics System, Schrödinger, LLC.

35. Kaminski, G. A., R. A. Friesner, J. Tirado-Rives, and W. L. Jorgensen. 2001. Evaluation and Reparametrization of the OPLS-AA Force Field for Proteins via Comparison with Accurate Quantum Chemical Calculations on Peptides. The Journal of Physical Chemistry B 105(28):6474-6487.

36. Joung, I. S., and T. E. Cheatham. 2008. Determination of Alkali and Halide Monovalent Ion Parameters for Use in Explicitly Solvated Biomolecular Simulations. The Journal of Physical Chemistry B 112(30):9020-9041.

37. Abraham, M. J., T. Murtola, R. Schulz, S. Páll, J. C. Smith, B. Hess, and E. Lindahl. 2015. GROMACS: High performance molecular simulations through multi-level parallelism from laptops to supercomputers. SoftwareX 1-2:19-25.

38. Aliev, A. E., M. Kulke, H. S. Khaneja, V. Chudasama, T. D. Sheppard, and R. M. Lanigan. 2014. Motional timescale predictions by molecular dynamics simulations: Case study using proline and hydroxyproline sidechain dynamics. Proteins: Structure, Function, and Bioinformatics 82(2):195-215.

39. Lindorff-Larsen, K., S. Piana, K. Palmo, P. Maragakis, J. L. Klepeis, R. O. Dror, and D. E. Shaw. 2010. Improved side-chain torsion potentials for the Amber ff99SB protein force field. Proteins 78(8):1950-1958.

40. Hess, B., H. Bekker, H. J. C. Berendsen, and J. G. E. M. Fraaije. 1997. LINCS: A linear constraint solver for molecular simulations. Journal of Computational Chemistry 18(12):1463-1472.

41. Jankauskaite, J., B. Jiménez-García, J. Dapkunas, J. Fernández-Recio, and I. H. Moal. 2019. SKEMPI 2.0: an updated benchmark of changes in protein-protein binding energy, kinetics and thermodynamics upon mutation. Bioinformatics 35(3):462-469.

42. John Towns, T. C., Maytal Dahan, Ian Foster, Kelly Gaither, Andrew Grimshaw, Victor Hazlewood, Scott Lathrop, Dave Lifka, Gregory D. Peterson, Ralph Roskies, J. Ray Scott, Nancy Wilkins-Diehr. 2014. XSEDE: Accelerating Scientific Discovery". Computing in Science & Engineering 16:62-74.

Supporting Figures

Figure S1. High β values cause low-energy frames to dominate the ΔG calculation.

The calculated ΔG values are shown as dot-dashed vertical lines, and the contribution of each portion of the ΔE distribution to the final ΔG value is shown as a dotted line. With the optimal β value of 0.002 (green), low-energy frames contribute appropriately to the overall ΔG value. Increasing β to 0.5 causes low-energy frames to dominate, resulting in inaccurate predicted ΔG values.

Figure S2. 3BT1 and 1BRS simulations sample approximately equivalent numbers of low-energy frames.

Comparison between the number of frames with an MDR-FEP predicted ΔE of less than -100 REU for mutations to 3BT1 and 1BRS in WT equilibrium (A) monomer and (B) dimer simulations.

D254A Monomer

ΔG = 61.5 REU
Contribution to ΔG

Figure S3. The MDR-FEP method calculates an inappropriately large negative ΔG value for the D254A mutation, even at low β values.

The ΔG value is calculated using the optimal parameter set of β = 0.002.0 and cutoff = 0.05. The fraction of the contribution (green dotted line) which falls within 99.9% of the main distribution (solid black line) is calculated to be 0.88.

Figure S4. The WT equilibrium simulation tends to sample fewer low-energy frames for negligible predicted ΔΔG mutations (gray) than mutations which have high predicted values (orange).

Mutation set, parameters used, and coloring scheme are based on Figure 3A/D.

Figure S5. WT equilibrium 3BT1 simulations looked less like mutant simulations for mutations which were incorrectly predicted to be stabilizing.

The mean RMSD of the WT equilibrium 3BT1 simulations to the average structure generated by mutant equilibrium simulation is plotted for the sets of mutations described by the coloring of Figure 3 A-C.