Mutational impact of APOBEC3B and APOBEC3A in a human cell line

Running title: Mutational signatures of APOBEC3B and APOBEC3A

Matthew C. Jarvis[1,2], Michael A. Carpenter[1,2,3], Nuri A. Temiz[2,4], Margaret R. Brown[1,2], Kyle A.

Richards[1,2,5], Prokopios P. Argyris[1,2,3], William L. Brown[1,2], Douglas Yee[2,6], Reuben S.

Harris[1,2,3,5,7,8]

[1] Department of Biochemistry, Molecular Biology and Biophysics, Masonic Cancer Center,

University of Minnesota, Minneapolis, MN 55455, USA

[2] Masonic Cancer Center, University of Minnesota, Minneapolis, MN 55455, USA

[3] Howard Hughes Medical Institute, University of Minnesota, Minneapolis, MN 55455, USA

[4] Institute for Health Informatics, University of Minnesota, Minneapolis, MN 55455, USA

[5] Department of Biochemistry and Structural Biology, University of Texas Health San Antonio,

San Antonio, TX 78229, USA

[6] Department of Medicine, University of Minnesota Medical School, Minneapolis, Minnesota

55455, USA

[7] Howard Hughes Medical Institute, University of Texas Health San Antonio, San Antonio, TX

78229, USA

[8] Correspondence to Reuben S. Harris, email: rsh@uthscsa.edu

Key Words: APOBEC mutation signature; APOBEC3A; APOBEC3B; cancer mutagenesis; DNA

damage response; DNA repair; human mutation reporter system; mutation signatures in cancer

Manuscript Information: 64 character title; 194 word abstract; 5900 word main text (excluding

methods, references, and legends); 5 Main Figures and 8 Supplementary Figures

Abstract

A prominent source of mutation in cancer is single-stranded DNA cytosine deamination by cellular APOBEC3 enzymes, which results in C-to-T and C-to-G mutations in TCA and TCT motifs. Although multiple enzymes have been implicated, reports conflict and it is unclear which enzyme(s) are responsible. Here we develop a selectable system to quantify genome mutation and compare the mutagenic activities of three leading candidates - APOBEC3A, APOBEC3B, and APOBEC3H. The human cell line, HAP1, was engineered to express the *thymidine kinase* (*TK*) gene of HSV-1, which confers sensitivity to ganciclovir. Clonal expression of APOBEC3A and APOBEC3B, but not catalytic mutant controls or APOBEC3H, triggered elevated DNA damage responses and increased frequencies of *TK* mutation. Mutant *TK* DNA sequences revealed nearly indistinguishable cytosine mutation patterns. Whole genome sequences from *TK* mutant clones confirmed these results and enabled broader bioinformatic analyses. Most importantly, comparisons of "pure" APOBEC3A- and APOBEC3B-inflicted mutation signatures from this system and the actual APOBEC3 signature from breast cancer indicated that most tumors manifest a composite signature. These studies help resolve a long-standing etiologic debate in the cancer field and indicate that future diagnostic and therapeutic efforts should focus on both APOBEC3A and APOBEC3B.

Introduction

Over the past decade, advances in DNA sequencing technologies and bioinformatics have helped to deconvolute a multitude of mutational processes that contribute to the genesis and evolution of cancer [see recent pan-cancer analysis by (Alexandrov et al., 2020) and reviews by (Koh, Degasperi, Zou, Momen, & Nik-Zainal, 2021; M. Petljak & Maciejowski, 2020; Venkatesan et al., 2018)]. Through these approaches and complementary bench experiments, the APOBEC3 family of single-stranded (ss)DNA cytosine deaminases has emerged as one of the top three sources of single base substitution (SBS) mutation in cancer with particularly large contributions

to tumors of the bladder, breast, cervix, lung, and head/neck. APOBEC3 signature mutations in

cancer are defined as C-to-T transitions and C-to-G transversions in 5'-TCW motifs (W = A or T;

SBS2 and SBS13, respectively) (Alexandrov et al., 2020; Alexandrov, Nik-Zainal, Wedge,

Aparicio, et al., 2013; Burns, Lackey, et al., 2013; Leonard et al., 2013; Nik-Zainal et al., 2012;

Nik-Zainal et al., 2016; Roberts et al., 2013). This definition is conservative because several

APOBEC3 enzymes can also accommodate 5'-NCG and 5'Nmethyl-CG ssDNA substrates

(Burns, Lackey, et al., 2013; Carpenter et al., 2012; Hultquist et al., 2011; Ito, Fu, Kao, Yang, &

Chen, 2017; Ito et al., 2018; Leonard et al., 2013; Schutsky, Nabel, Davis, DeNizio, & Kohli, 2017;

Shi et al., 2017; Suspène, Aynaud, Vartanian, & Wain-Hobson, 2013), which can also lead to C-

to-T transition mutations and overlap with the mutation signature attributable to spontaneous

water-mediated deamination of cytosine and methyl-cytosine nucleotides. Spontaneous methyl-

cytosine deamination is a clock-like mutation process that occurs predominantly in 5'Nmethyl-CG

motifs and associates positively with a patient's biological age (Alexandrov et al., 2015) (ageing

signature), whereas APOBEC-catalyzed deamination is not evident in normal tissues, not

associated with ageing, present in many primary tumors, and often enriched in metastases

(Alexandrov et al., 2015; Alexandrov et al., 2020; Angus et al., 2019; Burns, Lackey, et al., 2013;

Burns, Temiz, & Harris, 2013; Moore et al., 2021; Nik-Zainal et al., 2016; Roberts et al., 2013;

Sieuwerts et al., 2017; Sieuwerts et al., 2014).

The human APOBEC3 (A3) family is comprised of seven different enzymes with extensive

homology and overlapping activities [reviewed by (Harris & Dudley, 2015; Silvas & Schiffer, 2019;

Simon, Bloch, & Landau, 2015)], and it is unclear how much (or little) each contributes to the

composite APOBEC3 mutation signature evident in tumor DNA sequences. The bulk of evidence

favors two enzymes, APOBEC3A (A3A) and APOBEC3B (A3B), though to wildly different degrees

depending on the study, and additional work has also implicated APOBEC3H (A3H). Evidence

for A3A includes an intrinsic preference for 5'-TC substrates, high catalytic activity (highest of any

human DNA deaminase), cell-wide localization, DNA damage responses, induction by the tumor

virus HPV, and positive correlations between mRNA levels and tumor APOBEC mutation loads

(Carpenter et al., 2012; H. Chen et al., 2006; Cortez et al., 2019; Faden et al., 2017; Green et al.,

2016; Ito et al., 2017; Kouno et al., 2017; Shi et al., 2017; Taylor et al., 2013; Warren et al., 2015;

Westrich et al., 2018). Evidence for A3B includes all the same points except this enzyme is

several-fold less active, localizes constitutively to the nuclear compartment, is induced by multiple

DNA tumor viruses (HPV and polyomaviruses), and associates at both mRNA and protein levels

with clinical outcomes (Argyris et al., 2020; Burns, Lackey, et al., 2013; Burns, Temiz, et al., 2013;

de Bruin et al., 2014; Ito et al., 2017; Roelofs et al., 2020; Salamango et al., 2018; Shi et al., 2017;

Sieuwerts et al., 2017; Sieuwerts et al., 2014; Starrett et al., 2019; Taylor et al., 2013; Verhalen,

Starrett, Harris, & Jiang, 2016; Vieira et al., 2014; Walker et al., 2015; Warren et al., 2015;

Yamazaki et al., 2019; Yamazaki et al., 2020). Importantly, however, the APOBEC3 mutation

signature can still accumulate in tumors completely lacking *A3B* due to a naturally occurring full

gene deletion allele (T. W. Chen et al., 2017; Nik-Zainal et al., 2014; Starrett et al., 2016). This

observation with *A3B*-null tumors helped implicate a cell-wide, 5'-TC preferring variant of A3H

(haplotype I) in causing an APOBEC mutation signature (Feng et al., 2015; Hix, Wong, Flath,

Chelico, & Cisneros, 2020; Starrett et al., 2016). Another consideration is that A3A and A3B have

been reported to elicit broader tetranucleotide preferences upon overexpression in yeast, 5'-

YTCW vs RTCW, respectively (Buisson et al., 2019; Chan et al., 2015; Hoopes et al., 2016).

Here, we report a human cellular system for mutation research and use it to perform

unbiased comparisons of the mutagenic potential of A3A, A3B, and A3H. The human cell line

HAP1 was engineered to express a single copy of the HSV-1 *thymidine kinase* (*TK*) gene, which

enables the drug ganciclovir to be used to select rare *TK* mutants and quantify mutation

frequencies. Moreover, the *TK* gene can be amplified readily from ganciclovir-resistant (Gan[R])

clones by high-fidelity PCR and sequenced to provide initial assessments of mutation spectra

(signatures) prior to undertaking additional experiments such as whole genome sequencing

(WGS). Using this system, only expression of A3A and A3B caused significant increases in DNA

damage signaling (γ-H2AX positivity), DNA breakage (alkaline COMETs), and Gan^R mutation

frequencies. Sanger sequences from panels of individual *TK* mutant clones revealed a clear 5'-

TC-biased mutation pattern including two hotspots, which had no obvious stem-loop secondary

structures. A global perspective was obtained by analyzing the full genomic DNA sequences of

TK mutant clones. These WGS results demonstrated that both A3A and A3B can generate the

APOBEC3 mutational signatures SBS2 and SBS13. Moreover, although these "pure" A3A- and

A3B-inflicted mutation signatures each cluster with the APOBEC3 signature extracted from

different subsets of breast tumors, neither the A3A- nor the A3B-preferred motifs from the HAP1-

TK system are able to fully explain the observed composite APOBEC3 mutation signature in

cancer suggesting that both enzymes may be acting in concert in most tumors.

Results

HAP1-TK-M9 – a human cellular system to report DNA damage and mutagenesis

Model organisms such as *E. coli* and yeast are powerful systems for studying mutagens

including DNA deaminases [*e.g.*, original studies with A3 enzymes (Chan et al., 2012; Harris,

Petersen-Mahrt, & Neuberger, 2002; Mayorov et al., 2005; Petersen-Mahrt, Harris, & Neuberger,

2002; Schumacher, Nissley, & Harris, 2005)]. However, these model organisms do not

recapitulate all of the DNA repair and regulatory mechanisms found in human cells. We therefore

sought to combine strengths of both approaches by introducing a single copy of a selectable

reporter, the HSV-1 *thymidine kinase* (*TK*) gene, into the genome of the human cell line HAP1.

TK confers exquisite sensitivity to the drug ganciclovir and, like many antimicrobial agents, only

TK-mutant, ganciclovir-resistant (Gan^R) cells are able to survive selection by this drug. The vast

majority of Gan^R mutants can be characterized rapidly by conventional Sanger DNA sequencing

because *TK* is a single open reading frame. Moreover, once informative Gan^R mutants are

revealed by selection, secondary analyses including WGS can be used to uncover additional and

potentially global features of a given mutation process.

The overall experimental workflow is shown in **Figure 1A**. To generate a genetically clean

"mother" clone of the commercially available HAP1 cell line, Sleeping Beauty (SB)-mediated

transposition was used to introduce a single copy of a *TK-Neo* cassette into the genome (Burns,

Lackey, et al., 2013). Neo^R clones were selected with G418, expanded into healthy clonal

populations (ca. 10^6 cells/ml), and screened for ganciclovir sensitivity (Gan^S). One mother clone,

HAP1-TK-M9, was selected for further studies because it is Gan^S (apart from rare resistant

mutants) and mostly diploid (apart from some visible chromosome aberrations and S/G2/M cells),

it cultures, engineers, and clones well (below), and it shows a favorable *A3* expression profile

(**Figure 1B**; **Figure 1 – Figure Supplement 1**). In particular, RT-qPCR showed that its *A3A* and

A3B mRNA levels are lower than those of the original parent line and that *A3H* mRNA levels are

very low and near the detection threshold. Accordingly, virtually no ssDNA cytosine deaminase

activity is detected in extracts (vector control lanes in **Figure 1C-E**). In addition, genomic DNA

sequencing showed that the only *A3H* allele is haplotype III (ΔAsn15), which is known to produce

an unstable protein (Chesarino & Emerman, 2020; Ebrahimi et al., 2018; Refsland et al., 2014).

DNA damage phenotypes of HAP1-TK-M9 overexpressing A3A, A3B, and A3H

A panel of MLV-based human A3 expression constructs was assembled and used to

produce viral supernatants and transduce the HAP1-TK-M9 mother clone (**Methods**). Uniform

A3-expressing pools were selected with puromycin. A3A expression was demonstrated by

immunoblotting and immunofluorescent (IF) microscopy using a newly developed rabbit anti-

human A3A monoclonal antibody, UMN13, that specifically binds this enzyme through a unique

N-terminal epitope (**Figure 1C**, **1F**; **Figure 1 – Figure Supplement 2**). Expression of A3B was

similarly shown by immunoblotting and IF microscopy using a rabbit anti-human A3B monoclonal

antibody 5210-87-13 (Brown et al., 2019) (**Figure 1D**, **1F**). Expression of A3H haplotypes I and II

was shown by immunoblotting and IF microscopy with a rabbit polyclonal antibody (Refsland et

al., 2014; Shaban et al., 2018) (**Figure 1E**, **1F**). Endogenous A3A, A3B, and A3H proteins were

undetectable as expected based on low mRNA levels (above). In addition, whole cell lysates of

A3A-, A3B-, and A3H-expressing HAP1-TK-M9 cells elicited ssDNA cytosine deaminase activity,

well above background levels from vector control or catalytic mutant pools constructed in parallel

(**Figure 1C-E**).

We next asked whether these A3 enzymes trigger a DNA damage response and cause

DNA breakage in HAP1-TK-M9 cells. The populations described above were stained for the DNA

damage marker γ-H2AX and subjected to IF microscopy (representative images in **Figure 1F**,

quantification in **Figure 1G**). Only A3A and A3B triggered elevated levels of DNA damage as

characterized by γ-H2AX staining. Catalytic mutant derivatives of A3A and A3B as well as active

A3H-I and A3H-II showed vector control levels of γ-H2AX staining. A3A and A3B also elicited DNA

breaks as evidenced by larger tail moments in alkaline COMET assays (representative images in

Figure 1F, quantification in **Figure 1H**). These results combined to demonstrate that both A3A

and A3B are enzymatically active and capable of exerting DNA damage in HAP1-TK-M9 cells,

consistent with prior DNA damage studies using different cell lines (Akre et al., 2016; Burns,

Lackey, et al., 2013; Landry, Narvaiza, Linfesty, & Weitzman, 2011; Mussil et al., 2013; Nikkila et

al., 2017; Yamazaki et al., 2020). Despite showing cell-wide (including nuclear) localization and

measurable DNA deaminase activity, A3H-I did not trigger significant increases in DNA damage

or breakage. This result differs from an earlier report (Hix et al., 2020), which may have expressed

A3H-I to higher levels with an inducible promoter in comparison to the stable levels established

here with an MND promoter from a single-copy, integrated MLV-based construct.

***TK* mutation spectra of HAP1-TK-M9 with A3A, A3B, and A3H**

To directly test which A3 enzymes cause genomic mutation in the HAP1-TK-M9 system,

24 independent single-cell derived daughter clones were obtained for A3A, A3B, A3H-I, and A3H-

II expressing conditions as well as catalytic mutant controls. A classical fluctuation analysis was

performed by growing each single cell clone to >10^7 cells, subjecting each population to selection

183 by ganciclovir, and allowing time for single Gan^R mutant cells to grow into countable colonies.

184 Vector control conditions yielded a median Gan^R mutation frequency of 3 mutants per 5 million

185 cells (mean = 3.6, SD = 2.8). In contrast, A3A- and A3B-expressing clones caused median Gan^R

186 mutation frequencies to rise above 10 mutants per 5 million cells (A3A WT: median = 16, mean =

187 14, SEM = 2.2, p-value range = 1.7 x 10^{-4} to 3.1 x 10^{-5}; A3B WT: median = 11, mean = 12, SEM

188 = 1.8, p-value range = 2.2 x 10^{-4} to 1.9 x 10^{-5} by Welsh's t-test; **Figure 2A**). In comparison,

189 expression of A3H-I or A3H-II or catalytic mutant derivatives of any of these DNA enzymes failed

190 to trigger increased Gan^R mutation frequencies (A3H-I WT: median = 3.0, mean = 3.6, SEM =

191 0.71, p-value range = 0.38 to 0.60; A3H-II WT: median = 2.0, mean = 2.8, SEM = 0.58, p-value

192 range = 0.74 to 0.89 by Welsh's t-test; **Figure 2A**).

193  We next asked what types of genetic alterations led to inactivation of the *TK* gene in Gan^R

194 granddaughter clones derived from the different A3-expressing and control conditions. The *TK*

195 gene was PCR-amplified from genomic DNA of Gan^R granddaughter clones and sequenced via

196 Sanger sequencing. C-to-T and C-to-G mutations in an APOBEC-signature trinucleotide motif

197 (APOBEC), all other single base substitution mutations (other SBS), and all insertion/deletion

198 mutations (INDEL) were placed into groups for comparison (red, black, and blue tics in **Figure**

199 **2B**, respectively; individual sequence schematics in **Figure 2 – Figure Supplement 1**). *TK*

200 sequences derived from A3A-expressing granddaughter clones harbored a greater number of

201 APOBEC mutations [12/20 clones contained at least 1 APOBEC mutation, 22 T(C>T/G)W

202 mutations total, range of 0 to 3 SBS per sequence] relative to catalytic mutant control clones [3/19

203 clones contained 1 APOBEC mutation, 3 T(C>T/G)W mutations total, range of 0 to 1 SBS per

204 sequence]. Similarly, *TK* sequences derived from A3B-expressing granddaughter clones also

205 harbored a greater number of APOBEC mutations [11/20 clones contained at least 1 APOBEC

206 mutation, 19 T(C>T/G)W mutations total, range of 0 to 3 SBS per sequence] relative to catalytic

207 mutant control clones [2/18 clones contained 1 APOBEC mutation, 2 T(C>T/G)W mutations total,

208 range of 0 to 1 SBS per sequence] (**Figure 2B; Figure 2 – Figure Supplement 1**). Additionally,

TK cytosine bases 22 (Q8X) and 635 (R212K) emerged as mutational hotspots in both the A3A-

and A3B-expressing clones (**Figure 2B**; considered further in **Discussion**). No differences were

found in the number of other SBS or INDEL mutations between A3A- or A3B-expressing clones

and controls.

We next examined the sequence context of the 22 A3A- and 19 A3B-induced APOBEC

signature mutations that occurred at 5'-TC dinucleotides in *TK* (**Figure 2C-D**). In both instances,

A was preferred over T at the +1 nucleobase position relative to the mutated C, although this bias

was not significantly different between the two enzymes (68% for A3A and 74% for A3B; p=0.367

by Fisher's exact test). Similarly, no obvious biases were evident at the +2 or -2 nucleobase

positions with all four nucleotides observed at similar frequencies for both enzymes. Moreover,

even when pyrimidines and purines were grouped for comparison, A3A did not show an overt

preference for C/T (Y) or A/G (R) at the -2 or +2 nucleobase positions (51% vs 49% and 52% vs

48%, respectively). Likewise, A3B also failed to show an overt preference for C/T (Y) or A/G (R)

at the -2 or +2 nucleobase positions (45% vs 55% and 53% vs 47%, respectively). These

similarities underscore the fact that small mutation numbers are primarily useful for delineating

major differences such as the shifts described above from a heterogeneous pattern in catalytic

mutant- or vector control-expressing cells towards a predominantly 5'TC-focused SBS mutation

pattern in A3A- and A3B-expressing cells.

WGS of A3A and A3B clones reveals genome-wide SBS mutation signatures including

biases unique to each enzyme

To prepare for WGS of Gan^R granddaughter clones, RNA sequencing was done to confirm

expression of each exogenously expressed *A3* construct and compare mRNA levels relative to

established breast cell lines and primary breast tumors. All expression values were determined

relative to those of the conserved housekeeping gene *TBP* in order to be able to compare with

RNAseq data from different cell lines and tumors. Interestingly, the average *A3A* and *A3A-E72A*

mRNA levels in Gan^R granddaughter clones was over 5-fold higher than the average endogenous

A3A expression levels of APOBEC3-signature enriched breast cancer cell lines BT474 and MDA-

MD-453, breast cancer cell lines of the CCLE, or breast tumors of TCGA (**Figure 3 – Figure**

Supplement 1A). In other words, A3A levels were high in this system due to difficulty identifying

a sufficiently weak promoter for constitutive expression. In comparison, *A3B* and *A3B-E255A*

mRNA levels in Gan^R granddaughter clones were comparable to the averages reported for breast

cancer cell lines of the CCLE and breast tumors of the TCGA, and approximately 2-fold lower

than those of BT474 and MDA-MD-453 (**Figure 3 – Figure Supplement 1A**). *A3H* (haplotype I)

mRNA levels were higher but only two clones were analyzed by RNAseq and WGS due to

negative results above with the *TK* mutation analysis.

The mRNA expression levels of the other four *A3* genes, as well as *AICDA* (*AID*),

APOBEC1, *APOBEC2*, and *APOBEC4*, were also quantified and compared with those of *A3A*,

A3B, and *A3H* (**Figure 3 – Figure Supplement 1B**). Endogenous *A3C* was expressed at similarly

high levels in all granddaughter clones, providing a robust internal control. Endogenous *A3F* and

A3G were expressed at lower but still detectable levels, and endogenous *AICDA*, *APOBEC1*,

APOBEC2, *APOBEC4*, and *A3D* were expressed at very low or undetectable levels. As expected,

levels of ectopically expressed *A3A*, *A3B*, and *A3H* mRNA exceeded those in the HAP1-TK-M9

parent clone as well as those in vector expressing granddaughter controls. In addition, protein

level expression of A3A, A3B, and A3H was confirmed in *TK* mutant granddaughter clones by

immunoblotting and activity by ssDNA deamination assays (**Figure 3 – Figure Supplement 2**).

Finally, the integration site of the *TK* reporter was determined using WGS reads that spanned

both the 5' and 3'-ends of the integrated *TK* construct and the flanking human genome sequence.

Data from multiple clones demonstrated a single integration site in chromosome 3, which was

~35 kbp from the nearest annotated coding or regulatory element.

To investigate mutational differences genome-wide, Illumina short-read WGS was done

for randomly selected Gan^R granddaughter clones (specified in **Figure 2 – Figure Supplement**

1). Mutations unique to each granddaughter were identified by calling SBS variations versus the

genomic DNA sequence of the HAP1-TK-M9 mother clone. This approach eliminated any somatic

variation that accumulated in the Gan^S mother clone prior to transduction with each of the A3 or

control expression constructs. Thus, all new SBS mutations were present in a significant

proportion of reads from the Gan^R granddaughter clones and absent from the reads from the Gan^S

mother clone and, as such, must have accumulated in the presence of an active A3 enzyme or a

catalytic mutant control.

In A3A-expressing Gan^R granddaughter clones, the total number of unique SBSs ranged

from 2595 to 5821 (n=6, median=3784, mean=4191, SD=1179). The total number of SBSs in

A3B-expressing clones was similar, ranging from 2794 to 2990 (n=5, median=2951, mean=2921,

SD=69). In comparison, the total number of SBSs in catalytic mutant control Gan^R granddaughter

clones ranged from 2094 to 2825 (n=4, mean=2526, SEM=273). Most importantly, analyses of

the trinucleotide contexts of all unique SBS mutations revealed strong C-to-T and C-to-G mutation

biases in 5'-TCA and 5'-TCT motifs in A3A-expressing clones and weaker, but still significant,

mutation biases in the same motifs in A3B-expressing clones (**Figure 3A**; **Figure 3 – Figure**

Supplement 3). In other words, only A3A- and A3B-expressing granddaughter clones elicited

APOBEC3 signature mutations. This important result was confirmed using an independent metric,

APOBEC3 signature enrichment scores (Chan, 2015 #5423;Jarvis, 2018 #6333}, which indicated

that 6/6 A3A-expressing clones and 4/5 A3B-expressing clones had significant enrichments for

APOBEC3 signature mutations, whereas clones expressing catalytically inactive A3A or A3B, as

well as clones expressing A3H-I or vector control had none (**Figure 3 – Figure Supplement 4A**).

An orthologous bioinformatics approach, non-negative matrix factorization [NMF (Alexandrov,

Nik-Zainal, Wedge, Campbell, & Stratton, 2013)], yielded similar results with "signature A"

resembling SBS2 and SBS13 in A3A- and A3B-expressing clones (*i.e.*, APOBEC3 signature) and

"signature B" occurring in all clones regardless of A3 functionality (**Figure 3 – Figure Supplement**

4B). Taken together, these experiments constitute the first cause-and-effect demonstration in a

human cell line that both A3A and A3B are capable of eliciting a clear APOBEC3 mutation

signature.

We next analyzed the broader pentanucleotide contexts of the APOBEC3 signature

mutations that accumulated in A3A- and A3B-expressing clones (n=6520 and n=1590 C-to-T or

C-to-G mutations in 5'-TCW motifs, respectively; **Figure 3B-C**). First, a bias for +1 A over +1 T

emerged in A3A-expressing clones (43.3% A vs 39.6% T), whereas an opposite bias was evident

in A3B-expressing clones (32.5% A vs 43.2% T). Second, no significant bias was noted at the +2

position. Third, an exceptionally strong bias for a pyrimidine nucleobase (C or T) occurred at the

-2 position in A3A-expressing clones (69.1% YTCW vs 30.9% RTCW). These A3A results

resemble the strong -2 pyrimidine bias reported for human A3A mutations in murine hepatocellular

carcinomas (70% YTCW vs 30% RTCW) (Law et al., 2020) and for human A3A expression in

yeast (Chan et al., 2015; Hoopes et al., 2016). Fourth, a weaker bias was apparent for pyrimidines

at the -2 position in A3B-expressing clones (51.6% YTCW vs 48.4% RTCW). This latter result

opposes prior results from yeast, where human A3B appears to cause a bias toward RTCW motifs

[*i.e.*, RTCW > YTCW (Chan et al., 2015; Hoopes et al., 2016)]. Possible explanations for these

contrasting results are mentioned in **Discussion**).

Mesoscale features of A3A and A3B mutagenesis in the HAP1-TK-M9 system

X-ray structures have revealed a U-shaped bend in ssDNA substrates bound by A3A and

A3B (Kouno et al., 2017; Shi et al., 2017), and other studies have indicated that similarly bent

ssDNA loop regions of hairpins (*i.e.*, DNA cruciform structures) may be preferred substrates for

deamination by A3A but not A3B (Buisson et al., 2019; Jalili et al., 2020; Langenbucher et al.,

2021). To ask whether this might be a global feature, we analyzed our A3A and A3B *TK* PCR

sequences and granddaughter clone WGS data for evidence of mutagenesis in the single-

stranded loop regions of DNA hairpin structures. First, neither of the two A3A/B mutation hotspots

in the *TK* gene reported above are part of obvious stem-loop structures. Second, none of the top-

100 cruciform structures reported previously to harbor recurring APOBEC3 signature mutations

in tumors (Buisson et al., 2019) were hotspots in our HAP1 WGS data sets. Third, in global

comparisons of APOBEC3 signature mutations in predicted loop regions of stem-loop structures

versus APOBEC3 signature mutations in non-cruciform structures, neither the A3A- nor the A3B-

expressing conditions showed an increased proportion of TCW mutations in stem-loop regions

relative to control conditions (Fisher's exact test, q = 1 for all comparisons; **Figure 4 – Figure**

Supplement 1).

To assess relative rates of A3A and A3B-catalyzed deamination of hairpin versus non-

hairpin substrates, we used ssDNA substrates representing two previously reported A3A

mutational hotspots in *SDHB* and *NUP93* (Buisson et al., 2019; Sharma et al., 2015). A3A and

A3B were affinity-purified from human cells and incubated with these C-containing hairpin

substrates over time. First, both A3A and A3B showed a strong preference for deaminating the

SDHB hairpin substrate in comparison to a linear control with the same nucleobase content

scrambled (~4- and ~8-fold preference, respectively; **Figure 4A**). In addition, A3A shows higher

rates of deamination than A3B on both the hairpin and the linear substrate in agreement with prior

studies (Buisson et al., 2019). Thus, the relative deamination rates for *SDHB* substrates are:

A3A/hairpin > A3B/hairpin > A3A/linear > A3B linear (120, 41, 31, and 5.1 nM/min, respectively).

However, a different picture emerged from analyses of deamination of *NUP93*-based

substrates **Figure 4B**). Rates of A3A-catalyzed deamination were similarly high for the *NUP93*

hairpin and linear control with the same nucleobase content scrambled. A3B was able to

deaminate the hairpin substrate but at low efficiencies and linear kinetics. In contrast, A3B showed

higher rates of deamination of the linear substrate. Thus, the relative deamination rates for *NUP93*

substrates are: A3A/hairpin = A3A/linear > A3B linear >> A3B/hairpin (93, 83, 7.7, and 1.3 nM/min,

respectively). These results show that reaction rates can vary massively between substrates and

suggest that caution should be used when attempting to extend individual structural and

biochemical preferences to whole genomes. Nevertheless, these data conservatively show that

339 *both* A3A and A3B can deaminate hairpin and linear substrates and, as proposed (Buisson et al.,

340 2019), it is possible to find sites such as the *NUP93* hairpin that are strongly (but not exclusively)

341 preferred by a single A3 enzyme.

342  Another mesoscale feature of APOBEC3 mutagenesis in human cancer is clusters of

343 strand-coordinated cytosine SBS mutations in TCA and TCT motifs most likely caused by

344 processive deamination of exposed tracts of ssDNA (aka. *kataegis*) (Alexandrov, Nik-Zainal,

345 Wedge, Aparicio, et al., 2013; Alexandrov, Nik-Zainal, Wedge, Campbell, et al., 2013; Nik-Zainal

346 et al., 2012; Roberts et al., 2012). No APOBEC3 signature *kataegic* events were observed in

347 control conditions. However, several APOBEC3 signature *kataegic* events were evident in the

348 genomic DNA of both A3A and A3B expressing granddaughter clones (**Figure 4C**). For instance,

349 one A3A-attributable *kataegic* event was comprised of 7 T(C>T/G)A/T events within a 10 kbp

350 window, and an A3B-attributable *kataegic* event included 8 T(C>T/G)A/T events within a 10 kbp

351 window. Interestingly, the frequency of *kataegic* events did not differ significantly between A3A-

352 and A3B-expressing granddaughter clones (A3A WT: median = 30, mean = 35 events, SEM =

353 8.3; A3B WT: median = 12, mean = 15, SEM = 6.9; p = 0.28 by Welch's t-test). These results are

354 the first to indicate in a human cell line that *both* A3A and A3B can cause kataegis.

356 **APOBEC signature etiology in primary breast tumors**

357  Sequencing data from model systems such as HAP1-TK-M9 are powerful because the

358 resulting mutation signatures can be considered relatively pure and, therefore, can help to

359 establish cause-and-effect relationships for comparison to more complex tumor WGS data sets

360 to identify similarities and, potentially, to infer which enzyme(s) generated the observed mutation

361 signature in a given tumor. We therefore performed a Euclidean distance analysis to compare the

362 pentanucleotide cytosine mutation signatures derived from sequencing A3A, A3B, and A3H

363 expressing HAP1 clones and those from 794 primary breast tumors with WGS available through

364 the ICGC data portal resource. This analysis revealed three distinct tumor groups with respect to

APOBEC3 signature mutations: 1) a group that showed similarity to A3A-expressing HAP1-TK-

M9 clones, 2) a group that showed similarity to A3B-expressing HAP1-TK-M9 clones, and 3) a

group that showed little to no significant APOBEC3 mutation signature (**Figure 5**). As expected

from a lack of substantial APOBEC signature mutations above in the *TK* reporter or in

representative WGS, A3H-I expressing clones and catalytic-inactive clones clustered with the

non-APOBEC signature tumors.

Importantly, both the A3A-like and A3B-like groups were comprised of tumors that showed

high levels of APOBEC3 signature mutations and correspondingly high enrichment scores.

However, even in the A3A-like group, none of the breast tumors showed $\geq$70% APOBEC3

signature mutations in YTCW motifs [the overall bias observed here in the HAP1-TK-M9 system

and previously in murine tumors (Law et al., 2020)]. This key result strongly suggests that the

observed APOBEC3 mutation signatures in most breast tumors may be a composite resulting

from both A3A and A3B activity. In comparison, the homologous recombination repair deficiency

signature (SBS3) appeared underrepresented in both the A3A- and A3B-like tumor groups, and

the ageing signature (SBS1) occurred in all 3 groups regardless of A3 mutation status.

Discussion

Here we report the development and implementation of a novel system to investigate

mutational processes in human cells (**Figure 1**). Like many bacterial and yeast model systems,

the HAP1-TK-M9 system enables a uniform cytotoxic selection with ganciclovir such that only *TK*

mutant cells survive. An analysis of clonally derived, A3A and A3B expressing *TK* mutants by

Sanger sequencing of high-fidelity PCR amplicons demonstrates a strong shift in the mutational

pattern from a variety of different base substitution mutations in control conditions to a strongly

5'TCW-biased pattern (**Figure 2**). Interestingly, the *TK* mutation spectra inflicted by A3A and A3B

are virtually indistinguishable including two shared hotspots (Q8X, R212K) and no obvious

mesoscale features. This may be due to the limited number of mutable cytosines in *TK* that confer

resistance to ganciclovir (local base composition) and/or to selective pressure. However, a striking

analogy can be made with the mutational spectrum of the *PIK3CA* gene in breast, head/neck, and

others cancers, which has two prominent APOBEC3 signature hotspots (E542K, E545K) and no

obvious mesoscale structures (Buisson et al., 2019; Henderson, Chakravarthy, Su, Boshoff, &

Fenton, 2014). These observations combine to suggest that selective pressure has the potential

to overshadow the intrinsic preferences of the different APOBEC3 enzymes and complicate

assignment of direct cause-and-effect relationships. Drawing connections between A3A and/or

A3B and a given mutation, even a prominent hotspot, is additionally challenging due to the fact

that both enzymes can deaminate DNA cytosines in both linear substrates as well as single-

stranded loop regions of stem-loop structures [*e.g.*, **Figure 4** and prior biochemical studies

(Buisson et al., 2019; Jalili et al., 2020; Langenbucher et al., 2021)]. Thus, the *TK*-based system

described here is capable of yielding informative, rapid, and inexpensive mutation data sets with

positive results motivating genome-wide analyses where the vast majority of mutations will be

unselected and the larger mutation data sets will enable even stronger conclusions. Earlier work

also leveraged HAP1 cells to characterize a variety of different mutation sources but these studies

did not incorporate a selectable reporter nor did they address the enzyme(s) responsible for

APOBEC3 signature mutations (Zou et al., 2021; Zou et al., 2018).

The studies here are the first to demonstrate unambiguously that both A3A and A3B can

inflict an APOBEC3 mutation signature in human genomic DNA with, in both instances, ssDNA

deamination events immortalizing predominantly as C-to-T and C-to-G mutations in TCA and TCT

trinucleotide motifs (**Figure 3**). Over identical timeframes, A3A causes 4.1-fold more APOBEC3

signature mutations in comparison to A3B (6520 vs 1590 mutations in 6 and 5 subclone WGSs,

respectively). This difference can be explained by higher A3A expression levels in the HAP1-TK-

M9 system (beyond levels in tumor) and the higher intrinsic activity of this enzyme biochemically

in comparison to A3B. Regardless, both A3A and A3B yielded thousands of TC-focused signature

mutations and the most striking difference between these enzymes is the stronger preference of

A3A for a pyrimidine at the -2 position (70.5% YTCW for A3A vs 54.3% YTCW for A3B). This -2

pyrimidine bias mirrors recent WGS results from human A3A-induced murine hepatocellular

carcinomas [70% YTCW (Law et al., 2020)], and it also resembles the bias reported originally for

human A3A expression in yeast (Chan et al., 2015; Hoopes et al., 2016). It also mirrors a strong

-2 pyrimidine bias reported for A3A in the chicken B cell line DT40 [(DeWeerd et al., 2022) that

was published during preparation of this manuscript]. However, importantly, A3B also shows a

YTCW bias here in human cells, which is opposite to the RTCW bias reported in yeast (Chan et

al., 2015; Hoopes et al., 2016). This conflicting result may be due to differences in genetic and/or

epigenetic factors including but not limited to base content, chromatin state, DNA repair

processes, and/or regulatory factors specific to human A3B expression in human cells.

Regardless of the precise explanation(s), the fact that both A3A and A3B can elicit YTCW

mutational biases in human cells helps to inform interpretations of the APOBEC3 enzyme

responsible for the overall APOBEC3 mutation program in cancer. For instance, the observed

APOBEC3 mutation signature in tumors is likely to be a composite of A3A and A3B because very

few (if any) human cancers elicit a 70% YTCW mutation biases indicative of A3A exclusivity or a

52% YTCW bias indicative of A3B exclusivity. In support of this inference, our Euclidean distance

analysis APOBEC3-signature motifs from HAP1-TK-M9 WGSs here and nearly 800 breast tumor

WGSs identified both A3A- and A3B-like subgroups, with neither enzyme's preferred motif fully

explaining the composite signature across breast cancer (**Figure 5**).

In addition to selective pressures and mesoscale features, additional factors are likely to

influence the APOBEC3 signature component of an overall tumor mutational landscape including

whether A3A and/or A3B is expressed, expression level, duration of expression, intrinsic activity,

and accessibility of chromosomal DNA (replication stress, R-loop levels, chromatin state, *etc*.).

With regards to studies here with the HAP1-TK-M9 system, both A3A and A3B are expressed

constitutively from the same construct for identical durations prior to ganciclovir selection, A3A is

intrinsically more active than A3B, A3A is cell-wide and A3B predominantly nuclear, and yet these

enzymes and cellular factors combine to yield remarkably similar *TK* mutation frequencies and

only a 4-fold difference in overall genome-wide SBS mutation level. With respect to cancer, the

A3A gene is expressed at lower levels than *A3B* in almost all cell lines and tumors, A3A is cell-

wide or predominantly cytoplasmic where A3B is constitutively nuclear, and A3A has higher

enzymatic activity that can vary from 2- to 100-fold above that of A3B depending on substrate

[*e.g.*, **Figure 4** and prior biochemical studies (Buisson et al., 2019; Ito et al., 2017)]. It is therefore

remarkable here that the overall genome-wide level of APOBEC3 signature mutation from A3A is

only ~4-fold higher than that attributable to A3B. Endogenous *A3A* and *A3B* also have both distinct

and overlapping transcription programs, and both genes can be induced by a variety of conditions

including viral infection and inflammation (Argyris et al., 2020; Cho et al., 2018; Leonard et al.,

2015; Maruyama et al., 2016; Oh et al., 2021; Periyasamy et al., 2015; Periyasamy et al., 2017;

Roelofs et al., 2020; Starrett et al., 2019; Verhalen et al., 2016; Vieira et al., 2014; Warren,

Westrich, Doorslaer, & Pyeon, 2017; Warren et al., 2015; Westrich et al., 2018). *In vivo*, *A3A* and

A3B gene expression is also likely to be affected by the local tumor microenvironment, which can

vary both between and within cancer types, as well as by a patient's global state of health. Taken

together with unknown and likely lengthy multi-year durations of tumor development prior to

clinical manifestation, deducing the precise fractions of mutations attributable to A3A and/or A3B

may be fruitless.

Independent whole genome sequencing experiments have provided additional information

on the APOBEC3 mutation process. Initial studies induced overexpression of A3B in HEK-293-

derived cell lines, documented the resulting DNA damage responses, and performed WGS to

assess genome-wide associations (Akre et al., 2016; Nikkila et al., 2017). However, an

unambiguous APOBEC3 mutation signature was difficult to extract from these whole genome

sequences due to large numbers of mutations attributable to compromised mismatch repair (Akre

et al., 2016; Nikkila et al., 2017). A more recent study compared *de novo* mutations occurring in

APOBEC3 signature positive cell lines during multiple generations of clonal outgrowth (M. Petljak

et al., 2019). An intriguing finding from this work is that APOBEC3 signature mutations may be

able to occur in an episodic manner, accumulating in some generations and not others, consistent

with evidence discussed above that A3A and A3B expression can be induced by multiple signal

transduction pathways. Episodic mutagenesis, however, is unexpected in cell-based systems in

which stochastic mutagenesis should predominate given defined media and well-controlled

growth conditions. These studies were followed-up more recently by WGS comparisons of

subclones of the same cancer cell lines CRISPR-engineered to lack A3A or A3B (M. Petljak et

al., 2021). Interestingly, disruption of A3A but not A3B caused a near-complete loss of SBS2 and

SBS13 and a strong diminution of YTCA- and YTCN-focused APOBEC3 signature mutations. A

simplistic interpretation of these results is that A3A is the primary enzyme responsible for

APOBEC3 signature mutagenesis in these cancer cell lines. However, as discussed above,

conditions in culture are necessarily controlled and, at best, are likely to only recapitulate a subset

of the more complex local and global conditions of A3 expression, mutagenesis, and tumor

development *in vivo*. Thus, roles for A3B may be underestimated under such experimental

conditions.

A role for A3H, haplotypes I or II, in cancer is disfavored by our results here showing that

these enzymes are incapable of eliciting DNA damage responses or increasing the *TK* mutation

frequency. Two A3H-I expressing *TK* mutant clones were subjected to WGS and no APOBEC3

mutation signature was evident. In addition, no evidence for A3H emerged from sequencing

clonally-derived breast and B cell lymphoma cancer cell lines (M. Petljak et al., 2021). However,

all of these cell-based studies have limitations as discussed above and have yet to fully eliminate

A3H as a source of APOBEC3 signature mutations in cancer. For instance, A3H-I may take more

time to inflict detectable levels of mutation, it may be subject to different transcriptional and post-

transcriptional regulatory processes, and/or it may only be mutagenic in a subset of cancer types

subject to different stresses and different selective pressures.

Ultimately, the studies here show that both A3A and A3B are capable of triggering overt

DNA damage responses and inflicting a robust APOBEC3 mutation signature in the HAP1-TK-

M9 system and, taken together with other work summarized above, support a model in which both

of these enzymes contribute to the composite APOBEC3 mutation signature reported in many

different tumor types. This conclusion is supported by clinical studies implicating A3A and/or A3B

in a variety of different tumor phenotypes including drug resistance/susceptibility, metastasis, and

immune responsiveness (Argyris et al., 2020; Burns, Lackey, et al., 2013; Cannataro et al., 2019;

T. W. Chen et al., 2017; Z. Chen et al., 2019; Cortez et al., 2019; Driscoll et al., 2020; Faden et

al., 2017; Glaser et al., 2018; Law et al., 2016; Middlebrooks et al., 2016; Sieuwerts et al., 2017;

Sieuwerts et al., 2014; Tsuboi et al., 2016). Thus, efforts to diagnose and treat APOBEC3

signature-positive tumors should endeavor to take into account both enzymes, not simply one or

the other. Such longer-term goals are not trivial given the high degree of identity between A3A

and the A3B catalytic domain (>90%), the related difficulty of developing specific and versatile

antibodies for detecting each enzyme, and the fact that each can be regulated differentially by a

wide variety of common factors including virus infection and inflammation. Thus, we are hopeful

that the HAP1-TK-M9 system and A3A-specific rabbit monoclonal antibody described here will

help to expedite the achievement of these goals.

Materials and Methods

Key resources table

Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
Cell line (*homo sapiens,* male*)*	HAP1	Horizon	Cat#: C859	

Cell line (*homo sapiens,* male*)*	HAP1-TK-M9	Original clone, this study		Near diploid clone; expresses HSV-1 *TK* and is Gan^S
Cell line (*homo sapiens,* female*)*	293T	ATCC	Cat#: CRL-3216 RRID: CVCL_0063	
Cell line (*homo sapiens,* male*)*	THP1	ATCC	Cat#: TIB-202 RRID: CVCL_0006	
Antibody	Anti-A3A (rabbit Monoclonal)	This study	UMN13	WB (20 ng/mL)
Antibody	Anti-A3B (rabbit monoclonal)	(Brown et al., 2019)	Cat#:12397 RRID:AB_2721202	WB (1:1,000) IF (1:300)
Antibody	Anti-A3H (rabbit polyclonal)	Novus ARP10	Cat#: NBP1-91682 RRID: AB_2057523	WB (1:1,000) IF (1:300)
Antibody	Anti-γ-H2AX (S139) (rabbit polyclonal)	Millipore Sigma JW301	Cat#: H5912 RRID: AB_310406	IF (1:300)
Antibody	Anti-Tubulin (mouse monoclonal)	Sigma-Aldrich	Cat#: T5168 RRID: AB_477579	WB (1:20,000)

Recombinant DNA reagent	pQCXIP MLV vector	NovoPro	Cat#: V010396#	Parental transduction vector
Recombinant DNA reagent	pQCXIP-MND-A3Ai-IRES-Puro	This study	A3A with intron (pRH9978)	Request by contacting RSH
Recombinant DNA reagent	pQCXIP-MND-A3Ai-E72A-IRES-Puro	This study	A3A-E72A with intron (pRH9979)	Request by contacting RSH
Recombinant DNA reagent	pQCXIP-MND-A3Bi-IRES-Puro	This study	A3B with intron (pRH9980)	Request by contacting RSH
Recombinant DNA reagent	pQCXIP-MND-A3Bi-E255A-IRES-Puro	This study	A3B E255A with intron (pRH9981)	Request by contacting RSH
Recombinant DNA reagent	pQCXIP-MND-eGFP-IRES-Puro	This study	eGFP vector control (pRH9977)	Request by contacting RSH
Recombinant DNA reagent	pQCXIP-MND-A3H-Ii-IRES-Puro	This study	A3H hapI with intron (pRH9984)	Request by contacting RSH
Recombinant DNA reagent	pQCXIP-MND-A3H-IIi-IRES-Puro	This study	A3H hapII with intron (pRH9985)	Request by contacting RSH
Recombinant DNA reagent	pQCXIP-MND-A3H-IIi-E56A-IRES-Puro	This study	A3H hapII E56A with intron (pRH9986)	Request by contacting RSH
Software, algorithm	Burrows-Wheeler Aligner (BWA)		RRID: SCR_010910	

Software, algorithm	Fiji	Fiji	RRID: SCR_002285	
Software, algorithm	GraphPad Prism 6	GraphPad	RRID: SCR_002798	
Software, algorithm	ImageQuant	GE Healthcare	RRID: SCR_014246	
Software, algorithm	Image Studio	LI-COR Biosciences	RRID: SCR_015795	
Software, algorithm	MaxQuant version 1.5.2.8	MaxQuant	RRID: SCR_014485	
Software, algorithm	R for Statistical Computing		RRID: SCR_001905	
Software, algorithm	SpeedSeq		RRID: SCR_000469	
Software, algorithm	VarScan2		RRID: SCR_006849	
Other	BD LSRFortessa Flow Cytometer	BD Biosciences	RRID: SCR_019600	PI stain quantification
Other	LI-COR Odyssey FC	LI-COR	Cat#: 2800	WB imaging

Other	Typhoon FLA 7000	GE Healthcare	Cat#: 29-0044-13	Oligo cleavage assay imaging
Other	MiSeq Sequencing System	Illumina	RRID: SCR_016379	WGS library balancing validation
Other	NovaSeq 6000 Sequencing System	Illumina	RRID: SCR_016387	WGS

Cell lines and culture conditions

All cells lines were cultured at 37°C under 5% CO_2. HAP1 cells and derivatives were grown in IMDM (Invitrogen) supplemented with 10% fetal bovine serum (Sigma), penicillin (100 U/mL), and streptomycin (100 µg/mL). 293T cells were cultured in DMEM (Invitrogen) with 10% fetal bovine serum (Sigma), penicillin (100 U/mL), and streptomycin (100 µg/mL). THP1 cells were cultured in RPMI (ThermoFisher) supplemented with 10% fetal bovine serum (Sigma), penicillin (100 U/mL), and streptomycin (100 µg/mL). TransIT-LT1 (Mirus) was used for all transfections. All parent and clonal lineage cell lines tested negative for mycoplasma using a PCR-based assay (Uphoff & Drexler, 2011). Puromycin (ThermoFisher) and neomycin (ThermoFisher) were used at 1 µg/mL and 50 µg/mL, respectively. Ganciclovir (ThermoFisher) was used at 5 µM to select *TK* mutant clones.

Creating the HAP1-TK-M9 system

The HAP1-TK-M9 system was generated by co-transfecting HAP1 parent cells with a plasmid expressing the Sleeping Beauty transposase and a separate plasmid with *TK-Neo* coding sequences flanked by SB recognition sites (Burns, Lackey, et al., 2013; Stenglein, Burns, Li, Lengyel, & Harris, 2010). Semi-confluent cells in 6 well plates were transfected, treated 24 hrs

later with G418 (1 μg/mL), and subcloned by limiting dilution in 96 well plates to create single cell

derivatives. Single cell clones were then expanded and characterized as described in the main

text. The *A3H* genotype was determined by sequencing exon-specific PCR amplicons (Refsland

et al., 2010; Wang et al., 2019) and further confirmed by WGS (below).

Standard molecular cloning procedures were used to create derivatives of MLV pQCXIP

for expressing each A3 protein. First, pQCXIP was cut with MluI and PacI to excise the strong

CMV promoter and replace it with a weaker MND promoter (a synthetic promoter containing

regions of both the MLV LTR and the myeloproliferative sarcoma virus enhancer). Second, this

new construct was cut with SfiI and BsiWI to insert intron-disrupted A3 coding sequences (Law et

al., 2020). This was done for *A3A*, *A3B*, *A3H* haplotype-I, *A3H* haplotype-II, and appropriate

catalytic mutant derivatives (E-to-A). An eGFP expressing construct was generated in parallel to

use as a control in various experiments. All new constructs were confirmed by Sanger

sequencing. Each MLV-based construct was co-transfected into 293T cells with appropriate

packaging vectors and 48 hrs later the resulting viral supernatants were filtered (0.2 μm) and used

to transduce semi-confluent HAP1-TK-M9 cells. After an additional 48 hrs incubation, transduced

cells were selected with puromycin (1 μg/mL) and subcloned by limiting dilution to obtain A3

expressing daughter clones. These A3 expressing and control daughter clones were expanded

and characterized as described in the main text. Mutation frequencies were determined by plating

5×10^6 cells in 96 well flat bottom plates, treating with 5 μM ganciclovir (ThermoFisher), and after

14 days incubation counting the number of *TK* mutant colonies that survived selection. Single

Gan[R] granddaughter clones were counted using a light microscope and expanded and

characterized as described in the main text including immunoblotting, DNA deaminase activity

assays, *TK* sequencing, and WGS.

Immunoblots (IB)

Cells were treated with trypsin EDTA and collected, washed in 1X PBS, and re-suspended

in 100 µL of reducing sample buffer per one million cells [0.5 M Tris-HCl pH 6.8, 1% 2-

mercaptoethanol, 10% sodium dodecyl sulfate (SDS), 50% glycerol]. Proteins were denatured by

boiling samples for 20 min and resolved by SDS-polyacrylamide gel electrophoresis. Proteins

were transferred to a PVDF-FL membrane (Millipore Sigma) and blocked in 5% milk in 1X PBS.

Primary and secondary antibodies were incubated in blocking buffer, with the addition of 0.2%

SDS for fluorescent antibodies. The primary antibodies used were anti-A3A/B [5210-87-13, IB:

1:1,000, IF: 1:300 (Brown et al., 2019)], anti-A3H [Novus, IB: 1:1,000, IF: 1:300 (Refsland et al.,

2014)], anti-Tubulin (Sigma Aldrich, 1:20,000), and anti-γ-H2AX (S139) (Millipore Sigma JBW301,

1:300). The secondary antibodies used were anti-rabbit HRP-linked (CST 7074, 1:2,000), IRDye

800CW goat anti-mouse (LI-COR Biosciences, 1:10,000), Alexa Fluor 680 goat anti-rabbit

(Molecular Probes, Eugene, OR, USA, 1:10,000), and Alexa Fluor 680 goat anti-mouse

(Molecular Probes, Eugene, OR, USA, 1:10,000). Membranes were imaged using a LI-COR

Odyssey instrument or LI-COR Odyssey-Fc instrument for HRP visualization (LI-COR

Biosciences).

DNA deaminase activity assays

Whole cell extract (WCE) assays: ssDNA deamination activities were measured using

WCE prepared using 100 µL of HED lysis buffer per 1 million cells (25 mM HEPES, 5 mM EDTA,

10% Glycerol, 1 mM DTT, and 1 protease inhibitor tablet). The samples were sonicated 3 times

for 5 sec to ensure complete lysis. A3-containing lysates were incubated at 37°C for 2 hrs with

purified human UNG2 and a single-stranded DNA substrate containing a single TCA trinucleotide

motif and 3' FAM label (5'-ATTATTATTATTCAAATGGATTTATTTATTTATTTATTTATTT-FAM-

3') following established protocols (Burns, Lackey, et al., 2013; Lackey et al., 2012). After this

initial incubation, the reaction was treated with 1 mM NaOH for 10 min at 98°C. The reaction was

run out on a 15% TBE acrylamide urea gel to separate substrate oligo from cleaved product oligo

and imaged on a Typhoon FLA 7000 with ImageQuant TL 8.2.0 software (GE Healthcare).

Recombinant enzyme assays: A3A- and A3B-mycHis were prepared from transfected

293T as reported (Carpenter et al., 2012; McCann et al., 2021; Shaban et al., 2022; Shi et al.,

2017; Shi, Carpenter, Kurahashi, Harris, & Aihara, 2015). 25 nM of each protein was incubated

for the indicated times with 800 nM substrate in HEPES buffer (25 nM HEPES pH 7.4, 50 mM

NaCl, 5 mM imidazole) at 37°C. Reactions were stopped by freezing in liquid nitrogen and then

were heated to 95°C to denature the enzymes. Reactions were then treated with 0.5 U/reaction

Uracil DNA Glycosylase (NEB, USA) for 10 min at 37°C. Sodium hydroxide was added to each

reaction to 100 mM and reactions were heated to 95°C for 5 minutes to cleave the DNA. Products

were separated by 20% TBE-Urea PAGE, imaged on a Typhoon FLA-7000 (GE Healthcare,

USA), and quantified using ImageQuant TL 8.2.0 software (GE Healthcare). Deamination of the

target cytosine was calculated by dividing the total reaction product (and any subsequent off-

target cytosine deamination events) by the total amount of starting substrate.

PI staining and flow cytometry

Propidium iodide staining was used to assess the ploidy of HAP1 clones relative to THP1

as a confirmed diploid control. Cells were trypsinized and suspended in 100 µL of 1X PBS per 1

million cells. Then, 500 µL ice-cold ethanol was added to cell suspensions and incubated at -20°C

for 1 hour to fix the cells. After fixation, cells were pelleted and washed in 1X PBS. Cells were

finally suspended in 500 µL of FxCycle PI stain (Invitrogen) and incubated for 30 minutes at RT

in the dark to stain the cells. Cells were spun down and resuspended in 300 µL of the PI stain

solution and placed in a 96 well round bottom plate for flow cytometry analysis using a BD

LSRFortessa flow cytometer (with high-throughput 96 well adapter system). A minimum of ten

thousand events were acquired for each cell line.

RT-qPCR

Total RNA was extracted using the High Pure RNA isolation kit (Roche). cDNA was

synthesized using SuperScript First-Strand RT (ThermoFisher). Quantification of mRNA was

done using validated primer sets for all human *A3* genes relative to the housekeeping gene *TBP*

(Burns, Lackey, et al., 2013; Leonard et al., 2015; Refsland et al., 2010; Stenglein et al., 2010).

All RT-qPCR reaction were performed using SsoFast SYBR Green mastermix (Bio-Rad) in 384-

well plates on a LightCycler 480 (Roche) following the manufacturer's protocol. Statistical

analyses were done using GraphPad Prism 6 and R.

Immunofluorescent microscopy (IF)

IF was done as described (Lackey, Law, Brown, & Harris, 2013; McNamara, McCann,

Gudipaty, & D'Orso, 2013). Cells were grown in 6 well tissue culture plates at low density prior to

fixation. Cells were fixed with 4% paraformaldehyde in PBS for 15 minutes at RT. Cells were then

permeabilized using PBS containing 0.2% Triton X-100 (Sigma Aldrich), then rinsed with PBS.

Cells were blocked using IF blocking solution and 0.1% Triton X-100 for 1 hr at RT, then incubated

with primary antibody overnight at 4°C. The primary antibodies used were anti-A3A/B (5210-87-

13, 1:300) (Brown et al., 2019), anti-A3H (Novus, 1:300) (Refsland et al., 2014), and anti-γ-H2AX

(JBW301, Millipore Sigma, 1:300). Cells were then washed with PBS and incubated with a fluoro-

conjugated secondary antibody for 2 hrs at RT in the dark. The secondary antibodies used were

IRDye 800CW goat anti-mouse (LI-COR Biosciences, 1:10,000) and Alexa Fluor 680 goat anti-

rabbit (Molecular Probes, Eugene, OR, USA, 1:10,000). Primary and secondary antibodies were

diluted in blocking buffer. Hoescht 33342 (Mirus) was used at a final concentration of 1 μg/mL to

stain nuclei.

COMET assays

HAP1-TK-M9 clones expressing individual A3 enzymes were harvested at 70%

confluency and resuspended in ice cold 1X phosphor-buffered saline (PBS, Ca^{2+} and Mg^{2+} free,

$1.0x10^5$ cells/mL). The CometAssay kit and manufacturer's protocol were used for all alkaline

comet assays (Trevigen). Cells were spread at low density on a glass slide and covered in low-

melt agarose to keep the cells in position for lysis. After cell lysis, slides were washed 2 times with

1X UDG buffer (New England Biolabs), and then incubated with purified human UNG2 in 1X UDG

buffer (4 ng/µL) for 1 hour at 37°C. After incubation, DNA unwinding and electrophoresis were

done according to the manufacturer's alkaline protocol. Comets were analyzed using the

OpenComet plugin for Image J (Gyori, Venkatachalam, Thiagarajan, Hsu, & Clement, 2014) and

statistical analyses were done in GraphPad Prism 6 and R.

TK sequencing

Cells were harvested and genomic DNA was isolated using the Puregene DNA isolation

protocol.   The   *TK*   cassette   was   amplified   from   genomic   DNA   using   5'-

ATCTTGGTGGCGTGAAACTC-3'   and   5'-CTTCCGGTATTGTCTCCTTCC-3'.   PCR   products

were cleaned-up using the GeneJet Gel Extraction Kit (Thermo Scientific) and Sanger sequenced

with 4 different primers to cover the full open reading frame (5'-ATCTTGGTGGCGTGAAACTC-

3',   5'-GGTCATGCTGCCCATAAGGTA-3',   5'-CCGTTCTGGCTCCTCATATC-3',   and   5'-

CTTCCGGTATTGTCTCCTTCC-3'.

Whole genome sequencing (WGS)

Genomic DNA was prepared from cell pellets (1 million cells) using Allprep DNA/RNA mini

kit (Qiagen). Whole genome libraries were sequenced 150x2 bp on a NovaSeq 6000 (Illumina) to

a target read depth of 30X coverage for all granddaughter clones as well as the parental HAP1-

TK-M9 mother clone. Resulting sequences were aligned to the human genome (hg38) using

SpeedSeq (Chiang et al., 2015), which relies on the Burrows-Wheeler Aligner, BWA (version

0.7.17). PCR duplicates were removed using Picard (version 2.18.16). Reads were locally

realigned around InDels using GATK3 (version 3.6.0) tools RealignerTargetCreator to create

intervals, followed by IndelRealigner on the aligned bam files. Single base substitutions and small

InDels were called in each clone relative to the bam file generated from the HAP1-TK-M9 mother

clone using Mutect2 from GATK3 (version 3.6.0). SBSs that passed the internal GATK3 filter with

minimum 4 reads supporting each variant, minimum 20 total reads at each variant site and a

variant allele frequency over 0.05 were used for downstream analysis. SBSs were analyzed in R

(version 4.0.5) using the MutationalPatterns (Blokzijl, Janssen, van Boxtel, & Cuppen, 2018) and

deconstructSigs R packages [version 1.8.0 (Rosenthal, McGranahan, Herrero, Taylor, &

Swanton, 2016)]. All visualizations were generated using the ggplot2 package (version 3.3.5).

COSMIC single base substitution mutation signatures (v3 – May 2019

https://cancer.sanger.ac.uk/cosmic/signatures/SBS/) were obtained from

https://www.synapse.org/#!Synapse:syn11738319. *De-novo* non-negative matrix factorization of

mutational signatures was performed with the "extract_signature" command from the

MutationalPatterns package, with a rank of 2 and 100 iterations. TCW mutation enrichment scores

were calculated as described (Chan et al., 2015; Jarvis, Ebrahimi, Temiz, & Harris, 2018).

Sequence logos of -2 to +2 sequence surrounding C-to-T mutations were created using the

ggseqlogo (version 0.1) package.

APOBEC expression and mutation signature analyses in TCGA and ICGC data sets

TCGA primary breast tumors represented by both RNA-seq and whole exome sequencing

were downloaded from the Firehose GDAC resource through the Broad Institute pipeline

(http://gdac.broadinstitute.org/) for multiple tumor tissue types. APOBEC mutation signatures

were determined as described (Alexandrov, Nik-Zainal, Wedge, Aparicio, et al., 2013; Jarvis et

al., 2018) using the deconstructSigs R package (Rosenthal et al., 2016). APOBEC mutation

enrichment scores were calculated using the hg19 reference genome and published methods

(Chan et al., 2015). Enrichment score significance was assessed using a Fisher exact test with

Benjamini-Hochberg false discovery rate (FDR) correction. All dataset analyses and

visualizations were conducted using R and the ggplot2 package (https://www.R-project.org/).

TK **integration site determination**

To determine the integration site of the single copy *TK-Neo* construct, a TK reference

sequence was provided as an additional chromosome during alignment of the WGS reads to the

reference genome (hg38). Reads that mapped to this region were then categorized as discordant

and realigned using GRIDSS (v2.2) (Cameron et al., 2017) to determine site of integration.

Euclidean distance analysis

All primary breast tumor whole genome sequencing variant information from International

Cancer Genome Consortium (ICGC) was downloaded from the ICGC data portal

(https://dcc.icgc.org). SBSs that were used in these analyses included only C-to-T variants in a

TC dinucleotide, excluding all mutations as CG motifs. SBSs meeting these inclusion criteria from

all clones expressing A3A, A3B, A3A-E72A, A3B-E255A, and A3H-I were pooled per condition

for this analysis. A matrix comprised of the number of mutations within a tetranucleotide across

all samples within a cancer type was generated, and counts were normalized to frequency within

each cancer type. This matrix was converted to a hierarchical distance matrix using the hclust

function in R, which was then plotted as dendrograms. Mutation signatures were calculated using

deconstructSigs as above.

Data and code availability

All alignment files (FASTQ and BAM format) are available through the Sequence Read

Archive under the BioProject accession number PRJXXXXXXXX.

Author contributions

R.S. Harris and M.C. Jarvis conceptualized and designed the overall project. M.C. Jarvis, M.A. Carpenter, M.R. Brown, K. Richards, P.P. Argyris, and W.L. Brown performed experiments. M.C. Jarvis, M.A. Carpenter, N.A. Temiz, M.R. Brown, P.P. Argyris, and W.L. Brown did formal data analysis. R.S. Harris, M.C. Jarvis, and D. Yee contributed to funding acquisition. R.S. Harris and M.C. Jarvis drafted the manuscript, and all authors contributed to manuscript proofing and revision.

Additional contributions

We thank Arad Moghadasi and Sofia Moraes for assistance with flow cytometry and immunofluorescence microscopy, respectively. We also thank the UMN Cytogenetics Core for expert assistance with metaphase chromosome analyses and Scott McIvor for sharing a plasmid construct with the MND promoter. The results presented here are in part based upon data generated by the TCGA Research Network: http://www.cancer.gov/tcga. Additionally, data from the International Cancer Genome Consortium (ICGC) were used in these analyses: https://dcc.icgc.org/.

Additional information

Funding

These studies were supported in part by NCI P01-CA234228 (to RSH). Salary support for MCJ was provided in part by T32-CA009138 and subsequently F31-CA243306. RSH is the Ewing Halsell President's Council Distinguished Chair at University of Texas San Antonio and an Investigator of the Howard Hughes Medical Institute.

Compliance with ethical standards

Conflict of interest: The authors have no conflicts to declare.

References

Akre, M. K., Starrett, G. J., Quist, J. S., Temiz, N. A., Carpenter, M. A., Tutt, A. N., . . . Harris, R.

S. (2016). Mutation processes in 293-based clones overexpressing the DNA cytosine

deaminase apobec3b. *PLoS One, 11*(5), e0155391. doi:10.1371/journal.pone.0155391

Alexandrov, L. B., Jones, P. H., Wedge, D. C., Sale, J. E., Campbell, P. J., Nik-Zainal, S., &

Stratton, M. R. (2015). Clock-like mutational processes in human somatic cells. *Nat Genet,*

47(12), 1402-1407. doi:10.1038/ng.3441

Alexandrov, L. B., Kim, J., Haradhvala, N. J., Huang, M. N., Tian Ng, A. W., Wu, Y., . . .

Consortium, P. (2020). The repertoire of mutational signatures in human cancer. *Nature,*

578(7793), 94-101. doi:10.1038/s41586-020-1943-3

Alexandrov, L. B., Nik-Zainal, S., Wedge, D. C., Aparicio, S. A., Behjati, S., Biankin, A. V., . . .

Stratton, M. R. (2013). Signatures of mutational processes in human cancer. *Nature,*

500(7463), 415-421. doi:10.1038/nature12477 [pii]

Alexandrov, L. B., Nik-Zainal, S., Wedge, D. C., Campbell, P. J., & Stratton, M. R. (2013).

Deciphering signatures of mutational processes operative in human cancer. *Cell Rep,*

3(1), 246-259. doi:10.1016/j.celrep.2012.12.008

Angus, L., Smid, M., Wilting, S. M., van Riet, J., Van Hoeck, A., Nguyen, L., . . . Martens, J. W.

M. (2019). The genomic landscape of metastatic breast cancer highlights changes in

mutation and signature frequencies. *Nat Genet, 51*(10), 1450-1458. doi:10.1038/s41588-

019-0507-7

Argyris, P. P., Wilkinson, P. E., Jarvis, M. C., Magliocca, K. R., Patel, M. R., Vogel, R. I., . . .

Harris, R. S. (2020). Endogenous APOBEC3B overexpression characterizes HPV-positive

and HPV-negative oral epithelial dysplasias and head and neck cancers. *Mod Pathol*.

doi:10.1038/s41379-020-0617-x

Blokzijl, F., Janssen, R., van Boxtel, R., & Cuppen, E. (2018). Mutationalpatterns: Comprehensive

genome-wide analysis of mutational processes. *Genome Med, 10*(1), 33.

doi:10.1186/s13073-018-0539-0

Brown, W. L., Law, E. K., Argyris, P. P., Carpenter, M. A., Levin-Klein, R., Ranum, A. N., . . .

Harris, R. S. (2019). A rabbit monoclonal antibody against the antiviral and cancer

genomic DNA mutating enzyme APOBEC3B. *Antibodies (Basel), 8*(3).

doi:10.3390/antib8030047

Buisson, R., Langenbucher, A., Bowen, D., Kwan, E. E., Benes, C. H., Zou, L., & Lawrence, M.

S. (2019). Passenger hotspot mutations in cancer driven by APOBEC3A and mesoscale

genomic features. *Science, 364*(6447). doi:10.1126/science.aaw2872

Burns, M. B., Lackey, L., Carpenter, M. A., Rathore, A., Land, A. M., Leonard, B., . . . Harris, R.

S. (2013). APOBEC3B is an enzymatic source of mutation in breast cancer. *Nature,*

494(7437), 366-370. doi:10.1038/nature11881 [pii]

Burns, M. B., Temiz, N. A., & Harris, R. S. (2013). Evidence for APOBEC3B mutagenesis in

multiple human cancers. *Nat Genet, 45*(9), 977-983. doi:10.1038/ng.2701 [pii]

Cameron, D. L., Schroder, J., Penington, J. S., Do, H., Molania, R., Dobrovic, A., . . . Papenfuss,

A. T. (2017). Gridss: Sensitive and specific genomic rearrangement detection using

positional de bruijn graph assembly. *Genome Res, 27*(12), 2050-2060.

doi:10.1101/gr.222109.117

Cannataro, V. L., Gaffney, S. G., Sasaki, T., Issaeva, N., Grewal, N. K. S., Grandis, J. R., . . .

Townsend, J. P. (2019). APOBEC-induced mutations and their cancer effect size in head

and neck squamous cell carcinoma. *Oncogene, 38*(18), 3475-3487. doi:10.1038/s41388-

018-0657-6

Carpenter, M. A., Li, M., Rathore, A., Lackey, L., Law, E. K., Land, A. M., . . . Harris, R. S. (2012).

Methylcytosine and normal cytosine deamination by the foreign DNA restriction enzyme

APOBEC3A. *J Biol Chem, 287*(41), 34801-34808. doi:10.1074/jbc.M112.385161 [pii]

Chan, K., Roberts, S. A., Klimczak, L. J., Sterling, J. F., Saini, N., Malc, E. P., . . . Gordenin, D. A.

(2015). An APOBEC3A hypermutation signature is distinguishable from the signature of

background mutagenesis by APOBEC3B in human cancers. *Nat Genet, 47*(9), 1067-1072.

doi:10.1038/ng.3378

Chan, K., Sterling, J. F., Roberts, S. A., Bhagwat, A. S., Resnick, M. A., & Gordenin, D. A. (2012).

Base damage within single-strand DNA underlies in vivo hypermutability induced by a

ubiquitous   environmental   agent.   *PLoS*   *Genet,*   *8*(12),   e1003149.

doi:10.1371/journal.pgen.1003149

Chen, H., Lilley, C. E., Yu, Q., Lee, D. V., Chou, J., Narvaiza, I., . . . Weitzman, M. D. (2006).

APOBEC3A is a potent inhibitor of adeno-associated virus and retrotransposons. *Curr*

Biol, 16(5), 480-485.

Chen, T. W., Lee, C. C., Liu, H., Wu, C. S., Pickering, C. R., Huang, P. J., . . . Chang, Y. S. (2017).

APOBEC3A is an oral cancer prognostic biomarker in Taiwanese carriers of an APOBEC

deletion polymorphism. *Nat Commun, 8*(1), 465. doi:10.1038/s41467-017-00493-9

Chen, Z., Wen, W., Bao, J., Kuhs, K. L., Cai, Q., Long, J., . . . Guo, X. (2019). Integrative genomic

analyses of APOBEC-mutational signature, expression and germline deletion of apobec3

genes, and immunogenicity in multiple cancer types. *BMC Med Genomics, 12*(1), 131.

doi:10.1186/s12920-019-0579-3

Chesarino, N. M., & Emerman, M. (2020). Polymorphisms in human APOBEC3H differentially

regulate ubiquitination and antiviral activity. *Viruses, 12*(4). doi:10.3390/v12040378

Chiang, C., Layer, R. M., Faust, G. G., Lindberg, M. R., Rose, D. B., Garrison, E. P., . . . Hall, I.

M. (2015). Speedseq: Ultra-fast personal genome analysis and interpretation. *Nat*

Methods, 12(10), 966-968. doi:10.1038/nmeth.3505

Cho, R. J., Alexandrov, L. B., den Breems, N. Y., Atanasova, V. S., Farshchian, M., Purdom, E.,

. . . South, A. P. (2018). APOBEC mutation drives early-onset squamous cell carcinomas

in recessive dystrophic epidermolysis bullosa. *Sci Transl Med, 10*(455). doi:10.1126/scitranslmed.aas9668

Cortez, L. M., Brown, A. L., Dennis, M. A., Collins, C. D., Brown, A. J., Mitchell, D., . . . Roberts, S. A. (2019). APOBEC3A is a prominent cytidine deaminase in breast cancer. *PLoS Genet, 15*(12), e1008545. doi:10.1371/journal.pgen.1008545

de Bruin, E. C., McGranahan, N., Mitter, R., Salm, M., Wedge, D. C., Yates, L., . . . Swanton, C. (2014). Spatial and temporal diversity in genomic instability processes defines lung cancer evolution. *Science, 346*(6206), 251-256. doi:10.1126/science.1253462

DeWeerd, R. A., Nemeth, E., Poti, A., Petryk, N., Chen, C. L., Hyrien, O., . . . Green, A. M. (2022). Prospectively defined patterns of APOBEC3A mutagenesis are prevalent in human cancers. *Cell Rep, 38*(12), 110555. doi:10.1016/j.celrep.2022.110555

Driscoll, C. B., Schuelke, M. R., Kottke, T., Thompson, J. M., Wongthida, P., Tonne, J. M., . . . Vile, R. G. (2020). APOBEC3B-mediated corruption of the tumor cell immunopeptidome induces heteroclitic neoepitopes for cancer immunotherapy. *Nat Commun, 11*(1), 790. doi:10.1038/s41467-020-14568-7

Ebrahimi, D., Richards, C. M., Carpenter, M. A., Wang, J., Ikeda, T., Becker, J. T., . . . Harris, R. S. (2018). Genetic and mechanistic basis for APOBEC3H alternative splicing, retrovirus restriction, and counteraction by HIV-1 protease. *Nat Commun, 9*(1), 4137. doi:10.1038/s41467-018-06594-3

Faden, D. L., Thomas, S., Cantalupo, P. G., Agrawal, N., Myers, J., & DeRisi, J. (2017). Multi-modality analysis supports APOBEC as a major source of mutations in head and neck squamous cell carcinoma. *Oral Oncol, 74*, 8-14. doi:10.1016/j.oraloncology.2017.09.002

Feng, Y., Love, R. P., Ara, A., Baig, T. T., Adolph, M. B., & Chelico, L. (2015). Natural polymorphisms and oligomerization of human APOBEC3H contribute to single-stranded DNA scanning ability. *J Biol Chem, 290*(45), 27188-27203. doi:10.1074/jbc.M115.666065

Glaser, A. P., Fantini, D., Wang, Y., Yu, Y., Rimar, K. J., Podojil, J. R., . . . Meeks, J. J. (2018).

APOBEC-mediated mutagenesis in urothelial carcinoma is associated with improved

survival, mutations in DNA damage response genes, and immune response. *Oncotarget,*

9(4), 4537-4548. doi:10.18632/oncotarget.23344

Green, A. M., Landry, S., Budagyan, K., Avgousti, D. C., Shalhout, S., Bhagwat, A. S., &

Weitzman, M. D. (2016). APOBEC3A damages the cellular genome during DNA

replication. *Cell Cycle, 15*(7), 998-1008. doi:10.1080/15384101.2016.1152426

Gyori, B. M., Venkatachalam, G., Thiagarajan, P. S., Hsu, D., & Clement, M. V. (2014).

OPENCOMET: An automated tool for comet assay image analysis. *Redox Biol, 2*, 457-

465. doi:10.1016/j.redox.2013.12.020

Harris, R. S., & Dudley, J. P. (2015). APOBECs and virus restriction. *Virology, 479-480C*, 131-

145. doi:10.1016/j.virol.2015.03.012

Harris, R. S., Petersen-Mahrt, S. K., & Neuberger, M. S. (2002). RNA editing enzyme apobec1

and some of its homologs can act as DNA mutators. *Mol Cell, 10*(5), 1247-1253.

Henderson, S., Chakravarthy, A., Su, X., Boshoff, C., & Fenton, T. R. (2014). Apobec-mediated

cytosine deamination links PIK3CA helical domain mutations to human papillomavirus-

driven tumor development. *Cell Rep, 7*(6), 1833-1841. doi:10.1016/j.celrep.2014.05.012

Hix, M. A., Wong, L., Flath, B., Chelico, L., & Cisneros, G. A. (2020). Single-nucleotide

polymorphism of the DNA cytosine deaminase APOBEC3H haplotype I leads to enzyme

destabilization and correlates with lung cancer. *NAR Cancer, 2*(3), zcaa023.

doi:10.1093/narcan/zcaa023

Hoopes, J. I., Cortez, L. M., Mertz, T. M., Malc, E. P., Mieczkowski, P. A., & Roberts, S. A. (2016).

APOBEC3A and APOBEC3B preferentially deaminate the lagging strand template during

DNA replication. *Cell Rep, 14*(6), 1273-1282. doi:10.1016/j.celrep.2016.01.021

Hultquist, J. F., Lengyel, J. A., Refsland, E. W., LaRue, R. S., Lackey, L., Brown, W. L., & Harris,

R. S. (2011). Human and rhesus APOBEC3D, APOBEC3F, APOBEC3G, and

APOBEC3H demonstrate a conserved capacity to restrict Vif-deficient HIV-1. *J Virol,*

85(21), 11220-11234.

Ito, F., Fu, Y., Kao, S. A., Yang, H., & Chen, X. S. (2017). Family-wide comparative analysis of

cytidine and methylcytidine deamination by eleven human APOBEC proteins. *J Mol Biol,*

429(12), 1787-1799. doi:10.1016/j.jmb.2017.04.021

Ito, F., Yang, H., Xiao, X., Li, S. X., Wolfe, A., Zirkle, B., . . . Chen, X. S. (2018). Understanding

the structure, multimerization, subcellular localization and MC selectivity of a genomic

mutator and anti-HIV factor APOBEC3H. *Sci Rep, 8*(1), 3763. doi:10.1038/s41598-018-

21955-0

Jalili, P., Bowen, D., Langenbucher, A., Park, S., Aguirre, K., Corcoran, R. B., . . . Buisson, R.

(2020). Quantification of ongoing APOBEC3A activity in tumor cells by monitoring rna

editing at hotspots. *Nat Commun, 11*(1), 2971. doi:10.1038/s41467-020-16802-8

Jarvis, M. C., Ebrahimi, D., Temiz, N. A., & Harris, R. S. (2018). Mutation signatures including

APOBEC in cancer cell lines. *JNCI Cancer Spectr, 2*(1). doi:10.1093/jncics/pky002

Koh, G., Degasperi, A., Zou, X., Momen, S., & Nik-Zainal, S. (2021). Mutational signatures:

Emerging concepts, caveats and clinical applications. *Nat Rev Cancer, 21*(10), 619-637.

doi:10.1038/s41568-021-00377-7

Kouno, T., Silvas, T. V., Hilbert, B. J., Shandilya, S. M. D., Bohn, M. F., Kelch, B. A., . . . Schiffer,

C. A. (2017). Crystal structure of APOBEC3A bound to single-stranded DNA reveals

structural basis for cytidine deamination and specificity. *Nat Commun, 8*, 15024.

doi:10.1038/ncomms15024

Lackey, L., Demorest, Z. L., Land, A. M., Hultquist, J. F., Brown, W. L., & Harris, R. S. (2012).

APOBEC3B and AID have similar nuclear import mechanisms. *J Mol Biol, 419*(5), 301-

314. doi:10.1016/j.jmb.2012.03.011

Lackey, L., Law, E. K., Brown, W. L., & Harris, R. S. (2013). Subcellular localization of the apobec3

proteins during mitosis and implications for genomic DNA deamination. *Cell Cycle, 12*(5),

762-772. doi:10.4161/cc.23713 [pii]

Landry, S., Narvaiza, I., Linfesty, D. C., & Weitzman, M. D. (2011). APOBEC3A can activate the

DNA damage response and cause cell-cycle arrest. *EMBO Rep, 12*(5), 444-450.

Langenbucher, A., Bowen, D., Sakhtemani, R., Bournique, E., Wise, J. F., Zou, L., . . . Lawrence,

M. S. (2021). An extended APOBEC3A mutation signature in cancer. *Nat Commun, 12*(1),

1602. doi:10.1038/s41467-021-21891-0

Law, E. K., Levin-Klein, R., Jarvis, M. C., Kim, H., Argyris, P. P., Carpenter, M. A., . . . Harris, R.

S. (2020). APOBEC3A catalyzes mutation and drives carcinogenesis *in vivo*. *J Exp Med,*

217(12). doi:10.1084/jem.20200261

Law, E. K., Sieuwerts, A. M., LaPara, K., Leonard, B., Starrett, G. J., Molan, A. M., . . . Harris, R.

S. (2016). The DNA cytosine deaminase APOBEC3B promotes tamoxifen resistance in

ER-positive breast cancer. *Sci Adv, 2*(10), e1601737. doi:10.1126/sciadv.1601737

Leonard, B., Hart, S. N., Burns, M. B., Carpenter, M. A., Temiz, N. A., Rathore, A., . . . Harris, R.

S. (2013). APOBEC3B upregulation and genomic mutation patterns in serous ovarian

carcinoma. *Cancer Res, 73*(24), 7222-7231. doi:10.1158/0008-5472.CAN-13-1753

Leonard, B., McCann, J. L., Starrett, G. J., Kosyakovsky, L., Luengas, E. M., Molan, A. M., . . .

Harris, R. S. (2015). The PKC/NF-kappaB signaling pathway induces APOBEC3B

expression in multiple human cancers. *Cancer Res, 75*(21), 4538-4547.

doi:10.1158/0008-5472.CAN-15-2171-T

Maruyama, W., Shirakawa, K., Matsui, H., Matsumoto, T., Yamazaki, H., Sarca, A. D., . . . Takaori-

Kondo, A. (2016). Classical NF-kappaB pathway is responsible for APOBEC3B

expression in cancer cells. *Biochem Biophys Res Commun, 478*(3), 1466-1471.

doi:10.1016/j.bbrc.2016.08.148

Mayorov, V. I., Rogozin, I. B., Adkison, L. R., Frahm, C., Kunkel, T. A., & Pavlov, Y. I. (2005). Expression of human aid in yeast induces mutations in context similar to the context of somatic hypermutation at G-C pairs in immunoglobulin genes. *BMC Immunol, 6*(1), 10.

McCann, J. L., Cristini, A., Law, E. K., Lee, S. Y., Tellier, M., Carpenter, M. A., . . . Harris, R. S. (2021). R-loop homeostasis and cancer mutagenesis promoted by the DNA cytosine deaminase APOBEC3B. *BioRxiv*.

McNamara, R. P., McCann, J. L., Gudipaty, S. A., & D'Orso, I. (2013). Transcription factors mediate the enzymatic disassembly of promoter-bound 7SK SNRNP to locally recruit p-tefb for transcription elongation. *Cell Rep, 5*(5), 1256-1268. doi:10.1016/j.celrep.2013.11.003

Middlebrooks, C. D., Banday, A. R., Matsuda, K., Udquim, K. I., Onabajo, O. O., Paquin, A., . . . Prokunina-Olsson, L. (2016). Association of germline variants in the APOBEC3 region with cancer risk and enrichment with apobec-signature mutations in tumors. *Nat Genet, 48*(11), 1330-1338. doi:10.1038/ng.3670

Moore, L., Cagan, A., Coorens, T. H. H., Neville, M. D. C., Sanghvi, R., Sanders, M. A., . . . Rahbari, R. (2021). The mutational landscape of human somatic and germline cells. *Nature, 597*(7876), 381-386. doi:10.1038/s41586-021-03822-7

Mussil, B., Suspene, R., Aynaud, M. M., Gauvrit, A., Vartanian, J. P., & Wain-Hobson, S. (2013). Human APOBEC3A isoforms translocate to the nucleus and induce DNA double strand breaks leading to cell stress and death. *PLoS One, 8*(8), e73641. doi:10.1371/journal.pone.0073641

Nik-Zainal, S., Alexandrov, L. B., Wedge, D. C., Van Loo, P., Greenman, C. D., Raine, K., . . . Stratton, M. R. (2012). Mutational processes molding the genomes of 21 breast cancers. *Cell, 149*(5), 979-993. doi:S0092-8674(12)00528-4 [pii]

Nik-Zainal, S., Davies, H., Staaf, J., Ramakrishna, M., Glodzik, D., Zou, X., . . . Stratton, M. R.

(2016). Landscape of somatic mutations in 560 breast cancer whole-genome sequences.

Nature, 534(7605), 47-54. doi:10.1038/nature17676

Nik-Zainal, S., Wedge, D. C., Alexandrov, L. B., Petljak, M., Butler, A. P., Bolli, N., . . . Stratton,

M. R. (2014). Association of a germline copy number polymorphism of APOBEC3A and

APOBEC3B with burden of putative APOBEC-dependent mutations in breast cancer. *Nat*

Genet, 46(5), 487-491. doi:10.1038/ng.2955

Nikkila, J., Kumar, R., Campbell, J., Brandsma, I., Pemberton, H. N., Wallberg, F., . . . Lord, C. J.

(2017). Elevated APOBEC3B expression drives a kataegic-like mutation signature and

replication stress-related therapeutic vulnerabilities in p53-defective cells. *Br J Cancer,*

117(1), 113-123. doi:10.1038/bjc.2017.133

Oh, S., Bournique, E., Bowen, D., Jalili, P., Sanchez, A., Ward, I., . . . Buisson, R. (2021).

Genotoxic stress and viral infection induce transient expression of APOBEC3A and pro-

inflammatory genes through two distinct pathways. *Nat Commun, 12*(1), 4917.

doi:10.1038/s41467-021-25203-4

Periyasamy, M., Patel, H., Lai, C. F., Nguyen, V. T., Nevedomskaya, E., Harrod, A., . . . Ali, S.

(2015). APOBEC3B-mediated cytidine deamination is required for estrogen receptor

action in breast cancer. *Cell Rep, 13*(1), 108-121. doi:10.1016/j.celrep.2015.08.066

Periyasamy, M., Singh, A. K., Gemma, C., Kranjec, C., Farzan, R., Leach, D. A., . . . Ali, S. (2017).

P53 controls expression of the DNA deaminase APOBEC3B to limit its potential mutagenic

activity in cancer cells. *Nucleic Acids Res, 45*(19), 11056-11069. doi:10.1093/nar/gkx721

Petersen-Mahrt, S. K., Harris, R. S., & Neuberger, M. S. (2002). AID mutates *E. coli* suggesting

a DNA deamination mechanism for antibody diversification. *Nature, 418*(6893), 99-103.

Petljak, M., Alexandrov, L. B., Brammeld, J. S., Price, S., Wedge, D. C., Grossmann, S., . . .

Stratton, M. R. (2019). Characterizing mutational signatures in human cancer cell lines

reveals episodic APOBEC mutagenesis. *Cell, 176*(6), 1282-1294 e1220. doi:10.1016/j.cell.2019.02.012

Petljak, M., Chu, K., Dananberg, A., Bergstrom, E. N., von Morgen, P., Alexandrov, L. B., . . . Maciejowski, J. (2021). The APOBEC3A deaminase drives episodic mutagenesis in cancer cells. *BioRxiv,* doi: https://doi.org/10.1101/2021.02.14.431145.

Petljak, M., & Maciejowski, J. (2020). Molecular origins of APOBEC-associated mutations in cancer. *DNA Repair (Amst), 94*, 102905. doi:10.1016/j.dnarep.2020.102905

Refsland, E. W., Hultquist, J. F., Luengas, E. M., Ikeda, T., Shaban, N. M., Law, E. K., . . . Harris, R. S. (2014). Natural polymorphisms in human APOBEC3H and HIV-1 Vif combine in primary T lymphocytes to affect viral G-to-A mutation levels and infectivity. *PLoS Genet, 10*(11), e1004761. doi:10.1371/journal.pgen.1004761

Refsland, E. W., Stenglein, M. D., Shindo, K., Albin, J. S., Brown, W. L., & Harris, R. S. (2010). Quantitative profiling of the full *APOBEC3* mRNA repertoire in lymphocytes and tissues: Implications for hiv-1 restriction. *Nucleic Acids Res, 38*, 4274-4284.

Roberts, S. A., Lawrence, M. S., Klimczak, L. J., Grimm, S. A., Fargo, D., Stojanov, P., . . . Gordenin, D. A. (2013). An APOBEC cytidine deaminase mutagenesis pattern is widespread in human cancers. *Nat Genet, 45*(9), 970-976. doi:10.1038/ng.2702

Roberts, S. A., Sterling, J., Thompson, C., Harris, S., Mav, D., Shah, R., . . . Gordenin, D. A. (2012). Clustered mutations in yeast and in human cancers can arise from damaged long single-strand DNA regions. *Mol Cell, 46*(4), 424-435. doi:10.1016/j.molcel.2012.03.030

Roelofs, P. A., Goh, C. Y., Chua, B. H., Jarvis, M. C., Stewart, T. A., McCann, J. L., . . . Harris, R. S. (2020). Characterization of the mechanism by which the RB/E2F pathway controls expression of the cancer genomic DNA deaminase APOBEC3B. *Elife, 9.* doi:10.7554/eLife.61287

Rosenthal, R., McGranahan, N., Herrero, J., Taylor, B. S., & Swanton, C. (2016). Deconstructsigs: Delineating mutational processes in single tumors distinguishes DNA repair deficiencies

and patterns of carcinoma evolution. *Genome Biol, 17*, 31. doi:10.1186/s13059-016-0893-

4

Salamango, D. J., McCann, J. L., Demir, O., Brown, W. L., Amaro, R. E., & Harris, R. S. (2018).

APOBEC3B nuclear localization requires two distinct N-terminal domain surfaces. *J Mol*

Biol, 430(17), 2695-2708. doi:10.1016/j.jmb.2018.04.044

Schumacher, A. J., Nissley, D. V., & Harris, R. S. (2005). APOBEC3G hypermutates genomic

DNA and inhibits TY1 retrotransposition in yeast. *Proc Natl Acad Sci U S A, 102*(28), 9854-

9859.

Schutsky, E. K., Nabel, C. S., Davis, A. K. F., DeNizio, J. E., & Kohli, R. M. (2017). APOBEC3A

efficiently deaminates methylated, but not tet-oxidized, cytosine bases in DNA. *Nucleic*

Acids Res, 45(13), 7655-7665. doi:10.1093/nar/gkx345

Shaban, N. M., Shi, K., Lauer, K. V., Carpenter, M. A., Richards, C. M., Salamango, D., . . . Harris,

R. S. (2018). The antiviral and cancer genomic DNA deaminase APOBEC3H is regulated

by an RNA-mediated dimerization mechanism. *Mol Cell, 69*(1), 75-86 e79.

doi:10.1016/j.molcel.2017.12.010

Shaban, N. M., Yan, R., Shi, K., Moraes, S. N., Cheng, A. Z., Carpenter, M. A., . . . Harris, R. S.

(2022). Cryo-em structure of the EBV ribonucleotide reductase BORF2 and mechanism

of APOBEC3B inhibition. *Science Advances, In press*.

Sharma, S., Patnaik, S. K., Taggart, R. T., Kannisto, E. D., Enriquez, S. M., Gollnick, P., & Baysal,

B. E. (2015). APOBEC3A cytidine deaminase induces RNA editing in monocytes and

macrophages. *Nat Commun, 6*, 6881. doi:10.1038/ncomms7881

Shi, K., Carpenter, M. A., Banerjee, S., Shaban, N. M., Kurahashi, K., Salamango, D. J., . . .

Aihara, H. (2017). Structural basis for targeted DNA cytosine deamination and

mutagenesis by APOBEC3A and APOBEC3B. *Nat Struct Mol Biol, 24*(2), 131-139.

doi:10.1038/nsmb.3344

Shi, K., Carpenter, M. A., Kurahashi, K., Harris, R. S., & Aihara, H. (2015). Crystal structure of

the DNA deaminase APOBEC3B catalytic domain. *J Biol Chem, 290*(47), 28120-28130.

doi:10.1074/jbc.M115.679951

Sieuwerts, A. M., Schrijver, W. A., Dalm, S. U., de Weerd, V., Moelans, C. B., Ter Hoeve, N., . . .

van Deurzen, C. H. (2017). Progressive APOBEC3B mRNA expression in distant breast

cancer metastases. *PLoS One, 12*(1), e0171343. doi:10.1371/journal.pone.0171343

Sieuwerts, A. M., Willis, S., Burns, M. B., Look, M. P., Meijer-Van Gelder, M. E., Schlicker, A., . .

. Martens, J. W. (2014). Elevated APOBEC3B correlates with poor outcomes for estrogen-

receptor-positive breast cancers. *Horm Cancer, 5*(6), 405-413. doi:10.1007/s12672-014-

0196-8

Silvas, T. V., & Schiffer, C. A. (2019). APOBEC3s: DNA-editing human cytidine deaminases.

Protein Sci, 28(9), 1552-1566. doi:10.1002/pro.3670

Simon, V., Bloch, N., & Landau, N. R. (2015). Intrinsic host restrictions to HIV-1 and mechanisms

of viral escape. *Nat Immunol, 16*(6), 546-553. doi:10.1038/ni.3156

Starrett, G. J., Luengas, E. M., McCann, J. L., Ebrahimi, D., Temiz, N. A., Love, R. P., . . . Harris,

R. S. (2016). The DNA cytosine deaminase APOBEC3H haplotype I likely contributes to

breast and lung cancer mutagenesis. *Nat Commun, 7*, 12918. doi:10.1038/ncomms12918

Starrett, G. J., Serebrenik, A. A., Roelofs, P. A., McCann, J. L., Verhalen, B., Jarvis, M. C., . . .

Harris, R. S. (2019). Polyomavirus T antigen induces APOBEC3B expression using an

LXCXE-dependent and TP53-independent mechanism. *MBio, 10*(1).

doi:10.1128/mBio.02690-18

Stenglein, M. D., Burns, M. B., Li, M., Lengyel, J., & Harris, R. S. (2010). APOBEC3 proteins

mediate the clearance of foreign DNA from human cells. *Nat Struct Mol Biol, 17*(2), 222-

229.

Suspène, R., Aynaud, M. M., Vartanian, J. P., & Wain-Hobson, S. (2013). Efficient deamination

of 5-methylcytidine and 5-substituted cytidine residues in DNA by human APOBEC3A

cytidine deaminase. *PLoS One, 8*(6), e63461. doi:10.1371/journal.pone.0063461 [pii]

Taylor, B. J., Nik-Zainal, S., Wu, Y. L., Stebbings, L. A., Raine, K., Campbell, P. J., . . . Neuberger,

M. S. (2013). DNA deaminases induce break-associated mutation showers with

implication of APOBEC3B and 3A in breast cancer kataegis. *Elife, 2*, e00534.

doi:10.7554/eLife.00534 [pii]

Tsuboi, M., Yamane, A., Horiguchi, J., Yokobori, T., Kawabata-Iwakawa, R., Yoshiyama, S., . . .

Nishiyama, M. (2016). APOBEC3B high expression status is associated with aggressive

phenotype in japanese breast cancers. *Breast Cancer, 23*(5), 780-788.

doi:10.1007/s12282-015-0641-8

Uphoff, C. C., & Drexler, H. G. (2011). Detecting mycoplasma contamination in cell cultures by

polymerase chain reaction. *Methods Mol Biol, 731*, 93-103. doi:10.1007/978-1-61779-

080-5_8

Venkatesan, S., Rosenthal, R., Kanu, N., McGranahan, N., Bartek, J., Quezada, S. A., . . .

Swanton, C. (2018). Perspective: APOBEC mutagenesis in drug resistance and immune

escape in HIV and cancer evolution. *Ann Oncol, 29*(3), 563-572.

doi:10.1093/annonc/mdy003

Verhalen, B., Starrett, G. J., Harris, R. S., & Jiang, M. (2016). Functional upregulation of the DNA

cytosine deaminase APOBEC3B by polyomaviruses. *J Virol, 90*(14), 6379-6386.

doi:10.1128/JVI.00771-16

Vieira, V. C., Leonard, B., White, E. A., Starrett, G. J., Temiz, N. A., Lorenz, L. D., . . . Harris, R.

S. (2014). Human papillomavirus E6 triggers upregulation of the antiviral and cancer

genomic DNA deaminase APOBEC3B. *MBio, 5*(6). doi:10.1128/mBio.02234-14

Walker, B. A., Wardell, C. P., Murison, A., Boyle, E. M., Begum, D. B., Dahir, N. M., . . . Morgan,

G. J. (2015). APOBEC family mutational signatures are associated with poor prognosis

translocations in multiple myeloma. *Nat Commun, 6*, 6997. doi:10.1038/ncomms7997

Wang, J., Becker, J. T., Shi, K., Lauer, K. V., Salamango, D. J., Aihara, H., . . . Harris, R. S.

(2019). The role of RNA in HIV-1 Vif-mediated degradation of APOBEC3H. *J Mol Biol,*

431(24), 5019-5031. doi:10.1016/j.jmb.2019.09.014

Warren, C. J., Westrich, J. A., Doorslaer, K. V., & Pyeon, D. (2017). Roles of APOBEC3A and

APOBEC3B in human papillomavirus infection and disease progression. *Viruses, 9*(8).

doi:10.3390/v9080233

Warren, C. J., Xu, T., Guo, K., Griffin, L. M., Westrich, J. A., Lee, D., . . . Pyeon, D. (2015).

APOBEC3A functions as a restriction factor of human papillomavirus. *J Virol, 89*(1), 688-

702. doi:10.1128/JVI.02383-14

Westrich, J. A., Warren, C. J., Klausner, M. J., Guo, K., Liu, C. W., Santiago, M. L., & Pyeon, D.

(2018). Human papillomavirus 16 E7 stabilizes APOBEC3A protein by inhibiting CULLIN

2-dependent protein degradation. *J Virol, 92*(7). doi:10.1128/JVI.01318-17

Yamazaki, H., Shirakawa, K., Matsumoto, T., Hirabayashi, S., Murakawa, Y., Kobayashi, M., . . .

Takaori-Kondo, A. (2019). Endogenous APOBEC3B overexpression constitutively

generates DNA substitutions and deletions in myeloma cells. *Sci Rep, 9*(1), 7122.

doi:10.1038/s41598-019-43575-y

Yamazaki, H., Shirakawa, K., Matsumoto, T., Kazuma, Y., Matsui, H., Horisawa, Y., . . . Takaori-

Kondo, A. (2020). APOBEC3B reporter myeloma cell lines identify DNA damage response

pathways leading to APOBEC3B expression. *PLoS One, 15*(1), e0223463.

doi:10.1371/journal.pone.0223463

Zou, X., Koh, G. C. C., Nanda, A. S., Degasperi, A., Urgo, K., Roumeliotis, T. I., . . . Nik-Zainal,

S. (2021). A systematic CRISPR screen defines mutational mechanisms underpinning

signatures caused by replication errors and endogenous DNA damage. *Nat Cancer, 2*(6),

643-657. doi:10.1038/s43018-021-00200-0

Zou, X., Owusu, M., Harris, R., Jackson, S. P., Loizou, J. I., & Nik-Zainal, S. (2018). Validating

the concept of mutational signatures with isogenic cell models. *Nat Commun, 9*(1), 1744.

doi:10.1038/s41467-018-04052-8

1092

Main and Supplementary Figure Legends

Figure 1. A3 activity in the HAP1-TK-M9 mutation reporter system.

(**A**) Schematic of the construction of the HAP1-TK-M9 system and overall experimental workflow (see **Figure 1 – Figure Supplement 1** for representative HAP1-TK-M9 karyotype images).

(**B**) *A3* mRNA levels in parental HAP1 cells in comparison to the HAP1-TK-M9 daughter clone by RT-qPCR (*** = $p < 0.001$, NS = non-significant, Welsh's t-test).

(**C-E**) Immunoblots of A3A, A3B, and A3H following transduction of HAP1-TK-M9 cells. Tubulin (Tub) is a loading control (see **Figure 1 – Figure Supplement 2** for anti-A3A mAb validation). The lower images show ssDNA deaminase activity of extracts from the same cells (S, substrate; P, product).

(**F**) Immunofluorescent microscopy images of HAP1-TK-M9 cells expressing the indicated A3 enzymes (see methods for primary antibodies). A3 staining is green, γ-H2AX staining is red, and nuclei are blue from Hoechst. A3A is cell wide, A3B is predominantly nuclear, and A3H-I and -II are cell wide with nucleolar concentrations (scare bar = 25 µm). The lower row shows representative COMET images visualized with SYBR Gold (1:10,000) in the GFP channel.

(**G-H**) Quantification of γ-H2AX staining and COMET moments in experiments shown in panel F (each data point represents an independent cell; p-values determined with Welch's t-test).

Figure 1 – Supplementary Figure 1. HAP1-TK-M9 karyotype analysis.

Representative G-band and spectral karyotype (SKY) images of HAP1-TK-M9 M-phase chromosomes showing a near diploid DNA content and previously reported aberrations including the reciprocal chromosome 9:22 translocation (Philadelphia chromosome) characteristic of CML tumor cells. The Y-chromosome is missing, as reported previously in the KBM7 parent line of HAP1.

Figure 1 – Supplementary Figure 2. Validation of a custom rabbit-anti human APOBEC3A mAb UMN-13.

(**A**) Schematic of human A3A, A3B, and A3G indicating the unique N-terminal epitope used here to generate an A3A-specific mAb UMN-13. The schematic also shows the C-terminal epitope used previously to generate the versatile 5210-87-13 mAb that recognizes these three enzymes.

(**B**) Comparative immunoblots of whole cell extracts from 293T cells expressing each of the 7 human A3 family members and probed with either our custom rabbit anti-human A3A mAb UMN-13 (left) or a commercial anti-HA mAb as an expression control (right).

(**C**) Comparative immunoblots of whole cell extracts from the monocytic cell line THP-1 and a clonal derivative lacking A3A-through-A3G (ΔA-G) plus/minus treatment with LPS/IFN-α to induce expression of multiple A3s including A3A and A3G. The UMN-13 mAb blot on the left shows a single band representing full-length A3A (starting at Met1), and the 5210-87-13 mAb blot on the right shows A3G (strong top band), A3B (weak band just below A3G), and both A3A translation products (strong faster-migrating bands for full-length A3A starting at Met1 and the shorter isoform starting at Met13).

Figure 2. Characterization of *TK* mutations in ganciclovir-resistant clones.

(**A**) A dot plot of Gan^R colonies generated under the indicated A3 expression conditions. Each data point represents the number of Gan^R mutants in a single clonal culture (mean +/- SD shown with significance assessed using Welch's t-test).

(**B**) Schematics representing all *TK* mutations observed under the indicated A3 expression conditions (APOBEC3 signature T(C>T/G)W mutations in red, other SBSs in black, and InDels in blue; see **Figure 2 – Figure Supplement 1** for schematics of individual *TK* mutants for these and vector control conditions). Q8X and R212K mutation hotspots are labeled.

(**C-D**) Pentanucleotide logos depicting -2 and +2 sequence preferences flanking all T(C>T/G)W mutations observed under the indicated conditions.

1145

Figure 2 – Supplementary Figure 1. Mutations in TK genes derived from ganciclovir-resistant clones.

Schematics of TK mutation frequencies in ganciclovir-resistant clones per individual clone. T[C>G/T]W mutations are shown in red, and other SBSs are shown in black. InDels are shown in blue. Composite sequence mutation schematics are shown below each list of sequences per condition.

Figure 3. Single base substitution mutation signatures in ganciclovir-resistant clones by whole genome sequencing.

(**A**) Trinucleotide profiles of pooled SBSs across all clones sequenced for each listed experimental condition (A3A WT, n=6; A3A E72A, n=2; A3B WT, n=5; A3B E255A, n=2. APOBEC3 signature T(C>T/G)W mutations are highlighted by red dashed-line boxes. See **Figure 3 – Figure Supplement 1 and 2** for mRNA and protein level expression confirmation, respectively, and **Figure 3 – Figure Supplement 3** for SBS profiles from each WGS).

(**B-C**) Pentanucleotide logos depicting -2 and +2 sequence preferences flanking all T(C>T/G)W mutations attributable to A3A and A3B, respectively.

Figure 3 – Supplementary Figure 1. *A3* mRNA expression in the HAP1-TK-M9 system.

(**A**) *A3A*, *A3B*, and *A3H* mRNA expression levels relative to those of the housekeeping gene *TBP* for the indicated HAP1-TK-M9 conditions (RNA-seq FKPM from n$\geq$2 Gan^R clones for each condition; mean +/- SD shown). RNA-seq data from A3 signature-high breast cancer cell lines (BT-474 and MDA-MB-453), CCLE breast-derived cell lines (n=52), and TGCA primary breast cancers (n=1093) are presented alongside for comparison (mean +/- SD).

(**B**) A heatmap depicting mean expression levels of all 7 human *APOBEC3* family members, in addition to *APOBEC1*, *APOBEC2*, *APOBEC3*, and *AICDA*, relative to those of the housekeeping

1171   gene *TBP* (RNA-seq values are FKPM; n$\geq$2 for each condition to provide matching data sets for

1172   the Gan^R clones subjected to WGS). Endogenous *A3C* provides a robust internal control.

1173

1174   **Figure 3 – Supplementary Figure 2. A3 protein expression in Gan^R clones.**

1175   (**A**) Immunoblots of A3A, A3B, and A3H in the indicated Gan^R clones. Tubulin (TUB) is a loading

1176   control.

1177   (**B**) Deaminase activity of WCE on ssDNA from the same clones (S, substrate; P, product).

1178

1179   **Figure 3 – Supplementary Figure 3. Single base substitution mutation signatures in**

1180   **individual ganciclovir-resistant clones by whole genome sequencing.**

1181   Trinucleotide profiles of the single base substitution mutations found in WGSs from the indicated

1182   Gan^R clones. Composite profiles for each condition are shown in **Figure 3**.

1183

Figure 3 – Supplementary Figure 4. APOBEC3 mutation signature extracted from WGSs

using complementary methods.

(**A**) Mutation signature profiles extracted from Gan^R clone WGS using an NMF-based approach.

For each clone, all SBSs resolved into 2 signatures - an APOBEC-like signature shown in red and

a background signature in blue. APOBEC3 mutation enrichment scores are shown above each

bar, with significantly enriched values shown in red (BH FDR corrected q-value < 0.05).

(**B**) Trinucleotide profiles of the two signatures derived from NMF in panel A clearly showing

APOBEC3 signature T(C>T/G)W mutations.

1192

Figure 4. Mesoscale properties of A3A and A3B *in vitro* and in the HAP1-TK-M9 system.

(**A-B**) DNA deamination kinetics of A3A and A3B using *SDHB* and *NUP93* hairpin substrates in

comparison to corresponding linear controls made by scrambling the 5' or 3' portion of the stem,

respectively. See text for full description and **Figure 4 – Figure Supplement 1** for a genome-

wide analysis.

(**C**) Rainfall plots of genome-wide intermutation distances showing APOBEC3 signature kataegic

tracts (red arrows) in representative A3A- and A3B-expressing Gan^R clones (C>T mutations are

red, C>G black, C>A blue, T>A purple, T>C yellow, and T>G green).

1201

Figure 4 – Figure Supplement 1. Hairpin mutation analysis in HAP1-TK-M9 WGS data.

Pie charts representing the fraction of APOBEC3 signature mutations and other types of single

base substitution mutations found in predicted hairpin substrates (red) versus non-hairpin

contexts (grey). The total number of APOBEC3 signature T(C>T/G)W mutations is shown in the

center of each pie chart. Hairpin stems and loops were defined liberally as any feature between

3 and 100 nucleotides, and a hairpin is considered mutated if it has a SBS mutation in any position

of the loop region.

1209

1210   **Figure 5. A composite origin of APOBEC3 signature mutations in breast cancer.**

1211   A Euclidean distance analysis of similarity between the SBS profiles from WGSs of the A3A, A3B,

1212   and control Gan^R clones described here versus primary breast tumor whole-genome sequencing

1213   data (ICGC, n=794). The APOBEC3 signature is represented by both enrichment score and

1214   SBS2+13 (red), HRD signature as SBS3 (blue), and ageing signature as SBS1 (gray).

1215

Figure 1

Figure 1 - Supp 1

Figure 2

Figure 2 - Supp 1

Figure 3

Figure 3 - Supp 1

Figure 3 - Supp 2

A

B

	Control	A3A					A3A E72A		A3B					A3B E255A		A3H-I		Vec.	
		D06	D07	D09	D12	D21	D28	D29	D51	D52	D54	D59	D62	D76	D77	D99	D100	D168	D169
S																			
P																			

Figure 3 - Supp 3

Figure 3 - Supp 4

A

B

Figure 4

A

SDHB hairpin

5'-TG_{CCATC} T A
FAM-AG_{GGTAG} **C** T

5'-TGCCATCTAT**C**GATGGGA-FAM

SDHB linear

5'-TGTAGCCTAT**C**GATGGGA-FAM

B

NUP93 hairpin

5'-FAM-_{GCAAGCT} G T
AGT_{CGTTCGA} **C** T

5'-FAM-GCAAGCTGTT**C**AGCTTGCTGA

NUP93 linear

5'-FAM-GCAAGCTGTT**C**ACGTCTGGTA

C

Figure 4 - Supp 1

	A3A WT	A3A E72A	A3B WT	A3B E255A	A3H-I WT
APOBEC	62 (2.2%) 2830	2 (1.7%) 122	18 (2.6%) 691	12 (8.3%) 144	0 (0%) 199
Other	613 (2.5%) 24486	121 (2.6%) 4518	380 (2.3%) 16240	148 (2.8%) 5252	173 (2.8%) 6265

Figure 5