1 How to measure response diversity

Samuel R.P-J. Ross[1*], Owen L. Petchey[2], Takehiro Sasaki[3], David W. Armitage[1]

[1]Integrative Community Ecology Unit, Okinawa Institute of Science and Technology Graduate

University, Onna-son, Okinawa, 904-0495 Japan.

[2]Department of Evolutionary Biology and Environmental Studies, University of Zurich,

Winterthurerstrasse 190, 8057 Zürich, Switzerland.

[3]Graduate School of Environment and Information Sciences, Yokohama National University, 79-7

Tokiwadai, Hodogaya, Yokohama, 240-8501 Japan.

Email: s.ross.res@outlook.com (SRP-JR); owen.petchey@ieu.uzh.ch (OLP); sasaki-takehiro-

kw@ynu.ac.jp (TS); david.armitage@oist.jp (DWA).

ORCID: 0000-0001-9402-9119 (SRP-JR); 0000-0002-7724-1633 (OLP); 0000-0001-8727-9152 (TS);

0000-0002-5677-0501 (DWA).

CRediT Author Contribution Statement| SRP-JR: Conceptualisation, Software, Formal analysis,

Funding acquisition, Validation, Investigation, Visualisation, Methodology, Writing – original draft.

OLP: Conceptualisation, Data curation, Software, Validation, Methodology, Writing – reviewing and

editing. **TS:** Conceptualisation, Methodology, Writing – reviewing and editing. **DWA:**

Conceptualisation, Software, Formal Analysis, Methodology, Writing – reviewing and editing.

Running title: How to measure response diversity

Keywords: diversity-stability debate; functional diversity; resilience; functional trait; asynchrony;

environmental change; temporal variability; functional response; insurance hypothesis; nonlinear

responses.

Article type: Synthesis

Words: Abstract 198 words; main text 4,765 words; boxes 1,009 words.

References: 134 references.

Display items: 4 figures (+1 in Box 2); 0 tables; 2 text boxes.

*Correspondence: Samuel R.P-J. Ross, Integrative Community Ecology Unit, Okinawa Institute of

Science and Technology Graduate University, 1919-1 Tancha, Onna-son, Kunigami-gun, Okinawa, 904-

0495 Japan. E: s.ross.res@outlook.com

35 Abstract

Taxonomic diversity metrics are, at best, proxies for the inter- and intraspecific trait differences that

mechanistically underlie the insurance effect of biodiversity—that diversity enhances and stabilises

aggregate ecosystem properties. In theory, variation in functional (*e.g.*, biomass) responses to

environmental change can buffer and maintain ecosystem functioning. This variation, termed

response diversity, is therefore a potentially critical determinant of ecological stability. However,

response diversity has yet to be widely quantified, possibly due to difficulties in its measurement.

Even when it has been measured, approaches have varied. Here, we review methods for measuring

response diversity and propose a methodological framework for quantifying response diversity from

experimental or observational data, which can be practically applied in lab and field settings across a

range of taxa. Our approach, which is based on environment-dependent functional responses to any

biotic or abiotic environmental variable, is conceptually simple and robust to any form of

environmental response, including nonlinear responses. By capturing even subtle inter- or

intraspecific variation in environmental responses, and environment-dependencies in response

diversity, we hope this framework will motivate tests of the diversity-stability relationship from a new

perspective, and provide an approach for mapping, monitoring, and protecting this critical element of

biodiversity.

Introduction

Ecological stability has been a core focus of ecology since the 1950s when interest in the

relationship between diversity and stability first arose (McCann 2000). Ecological stability is a

multidimensional concept encompassing a range of metrics for measuring resistance to and recovery

from disturbance, and variability in time and space (Pimm 1984; Donohue *et al.* 2013). Early

observations suggested that more diverse communities are more stable in their ability to resist

invasions or in the magnitude of their fluctuations (Odum 1953; MacArthur 1955; Elton 1958). Using

randomly structured model communities, May (1973) then demonstrated that diversity *per se* did not

increase stability (*i.e.*, did not dampen population fluctuations), but rather decreased it (see also

Pimm & Lawton 1978; Yodzis 1981). Though debate continues (Jacquet *et al.* 2016; Pennekamp *et al.*

2018), subsequent developments (*e.g.*, McCann *et al.* 1998; Kondoh 2003; Ives & Carpenter 2007)

have led to the consensus that in more complex models (that include for example, variable

environments, interaction strengths, or behaviours), species-rich systems will exhibit more stable (less

variable) aggregate ecosystem properties compared to species-poor systems (Ives *et al.* 1999). This is

because declines in one species' abundance or performance—that is, any aspect of an organism's

higher- or lower-level functional attributes (traits) that contributes to individual fitness or higher-level

(ecosystem) functioning (Hooper *et al.* 2005)—will be offset by neutral or positive responses of other

species or phenotypes (reviewed by Loreau *et al.* 2021). Together, these ideas encapsulate the

insurance effect of biodiversity (Yachi & Loreau 1999)—that biodiversity both enhances and stabilises

ecosystem functioning. However, despite biodiversity being only the "*passive recipient*" through

which underlying ecological mechanisms drive stability (McCann 2000; Shade 2017), attempts to find

a general and empirically quantifiable dimension of biodiversity that reliably predicts stability across

contexts have had limited success. Various mechanisms giving rise to the insurance effect have been

proposed, including: variable interaction strengths (*e.g.*, Yodzis 1981; McCann *et al.* 1998) or mixtures

of interaction types (*e.g.*, Mougi & Kondoh 2012; Hammill *et al.* 2015); food web nestedness (*e.g.*,

Thebault & Fontaine 2010); trophic flexibility and food web rewiring (*e.g.*, Kondoh 2003); dominant,

rare, or otherwise key species (*e.g.*, Walker *et al.* 1999; Sasaki & Lauenroth 2011; Arnoldi *et al.* 2019;

Ross *et al.* 2022); and asynchronous or compensatory dynamics (Gonzalez & Loreau 2009; Wang *et al.*

2021) in time (*e.g.*, Craven *et al.* 2018; Sasaki *et al.* 2019) or space (*e.g.*, Loreau *et al.* 2003; Wang &

Loreau 2016). Recently, the concept of *response diversity* has also emerged as a potentially key, yet

under-explored driver of stability (Pennekamp *et al.* 2018; Kahiluoto *et al.* 2019; Sasaki *et al.* 2019).

Response diversity characterises the range of responses to the environment displayed among

members (e.g., individuals, species) of a focal group such as a population, guild, or community

(Elmqvist *et al.* 2003; Nyström 2006; Mori *et al.* 2013). The concept was co-opted from elsewhere in

the biological sciences, where response diversity describes, for example, the variation in tumour

responses to radiation therapy (Hlatky *et al.* 1994) or the range of cellular reactions to stimuli in the

brain (Burrows *et al.* 1997). In ecology, Ives *et al.* (1999) posited that differences in how species

respond to environmental change (that is, response diversity) may explain species diversity's

stabilising effect on community stability. In a later study, Ives & Carpenter (2007) found that, in a

model of randomly structured competitive communities, response diversity's stabilising effect on

communities negated the destabilising effects of interspecific interactions. The authors conclude that

species-environment relationships were more important for stability than were interspecific

interactions (Ives & Carpenter 2007). Interspecific variation in species-environment relationships is

critical for the maintenance of diversity in variable environments (Chesson 2000) and species-rich

systems are thus anticipated to exhibit a wider diversity of species-environment interactions

compared to species-poor systems (Vogel *et al.* 2019). This response diversity may, in turn, support a

variety of stabilising mechanisms acting on the aggregate properties of communities and ecosystems.

For example, response diversity may be the ecological mechanism underpinning many of the

statistical explanations of the diversity-stability relationship, such as the portfolio effect (Tilman *et al.*

1998; Loreau 2010; Loreau *et al.* 2021). Moreover, response diversity can promote asynchrony in

species' biomass or abundance through time (Sasaki *et al.* 2019) and species-specific tolerance to

environmental conditions may be partly responsible for temporal and functional complementarity

among species (Petchey 2003; Loreau & de Mazancourt 2013). Indeed, asynchrony may often be an

outcome of response diversity (*e.g.*, Sasaki *et al.* 2019), suggesting that the often-observed stabilising

effect of asynchrony on aggregate community or ecosystem properties (*e.g.*, Morin *et al.* 2014;

Hector *et al.* 2015), may in fact be a by-product of underlying differences in species-environment

relationships. It is thus beneficial to measure response diversity directly instead of (or as well as)

asynchrony, since response diversity may capture properties of underlying species-environment

relationships that do not manifest as asynchrony of biomass (or other functional attributes) through

time. Further, given the challenge of statistically separating asynchrony among populations from

asynchrony in species-environment responses (de Mazancourt *et al.* 2013), it may be conceptually

and empirically simpler to focus efforts on the ultimate rather than proximate causal mechanism:

response diversity.

There are still relatively few empirical studies that explicitly measure response diversity.

Though we recognise that many studies may measure response diversity without labelling it such, a

systematic review of the literature returned 47 ecology papers that empirically measured response

diversity (see Fig. 1). These empirical studies used various methods to measure response diversity.

The majority (*n* = 28 studies, ~60%) measured response diversity using functional traits (see Box 1 for

discussion), where the diversity of response traits— that is, those traits that predict some aspect of

how a species responds to the environment (Suding *et al.* 2008)—represents response diversity,

measured using the functional dispersion [FDis] index (*e.g.,* Spasojevic *et al.* 2016; Hordley *et al.* 2021;

Schnabel *et al.* 2021). Some studies measuring the dispersion of functional response traits explicitly

do so after defining functional effect groups (Laliberté *et al.* 2010; Thornhill *et al.* 2018; Sasaki *et al.*

2019; Morel *et al.* 2020), producing a response diversity measure closely aligned to the Elmqvist *et al.*

(2003) definition, which specifies that response diversity should be measured among species

contributing to the same ecosystem function (though subsequent definitions relax this requirement;

e.g., Mori *et al.* 2013). Most empirical studies measuring response trait diversity do so for trees (*e.g.*,

Craven *et al.* 2016; Altomare *et al.* 2021; Schnabel *et al.* 2021) and other groups of plants (*e.g.*,

Mandle & Ticktin 2015; Döbert *et al.* 2017; Morel *et al.* 2020), or for freshwater or terrestrial

invertebrates (*e.g.*, Mumme *et al.* 2015; Thornhill *et al.* 2018).

132   Another approach to measuring response diversity empirically is to define a species-specific

interaction term in multi-species models of, for example, abundance as a function of the environment

(*n* = 8 studies; Fig. 1). If the species-specific interaction term passes some significance or model

selection threshold, then species are claimed to differ in their abundance ~ environment slopes; that

is, they exhibit response diversity (Winfree & Kremen 2009). This approach to measuring response

diversity has emerged primarily from the pollination literature (Winfree & Kremen 2009; Bartomeus

et al. 2013; Cariveau *et al.* 2013; Stavert *et al.* 2017), but has been used elsewhere (Fauchald *et al.*

2011; Malyshev *et al.* 2016), and is based on the assumptions of linear regression models, meaning

nonlinear responses are not captured. Another related but less common approach to measuring

response diversity (*n* = 2 studies) is to measure some aspect of a population's performance, such as

its intrinsic rate of increase (*e.g.*, McCann 2016) or biomass change in response to the environment,

and take the range of performance-environment model slopes as a direct measure of response

diversity. Leary & Petchey (2009) did this with four ciliate species by modelling their intrinsic rates of

increase against temperature when in isolation, first determining species-specific environmental

responses. They then assembled experimental communities with different combinations of these

species, producing different levels of response diversity. Using observational time series of reef fish

abundance, Thibaut *et al.* (2012) similarly quantified response diversity as the average correlation

coefficient between pairwise combinations of species in terms of their intrinsic rates of increase as a

function of the environment. These various methods for measuring response diversity have emerged

independently, resulting in a disconnect among empirical response diversity studies, and a general

methodological framework for measuring response diversity is lacking.

Figure 1. PRISMA diagram detailing the results of our systematic review. On 17 Dec 2021, we conducted a Web

of Science search for "response diversity" in all fields within the "Ecology" Web of Science category. We

supplemented the resulting 175 papers with four that we knew to be relevant but were not included in the

initial results. We first excluded 49 articles that were not empirical studies (reviews, perspectives, *etc*.). Next, we

screened the full-text articles and excluded 83 papers that did not measure response diversity (whether termed

response diversity or not) based on the broad definition that response diversity should in some way characterise

the range of species responses to environmental conditions. The resulting 47 empirical response diversity

papers measured response diversity using various methods, the majority using functional response traits,

followed by species-specific interaction terms in multi-species models of responses to the environment. We

additionally searched using the term "response variation" which returned nine hits, of which eight were

excluded and one was duplicate with our initial search. For consistency in categorisation and exclusion criteria,

all screening was conducted by a single researcher (S.R.P-J.R.).

Measuring response diversity empirically

Here, we propose a framework for empirically measuring response diversity using a

conceptually simple and empirically tractable approach that can be applied in observational and

experimental settings in both the laboratory and field. We extend the models describing

performance-environment relationships presented above to include any functional attribute that

represents a species' functional response along a continuous gradient of abiotic or biotic

environmental condition. In contrast to most previous studies of response diversity which focus on

'low-level' functional traits such as nitrogen content or specific leaf area (Box 1), our approach

considers response diversity a 'high-level' trait of *per capita* growth rates for instance, tying it more

directly to species performance. We discuss the strong potential for non-linearity in responses leading

to environment-dependent response diversity, and the importance of divergent responses to

environmental change in stabilising aggregate ecosystem properties. We also discuss the importance

of assessing if the diversity of responses includes responses that differ in direction. We then

demonstrate the ease with which response diversity can be measured and manipulated for use in

empirical tests of the diversity-stability relationship.

We propose to extend the performance-environment models used elsewhere (Leary &

Petchey 2009; Thibaut *et al.* 2012) into a generalisable framework for measuring response diversity.

Despite rarely being used to date, this method is easily tractable in experimental and observational

studies (Thibaut *et al.* 2012), provides a more direct measurement of how species respond to the

environment than, for example, functional response traits (Box 1), and allows identification of key

contributors to response diversity (Leary & Petchey 2009). In such models the response variable need

not be intrinsic rate of increase, but could in fact be any relevant functional variable, such as biomass,

abundance, or an ecosystem function to which the species contributes (Fig. 2). Similarly, the

'*environment*' to which species are responding does not have to be an abiotic feature such as

temperature; it can be any continuous biotic or abiotic condition. For example, one could model the

responses along a land-use intensification gradient (*e.g.*, Winfree & Kremen 2009; Moore & Olden

2017; Stavert *et al.* 2017; Fig. 2b), or in response to the abundance of a predator, competitor, or the

availability of a prey species (*e.g.*, Fauchald *et al.* 2011; Fig. 3). The latter approach, using biotic

conditions as the environment to which a species responds, is conceptually similar to modelling the

functional response (Holling 1966), which in turn may scale with abiotic conditions such as

temperature (Daugaard *et al.* 2019; Sohlström *et al.* 2021). Similarly, modelling biomass change in

response to a competitor is akin to concepts in coexistence theory such as negative-frequency

dependent growth (*e.g.*, Armitage & Jones 2019) or invasion growth rate (Grainger *et al.* 2019).

From such models, we suggest measuring response diversity as the variation in the slope of a

performance-environment function evaluated over the environmental gradient in question.

Specifically, we suggest using Generalised Additive Models (GAMs) to fit performance-environment

relationships individually for each species using smoothing parameters as appropriate (Fig. 2a-c), then

taking the first derivatives of these models to estimate model slopes along the environmental axis

(Fig. 2d-f). GAMs are free from many of the assumptions of linear regressions and can be used to

evaluate both linear and nonlinear responses, or a mixture of the two (Hastie & Tibshirani 1987; see

Github). Response diversity can then be measured as the variation in the distribution of these

derivatives to capture the variation in species-specific environmental responses (Fig. 2g-i). In practice,

this variation could be described using any suitable estimator, but here we use a two-dimensional

measure.

First, we use a similarity-scaled measure of diversity introduced by Leinster & Cobbold (2012).

Briefly, similarity-based diversity (Eq. 1 in Leinster & Cobbold 2012) is calculated based on pairwise

Euclidean distances (that is, *dissimilarity*) in performance-environment relationships between all pairs

of species in the community. The metric is rooted in Hill numbers, with an exponent (q) defining the

relative weighting of richness and evenness (Leinster & Cobbold 2012). Here, we fix q at zero, to focus

on species richness, though our measure of dissimilarity includes information on species relative

abundances within the S x S similarity matrix. Thus, we can use this similarity-based diversity as a

metric capturing information on both richness and similarity in species' responses (see Leinster &

Cobbold 2012; Github). We chose this estimator since it has several desirable properties for

describing response diversity: it cannot decrease with species additions; does not increase with

additions of redundant species responses; and captures even subtle differences in the distribution of

different responses within the overall range. Response diversity should therefore be lowest (1) when

all species respond identically (Fig. 2g), and highest when species responses are as dissimilar as

possible (the estimator converges to species richness as performance-environment slopes diverge to

infinity and –infinity).

Critically, whether the first derivatives of performance-environment relationships span zero is

a key indicator of the stabilising potential of response diversity under a given environmental

condition. As such, we also isolate and measure this property of response diversity using the formula

$\frac{\text{range}(x_e)- \left\lVert \max(x_e)\rvert-\lvert\min(x_e)\right\rVert}{\text{range}(x_e)}$, where $\mathbf{x_e}$ represents the derivative of a performance-environment

relationship for environmental condition (**e**). If the range of first derivates does not span zero (*e.g.*,

Fig. 2d,e), then all species respond in the same direction to the environment (albeit at varying

magnitudes). In such cases, there is no response diversity along this sign-sensitive axis of species

responses (the estimator is zero). In contrast, when first derivatives are perfectly symmetric around

zero, the value is one. Without interspecific interactions, even large differences in the magnitude of

response (that is, the dimension of response diversity discussed above) cannot be stabilising (Ives *et*

al. 1999; Mougi & Kondoh 2012). Conversely, when the range of first derivatives spans zero for a

given environmental state (*e.g.*, Fig. 2f), species responses are diverging (at least one species is

decreasing while another is increasing), which we expect *a priori* to be stabilising. Our estimator is

largest (1) when $\min(x_e) = -\max(x_e)$. If observing stable aggregate ecosystem properties (*e.g.*,

biomass) when the range of first derivatives does not span zero, this suggests response diversity is not

the mechanism driving stability in that case. Together, our chosen similarity-based diversity measure

and this sign-sensitive index make up the two core dimensions of response diversity of interest. The

specific metrics we use here may not be ideal for all use cases. Further work should thus consider the

need to develop suitable estimators of response diversity when those proposed fall short, including

the possibility of uniting the two dimensions of response diversity discussed here into a single metric.

That said, we demonstrate here the utility of these estimators for quantifying response diversity in a

range of contexts (Figures 2 and 3), including applications to real data (Box 2).

Figure 2. Measuring response diversity for linear species-environment relationships. We reimagined the original

analyses of three studies, using standardised, simulated data to demonstrate how response diversity could be

measured practically under our framework. (**a**) We propose a hypothetical case (based on Malyshev *et al.* 2016),

where *Chlorophyll* production (which is often an indicator of biomass, particularly for aquatic plants) declines

linearly across several species [coloured lines] of plants with spring frost intensity, with no difference among

species in the slope of this decline. (**d**) We can calculate the first derivatives of these responses, which in this

case would be identical for all species. (**g**) Then, we can measure response diversity using our proposed

similarity-based diversity measure (see main text), and find it to be one (its lower bound), since all species

respond identically to spring frost. Our sign-sensitive index is zero in this case, as the first derivatives do not

span zero. (**b**) Next consider a case (based on Thornhill *et al.* 2018), where the abundance of several species of

macroinvertebrates decline linearly, but at different magnitudes, along an urbanisation gradient. (**e**) Here, the

first derivatives differ (as a result of the differing model slopes), resulting in a higher dissimilarity value (**h**).

Again, the derivatives do not span zero, so the sign-sensitive index is zero. (**c**) Finally, we might be interested in

the decline in mass of several bee species as a function of winter temperature (see Fründ *et al.* 2013). In this

example, species responses to winter temperature diverge (one species increases, one decreases, and one is

unaffected), but all such responses are linear, and the magnitude of the responses are not as extreme as in (**b**).

(**f**) Thus, the first derivatives are less spread out than in (**e**), but span zero. (**i**) Accordingly, our dissimilarity value

is lower than for (**h**), but in this case, our sign-sensitive index is at its maximum value, 1, across the entire range

of winter temperature values (the derivatives span zero at any temperature). Dashed lines indicate zero in

panels **a**–**f**, and the median value of the similarity-based diversity index (dissimilarity) along the X-axis in panels

g–**I** (here, equivalent to any value along the X-axis since response diversity is constant).

Ecological responses to the environment are often nonlinear, necessitating a wide range of

statistical tools for considering such responses (Fraterrigo & Rusak 2008; Ross *et al.* 2021b). When

species' responses are nonlinear (*e.g.*, Leary & Petchey 2009), quantifying response diversity is not as

simple as measuring the range of model slopes. However, the statistical framework we propose for

measuring response diversity (that is, measuring variation in performance-environment first

derivatives) is robust to nonlinearities such as species' functional responses to competition,

predation, or resource availability (Fig. 3) or when tipping points are detected in empirical data

(Berdugo *et al.* 2020). Figure 3 illustrates three such cases: one where the intrinsic rate of increase of

flowering plant species responds nonlinearly but with low response diversity to total flowering plant

abundance (Fig. 3a); one where seedling establishment responds to grazing pressure with higher

response diversity among species (Fig. 3b); and one where trees respond nonlinearly to fire

frequency, and differ in the form of this relationship (Fig. 3c)—our framework uses GAMs as they are

robust even to cases where the form of the performance-environment relationship differs among

species (Hastie & Tibshirani 1987; Fig. 3f,i). Where responses are nonlinear, the first derivatives of

performance-environment responses vary as a function of the environmental variable (along the X-

axis in Figures 3d-f). In turn, this produces a measure of response diversity that also varies as a

function of the environment (Fig. 3g-i). In cases where there is no need to treat response diversity as

environment dependent, one could estimate a single response diversity value across the

environmental axis using any summary statistic (see median response diversity in Figures 3g-i),

though it may be appropriate to weight such a summary statistic by the distribution of the

environmental variable. In linear cases, response diversity does not vary as a function of environment,

so summary statistics are not required (Fig. 2).

Figure 3. Measuring response diversity for nonlinear species-environment relationships. We reimagined the

original analyses of three studies, using standardised, simulated data to demonstrate how response diversity

could be measured practically under our framework. (**a**) We propose a hypothetical case (based on Ishii &

Masahiko 2001), where flowering plant intrinsic rate of increase changes as a function of total flowering plant

abundance. Here, responses may be nonlinear but similar in form, as competition is limiting at high abundance

while pollinator visitations might be limiting where total flowering plant abundance is low (Ishii & Masahiko

2001; Reilly *et al.* 2020). (**d**) In this case, the first derivatives of the intrinsic rates of increase vary as a function

of flowering plant abundance, but do not differ much among species. (**g**) This results in a dissimilarity value that

is close to its minimum (1) but varies a little with total flowering plant abundance. (**b**) Next consider a case

(based on Ford *et al.* 2018), where seedling establishment of several plant species respond nonlinearly to

grazing pressure. This might be the case if, for example, one species is a shade-tolerant understorey specialist

[light blue] that does well when grazing pressure is low but not when high, while another species is a shade-

intolerant disturbance specialist [dark blue] (Ford *et al.* 2018). (**e**) These divergent nonlinear responses produce

a range of first derivatives that change as a function of grazing pressure, in turn producing (**g**) grazing pressure-

dependent response diversity which is on, average, high in terms of dissimilarity, and is zero-spanning at

intermediate grazing pressure. (**c**) Finally, consider a case (based on Altomare *et al.* 2021), where the density of

several tree species responds nonlinearly to fire frequency with different functional forms. This might occur if

some species are highly sensitive to burning [brown line] while others are disturbance specialists that require

nutrient inputs from fire or reduced competition from dominant fire-sensitive species (Altomare *et al.* 2021). (**f**)

Here, the first derivatives of tree density differ considerably and vary as a function of fire frequency, producing

(**i**) generally high response diversity—in both dissimilarity and sign—that also varies with fire frequency. Here,

response diversity is largest at low and high fire frequency (when specialists respond strongly) and is lowest at

intermediate fire frequency. Dashed lines indicate zero in panels **a**–**f**, and the median value of response diversity

along the X-axis in panels **g**–**i**.

By modelling individual species responses to environmental conditions with GAMs, this

approach is conceptually similar to the species-specific interaction term described above (Winfree &

Kremen 2009; Cariveau *et al.* 2013). However, the framework we describe provides a quantitative

estimate of response diversity, identifying precisely the relevant differences in the form of species

environmental responses (divergence, magnitude, *etc*.) that have the potential to drive ecological

stability, and allowing consideration of response diversity as a function of the environment. This

method is flexible enough to be used on traits, rates (first derivatives), or sensitivities of rates (second

derivatives). We demonstrate cases taking first derivatives of traits before calculating response

diversity (*e.g.*, Fig. 2b), thus producing *rates* from which we calculate response diversity (Fig. 2e).

However, it is also possible to estimate derivatives of rates (*e.g.*, Fig. 3d and Box 2), producing a

measure of sensitivity of those rates to environmental change, which can be ecologically meaningful

when measured over environmental gradients (Shyu & Caswell 2014). However, study designs often

estimate intrinsic rates of increase, for instance, under only two environmental contexts (*e.g.*,

disturbed versus in-tact habitats). In such cases, it may be more meaningful to estimate response

diversity based directly on the intrinsic rates of increase (rather than transforming them to second

derivatives of abundance). Individual use cases will differ, so we suggest response diversity studies

carefully consider whether response diversity should be measured based on traits, rates, or

sensitivities, depending on available data and study goals. Practical users of response diversity can

make use of the *mgcv* and *gratia* packages in *R* to fit GAMs and calculate derivatives, respectively

(Wood *et al.* 2016; Simpson 2021), and we provide *R* code to calculate response diversity for the

simulated case studies from Figures 2 and 3 (available through Github).

Using the framework for measuring response diversity we present here, one can measure

response diversity empirically in the laboratory or field. The key advantage of a conceptually simple

and empirically tractable measure of response diversity is that it unlocks a range of questions that can

be experimentally tested in controlled systems. For example, once the range of response trajectories

among species—or individuals/populations for cases of intraspecific response diversity—is

established, communities encompassing differing amounts of response diversity can be manually

assembled (Leary & Petchey 2009; Baskett *et al.* 2014). This, in turn, allows testing a variety of

questions with response diversity as the predictor variable, such as whether experimentally

assembled communities with higher response diversity are more stable in terms of, for example, the

temporal variability of total community biomass. In theory, performance-environment relationships

can be established in isolation and then communities assembled from individual species responses.

However, this approach neglects any interspecific interactions within multispecies communities. As

such, it is likely more relevant to model species responses (individually, using species-specific GAMs;

Github) within a multispecies community; thus allowing species-environment relationships to interact

with species-species relationships as they would in nature (Ives & Carpenter 2007; Armitage & Jones

2020).

Observational time series can also contain rich information on, for instance, community

dynamics or environmental responses (Ives *et al*. 2003; Ushio *et al*. 2018; Shoemaker *et al*. 2022).

Though methods for inferring causality from time series are emerging (*e.g.*, *empirical dynamic*

modelling, see Ushio *et al*. 2018; Ross *et al.* 2021b), such methods are often complex, and results

must be interpreted carefully. In principle, our approach for measuring response diversity can also be

applied to observational time series. For response diversity to be measurable in this way, time series

should comprise higher-level traits as described above (*e.g.*, measuring intrinsic rather than realised

rate of increase). Moreover, a high sampling resolution along environmental gradients is required to

accurately differentiate between linear and nonlinear performance-environment relationships; longer

or higher-resolution time series are better able to reveal nonlinearities in data, and response diversity

experiments should be designed with sampling resolution in mind to sufficiently capture all possible

forms of ecological responses. While the methods we describe here are useful for measuring multiple

dimensions of stability from experimental or observational time series, response diversity could

(perhaps more easily) also be measured over spatial gradients to assess spatial variation in the

stability and synchrony of functional properties such as primary productivity among habitat patches

(Loreau *et al.* 2003; Lamy *et al.* 2019; Wang *et al.* 2019, 2021).

Future directions for empirical studies of response diversity

The framework for measuring response diversity introduced above is conceptually simple and

applicable across contexts. Using this approach, response diversity can be easily measured in

empirical settings (see Box 2 for reanalysis of data from Leary & Petchey 2009), and experimentally

manipulated, providing clear advantages over the functional trait approach dominant in the literature

(Box 1). Owing to its simplicity, our framework can be extended in several promising directions (Fig.

4).

First, rather than using response diversity solely as a predictor variable for estimating stability

responses, a system's response diversity to a given stressor could be measured as a function of

environmental change; that is, its magnitude at any given environmental state can be assessed and

compared to other treatments or replicates across any environmental ranges of interest. Such an

approach also permits quantification of response diversity to multiple environmental stressors (Fig.

4a), which often occur simultaneously (Garnier *et al.* 2017; Bowler *et al.* 2020). The change in

response diversity to one environmental condition as a function of a second then allows study of co-

tolerance to multiple stressors, and stress-induced community sensitivity (see Vinebrooke *et al.*

2004). It should also be possible to measure multivariate response diversity in the same ways that

multivariate functional trait diversity is measured (Petchey & Gaston 2006).

Second, given the multidimensionality of the ecological stability concept (Pimm 1984;

Donohue *et al.* 2013; Hillebrand *et al.* 2018), the predictive capacity of response diversity can be

tested as a driver of different stability dimensions (that is, response diversity as a predictor of

variability, resistance, resilience *etc.*; Fig. 4b). Another extension of this framework would consider

the relative importance of each species to community-level response diversity by, for example,

measuring the median dissimilarity in performance-environment model slopes between each species

and all others in the community (Fig. 4c). In this case, the species with the greatest dissimilarity from

the others should be the largest contributor to response diversity, and thus be that species whose

loss from the community should most greatly reduce response diversity, and hence stability (provided

the two are mechanistically linked). Such an approach complements studies considering the

contribution of individual species to ecological responses and stability (*e.g.*, Donohue *et al.* 2017;

White *et al.* 2020; Ross *et al.* 2021a, 2022), in turn, aiding progress towards targeted conservation

management decisions based on the quantifiable importance of individual taxa. Likewise, the

similarity-based diversity measure we used to estimate response diversity is based on hill numbers,

with an initial value of $q = 0$ indicating a focus on species richness. However, by increasing q to

higher values, it should be straightforward to also incorporate relative species abundances (Chao *et*

al. 2014), and hence the potential role of dominance or rarity (*e.g.*, Sasaki & Lauenroth 2011).

Importantly, the subjects of a response diversity analysis need not be different species; individuals

can also exhibit diversity in their response traits, further underscoring the importance of intraspecific

variation in ecological responses to the environment (Barabás & D'Andrea 2016; Mimura *et al.* 2017;

Ross *et al.* 2017; Herrando-pérez *et al.* 2019). Finally, if response diversity can be measured from

observational time series, it should be conceptually straightforward to harness global datasets (*e.g.*,

BioTIME; Dornelas *et al.* 2018) to unveil macroecological patterns and hotspots of ecological stability

using response diversity (Capdevila *et al.* 2022; Fig. 4d).

Figure 4. Future extensions to the response diversity framework. (a) After measuring response diversity to an

environmental condition (*e.g.*, noise pollution), the change in response diversity to the first stressor could be

modelled against a second stressor (*e.g.*, light pollution) to assess whether communities exhibit positive or

negative co-tolerance to multiple stressors (Vinebrooke *et al.* 2004). It should then also be possible to develop

multivariate response diversity metrics capable of predicting simultaneous responses to concurrent

environmental stressors. (**b**) Ecological stability is a multidimensional concept (Donohue *et al.* 2013); multiple

stability metrics (*e.g.*, temporal variability [1], resistance [2], recovery time [3], and resilience [4]) can be

calculated from a single response variable to a known disturbance. In such cases, response diversity can be used

as a predictor of stability beyond just temporal variability, by extending tests of response diversity's predictive

capacity to other dimensions of stability (here resistance, recovery time, and resilience), and response diversity

may exhibit a different relationship with each stability dimension. (**c**) The relative contribution of species within

a community to response diversity can be captured by considering how different each species is from the others

under a given environmental condition. Here, we see the dark brown species is most different in its

performance-environment relationship, resulting in the highest relative contribution (importance) of this

species for response diversity. (**d**) With replicated global experiments or by exploiting global datasets of

assemblage time series (*e.g.*, Dornelas *et al.* 2018), it should be possible to construct macroecological

relationships for response diversity. For example, latitudinal response diversity gradients (coloured lines) should

tell us something about whether certain communities or biomes may be 'hotspots' of resilience to

environmental change.

That response diversity can be environment-dependent in cases where species respond

nonlinearly to the environment has considerable implications for empirical studies of response

diversity and for our understanding of ecological stability. Though threshold responses may be rare

and/or difficult to detect in experiments (Hillebrand *et al.* 2020), nonlinear responses to many

variables are common, such as Type II or III functional responses to prey availability (Holling 1966;

Daugaard *et al.* 2019), or light or temperature response curves (*e.g.*, Peek *et al.* 2002; Payne *et al.*

2016). Nonlinear responses to the environment may result in environment-dependencies in species

asynchrony (Shoemaker *et al.* 2022)—recently described using tail association methods (Ghosh *et al.*

2020a, 2020b; Walter *et al.* 2022)—in turn, producing environment-dependencies in ecosystem

multifunctionality and stability (Morin *et al.* 2014; Sasaki *et al.* 2019). Environment-dependent

response diversity may therefore explain inconsistencies in the reported effects of biodiversity on

ecological stability across studies (Jacquet *et al.* 2016; Pennekamp *et al.* 2018). Importantly, the

empirical framework we propose allows identification of the specific environmental conditions under

which response diversity should or should not be stabilising. For a nonlinear environmental response,

r(**e**)*,* one can examine the values of **e** (the 'environment') for which the first derivatives of species-

environment relationships, r'(e)*,* including their confidence regions, span zero. In other words, our

framework makes it conceptually and practically simple to identify *a priori* hypotheses regarding the

stabilising effect of response diversity based on environmental conditions (Box 2). This should

complement existing studies of the impact of environmental conditions on ecological stability (*e.g.*,

Hillebrand *et al.* 2018; Ross *et al.* 2022). Moreover, in cases where response diversity varies as a

function of the environment (that is, where species-environment relationships are nonlinear), it is

possible to manipulate response diversity without altering community composition. It might then be

possible to identify response diversity as a stabilising mechanism by comparing the stability of

aggregate ecosystem properties between communities of the same species under different (fixed)

environmental contexts. Such an approach sidesteps the challenges associated with causal inference

when comparing communities of different species; by controlling species identities, there are fewer

potential confounding factors driving community-level responses (though environmental context is

added as an extrinsic driver). For example, one could measure the diversity-stability relationship for a

plant community with known performance responses to drought under drought and non-drought

conditions to mechanistically consider how drought relate to stability. There is much still to learn

about context-dependent ecological stability as it relates to nonlinear performance-environment

relationships and environment-dependent response diversity, and we suggest this as a promising

avenue of future work.

Since the observation that diversity alone is insufficient to explain the stabilising effect of

species richness on aggregate ecosystem properties (May 1973; Ives *et al.* 1999), ecologists have

been searching for a general mechanism explaining the insurance effect of biodiversity across

contexts (Yachi & Loreau 1999). Here we proposed response diversity as a promising but

understudied mechanism for predicting ecological responses to environmental conditions (Elmqvist *et*

al. 2003; Nyström 2006; Mori *et al.* 2013). We found that most empirical studies of response diversity

measure the diversity of functional response traits (*e.g.*, Laliberté *et al.* 2010; Morel *et al.* 2020).

However, given the challenges associated with reliably producing functional response diversity

estimates from trait data (Box 1), we suggest measuring response diversity as the variation in

responses to environmental conditions (Leary & Petchey 2009; Thibaut *et al.* 2012)—unless using

functional traits empirically demonstrated to encode sufficient information about responses to

environmental change (*e.g.*, Kandlikar *et al.* 2022). The framework we propose herein provides a

direct, empirically tractable, and conceptually simple measure of response diversity that should

facilitate robust empirical tests of the diversity-stability relationship across taxa and environments.

Such tests have the potential to identify the long sought-after mechanisms underpinning

biodiversity's role in buffering ecosystems against environmental change.

Box 1. Response trait diversity: a reliable measure?

Functional diversity measures the extent of functional differences among species in a given

community (Tilman 2001), and myriad functional diversity metrics are now used to estimate

ecosystem functioning using functional traits, which represent a species' contribution to functioning

(Díaz & Cabido 2001; Petchey & Gaston 2006; Cadotte *et al.* 2011). Given their effect on ecosystem

processes and functions, such traits are now termed *effect* traits, while *response* traits aim to capture

information about how a species responds to environmental change (Suding *et al.* 2008; Oliver *et al.*

2015), though such information is not always well captured by putative functional traits (Bartomeus

et al. 2017). Examples of response traits include life-form and seed mass in plants (Sasaki *et al.* 2019),

aquatic stage and reproductive strategy in freshwater macroinvertebrates (Thornhill *et al.* 2018), and

clutch size and thermal maximum in birds (Hordley *et al.* 2021). Response trait diversity is then

typically measured as the functional dispersion (FDis) of the response trait space (Laliberté *et al.*

2010; Rader *et al.* 2013), or sometimes using the closely related Rao's Quadratic Entropy (RaoQ) index

(Chillo *et al.* 2011; Correia *et al.* 2018). Some studies alternatively group species into functional

response groups (*e.g.*, Joseph *et al.* 2014; Schweiger *et al.* 2015), where species within a functional

group are posited to respond similarly to environmental change based on their response traits, but

this approach ignores a suite of functional differences between species; functional grouping is rarely

ecologically meaningful (Petchey & Gaston 2006).

Considerable challenges remain when quantifying functional diversity. For example, there is

still no consensus on which traits are 'functional' in a range of systems (Violle *et al.* 2007), and many

traits can be highly correlated with, for example, body size (Koziowski & Weiner 1997). When

attempting to measure response trait diversity, it can be difficult to delineate responses from effects;

many traits act both as response and effect traits (Hooper *et al.* 2005; Suding *et al.* 2008). Moreover,

it is often practically challenging to measure functional traits in the field, particularly for mobile

organisms (Pakeman & Quested 2007; Moretti *et al.* 2017). When measuring functional diversity from

species traits, there is no single best metric for all scenarios, as most functional diversity indices

capture different properties of the functional differences within a community (Villéger *et al.* 2008;

Schleuter *et al.* 2010). Other methodological considerations include whether and how to abundance-

weight functional diversity indices to account for rare species (Lavorel *et al.* 2008), and how to handle

intraspecific trait variation given its ecological importance (de Bello *et al.* 2011; Albert *et al.* 2012;

Ross *et al.* 2017). Together, these challenges hinder the empirical quantification of response diversity

using functional response traits. As such, we suggest caution in measuring response trait diversity

unless using well-defined and empirically tested response traits (*e.g.*, Kandlikar *et al.* 2022).

Box 2. Re-analysing response diversity for experimental aquatic communities

Here, we demonstrate the ease with which our statistical framework can be applied to

empirical data, by reanalysing the variation in performance-environment relationships among aquatic

microorganisms first presented by Leary and Petchey (2009). The original study assembled ciliate

communities comprising four species (*Colpidium striatum*, *Loxocephallus* sp., *Paramecium caudatum*,

and *Tetrahymena pyriformis*) with measured responses to temperature across five levels (18, 20, 22,

24, and 26°C; Box Figure 1a,b). The authors then measured response diversity as the correlation

between pairwise species responses to temperature, measured using two higher-level traits—intrinsic

rate of increase (r) and carrying capacity ($\log_{10} K$)—and used this response diversity as a predictor of,

among other things, community biomass (Leary & Petchey 2009). Here, we took a different approach

to measure response diversity. We modelled each species' performance (r or $\log_{10}K$) individually using

Generalized Additive Models (GAMs), and calculated the first derivates of each performance-

temperature relationship (Box Fig. 1c,d); here producing *second* derivatives of abundance that

indicate sensitivity to environmental change (Shyu & Caswell 2014). We then estimated the two

dimensions of response diversity described above; we measured the dissimilarity of these first

derivatives (Leinster & Cobbold 2012) and their sign-sensitivity (that is, the extent to which the first

derivatives span zero for a given environmental context). In doing this as a function of temperature,

we produced temperature-dependent estimates of response diversity, which was highest for r at

around 23.5°C when performance-environment responses were divergent and zero-spanning (Box

Fig. 1e), and highest for $\log_{10}K$ at the maximum temperature as the carrying capacity of *C. striatum*

declines while the other species remain fairly constant (suggesting *C. striatum* has a disproportionate

contribution to dissimilarity here). Note, however, that the sign-sensitivity of the derivatives is low at

this temperature, so we might predict any community-level properties driven by carrying capacity to

be unstable at high temperatures (Box Fig. 1f). Such temperature-dependent response diversity

values provide an added layer of information—and hence predictive capacity—compared to the

original analyses based on correlated performance-environment responses (Leary & Petchey 2009).

550
Box Figure 1. Re-   herein. Using
empirical data from Leary & Petchey (2009), we measured performance-environment relationships for four
aquatic ciliate species [coloured lines] as (**a**) the intrinsic rate of increase [*r*] and (**b**) the carrying capacity [$\log_{10}K$]
along a temperature gradient (18-26°C). Data points are from a single population (see Figs. 1a and 2a in Leary &
Petchey 2009). (**c**,**d**) We calculated the first derivatives of the performance-temperature relationships, which
change as a function of temperature due to nonlinear responses. (**e**,**f**) We then measured the similarity-based
diversity and sign-sensitivity of the derivatives of all species in the community for both (**e**) *r* and (**f**) $\log_{10}K$. (**e**) For
r we found that both indices peaked around 22-24°C. (**f**) For $\log_{10}K$, dissimilarity peaked at the upper
temperature limit, as the response of one species (*C. striatum*) diverged from the others (**b**,**d**), while the sign-
sensitive index was mostly zero across the temperature range. Dashed lines indicate zero in panels **c** and **d**, and
the median value of response diversity along the X-axis in panels **e** and **f**.

Acknowledgements

We thank Dan Leary for generating the empirical data we reanalysed in this study (Box 2). Thanks also
to Jean-François Arnoldi and Ian Donohue for earlier discussion. This study was supported by a
SHINKA collaboration grant awarded to SRP-JR by the Okinawa Institute of Science and Technology

Graduate University (OIST). SRP-JR and DWA were supported by subsidy funding to OIST. OLP was

supported by the University of Zurich Research Priority Programme in Global Change and Biodiversity

(URPP GCB) and by SNF project 310030_188431. TS was supported by a Fostering Joint International

Research A (no. 19KK0393) from the Japanese Ministry of Education, Culture, Sports, Science and

Technology.

Data availability statement

All data and R code necessary to reproduce the analyses and figures will be available in the Zenodo

digital repository (and assigned a DOI) upon acceptance. For now, the GitHub repository can be

accessed directly at:

https://github.com/opetchey/response_diversity_how_to_measure/releases/tag/v1.0-review

References

Albert, C.H., de Bello, F., Boulangeat, I., Pellet, G., Lavorel, S. & Thuiller, W. (2012). On the importance of

581   intraspecific variability for the quantification of functional diversity. *Oikos*, 121, 116–126.

582   Altomare, M., Vasconcelos, H.L., Raymundo, D., Lopes, S., Vale, V. & Prado-Junior, J. (2021). Assessing the fire

583   resilience of the savanna tree component through a functional approach. *Acta Oecologica*, 111,

584   103728.

585   Armitage, D.W. & Jones, S.E. (2019). Negative frequency-dependent growth underlies the stable coexistence of

586   two cosmopolitan aquatic plants. *Ecology*, 100, e02657.

587   Armitage, D.W. & Jones, S.E. (2020). Coexistence barriers confine the poleward range of a globally distributed

588   plant. *Ecol. Lett.*, 23, 1838–1848.

589   Arnoldi, J.F., Loreau, M. & Haegeman, B. (2019). The inherent multidimensionality of temporal variability: how

590   common and rare species shape stability patterns. *Ecol. Lett.*, 22, 1557–1567.

591   Barabás, G. & D'Andrea, R. (2016). The effect of intraspecific variation and heritability on community pattern

592   and robustness. *Ecol. Lett.*, 19, 977–986.

593   Bartomeus, I., Cariveau, D.P., Harrison, T. & Winfree, R. (2017). On the inconsistency of pollinator species traits

594   for predicting either response to land-use change or functional contribution. *Oikos*, 127, 306–315.

595   Bartomeus, I., Park, M.G., Gibbs, J., Danforth, B.N., Lakso, A.N. & Winfree, R. (2013). Biodiversity ensures plant–

596   pollinator phenological synchrony against climate change. *Ecol. Lett.*, 16, 1331–1338.

597   Baskett, M.L., Fabina, N.S. & Gross, K. (2014). Response diversity can increase ecological resilience to

598   disturbance in coral reefs. *Am. Nat.*, 184, E16–E31.

599   de Bello, F., Lavorel, S., Albert, C.H., Thuiller, W., Grigulis, K., Dolezal, J., *et al.* (2011). Quantifying the relevance

600   of intraspecific trait variability for functional diversity. *Methods Ecol. Evol.*, 2, 163–174.

601   Berdugo, M., Delgado-Baquerizo, M., Soliveres, S., Hernández-Clemente, R., Zhao, Y., Gaitán, J.J., *et al.* (2020).

602   Global ecosystem thresholds driven by aridity. *Science*, 367, 787–790.

Bowler, D.E., Bjorkman, A.D., Dornelas, M., Myers-Smith, I.H., Navarro, L.M., Niamir, A., *et al.* (2020). Mapping

human pressures across the planet uncovers anthropogenic threat complexes. *People Nat.*, 2, 380–

394.

Burrows, R.C., Wancio, D., Levitt, P. & Lillien, L. (1997). Response Diversity and the Timing of Progenitor Cell

Maturation Are Regulated by Developmental Changes in EGFR Expression in the Cortex. *Neuron*, 19,

251–267.

Cadotte, M.W., Carscadden, K. & Mirotchnick, N. (2011). Beyond species: Functional diversity and the

maintenance of ecological processes and services. *J. Appl. Ecol.*, 48, 1079–1087.

Capdevila, P., Noviello, N., McRae, L., Freeman, R. & Clements, C.F. (2022). Global patterns of resilience decline

in vertebrate populations. *Ecol. Lett.*, 25, 240–251.

Cariveau, D.P., Williams, N.M., Benjamin, F.E. & Winfree, R. (2013). Response diversity to land use occurs but

does not consistently stabilise ecosystem services provided by native pollinators. *Ecol. Lett.*, 16, 903–

911.

Chao, A., Gotelli, N.J., Hsieh, T.C., Sander, E.L., Ma, K.H., Colwell, R.K., *et al*. (2014). Rarefaction and

extrapolation with Hill numbers: a framework for sampling and estimation in species diversity studies.

Ecol. Monogr., 84, 45–67.

Chesson, P. (2000). Mechanisms of Maintenance of Species Diversity. *Annu. Rev. Ecol. Syst.*, 31, 343–366.

Chillo, V., Anand, M. & Ojeda, R.A. (2011). Assessing the Use of Functional Diversity as a Measure of Ecological

Resilience in Arid Rangelands. *Ecosystems*, 14, 1168–1177.

Correia, D.L.P., Raulier, F., Bouchard, M. & Filotas, É. (2018). Response diversity, functional redundancy, and

post-logging productivity in northern temperate and boreal forests. *Ecol. Appl.*, 28, 1282–1291.

Craven, D., Eisenhauer, N., Pearse, W.D., Hautier, Y., Isbell, F., Roscher, C., *et al.* (2018). Multiple facets of

biodiversity drive the diversity–stability relationship. *Nat. Ecol. Evol.*, 2, 1579–1587.

Craven, D., Filotas, E., Angers, V.A. & Messier, C. (2016). Evaluating resilience of tree communities in fragmented

landscapes: linking functional response diversity with landscape connectivity. *Divers. Distrib.*, 22, 505–

518.

Daugaard, U., Petchey, O.L. & Pennekamp, F. (2019). Warming can destabilize predator–prey interactions by

shifting the functional response from Type III to Type II. *J. Anim. Ecol.*, 88, 1575–1586.

Díaz, S. & Cabido, M. (2001). Vive la différence: Plant functional diversity matters to ecosystem processes.

Trends Ecol. Evol., 16, 646–655.

Döbert, T.F., Webber, B.L., Sugau, J.B., Dickinson, K.J.M. & Didham, R.K. (2017). Logging increases the functional

and phylogenetic dispersion of understorey plant communities in tropical lowland rain forest. *J. Ecol.*,

105, 1235–1245.

Donohue, I., Petchey, O.L., Kéfi, S., Génin, A., Jackson, A.L., Yang, Q., *et al.* (2017). Loss of predator species, not

intermediate consumers, triggers rapid and dramatic extinction cascades. *Glob. Change Biol.*, 23, 2962–

2972.

Donohue, I., Petchey, O.L., Montoya, J.M., Jackson, A.L., Mcnally, L., Viana, M., *et al.* (2013). On the

dimensionality of ecological stability. *Ecol. Lett.*, 16, 421–429.

Dornelas, M., Antão, L.H., Moyes, F., Bates, A.E., Magurran, A.E., Adam, D., *et al.* (2018). BioTIME: A database of
642   biodiversity time series for the Anthropocene. *Glob. Ecol. Biogeogr.*, 27, 760–786.

643   Elmqvist, T., Folke, C., Nyström, M., Peterson, G., Bengtsson, J., Walker, B., *et al.* (2003). Response diversity,
644   ecosystem change, and resilience. *Front. Ecol. Environ.*, 1, 488–494.

645   Elton, C.S. (1958). *Ecology of Invasions by Animals and Plants*. Chapman & Hall, London.

646   Fauchald, P., Skov, H., Skern-Mauritzen, M., Hausner, V.H., Johns, D. & Tveraa, T. (2011). Scale-dependent
647   response diversity of seabirds to prey in the North Sea. *Ecology*, 92, 228–239.

648   Ford, H., Healey, J.R., Markesteijn, L. & Smith, A.R. (2018). How does grazing management influence the
649   functional diversity of oak woodland ecosystems? A plant trait approach. *Agric. Ecosyst. Environ.*, 258,
650   154–161.

651   Fraterrigo, J.M. & Rusak, J.A. (2008). Disturbance-driven changes in the variability of ecological patterns and
652   processes. *Ecol. Lett.*, 11, 756–770.

653   Fründ, J., Zieger, S.L. & Tscharntke, T. (2013). Response diversity of wild bees to overwintering temperatures.
654   *Oecologia*, 173, 1639–1648.

655   Garnier, A., Pennekamp, F., Lemoine, M. & Petchey, O.L. (2017). Temporal scale dependent interactions
656   between multiple environmental disturbances in microcosm ecosystems. *Glob. Change Biol.*, 23, 5237–
657   5248.

658   Ghosh, S., Sheppard, L.W., Reid, P.C. & Reuman, D. (2020a). A new approach to interspecific synchrony in
659   population ecology using tail association. *Ecol. Evol.*, 10, 12764–12776.

660   Ghosh, S., Sheppard, L.W. & Reuman, D.C. (2020b). Tail associations in ecological variables and their impact on
661   extinction risk. *Ecosphere*, 11, e03132.

662   Gonzalez, A. & Loreau, M. (2009). The Causes and Consequences of Compensatory Dynamics in Ecological
663   Communities. *Annu. Rev. Ecol. Evol. Syst.*, 40, 393–414.

664   Grainger, T.N., Levine, J.M. & Gilbert, B. (2019). The Invasion Criterion: A Common Currency for Ecological
665   Research. *Trends Ecol. Evol.*, 34, 925–935.

666   Hammill, E., Kratina, P., Vos, M., Petchey, O.L. & Anholt, B.R. (2015). Food web persistence is enhanced by non-
667   trophic interactions. *Oecologia*, 178, 549–556.

668   Hastie, T. & Tibshirani, R. (1987). Generalized additive models: some applications. *J. Am. Stat. Assoc.*, 82, 371–
669   386.

670   Hector, A.A., Hautier, Y., Saner, P., Wacker, L., Bagchi, R., Joshi, J., *et al.* (2015). General stabilizing effects of
671   plant diversity on grassland productivity through population asynchrony and overyielding. *Ecology*, 91,
672   2213–2220.

673   Herrando-pérez, S., Vieites, D.R., Monasterio, F.F.C., Belliure, J., Chown, S.L., Beukema, W., *et al.* (2019).
674   Intraspecific variation in lizard heat tolerance alters estimates of climate impact. *J. Anim. Ecol.*, 88,
675   247–257.

676   Hillebrand, H., Donohue, I., Harpole, W.S., Hodapp, D., Kucera, M., Lewandowska, A.M., *et al.* (2020). Thresholds
677   for ecological responses to global change do not emerge from empirical data. *Nat. Ecol. Evol.*, 4, 1502–
678   1509.

Hillebrand, H., Langenheder, S., Lebret, K., Lindström, E., Östman, Ö. & Striebel, M. (2018). Decomposing
multiple dimensions of stability in global change experiments. *Ecol. Lett.*, 21, 21–30.

Hlatky, L.R., Hahnfeldt, P. & Sachs, R.K. (1994). Influence of time-dependent stochastic heterogeneity on the
radiation response of a cell population. *Math. Biosci.*, 122, 201–220.

Holling, C.S. (1966). The Functional Response of Invertebrate Predators to Prey Density. *Mem. Entomol. Soc.*
Can., 98, 5–86.

Hooper, D.U., Chapin, F.S., Ewel, J.J., Hector, A., Inchausti, P., Lavorel, S., *et al.* (2005). Effects of biodiversity on
ecosystem functioning: A consensus of current knowledge. *Ecol. Monogr.*, 75, 3–35.

Hordley, L.A., Gillings, S., Petchey, O.L., Tobias, J.A. & Oliver, T.H. (2021). Diversity of response and effect traits
provides complementary information about avian community dynamics linked to ecological function.
Funct. Ecol., 35, 1938–1950.

Ishii, R. & Masahiko, M. (2001). Coexistence induced by pollen limitation in flowering-plant species. *Proc. R. Soc.*
Lond. B Biol. Sci., 268, 579–585.

Ives, A.R. & Carpenter, S.R. (2007). Stability and diversity of ecosystems. *Science*, 317, 58–62.

Ives, A.R., Dennis, B., Cottingham, K.L. & Carpenter, S.R. (2003). Estimating community stability and ecological
interactions from time‐series data. *Ecol. monogr.*, 73, 301–330.

Ives, A.R., Gross, K. & Klug, J.L. (1999). Stability and Variability in Competitive Communities. *Science*, 286, 542–
544.

Jacquet, C., Moritz, C., Morissette, L., Legagneux, P., Massol, F., Archambault, P., *et al.* (2016). No complexity–
stability relationship in empirical ecosystems. *Nat. Commun.*, 7, 12573.

Joseph, G.S., Seymour, C.L., Cumming, G.S., Cumming, D.H.M. & Mahlangu, Z. (2014). Termite Mounds Increase
Functional Diversity of Woody Plants in African Savannas. *Ecosystems*, 17, 808–819.

Kahiluoto, H., Kaseva, J., Olesen, J.E., Kersebaum, K.C., Ruiz-Ramos, M., Gobin, A., *et al.* (2019). Genetic response
diversity to provide yield stability of cultivar groups deserves attention. *Proc. Natl. Acad. Sci. U.S.A.*,
166, 10627–10629.

Kandlikar, G.S., Kleinhesselink, A.R. & Kraft, N.J.B. (2022). Functional traits predict species responses to
environmental variation in a California grassland annual plant community. *J. Ecol.* [In press].

Kondoh, M. (2003). Foraging Adaptation and the Relationship Between Food-Web Complexity and Stability.
Science, 299, 1388–1391.

Koziowski, J. & Weiner, J. (1997). Interspecific allometries Are by-products of body size optimization. *Am. Nat.*,
149, 352–380.

Laliberté, E., Wells, J.A., Declerck, F., Metcalfe, D.J., Catterall, C.P., Queiroz, C., *et al.* (2010). Land-use
intensification reduces functional redundancy and response diversity in plant communities. *Ecol. Lett.*,
13, 76–86.

Lamy, T., Wang, S., Renard, D., Lafferty, K.D., Reed, D.C. & Miller, R.J. (2019). Species insurance trumps spatial
insurance in stabilizing biomass of a marine macroalgal metacommunity. *Ecology*, 100, e02719.

Lavorel, S., Grigulis, K., McIntyre, S., Williams, N.S.G., Garden, D., Dorrough, J., *et al.* (2008). Assessing functional
diversity in the field - Methodology matters! *Funct. Ecol.*, 22, 134–147.

Leary, D.J. & Petchey, O.L. (2009). Testing a biological mechanism of the insurance hypothesis in experimental
718   aquatic communities. *J. Anim. Ecol.*, 78, 1143–1151.

719   Leinster, T. & Cobbold, C.A. (2012). Measuring diversity: the importance of species similarity. *Ecology*, 93, 477–
720   489.

721   Loreau, M. (2010). Stability and Complexity of Ecosystems: New Perspectives on an Old Debate. In: *From
722   Populations to Ecosystems*, Theoretical Foundations for a New Ecological Synthesis (MPB-46).
723   Princeton University Press, pp. 123–163.

724   Loreau, M., Barbier, M., Filotas, E., Gravel, D., Isbell, F., Miller, S.J., *et al.* (2021). Biodiversity as insurance: from
725   concept to measurement and application. *Biol. Rev.*, 96, 2333–2354.

726   Loreau, M. & de Mazancourt, C. (2013). Biodiversity and ecosystem stability: A synthesis of underlying
727   mechanisms. *Ecol. Lett.*, 16, 106–115.

728   Loreau, M., Mouquet, N., Gonzalez, A. & Mooney, H.A. (2003). Biodiversity as spatial insurance in
729   heterogeneous landscapes. *Proc. Natl. Acad. Sci. U.S.A.*, 100, 12765–12770.

730   MacArthur, R. (1955). Fluctuations of Animal Populations and a Measure of Community Stability. *Ecology*, 36,
731   533–536.

732   Malyshev, A.V., Arfin Khan, M.A.S., Beierkuhnlein, C., Steinbauer, M.J., Henry, H.A.L., Jentsch, A., *et al.* (2016).
733   Plant responses to climatic extremes: within-species variation equals among-species variation. *Glob.
734   Change Biol.*, 22, 449–464.

735   Mandle, L. & Ticktin, T. (2015). Moderate land use changes plant functional composition without loss of
736   functional diversity in India's Western Ghats. *Ecol. Appl.*, 25, 1711–1724.

737   May, R.M. (1973). *Stability and complexity in model ecosystems*. Princeton University Press.

738   de Mazancourt, C., Isbell, F., Larocque, A., Berendse, F., De Luca, E., Grace, J.B., *et al.* (2013). Predicting
739   ecosystem stability from community composition and biodiversity. *Ecol. Lett.*, 16, 617–625.

740   McCann, K., Hastings, A. & Huxel, G.R. (1998). Weak trophic interactions and the balance of nature. *Nat. 1998
741   3956704*, 395, 794–798.

742   McCann, K.S. (2000). The diversity-stability debate. *Nature*, 405, 228–233.

743   McCann, M.J. (2016). Response diversity of free-floating plants to nutrient stoichiometry and temperature:
744   growth and resting body formation. *PeerJ*, 4, e1781.

745   Mimura, M., Yahara, T., Faith, D.P., Vázquez-Domínguez, E., Colautti, R.I., Araki, H., *et al.* (2017). Understanding
746   and monitoring the consequences of human impacts on intraspecific variation. *Evol. Appl.*, 10, 121–
747   139.

748   Moore, J.W. & Olden, J.D. (2017). Response diversity, nonnative species, and disassembly rules buffer
749   freshwater ecosystem processes from anthropogenic change. *Glob. Change Biol.*, 23, 1871–1880.

750   Morel, L., Barbe, L., Jung, V., Clément, B., Schnitzler, A. & Ysnel, F. (2020). Passive rewilding may (also) restore
751   phylogenetically rich and functionally resilient forest plant communities. *Ecol. Appl.*, 30, e02007.

752   Moretti, M., Dias, A.T.C., de Bello, F., Altermatt, F., Chown, S.L., Azcárate, F.M., *et al.* (2017). Handbook of
753   protocols for standardized measurement of terrestrial invertebrate functional traits. *Funct. Ecol.*, 31,
754   558–567.

Mori, A.S., Furukawa, T. & Sasaki, T. (2013). Response diversity determines the resilience of ecosystems to
environmental change. *Biol. Rev.*, 88, 349–364.

Morin, X., Fahse, L., de Mazancourt, C., Scherer-Lorenzen, M. & Bugmann, H. (2014). Temporal stability in forest
productivity increases with tree diversity due to asynchrony in species dynamics. *Ecol. Lett.*, 17, 1526–
1535.

Mougi, A. & Kondoh, M. (2012). Diversity of Interaction Types and Ecological Community Stability. *Science*, 337,
349–351.

Mumme, S., Jochum, M., Brose, U., Haneda, N.F. & Barnes, A.D. (2015). Functional diversity and stability of
litter-invertebrate communities following land-use change in Sumatra, Indonesia. *Biol. Conserv.*, 191,
750–758.

Nyström, M. (2006). Redundancy and Response Diversity of Functional Groups: Implications for the Resilience of
Coral Reefs. *AMBIO J. Hum. Environ.*, 35, 30–35.

Odum, E.P. (1953). *Fundamentals of Ecology*. Saunders, Philadelphia.

Oliver, T.H., Heard, M.S., Isaac, N.J.B., Roy, D.B., Procter, D., Eigenbrod, F., *et al.* (2015). Biodiversity and
Resilience of Ecosystem Functions. *Trends Ecol. Evol.*, 30, 673–684.

Pakeman, R.J. & Quested, H.M. (2007). Sampling plant functional traits: What proportion of the species need to
be measured? *Appl. Veg. Sci.*, 10, 91–96.

Payne, N.L., Smith, J.A., van der Meulen, D.E., Taylor, M.D., Watanabe, Y.Y., Takahashi, A., *et al.* (2016).
Temperature dependence of fish performance in the wild: links with species biogeography and
physiological thermal tolerance. *Funct. Ecol.*, 30, 903–912.

Peek, M.S., Russek-Cohen, E., Wait, A.D. & Forseth, I.N. (2002). Physiological response curve analysis using
nonlinear mixed models. *Oecologia*, 132, 175–180.

Pennekamp, F., Pontarp, M., Tabi, A., Altermatt, F., Alther, R., Choffat, Y., *et al.* (2018). Biodiversity increases
and decreases ecosystem stability. *Nature*, 563, 109–112.

Petchey, O.L. (2003). Integrating methods that investigate how complementarity influences ecosystem
functioning. *Oikos*, 101, 323–330.

Petchey, O.L. & Gaston, K.J. (2006). Functional diversity: Back to basics and looking forward. *Ecol. Lett.*, 9, 741–
758.

Pimm, S.L. (1984). The complexity and stability of ecosystems. *Nature*, 307, 321–326.

Pimm, S.L. & Lawton, J.H. (1978). On feeding on more than one trophic level. *Nature*, 275, 542–544.

Rader, R., Reilly, J., Bartomeus, I. & Winfree, R. (2013). Native bees buffer the negative impact of climate
warming on honey bee pollination of watermelon crops. *Glob. Change Biol.*, 19, 3103–3110.

Reilly, J.R., Artz, D.R., Biddinger, D., Bobiwash, K., Boyle, N.K., Brittain, C., *et al.* (2020). Crop production in the
USA is frequently limited by a lack of pollinators. *Proc. R. Soc. B Biol. Sci.*, 287, 20200922.

Ross, S.R.P.-J., Arnoldi, J.-F., Loreau, M., White, C.D., Stout, J.C., Jackson, A.L., *et al.* (2021a). Universal scaling of
robustness of ecosystem services to species loss. *Nat. Commun.*, 12, 5167.

Ross, S.R.P.-J., García Molinos, J., Okuda, A., Johnstone, J., Atsumi, K., Futamura, R., *et al.* (2022). Predators

mitigate the destabilising effects of heatwaves on multitrophic stream communities. *Glob. Change*

Biol., 28, 403–416.

Ross, S.R.P.-J., Hassall, C., Hoppitt, W.J.E., Edwards, F.A., Edwards, D.P. & Hamer, K.C. (2017). Incorporating

intraspecific trait variation into functional diversity: Impacts of selective logging on birds in Borneo.

Methods Ecol. Evol., 8, 1499–1505.

Ross, S.R.P.-J., Suzuki, Y., Kondoh, M., Suzuki, K., Villa Martín, P. & Dornelas, M. (2021b). Illuminating the

intrinsic and extrinsic drivers of ecological stability across scales. *Ecol. Res.*, 36, 364–378.

Sasaki, T. & Lauenroth, W.K. (2011). Dominant species, rather than diversity, regulates temporal stability of

plant communities. *Oecologia*, 166, 761–768.

Sasaki, T., Lu, X., Hirota, M. & Bai, Y. (2019). Species asynchrony and response diversity determine

multifunctional stability of natural grasslands. *J. Ecol.*, 107, 1862–1875.

Schleuter, D., Daufresne, M., Massol, F. & Argillier, C. (2010). A user's guide to functional diversity indices. *Ecol.*

Monogr., 80, 469–484.

Schnabel, F., Liu, X., Kunz, M., Barry, K.E., Bongers, F.J., Bruelheide, H., *et al.* (2021). Species richness stabilizes

productivity via asynchrony and drought-tolerance diversity in a large-scale tree biodiversity

experiment. *Sci. Adv.*, 7, eabk1643.

Schweiger, A.H., Audorff, V. & Beierkuhnlein, C. (2015). The acid taste of climate change: 20th century

acidification is re-emerging during a climatic extreme event. *Ecosphere*, 6, art94.

Shade, A. (2017). Diversity is the question, not the answer. *ISME J.*, 11, 1–6.

Shoemaker, L.G., Hallett, L.M., Zhao, L., Reuman, D.C., Wang, S., Cottingham, K.L., *et al.* (2022). The long and the

short of it: Mechanisms of synchronous and compensatory dynamics across temporal scales. *Ecology*,

e3650.

Shyu, E., & Caswell, H. (2014). Calculating second derivatives of population growth rates for ecology and

evolution. *Methods Ecol. Evol.*, 5, 473–482.

Simpson, G.L. (2021). gratia: Graceful 'ggplot'-Based Graphics and Other Functions for GAMs Fitted Using 'mgcv'.

R package version 0.6.0. https://CRAN.R-project.org/package=gratia

Sohlström, E.H., Archer, L.C., Gallo, B., Jochum, M., Kordas, R.L., Rall, B.C., *et al.* (2021). Thermal acclimation

increases the stability of a predator–prey interaction in warmer environments. *Glob. Change Biol.*, 27,

3765–3778.

Spasojevic, M.J., Bahlai, C.A., Bradley, B.A., Butterfield, B.J., Tuanmu, M.-N., Sistla, S., *et al.* (2016). Scaling up the

diversity–resilience relationship with trait databases and remote sensing data: the recovery of

productivity after wildfire. *Glob. Change Biol.*, 22, 1421–1432.

Stavert, J.R., Pattemore, D.E., Gaskett, A.C., Beggs, J.R. & Bartomeus, I. (2017). Exotic species enhance response

diversity to land-use change but modify functional composition. *Proc. R. Soc. B Biol. Sci.*, 284,

20170788.

Suding, K.N., Lavorel, S., Chapin, F.S., Cornelissen, J.H.C., Díaz, S., Garnier, E., *et al.* (2008). Scaling environmental
change through the community-level: A trait-based response-and-effect framework for plants. *Glob.*
Change Biol., 14, 1125–1140.

Thebault, E. & Fontaine, C. (2010). Stability of Ecological Communities and the Architecture of Mutualistic and
Trophic Networks. *Science*, 329, 853–856.

Thibaut, L.M., Connolly, S.R. & Sweatman, H.P.A. (2012). Diversity and stability of herbivorous fishes on coral
reefs. *Ecology*, 93, 891–901.

Thornhill, I., Biggs, J., Hill, M., Briers, R., Gledhill, D., Wood, P., *et al.* (2018). The functional response and
resilience in small waterbodies along land-use and environmental gradients. *Glob. Change Biol.*, 24,
3079–3092.

Tilman, D. (2001). Functional Diversity. In: *Encyclopedia of Biodiversity*. Elsevier Inc., pp. 587–596.

Tilman, D., Lehman, C.L. & Bristow, C.E. (1998). Diversity-stability relationships: statistical inevitability or
ecological consequence? *Am. Nat.*, 151, 277–82.

Ushio, M., Hsieh, C.H., Masuda, R., Deyle, E.R., Ye, H., Chang, C.W., *et al*. (2018). Fluctuating interaction network
and time-varying stability of a natural fish community. *Nature*, 554, 360–363.

Villéger, S., Mason, N.W.H. & Mouillot, D. (2008). New multidimensional functional diversity indices for a
multifaceted framwork in functional ecology. *Ecology*, 89, 2290–2301.

Vinebrooke, R.D., Cottingham, K.L., Norberg, J., Scheffer, M., Dodson, S.I., Maberly, S.C., *et al.* (2004). Impacts of
multiple stressors on biodiversity and ecosystem functioning: The role of species co-tolerance. *Oikos*,
104, 451–457.

Violle, C., Navas, M.L., Vile, D., Kazakou, E., Fortunel, C., Hummel, I., *et al.* (2007). Let the concept of trait be
functional! *Oikos*, 116, 882–892.

Vogel, A., Manning, P., Cadotte, M.W., Cowles, J., Isbell, F., Jousset, A.L.C., *et al.* (2019). *Lost in trait space:*
species-poor communities are inflexible in properties that drive ecosystem functioning. *Adv. Ecol. Res.*
1st edn. Elsevier Ltd.

Walker, B., Kinzig, A. & Langridge, J. (1999). Plant Attribute Diversity, Resilience, and Ecosystem Function: The
Nature and Significance of Dominant and Minor Species. *Ecosystems*, 2, 95–113.

Walter, J.A., Castorani, M.C.N., Bell, T.W., Sheppard, L.W., Cavanaugh, K.C. & Reuman, D.C. (2022). Tail-
dependent spatial synchrony arises from nonlinear driver–response relationships. *Ecol. Lett.*,
ele.13991.

Wang, S. & Loreau, M. (2016). Biodiversity and ecosystem stability across scales in metacommunities. *Ecol. Lett.*,
19, 510–518.

Wang, S., Loreau, M., de Mazancourt, C., Isbell, F., Beierkuhnlein, C., Connolly, J., *et al.* (2021). Biotic
homogenization destabilizes ecosystem functioning by decreasing spatial asynchrony. *Ecology*, 102, 1–
10.

Wang, Y., Cadotte, M.W., Chen, Y., Fraser, L.H., Zhang, Y., Huang, F., *et al.* (2019). Global evidence of positive
biodiversity effects on spatial ecosystem stability in natural grasslands. *Nat. Commun.*, 10, 3207–3207.

White, L., O'Connor, N.E., Yang, Q., Emmerson, M.C. & Donohue, I. (2020). Individual species provide

multifaceted contributions to the stability of ecosystems. *Nat. Ecol. Evol.*, 4, 1594–1601.

Winfree, R. & Kremen, C. (2009). Are ecosystem services stabilized by differences among species? A test using

crop pollination. *Proc. R. Soc. B Biol. Sci.*, 276, 229–237.

Wood, S.N., Pya, N. & Säfken, B. (2016). Smoothing Parameter and Model Selection for General Smooth Models.

J. Am. Stat. Assoc., 111, 1548–1563.

Yachi, S. & Loreau, M. (1999). Biodiversity and ecosystem productivity in a fluctuating environment: The

insurance hypothesis. *Proc. Natl. Acad. Sci.*, 96, 1463–1468.

Yodzis, P. (1981). The stability of real ecosystems. *Nature*, 289, 674–676.