RESEARCH ARTICLE

Purification of mitochondria from skeletal muscle tissue for transcriptomic analyses reveals
localisation of nuclear-encoded non-coding RNAs

Authors:

Jessica Silver[1] and Adam J. Trewin[1] (co-first authors), Stella Loke[2], Larry Croft[2], Mark Ziemann[3],
Hayley Dillon[1,4], Søren Nielsen[5], Séverine Lamon[1], Glenn. D. Wadley[1]

Affiliations:

[1]Institute for Physical Activity and Nutrition, School of Exercise and Nutrition Sciences, Deakin
University, Geelong, Victoria, Australia.

[2]Genomics Centre, School of Life and Environmental Sciences, Deakin University, Geelong, Victoria,
Australia.

[3]School of Life and Environmental Sciences, Deakin University, Geelong, Victoria, Australia.

[4]Human Integrated Physiology and Sports Cardiology Laboratory, Baker Heart and Diabetes Institute,
Melbourne, Australia.

[5]The Centre of Inflammation and Metabolism and the Centre for Physical Activity Research,
Rigshospitalet, University of Copenhagen, Copenhagen, Denmark.

Corresponding author:

Prof. Glenn Wadley

Deakin University

Institute for Physical Activity and Nutrition,

School of Exercise and Nutrition Sciences,

Australia

3125

Email: glenn.wadley@deakin.edu.au

ABSTRACT

Mitochondria are central to cellular function, particularly in metabolically active tissues such as

skeletal muscle. Non-coding RNAs (ncRNAs) typically localise within the nucleus and cytosol but

may also translocate to subcellular compartments such as mitochondria. We aimed to investigate the

nuclear-encoded ncRNAs that localise within the mitochondria of skeletal muscle cells and tissue.

Intact mitochondria were isolated via immunoprecipitation and an enzymatic digestion approach was

optimised to remove transcripts located exterior to mitochondria, making it amenable for high-

throughput transcriptomic sequencing. Small-RNA sequencing libraries were successfully constructed

from as little as 1.8ng mitochondrial RNA input. Small-RNA and whole transcriptome sequencing of

mitochondria reveals the enrichment of over 200 miRNAs and 200 lncRNAs that have not previously

been observed within skeletal muscle mitochondria. In summary, we describe a novel, powerful

sequencing approach applicable to animal and human tissues and cells that reveals the unexpected

diversity of nuclear-encoded ncRNA transcripts localised within skeletal muscle mitochondria.

152 words

INTRODUCTION

Mitochondria are subcellular organelles essential for numerous cellular processes including energy metabolism, signalling, ion and redox homeostasis as well as regulation of apoptosis (Spinelli and Haigis, 2018). Mitochondria are comprised of proteins encoded by both the nuclear and mitochondrial genomes, but nuclear-encoded RNAs are not usually localised within mitochondria. A small number of specific RNAs, such as microRNAs (miRNAs) (Barrey et al., 2011, Sripada et al., 2012, Das et al., 2012) and long non-coding RNAs (lncRNAs) (Noh et al., 2016) can however be imported into mitochondria (Silver et al., 2018) via the polynucleotide phosphorylase (Wang et al., 2010) and other putative transport mechanisms (reviewed in Silver et al., 2018). Because ncRNAs have the ability to regulate various transcriptional and translational processes relating to mitochondrial function, it is of interest to determine which of these may localise within mitochondria in order to understand the nature of their potential regulatory actions.

Investigations into mitochondria-localised nuclear-encoded ncRNAs have been constrained, first, by the technical challenges of obtaining pure mitochondrial preparations free of confounding RNAs peripheral to isolated mitochondria and, second, by the significantly lower RNA yields from subcellular fractions when compared to whole tissue homogenates. Early studies providing evidence for the localisation of ncRNAs within mitochondria stem from micro-array analysis and are routinely validated by single qPCR assays or *in situ* hybridisation (Barrey et al., 2011, Sripada et al., 2012, Das et al., 2012). However, the overall population of ncRNAs that localise within mitochondria and their regulatory actions remain unknown.

Skeletal muscle constitutes ~40% of human body mass and plays a central role in whole body metabolism (Zurlo et al., 1990). To meet these dynamic bioenergetic demands, skeletal muscle is densely populated with mitochondria (Russell et al., 2014). Thus, gaining a greater understanding of the RNA profile within muscle mitochondria may yield important insights into the cellular biology and aetiology of diverse metabolic diseases characterised by mitochondrial dysfunction, such as diabetes mellitus and chronic myopathies (Silver et al., 2018). Here, we report a method for isolating and optimising the enzymatic purification of mitochondria from rodent muscle cells or tissues. We provide direct evidence for the purity of this mitochondrial fraction, making it amenable for downstream high-throughput analyses. Finally, we describe a method suitable for small-RNA and whole transcriptome sequencing of mitochondria isolated from skeletal muscle cell and tissue samples from animals and humans, and report the enrichment of numerous nuclear-encoded RNAs that have not previously been observed within skeletal muscle mitochondria.

RESULTS

Protease purification of isolated mitochondria

We first isolated mitochondria from cultured L6 rat myoblasts using an antibody against an outer mitochondrial membrane protein (TOMM22) conjugated to magnetic beads allowing for precipitation of intact isolated mitochondria (Figure 1). Immunoblot of the post-precipitation mitochondrial fraction revealed that a small amount of α-tubulin and Lamin B1 (used as protein markers of the cytosol and nucleus, respectively) were present in the isolated mitochondria fraction (Figure 2, lane 1). The isolated mitochondrial fraction was then incubated with a serine protease (proteinase-K) to digest proteins not located within the mitochondrial intermembrane space (IMS) or matrix compartments. This led to depletion of α-tubulin and Lamin B1, in addition to a marker of the outer mitochondrial membrane (TOMM20). Intermembrane space (cytochrome-*c*) and matrix protein markers (citrate synthase) were protected from digestion (Figure 2, lane 2). All proteins probed were susceptible to proteinase-K when mitochondrial membranes were dissolved in the presence of detergent (triton-X, Figure 2, lane 3). Together, this demonstrates that mitochondrial isolation by TOMM22-immunoprecipitation followed by enzymatic treatment yields a purified, intact mitochondrial fraction suitable for downstream protein analyses. Based on this, we hypothesised that an analogous approach could be used with a ribonuclease to remove RNA located outside of isolated mitochondria.

Ribonuclease purification of isolated mitochondria

Next, mitochondria isolated from L6 myoblasts were exposed to a range of concentrations of RNase-A (Figure 3), an endonuclease that cleaves phosphodiester bonds after pyrimidine residues in the RNA molecule (Witzel, 1963). RNase-A-treated isolated mitochondria were assessed by RT-qPCR for abundance of mtDNA-encoded mRNA (*Mt-co3*) that localises exclusively within the mitochondrial matrix compartment, and a nuclear-encoded mRNA (*Cox4i1*) that localises outside mitochondria. We found that *Mt-co3* was protected from degradation at RNase concentrations up to at least 100 µg/mL, whereas partial *Cox4i1* degradation occurred at 10 µg/mL and was mostly degraded at 100 µg/mL (Figure 3). In the presence of detergent, 1 µg/mL RNase-A was sufficient to digest all *Cox4i1* and *Mt-co3*, as well as an exogenous mRNA spike-in control (Figure 3). This suggests that a certain proportion of cytosolic *Cox4i1* is protected from digestion at lower RNase-A concentrations, potentially as a result of being bound to ribosomes at the outer mitochondrial membrane (Gold et al., 2017). Taken together, these data suggest that TOMM22-IP mitochondrial isolation followed by RNase-A treatment is suitable for downstream mitochondrial RNA analyses provided the RNase-A concentration is optimised to remove transcripts not located within the mitochondrial membranes.

Concurrent investigations have confirmed that RNase-A concentrations exceeding 100 µg/mL are suitable for use in human tissue analysis. This approach successfully removes non-mitochondrial

RNA in isolated mitochondria from human skeletal muscle biopsy samples, and is suitable for the

detection of mitochondria-localised miRNAs (Supplementary Table 1).

Small RNA sequencing on purified isolated mitochondria from myoblast reveals mitochondria

localised miRNAs

Having established optimal RNase-A conditions to obtain purified mitochondria, we next optimized a

protocol to perform transcriptomic analysis of the small RNAs (sRNA) contained within this fraction.

We first prepared sRNA libraries from 120 ng and 60 ng mitochondrial RNA with 0.5X adapter

dilutions (NEBNext 3´ and 5' SR Adaptors, and RT primer for Ilumina), in line with the standard

manufacturer protocol (NEBNext Multiplex Small RNA Library Prep Set (New England Biolabs).

The standard protocol produced large amounts of adapter-dimer and was not sufficient to ligate

adapters to the target miRNAs within the sample (Figure 4 A, B). We next decreased the amount of

the 3' and 5' adapters to investigate if a lower molar ratio improved miRNA ligation efficiency. Both

the modified-0.3X (Figure 4 C) and modified-0.1X (data not shown) approaches produced

quantifiable amounts of the target miRNA library when prepared from 60 ng mitochondrial RNA,

although the total cDNA yield was approximately 7-fold higher from the modified-0.3X when

compared to modified-0.1X protocol. Both modified approaches did decrease, but not prevent,

adapter-dimer formation.

While 60 ng mitochondrial RNA is below the recommended library RNA input (>100 ng, NEBiolabs,

Inc.), mitochondrial RNA yield from cell and tissue samples is often considerably below this

threshold. Thus, it was necessary to investigate if the modified-0.3X and modified-0.1X approach

could similarly produce the target miRNA library when prepared from much lower amounts of

mitochondrial RNA. The modified-0.3X (Figure 4 C, D) and modified-0.1X (data not shown)

approaches both produced the target library when prepared from >1.8 ng (range 1.8-60 ng)

mitochondria RNA. Total cDNA yields were, on average, 4.3-fold higher ($p<0.05$) when using the

modified-0.3X protocol compared to the modified-0.1X. The presence of adapter-dimer dominated

each library, regardless of the mitochondrial RNA input amount used. To combat this, all uniquely

indexed libraries were combined in an equimolar pool and run across two lanes of a TBE

polyacrylamide gel. The gel fragment corresponding to the miRNA region was manually excised,

extracted from the gel and then sequenced. There were no differences in total RNA-seq reads ($p=0.79$)

and total mapped miRNA reads ($p=0.19$) between the modified-0.3X and modified-0.1X protocols

(Figure 5). Over 200 miRNAs were detected across all but the lowest (1.8ng, modified-0.1X)

mitochondrial RNA inputs (Figure 6). The number of miRNAs detected appeared to plateau when

libraries were prepared from 15-60ng mitochondrial RNA (Figure 6B) and suggests that RNA inputs

of at least 15ng are sufficient to detect most miRNAs within a sample. Ultimately, either modified

approach is suitable for the production of miRNA libraries. However, the significantly higher library

yields make the modified-0.3X approach amenable to library pooling, gel excision, and subsequent

sequencing.

We next investigated the relative abundance of miRNAs sequenced from serial dilutions of

mitochondrial RNA input (1.8-30 ng), when compared to the highest mitochondria RNA input (60

ng). Most miRNAs detected were common between libraries prepared from 60 ng mitochondria RNA

and the lower mitochondria RNA inputs (Figure 7A-C). Further to this, the relative abundance of

miRNAs was comparable between libraries prepared from the two highest RNA inputs (60 vs 30 ng;

Figure 7F). A small number of miRNA species were over- (8 miRNAs) and under-represented (5

miRNAs) within libraries prepared from the highest RNA input (60 ng) when compared to lowest

RNA input (1.8 ng; Figure 7D). MiRNAs detected in the highest, but not lowest, RNA input displayed

low expression levels within the 60 ng sample, and together constituted a small proportion (1069 rpm)

of total reads. Interestingly, the skeletal muscle-enriched miR-1, miR-133a, miR-133b and miR-16

and miR-486, predicted to have roles in the regulation of mitochondrial function (Nielsen et al., 2010,

Russell et al., 2013), were detected at higher mitochondria RNA inputs (60 and 30 ng), but not lower

inputs (1.8-15 ng; Figure 7). In contrast, the let-7 family as well as the skeletal muscle-enriched miR-

206 were highly abundant in all samples, and together constituted 63±8% of all mapped miRNA

reads. Importantly, the relative distribution of miRNAs within each sample provides adequate

representation across serial dilutions of mitochondrial RNA, making it possible to accurately

investigate differential miRNA expression across unique cell and tissue samples with variable

mitochondrial RNA yields.

Together, these data suggest that stringent optimization is required when preparing sRNA libraries

from low amounts of RNA from purified mitochondria. To minimize adapter-dimer presence in sRNA

libraries, gel excision and purification of the target miRNA fraction is recommended. Importantly, our

data demonstrate that RNA-seq results were comparable across serial dilutions of mitochondrial RNA

input, suggesting that low mitochondria RNA input does not limit the number nor the relative

distribution of miRNA species that can be detected.

<u>Total RNA-sequencing of isolated mitochondria from rat skeletal muscle tissue reveals selective</u>

<u>enrichment of nuclear-encoded non-coding RNAs</u>

To gain further insight into the RNA landscape within mammalian muscle mitochondria, we then

applied the mitochondrial isolation and purification strategies described above to the gastrocnemius

tissue of 9-week old rats. Whole-transcriptome RNA sequencing revealed that these mitochondria had

a high degree of purity as indicated by the following observations. Firstly, there was a low relative

proportion of nuclear-encoded mRNAs unrelated to mitochondria, such as genes associated with

muscle sarcomere function i.e. myosin light (e.g. *Myl1*) and heavy (e.g. *Myh4* and *Myh7*) isoforms, α-

actin (*Acta1*) and troponins (e.g. *Tnnc2*, *Tnnt3*; Figure 8A). Secondly, the mitochondrial fraction also

contained a low relative abundance of nuclear-encoded OXPHOS genes (including *Cox4i1)*, as
expected (Figure 8B). Thirdly, there was similar abundance of all mtDNA-encoded OXPHOS
transcripts including *Mt-co3* between the whole tissue and mitochondrial fractions (Figure 8C). This
serves as an important internal control because mtDNA transcripts should be present in both isolated
mitochondria and whole tissue in similar proportions. Based on these assumptions, we observed that
non-coding RNAs were enriched in the mitochondrial fraction relative to the respective whole tissue,
including 222 long non-coding RNAs and 102 pseudogenes, ((FDR ≤0.05, log2FC ≥1, Figure 8D and
Supplementary Figure 1). The vast majority of these non-coding RNA transcripts are of unknown
function. Additionally, there were 49 transcripts that encode miRNA genes (i.e. primary- or pre-miR
transcripts) that were also detected in the mitochondrial fraction (Figure 8D and Supplementary
Figure 1).

DISCUSSION

Studies of subcellular fractions using high-throughput analyses requires careful consideration of
purification strategies to avoid confounding effects of contaminants. Here we report an optimised
method for purification of isolated mitochondria for downstream transcriptomic analysis. Revealing
the utility of this approach, we confirmed that miRNAs can be reliably detected in the mitochondria of
L6 myoblasts even when libraries are prepared from RNA inputs well below the thresholds often
recommended by manufacturers. Following this, we report, for the first time, the selective enrichment
of various nuclear-encoded non-coding RNAs within rat skeletal muscle mitochondria.

Incubation of isolated mitochondria with RNase-A can be used to degrade contaminating, non-
mitochondrial RNA to ensure that only RNA species contained within the mitochondria membranes
are used in downstream analyses. This is important when investigating transcripts that do not localise
in one subcellular compartment, but instead may be actively transported between the mitochondria
and cytosol in response to various physiological stimuli. RNase-A protocols differ substantially
between studies that have investigated mitochondrial RNA expression. In previous studies, RNase-A
has been used at variable final concentrations (5-50 μg/mL) (Barrey et al., 2011, Mercer et al., 2011,
Zhang et al., 2014, Bandiera et al., 2011) or has been omitted entirely (Das et al., 2012, Dasgupta et
al., 2015, Lung et al., 2006, Sripada et al., 2012). Here, we demonstrate that RNase-A concentrations
of at least 100 µg/mL are required to digest the majority of nuclear-encoded transcripts (such as
Cox4i1) in rat myoblasts. Importantly, this approach is transferable between cell and tissue models
and is shown to effectively degrade potentially contaminating, non-mitochondrial RNA in
mitochondria isolated from human skeletal muscle tissue (Supplementary Table 1).

Investigations into the mitochondrial ncRNA population are often limited by technical challenges. For
example, the inherently low mitochondrial RNA yields compared to total cellular RNA are well below
the threshold required for commercially available small RNA library preparation protocols. This

challenge may be overcome by combining mitochondria RNA from biological samples into pools that represent a single 'intervention' or 'clinical' group, and the 'control' group (Jagannathan et al., 2015, Wang et al., 2017). While this effectively increases the amount of mitochondrial RNA available for sRNA library preparation, this approach introduces further limitations. First, the use of pooled samples requires a much larger sample size to achieve the same statistical power (Takele Assefa et al., 2020). Second, gene expression is highly variable between individuals; taking a pooled sample approach may mask potential individual variability in miRNA expression (Takele Assefa et al., 2020, Rajkumar et al., 2015). Our findings illustrate that small RNA libraries can be prepared from individual biological samples without the need to pool multiple samples, thereby increasing the statistical power available, especially for human studies with limited sample material and high participant burden.

A modified small RNA library preparation protocol increased ligation efficiency of the 3' and 5' adapters to miRNAs within mitochondrial samples. However, this modified approach was not able to prevent adapter-dimer formation entirely. As the adapter-dimer product is approximately 20 bp smaller than adapter-ligated miRNAs and preferentially binds to the flow cell during RNA-Seq, gel extraction of the target miRNA region is essential to maximise read counts that map to miRNAs rather than adapter-dimer (Shore et al., 2016). When used together, the modified-0.3X protocol plus gel extraction were successful in detecting over 200 miRNAs from L6 mitochondria RNA inputs as low as 1.8 ng. Importantly, the relative distribution of miRNAs within each sample was reasonably consistent as RNA input decreased. However, some miRNA species tended to be over-represented at the lowest mitochondria RNA inputs. This may be because there were less miRNA species detected in the library prepared from 1.8 ng (207 miRNAs) when compared to 60ng mitochondria RNA (266 miRNAs). Overall, there appears to be minimal bias towards over- or under-representing specific miRNA species at low RNA inputs.

Of note, the skeletal muscle-enriched miR-1, miR-133a and miR-133b were detected at high (30 and 60 ng) but not low (1.8-15 ng) mitochondrial RNA inputs. MiR-1 and miR-133a have previously been observed in mitochondria isolated from human skeletal and rodent cardiac muscle cells *in vitro* (Barrey et al., 2011, Jagannathan et al., 2015, Zhang et al., 2014). MiR-1 enhances the transcription of the mitochondrial-encoded mt*Cox1,* mt*Nd1* and mt*Cytb*, among others (Zhang et al., 2014), while miR-133a inhibits the transcription of nuclear-encoded *PGC-1* isoforms and *Tfam* (Nie et al., 2016), both of which are upstream regulators of mitochondrial biogenesis. The use of proliferating L6 myocytes in this study, compared with primary human myocytes (Barrey et al., 2011) and rat neonatal cardiomyocytes (Zhang et al., 2014) used previously, may partially account for the differences in miRNA species detected. Although L6 cells have reportedly higher rates of aerobic metabolism when compared to C2C12 cells and primary human myocytes, cell culture models are more dependent upon anaerobic glycolysis (Abdelmoez et al., 2020). Thus, mitochondrial abundance and function in

various cell culture models may differ to those observed with *in vivo* models. This study was primarily designed to optimise the ncRNA-Seq library preparation protocol rather than investigate the physiological relevance of specific miRNAs. Future *in vivo* studies are now required to profile mitochondrial miRNAs and their roles in tissues that have high bioenergetic requirements such as skeletal and cardiac muscle.

Finally, the enrichment of non-coding RNAs, including lncRNAs and pseudogenes, within mitochondria relative to whole skeletal muscle tissue suggests that a previously underappreciated scale of RNA import and/or regulation may occur at the subcellular level. Only a relatively small fraction (~5-10%) of all identified lncRNAs have been functionally characterized (Gao et al., 2020). LncRNA can have important regulatory roles (Wadley et al., 2019), mediating cellular processes by modulating the expression of genes, stability of transcripts and/or function of proteins (Long et al., 2017), including those that directly determine mitochondrial function. Individual validation and functional characterisation of the lncRNAs identified in the present study is required to understand the precise mechanisms involved and potential functional implications. This may also open new avenues of investigation into how they are imported into the mitochondrial compartment, what are their putative binding target(s) and whether they modulate expression levels of mtDNA encoded genes and/or function of mitochondrial proteins. This will be key to understanding the potential effects on mitochondrial parameters such as oxidative phosphorylation, ROS production, mitochondrial biogenesis, dynamics or mitophagy.

In summary, we report a method for obtaining a highly pure fraction of mitochondria from skeletal muscle tissue and optimised approach for high throughput transcriptomic sequencing on this subcellular fraction. Importantly, we reveal the presence of non-coding RNAs within mitochondria that were not previously known to reside within this subcellular compartment. Given the increasing interest in non-coding RNA mediated regulation of cellular function in health and disease, future investigations are warranted to better understand the potential biological and physiological significance of the mitochondrial localisation of these non-coding RNAs.

MATERIALS AND METHODS

Cell culture

L6 rat myoblast cells were obtained from a commercial vendor (ATCC Cat# CRL-1458, RRID: CVCL_0385). Cells were cultured at 37□ with 5% CO_2 in a humidified incubator in media consisting of Dulbecco's Modified Eagle Medium (DMEM; Gibco #11995-065) supplemented with 10% v/v FBS (Gibco #A3161001), 100 U/ml penicillin and 100 μg/ml streptomycin (Gibco #15140-122). Cells were grown in 150 mm dishes to ~90% confluence then washed with sterile PBS and detached (Gibco

TrypLE Express #LTS12604021). Cells used for all experiments were ≤9 passages from stock and

tested negative for mycoplasma contamination (Agilent MycoSensor QPCR Assay Kit, 302107).

<u>Myoblast mitochondrial isolation</u>

Mitochondria were isolated from L6 myoblasts via immunoprecipitation (Miltenyi Biotek #130-096-

946). Approximately 5×10^6 cells (suspended in a small volume of PBS after detachment) were

transferred to 1 mL ice-cold lysis buffer, then mechanically homogenised with 20 passes in a 7 mL

glass dounce homogeniser (Wheaton) on ice. Wash buffer (9 mL) was added to the lysate and

transferred to a falcon tube. Anti-TOMM22 antibody conjugated magnetic beads (50 µL) were added

to the 10 mL lysate and incubated with gentle inversion for 1 hour at 4□. Lysate was passed through a

pre-separation filter then flowed through a column mounted on a magnetic rack. Captured

mitochondria were washed 3 times before being removed from the magnetic rack for elution.

Mitochondria were pelleted via centrifugation at 13,000 x g for 2 minutes at 4□. Supernatant was then

removed and the mitochondrial pellet resuspended in storage buffer to be used for subsequent

purification.

<u>Enzymatic purification of myoblast mitochondria</u>

Protease treatment: To remove proteins not localised within intact mitochondria, aliquots of isolated

mitochondria (1 µg total protein each) were incubated in a 50 µL reaction with or without triton-x

(1% v/v) and with or without proteinase K (Qiagen #19131, final concentration 20 µg/mL) for 15

minutes at room temperature. Phenylmethylsulfonyl fluoride (20 mM) was added to inhibit

proteinase-K. Samples were then stored at -20□ until immunoblot analysis.

RNase treatment: To remove RNA not localised within intact mitochondria, in a 50 µL reaction,

aliquots of isolated mitochondria (1 µg total protein each) first had 100 ng mRNA spike-in control

added (StemMACS eGFP mRNA #130-101-114, Miltenyi Biotec), and were then incubated with or

without triton-x (1% v/v) and with or without RNase-A (Qiagen #19101, final concentration range 1-

1000 µg/mL) and incubated on ice for 30 minutes. Trizol reagent was then added to simultaneously

inactivate RNase activity and preserve RNA. Samples were then stored in Trizol at -20□ until RNA

isolation.

<u>Immunoblot analysis</u>

Mitochondrial samples from proteinase-K treatment were thawed and resuspended in RIPA buffer

(Millipore #20-188) containing protease inhibitors (Sigma #P8340). Protein concentration was

determined by BCA assay (Pierce #23225). Samples were mixed with reducing loading buffer (4x

Laemmli sample buffer with 10% 2-mercaptoethanol). Equal volumes of mitochondrial samples (0.2

µg total protein) were loaded into a 4-15% gel (Bio-Rad #5678085) in addition to whole cell lysate

(12 µg) and molecular weight marker (Bio-Rad #161-0373), which were then separated by

electrophoresis at 40 V for 30 min then 120 V for 60 min. The stain-free gel was UV-activated for 1

minute and imaged for total protein (BioRad Chemi Doc XR+, Hercules, CA, USA) with ImageLab

software (BioRad Image Lab v6). Protein was then transferred (Turbo Transfer system, Bio-Rad) to a

methanol soaked polyvinylidene difluoride (PVDF) membrane (Millipore Immobilon FL 0.45 µm

#IPFL00010). The membrane was then PBS washed, dried for 1 hour, re-wetted with methanol,

washed with PBS then blocked for 1 hour (Li-Cor Intercept PBS blocking buffer). The membrane was

then incubated with primary antibody in blocking buffer + 0.2% v/v Tween-20 overnight at 4°C.

Primary antibodies used were: anti-TOMM20 (Abcam #ab186735; 1:1000), anti-Cytochrome-c

(Abcam #ab133504; 1:1000), anti-Citrate Synthase (Cell Signaling Technology #14309; 1:1000),

anti-α-Tubulin (Cell Signaling Technology #3873; 1:1000), and anti-Lamin B1 (Cell Signaling

Technology #12586; 1:1000). Membranes were then washed with PBST and incubated with

secondary antibodies (Cell Signaling Technology) anti☐rabbit or anti☐mouse IgG Dylight® 680 nm

(Cell Signaling Technology #5366S, #5470S) or 800 nm (Cell Signaling Technology #5151S or

#5257S) at 1:10,000 in blocking buffer containing 0.2% Tween-20 and 0.01% SDS for 1 hour at room

temperature. Fluorescent signal was then digitally acquired (Licor Odyssey® Infrared Imaging

System, Lincoln, NE, USA).

<u>RNA isolation and quality control</u>

Myoblast mitochondria samples stored in Trizol were thawed on ice and centrifuged at 10,000 x g for

10 min at 4☐. An aliquot of the supernatant was then used for RNA purification with on-column

DNase-I treatment (Direct-Zol RNA Microprep R2060, Zymo Research). RNA concentration and

purity was determined by spectrophotometry (NanoDrop 1000, Thermo Fisher Scientific) while RNA

fragment size distribution was assessed by gel electrophoresis (Tapestation HS-RNA, Agilent

Technologies).

<u>Reverse transcription and quantitative PCR (RT-qPCR)</u>

Equal volumes of isolated RNA from the RNase-treated mitochondria samples (to account for

differences in total RNA yield between samples due to RNase treatment) were reverse transcribed to

first-strand cDNA in a 20 µL reaction along with no-template and no-RTase controls (Applied

Biosystems High Capacity RT kit #4368814). Quantitative PCR was performed in triplicate (Agilent

AriaMX G8830A) on 4 ng of cDNA in a 10 µL reaction consisting of SYBR green master mix

(Applied Biosystems #4367659) with 0.3 µM forward and reverse primers (Table 1).

Table 1. List of RT qPCR primers

Target	Forward (5' -3')	Reverse (5' -3')	Accession ID
Mt-co3	GACGGAATTTACGGCTCAACAT	AATTAGGAAAGTTGAGCCAATAATTACG	>NC_001665.2 Rattus norvegicus strain BN/SsNHsdMCW mitochondrion, complete genome

	Forward	Reverse	Reference
Cox4i1	GTGCTGATCTGGGAGAAGAGCTA	GGTTGACCTTCATGTCCAGCAT	>NM_017202.1 Rattus norvegicus cytochrome c oxidase subunit 4i1 (Cox4i1), mRNA; nuclear gene for mitochondrial product
eGFP	ACTTCTTCAAGTCCGCCATG	AGCTCGATGCGGTTCACCAG	>XM_013480425.1 Eimeria maxima Green fluorescent protein, related partial mRNA

Thermal conditions used for qPCR were 10 min at 95□ (activation) then 40 cycles of 15 s at 95°C

(denature) and 60 s at 55-60□ (anneal/extend). A dissociation curve was performed to confirm the

amplification of a single product. Quantification cycle (C_q) thresholds were calculated using software

(Agilent Aria v1.5) expressed after linear transformation.

<u>Myoblast small RNA sequencing and bioinformatics</u>

Total RNA isolated from individual (n=7) L6 mitochondria samples was pooled into a single tube,

from which serial dilutions ranging from 0.3-20 ng/µL were prepared. Complementary DNA libraries

were then prepared using a range of mitochondrial RNA inputs (final input range 1.8-120 ng) using

the NEBNext Multiplex Small RNA Library Prep Set (New England Biolabs.). To investigate the

efficiency of small RNA library preparation across a range of mitochondrial RNA inputs, libraries

were prepared either according to the manufacturer's recommendations for low RNA inputs

(NEBNext 3´ and 5' SR Adaptors, and RT primer for Ilumina diluted 0.5X, 15 PCR cycles) or with

modifications (NEBNext 3´ and 5' SR Adaptors, and RT primer for Ilumina diluted 0.3X or 0.1X, 20

PCR cycles). cDNA concentration was determined by fluorometric quantitation (Qubit 1xdsDNA,

Thermo Fisher Scientific) while cDNA fragment size distribution was assessed by gel electrophoresis

(Tapestation HS-D1000, Agilent Technologies).

An equimolar pool of all uniquely indexed small RNA libraries was prepared. Twenty-five µL of the

equimolar pool was loaded across two lanes on a 6% Novex TBE polyacrylamide gel (Thermo

Fischer Scientific), with 5 µL Quick-Load pBR322 DNA-MspI Molecular Marker (New England

Biolabs Inc.) in a separate lane. The buffer dam was filled with 600 mL 1X TBE running buffer and

the gel run at 120V for 1h. The gel was then incubated with 50 mL 1X TBE spiked with 1X SYBR

Gold Nucleic Acid Gel Stain (Thermo Fischer Scientific), before being visualised on a Safe Imager

2.0 Blue-Light Transilluminator (Invitrogen) following exposure to UV-Blue (470 nm) light.

Fragments corresponding to the 130-160 bp region were manually excised, before suspended and then

crushed in 250 µL gel elution buffer (New England Biolabs Inc.). To increase the recovery of low

concentrations of small nucleic acids, cDNA was precipitated overnight at -20°C with 0.3 M sodium

acetate (pH 5.5), 4X volumes of 100% ethanol and 1 µL linear acrylamide carrier. Following

overnight precipitation, the solution was centrifuged at 16,000x*g* for 60 minutes to pellet the cDNA

fragment. The cDNA pellet was washed twice with 80% ethanol and resuspend in 12 µL TE buffer.

Fragment size distribution of the gel-extracted cDNA pool was assessed by gel electrophoresis

(Tapestation HS-D1000, Agilent Technologies).

RNA sequencing was performed by the Deakin University Genomics Core on the Illumina MiniSeq

platform. A 1.3 pM loading pool was prepared with 20% Phi-X spike-in and was sequenced on a

single-end 75 bp run. Sequenced reads underwent quality checks with FastQC before the adapter and

reads <20nt were trimmed. Small RNAs, including miRNAs, are not well sequenced in the rat

genome but are conserved between rat and mouse species. Reads were mapped to known mature

mouse miRNAs accessible from miRbase v22.0. Raw read counts were normalised by the size of each

individual library. Counts were visualised using GraphPad Prism (v7).

Rodent study ethical approval and procedures

All experimental procedures were approved by the Deakin University Animal Ethics Committee

(G02-2019). All animals were housed and treated in accordance with standards set by Deakin

University's Animal Welfare Committee, which complies with the ethical and governing principles

outlined in the Australian code for the care and use of animals for scientific purposes. Wistar Kyoto

male rats were obtained from the Animal Resource Centre, Perth, Western Australia. Rats were

housed in pairs and maintained with a 12-hour light/dark cycle, constant temperature of $21 \pm 2°C$, and

humidity levels between 40 and 70%. Rats had *ad libitum* access to standard chow diet and tap water.

At 9-weeks of age, animals were humanely killed following dissection of the heart under heavy

anaesthesia using 5% isoflurane gas. The gastrocnemius muscle was rapidly excised then the inner red

portion was dissected. A portion of this was snap frozen in liquid nitrogen for subsequent whole

muscle RNA analyses, and a separate ~ 100 mg piece was taken for mitochondrial isolation. This was

immediately placed in 1 mL ice-cold lysis buffer and minced using fine scissors before mechanical

homogenisation with an ice-cold Teflon-tipped glass dounce homogeniser (10 passes with rotation at

350 rpm). Intact mitochondria were then isolated using the magnetic-bead immunoprecipitation

method as described above for L6 myoblasts except with the following RNase treatment conditions:

RNase-A 340 µg/mL in a reaction with ~20 µg total mitochondrial protein performed for 1 hour at

37□, after which 5 µL proteinase-K (stock concentration 600 mAU/mL) was then added to inactivate

RNase activity (Barrey et al., 2011). Each tube was briefly inverted before the mitochondria solutions

were then centrifuged at 8000 x g for 10 min at 4□. The supernatant was aspirated and the pellet was

washed twice with 100 µL ice-cold storage buffer. The RNase-treated mitochondrial pellet was

resuspended in 100 uL storage buffer then frozen at -80□ until RNA extraction.

RNA extraction from rat muscle tissue and isolated mitochondria

For RNA extraction from whole red gastrocnemius muscle, ~15 mg frozen tissue was placed in a liquid nitrogen pre-chilled cyrotube with a 5 mm stainless steel bead. The tissue was then mechanically disrupted (2 cycles at 4m/s for 10 s, MP Biomedical FastPrep). Tri-reagent (600 uL) was added, then RNA extracted (Zymo DirectZol RNA Miniprep #R2050). For RNA extraction from isolated mitochondria from the respective red portion of gastrocnemius muscle, frozen samples were thawed in the presence of 5 volumes of TRI-reagent (Qiagen), pipette mixed thoroughly then centrifuged at 16,000 x g for 1 min. RNA extraction was then performed with on-column DNase I treatment (Zymo Research DirectZol RNA Mircroprep #2060). Extracted RNA was tested for purity on Nanodrop, concentration with Qubit HS RNA assay (Thermo Fisher) then RNA integrity number (RIN) was determined using a Tapestation (Agilent) with High Sensitivity RNA reagents. All RINs from whole tissue RNA were ≥7.0.

Rat muscle tissue and isolated mitochondria total RNA sequencing

Total RNA from whole-tissue (50 ng) and isolated mitochondria (10 ng) were converted to cDNA libraries (Zymo-Seq RiboFree Total RNA #R300, Zymo Research). Briefly, ribosomal RNA was depleted, then remaining transcripts were fragmented and converted to cDNA using random priming. After second strand synthesis, the ends of the cDNA were enzymatically repaired and Illumina-compatible sequencing adaptors were ligated. Library size (TapeStation, Agilent) and concentration (Qubit) was assessed prior to sequencing. RNA sequencing was performed by the Deakin University Genomics Core on the Illumina NovaSeq 6000 platform. Reads underwent adapter trimming and quality check with Skewer then mapped to the rat genome (Ensembl version 99) with STAR aligner v2.7.2a. STAR generated read counts were collated in R version 4.0.3 (R Development Core Team, 2010). Normalised counts were visualised using GraphPad Prism (v7). Analysis of differential expression was performed using Voom/Limma in Degust v 4.1.5 (Powell, 2019). Transcripts with a false discovery rate (FDR) <0.05 were considered differentially expressed.

Statistical analysis

Small RNA-seq data was analysed by paired t-test. RNase-A dose-response data were tested for normality of variance and then analysed by one- or two-way ANOVA with Tukey's post hoc test for multiple comparisons using GraphPad Prism (v7). All data are presented as mean (SD) with significance accepted at $p < 0.05$.

Acknowledgments

We acknowledge Gisella Mazzarino and Amandi Gunawardena for their assistance with animal experiments and tissue collection. We acknowledge Sarah Alexander for her assistance with female participant recruitment. Severine Lamon is supported by an Australian Research Council Future Fellowship (FT10100278).

Author contributions

Conceptualization: JLS, AJT, SN, S.Lamon, GDW

Conducted experiments: JLS, AJT, GDW, S.Loke

Conducted human studies: JLS, GDW, S.Lamon, HD

Data analysis (bioinformatics): MZ, LC

Data analysis and visualisation: JLS, AJT

Manuscript – original draft: JLS & AJT

Manuscript– review and editing: all authors.

Approved final version of manuscript: all authors.

REFERENCES

ABDELMOEZ, A. M., SARDÓN PUIG, L., SMITH, J. A. B., GABRIEL, B. M., SAVIKJ, M., DOLLET, L., CHIBALIN, A. V., KROOK, A., ZIERATH, J. R. & PILLON, N. J. 2020. Comparative profiling of skeletal muscle models reveals heterogeneity of transcriptome and metabolism. *Am J Physiol Cell Physiol,* 318, C615-c626.

BANDIERA, S., RUBERG, S., GIRARD, M., CAGNARD, N., HANEIN, S., CHRETIEN, D., MUNNICH, A., LYONNET, S. & HENRION-CAUDE, A. 2011. Nuclear outsourcing of RNA interference components to human mitochondria. *PLoS One,* 6, e20746.

BARREY, E., SAINT-AURET, G., BONNAMY, B., DAMAS, D., BOYER, O. & GIDROL, X. 2011. Pre-microRNA and mature microRNA in human mitochondria. *PLoS One,* 6, e20220.

DAS, S., FERLITO, M., KENT, O. A., FOX-TALBOT, K., WANG, R., LIU, D., RAGHAVACHARI, N., YANG, Y., WHEELAN, S. J., MURPHY, E. & STEENBERGEN, C. 2012. Nuclear miRNA regulates the mitochondrial genome in the heart. *Circ Res,* 110, 1596-603.

DASGUPTA, N., PENG, Y., TAN, Z., CIRAOLO, G., WANG, D. & LI, R. 2015. miRNAs in mtDNA-less cell mitochondria. *Cell Death Discov,* 1, 15004.

GAO, F., CAI, Y., KAPRANOV, P. & XU, D. 2020. Reverse-genetics studies of lncRNAs—what we have learnt and paths forward. *Genome Biology,* 21, 93.

GOLD, V. A., CHROSCICKI, P., BRAGOSZEWSKI, P. & CHACINSKA, A. 2017. Visualization of cytosolic ribosomes on the surface of mitochondria by electron cryo-tomography. *EMBO Rep,* 18, 1786-1800.

JAGANNATHAN, R., THAPA, D., NICHOLS, C. E., SHEPHERD, D. L., STRICKER, J. C., CROSTON, T. L., BASELER, W. A., LEWIS, S. E., MARTINEZ, I. & HOLLANDER, J. M. 2015. Translational Regulation of the Mitochondrial Genome Following Redistribution of Mitochondrial MicroRNA (MitomiR) in the Diabetic Heart. *Circulation. Cardiovascular genetics,* 8, 785-802.

LONG, Y., WANG, X., YOUMANS, D. T. & CECH, T. R. 2017. How do lncRNAs regulate transcription? *Science Adv,* 3, eaao2110.

LUNG, B., ZEMANN, A., MADEJ, M. J., SCHUELKE, M., TECHRITZ, S., RUF, S., BOCK, R. & HUTTENHOFER, A. 2006. Identification of small non-coding RNAs from mitochondria and chloroplasts. *Nucleic Acids Res,* 34, 3842-52.

MERCER, T. R., NEPH, S., DINGER, M. E., CRAWFORD, J., SMITH, M. A., SHEARWOOD, A. M., HAUGEN, E., BRACKEN, C. P., RACKHAM, O., STAMATOYANNOPOULOS, J. A., FILIPOVSKA, A. & MATTICK, J. S. 2011. The human mitochondrial transcriptome. *Cell,* 146, 645-58.

NIE, Y., SATO, Y., WANG, C., YUE, F., KUANG, S. & GAVIN, T. P. 2016. Impaired exercise tolerance, mitochondrial biogenesis, and muscle fiber maintenance in miR-133a-deficient mice. *Faseb j,* 30, 3745-3758.

NIELSEN, S., SCHEELE, C., YFANTI, C., AKERSTROM, T., NIELSEN, A. R., PEDERSEN, B. K. & LAYE, M. J. 2010. Muscle specific microRNAs are regulated by endurance exercise in human skeletal muscle. *J Physiol,* 588, 4029-37.

NOH, J. H., KIM, K. M., ABDELMOHSEN, K., YOON, J.-H., PANDA, A. C., MUNK, R., KIM, J., CURTIS, J., MOAD, C. A. & WOHLER, C. M. 2016. HuR and GRSF1 modulate the nuclear export and mitochondrial localization of the lncRNA RMRP. *Genes Dev,* 30, 1224-1239.

POWELL, D. 2019. Degust v4.1.1. Zenodo. doi.org/10.5281/zenodo.3501067.

RAJKUMAR, A. P., QVIST, P., LAZARUS, R., LESCAI, F., JU, J., NYEGAARD, M., MORS, O., BØRGLUM, A. D., LI, Q. & CHRISTENSEN, J. H. 2015. Experimental validation of methods for differential gene expression analysis and sample pooling in RNA-seq. *BMC Genomics,* 16, 548.

RUSSELL, A. P., FOLETTA, V. C., SNOW, R. J. & WADLEY, G. D. 2014. Skeletal muscle mitochondria: a major player in exercise, health and disease. *Biochim Biophys Acta,* 1840, 1276-84.

RUSSELL, A. P., LAMON, S., BOON, H., WADA, S., GULLER, I., BROWN, E. L., CHIBALIN, A. V., ZIERATH,
J. R., SNOW, R. J., STEPTO, N., WADLEY, G. D. & AKIMOTO, T. 2013. Regulation of miRNAs in
human skeletal muscle following acute endurance exercise and short-term endurance
training. *J Physiol,* 591**,** 4637-53.
SHORE, S., HENDERSON, J. M., LEBEDEV, A., SALCEDO, M. P., ZON, G., MCCAFFREY, A. P., PAUL, N. &
HOGREFE, R. I. 2016. Small RNA Library Preparation Method for Next-Generation Sequencing
Using Chemical Modifications to Prevent Adapter Dimer Formation. *PLoS One,* 11**,** e0167009.
SILVER, J., WADLEY, G. & LAMON, S. 2018. Mitochondrial regulation in skeletal muscle: A role for
non-coding RNAs? *Experimental Physiology* 103**,** 1132-1144.
SPINELLI, J. B. & HAIGIS, M. C. 2018. The multifaceted contributions of mitochondria to cellular
metabolism. *Nat Cell Biol,* 20**,** 745-754.
SRIPADA, L., TOMAR, D., PRAJAPATI, P., SINGH, R., SINGH, A. K. & SINGH, R. 2012. Systematic
Analysis of Small RNAs Associated with Human Mitochondria by Deep Sequencing: Detailed
Analysis of Mitochondrial Associated miRNA. *PLoS ONE,* 7**,** e44873.
TAKELE ASSEFA, A., VANDESOMPELE, J. & THAS, O. 2020. On the utility of RNA sample pooling to
optimize cost and statistical power in RNA sequencing experiments. *BMC Genomics,* 21**,** 312.
WADLEY, G. D., LAMON, S., ALEXANDER, S. E., MCMULLEN, J. R. & BERNARDO, B. C. 2019. Non-
coding RNAs regulating cardiac muscle mass. *Journal of Applied Physiology,* 127**,** 633-644.
WANG, G., CHEN, H. W., OKTAY, Y., ZHANG, J., ALLEN, E. L., SMITH, G. M., FAN, K. C., HONG, J. S.,
FRENCH, S. W., MCCAFFERY, J. M., LIGHTOWLERS, R. N., MORSE, H. C., 3RD, KOEHLER, C. M.
& TEITELL, M. A. 2010. PNPASE regulates RNA import into mitochondria. *Cell,* 142**,** 456-67.
WANG, X., SONG, C., ZHOU, X., HAN, X., LI, J., WANG, Z., SHANG, H., LIU, Y. & CAO, H. 2017.
Mitochondria Associated MicroRNA Expression Profiling of Heart Failure. *Biomed Res Int,*
2017**,** 4042509.
WITZEL, H. 1963. The Function of the Pyrimidine Base in the Ribonuclease Reaction. *In:* DAVIDSON, J.
N. & COHN, W. E. (eds.) *Progress in Nucleic Acid Research and Molecular Biology.* Academic
Press.
ZHANG, X., ZUO, X., YANG, B., LI, Z., XUE, Y., ZHOU, Y., HUANG, J., ZHAO, X., ZHOU, J., YAN, Y.,
ZHANG, H., GUO, P., SUN, H., GUO, L., ZHANG, Y. & FU, X. D. 2014. MicroRNA directly
enhances mitochondrial translation during muscle differentiation. *Cell,* 158**,** 607-19.
ZURLO, F., LARSON, K., BOGARDUS, C. & RAVUSSIN, E. 1990. Skeletal muscle metabolism is a major
determinant of resting energy expenditure. *J Clin Invest,* 86**,** 1423-7.

FIGURE LEGENDS

Figure 1. Overview of TOMM22 immunoprecipitation based method for mitochondrial isolation

Figure 2. Protease treatment of mitochondria isolated from L6 myocytes. Isolated mitochondria were incubated with or without proteinase-K (20 µg/mL) and detergent (1% triton-X). Marker proteins of subcellular compartments were then probed by immunoblot. Blots are representative of n=2 technical replicates from a single mitochondrial isolation preparation.

Figure 3. RNase-A treatment of mitochondria isolated from L6 myocytes. Isolated mitochondria were incubated with or without RNase-A (1-1000 µg/mL) in the presence or absence of detergent (1% triton-X). The levels of representative mtDNA (*Mtco3*) and nuclear (*Cox4i1*) encoded genes were then assessed by RT-qPCR in addition to an exogenous mRNA spike-in control (*eGFP*) added to isolated mitochondria (but not to the whole cell lysate). Data are mean(SD) for N=3 technical replicates from a single mitochondrial isolation preparation.

Figure 4. Small RNA-seq library preparation from mitochondrial RNA. Representative electropherogram traces of libraries generated with the standard 0.5X dilution of 3' and 5' adapters produces large amounts of adapter-dimer (red box) and does not ligate to mitochondria miRNAs with either A) 120 ng or B) 60 ng mitochondrial RNA input. Lower molar ratios (0.3X dilution) of 3' and 5' adapters produces the target miRNA library (green box) with C) 60ng and D) 1.8ng mitochondria RNA. Peaks at 25 and 1500 bp are internal size standards.

Figure 5. Small RNA-Seq on isolated mitochondria from L6 myocytes. Small RNA libraries were prepared from 1.8 to 60 ng mitochondria RNA with modified-0.3X (n=6) or modified-0.1X (n=4) protocols. The total number of reads (p=0.79) and total mapped miRNA reads (p=0.19) were not different between protocols. Student's paired t-test, n=1 at each RNA input.

Figure 6. Neither mitochondrial RNA input nor library preparation protocols limit the number of mature miRNAs that can be detected from mitochondrial RNA. A) The number of mature miRNA species detected across all mitochondria RNA inputs was not different (p=0.60, student's paired t-test) following modified-0.3X (m-0.3X; n=6) and modified-0.1X (m-0.1X; n=4) protocols. Each point represents an independent library prepared from 1.8-60ng mitochondrial RNA.

Figure 7. The relative abundance of miRNAs within each sample is comparable between small RNA libraries prepared from high, moderate and low mitochondrial RNA inputs. Venn diagrams depicting unique and shared miRNAs detected from 60ng mitochondria RNA when compared to **A)** 1.8ng, **B)** 7.5ng and **C)** 30ng mitochondria RNA, prepared using the modified-0.3X library preparation approach. The relative distribution of miRNAs detected at each RNA input was expressed relative to mmu-miR-let-7c-2. Differences in the relative distribution of miRNAs between the highest RNA input and serial RNA dilutions (**D)** 1.8ng, **E)** 7.5ng and **F)** 30ng) are depicted by grey dots (<5-fold change when compared to 60ng), red dots (>5-fold change when compared to 60ng) and green dots (>10-fold change when compared to 60ng). Highly abundant miRNAs across all RNA inputs (including the let-7 family and mmu-miR-206) are shown in orange dots.

Figure 8. RNase-treated mitochondria isolated from rat skeletal muscle tissue contains nuclear-encoded non-coding RNAs. Whole-transcriptome RNA sequencing of isolated mitochondria and respective whole muscle tissue (rat gastrocnemius) shows that the isolated mitochondria were free of **A)** muscle sarcomere related mRNAs **B)** and nuclear-encoded mRNAs involved in the mitochondrial electron transport chain, while **C)** mtDNA encoded mitochondrial electron transport chain genes were present. **D)** Scatterplot depicting the abundance of RNA classes in isolated mitochondria relative to whole muscle; each point represents the mean normalised abundance (reads per million, RPM) of an individual gene from *n*=6 biological replicates.

Figure 1.

Isolated mitochondria   **Whole cell**

Proteinase-K   -   +   +   -
Triton-X   -   -   +   -

Ladder

16 kDa — TOMM20

14 kDa — Cytochrome-*c*

45 kDa — Citrate Synthase

52 kDa — α-tubulin

68 kDa — Lamin B1

100 kDa — Stain-free total protein

15 kDa

Figure 2.

Figure 3.

Figure 4.

Figure 5.

Figure 6.

Figure 7.

Figure 8.