Mutation rates and adaptive variation among the clinically dominant clusters of

Mycobacterium abscessus

Nicoletta Commins[1], Mark R. Sullivan[2], Kerry McGowen[2], Evan Koch[1], Eric J. Rubin[2,3], Maha

Farhat[1,4] *

[1] Department of Biomedical Informatics, Harvard Medical School, Boston, MA, 02115, USA.

[2] Department of Immunology and Infectious Diseases, Harvard T. H. Chan School of Public

Health, Boston, MA, 02115, USA.

[3] Department of Microbiology, Harvard Medical School, Boston, MA, 02115, USA.

[4] Division of Pulmonary and Critical Care Medicine, Massachusetts General Hospital, Boston,

MA, 02114, USA.

* Correspondence: maha_farhat@hms.harvard.edu

Abstract

Mycobacterium abscessus (*Mab*) is a multi-drug resistant pathogen increasingly responsible for

severe pulmonary infections. Analysis of whole genome sequences (WGS) of *Mab* demonstrates

dense genetic clustering of clinical isolates collected from disparate geographic locations. This

has been interpreted as supporting patient-to-patient transmission, but epidemiological studies

have contradicted this interpretation. Here we present evidence for a slowing of the *Mab*

molecular clock rate coincident with the emergence of phylogenetic clusters. We find that

clustered isolates are enriched in mutations affecting DNA repair machinery and have lower

spontaneous mutation rates in *in vitro*. We propose that *Mab* adaptation to the host environment

through variation in DNA repair genes affects the organism's mutation rate and that this manifests

as phylogenetic clustering. These results inform our understanding of niche switching for

facultative pathogens, and challenge the model of transmission as the major mode of

dissemination of clinically dominant *Mab* clusters.

Introduction

Mycobacterium abscessus (*Mab*) is an emerging, multi-drug resistant pathogen that is

increasingly responsible for opportunistic pulmonary disease, most commonly in patients with

underlying structural lung conditions such as cystic fibrosis (CF).[1] *Mab* is divided into three

subspecies: *M. a. abscessus*, *M. a. massiliense*, and *M. a. bolletii,* the former two being the most

clinically relevant subspecies.[2] Infections with *Mab* are associated with accelerated decline in lung

function and can result in severe disseminated disease and/or loss of eligibility for lung

transplantation. *Mab* infections are difficult to treat, requiring a prolonged antibiotic regimen with

high rates of treatment failure.[3]

Until recently, *Mab* infections were thought to be acquired independently by each patient

from environmental reservoirs in soil or water. Large scale analyses of whole genome sequencing

(WGS) of Mab clinical isolates have revealed that a large proportion of clinical isolates fall into

several phylogenetically characterized clusters, often termed dominant circulating clones.[4]

Isolates within these clusters have high core genome similarity that is not explained by point

source outbreaks as they were sampled from diverse geographic origins. These observations

have been interpreted as supporting widespread recent transmission, either through direct contact

or indirectly through fomites. However, a number of epidemiological studies have refuted person-

to-person transmission as a major mode of *Mab* dissemination.[5–11] Reconciling these conflicting

lines of evidence is important as they dictate public health practice and allocation of resources to

protect vulnerable populations from infection. However, it is still unclear what role transmission

plays in the emergence of *Mab* clusters.

While widespread transmission is one proposed explanation for the phylogenetic structure

observed in *Mab*, it is important to consider other factors in the species' evolutionary history that

may contribute. For example, ecological niche switching from environmental sources to a human

host may introduce bottlenecks resulting in low diversity and effective population size. Another

explanation is a change in mutation rates, either through changes in bacterial generation time or

through adaptive genetic variation. A third consideration is sampling bias, where particular *Mab*

lineages that cause more frequent, severe and/or persistent disease are more likely to be isolated

and sequenced. Consistent with the latter hypothesis, previous work has shown that isolates

within clusters are more likely to contain point mutations associated with antibiotic resistance and

are associated with greater bacterial burden and lung inflammation in mice.[4]

Interpreting tree structures in the context of population history is aided by phylogenetic

dating, which translates genetic distances into time scale. Previous studies[4,9,12,13] have estimated

molecular clock rates in *Mab* but these have been limited to subpopulations within clusters. Here

we report on molecular clock rate estimation for the deep branches of the *M. a. abscessus*

phylogeny and for within-host isolates belonging to clusters. We provide evidence for an

evolutionary slow-down coincident with the expansion of clinically dominant clusters and propose

that the emergence of dense phylogenetic clustering in *Mab* may result from genetically encoded

changes to the mutation rate. These results indicate that transmission is not the only plausible

explanation for the dense clustering observed in the *Mab* phylogeny, and that more research is

needed to clarify the transmission dynamics of this emerging pathogen.

Results

<u>Phylogenetic analysis confirms the presence of large, dominant clusters (DCs) spanning multiple</u>

<u>geographies</u>. We identified 1,629 *M. abscessus* isolates with publicly available whole genome

sequencing (WGS) data and associated dates of collection (Table S1). We excluded isolates with

likely contamination or low-quality assemblies (Methods). Among the 1,461 isolates passing

quality filters, the majority (80.1%) were derived from *in vitro* culture of pulmonary samples. Other

samples were derived from a skin swab or biopsy, other body sites, and environmental sources

(Table S2). Of the 1,181 pulmonary isolates, 98.3% (1,161/1,181) were collected from patients

with cystic fibrosis (CF), while the rest were collected from patients with no documented diagnosis.

The dataset represented 483 patients with unique patient identifiers, with 59.6% (288/483) of

patients contributing a single isolate, and the remaining contributing between 2 to 24 isolates

sampled over time. Isolates originated from nine countries and were collected between 1998-

2017, with the exception of the reference strain ATCC 19977 which was collected in 1957 (Figure

1B).

The majority (91.7%) of samples had ≥98% average nucleotide identity (ANI) with one of

three reference genomes (Table S3) representing each of the three Mab subspecies, consistent

with the commonly used ANI threshold for bacterial subspecies. Isolates with <98% ANI with any

subspecies were excluded from downstream analysis. The resulting dataset overrepresented

clinically relevant subspecies of *M. abscessus*, with 93.9% belonging to either *M. a. abscessus*

or *M. a. massiliense*. To better resolve true evolutionary relationships among samples, we defined

and restricted analysis to the core genome for each subspecies, then further excluded predicted

recombination events (Methods). We performed maximum likelihood (ML) phylogenetic analysis

for each of the three *Mab* subspecies, using only the first isolate from each patient to focus on

between-host diversity. The resulting subspecies trees confirmed the presence of multiple,

globally distributed clades with dense phylogenetic clustering. Clusters were defined using a one-

dimensional scanning statistic that compared the branch lengths of a clade to the distribution of

branch lengths across the entire subspecies tree (Methods). Using this approach, we initially

identified 30 clusters. We further excluded clustering due to potential point source outbreaks by

limiting to clusters containing isolates from more than one country and a minimum of three isolates

for downstream analysis. In total, we confirmed 16 clusters (Figure S2A) including four clusters

with > 50 isolates, which we refer to here as dominant clusters (DCs). Isolates within clusters

comprised 76.4% of isolates included in the phylogenetic analysis. The median pairwise distance

in *M. a. abscessus* DC 1 was 68 SNPs. For comparison, the mean pairwise distance across the

subspecies trees were 5537, 4100, and 8034 SNPs for *M. a. abscessus*, *M. a. massiliense*, and

M. a. bolletii, respectively. All four DCs contained isolates originating from three continents (North

America, Europe, and Australia). Clustered isolates also spanned the full range of isolation times,

including ATCC 19977, which was placed within *M. a. abscessus* DC 1 despite having been

collected at least 40 years prior to any other isolates in the tree (Figure 1A-B).

<u>Coalescent analysis reveals a faster long-term evolutionary rate than observed in DCs</u>. To aid in

transmission inference and to reconstruct the evolutionary history of *Mab*, we performed a

coalescent analysis using our SNP alignment. Previous work has reported molecular clock rates

within *Mab* clusters, but to our knowledge no study has successfully dated any of the full *Mab*

subspecies phylogenies. To evaluate the presence of temporal signal, we performed root-to-tip

regression for each subspecies tree but in each case did not observe a positive correlation

between sampling date and root-to-tip distance. We reasoned that the high degree of population

structure in the *Mab* phylogeny may obscure the genetic-temporal signal. To test this hypothesis,

we pruned each subspecies tree until it contained the minimal number of samples required to

capture 95% of the genetic diversity in the original tree (Methods). Following pruning of the *M. a.*

abscessus subspecies tree, root-to-tip regression demonstrated the presence of positive temporal

signal (Figure 1C, D). Evidence of temporal signal in *M. a. abscessus* was further strengthened

using a permutation test and a Bayesian evaluation of temporal signal (Figure S2C, Table S4).

We used a Bayesian implementation of coalescent analysis, BEAST[14], to estimate the

molecular clock rate of the pruned *M. a. abscessus* subspecies tree assuming a relaxed clock

model, which allows each branch of the tree to have its own clock rate. To explore the effect of

the tree prior on molecular clock estimation, we ran analyses using two different tree priors,

assuming either a constant coalescent model, which assumes constant population size over time,

or a Bayesian skyline coalescent model, which allows the population size to change over time.

While the Bayesian skyline coalescent had a slightly better marginal likelihood, we found no

significant difference between the clock rates estimated by the two models (Table 1, Figure 2C).

Our estimates for the clock rate in the long, internal branches of the tree were approximately 10-

fold faster than nearly all previously reported estimates of the clock rate within *Mab* clusters

(Figure 2C). We next attempted to date the full (unpruned) *M. a. abscessus* phylogeny using the

faster rate inferred from our coalescent analysis of the pruned tree. These chains suffered from

poor convergence and sampled trees with a distorted topology, elongating the short branches

and eliminating the densely clustered structure observed in the ML phylogeny (data not shown).

We therefore hypothesized that the lack of temporal signal in the full *M. a. abscessus* dataset

may result from a difference between the long-term clock rate and the more recent evolutionary

rate within phylogenetic clusters.

Analysis of longitudinally sampled within-host isolates supports a slower clock rate in DCs. To

further investigate the differences in clock rate between long-term historical time scales and short-

term contemporary time, we used pairs of isolates sampled from the same patient over time to

estimate the mutation rate. We focused on estimating a within-host clock rate for *M. a. abscessus*

to compare directly with the estimate from our coalescent analysis and because we had the most

longitudinal samples for this subspecies. Because polyclonal *Mab* infection is common in CF

patients, we first excluded any isolate pairs with a SNP distance > 20. This threshold was selected

by comparing the distribution of all possible within-host isolate pairs to the distribution of pairwise

SNP distances within *Mab* clusters (Figure 2A, Figure S3). Regressing the SNP distance between

isolate pairs on the time between sample pairs yielded a mutation rate of 2.21 SNPs/year [1.06 –

3.37 95% CI], or 5.17×10^{-7} SNPs/site/year [2.48×10^{-7} – 7.92×10^{-7} 95% CI]. This value is more

comparable to previously reported molecular clock rate estimates within *Mab* clusters[4,9,12,13] than

to those estimated in the long branches of the phylogeny (Figure 2C).

<u>Simulated ancestries support mutation rate slow-down over population size effects as the major</u>

<u>driver of phylogenetic clustering</u>. The high core genome similarity of isolates within clusters has

been interpreted as supporting rapid global dissemination of *Mab* clones through direct

transmission, but may instead relate to population bottlenecks, oversampling, or a relatively slow

rate of molecular evolution in human-adapted clones. To test whether sampling from clades with

low genetic diversity is sufficient to explain the degree of clustering observed in the ML phylogeny,

or whether changes in the molecular clock rate are a better explanation for the observed

phylognetic data, we performed coalescent simulations of ancestries with one clustered

subpopulation (A) and one unclustered subpopulation (B). Under a range of effective population

sizes (N_e) and mutation rates (μ), we simulated ancestries through time with 1000 replicates per

simulation. For each replicate we assessed the degree of phylogenetic clustering by measuring

d_A/d_B where d_i is the mean pairwise SNP distance within subpopulation i. The total N_e (N_e^T),

number of samples, and sampling dates for subpopulations A and B were drawn from the true

values for *M. a. abscessus* DC 1 and from all "unclustered" isolates in our dataset, respectively

(Figure 3A). We first tested whether a smaller population size for the clustered subpopulation A

(N_e^A) could explain the degree of phylogenetic clustering observed. We estimated a range of

realistic values for N_e^A using BEAST by assuming a range of possible clock rates (Table S5).

These rates spanned a published clock rate estimate for *M. a. abscessus* DC1 to a clock rate that

is 50-fold higher, far exceeding the upper 95% highest posterior density (HPD) interval for the

published value.[13] Under the the simulated conditions of equal mutation rate between

subpopulation A and B, phylogenetic clustering to the extent observed in the clinical isolate

phylogeny was not a likely outcome (Figure 3B). We confirmed these finding in simulations where

the population sizes N_e^T spanned the 95% confidence intervals under both tree priors shown in

Table 1, except in the scenario where $N_e^T \gg N_e^A$ (Figure S4). We next tested whether a change

in the mutation rate at the root of supopulation A could better explain the observed phylogenetic

structure in *Mab*. The simulated ancestries where N_e^A and N_e^B are fixed and the mutation rate

slowdown is between 10- and 20-fold better recapitulated the observed phylognetic structure

(Figure 3C). This result was robust to changes in N_e^A and N_e^T across the 95% HPD for each

parameter (Figure S5). These results indicate that for person-to-person transmission to drive

phylogenetic clustering in *Mab*, the total N_e must be very large compared to the N_e of the clusterd

subpopulation, an unlikely scenario given the observed genetic data and sampling times. By

contrast, a mutation rate change of 10- to 20- fold explains the degree of clustering observed in

Mab under more realistic population sizes given these data.

<u>UvrD/Rep family helicases are enriched in DCs in *M. a. abscessus*</u>. To identify genetic variants

under positive selection in *Mab* clusters, we searched for evidence of phylogenetic convergence

in the core genome of *M. a. abscessus* with regions of recombination excluded. Briefly, we

performed ancestral sequence reconstruction for the internal nodes of the subspecies tree and

counted the number of independent arisals of each SNP (Methods). Of the 65,202 SNPs identified

in *M. a. abscessus*, 34,813 (53.4%) of mutations occurred only once and were excluded. For the

remaining SNPs, we ranked each by its enrichment in DCs relative to other regions of the *M. a.*

abscessus tree (Figure 4A). The most significantly enriched variants in the *M. a. abscessus* DCs

included homologs of several known or putative virulence factors as well as a number of cluster-

enriched mutations in UvrD-like helicases *MAB_1054*, *MAB_3515c*, and *MAB_3516c* (Table 2,

Figure 4A). While the function of these genes is not well characterized in *M. abscessus*, all three

share homology with the UvrD/Rep family of helicases. This family of genes is involved in DNA

repair and has been associated with growth and persistence *in vivo*.[15,16]

<u>Isolates in *Mab* DCs are associated with a lower spontaneous mutation rate</u>. To test the

association between phylogenetic clustering and spontaneous mutation rate, we selected *M. a.*

abscessus and *M. a. massiliense* strains that were grouped within DCs or that were not present

in those clusters. Each of the strains present in the DCs has the derived (minor) allele at four sites

in the UvrD/Rep family genes *MAB_1054*, *MAB_3515c*, and *MAB_3516c*, and the strains outside

of DCs possess the ancestral (major) allele at those sites (Table 2, Figure 4A). The strains used

are described in Table S6 and their genotypes are described in Table S7. The spontaneous

mutation rate was estimated in all strains using a fluctuation assay[17] adapted for use in *M.*

abscessus. The fluctuation assay counts cells gaining resistance to the antibiotic amikacin during

growth in culture without antibiotic. By normalizing the number of mutational events to the

population size of the culture, the fluctuation assay estimates the spontaneous mutation rate per

generation per the number of bases that can be mutated to confer resistance to amikacin. The

strains chosen display similar growth rates (Figure S6A) and very low baseline resistance to

amikacin (Table S8) which allows them to be compared using a fluctuation assay. Among *M. a.*

abscessus strains, those present in DCs with the derived (minor) allele at the four UvrD/Rep loci

had lower spontaneous mutation rates than those with the ancestral (major) allele (Figure 4C). A

similar trend was observed in *M. a. massiliense* strains (Figure 4D). Further, the average mutation

rate was significantly lower in isolates present in DCs across both subspecies (Figure 4E),

consistent with the hypothesis that the emergence of dense clustering may be driven in part by

changes in mutation rate.

Discussion

In this study we have pooled a large global sample of *Mab* genome sequences and

examined evolutionary rates and positive selection with the goal of understanding the emergence

of *Mab* clusters. We present evidence supporting a slower evolutionary rate within DCs compared

to the long internal branches of the phylogeny and further data associating adaptive genetic

variation in DNA repair genes with the slower evolutionary rate. We posit that changes to the
mutation rate underlie the dense clustering in *Mab*, and that while other effects such as population
or transmission bottlenecks and sampling bias may contribute to the low genetic diversity within
clusters, they are not sufficient to explain the observed phylogenetic structure.

Previous studies have restricted molecular clock rate estimation to *Mab* clusters, most
likely because of the lack of temporal signal at the subspecies-level. By pruning the *Mab* tree
while preserving the majority of the subspecies diversity, we were able to measure a molecular
clock rate for *Mab* spanning the time scale from the common ancestor of the subspecies to the
present. Estimates of evolutionary rates are recognized to depend on the time scale of
measurement. In such examples, the clock rate usually decreases with increasing time depth.[19]
Here we observe the opposite effect where the clock rate increases with time depth. The pruned
Mab tree most likely reflects an evolutionary process taking place predominantly in environmental
reservoirs, in contrast to phylogenetic clusters where terminal branches reflect evolution within
the human lung environment. We also note that our measurement of within-host mutation rate in
Mab was independent of phylogenetic inference and the full temporal signal in our sample.
Nevertheless, the measured rate within-host was comparable to the clock rate measured
previously using coalescent analysis within clusters (Figure 2C).

The molecular clock rate is influenced by both the spontaneous mutation rate and the
generation time of an organism. Changes in generation time could result from physiological
constraints of the human lung environment. For example, differences in nutrient availability may
change the growth rate in the lung compared to those living in environmental reservoirs. Longer
generation times for bacteria growing in the lung would result in a decrease in the molecular clock
rate. Ruis et al. reported differences in mutational signatures along internal branches of the
subspecies phylogeny compared to internal branches of DCs, supporting a change in the
mutational process within some clades of the tree.[13] The authors attributed these changes to
differences in mutagen exposure across niches. Generation times and spontaneous mutation

rates can also be modified by genetic changes, for example to DNA replication or repair

machinery. The observation that some lineages have adapted to the host environment[4] raises the

possibility that genetic changes that underlie or associate with this adaptation affect mutation rate,

either directly by affecting DNA replication or repair machinery, or indirectly by affecting the growth

rate.

We scanned the core genomes for variants enriched in the DCs in *M. a. abscessus*, the

subspecies for which we have the most WGS data and for which there are the highest number of

clusters. Notably, we observed several cluster-enriched mutations in UvrD-like helicases. UvrD is

a superfamily 1 helicase that is involved in DNA repair and is widely conserved in gram-negative

bacteria. UvrD1 in *M. tuberculosis* plays an important role both in nucleotide excision repair (NER)

and in pathogenesis and persistence because its DNA repair activity confers tolerance to reactive

oxygen intermediates (ROI) and reactive nitrogen intermediates (RNI)[15]. Mutations that promote

the activity of UvrD1 and its homologs are likely advantageous in the context of lung infection and

may have the additional consequence of decreasing the spontaneous mutation rate. We

additionally observed mutations in UvrD-like helicases occurring along the cluster-defining branch

of the large *M. a. massiliense* cluster, further supporting the role of the NER pathway in *Mab*

pathogenesis.

Other factors may contribute to the low genetic diversity within clusters. Clinical and

experimental data have shown that clustered *Mab* isolates are more likely to cause chronic

disease, have point mutations associated with antibiotic resistance, have increased intracellular

survival in macrophages, and cause greater bacterial burden and inflammation in mice.[4] Mab

genotypes that cause more frequent, persistent, or severe disease are more likely to be sampled

and sequenced, and dense sampling of closely related isolates may contribute to the appearance

of extreme clustering in the *Mab* phylogeny. Similarly, population bottlenecks due to transmission

or other factors may purge diversity and result in the observation of closely related isolates. While

epidemiological evidence does not currently support transmission as a major driver of *Mab*

dissemination, isolated examples of putative recent transmission events have been described,

which may contribute to the observation of dense phylogenetic clustering independent of

bottlenecks. However, the results of our ancestry simulations suggest that population size or

sampling effects alone do not sufficiently account for the observed phylogenetic structure in *Mab*

and instead favor mutation rate changes as a more likely explanation.

Our study has several limitations. Our dataset overrepresented clinical isolates from the

UK where routine whole genome sequencing is part of *Mab* surveillance. Oversampling of some

lineages and geographies may mean that we have not adequately captured the global population

structure and that we may have focused on large phylogenetic clusters because of their

geographic distribution and not because of their clinical significance. While our study argues

against transmission as the main driver of *Mab* dissemination and the emergence of DCs, we do

not examine the geographic distribution of Mab within clusters. It is possible that future work

considering this data can highlight mechanisms of parallel evolution that explain why clustered

Mab isolates are detected in distant geographic locations. While our results support a faster

molecular clock rate in the long internal branches of the phylogeny compared to clustered

lineages, we are unable to distinguish between rate differences due to the spontaneous mutation

rate from generation time effects. Differences in growth conditions across environments is difficult

to quantify as *Mab* is ubiquitous in both natural and built environments, but nutrient availability

likely differs greatly between natural reservoirs such as soil and water compared to man-made

reservoirs such as showerheads. A healthy lung supports relatively little bacterial growth, but

inflammation increases nutrient availability by promoting mucus production and vascular leakage

into the airway.[20] Because there is little publicly available sequencing data from environmental

isolates, we were also unable to compare genotypes or mutation rates in environmental samples

compared to clinical isolates. Our analysis of variants associated with phylogenetic clustering is

also limited in its ability to identify true causal genotypes or those that are enriched in clusters due

to hitchhiking to other adaptive variants.

In summary, we present new evidence for changes in the molecular clock rate that

contribute to the dense phylogenetic clustering observed in *Mab*. These results argue against

person-to-person transmission as the primary driver of clustering in the *Mab* phylogeny.

Continuing genomic surveillance is needed to understand how *Mab* may be adapting to human

infection and to characterize the risk of transmission to at-risk patients. We argue further that the

phylogeographic structure of *Mab* is influenced by its complex ecological history as an emerging

facultative pathogen that should be considered carefully in future analysis of *Mab* genomic data.

Future work should extend these analyses by examining differences in mutation rates and

phenotypes between clinical isolates and environmental strains for evidence of host adaptation.

Acknowledgements

We thank members of the Farhat lab for feedback and discussion. We are grateful to Luca Freschi

for advice on processing WGS data, calling variants, and phylogenetic inference; Maximillian

Marin for input on calculating mappability pileup; and Roger Vargas for support in homoplasy

counting. We also thank Wilder Wohns for discussions on implementing ancestry simulations with

mutation rate changes through time. M.R.S. is a Merck Fellow of the Damon Runyon Cancer

Research Foundation, DRG-2415-20. Computational resources and support were provided by the

Orchestra High Performance Compute Cluster at Harvard Medical School, which is funded by the

NIH (NCRR 1S10RR028832-01).

Author Contributions

Conceptualization, N.C. and M.F.; Methodology – Computational analysis, N.C., E.K., and M.F.;

Methodology – Experimental studies, M.R.S., K.M.; Investigation – Computational analysis, N.C.;

Investigation – Experimental studies, M.R.S., K.M.; Data Curation, N.C.; Writing – Original Draft,

N.C. and M.R.S.; Writing – Review & Editing, N.C., M.R.S., E.K., M.F.; Supervision, E.J.R., M.F.

Declaration of Interests

The authors declare no competing interests.

360   **Tables**

362   **Table 1**. Coalescent analysis results and model comparison

Model	Marginal Log Likelihood (SD)	UCLD Rate (SNPs/site/year)	UCLD St. Dev.
Coalescent constant population	-7144301.64 (21.42)	2.82×10^{-6} [1.33×10^{-6}, 4.83×10^{-6}]	0.182 [0.135, 0.234]
Bayesian skyline coalescent	-7144118.76 (19.93)	2.63×10^{-6} [1.16×10^{-6} , 4.0×10^{-6}]	0.181 [0.138, 0.232]

Table 2. Summary of top 10 nonsynonymous variants enriched in DCs of *M. a. abscessus*

Gene	Gene Symbol	Description	Comments	Enrichment p-value	Number of clustered taxa containing variant	Number of unclustered taxa containing variant
MAB_3334c	*gatB* [c]	Aspartyl/glutamyl-tRNA (Asn/Gln) amidotransferase subunit B [c,d]	Essential for growth and survival *in vivo* in *M. tuberculosis* [17]	1.03×10^{-65}	240	1
MAB_3244		Mercury resistance transport protein/cytochrome c biogenesis protein [c]		1.72×10^{64}	189	3
MAB_3480		Putative short chain dehydrogenase/Reductase [c]		7.23×10^{-64}	232	13
MAB_4141[a]		PE/PPE family protein [c]		1.58×10^{-62}	189	2
MAB_3242	*fni* [c]	Isopentenyl-diphosphate delta-isomerase [c,d]		8.07×10^{-62}	189	3
MAB_3481	*fadE* [c]	Probable acyl-CoA dehydrogenase [c]	FadE mutations in *Mtb* result in decreased growth and virulence in macrophages and in mice[18]	2.10×10^{-61}	232	15
MAB_4057c	*mshA* [c]	D-inositol 3-phosphate glycosyltransferase [d]	Catalyzes biosynthesis of mycothiol, an important compound in resisting oxidative stress and detoxifying harmful agents such as antibiotics[19]	2.78×10^{-60}	189	2
MAB_3581c		Hypothetical protein		6.09×10^{-60}	169	3
MAB_1054[b]	*pcrA*[c] / *uvrD1*[d]	ATP-dependent DNA helicase [c,d]	Involved in nucelotide excision repair (NER), mismatch repair (MMR), homologous recombination (HR), and rolling circle replication[15]	2.72×10^{-58}	216	7
MAB_3515c	*rep* [c,d]	ATP-dependent DNA helicase [c,d]	Involved in NER, MMR, HR and rolling circle replication[15]	8.42×10^{-57}	189	7

Table 2 Legend:
[a] Multiple variants in *MAB_4141* have been collapsed and the lowest p-value is shown. See
Supplementary Material for full list of homoplasic SNPs ranked by their enrichment in clusters.
[b] Gram-negative bacteria possess UvrD and a closely related homolog, Rep, while gram-positive
bacteria possess a single homolog called PcrA.[18]
[c] Mycobrowser annotation
[d] prokka annotation
457

Figures

 Figure 1. Dominant clusters of *M. abscessus* and the presence of temporal signal in internal branches. A) Species-wide phylogeny of *M. abscessus* representing one isolate per patient to avoid within-patient clustering of samples. Each subspecies tree was constructed independently using a subspecies specific reference genome. The four largest phylogenetic clusters are highlighted. B) Histogram showing the distribution of collection dates of all clustered isolates v. all unclustered isolates in *M. a. abscessus*. C) Pruned *M. a. abscessus* phylogeny representing 95% of the original subspecies tree diversity. C) Root-to-tip regression showing evidence of positive temporal signal in the pruned tree shown in (B).

Figure 2. Longitudinally sampled within-host isolates and coalescent analysis support a slower clock rate within DCs. A) Clock rate estimation using isolate pairs sampled from the same patient over time. Left: distribution of SNP distances between all possible within-host isolate pairs compared to SNP distances between all possible isolate pairs within *M. a. abscessus* DC 1. The orange line represents the threshold of 20 SNPs used to exclude pairs representing independent infections. Right: regression of SNP distance on the time between isolates for each isolate pair. The slope of the regression line was 2.51 SNPs/year [1.24 – 3.78 95% CI], or roughly 5.93×10^{-7} SNPs/site/year [2.92×10^{-7} – 8.9×10^{-7} 95% CI]. B) UCLD clock rate estimates for the three models used in this study (two coalescent models shown in red and one regression model shown in blue) compared to mutation rate estimates from the within-host regression (blue) and published clock rates (grey).

Figure 3. Simulated ancestries support a mutation rate slow-down coincident with DC emergence. A) Example neighbor joining (NJ) tree from the simulation shown in B. The number of samples and sampling dates for subpopulations A and B are drawn from real data from *M. a. abscessus* cluster 1 and from all unclustered samples, respectively. The dotted line crosses the common ancestor node for subpopulation A, and all branches descended from this node (shown here in red) are subject to the change in mutation rate. B) Boxplot showing the degree of phylogenetic clustering in subpopulation A relative to subpopulation B over a range of effective population sizes of subpopulation A (N_e^A). The simulation assumes that N_e^T=8000. C) Boxplot showing the degree of phylogenetic clustering in subpopulation A relative to subpopulation B over a range of mutation rate changes. The simulation assumes that N_e^T=8000 and N_e^A=718. In both A) and B) the degree of clustering is estimated with metric $\log(\frac{d_A}{d_B})$ where d_i is the mean pairwise SNP distance among all sample pairs in subpopulation i. The red line represents the clustering metric estimated using the observed ML phylogeny.

Figure 4. Dominant clusters are associated with a decrease in the spontaneous mutation rate *in vitro*. A) Subspecies phylogeny for *M. a. abscessus* (left) and corresponding heatmap representing the presence of the derived allele (blue) or ancestral allele (grey) for each nonsynonymous variant. Variants are ranked in ascending order by their p-value derived from Fisher's tests for enrichment in the three largest clusters. The top 50 hits are shown. The genes where each variant is located are denoted. B) Subspecies phylogeny for *M. a. massiliense*. The branches where mutations have occurred in homologs of Rep and UvrD2 are shown with asterisks. Probability of acquiring an amikacin resistance mutation per generation in C) *M. a. abscessus* strains, D) *M. a. massiliense* strains, or D) in all strains tested that are part of a dominant cluster or not. Mean +/- SD is displayed. In C-D, n=3 biological replicates. In E, each point represents the average mutation probability for a single strain. p-value derived from two-tailed, unpaired t-test.

Methods

Data Sources and Description: WGS data were curated from published studies[4,7,10,12,21–23] and additional data were downloaded from the NCBI Sequence Read Archive (SRA). A search of "abscessus" was conducted on October 4, 2019 using the filters "DNA" and "Illumina". Sequences were selected that had an associated date of collection with the year at minimum. Collection dates were obtained either from the literature, SRA metadata table, or provided by the authors. A summary of all samples used in this study is provided in Table S1. Raw sequencing reads were downloaded using the NCBI SRA toolkit[24]. Our procedure for downloading and processing sequencing reads is available as a snakemake pipeline as well as all custom scripts are available at https://github.com/nicolettacommins/mab_mutation_rates_2022.

Quality control of Illumina reads: Reads were trimmed and filtered with trimmomatic[25]. To detect contaminated samples, the mean GC content of the reads were compared to a modelled normal distribution of GC content using fastqc[26]. If the sum of the deviations from the normal distribution represented more than 30% of the reads, the sequencing run was excluded from further analysis.

Subspecies assignment and genome assembly: Multiple sequencing runs corresponding to one biological sample were combined into one file. Reads were then assembled *de novo* using

SPAdes[27]. Assemblies were compared to a reference sequence for each of the three subspecies

(Table S3) and a whole-genome based average nucleotide identity (gANI) score was calculated

using fastANI[28]. The resulting alignments were assigned to a subspecies based on having gANI

of at least 98% with exactly one reference strain. Isolates that did not have at least 98% ANI with

any reference strain were excluded. No isolates matched more than one reference genome. After

assigning isolates to a subspecies, reads were mapped to the corresponding reference genome

using BWA MEM[29]. After alignment, isolates with <80% coverage of the reference genome at a

depth of at least 20x or with missing data at >30% of sites were excluded.

<u>Variant calling and SNP filtering</u>: Variants were called using Pilon[30]. We excluded ambiguous calls

as wells as indels and MNVs. We also excluded calls with low coverage using a minimum depth

of either 10% of the mean coverage, or 5, whichever was greater (default settings for Pilon –

mindepth). We used bcftools[31] to exclude calls with mapping quality or base quality scores < 20.

<u>Core genome inference</u>: Unlike the professional mycobacterial pathogens *M. tuberculosis* or *M.*

leprae, which have small accessory genomes, *M. abscessus* has a large, open accessory

genome.[32] To define the core genome for phylogenetic analysis and to exclude loci that are highly

divergent from the reference sequence, we masked all loci for downstream analysis where read

depth was <20x in more than 5% of the isolates in our sample set. The remaining loci were

compared to an estimate of the core genome calculated by Roary[33], and the two methods yielded

nearly identical results. We further excluded regions with average mapping quality scores or

average based quality scores <20. Finally, we used GenMap[34] to calculate mappability scores

across each reference genome as defined in reference[35]. We then calculated a mappability pileup

score for each base position as described in reference[36] and excluded loci where the mappability

pileup score was <95% from downstream analysis. In total, we excluded 14.4%, 13.5%, and

15.1% of the *M. a. abscessus*, *M. a. massiliense*, and *M. a. bolletii* reference genome length,

respectively.

<u>Phylogenetic tree inference</u>: A full-length sequence alignment was created using a custom script,

masking loci excluded by our filtering criteria described above. Gubbins[37] was used to predict and

mask regions of variation produced by recombination from the full-length alignment. A

recombination-free SNP alignment generated by Gubbins was then used to build a phylogeny for

each subspecies using IQ-TREE[38], using the ModelFinder Plus (-mfp) option and 1000 bootstrap

replicates. Clusters were defined as previously described[4] using a one-dimensional scanning

statistic to identify clades with a distribution of branch lengths that is shorter than the distribution

of branch lengths in the rest of the tree[39].

<u>Coalescent analysis</u>: Temporal signal was assessed using root-to-tip regression. In both *M. a.*

abscessus and *M. a. massiliense* subspecies trees and all three major clusters, a slightly negative

slope was observed, indicated lack of temporal signal. We recognized that the high degree of

clustering and short time span over which the clustered isolates were sampled may confound the

genetic temporal signal. To address this, we pruned the trees down to taxa that preserved 95%

of the relative tree length (RTL) of the original trees using Treemmer[40]. A threshold of 95%

eliminated most of the dense clustering, but preserved the long branches of the original tree

(Figure 2B, Figure S2B). Using the pruned tree, we observed evidence of temporal signal in the

MAB subspecies using root-to-tip regression. We generated a SNP alignment including only the

taxa present in the pruned *M. a. abscessus* tree and used this as input for coalescent analysis

using BEAST2. ModelTest-NG[41] was used to select the site model GTR+4. MegaX[42] was used to

perform a maximum likelihood test to determine that a relaxed clock model is best suited for our

data. This choice was confirmed using the nested sampling algorithm to compare marginal

likelihoods for a strict v. relaxed clock model. We then used nested sampling to compare various

tree priors. We using Bayesian Evaluation of Temporal Signal (BETS) to confirm the presence of

temporal signal by comparing marginal likelihoods when the tip dates were used to those when

all samples were assumed to be isochronous (Table S4). Based on marginal likelihood values,

we ran both a coalescent constant population model and a Bayesian skyline coalescent model,

assuming a relaxed clock and including the sampling dates for both models. We used the ML

phylogeny of the pruned dataset as a starting tree. For each model we ran two independent chains

for at least 100 million states to ensure convergence.

Phylogenetic dating: To attempt to date the full MAB phylogeny, we ran both coalescent constant

population and Bayesian skyline coalescent models assuming a relaxed clock without sampling

dates. We set the mean UCLD clock rate to the values estimated from our coalescent analysis of

the pruned tree. We ran each chain for at least 100 million states.

Estimation of mutation rate within-host: We identified 90 patients with two or more *M. a. abscessus*

isolates. Because coinfection with multiple *M. abscessus* clones is common in cystic fibrosis

patients, we first sought to exclude any isolate pairs that represented distinct clonal infections.

We selected a threshold of 20 SNPs by comparing the distribution of all possible within-host

isolate pairs to the distribution of pairwise SNP distances within *Mab* clusters. We further excluded

any isolates where >20% of the genome is predicted to be rearranged by recombination. For the

remaining isolate pairs we regressed the pairwise SNP distance between the first and last isolates

sampled from each patient on the time between the collection of each sample. The slope of the

regression line was used as an estimate of the number of SNPs accumulated over time.

Ancestry simulations: We used msprime[43] to simulate ancestries to model two subpopulations:

(1) a "clustered" subpopulation (A) and (2) an "unclustered" subpopulation (B). We assumed a

constant population size and initially set the total effective population size (N_e) to 8000, the N_e

estimated in our coalescent analysis of the pruned *Mab* tree. We chose a range of realistic N_e for

subpopulation A, N_e^A by running a separate coalescent analysis of *M. a. abscessus* cluster 1 using

BEAST, assuming a constant population size, relaxed clock, and a range of UCLD clock rates

spanning 1.7×10^{-7} SNPs/site/year, as previously reported for this cluster in ref[13], to 8.5×10^{-6}

SNPs/site/year, a value well above the 95% HPD interval for our estimate of the clock rate in the

pruned tree (Table S5). We assumed the N_e of subpopulation B to be: $N_e^B = N_e^{total}-N_e^A$. The number

of samples and sampling dates supplied to the model for subpopulations A and B are drawn from

the true values for *Mab* cluster 1 and from all "unclustered" isolates in the tree, respectively. We

assumed an initial mutation rate of 1×10^{-9} mutations/generation with a sequence length of 5Mb

and a divergence time of 30,000 generations between subpopulations A and B. We also assumed

the nucleotide subsitution model estimated using BEAST.

We first simulated a scenario where the mutation rate remains constant, varying N_e^A over

the range shown in Table S5. For each simulated ancestry we overlaid mutation events, supplying

the model with a starting substitution rate of 1×10^{-9} mutations/generation, a nucleotide substitution

model estimated using BEAST, and a sequence length of 5Mb. We then extracted the simulated

genotypes and constructed a neighbor joining (NJ) tree. The NJ tree was used to exclude any

simulations where the two subpopulations did not exhibit reciprocal monophyly. We used the

extracted genotypes to estimate the degree of clustering in each tree by comparing the mean

pairwise SNP distance in subpopulation A (d_A) to the mean pairwise SNP distance in

subpopulation B (d_B). For each simulated condition we performed 1000 replicates. To simulate

ancestries with a mutation rate change, we repeated the above procedure conservatively

assuming $N_e^A = 718$ (Table S5). We modeled a per-generation mutation rate change starting at

the common ancestor node for subpopulation A and ranging from 1-fold (no change) to 1/25-fold

spanning a range of reasonable rate changes based on the range of clock rate estimates

observed in this study and reported in the literature (Figure 2B). For all simulated conditions, we

the simulation over a range of values for N_e^{total} spanning the 95% HPD intervals for the N_e

estimates using the Treemmer tree under both the constant coalescent and coalescent skyline

tree priors.

<u>Scan for variants enriched in host-adapted lineages</u>: To search for variants associated with host-

adapted MAB lineages, we first inferred the number of independent arisals in the MAB phylogeny

for each SNP using SNPPar[44]. Briefly, SNPPar uses TreeTime to perform an ancestral sequence

reconstruction at each node, then infers mutation events arising on each branch of the tree. We

quantified the number of independent mutation events for each SNP. We considered only

mutation events where the derived call was the minor allele and the ancestral call was the major

allele. Because of the extreme phylogenetic clustering in *Mab*, we defined the major and minor

alleles using the pruned alignment generated by Treemmer. For all homoplasic SNPs (occurring

two or more times independently in the tree), we used Fisher's exact test to quantify the

enrichment of that variant within phylogenetic clusters. We conducted two separate analyses: a)

focusing on the three largest clusters in the *M. a. abscessus* tree (described in the main text), and

b) considering all clusters identified by this approach. In each case we obtained similar results.

For both analyses, full lists of variants ranked by their enrichment scores are provided in

Supplementary Data. We also used SNPPar to identify all mutation events occurring on the

branch ancestral to *M. a. massiliense* cluster 1.

M. abscessus <u>culture</u>: *M. abscessus* clinical isolates used for fluctuation assays were isolated

from patients in Taiwan and in Boston, MA, USA. All *M. abscessus* strains were cultured in

Middlebrook 7H9 broth (271310, BD Diagnostics, Franklin Lakes, NJ, USA) with 0.2% (v/v)

glycerol, 0.05% (v/v) Tween-80 (P1754, Sigma-Aldrich, St. Louis, MO, USA), and 10% (v/v)

OADC (90000-614, VWR, Radnor, PA, USA). Proliferation rates were determined by inoculating

5 mL media with 500,000 colony forming units (cfu), then measuring OD600 at 24 and 48 hours

post-inoculation. Proliferation rate was calculated as follows: proliferation rate in doublings per

day = $\log_2$(OD600 at 48 hours / OD600 at 24 hours).

Fluctuation Assay: Fluctuation assays were performed as described in reference[17] with

adaptations for *M. abscessus*. *M. abscessus* strains were grown to saturation, then diluted to an

OD600 < 0.01 and grown overnight until cultures reached an OD600 = 0.6-0.8. These cultures

were used for fluctuation assays as well as determination of baseline amikacin sensitivity.

Baseline amikacin sensitivity was tested by spreading dilutions of 10 million, 100 million, or 1

billion cfu on agar 7H10 (262710, BD Diagnostics) + OADC (90000-614, VWR) + 300 μg/mL

amikacin sulfate (A2324, Sigma-Aldrich) plates. For fluctuation assays, 5000 cfu were plated in 1

mL media in each of 21 wells of a 24-well culture plate (10861-558, VWR), and initial cultures

were plated for cfu on 7H10 + OADC to confirm number of cells in initial culture. 24-well plates

were incubated shaking at 150 rpm for 6 days at 37°C. 18 wells of each strain were plated onto

6-well plates containing 5 mL 7H10 agar + OADC + 300 μg/mL amikacin sulfate in each well.

Amikacin was used as the antibiotic selection for the assay because all tested strains had low

baseline resistance (Table S8), which is required for the validity of the fluctuation assay. The

remaining 3 wells from each strain were diluted and plated on 7H10 + OADC plates to enumerate

average final cfu. Colonies on all agar plates were counted after 5 days incubation at 37°C.

Mutation rate per generation per total number of bases that can be mutated to confer amikacin

resistance was calculated using the Shinyflan application for the flan R package[17] using the

Maximum Likelihood estimation method, assumption of unknown fitness effects of mutations, the

Exponential (LD model) distribution of mutant lifetime, and a Winsor parameter of 1024. One

strain (BWH-E) was excluded from analysis due to highly variable results across replicates (Figure

S6B).

References

1. Adjemian, J., Olivier, K. N. & Prevots, D. R. Epidemiology of Pulmonary Nontuberculous Mycobacterial Sputum Positivity in Patients with Cystic Fibrosis in the United States, 2010-2014. *Ann. Am. Thorac. Soc.* **15**, 817–826 (2018).

2. Lee, M.-R. *et al.* Mycobacterium abscessus Complex Infections in Humans. *Emerg. Infect. Dis.* **21**, 1638–1646 (2015).

3. Nessar, R., Cambau, E., Reyrat, J. M., Murray, A. & Gicquel, B. Mycobacterium abscessus: a new antibiotic nightmare. *J. Antimicrob. Chemother.* **67**, 810–818 (2012).

4. Bryant, J. M. *et al.* Emergence and spread of a human-transmissible multidrug-resistant nontuberculous mycobacterium. *Science* **354**, 751–757 (2016).

5. Bange, F. C., Brown, B. A., Smaczny, C., Wallace, R. J., Jr & Böttger, E. C. Lack of transmission of mycobacterium abscessus among patients with cystic fibrosis attending a single clinic. *Clin. Infect. Dis.* **32**, 1648–1650 (2001).

6. Olivier, K. N. *et al.* Nontuberculous mycobacteria. I: multicenter prevalence study in cystic fibrosis. *Am. J. Respir. Crit. Care Med.* **167**, 828–834 (2003).

7. Doyle, R. M. *et al.* Cross-transmission is not the source of new Mycobacterium abscessus infections in a multi-centre cohort of cystic fibrosis patients. *Clin. Infect. Dis.* (2019) doi:10.1093/cid/ciz526.

8. Sermet-Gaudelus, I. *et al.* Mycobacterium abscessus and children with cystic fibrosis. *Emerg. Infect. Dis.* **9**, 1587–1591 (2003).

9. Lipworth, S. *et al.* Epidemiology of Mycobacterium abscessus in England: an observational study. *The Lancet Microbe* (2021) doi:10.1016/S2666-5247(21)00128-2.

10. Harris, K. A. *et al.* Whole-genome sequencing and epidemiological analysis do not provide evidence for cross-transmission of mycobacterium abscessus in a cohort of pediatric cystic fibrosis patients. *Clin. Infect. Dis.* **60**, 1007–1016 (2015).

11. Tortoli, E. *et al.* Mycobacterium abscessus in patients with cystic fibrosis: low impact of inter-

human transmission in Italy. *Eur. Respir. J.* **50**, (2017).

12. Everall, I. *et al.* Genomic epidemiology of a national outbreak of post-surgical Mycobacterium

abscessus wound infections in Brazil. *Microb Genom* **3**, e000111 (2017).

13. Ruis, C. *et al.* Dissemination of Mycobacterium abscessus via global transmission networks.

Nature Microbiology 1–10 (2021).

14. Bouckaert, R. *et al.* BEAST 2.5: An advanced software platform for Bayesian evolutionary

analysis. *PLoS Comput. Biol.* **15**, e1006650 (2019).

15. Houghton, J. *et al.* Important role for Mycobacterium tuberculosis UvrD1 in pathogenesis and

persistence apart from its function in nucleotide excision repair. *J. Bacteriol.* **194**, 2916–2923

(2012).

16. Sinha, K. M., Unciuleac, M.-C., Glickman, M. S. & Shuman, S. AdnAB: a new DSB-resecting

motor-nuclease from mycobacteria. *Genes Dev.* **23**, 1423–1437 (2009).

17. Krašovec, R. *et al.* Measuring Microbial Mutation Rates with the Fluctuation Assay. *J. Vis.*

Exp. (2019) doi:10.3791/60406.

18. Curti, E., Smerdon, S. J. & Davis, E. O. Characterization of the helicase activity and substrate

specificity of Mycobacterium tuberculosis UvrD. *J. Bacteriol.* **189**, 1542–1555 (2007).

19. Ho, S. Y. W., Shapiro, B., Phillips, M. J., Cooper, A. & Drummond, A. J. Evidence for time

dependency of molecular rate estimates. *Syst. Biol.* **56**, 515–522 (2007).

20. Huffnagle, G. B., Dickson, R. P. & Lukacs, N. W. The respiratory tract microbiome and lung

inflammation: a two-way street. *Mucosal Immunol.* **10**, 299–306 (2017).

21. Bryant, J. M. *et al.* Whole-genome sequencing to identify transmission of Mycobacterium

abscessus between patients with cystic fibrosis: a retrospective cohort study. *Lancet* **381**,

1551–1560 (2013).

22. Hasan, N. A. *et al.* Population Genomics of Nontuberculous Mycobacteria Recovered from

United States Cystic Fibrosis Patients. *bioRxiv* 663559 (2019) doi:10.1101/663559.

23. Shaw, L. P. *et al.* Children with cystic fibrosis are infected with multiple subpopulations of

Mycobacterium abscessus with different antimicrobial resistance profiles. *Clin. Infect. Dis.*

(2019) doi:10.1093/cid/ciz069.

24. Download : Software : Sequence Read Archive : NCBI/NLM/NIH.

https://trace.ncbi.nlm.nih.gov/Traces/sra/sra.cgi?view=software.

25. Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina sequence

data. *Bioinformatics* **30**, 2114–2120 (2014).

26. Babraham Bioinformatics - FastQC A Quality Control tool for High Throughput Sequence

Data. https://www.bioinformatics.babraham.ac.uk/projects/fastqc/.

27. Prjibelski, A., Antipov, D., Meleshko, D., Lapidus, A. & Korobeynikov, A. Using SPAdes De

Novo Assembler. *Curr. Protoc. Bioinformatics* **70**, e102 (2020).

28. Jain, C., Rodriguez-R, L. M., Phillippy, A. M., Konstantinidis, K. T. & Aluru, S. High throughput

ANI analysis of 90K prokaryotic genomes reveals clear species boundaries. *Nat. Commun.*

9, 5114 (2018).

29. Li, H. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.

arXiv [q-bio.GN] (2013).

30. Walker, B. J. *et al.* Pilon: an integrated tool for comprehensive microbial variant detection

and genome assembly improvement. *PLoS One* **9**, e112963 (2014).

31. Li, H. A statistical framework for SNP calling, mutation discovery, association mapping and

population genetical parameter estimation from sequencing data. *Bioinformatics* **27**, 2987–

2993 (2011).

32. Choo, S. W. *et al.* Genomic reconnaissance of clinical isolates of emerging human pathogen

Mycobacterium abscessus reveals high evolutionary potential. *Sci. Rep.* **4**, 4061 (2014).

33. Page, A. J. *et al.* Roary: rapid large-scale prokaryote pan genome analysis. *Bioinformatics*

31, 3691–3693 (2015).

34. Pockrandt, C., Alzamel, M., Iliopoulos, C. S. & Reinert, K. GenMap: ultra-fast computation of

genome mappability. *Bioinformatics* **36**, 3687–3692 (2020).

35. Derrien, T. *et al.* Fast computation and applications of genome mappability. *PLoS One* **7**,

e30377 (2012).

36. Marin, M. *et al.* Benchmarking the empirical accuracy of short-read sequencing across the

M. tuberculosis genome. *Bioinformatics* (2022) doi:10.1093/bioinformatics/btac023.

37. Croucher, N. J. *et al.* Rapid phylogenetic analysis of large samples of recombinant bacterial

whole genome sequences using Gubbins. *Nucleic Acids Res.* **43**, e15 (2015).

38. Minh, B. Q. *et al.* IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference

in the Genomic Era. *Mol. Biol. Evol.* **37**, 1530–1534 (2020).

39. Harris, S. TreeGubbins. (2016).

40. Menardo, F. *et al.* Treemmer: a tool to reduce large phylogenetic datasets with minimal loss

of diversity. *BMC Bioinformatics* **19**, 164 (2018).

41. Darriba, D. *et al.* ModelTest-NG: A New and Scalable Tool for the Selection of DNA and

Protein Evolutionary Models. *Mol. Biol. Evol.* **37**, 291–294 (2020).

42. Kumar, S., Stecher, G., Li, M., Knyaz, C. & Tamura, K. MEGA X: Molecular Evolutionary

Genetics Analysis across Computing Platforms. *Mol. Biol. Evol.* **35**, 1547–1549 (2018).

43. Kelleher, J. & Lohse, K. Coalescent Simulation with msprime. in *Statistical Population*

Genomics (ed. Dutheil, J. Y.) 191–230 (Springer US, 2020).

44. Edwards, D. J., Duchêne, S., Pope, B. & Holt, K. E. SNPPar: identifying convergent evolution

and other homoplasies from microbial whole-genome alignments. *bioRxiv*

2020.07.08.194480 (2020) doi:10.1101/2020.07.08.194480.