Quantitative environmental DNA metabarcoding shows high potential as a novel approach

to quantitatively assess fish community

Satsuki Tsuji[1,2]*, Ryutei Inui[3], Ryohei Nakao[2], Seiji Miyazono[2], Minoru Saito[2,4], Takanori Kono[2,5], Yoshihisa

Akamatsu[2] (*corresponding author)

[1] Graduate School of Science, Kyoto University, Kitashirakawa-Oiwakecho, Sakyo-ku, Kyoto 606–8502, Japan

[2] Graduate School of Science and Technology for Innovation, Yamaguchi University, 2-16-1 Tokiwadai, Ube, Yamaguchi, 755–8611, Japan

[3] Faculty of Socio-Environmental Studies, Fukuoka Institute of Technology, Wajiro-higashi, Higashi-ku, Fukuoka 811–0295, Japan

[4] Fisheries Division, Japan International Research Center for Agricultural Sciences, 1-1, Ohwashi, Tsukuba, Ibaraki 305–8686, Japan

[5] Aqua Restoration Research Center, Public Works Research Institute, National Research and Development Agency, Kawashima, Kasada-machi,

Kakamigahara, Gifu 501–6021, Japan

Highlights

● eDNA analysis and capture surveys of fishes were conducted at a total of 21 sites.

● Quantitative eDNA metabarcoding (qMiSeq) successfully quantified the eDNA of fishes.

● For each site, the estimated eDNA conc. reflected the abundance and biomass of fishes.

● For each taxon, the estimated eDNA conc. were comparable among sites.

● qMiSeq is a suitable and useful tool for quantitative monitoring of fish community.

Keywords: Environmental DNA, Quantitative eDNA metabarcoding, qMiSeq, MiFish, Electrical

shocker, Fish community

Abstract

The simultaneous conservation of species richness and evenness is important to effectively

reduce biodiversity loss and keep ecosystem health. Environmental DNA (eDNA) metabarcoding

has been used as a powerful tool for identifying community composition, but it does not

necessarily provide quantitative information due to several methodological limitations. Thus, the

quantification of eDNA through metabarcoding is an important frontier of eDNA-based

biomonitoring. Particularly, the qMiSeq approach has recently been developed as a quantitative

metabarcoding method and has attracted much attention due to its usefulness. The aim here was

to evaluate the performance of the qMiSeq approach as a quantitative monitoring tool for fish

communities by comparing the quantified eDNA concentrations with the results of fish capture

surveys. The eDNA water sampling and the capture surveys using the electrical shocker were

conducted at a total of 21 sites in four rivers in Japan. As a result, we found significant positive

relationships between eDNA concentrations of each species quantified by qMiSeq and both the

abundance and biomass of each captured taxon at each site. Furthermore, for seven out of eleven

taxa, a significant positive relationship was observed between quantified DNA concentrations by

sample and the abundance and/or biomass. In total, our results demonstrated that eDNA

metabarcoding with the qMiSeq approach is a suitable and useful tool for quantitative monitoring

of fish communities. Due to the simplicity of the eDNA analysis, the eDNA metabarcoding with

qMiSeq approach would promote further growth of quantitative monitoring of biodiversity.

Introduction

Conspicuous loss in biological diversity is a current and fast-emerging global problem resulting in changes of ecosystem functions (Cardinale et al., 2012; Dornelas et al., 2014; Magurran et al., 2018). To effectively reduce biodiversity loss, a number of previous studies have argued for the importance of assessing and conserving ecosystem health using species abundance, evenness and richness as indicators (Blowes et al., 2022; Crowder et al., 2012; Hillebrand et al., 2008). Accordingly, researchers and resource managers have made efforts to quantitatively estimate biodiversity based on conventional survey methods, such as direct capture and visual census (Masuda et al., 2010; Miyazono et al., 2015; Yonekura et al., 2004). While these survey approaches provide us with valuable data, they also require the large amount of effort, time and expertise, which limits the feasibility and continuity of the research itself and the reliability of the data (Evans et al., 2017; Miya et al., 2020; Oka et al., 2021; Thomsen et al., 2012). Additionally, especially for endangered species, sampling activities in direct capture surveys may damage target-species populations and/or their habitats (Pimm et al., 2015; Rourke et al., 2022; Tsuji et al., 2018). To overcome these difficulties in the traditional survey methods, new approaches for accurate and effective biodiversity monitoring are being explored (Kissling et al., 2018; Rodríguez-Ezpeleta et al., 2021).

Environmental DNA (eDNA) analysis has been rapidly developed over the past decade as an alternative and/or complementary biomonitoring tool and has become widely used for organisms from various taxonomic groups (Boivin-Delisle et al., 2021; Deiner et al., 2017; Doi et al., 2021; Kelly, 2016; Kumar et al., 2020). Environmental DNA analysis enables us to estimate the presence of organisms via only the collection and detection of cellular materials shed from

them in environments, including soil, water, and air (Ficetola et al., 2008; Kuwae et al., 2020;

Lynggaard et al., 2022). The success and high expectations of biomonitoring based on eDNA

analysis can also be seen in the rapid growth of the number of publications (Rodríguez-Ezpeleta

et al., 2021; Tsuji et al., 2019). Moreover, a meta-analysis has shown that 90% of the 63 studies

reported the positive relationships between eDNA concentration in a water sample and the

abundance and/or biomass of aquatic organisms (Rourke et al., 2022). These results suggest that

the eDNA analysis can be further developed and used in the future as a quantitative estimation

tool for biodiversity.

Environmental DNA analysis for detecting macroorganisms can be technically

categorized into two methods, i.e., species-specific detection and metabarcoding; however, some

challenges remain in the quantitative detection of eDNA in both methods (Bylemans et al., 2019).

The species-specific detection method using species-specific primers/probe and a real-time PCR

system is currently a major eDNA quantitative method. Nevertheless, the development of a

species-specific detection system or multiplexing assays of them is time-consuming and costly

and requires prior knowledge and assumptions about the species present within a study site

(Bylemans et al., 2019; Tsuji et al., 2018; Wozney and Wilson, 2017). Thus, it is unsuitable for

the quantitative detection targeting multiple species. Against this background, researchers have

recently begun to explore the possibilities of quantitative analysis on eDNA metabarcoding using

universal primers targeting taxonomic groups (e.g. Evans et al., 2016; Fraija-Fernández et al.,

2020; Kelly et al., 2014; Thomsen et al., 2016). The eDNA metabarcoding enables us to identify

community composition including multiple target taxa. However, the number of reads for each

taxon output from the high-throughput sequencer is difficult to treat as quantitative information

because it can easily vary between samples due to several problems such as PCR inhibitions,

primer bias, and library preparation bias (Deiner et al., 2017; Lamb et al., 2019; Lim et al., 2016;

Rourke et al., 2022).

Challenge to accurate quantification of eDNA through metabarcoding is an important

frontier of eDNA-based biomonitoring, and some approaches have recently been developed

(Hoshino et al., 2021; Smets et al., 2016; Ushio et al., 2018). Particularly, the quantitative MiSeq

sequencing approach (hereafter, qMiSeq approach) developed by Ushio et al. (2018) allows us to

convert the sequence read numbers of detected taxa to DNA copy numbers based on a linear

regression between known DNA copy numbers and observed sequence reads of internal standard

DNAs. One of the advantages of the qMiSeq approach is that the copy number can be calculated

considering the sample-specific effects of PCR inhibition and library preparation bias because a

sample-specific standard line is obtained by adding internal standard DNAs to each sample.

Accordingly, Ushio et al. (2018) reported the successful quantitative monitoring of eDNA derived

from multiple fish species in coastal marine ecosystems by combining the qMiSeq approach and

the fish universal primer, MiFish-U (Miya et al., 2015). A further verification experiment also

showed a positive correlation between the estimated eDNA concentration and the sound intensity

of fish (Sato et al., 2021). These results suggested that the qMiSeq approach is a promising

technique for quantitative eDNA-based monitoring of fish community. However, none of the

previous studies have been able to compare the observation data with the estimated eDNA

concentrations by species and study site. The accumulation of comparative studies on the use of

the qMiSeq approach will contribute to the further development and application of quantitative

eDNA metabarcoding and maximize its potential as a quantitative monitoring tool for

biodiversity (Ushio, 2022).

The objective of this study was to evaluate the performance of the qMiSeq approach as a

quantitative monitoring tool for fish community. Here, we compared the qMiSeq approach to the

capture-based survey with an electrical shocker at four river systems. Specifically, the followings

were tested: 1) for each study site, whether there is a significant positive relationship between the

fish eDNA concentrations obtained by the qMiSeq approach and the capture data (fish abundance

and biomass), and 2) for each taxon, whether the capture data at each site have significant

relationships with on the eDNA concentrations. Based on the results, we discussed the usefulness

of the qMiSeq approach for quantitative monitoring of fish community.

2 Materials and Methods

2.1 Study site and overview of survey design

Field surveys including capture survey and water sampling were conducted in four rivers in

western Japan (Fig. 1a, Fig. S1); Yokomichi River (YK; 11th September 2019), Hisakane River

(HS; 12th September 2019), Fukuchi River (FK; 8th November 2019) and Ino River (IN; 15th

November 2019). The three (HS) or six (YK, FK and IN) study sites were set for each river.

Details of each survey site were shown in Table S1. The overview of the survey design is shown

in Fig. 1b. Water sampling for eDNA analysis and capture survey with an electrical shocker were

performed for each study site.

2.2 Water sampling and filtration for eDNA analysis and Capture survey by electrical shocker

At each study site, we collected 1 L of surface water using a bleached bottle from the center line

of stream at the downstream end of each site (a total of 21 field samples, Fig. 1b). We added

benzalkonium chloride (1 mL, 10% w/v; Yamanaka et al., 2017) (Fujifilm Wako Pure Chemical

Corporation, Osaka, Japan) to each water sample to preserve eDNA and transported the sample

bottles to the laboratory under refrigeration. As a cooler blank for each sampling day, 1 L of

deionized water was placed into the same cooler box and thereafter treated in the same way as the

collected water samples (a total of four cooler blank samples). The water samples were vacuum

filtered using GF/F glass fibre filters (diameter: 47 mm, mesh size: 0.7 μm; GE Healthcare Japan,

Tokyo, Japan) within 36 hours after sampling. All filter samples were stored at –20˚C until DNA

extraction.

After the water sampling, we conducted fish samplings (up to 45 minutes per site) using

backpack electroshocker (Smith-Root, LR-20B, USA) and dip nets. Three survey lines were set

at each survey site and electric shocks were applied along each line (Fig. 1b). The collected fishes

were identified and released within the study reach except for the fishes we could not identify in

the field.

2.3 DNA extraction from the filter sample

The DNA on each filter sample was extracted following the method described by Environmental

DNA Sampling and Experiment Manual ver 2.1 (Minamoto et al., 2021) with minor

modifications. First, each filter sample was placed in the upper part of the Salivette tube

(SARSTEDT AG & Co. KG, Nümbrecht, Germany). The 220 μL of extraction solution

containing 200 μL Buffer AL and 20 μL of proteinase K in DNeasy Blood & Tissue Kit (Qiagen,

Hilden, Germany) were added onto each filter and incubated at 56˚C for 30 min. After incubation,

the Salivette tube was centrifuged at 5,000 × g for 1 min. 220 μL Tris–EDTA (TE) buffer (pH 8.0;

Nippon Gene Co., Ltd., Tokyo, Japan) was added onto each filter and re-centrifuged at 5,000 × g

for 3 min. After that, 200 μL of ethanol was added to the solution in the bottom part of the

Salivette tube and mixed well by gently pipetting. The whole of the mixed solution was

transferred to a DNeasy Mini spin column, and the DNA was purified according to the

manufacturer's protocol. The DNA was finally eluted in 100 μL Buffer AE and stored at −20˚C.

2.4 eDNA quantification: Quantitative metabarcoding with qMiSeq approach

The qMiSeq approach (Ushio et al. 2018) and MiFish-U primers (Miya et al. 2015) was

performed to identify fish taxa and quantify their eDNA concentrations simultaneously. The

qMiSeq allows us to convert the number of sequence reads into DNA copies without being

affected by differences in PCR efficiency by obtaining a sample-specific standard line using

internal standard DNAs. The paired-end library preparation with a two-step PCR was performed

in 12 μL of reaction mixture according to the described method in Tsuji et al., (2022) (See

appendix for details). Briefly, in first-round PCR was performed with MiFish-U-F/R primers

(Miya et al. 2015, Table S2), which can amplify fish DNA and three internal standard DNAs (5,

25, 50 copies per reaction, respectively: Table S3). Internal standard DNAs were added to only

field samples and cooler blanks. PCR negative control with ultrapure water instead of both eDNA

sample and standard DNA mix was added in all first PCR runs. The second-round PCR was

performed to add index-sequence and adapter-sequence for the Illumina sequencing platform

(Table S2). The indexed products of the second-round PCR were pooled, and the target bands (ca.

370 bp) were excised using 2% E☐Gel SizeSelect Agarose Gels (Thermo Fisher Scientific). The

prepared DNA libraries were sequenced by 2×150 bp paired-end sequencing on the iSeq

platform using the iSeq 100 i1 Reagent v1 cartridge (Illumina, CA, USA) with 30% PhiX

spike-in.

The bioinformatics analysis was performed using the PMiFish pipeline (Miya et al.,

2020; https://github.com/rogotoh/PMiFish). Briefly, first, low-quality tails were trimmed from

each sequence read, and the paired-end reads were merged. Then, primer sequences were

removed, and identical sequences were merged using UCLUST (USEARCH v10.0.240, Edgar,

2010). The merged sequences with 10 or more reads were assigned to the taxonomy using local

BLASTN search with the reference database including all inhabiting freshwater fish taxa around

the study sites (MiFish DB ver. 37) and the sequence of used three internal standard DNAs. The

top BLAST hits with a sequence identity ≥ 98.5% were applied and used in further analyses. For

Odontobutis obscura only, sequences detected with sequence identity ≥ 90.0% were also used in

the analysis, as they are known to exist in highly genetically differentiated regional populations.

As all study sites were in freshwater areas, any saltwater fishes detected were excluded from

subsequent analyses. For internal standard DNAs, to obtain a sample-specific standard line, linear

regression analysis (lm function in R version 3.6.0 software) was performed using the obtained

sequence reads and their known copy numbers (the intercept was set as zero). For each sample,

the number of eDNA copies per litre for each taxon was calculated as qMiSeq eDNA

concentration using each sample-specific standard line: qMiSeq eDNA concentration = the

number of iSeq sequence reads/regression slope of the sample-specific standard line (Table S4).

2.5 eDNA quantification: real-time quantitative PCR (qPCR) for two taxa

The real-time qPCR for *Candidia temminckii* and *Cottus pollux* ME were performed using a

StepOne-Plus Real-Time PCR system (Applied Biosystems, FosterCity, CA, USA). The eDNA

samples from the Yokomichi River were omitted from the comparison examination because they

were used in another study and there was no remaining volume for analysis. Species-specific

primers for each target species were developed in this study (Table S2), and their specificity was

tested by in silico and in vitro tests. All qPCR reactions were performed in a total 15 µL volume

and triplicated (See appendix for details of real-time qPCR conditions and species-specific primer

development).

2.6 Statistical analyses

For the following analyses, the R ver. 3.6.0 software was used (R Core Team. R, 2019). The

significance level was set at 0.05 in all analyses. To examine whether the DNA concentrations

quantified by the qMiSeq approach reflect those quantified by species-specific qPCR, the eDNA

concentrations of *C. temminckii* and *C. pollux* ME quantified by the two methods were compared

using the linear regression analysis (lm function in R). The differences in fish community

compositions among sites were visualised using nonmetric multidimensional scaling (NMDS)

with 500 permutations and cluster analysis with group average method. The Bray-Curtis

dissimilarity index was used as the fish community dissimilarity in both NMDS and k-means

clustering to use the DNA copy number as quantitative information. Based on k-means clustering,

a total of 21 study sites were divided into five clusters (see Results). The number of cluster, five,

was determined on the basis of Calinski criterion values (Fig. S2). The differences in eDNA

release rate and capture efficiency are expected between study sites with largely different fish

community compositions due to differences in geographical distribution, ecology, and behaviour

of each species, which could distort the relationship between observed DNA concentrations and

capture data. Thus, qMiSeq eDNA concentrations and capture data (abundance and biomass of

each taxon) were compared using the Kendall-rank correlation test by classifying the data in two

patterns as follows: (1) using data from all survey sites and (2) using each cluster site data

classified by similarity of fish community. Moreover, for eleven taxa which were detected from

three and over sites by each of qMiSeq approach and electrical shocker among all study sites, the

pooled dataset of qMiSeq eDNA concentrations and capture data (abundance or biomass at each

survey site) was created for each taxa. For each taxa, qMiSeq eDNA concentrations and capture

data were compared using generalized linear models (GLM; the glm.nb function in the MASS

package in R, Venables and Ripley, 2002) assuming that eDNA copy numbers follow a negative

binomial distribution with log-link function (Table 1). Here we have chosen GLMs as a

number of previous studies have suggested that the relationship between abundance

or biomass of each fish species and eDNA concentration is non-linear (e.g. Coulter et

al., 2019; Doi et al., 2017). Additionally, the eleven taxa were divided into four benthic fish

taxa group and seven pelagic fish taxa group, and the same analysis was carried out for each

group as described above. Note, however, that data on *O. obscura* and *O. hikimius*, which are

closely related and have almost identical ecology, were merged and treated as a single taxon.

3. Results

3.1 Overview of qMiSeq approach results and the comparison of eDNA concentrations quantified

by the qMiSeq approach with those estimated by qPCR

The iSeq paired-end sequencing (2×150 bp) for the 27 libraries (21 field samples, four cooler

blanks, two PCR negative controls) yielded a total of 4.71 million reads (Q30 = 95.2; PF =

67.3 %). Negligible sequence reads were detected from PCR negative controls (maximum 75

reads; Table S5), which were ignored in the subsequence analyses. The sequence reads of internal

standard DNAs in field samples and cooler blanks had a significant positive relationship with the

copy numbers (lm, $P < 0.01$, $R^2 > 0.93$; Table S4). The coefficients of the linear regressions

varied from 174.1 (IN st. 4) to 900.8 (IN cooler blank) (Table S4), suggesting that sequence reads

are proportional to the DNA copy numbers in a single sample. Thus, we converted sequence

reads of detected taxa in each sample to the DNA copies using the coefficient of the

sample-specific regression line. In the comparison of DNA concentrations of the three taxa

calculated by qMiSeq approach and species-specific qPCR, a significant positive relationship was

found for each taxa (linear regression analysis; $P < 0.001$, $R^2 = 0.81$, estimate 0.32 for *C.*

temminckii; $P < 0.001$, $R^2 = 0.99$, estimate 0.58 for *C. pollux* ME; Fig. 2).

3.2 Comparison of species richness observed by qMiSeq and electrical shocker

The qMiSeq approach consistently detected more species than the capture-based survey using an

electrical shocker (Fig. 3). On the other hand, one or two taxa that were detected by the electrical

shocker were not detected by the qMiSeq approach (i.e., false negative) in several study sites: HS

st.3, *Liobagrus reini*; FK st. 4, *Opsariichthys platypus* and *Oryzias latipes*; FK st. 5, *Cobitis*

matsubarae; FK st.6, *Squalidus gracilis gracilis* and *Tachysurus nudiceps*; IN st. 5; *S. gracilis*

gracilis. The false negative result in qMiSeq results for *Cobitis matsubarae* at FK st. 5 was

excluded from all subsequent analyses as it was likely due to a lack of reference DNA database

(see Discussion for detail). On the other hand, no false negatives were observed at 16 out of a

total of 21 sites. At almost all sites, *Rhinogobius* spp. was detected by both or either of methods,

but it was excluded from subsequent analyses because they contain many unknown haplotypes

and many closely related species that cannot be discriminated and/or detected based on the target

region of MiFish primer. Also, at IN st. 5, *Cyprinus carpio* were detected by both methods, but

were excluded from subsequent analyses because five individuals (approximately 50 cm, total

length) were also visually identified immediately upstream of the study section. Their eDNA

almost certainly flowed into the study section and was considered to be noise in this study. The

fish taxa detected only by qMiSeq approach were the taxon collected from other sites in the same

river, a rare native species likely to be inhabited or an invasive species in the early stages of

invasion (Table S5).

3.3 Fish community structure

The dissimilarity of the fish community structure among study sites was shown in the nonmetric

multidimensional scaling (NMDS) biplot (Fig. 4a). A total of 20 research sites (FK st. 1 was

excluded due to absebce of detected taxa) were categorised into five clusters by k-means

clustering. The clusters tended to be divided by the down and up stream. Cluster 3 and 4

consisted only of the study sites at Yokomichi River.

3.4 Comparison of the qMiSeq eDNA concentrations with the capture data by electrical shocker

When all sites were used for analysis, both the number and biomass of captured fishes had

significant positive correlations with the qMiSeq eDNA concentrations (Kendall rank correlation

test, Ps < 0.0001, $\tau = 0.467$ for abundance and $\tau = 0.469$ for biomass; Fig. 4b, Table S6).

Additionally, the significant positive correlations were also found for all site clusters (Kendall

rank correlation test, both the number and biomass, $P < 0.01$ for all site clusters; Fig. 4b, Table

S6). The Kendall's τ value varied widely among clusters (0.315 to 0.598 for abundance; 0.359 to

0.601 for biomass). Moreover, for seven out of eleven taxa which were detected from three and

more sites by each of qMiSeq approach and electrical shocker, we found significant positive

relationships between the qMiSeq eDNA copy numbers at each site and the number of captured

individuals and/or the total biomass (GLM; Fig. 5, Table 1). When eleven taxa were grouped into

four benthic fish taxa and seven pelagic fish taxa, there were significant positive relationships

between the qMiSeq eDNA concentration and both the number of captured individuals and the

total biomass (GLM; Table 1).

4. Discussion

4.1 eDNA metabarcoding detected higher species richness than electrofishing survey

We found that eDNA metabarcoding with the qMiSeq approach could detect not only almost all

the taxa captured using the electric shocker but also other taxa that could be false negatives in the

capture surveys. This result is consistent with several previous studies that have reported

advantages of eDNA metabarcoding compared with traditional capture-based methods in terms of

detection sensitivity on biodiversity assessment (Andruszkiewicz et al., 2017; Bylemans et al.,

2018; Civade et al., 2016; Deiner et al., 2016; Hänfling et al., 2016; Miya et al., 2015; Nakagawa

et al., 2018; Sato et al., 2017; Shaw et al., 2016; Thomsen et al., 2012; Valentini et al., 2016;

Yamamoto et al., 2017). On the other hand, we should always be aware the possibility of false

negative. In this study, although almost all taxa were detected, a total of six taxa were missed by

eDNA metabarcoding at the sampling site level. Three main causes of false negative results have

been suggested in eDNA metabarcoding: (1) low concentration of target DNA, (2) PCR drop-out

because of primer mismatch and (3) incompleteness and inaccuracy of the reference sequence

database (Evans et al., 2017; Jane et al., 2015; Miya et al., 2020).

4.2 The significant positive correlation between eDNA concentrations quantified by qMiSeq and

qPCR

The significant positive relationship between eDNA concentrations quantified by the qMiSeq

approach and species-specific qPCR indicated that the qMiSeq approach successfully estimated

the number of sequence reads of each taxon to the eDNA copy number. For both taxa considered,

the eDNA copy numbers calculated by the qMiSeq approach tended to be smaller than those

estimated by qPCR. This observation was consistent with those reported in the previous study

(Ushio et al., 2018). Additionally, eDNA metabarcoding has been recognised to be less sensitive

than species-specific detection due to the effects of technical issues such as primer bias, PCR bias

and the potential reduced amplification efficiency in tailed-PCR (Bylemans et al., 2019; Harper et

al., 2018; Nichols et al., 2018). However, this characteristic would not be a serious problem when

we compare the eDNA concentrations quantified by the qMiSeq approach on a

species-by-species basis.

4.3 Calculated eDNA concentration by qMiSeq reflected both fish abundance and biomass in

each study site

It was worthy of note that both abundance and biomass of inhabiting fishes had positive

correlations with the qMiSeq eDNA concentrations. This study is the first to demonstrate the

relationships between the quantitative capture data of fishes and their eDNA concentrations

quantified using eDNA metabarcoding. Meanwhile, Kendall's rank correlation coefficients were

low in both abundance and biomass ($\tau < 0.5$), possibly because differences in eDNA release

and/or degradation rates among taxa inhabiting the study sites, capture efficiency with electric

shockers, and environmental conditions could affect the results.

The rank correlation coefficients (Kendall's τ) tended to be improved by dividing the

samples into site clusters based on the dissimilarity of fish community compositions, compared to

when all study sites were considered. The results suggested that the study site clustering based on

fish community compositions has mitigated the effects of the differences of the eDNA ecology

and/or dynamics such as release rate and/or degradation rates for each taxon and dispersion

processes on the relationship between qMiSeq eDNA concentration and fish capture data. In

contrast, at some clusters, Kendall's τ values were decreased. Particularly, in cluster 5 sites,

Kendall's τ values decreased for both abundance and biomass and were smallest in all site

clusters. The cluster 5 consisted only of downstream sites (IN st. 5, 6 and FK st. 5, 6), and

vegetation covered large areas of the channel, mainly along the river bank (Figure S1). Relatedly,

the taxa with highly biased relationships between qMiSeq eDNA concentrations and capture data

had ecological characteristics such as a preference for vegetation (*Pungtungia herzi*) and rock

crevices (*O. obscura*), hiding in the sandy bottom (*Pseudogobio esocinus esocinus*) and high

swimming ability (*C. temminckii*) (Hosoya, 2019; Nakabo, 2013). The environmental condition

and these ecological characteristics would have strongly influenced capture efficiency in the

electrical shocker survey. Taxa hiding in vegetation, rock crevices and sandy bottom were

difficult to capture because electric shocks cannot reach them, or they faint in unseen crevices.

Fish taxa with high swimming ability were able to quickly escape from the investigator's position

or hide under vegetation. In this study, it was difficult to conduct an exhaustive fish capture by

closing the study site section due to the limitation of labour, manpower and time for the survey.

These facts may indicate that differences in the capture efficiency of the electrical shocker survey

have influenced the results of this study, although the magnitude of the impact could depend on

the environmental conditions and the ecological characteristics of taxa that inhabit at each study

site.

4.4 qMiSeq eDNA concentration reflected both fish abundance and biomass in each detected

taxon

For seven taxa out of eleven taxa with higher detection frequency in both methods, we

demonstrated that the qMiSeq eDNA concentration was significantly related to the abundance

and/or biomass. Additionally, qMiSeq eDNA concentration and both the number of captured

individuals and total biomass were significantly related when the eleven taxa were grouped into

benthic or pelagic fish taxa. It was consistent with the results of many previous studies showing

positive relationships between eDNA concentration and the abundance and/or biomass of target

species (90% of relevant 63 papers published by 2020, Rourke et al., 2022). This observation

supports the potential and usefulness of the qMiSeq approach as a quantitative monitoring tool of

biodiversity beyond simply identifying species presence. Our result in this study was significant

for two reasons.

First, to the best of our knowledge, there have been no studies comparing eDNA

concentrations of various macroorganism taxa obtained by quantitative metabarcoding methods,

such as the qMiSeq approach, with quantitative capture data for each taxon. To the future

implementation of quantitative eDNA metabarcoding methods for quantitative biodiversity

monitoring, we should continue the detailed and repeated research efforts to determine whether

the results obtained from conventional survey methods and eDNA-based methods yield similar

measurements for the taxa of interest (Deiner et al., 2017). We believe that this study was the first

step in this process. Second, it is worth mentioning that the comparisons of eDNA concentrations

and capture data in this study have been performed using pooled eDNA concentration data for

each species derived from multiple different samples. In general, the direct comparison of

sequence reads count among different samples to quantitatively estimate biodiversity lead to

erroneous conclusions because the number of sequence reads will vary among samples depending

on the effects of PCR inhibition, primer bias, library preparation bias and sequencing depth, etc

(Hoshino et al., 2021; Ushio et al., 2018). In most cases, researchers used the proportional

abundance of sequence read counts in a sample for each detected taxon instead of a direct

comparison of their sequence read counts (e.g. Fraija-Fernández et al., 2020; Goutte et al., 2020;

Thomsen et al., 2016). However, as differences in the number of taxa detected per sample

inherently bias the proportion of sequence read counts (i.e. an increase in the total number of

species would reduce the proportional abundance), it is difficult to apply to quantitative

comparisons of target taxa among sites, seasons and years with different community

compositions (Bylemans et al., 2019). In contrast, although the qMiSeq approach also cannot

mitigate the effects of primer bias (see below), it allows avoiding the effects of PCR inhibition,

library preparation bias and sequencing depth (Ushio et al., 2018). We reasonably compared the

eDNA concentrations of each taxon from multiple samples and demonstrated the usefulness of

the analytical advantages of the qMiSeq method for quantitative eDNA metabarcoding for

macroorganisms.

Using the qMiSeq approach, although a significant relationship was observed between

qMiSeq eDNA concentration and capture data for 7 taxa and benthic or pelagic fish taxa group,

the obtained values varied widely between and within the taxon. It is obvious that the relationship

between qMiSeq eDNA concentration and abundance and/or biomass of target taxon observed by

field experiments can be obscured by biological and non-biological factors, which is the

limitation of analytical method and survey design (Deiner et al., 2017; Rourke et al., 2022). In

this study, the environmental conditions such as water temperatures and flow velocity at the time

of water sampling varied widely among study sites (Table S1), and there may have been

associated differences in the characteristics of the 'ecology of eDNA' within each taxon (e.g., its

origin, state, fate and transport) (Barnes and Turner, 2016; Collins et al., 2019; Jane et al., 2015;

Tsuji et al., 2017). Additionally, as mentioned above, the limitations of capture survey using

electrical shocker was likely to have caused significant fluctuations in the observed values.

Furthermore, we also need to mention the effect of primer bias, which is an analytical limitation

of eDNA metabarcoding. Primer bias is recognised as one of the main causes of the distortion of

the relative abundance of amplified DNA in a sample in DNA metabarcoding data (Elbrecht and

Leese, 2015; Lamb et al., 2019; Nester et al., 2020; Piñol et al., 2015). The MiFish-U primers

used in this study has been proven to have high primer universality by many previous studies (e.g.

Collins et al., 2019; Miya et al., 2020; Zhang et al., 2020). Thus, we believe that the distortion of

results due to primer bias was minimised in this study.

Conclusions

We here compared the eDNA concentrations quantified by the qMiSeq approach and the results

of the capture-based survey using an electrical shocker to evaluate the performance of the

qMiSeq approach as a quantitative monitoring tool for fish community. We found positive

relationships between the qMiSeq eDNA concentrations and both the abundance and biomass of

each captured taxon. Together, our results suggested that eDNA metabarcoding with the qMiSeq

approach is a suitable and useful tool for quantitative monitoring of fish community. The

simplicity of eDNA analysis will reduce barriers forassessing changes in species abundance,

evenness and richness and will facilitate the collection of valuable information for better

understanding biodiversity changes.

Authors' contributions

S.T., R.I. and Y.A. conceived and designed research. R.I, S.M. M.S. and T.K. performed a field

survey. S.T. and R.N. performed molecular experiments and data analysis. S.T. wrote the early

draft and completed it with significant inputs from all authors.

Data availability

Full details of the eDNA metabarcoding with qMiSeq approach and species-specific qPCR

results are available in the supporting information (Table S5 and S7). All raw sequences were

deposited in the DDBJ Sequence Read Archive (accession number: DRA013858).

Acknowledgements

We thank laboratory members of Akamatsu laboratory, Yamaguchi University, and Inui

Laboratory, Fukuoka Institute of Technology University, for help with the field sampling. We

thank Mr. S. Kunimatsu and Mr. Y. Fuke for providing fish pictures. We thank Dr. M. Ushio for

his valuable comments and advices on the draft. This is supported by the River Fund of The

River Foundation, Japan (2019-5211-030). This work was supported by YU Project for

Formation of the Core Research Center.

References

Andruszkiewicz, E.A., Sassoubre, L.M., Boehm, A.B., 2017. Persistence of marine fish
environmental DNA and the influence of sunlight. PLOS ONE 12, e0185043.
https://doi.org/10.1371/journal.pone.0185043
Barnes, M.A., Turner, C.R., 2016. The ecology of environmental DNA and implications for
conservation genetics. Conservation genetics 17, 1–17.
Blowes, S.A., Daskalova, G.N., Dornelas, M., Engel, T., Gotelli, N.J., Magurran, A.E., Martins,
I.S., McGill, B., McGlinn, D.J., Sagouis, A., Shimadzu, H., Supp, S.R., Chase, J.M., 2022.
Local biodiversity change reflects interactions among changing abundance, evenness, and
richness. Ecology online, e3820. https://doi.org/10.1002/ecy.3820
Boivin-Delisle, D., Laporte, M., Burton, F., Dion, R., Normandeau, E., Bernatchez, L., 2021.
Using environmental DNA for biomonitoring of freshwater fish communities: Comparison
with established gillnet surveys in a boreal hydroelectric impoundment. Environmental
DNA 3, 105–120.
Bylemans, J., Gleeson, D.M., Duncan, R.P., Hardy, C.M., Furlan, E.M., 2019. A performance
evaluation of targeted eDNA and eDNA metabarcoding analyses for freshwater fishes.
Environmental DNA 1, 402–414. https://doi.org/10.1002/edn3.41
Bylemans, J., Gleeson, D.M., Hardy, C.M., Furlan, E., 2018. Toward an ecoregion scale evaluation
of eDNA metabarcoding primers: A case study for the freshwater fish biodiversity of the
Murray–Darling Basin (Australia). Ecology and Evolution 8, 8697–8712.
https://doi.org/10.1002/ece3.4387
Cardinale, B.J., Duffy, J.E., Gonzalez, A., Hooper, D.U., Perrings, C., Venail, P., Narwani, A.,
Mace, G.M., Tilman, D., Wardle, D.A., Kinzig, A.P., Daily, G.C., Loreau, M., Grace, J.B.,
Larigauderie, A., Srivastava, D.S., Naeem, S., 2012. Biodiversity loss and its impact on
humanity. Nature 486, 59–67. https://doi.org/10.1038/nature11148
Civade, R., Dejean, T., Valentini, A., Roset, N., Raymond, J.-C., Bonin, A., Taberlet, P., Pont, D.,
2016. Spatial representativeness of environmental DNA metabarcoding signal for fish
biodiversity assessment in a natural freshwater system. PLOS ONE 11, e0157366.
https://doi.org/10.1371/journal.pone.0157366

Collins, R.A., Bakker, J., Wangensteen, O.S., Soto, A.Z., Corrigan, L., Sims, D.W., Genner, M.J.,
Mariani, S., 2019. Non-specific amplification compromises environmental DNA
metabarcoding with COI. Methods in Ecology and Evolution 10, 1985–2001.
https://doi.org/10.1111/2041-210X.13276
Coulter, D.P., Wang, P., Coulter, A.A., Susteren, G.E.V., Eichmiller, J.J., Garvey, J.E., Sorensen,
P.W., 2019. Nonlinear relationship between Silver Carp density and their eDNA
concentration in a large river. PLOS ONE 14, e0218823.
https://doi.org/10.1371/journal.pone.0218823
Crowder, D.W., Northfield, T.D., Gomulkiewicz, R., Snyder, W.E., 2012. Conserving and
promoting evenness: organic farming and fire-based wildland management as case
studies. Ecology 93, 2001–2007. https://doi.org/10.1890/12-0110.1
Deiner, K., Bik, H.M., Mächler, E., Seymour, M., Lacoursière-Roussel, A., Altermatt, F., Creer, S.,
Bista, I., Lodge, D.M., de Vere, N., Pfrender, M.E., Bernatchez, L., 2017. Environmental
DNA metabarcoding: Transforming how we survey animal and plant communities.
Molecular Ecology 26, 5872–5895. https://doi.org/10.1111/mec.14350
Deiner, K., Fronhofer, E.A., Mächler, E., Walser, J.-C., Altermatt, F., 2016. Environmental DNA
reveals that rivers are conveyer belts of biodiversity information. Nat Commun 7, 12544.
https://doi.org/10.1038/ncomms12544
Doi, H., Inui, R., Akamatsu, Y., Kanno, K., Yamanaka, H., Takahara, T., Minamoto, T., 2017.
Environmental DNA analysis for estimating the abundance and biomass of stream fish.
Freshwater Biology 62, 30–39. https://doi.org/10.1111/fwb.12846
Doi, H., Minamoto, T., Takahara, T., Tsuji, S., Uchii, K., Yamamoto, S., Katano, I., Yamanaka, H.,
2021. Compilation of real-time PCR conditions toward the standardization of
environmental DNA methods. Ecological Research 36, 379–388.
Dornelas, M., Gotelli, N.J., McGill, B., Shimadzu, H., Moyes, F., Sievers, C., Magurran, A.E., 2014.
Assemblage time series reveal biodiversity change but not systematic loss. Science 344,
296–299. https://doi.org/10.1126/science.1248484
Elbrecht, V., Leese, F., 2015. Can DNA-based ecosystem assessments quantify species
abundance? Testing primer bias and biomass—sequence relationships with an innovative
metabarcoding protocol. PLOS ONE 10, e0130324.
https://doi.org/10.1371/journal.pone.0130324
Evans, N.T., Olds, B.P., Renshaw, M.A., Turner, C.R., Li, Y., Jerde, C.L., Mahon, A.R., Pfrender,
M.E., Lamberti, G.A., Lodge, D.M., 2016. Quantification of mesocosm fish and amphibian
species diversity via environmental DNA metabarcoding. Molecular Ecology Resources 16,
29–41. https://doi.org/10.1111/1755-0998.12433
Evans, N.T., Shirey, P.D., Wieringa, J.G., Mahon, A.R., Lamberti, G.A., 2017. Comparative cost
and effort of fish distribution detection via environmental DNA analysis and
electrofishing. Fisheries 42, 90–99.
Ficetola, G.F., Miaud, C., Pompanon, F., Taberlet, P., 2008. Species detection using environmental
DNA from water samples. Biology letters 4, 423–425.
Fraija-Fernández, N., Bouquieaux, M.-C., Rey, A., Mendibil, I., Cotano, U., Irigoien, X., Santos,
M., Rodríguez-Ezpeleta, N., 2020. Marine water environmental DNA metabarcoding
provides a comprehensive fish diversity assessment and reveals spatial patterns in a

large oceanic area. Ecology and Evolution 10, 7560–7584.
https://doi.org/10.1002/ece3.6482

Goutte, A., Molbert, N., Guérin, S., Richoux, R., Rocher, V., 2020. Monitoring freshwater fish
communities in large rivers using environmental DNA metabarcoding and a long-term
electrofishing survey. Journal of Fish Biology 97, 444–452.
https://doi.org/10.1111/jfb.14383

Hänfling, B., Lawson Handley, L., Read, D.S., Hahn, C., Li, J., Nichols, P., Blackman, R.C., Oliver,
A., Winfield, I.J., 2016. Environmental DNA metabarcoding of lake fish communities
reflects long-term data from established survey methods. Molecular Ecology 25,
3101–3119. https://doi.org/10.1111/mec.13660

Harper, L.R., Lawson Handley, L., Hahn, C., Boonham, N., Rees, H.C., Gough, K.C., Lewis, E.,
Adams, I.P., Brotherton, P., Phillips, S., Hänfling, B., 2018. Needle in a haystack? A
comparison of eDNA metabarcoding and targeted qPCR for detection of the great crested
newt (*Triturus cristatus*). Ecology and Evolution 8, 6330–6341.
https://doi.org/10.1002/ece3.4013

Hillebrand, H., Bennett, D.M., Cadotte, M.W., 2008. Consequences of dominance: a review of
evenness effects on local and regional ecosystem processes. Ecology 89, 1510–1520.
https://doi.org/10.1890/07-1053.1

Hoshino, T., Nakao, R., Doi, H., Minamoto, T., 2021. Simultaneous absolute quantification and
sequencing of fish environmental DNA in a mesocosm by quantitative sequencing
technique. Scientific Report 11, 4372. https://doi.org/10.1038/s41598-021-83318-6

Hosoya, K., 2019. Sankei Handy Illustrated Book 15: Freshwater fish of Japan, enlarged and
revised edition. ed. Yama-kei Publishers co.,Ltd.

Jane, S.F., Wilcox, T.M., McKelvey, K.S., Young, M.K., Schwartz, M.K., Lowe, W.H., Letcher, B.H.,
Whiteley, A.R., 2015. Distance, flow and PCR inhibition: eDNA dynamics in two
headwater streams. Molecular Ecology Resources 15, 216–227.
https://doi.org/10.1111/1755-0998.12285

Kanno, K., Onikura, N., Kurita, Y., Koyama, A., Nakajima, J., 2018. Morphological, distributional,
and genetic characteristics of *Cottus pollux* in the Kyushu Island, Japan: indication of
fluvial and amphidromous life histories within a single lineage. Ichthyol Res 65, 462–470.
https://doi.org/10.1007/s10228-018-0637-4

Kelly, R.P., 2016. Making environmental DNA count. Molecular Ecology Resources 16, 10–12.
https://doi.org/10.1111/1755-0998.12455

Kelly, R.P., Port, J.A., Yamahara, K.M., Crowder, L.B., 2014. Using environmental DNA to census
marine fishes in a large mesocosm. PLoS ONE 9, e86175.
https://doi.org/10.1371/journal.pone.0086175

Kissling, W.D., Ahumada, J.A., Bowser, A., Fernandez, M., Fernández, N., García, E.A.,
Guralnick, R.P., Isaac, N.J.B., Kelling, S., Los, W., McRae, L., Mihoub, J.-B., Obst, M.,
Santamaria, M., Skidmore, A.K., Williams, K.J., Agosti, D., Amariles, D., Arvanitidis, C.,
Bastin, L., De Leo, F., Egloff, W., Elith, J., Hobern, D., Martin, D., Pereira, H.M., Pesole,
G., Peterseil, J., Saarenmaa, H., Schigel, D., Schmeller, D.S., Segata, N., Turak, E., Uhlir,
P.F., Wee, B., Hardisty, A.R., 2018. Building essential biodiversity variables (EBVs) of

species distribution and abundance at a global scale. Biological Reviews 93, 600–625.
https://doi.org/10.1111/brv.12359

Kumar, G., Eble, J.E., Gaither, M.R., 2020. A practical guide to sample preservation and pre-PCR
processing of aquatic environmental DNA. Molecular Ecology Resources 20, 29–39.
https://doi.org/10.1111/1755-0998.13107

Kuwae, M., Tamai, H., Doi, H., Sakata, M.K., Minamoto, T., Suzuki, Y., 2020. Sedimentary DNA
tracks decadal-centennial changes in fish abundance. Commun Biol 3, 1–12.
https://doi.org/10.1038/s42003-020-01282-9

Lamb, P.D., Hunter, E., Pinnegar, J.K., Creer, S., Davies, R.G., Taylor, M.I., 2019. How
quantitative is metabarcoding: A meta-analytical approach. Molecular Ecology 28,
420–430. https://doi.org/10.1111/mec.14920

Lim, N.K.M., Tay, Y.C., Srivathsan, A., Tan, J.W.T., Kwik, J.T.B., Baloğlu, B., Meier, R., Yeo,
D.C.J., 2016. Next-generation freshwater bioassessment: eDNA metabarcoding with a
conserved metazoan primer reveals species-rich and reservoir-specific communities. R.
Soc. open sci. 3, 160635. https://doi.org/10.1098/rsos.160635

Lynggaard, C., Bertelsen, M.F., Jensen, C.V., Johnson, M.S., Frøslev, T.G., Olsen, M.T., Bohmann,
K., 2022. Airborne environmental DNA for terrestrial vertebrate community monitoring.
Current Biology 32, 701-707.e5. https://doi.org/10.1016/j.cub.2021.12.014

Magurran, A.E., Deacon, A.E., Moyes, F., Shimadzu, H., Dornelas, M., Phillip, D.A.T., Ramnarine,
I.W., 2018. Divergent biodiversity change within ecosystems. Proceedings of the National
Academy of Sciences 115, 1843–1847. https://doi.org/10.1073/pnas.1712594115

Masuda, R., Shiba, M., Yamashita, Y., Ueno, M., Kai, Y., Nakanishi, A., Torikoshi, M., Tanaka, M.,
2010. Fish assemblages associated with three types of artificial reefs: density of
assemblages and possible impacts on adjacent fish abundance.

Minamoto, T., Miya, M., Sado, T., Seino, S., Doi, H., Kondoh, M., Nakamura, K., Takahara, T.,
Yamamoto, S., Yamanaka, H., Araki, H., Iwasaki, W., Kasai, A., Masuda, R., Uchii, K.,
2021. An illustrated manual for environmental DNA research: Water sampling guidelines
and experimental protocols. Environmental DNA 3, 8–13.
https://doi.org/10.1002/edn3.121

Miya, M., Gotoh, R.O., Sado, T., 2020. MiFish metabarcoding: a high-throughput approach for
simultaneous detection of multiple fish species from environmental DNA and other
samples. Fisheries Science 86, 939–970.

Miya, M., Sato, Y., Fukunaga, T., Sado, T., Poulsen, J.Y., Sato, K., Minamoto, T., Yamamoto, S.,
Yamanaka, H., Araki, H., 2015. MiFish, a set of universal PCR primers for metabarcoding
environmental DNA from fishes: detection of more than 230 subtropical marine species.
Royal Society open science 2, 150088.

Miyazono, S., Patiño, R., Taylor, C.M., 2015. Desertification, salinization, and biotic
homogenization in a dryland river ecosystem. Science of The Total Environment 511,
444–453. https://doi.org/10.1016/j.scitotenv.2014.12.079

Nakabo, T., 2013. Fishes of Japan with pictorial keys to the species (3-Volume Set). Tokai
University Press.

Nakagawa, H., Yamamoto, S., Sato, Y., Sado, T., Minamoto, T., Miya, M., 2018. Comparing
local-and regional-scale estimations of the diversity of stream fish using eDNA
metabarcoding and conventional observation methods. Freshwater Biology 63, 569–580.
Nester, G.M., De Brauwer, M., Koziol, A., West, K.M., DiBattista, J.D., White, N.E., Power, M.,
Heydenrych, M.J., Harvey, E., Bunce, M., 2020. Development and evaluation of fish eDNA
metabarcoding assays facilitate the detection of cryptic seahorse taxa (family:
Syngnathidae). Environmental DNA 2, 614–626. https://doi.org/10.1002/edn3.93
Nichols, R.V., Vollmers, C., Newsom, L.A., Wang, Y., Heintzman, P.D., Leighton, M., Green, R.E.,
Shapiro, B., 2018. Minimizing polymerase biases in metabarcoding. Molecular Ecology
Resources 18, 927–939. https://doi.org/10.1111/1755-0998.12895
Oka, S., Doi, H., Miyamoto, K., Hanahara, N., Sado, T., Miya, M., 2021. Environmental DNA
metabarcoding for biodiversity monitoring of a highly diverse tropical fish community in a
coral reef lagoon: Estimation of species richness and detection of habitat segregation.
Environmental DNA 3, 55–69. https://doi.org/10.1002/edn3.132
Pimm, S.L., Alibhai, S., Bergl, R., Dehgan, A., Giri, C., Jewell, Z., Joppa, L., Kays, R., Loarie, S.,
2015. Emerging technologies to conserve biodiversity. Trends in Ecology & Evolution 30,
685–696. https://doi.org/10.1016/j.tree.2015.08.008
Piñol, J., Mir, G., Gomez-Polo, P., Agustí, N., 2015. Universal and blocking primer mismatches
limit the use of high-throughput DNA sequencing for the quantitative metabarcoding of
arthropods. Molecular Ecology Resources 15, 819–830.
https://doi.org/10.1111/1755-0998.12355
R Core Team. R, 2019. A language and environment for statistical computing.
http://www.R-project.org.
Rodríguez-Ezpeleta, N., Zinger, L., Kinziger, A., Bik, H.M., Bonin, A., Coissac, E., Emerson, B.C.,
Lopes, C.M., Pelletier, T.A., Taberlet, P., Narum, S., 2021. Biodiversity monitoring using
environmental DNA. Molecular Ecology Resources 21, 1405–1409.
https://doi.org/10.1111/1755-0998.13399
Rourke, M.L., Fowler, A.M., Hughes, J.M., Broadhurst, M.K., DiBattista, J.D., Fielder, S., Wilkes
Walburn, J., Furlan, E.M., 2022. Environmental DNA (eDNA) as a tool for assessing fish
biomass: A review of approaches and future considerations for resource surveys.
Environmental DNA 4, 9–33. https://doi.org/10.1002/edn3.185
Sato, H., Sogo, Y., Doi, H., Yamanaka, H., 2017. Usefulness and limitations of sample pooling for
environmental DNA metabarcoding of freshwater fish communities. Sci Rep 7, 14860.
https://doi.org/10.1038/s41598-017-14978-6
Sato, M., Inoue, N., Nambu, R., Furuichi, N., Imaizumi, T., Ushio, M., 2021. Quantitative
assessment of multiple fish species around artificial reefs combining environmental DNA
metabarcoding and acoustic survey. Scientific reports 11, 1–14.
Shaw, J.L.A., Clarke, L.J., Wedderburn, S.D., Barnes, T.C., Weyrich, L.S., Cooper, A., 2016.
Comparison of environmental DNA metabarcoding and conventional fish survey methods
in a river system. Biological Conservation 197, 131–138.
https://doi.org/10.1016/j.biocon.2016.03.010
Smets, W., Leff, J.W., Bradford, M.A., McCulley, R.L., Lebeer, S., Fierer, N., 2016. A method for
simultaneous measurement of soil bacterial abundances and community composition via

16S rRNA gene sequencing. Soil Biology and Biochemistry 96, 145–151.
https://doi.org/10.1016/j.soilbio.2016.02.003

Thomsen, P.F., Kielgast, J., Iversen, L.L., Wiuf, C., Rasmussen, M., Gilbert, M.T.P., Orlando, L.,
Willerslev, E., 2012. Monitoring endangered freshwater biodiversity using environmental
DNA. Molecular Ecology 21, 2565–2573. https://doi.org/10.1111/j.1365-294X.2011.05418.x

Thomsen P.F., Møller P.R., Sigsgaard E.E., Knudsen S.W., Jørgensen O.A., Willerslev E., 2016.
Environmental DNA from seawater samples correlate with trawl catches of subarctic,
deepwater fishes. PLOS ONE 11, e0165252. https://doi.org/10.1371/journal.pone.0165252

Tsuji, S., Iguchi, Y., Shibata, N., Teramura, I., Kitagawa, T., Yamanaka, H., 2018. Real-time
multiplex PCR for simultaneous detection of multiple species from environmental DNA:
an application on two Japanese medaka species. Scientific reports 8, 1–8.

Tsuji, S., Nakao, R., Saito, M., Minamoto, T., Akamatsu, Y., 2022. Pre-centrifugation before DNA
extraction mitigates extraction efficiency reduction of environmental DNA caused by the
preservative solution (benzalkonium chloride) remaining in the filters. Limnology 23,
9–16. https://doi.org/10.1007/s10201-021-00676-w

Tsuji, S., Takahara, T., Doi, H., Shibata, N., Yamanaka, H., 2019. The detection of aquatic
macroorganisms using environmental DNA analysis—A review of methods for collection,
extraction, and detection. Environmental DNA 1, 99–108.

Tsuji, S., Ushio, M., Sakurai, S., Minamoto, T., Yamanaka, H., 2017. Water
temperature-dependent degradation of environmental DNA and its relation to bacterial
abundance. PLOS ONE 12, e0176608. https://doi.org/10.1371/journal.pone.0176608

Ushio, M., 2022. Interaction capacity as a potential driver of community diversity. Proceedings of
the Royal Society B: Biological Sciences 289, 20212690.
https://doi.org/10.1098/rspb.2021.2690

Ushio, M., Murakami, H., Masuda, R., Sado, T., Miya, M., Sakurai, S., Yamanaka, H., Minamoto,
T., Kondoh, M., 2018. Quantitative monitoring of multispecies fish environmental DNA
using high-throughput sequencing. Metabarcoding and Metagenomics 2, e23297.
https://doi.org/10.3897/mbmg.2.23297

Valentini, A., Taberlet, P., Miaud, C., Civade, R., Herder, J., Thomsen, P.F., Bellemain, E.,
Besnard, A., Coissac, E., Boyer, F., Gaboriaud, C., Jean, P., Poulet, N., Roset, N., Copp,
G.H., Geniez, P., Pont, D., Argillier, C., Baudoin, J.-M., Peroux, T., Crivelli, A.J., Olivier, A.,
Acqueberge, M., Le Brun, M., Møller, P.R., Willerslev, E., Dejean, T., 2016.
Next-generation monitoring of aquatic biodiversity using environmental DNA
metabarcoding. Molecular Ecology 25, 929–942. https://doi.org/10.1111/mec.13428

Venables, W.N., Ripley, B.D., 2002. Modern Applied Statistics with S, Fourth. ed. Springer, New
York.

Wozney, K.M., Wilson, C.C., 2017. Quantitative PCR multiplexes for simultaneous multispecies
detection of Asian carp eDNA. Journal of Great Lakes Research 43, 771–776.
https://doi.org/10.1016/j.jglr.2017.05.001

Yamamoto, S., Masuda, R., Sato, Y., Sado, T., Araki, H., Kondoh, M., Minamoto, T., Miya, M.,
2017. Environmental DNA metabarcoding reveals local fish communities in a species-rich
coastal sea. Sci Rep 7, 40368. https://doi.org/10.1038/srep40368

Yamanaka, H., Minamoto, T., Matsuura, J., Sakurai, S., Tsuji, S., Motozawa, H., Hongo, M., Sogo,
Y., Kakimi, N., Teramura, I., Sugita, M., Baba, M., Kondo, A., 2017. A simple method for
preserving environmental DNA in water samples at ambient temperature by addition of
cationic surfactant. Limnology 18, 233–241. https://doi.org/10.1007/s10201-016-0508-5
Yonekura, R., Kita, M., Yuma, M., 2004. Species diversity in native fish community in Japan:
comparison between non-invaded and invaded ponds by exotic fish. Ichthyological
Research 51, 176–179. https://doi.org/10.1007/s10228-003-0200-8
Zhang, S., Zhao, J., Yao, M., 2020. A comprehensive and comparative evaluation of primers for
metabarcoding eDNA from fish. Methods in Ecology and Evolution 11, 1609–1625.
https://doi.org/10.1111/2041-210X.13485
714
715

Table

Table 1 Results of GLM for relationships between qMiSeq eDNA concentration (copies/L) and
abundance or biomass for each taxon. The scientific name basically followed Nakabo
(2013), but *Cottus* sp. were denoted as *Cottus pollux* ME because it was known that
Cottus sp. in the study sites have mitochondria sequences consistent with *Cottus pollux*
ME regardless of morphology (Kanno et al., 2018).

Taxon	No. of individuals				Biomass (g)			
	Estimate	SE	z	P	Estimate	SE	z	P
Four benthic fish taxa	0.085	0.018	4.86	**< 0.0001**	0.015	0.003	5.39	**< 0.0001**
Odontobutis obscura and *O. hikimius*	0.082	0.022	3.67	**< 0.001**	0.012	0.004	3.13	**< 0.01**
Tridentiger spp.	0.034	0.023	1.51	0.13	0.012	0.008	1.46	0.14
Cottus pollux ME	0.073	0.018	4.04	**< 0.0001**	0.016	0.008	1.92	0.06
Pseudogobio esocinus esocinus	-0.117	0.097	-1.21	0.23	-0.005	0.004	-1.29	0.20
Seven pelagic fish taxa	0.030	0.005	5.74	**< 0.0001**	0.005	0.001	4.50	**< 0.0001**
Carrassius spp.	0.039	0.028	1.42	0.16	0.002	0.001	1.57	0.12
Candidia temminckii	0.011	0.006	2.02	**< 0.05**	0.004	0.002	2.61	**< 0.01**
Pungtungia herzi	0.033	0.019	1.71	0.09	0.008	0.004	2.21	**< 0.05**
Opsariichthys platypus	0.040	0.000	88.77	**< 0.0001**	0.010	0.008	1.29	0.20
Salvelinus sp.	0.474	0.099	4.79	**< 0.0001**	0.018	0.005	3.61	**< 0.0001**
Oryzias latipes	0.209	0.156	1.34	0.18	1.458	0.952	1.53	0.13
Rhynchocypris oxycephalus jouyi	0.099	0.033	3.01	**< 0.01**	0.011	0.004	2.62	**< 0.01**

Fig. 1. Overview of (a) the study sites and (b) survey design in this study. The bar and arrow in the bottom right-hand corner of the detailed maps in (a) indicate 2 km distance and water flow direction, respectively.

Fig. 2. Relationship between the eDNA concentration quantified by qMiSeq approach and that by species-specific qPCR for (a) *C. temminckii* and (b) *C. pollux* ME. Yellow circles (FK), pink triangles (HS) and blue squares (IN) indicate each study river, respectively. The eDNA samples from the Yokomichi River were omitted from the comparison examination because they were used in another study and there was no remaining volume for analysis. Solid and dashed lines indicate linear regression lines (all regression lines are significant, P < 0.001) and 1:1 line, respectively. The shaded area around each linear regression lines corresponds to the 95% confidence intervals.

Fig. 3. The number of taxa detected by eDNA metabarcoding with qMiSeq (blue), a capture-based survey using electrical shocker (pink) or both methods (grey) at each study site of each river.

Fig. 4. Visualisation of fish communities detected by qMiSeq approach at each study site: (a) Nonmetric multidimensional scaling (NMDS) ordination plot of fish communities and (b) relationship between the qMiSeq eDNA concentration and the abundance or biomass of each taxon for all study sites and study sites included in each site cluster. NMDS ordination plot and site clustering in (a) panel were based on Bray-Curtis dissimilarity index and k-means clustering. Solid black line in (b) panels indicates a linear regression line obtained using all study sites data. Coloured dashed lines indicate linear regression lines obtained using each cluster site data. Coloured letters next to regression lines indicate the rank correlation coefficients (Kendall's τ). If the Kendall's τ value obtained using data from each cluster site was greater than that obtained using data from all study sites, the rank correlation coefficients were expressed in bold.

Fig. 5. Relationship between the qMiSeq eDNA concentrations quantified and capture data
(abundance or biomass) for each taxon at each study site. Each taxa was allocated to
either upper or lower panels according to the abundance/biomass level. Each coloured
solid line indicates nonlinear regressions of a generalized linear model (GLM) based on
negative binomial distribution. Photo copyright: *O. obscura*, *C. temminckii*, *P herzi*, *C.
pollux* ME, *P. esocinus esocinus* and *R. oxycephalus jouyi* for Mr. S. Kunumatsu;
Carassius spp. and *O. latipes* for Mr. Y. Fuke; *O. platypus* and *Salvenius* sp. for ffish.asia
(https://ffish.asia/, 2022.04.18 downloaded); *Tridentiger* spp. for S.T.

Fig. S1 Picture of each survey site.

Fig. S2 (a) K-means partitions comparison and (b) Calinski values.