1

20 years of misunderstanding: Compensatory traits can explain U-shaped cost function of purely sexual traits

Running title: **Performance, ornaments, and compensatory traits**

Masaru HASEGAWA[1]

[1] *Department of Environmental Science, Ishikawa Prefectural University,*

Nonoichi, Ishikawa 921-8836, Japan

Number of table 1

Number of figures 2

Number of supplementary material 2

Author for corresponding

Dr Masaru Hasegawa

E-mail: perorobomusadiobe@gmail.com

Tel: (+81)- 46-858-1562

Acknowledgments

I thank Dr. Emi Arai, Dr. Shumpei Kitamura and his lab members at Ishikawa Prefectural University for their kindest advices and supports. I also thank Prof. Michael D. Jennions, reviewers, and English-proofing company (Enago) for their helpful comments. MH was supported by the KAKENHI grant of the Japan Society for the Promotion of Science (JSPS, 19K06850).

Data Availability Statement

Data attached as Table S1 will be deposited into osf.io once accepted.

Compliance with ethical standards

Conflict if interests   The author declares no conflict of interest.

Ethical approval   This study does not include any treatments of animals, as all the information was gathered from model experiments.

Author Contribution

I performed data analysis and wrote the manuscript alone.

51 # 20 years of misunderstanding: Compensatory traits can
52 ## explain U-shaped cost function of purely sexual traits
53

The cost of ornamentation is often measured experimentally to study the

relative importance of sexual and viability selection for ornamentation, but

these experiments can lead to a misleading conclusion when compensatory

trait is ignored. A famous, classic experiment on the outermost tail feathers

in the barn swallow *Hirundo rustica*, explains that the U-shaped

aerodynamic performance of the outermost tail feathers would be the

evolutionary outcome mainly through viability selection for optimal tail

length, but this conclusion depends on the assumption that compensatory

traits do not cause reduced performance. Using a simple model experiment, I

demonstrated that ornamentation evolved purely though sexual selection can

produce a U-shaped cost function under the presence of compensatory traits.

Therefore, U-shaped cost function (and the low performance of individuals

with reduced ornaments) cannot be used to infer the evolutionary force

favoring ornamentation, due to a previously overlooked concept,

"overcompensation", which can increase, rather than decrease, the apparent

cost of ornamentation.

Keywords: cost-reducing traits, cost of ornamentation, viability selection,

sexual selection

Animals sometimes possess ornamentation that seem to have negative effects

on survivorship, such as long tails and colorful plumage in birds (Andersson

1994). Many empirical studies have shown sexual selection for ornaments,

explaining the evolution of seemingly costly ornaments (e.g., see Hill &

McGraw 2006 for a review on birds). However, the mere presence of sexual

selection does not necessarily mean that sexual selection has been the major

selection force causing ornament elaboration, because focal traits can have

mainly evolved via viability selection and sexual selection may have caused

ornament elaboration only slightly beyond viability optimum (i.e., possible

exaptation; Bergstrom & Dugatkin 2016). Then, it is not surprising to see that

researchers have alternatively focused on the cost of ornamentation (e.g.,

Evans 1998; see below), because the ornaments' cost function would clarify

the relative importance of sexual and viability selection on the focal traits.

A famous example is those testing aerodynamic performance of long

outermost tail feathers in the barn swallows *Hirundo rustica* (Fig. 1; e.g.,

Evans 1998; Buchnan & Evans 2000; Bro-Jørgensen *et al.* 2007). Although

long outermost tail feathers have been shown to be sexually selected (e.g.,

Møller 1988; reviewed in Møller 1994; Turner 2006; Romano *et al.* 2016), it

remains unclear how and why long tails evolved due to the possible

aerodynamic function of long tails (Norberg 1994). Evans and colleagues have

experimentally shortened outermost tail feather length and measured

aerodynamic performances of swallows (e.g., Evans 1998; Buchanan & Evans

2000; Rowe *et al.* 2001; also see Evans & Thomas 1997; Thomas & Rowe 1997

for the detailed descriptions of predictions). They predicted that, if long

outermost tail feathers evolved purely through sexual selection, shortening

the length should produce a decrease in aerodynamic cost (Fig. 1 upper left).

Or, if it evolved purely through viability selection due to the aerodynamic

function, shortening the length should increase the aerodynamic cost (Fig. 1 upper right). Lastly, if viability and sexual selection together favors the evolution of long tails, shortening the tail would decrease the aerodynamic cost first, and then increase the cost beyond the aerodynamic optimum (Fig. 1 upper middle). Their results were consistent with the last prediction with estimated peak values located around 10 mm shorter from the current tail length. Therefore, they concluded that tail feathers mainly evolved through viability selection and sexual selection would have elongated tails only around 10 mm (9–20%) beyond the aerodynamic optimum (and thus many ornithological books follow their conclusions; e.g., Fjeldsa et al. 2020). However, this argument is problematic, because they do not consider compensatory traits, which have been reported to affect aerodynamic cost of long tails in many bird species including barn swallows (e.g., Balmford *et al.* 1994; Barbosa & Møller 1995; Møller *et al.* 1995; Møller 1996). In fact, several compensatory traits have been reported in barn swallows, including greater wingspans, wing area, aspect ratios, reduced wing loading, and so on (e.g., Møller *et al.* 1995; reviewed in Husak & Swallow 2011; also see Moreno & Møller 1996; Saino *et al.* 1997; Tubaro 2003 for other kinds of compensation). Animal performance could be affected by compensatory traits, which is thought to be the case in the peacock *Pavo cristatus*, another model species of sexual selection: peacocks with fully expressed trains, i.e., a costly trait, have a lower metabolic cost of locomotion than those with rudimentary trains possibly due to the presence of compensatory traits (Thavarajah *et al.* 2016). Similar but less striking results, such as no detectable performance cost of gorgeous ornaments, are repeatedly reported in ornamented animals (e.g., Baumgartner *et al.* 2011; Trappet *et al.* 2013; Kojima & Lin 2019).

The importance of compensatory traits has repeatedly been advocated

(e.g., Møller 1996; Oufiero & Garland 2007; Husak *et al.* 2015). Using a

hypothetical data set, Oufiero & Garland (2007) demonstrated that ignoring

a compensatory trait led to an incorrect (and opposite) conclusion. Likewise,

Husak & Swallows (2011) reviewed compensatory traits and proposed that a

simple test of relationships between ornamentation and performance can lead

to misleading conclusions (also see Husak *et al.* 2015 for an updated review).

However, although these studies stress that manipulation experiments

should consider compensatory traits carefully (e.g., Husak & Swallow 2011),

it is still unclear whether and how compensatory traits affect cost function

concerning ornament manipulation. Particularly, whether the low

performance of individuals with reduced ornaments (and hence U-shaped cost

function; see preceding paragraph) can be explained by compensatory traits

is not known. As Oufiero & Garland (2007) used a hypothetical data set, a

simple model experiment rather than complicated empirical data, would be

helpful to demonstrate the potential influence of compensatory traits.

Here, using a simple "toy" model, I examined whether the U-shaped

cost function (see Fig. 1) could be produced when the ornament has evolved

purely through sexual selection for exaggerated ornamentation with the

coevolution of compensatory traits. Although I verbally explained potential

confounding effects of compensatory traits above, our simple model

experiment would directly demonstrate how compensatory traits affects the

performance of (virtual) animals with ornamentation. For this purpose, I used

"chuonchuon" (Fig. 2 upper panel), a traditional balancing toy in Vietnam, as

a model system. Chuonchuon is a suitable model, because its balance is

determined by the whole phenotype, as in the locomotor performance of

animals (e.g., see Husak & Swallow 2011). Different from purely verbal

models, simulations, or mathematical models, this kind of simple, physical

toy model would be intuitive (and thus suitable) for empirical researchers of

animal performance to understand the importance of compensatory traits.

Clearly, this toy model is NOT to test the function of particular traits (e.g.,

swallows' tails), but to test whether compensatory traits can cause reduced

performance under an experimental reduction of ornamental traits that has

evolved purely through sexual selection. Rather than testing the performance

of models with arbitrary values of ornamental and compensatory traits, I first

determined an evolutionary stable form of chuonchuon based on game theory

under the setting that long tails have been sexually selected (i.e., providing

reproductive advantages) and its negative effects on balance (i.e., survival of

chuonchuon) could be reduced by a compensatory trait, extended wings. Then,

I tested the cost function of ornamentation by experimentally shortening tail

length, as in the previous manipulation experiments in swallows listed above.

I discussed the evolutionary implications of the observed pattern.

METHODS

Experimental setup

I used four commercially available "chuonchuon" for my experiment (ca. 7 cm

length: Tirakita, Japan). I used unmanipulated chuonchuon as an ancestral

state, in which neither sexually selected ornamentation nor compensatory

traits have yet been evolved. Then, I used a plastic straw (0.4 g, 4 mm width

and 160 mm length; Strix design, Japan) as an elongated tail. As a

compensatory trait, I made extended wings using clay (0.8 g per wing, i.e.,

twice as heavy as a plastic straw to account for the lever principle; Fig. 2

middle panel).

I here simply set the situation with a set of dichotomous events, that

is, only balanced chuonchuon can "survive," and only longer tailed

chuonchuon in the population can "reproduce." Then, the ancestral form can

survive anytime, but the tail-elongated form cannot survive when elongated

tails make them unbalanced (I used a cut-off point as 30° deviation from the

horizontal plane, here). Tail elongation made models unbalanced in the range

of the current study (i.e., maximum length to −80 mm shortened tails; see Fig.

1) and thus they cannot survive without compensatory traits, i.e., extended

wings. Even when it survived, the ancestral form cannot have descendants

when any derived forms with elongated tails (i.e., those with longer tails than

ancestral form) exist. Because my interest here is not in the actual

evolutionary trajectory but the cost function of the derived form, I obtained

the evolutionary stable form by comparing the "fitness" of each form. For

simplicity, I assumed continuous variation in tail length but assumed two

discrete states of the extended wings (i.e., with and without extended wings).

In this case, the evolutionary stable form is a chuonchuon with maximum

elongated tails with extended wings. This is because chuonchuon with

maximum elongated tails and extended wings can survive (i.e., balanced) and

thus reproduce under any circumstances (i.e., they have the longest tails

compared to all other forms). I did not subdivide compensatory traits (e.g.,

small/medium/large), because the purpose of the current study is not to

quantify the exact peak performance or identify the best size of compensatory

traits but to examine the effect of compensatory traits.

Measuring performance

As in manipulation experiments in the barn swallow (Buchanan & Evans

2000; Rowe *et al.* 2001), I shortened the extended tails of chuonchuon models

(20 mm/40 mm/60 mm/80 mm, here) and took two pictures each time. From

each of the two pictures, I measured the absolute angle of chuonchuon

deviated from the horizontal plane as stability (or balance-ability: Fig. 2 middle panel) using ImageJ software. Then, I averaged the two measurements (repeatability = 0.99, $F_{19,20}$ = 270.24, P < 0.0001; Lessells & Boag 1987) to have a representative estimate of stability for each treatment.

Statistics

To account for individual variation in performance, I used a Bayesian linear mixed-effects model with a normal error distribution to examine the stability of chuonchuon given tail shortening. Individual identity was used as a random effect. By calculating the Brooks-Gelman-Rubin statistic (Rhat), which must be <1.2 for all parameters (Kass *et al.* 1998), the reproducibility of the MCMC simulation was confirmed. Data analyses were conducted using the R statistical package (ver. 4.1.0; R Core Team 2021), using the function "MCMCglmm" (with its default setting) in the package "MCMCglmm" (Hadfield 2010).

RESULTS

I found a significant quadratic relationship between tail length manipulation and stability of the chuonchuon (Table 1; Fig. 2 lower panel): the stability of the chuonchuon, measured as an absolute angle deviated from the horizontal plane, increased with decreasing tail length until the peak value where chuonchuon had the maximum estimated stability and further tail shortening beyond the peak value decreased stability.

DISCUSSION

The main finding from the chuonchuon experiment is that the apparent U-shaped cost function can evolve purely through sexual selection in the

presence of compensatory traits (i.e., with no viability advantage of

ornamentation). The importance of compensatory traits has repeatedly been

advocated (reviewed in Møller 1996; Husak & Swallow 2010; Husak *et al.*

2015), but previous studies assume that compensatory traits still function as

"cost-reducing" traits regardless of the situation (e.g., see Husak *et al.* 2015,

p. 18). Here, I showed that "overcompensation" could decrease performance,

producing a U-shaped cost function. As an old Chinese proverb say, "too much

is as bad as too little" at least in some cases. The assumption that the cost

function can reveal the relative importance of sexual and viability selection

(e.g., Buchanan & Evans 2000) is therefore unreliable. In other words, the

prediction of sexual selection (i.e., Fig. 1C) is no longer valid, and can be

replaced by U-shaped cost function as in Fig. 1B, indicating that the

presence/absence of viability selection cannot be detected (see Fig. S1 for the

corrected prediction).

Of course, the current experiment using chuonchuon models does not

mean that sexual selection always produces a U-shaped cost function (and

does not mean that the same physical property, i.e., stability, determined the

evolution of ornamentation, such as swallows' tail, and compensatory traits).

Still, even without formal mathematical model, it seems likely that

overcompensation will often reduce locomotor performance, particularly

when locomotion requires co-opted, integrated phenotype (e.g., aerial

locomotion). In fact, concerning the function of long tails in barn swallows,

Norberg (1994) mentioned that "*any experimental shortening and*

lengthening of the outer tail feathers is likely to upset an original co-adapted

character set..." (p. 231). When compensatory traits co-evolved with the

sexually selected tail length to minimize the aerodynamic cost of tail length

(Møller 1996), an experimental reduction in tail length alone cannot clarify

the aerodynamic cost of tail length, because the manipulation did not follow the actual evolutionary pathway (see Fig. 3 for a schematic illustration; also see Fig. S2 for an alternative scenario in which long tails have some aerodynamic function). Also, all macroevolutionary studies of these aerial insectivores so far have supported sexual (rather than viability) selection explanation (e.g., Hasegawa *et al.* 2016; Hasegawa & Arai 2017, 2018, 2020a,b, 2021, 2022), demanding further experimental studies, possibly using a smaller amount of manipulation (so that overcompensation can be avoided) in many species with various ornamentation (i.e., phylogenetic comparative "experiment"). An alternative approach, the manipulation of compensatory traits (sensu Møller 1996; see also Fig. 3) would in theory be valuable to show the effect of overcompensation, but is impractical due to the multi-dimensionality of compensatory traits (e.g., manipulations of wing size and shape are often difficult to conduct).

In addition, the current experiment provides useful insights. First, the cost of ornamentation can be transformed from one type to another. Extended wings enhanced the stability of chuonchuon with elongated tails, though they required additional investment (i.e., a production cost) instead (also see Møller 1996 for a similar argument). Furthermore, the production cost of compensatory traits can be much higher than the production cost of the ornament itself (see Methods). It is straightforward to conclude that a single measure of cost (e.g., aerodynamic cost) is inappropriate to approximate the total cost (and thus viability disadvantage) of ornamentation, even when the production cost of ornamentation itself seems negligible at first glance, as is the case for the outermost tail feathers in barn swallows (Fig. 1).

In summary, the current model experiment demonstrated that a U-

shaped cost function, which is often thought to result from viability advantage

of focal ornaments, can be observed when sexual selection has favored the

evolution of the costly ornaments, due to the presence of compensatory traits.

Given that compensatory traits coevolved with sexual traits (Møller 1996),

the manipulation of sexual traits alone would not clarify the evolutionary

force on sexual traits (see Figs. 3 & S2). Future studies should consider that

overcompensation can be detrimental and that more sophisticated

experimental design is needed when inferring selection pressure on (and the

evolutionary history of) ornamentation.

Acknowledgment   See Title page.

Author Contribution   See Title page.

Conflict statement   See Title page.

Permits   See Title page

Data availability   See Title page
306

REFERENCES

Andersson, M. 1994. *Sexual selection*. Princeton: Princeton Univ. Press.

Barbosa, A. & Møller, A.P. 1999. Aerodynamic costs of long tails in male

barn swallows *Hirundo rustica* and the evolution of sexual size

dimorphism. *Behav. Ecol.* **10**: 128–135.

https://doi.org/10.1093/beheco/10.2.128

Balmford, A., Jones, I.L. & Thomas, A.L.R. 1994. How to compensate for

costly sexually selected tails: the origin of sexually dimorphic wings in

long-tailed birds. *Evolution* **48**: 1062–1070.

https://doi.org/10.2307/2410366

Baumgartner, A., Coleman, S. & Swanson, B. 2011. The Cost of the Sword:

Escape Performance in Male Swordtails. *PLoS ONE* **6**: e15837.

https://doi.org/10.1371/journal.pone.0015837

Bergstrom, C.T. & Dugatkin, L.A. 2016. *Evolution*. New York: WW Norton &

Co. Inc.

Bro-Jørgensen, J., Johnstone, R.A. & Evans, M.R. 2007. Uninformative

exaggeration of male sexual ornaments in barn swallows. *Curr. Biol.* **17**:

850–855. https://doi.org/10.1016/j.cub.2007.03.042

Buchanan, K.L. & Evans, M.R. 2000. The effect of tail streamer length on

aerodynamic performance in the barn swallow. *Behav. Ecol.* **11**: 228–238.

https://doi.org/10.1093/beheco/11.2.228

Evans, M.R. 1998. Selection on swallow tail streamers. *Nature* **394**: 233–

234. https://doi.org/10.1038/28297

Evans, M.R. & Thomas, A.L.R. 1997. Testing the functional significance of

tail streamers. *Proc. R. Soc. Lond. B* **264**: 211–217.

https://doi.org/10.1098/rspb.1997.0030

Fjeldsa, J., Christidis, L., Ericson, P.G.P. 2022. *The largest avian radiation:*

the evolution of perching birds, or the order Passeriformes. Lynx

edicions, Barcelona.

Hadfield, J.D. 2010. MCMC methods for multi-response generalized linear

mixed models: The MCMCglmm R package. *J. Stat. Soft.* **33**: 1–22.

https://doi.org/10.18637/ jss.v033.i02

Hasegawa, M., Arai, E. & Kutsukake, N. 2016. Evolution of tail fork depth

in genus *Hirundo. Ecol. Evol.* **6**: 851–858.

https://doi.org/10.1002/ece3.1949

Hasegawa, M. & Arai, E. 2017. Egg size decreases with increasing female

tail fork depth in family Hirundinidae. *Evol. Ecol.* **31**: 559–569.

https://doi.org/10.1007/s1068 2-017-9895-2

Hasegawa, M. & Arai, E. 2018. Convergent evolution of the tradeoff between

egg size and tail fork depth in swallows and swifts. *J. Avian Biol.* **49**:

e01684. https://doi.org/10.1111/jav.01684

Hasegawa, M. & Arai, E. 2020a. Fork tails evolved differently in swallows

and swifts. *J. Evol. Biol.* **33**: 911–919. https://doi.org/10.1111/jeb.13622

Hasegawa, M. & Arai, E. 2020b. Correlated evolution of biparental

incubation and sexual tail monomorphism in swallows and martins

(Aves: Hirundinidae). *Evol. Ecol.* **34**: 777–788.

https://doi.org/10.1007/s10682-020-10066-w

Hasegawa, M. & Arai, E. 2021. Opposing population trends of fork-tailed

swallows and reddish-coloured swallows in our changing world. *J. Evol.*

Biol. **34**: 331–338. https://doi.org/10.1111/jeb.13736

Hasegawa, M. & Arai, E. 2022. Extrapair mating opportunity explains the

evolutionary transition between the presence and absence of tail

ornamentation in swallows. *Evol. Ecol.* 36: 899–913. doi:

10.1007/210682-022-10196-3

Hill, G.E. & McGraw, K.J. 2006. *Bird coloration II: function and evolution.*

Cambridge: Harvard Univ. Press.

Husak, J.F. & Swallow, J.G. 2011. Compensatory traits and the evolution of

male ornaments. *Behaviour* **148**: 1–29.

https://doi.org/10.1163/000579510X541265

Husak, J.F., Henningsen, J.P., Vanhooydonck, B. & Irschick, D.J. 2015. A

performance-based approach to studying costs of reliable signals. In:

Irschick, D.J. Briffa, M. & Podos, J. (eds) *Animal signaling and function:*

an integrative approach. Hoboken: Wiley Blackwell, pp. 47–74.

Kass, R.E., Carlin, B.P., Gelman, A. & Neal, R. 1998. Markov chain Monte

Carlo in practice: A round table discussion. *Amer. Stat.* **52**: 93–100.

https://doi.org/10.2307/2685466

Kojima, W. & Lin, C.P. 2019. Sprint speed is not reduced by exaggerated

male weapons in a flower beetle *Dicronocephalus wallichii. Ethology*

125: 47–56. https://doi.org/10.1111/eth.12824

Lessells, C.M. & Boag, P.T. 1987. Unrepeatable repeatabilities: A common

mistake. *Auk* **104**: 116–121. https://doi.org/10.2307/4087240

Matyjasiak, P., Marzal, A., Navarro, C., de Lope, F. & Møller, A.P. 2009. Fine

morphology of experimental tail streamers and flight manoeuvrability in

the house martin *Delichon urbica. Funct. Ecol.* **23**: 389–396.

https://doi.org/10.1111/j.1365-2435.2008.01490.x

Moreno, E. & Møller, A.P. 1996. Morphological aspects of avian tail

movements: a functional approach in hirundines. *Auk* **113**: 647–654.

https://doi.org/10.2307/4088985

Møller, A.P. 1988. Female choice selects for male sexual tail ornaments in

the monogamous swallow. *Nature* **332**: 640–642.

https://doi.org/10.1038/332640a0

Møller, A.P. 1994. *Sexual selection and the barn swallow*. Oxford: Oxford

Univ. Press.

Møller, A.P. 1996. The cost of secondary sexual characters and the evolution

of cost-reducing traits. *Ibis* **138**: 112–119. https://doi.org/10.1111/j.1474-

919X.1996.tb04317.x

Møller, A.P., de Lope, F. & Saino, N. 1995. Sexual selection in the barn

swallow *Hirundo rustica*. VI. Aerodynamic adaptations. *J. Evol. Biol.* **8**:

671–687. https://doi.org/10.1046/j.1420-9101.1995.8060671.x

Norberg, R. 1994. Swallow tail streamer is a mechanical device for

selfdeflection of tail leading edge enhancing aerodynamic efficiency and

flight manoeuvrability. *Proc. R. Soc. Lond. B* **257**: 227–233.

https://doi.org/10.1098/rspb.1994.0119

Oufiero, C.E. & Garland, Jr T. 2007. Evaluating performance costs of

sexually selected traits. *Funct. Ecol.* **21**: 676–689.

https://doi.org/10.1111/j.1365-2435.2007.01259.x

R Core Team. 2021. R: A language and environment for statistical

computing. R Foundation for Statistical Computing, Vienna, Austria.

Available at http://www.R-project.org/ (accessed 11 September 2021)

Romano, A., Constanzo, A., Rubolini, D., Saino, N. & Møller, A.P. 2017.

Geographical and seasonal variation in the intensity of sexual selection

in the barn swallow *Hirundo rustica*: A metaanalysis. *Biol. Rev.* **92**:

1582–1600. https://doi.org/10.1111/brv.12297

Rowe, L.V., Evans, M.R. & Buchanan, K.L. 2001. The function and evolution

of the tail streamer in hirundines. *Behav. Ecol.* **12**: 157–163.

https://doi.org/10.1093/beheco/12.2.157

Saino, N., Cuervo, J.J., Ninni, P., de Lope, F. & Møller, A.P. 1997.

Haematocrit correlates with tail ornament size in three populations of

barn swallow (*Hirundo rustica*). *Funct. Ecol.* **11**: 604–610.

https://doi.org/10.1046/j.1365-2435.1997.00131.x

Thavarajah, N.K., Tickle, P.G., Nudds, R.L. & Codd, J.R. 2016. The peacock

train does not handicap cursorial locomotor performance. *Sci. Rep.* **6**:

36512. https://doi.org/10.1038/srep36512

Thomas, A.L.R. & Rowe, L.V. 1997. Experimental tests on tail elongation

and sexual selection in swallows (*Hirundo rustica*) do not affect the tail

streamer and cannot test its function. *Behav. Ecol.* **8**: 580–581.

https://doi.org/10.1093/beheco/8.5.580

Trappett, A., Condon, C.H., White, C., Matthew, P. & Wilson, R.S. 2013.

Extravagant ornaments of male threadfin rainbowfish (*Iriatherina*

werneri) are not costly for swimming. *Funct. Ecol.* **27**: 1034–1041.

https://doi.org/10.1111/1365-2435.12097

Tubaro, P.L. 2003. A comparative study of aerodynamic function and flexural

stiffness of outer tail feathers in birds. *J. Avian Biol.* **34**: 243–250.

https://doi.org/10.1034/j.1600-048X.2003.03084.x

Turner, A.K. 2006. *The barn swallow.* London: T & AD Poyser.

Table 1.   Bayesian linear mixed-effects model of stability of chuonchuon

in relation to experimental manipulation of elongated tails (N_{ID} = 4, $Ntotal$ =

20).

Variable	Coefficient [95% CI]	P_{MCMC}
Intercept	**22.37 [18.95, 25.97]**	**<0.0001**
Linear term of manipulation	**0.96 [0.71, 1.23]**	**<0.0001**
Quadratic term of manipulation	**8.36 [6.21, 10.51]**	**<0.0001**

438 The dependent variable is the absolute angle of chuonchuon deviated from

439 the horizontal plane. I included chuonchuon ID as a random factor (variance

440 = 0.20) in addition to units (variance = 15.39).

441 Significant test results (i.e. P < 0.05) are indicated in bold.

443 Figure legends

Figure 1.   Details of manipulation experiment of Evans and colleagues

(see Evans 1998; Buchanan & Evans 2000; Rowe *et al.* 2001 for original

figures; also see Evans & Thomas 1997; Thomas & Rowe 1997 for similar

predictions). The vertical line in each plot indicates current ornament

expression, and the triangle indicates ornament expression providing peak

performance (note that lower values indicate higher performance here).

Although they originally called deviation of outermost tail feathers (or

possibly, narrowed part of outermost tail feather length) as a "streamer," I

avoided this wording, because it implies some kinds of aerodynamic function

of the ornaments, and because the formal definition of the streamer is unclear

(and not to confound size and shape: Matyjasiak *et al.* 2009). Likewise, I used

viability selection rather than natural selection to be precise.

Figure 2.   Chuonchuon (upper panel), manipulated chuonchuon with elongated tails and extended wings (middle panel), and results of chuonchuon experiment (lower panel). A simple quadratic regression line is denoted (see Table 1 for a formal analysis). Individual identity was denoted by different coloration. Mean ± SE values of stability, measured by absolute angle deviated from horizontal plane, in ancestral state shown in upper panel was 4.87 ± 1.78 (range: 1.38–8.87), and, not surprisingly, tail elongation without extended wings made models unbalanced in the range of the current study (e.g., when elongated tails = 30 mm, which corresponds to the manipulation = −80 mm, mean stability ± SE = 36.60 ± 2.30, range = 32.69–42.93).

Figure 3.   Schematic illustration of conceptual fitness landscape measured

as flight performance in relation to tail length and phenotypic expression of

compensatory traits (upper panel). Darker background coloration indicates

lower flight performance (note that the best flight performance is attained in

the lower left corner). Dark arrow indicates hypothetical evolutionary

pathway to the current state (filled circle). Light arrow indicates phenotypic

change when researchers experimentally reduced tail length (in which

compensatory traits remain unchanged). Lower panel shows flight cost

measurement in relation to tail length along with dark or light arrows. The

evolutionary pathway on this fitness landscape is biased toward right side

because of intense sexual selection favoring long tails (i.e., actual

evolutionary pathway depends on sexual selection in addition to viability

selection due to flight cost and other costs, such as production costs of tail

length and compensatory traits; see text). Narrow contour width indicates

that small deviation from integrated, co-opted trait sets had a strong negative

effect on flight performance, as predicted by Norberg (1994).

525 **Figure S1.** The "corrected" predictions for manipulation experiment of Evans

526 and colleagues under the presence of compensatory traits. Note that two type

527 of scenarios (B and C) cannot be distinguished. See text and Fig. 1 for detailed

528 explanations (also see Figs. 3 & S2).

Figure S2.   Schematic illustration of conceptual fitness landscape

measured as flight performance in relation to tail length and phenotypic

expression of compensatory traits (upper panel). Darker background

coloration indicates lower flight performance (note that the lightest

background located near the current state, shown as a filled circle, in this

example). Dark arrow indicates hypothetical evolutionary pathway to the

current state. Light arrow indicates phenotypic change when researchers

experimentally reduced tail length (in which compensatory traits remain

unchanged). Lower panel shows flight cost measurement in relation to tail

length along with dark or light arrows. The evolutionary pathway on this

fitness landscape is biased toward right side because of intense sexual

selection favoring long tails (i.e., actual evolutionary pathway depends on

sexual selection in addition to viability selection due to flight cost and other

costs, such as production costs of tail length and compensatory traits; see text).

Narrow contour width indicates that small deviation from integrated, co-

opted character sets had a strong negative effect on flight performance, as

predicted by Norberg (1994).