Compensatory traits can explain concave cost function of purely sexual traits

Short title: **Performance, ornaments, and compensatory traits**

Masaru HASEGAWA[1]*

[1] Department of Environmental Science, Ishikawa Prefectural University, Nonoichi, Ishikawa 921-8836, Japan

Number of table 1

Number of figures 3

Number of supplementary materials 5

*Corresponding author

Dr Masaru Hasegawa

E-mail: perorobomusadiobe@gmail.com

Tel: (+81)- 46-858-1562

Abstract

The cost of ornamentation is often measured experimentally to study the relative importance of sexual and viability selection for ornamentation, but these experiments can lead to a misleading conclusion when compensatory trait is ignored. For example, a classic experiment on the outermost tail feathers in the barn swallow *Hirundo rustica*, explains that the concave (or U-shaped) aerodynamic performance cost of the outermost tail feathers would be the evolutionary outcome through viability selection for optimal tail length, but this conclusion depends on the assumption that compensatory traits do not cause reduced performance. Using a simple "toy model" experiment, I demonstrated that ornamentation evolved purely though sexual selection can produce a concave cost function under the presence of compensatory traits, which was further reinforced by a simple mathematical model. Therefore, concave cost function (and the low performance of individuals with reduced ornaments) cannot be used to infer the evolutionary force favoring ornamentation, due to a previously overlooked concept, "overcompensation", which can increase the apparent cost of ornamentation.

Keywords

cost-reducing traits, cost of ornamentation, viability selection, sexual selection.

Introduction

Animals sometimes possess ornamentation that seem to have negative effects on survivorship, such as long tails and colorful plumage in birds (Andersson, 1994). Many empirical studies have shown sexual selection for ornaments, explaining the evolution of seemingly costly ornaments (e.g., see Hill & McGraw, 2006 for a review on birds). However, the mere presence of sexual selection does not necessarily mean that sexual selection has been the major selection force causing ornament elaboration, because focal traits can have mainly evolved via viability selection and sexual selection may have caused ornament elaboration only slightly beyond viability optimum (i.e., possible exaptation; Bergstrom & Dugatkin, 2016). Then, it is not surprising to see that researchers have alternatively focused on the cost of ornamentation (e.g., Evans, 1998; see below), because the ornaments' cost function would clarify the relative importance of sexual and viability selection on the focal traits.

A famous example is those testing aerodynamic performance of long outermost tail feathers in the barn swallows *Hirundo rustica* (Fig. 1; e.g., Evans, 1998; Buchnan & Evans, 2000; Bro-Jørgensen et al., 2007). Although long outermost tail feathers have been shown to be sexually selected (e.g., Møller, 1988; reviewed in Møller, 1994; Turner, 2006; Romano et al., 2016), it remains unclear how and why long tails evolved due to the possible aerodynamic function of long tails (Norberg, 1994). Evans and colleagues have experimentally shortened outermost tail feather length and measured aerodynamic performances of swallows (e.g., Evans, 1998; Buchanan & Evans, 2000; Rowe et al., 2001; also see Evans & Thomas, 1997; Thomas & Rowe, 1997 for the detailed descriptions of predictions). They predicted that, if long outermost tail feathers evolved purely through sexual selection,

shortening the length should produce a decrease in aerodynamic cost (Fig. 1 upper left). Or, if it evolved purely through viability selection due to the aerodynamic function, shortening the length should increase the aerodynamic cost (Fig. 1 upper right). Lastly, if viability and sexual selection together favors the evolution of long tails, shortening the tail would decrease the aerodynamic cost first, and then increase the cost beyond the aerodynamic optimum (Fig. 1 upper middle). Their results were consistent with the last prediction with estimated peak values located around 10 mm shorter from the current tail length. Therefore, they concluded that tail feathers mainly evolved through viability selection and sexual selection would have elongated tails only around 10 mm (9–20%) beyond the aerodynamic optimum (and thus many ornithological books follow their conclusions; e.g., Fjeldsa et al., 2020). However, this argument is problematic, because they do not consider compensatory traits (also known as cost-reducing traits), which have been reported to affect aerodynamic cost of long tails in many bird species including barn swallows (e.g., Balmford et al., 1994; Barbosa & Møller, 1995; Møller et al., 1995; Møller, 1996). In fact, several compensatory traits have been reported in barn swallows, including greater wingspans, wing area, aspect ratios, reduced wing loading, and so on (e.g., Møller et al., 1995; reviewed in Husak & Swallow, 2011; also see Moreno & Møller, 1996; Saino et al., 1997; Tubaro, 2003 for other kinds of compensation). Animal performance could be affected by compensatory traits, which is thought to be the case in the peacock *Pavo cristatus*, another model species of sexual selection: peacocks with fully expressed trains, i.e., a costly trait, have a lower metabolic cost of locomotion than those with rudimentary trains possibly due to the presence of compensatory traits (Thavarajah et al., 2016). Similar but less striking results, such as no detectable performance

cost of gorgeous ornaments, are repeatedly reported in ornamented animals (e.g., Baumgartner et al., 2011; Trappet et al., 2013; Kojima & Lin, 2019). Assuming costly nature of ornamental traits, these findings are puzzling (although the possibility that all these seeming costly ornaments are virtually cost-free cannot be excluded).

The importance of compensatory traits has repeatedly been advocated (e.g., Møller, 1996; Oufiero & Garland, 2007; Husak et al., 2015). Using a hypothetical data set, Oufiero & Garland (2007) demonstrated that ignoring a compensatory trait led to an incorrect (and opposite) conclusion, at least in a correlational study, because the focal costly trait appeared to have a positive effect on performance if a compensatory trait was not taken into account. Likewise, Husak & Swallow (2011) reviewed compensatory traits and proposed that a simple test of relationships between ornamentation and performance can lead to misleading conclusions (also see Husak et al., 2015 for an updated review). However, although these studies stress that correlational and even manipulation experiments should consider compensatory traits carefully (e.g., Husak & Swallow, 2011), it is still unclear whether and how compensatory traits affect cost function concerning ornament manipulation. Particularly, whether the low performance of individuals with reduced ornaments (and hence concave cost function; see preceding paragraph) can be explained by compensatory traits is not known. As Oufiero & Garland (2007) used a hypothetical data set, a simple model experiment rather than complicated empirical data, would be helpful to demonstrate the potential influence of compensatory traits.

Here, using a simple "toy" model, I examined whether the concave cost function (see Fig. 1) could be produced when the ornament has evolved purely through sexual selection for exaggerated ornamentation with the

coevolution of compensatory traits. Although I verbally explained potential confounding effects of compensatory traits above, our simple model experiment would directly demonstrate how compensatory traits affects the performance of (virtual) animals with ornamentation. For this purpose, I used "chuonchuon" (Fig. 2 upper panel), a traditional balancing toy in Vietnam, as a model system. Chuonchuon is a suitable model, because its balance is determined by the whole phenotype, as in the locomotor performance of animals (e.g., see Husak & Swallow, 2011). Different from purely verbal models, simulations, or mathematical models, this kind of simple, physical toy model would be intuitive (and thus suitable) for empirical researchers of animal performance to understand the importance of compensatory traits. Clearly, this toy model is NOT to test the function of particular traits (e.g., swallows' tails), but to test whether compensatory traits can cause reduced whole-body performance under an experimental reduction of ornamental traits that has evolved purely through sexual selection. Rather than testing the performance of models with arbitrary values of ornamental and compensatory traits, I first determined an evolutionary stable form of chuonchuon based on game theory under the setting that long tails have been sexually selected (i.e., providing reproductive advantages) and its negative effects on balance (i.e., survival of chuonchuon) could be reduced by a compensatory trait, extended wings. Then, I tested the cost function of ornamentation by experimentally shortening tail length, as in the previous manipulation experiments in swallows listed above. To test the generality of the finding of chuonchuon, I also provided a simple (or even rudimentary) mathematical model. Even such a simple model would be useful to determine whether and, if any, in which condition the apparent concave cost function of ornament can be found.

I discussed the evolutionary implications of the observed pattern.

Material and methods

Experimental setup

I used four commercially available "chuonchuon" for my experiment (ca. 7 cm

length: Tirakita, Japan). I used unmanipulated chuonchuon as an ancestral

state, in which neither sexually selected ornamentation nor compensatory

traits have yet been evolved. Then, I used a plastic straw (0.4 g, 4 mm width

and 160 mm length; Strix design, Japan) as an elongated tail. As a

compensatory trait, I made extended wings using clay (0.8 g per wing, i.e.,

twice as heavy as a plastic straw to account for the lever principle; Fig. 2

middle panel).

I here simply set the situation with a set of dichotomous events, that

is, only balanced chuonchuon can "survive," and only longer tailed

chuonchuon in the population can "reproduce." Then, the ancestral form can

survive anytime, but the tail-elongated form cannot survive when elongated

tails make them unbalanced (I used a cut-off point as 30° deviation from the

horizontal plane, here). Tail elongation made models unbalanced in the

range of the current study (i.e., maximum length to −80 mm shortened tails;

see Fig. 1) and thus they cannot survive without compensatory traits, i.e.,

extended wings. Even when it survived, the ancestral form cannot have

descendants when any derived forms with elongated tails (i.e., those with

longer tails than ancestral form) exist. Because my interest here is not in the

actual evolutionary trajectory but the cost function of the derived form, I

obtained the evolutionary stable form by comparing the "fitness" of each

form. For simplicity, I assumed continuous variation in tail length but

assumed two discrete states of the extended wings (i.e., with and without

extended wings). In this case, the evolutionary stable form is a chuonchuon with maximum elongated tails with extended wings. This is because chuonchuon with maximum elongated tails and extended wings can survive (i.e., balanced) and thus reproduce under any circumstances (i.e., they have the longest tails compared to all other forms). I did not subdivide compensatory traits (e.g., small/medium/large), because the purpose of the current study is not to quantify the exact peak performance or identify the best size of compensatory traits but to examine the effect of compensatory traits.

Measuring performance

As in manipulation experiments in the barn swallow (Buchanan & Evans, 2000; Rowe et al., 2001), I shortened the extended tails of the evolutionary stable form (i.e., a chuonchuon with maximum elongated tails as ornamentation with extended wings as a compensatory trait; see above) of chuonchuon models by 0 mm, 20 mm, 40 mm, 60 mm, and 80 mm, and took two pictures each time. From each of the two pictures per individual chuonchuon, I measured the absolute angle of chuonchuon deviated from the horizontal plane as stability (or balance-ability: Fig. 2 middle panel) using ImageJ software. Then, I averaged the two measurements (repeatability = 0.99, $F_{19,20}$ = 270.24, P < 0.0001; Lessells & Boag, 1987) to have a representative estimate of stability for each treatment (i.e., tail shortened by 0 mm/20 mm/40 mm/60 mm/80 mm) in each chuoncuhon.

Statistics

To account for individual variation in performance, I used a Bayesian linear mixed-effects model with a normal error distribution to examine the stability

of chuonchuon given tail shortening. Here, stability, measured as the absolute angle of chuonchuon (see above), was used as a response variable. Individual identity was used as a random effect. By calculating the Brooks-Gelman-Rubin statistic (Rhat), which must be <1.2 for all parameters (Kass et al., 1998), the reproducibility of the MCMC simulation was confirmed. Data analyses were conducted using the R statistical package (ver. 4.1.0; R Core Team, 2021), using the function "MCMCglmm" (with its default setting) in the package "MCMCglmm" (Hadfield, 2010).

Mathematical model

To test whether a quadratic cost function of purely sexual traits (see above) can be found in the presence of compensatory traits, I used a simple mathematical model. I used here an equation for a bivariate quadratic selection surface, characterized by slope and curvilinear terms, as commonly used in evolutionary biology (cf. Lande & Arnold, 1983; Arnold et al., 2001; Arnold, 2003; Stinchcombe et al., 2008; see Results). This selection surface can be used to approximate complex empirical performance surfaces regardless of the shape of the actual performance surface (Arnold, 2003). I used the same symbols for each term as those used in the literature on selection surface (e.g., $0.5*\gamma_{ii}$, instead of q_{ii}, though the latter is often used in multiple regression models; Stinchcombe et al., 2008).

Results

Toy model

I found a significant quadratic relationship between tail length manipulation and stability of the chuonchuon (Table 1; Fig. 2 lower panel): the stability of the chuonchuon, measured as an absolute angle deviated from the horizontal

plane, increased with decreasing tail length until the peak value where

chuonchuon had the maximum estimated stability and further tail

shortening beyond the peak value decreased stability.

Mathematical model

Suppose that animal performance can be approximated using the following

equation (cf. Lande & Arnold, 1983; Arnold et al., 2001; Arnold, 2003;

Stinchcombe et al., 2008):

$$\text{Performance} = 0.5*\gamma_{11}*x_1^2 + \beta_1*x_1 + 0.5*\gamma_{22}*x_2^2 + \beta_2*x_2 + \gamma_{12}*x_1*x_2 + \alpha,$$

where x_1 denotes the expression of the ornament, x_2 denotes the expression of

compensatory trait(s), and γ_{11}, γ_{22}, γ_{12}, β_1, and β_2 are coefficients for each

variable with α as a constant.

For simplicity, imagine an elliptical performance surface where the

optimal performance can be set at $(x_1,x_2) = (0,0)$, so that any additions of the

ornament and compensatory traits decrease performance, which is

accomplished by the condition, $\beta_1 = \beta_2 = 0$, with $\gamma_{11} < 0$, $\gamma_{22} < 0$, and $\gamma_{12}^2 <$

$\gamma_{11}*\gamma_{22}$ (note that, when $\gamma_{12}^2 > \gamma_{11}*\gamma_{22}$, there will be a saddle in the performance

surface: Stinchcombe et al. 2008). Assuming that the performance surface

has a hill with an upward tilting axis (i.e., for each ornament expression, the

decline of performance can be minimized by the presence of certain amounts

of compensatory traits, which is accomplished by $\gamma_{12} > 0$; see Fig. 3 for an

example), the above equation can be reduced as follows:

$$\text{Performance} = 0.5*\gamma_{11}* x_1^2 + 0.5*\gamma_{22}*x_2^2 + \gamma_{12}*x_1*x_2 + \alpha,$$

with the condition, $\gamma_{11} < 0$, $\gamma_{22} < 0$, $\gamma_{12}^2 < \gamma_{11}*\gamma_{22}$, and $\gamma_{12} > 0$. In this bivariate

performance surface, the experimental manipulation of the ornament is

equivalent to changing the x_1 value while x_2 value remains unchanged from

the current value of compensatory traits (say, k; note that k > 0 under the

presence of compensatory traits). Then, by substituting y = k into the above

equation, the quadratic function of x_1 is obtained:

$$\text{Performance} = 0.5*\gamma_{11}*x_1^2 + \gamma_{22}*k^2 + \gamma_{12}*k*x_1 + \alpha.$$

This can be re-written as follows:

$$\text{Performance} = 0.5*\gamma_{11}(x_1 + k*\gamma_{12}/\gamma_{11})^2 + \text{Constant}.$$

Now, it is clear that we can have a convex performance function (and hence,

concave cost function) of ornament, because $k*\gamma_{12}/\gamma_{11} < 0$ (note that k > 0, $\gamma_{12} >$

0, and $\gamma_{11} < 0$; see above). Given that sexual selection favors the exaggeration

of the ornament (i.e., the bivariate sexual selection surface can be an

increasing function of x_1), the current value of the ornament should be

located on the right side of the ridge line (i.e., beyond the upward tilting axis

of the hill; see above; also see Fig. 3) where sexually selected advantages offset

low performance of ornamented animals. When experimentally reducing the

ornament expression, we find that performance increases until the ornament

is reduced to be $k*\gamma_{12}/\gamma_{11}$, and then decreases (Fig. 3).

Retrospectively, this equation indicates that an incremental cost

function of sexual traits, which is supposed to be common under

manipulative experiments of the ornament (see Fig. 1C), can be predicted

only when $k*\gamma_{12}/\gamma_{11} \geq 0$. This can be accomplished when k = 0 (i.e., when

compensatory traits are absent for some reason, such as lack of genetic

variation) or when $\gamma_{12} \leq 0$ (i.e., when seemingly compensatory traits are not

actually compensatory traits but just costly traits like ornament with their

joint costs being additive or multiplicative). Although this condition is

confined to an elliptical performance surface as assumed above (i.e., when

$\gamma_{12}^2 > \gamma_{11}{*}\gamma_{22}$), this is a reasonable assumption. When there is a saddle in the

performance surface (e.g., see Fig. S1 left panel), there is no optimal

performance different from elliptical performance surface. Furthermore, it

accompanies the changing function of the ornament, in which ornament is

not a purely sexual trait but can be a functional trait in some ranges of

variables, which is beyond the scope of the current study (because here we

focused on purely sexual traits; see Introduction). Under the condition $\gamma_{12}^2 =$

$\gamma_{11}{*}\gamma_{22}$, the cost of the ornament is perfectly compensated by compensatory

traits, and thus animals can have any ornament value together with the

corresponding compensatory traits without performance costs (Fig. S1 right

panel). In other words, the cost of ornament can be virtually absent under

this condition. As in elliptical performance surfaces (see above), performance

increases until the ornament is reduced to $k{*}\gamma_{12}/\gamma_{11}$, and then decreases in

this case.

Discussion

The main finding from the chuonchuon experiment is that the apparent

concave cost function can evolve purely through sexual selection in the

presence of compensatory traits (i.e., with no viability advantage of

ornamentation), which is further reinforced by a mathematical model. The

importance of compensatory traits has repeatedly been advocated (reviewed

in Møller, 1996; Husak & Swallow, 2010; Husak et al., 2015), but previous

studies assume that compensatory traits still function as "cost-reducing" traits regardless of the situation (e.g., see Husak et al., 2015, p. 18). Here, I showed that "overcompensation" could decrease performance, producing a concave cost function, which is not surprising as individual performance is determined not by single traits but by the whole phenotype. In the current toy model, extended wings as a compensatory trait worsened the whole-body performance measured as stability, once the part of the elongated tail was removed. As an old Chinese proverb say, "too much is as bad as too little" at least in some cases. The assumption that the cost function can reveal the relative importance of sexual and viability selection (e.g., Buchanan & Evans, 2000) is therefore unreliable. In other words, the prediction of sexual selection (i.e., Fig. 1C) is no longer valid and can be replaced by a concave cost function (as in Fig. 1B) at least under the presence of compensatory traits, indicating that the presence/absence of viability selection is undetectable (see Fig. S2 for the corrected prediction).

Of course, the current experiment using chuonchuon models does not mean that sexual selection always produces a concave cost function (and does not mean that the same physical property, i.e., stability, determined the evolution of ornamentation, such as swallows' tail, and compensatory traits). The actual compensatory traits would include multiple traits (see Introduction), and hence the current balancing toy model should be regarded as a "basic" model possessing single compensatory trait (or, more accurately, a set of compensatory traits, since extended wings in fact alter multiple dimensions of the wing morphology). Still, even without mathematical model (see results), it seems likely that overcompensation will often reduce locomotor performance, particularly when locomotion requires co-opted, integrated phenotype (e.g., aerial locomotion). In fact, concerning the

function of long tails in barn swallows, Norberg (1994) mentioned that "*any experimental shortening and lengthening of the outer tail feathers is likely to upset an original co-adapted character set...*" (p. 231). When compensatory traits co-evolved with the sexually selected tail length to minimize the aerodynamic cost of tail length (Møller, 1996), an experimental reduction in tail length alone cannot clarify the aerodynamic cost of tail length, because the manipulation did not follow the actual evolutionary pathway (see Fig. 3; also see Fig. S3 for an alternative scenario in which ornament has some aerodynamic function). Also, all macroevolutionary studies of these aerial insectivores so far have supported sexual (rather than viability) selection explanation (e.g., Hasegawa et al., 2016; Hasegawa & Arai, 2017, 2018, 2020a,b, 2021, 2022), demanding further experimental studies, possibly using a smaller amount of manipulation (so that overcompensation can be avoided) in many species with various ornamentation (i.e., phylogenetic comparative "experiment"; Fig. S4). An alternative approach, the manipulation of compensatory traits (sensu Møller, 1996; see also Fig. 3) would in theory be valuable to show the effect of overcompensation, but is impractical due to the multi-dimensionality of compensatory traits (e.g., manipulations of wing size and shape are often difficult to conduct).

In addition, the current experiment provides useful insights. First, the cost of ornamentation can be transformed from one type to another. Extended wings enhanced the stability of chuonchuon with elongated tails, though they required additional investment (i.e., a production cost) instead (also see Møller, 1996 for a similar argument). Furthermore, the production cost of compensatory traits can be much higher than the production cost of the ornament itself (see Methods). It is straightforward to conclude that a single measure of cost (e.g., aerodynamic cost) is inappropriate to

approximate the total cost (and thus viability disadvantage) of

ornamentation, even when the production cost of ornamentation itself seems

negligible at first glance, as is the case for the outermost tail feathers in barn

swallows (Fig. 1). Although we discussed here about a simple compensatory

trait (i.e., extended wings) with a production cost, alternative, but not

mutually exclusive, kinds of costs can be involved: for example, by changing

the position of pivot point via the change of the body shape in the toy model

(i.e., possible developmental cost). Such morphological changes will

accompany several physiological and behavioral changes (and thus may also

require some form of maintenance cost) in living animals. We therefore

should keep in mind that any, or at least many, kinds of costs can be involved

to compensate costly ornaments, and that the considerations of all these

costs might be impractical.

In summary, the current model experiment demonstrated that a

concave cost function, which is often thought to result from viability

advantage of focal ornaments, can be observed when sexual selection has

favored the evolution of the costly ornaments, due to the presence of

compensatory traits. Given that compensatory traits coevolved with sexual

traits (Møller, 1996), the manipulation of sexual traits alone would not

clarify the evolutionary force on sexual traits (see Figs. 3 & S3). Future

studies should consider that overcompensation can be detrimental and that

more sophisticated experimental design is needed when inferring selection

pressure on (and the evolutionary history of) ornamentation.

Acknowledgments

I thank Dr. Emi Arai, Dr. Shumpei Kitamura and his lab members at

Ishikawa Prefectural University for their kindest advices and supports. I

also thank Prof. Michael D. Jennions, reviewers, and English-proofing

company (Enago) for their helpful comments. MH was supported by the

KAKENHI grant of the Japan Society for the Promotion of Science (JSPS,

19K06850). I am grateful to valuable comments by Dr Alex McQueen and

anonymous reviewers, which greatly improve the quality of the manuscript.

Data Availability Statement

Data attached as Table S1 will be deposited into osf.io once accepted.

Compliance with ethical standards

Conflict if interests   The author declares no conflict of interest.

Ethical approval   This study does not include any treatments of

animals, as all the information was gathered from model experiments.

Author Contribution

I performed data analysis and wrote the manuscript alone.

References

Andersson, M. (1994). Sexual selection. − Princeton Univ. Press, Princeton.

Arnold, S.J. (2003). Performance surfaces and adaptive landscapes. − Integr.

Comp. Biol. 43:367–375. https://doi.org/10.1093/icb/43.3.367

Arnold, S.J., Pfrender, M.E. & Jones, A.G. (2001). The adaptive landscape as

a conceptual bridge between micro- and macroevolution. − Genetica

112–113: 9–32. https://doi.org/10.1023/A:1013373907708

Barbosa, A. & Møller, A.P. (1999). Aerodynamic costs of long tails in male

barn swallows *Hirundo rustica* and the evolution of sexual size

dimorphism. − Behav. Ecol. 10: 128–135.

https://doi.org/10.1093/beheco/10.2.128

Balmford, A., Jones, I.L. & Thomas, A.L.R. (1994). How to compensate for

costly sexually selected tails: the origin of sexually dimorphic wings in

long-tailed birds. −Evolution 48: 1062−1070.

https://doi.org/10.2307/2410366

Baumgartner, A., Coleman, S. & Swanson, B. (2011). The cost of the sword:

escape performance in male swordtails. − PLoS ONE 6: e15837.

https://doi.org/10.1371/journal.pone.0015837

Bergstrom, C.T. & Dugatkin, L.A. (2016). Evolution. −WW Norton & Co. Inc,

NY.

Bro-Jørgensen, J., Johnstone, R.A. & Evans, M.R. (2007). Uninformative

exaggeration of male sexual ornaments in barn swallows. − Curr. Biol.

17: 850−855. https://doi.org/10.1016/j.cub.2007.03.042

Buchanan, K.L. & Evans, M.R. (2000). The effect of tail streamer length on

aerodynamic performance in the barn swallow. −Behav. Ecol. 11:

228−238. https://doi.org/10.1093/beheco/11.2.228

Evans, M.R. (1998). Selection on swallow tail streamers. −Nature 394:

233−234. https://doi.org/10.1038/28297

Evans, M.R. & Thomas, A.L.R. (1997). Testing the functional significance of

tail streamers. −Proc. R. Soc. Lond. B 264: 211−217.

https://doi.org/10.1098/rspb.1997.0030

Fjeldsa, J., Christidis, L., Ericson, P.G.P. (2022). The largest avian radiation:

the evolution of perching birds, or the order Passeriformes. −Lynx

edicions, Barcelona.

Hadfield, J.D. (2010). MCMC methods for multi-response generalized linear

mixed models: The MCMCglmm R package. −J. Stat. Soft. 33: 1−22.

https://doi.org/10.18637/ jss.v033.i02

Hasegawa, M., Arai, E. & Kutsukake, N. (2016). Evolution of tail fork depth

in genus *Hirundo*. − Ecol. Evol. 6: 851–858.

https://doi.org/10.1002/ece3.1949

Hasegawa, M. & Arai, E. (2017). Egg size decreases with increasing female

tail fork depth in family Hirundinidae. − Evol. Ecol. 31: 559–569.

https://doi.org/10.1007/s1068 2-017-9895-2

Hasegawa, M. & Arai, E. (2018). Convergent evolution of the tradeoff

between egg size and tail fork depth in swallows and swifts. − J. Avian

Biol. 49: e01684. https://doi.org/10.1111/jav.01684

Hasegawa, M. & Arai, E. (2020a). Fork tails evolved differently in swallows

and swifts. J. Evol. Biol. 33: 911–919. https://doi.org/10.1111/jeb.13622

Hasegawa, M. & Arai, E. (2020b). Correlated evolution of biparental

incubation and sexual tail monomorphism in swallows and martins

(Aves: Hirundinidae). − Evol. Ecol. 34: 777–788.

https://doi.org/10.1007/s10682-020-10066-w

Hasegawa, M. & Arai, E. (2021). Opposing population trends of fork-tailed

swallows and reddish-coloured swallows in our changing world. − J. Evol.

Biol. 34: 331–338. https://doi.org/10.1111/jeb.13736

Hasegawa, M. & Arai, E. (2022). Extrapair mating opportunity explains the

evolutionary transition between the presence and absence of tail

ornamentation in swallows. −Evol. Ecol. 36: 899–913. doi:

10.1007/210682-022-10196-3

Hill, G.E. & McGraw, K.J. (2006). Bird coloration II: function and evolution.

— Harvard Univ. Press, Cambridge.

Husak, J.F. & Swallow, J.G. (2011). Compensatory traits and the evolution of

male ornaments. − Behaviour 148: 1–29.

https://doi.org/10.1163/000579510X541265

Husak, J.F., Henningsen, J.P., Vanhooydonck, B. & Irschick, D.J. (2015). A

performance-based approach to studying costs of reliable signals. – In:

Animal signaling and function: an integrative approach (Irschick, D.J.

Briffa, M. & Podos, J., eds). Hoboken: Wiley Blackwell, pp. 47–74.

Kass, R.E., Carlin, B.P., Gelman, A. & Neal, R. (1998). Markov chain Monte

Carlo in practice: A round table discussion. – Amer. Stat. 52: 93–100.

https://doi.org/10.2307/2685466

Kojima, W. & Lin, C.P. (2019). Sprint speed is not reduced by exaggerated

male weapons in a flower beetle *Dicronocephalus wallichii*. – Ethology

125: 47–56. https://doi.org/10.1111/eth.12824

Lande, R. & Arnold, S.J. (1983). The measurement of selection on correlated

characters. – Evolution 37:1210–1226. https://doi.org/10.2307/2408842

Lessells, C.M. & Boag, P.T. (1987). Unrepeatable repeatabilities: A common

mistake. – Auk 104: 116–121. https://doi.org/10.2307/4087240

Matyjasiak, P., Marzal, A., Navarro, C., de Lope, F. & Møller, A.P. (2009).

Fine morphology of experimental tail streamers and flight

manoeuvrability in the house martin *Delichon urbica*. – Funct. Ecol. 23:

389–396. https://doi.org/10.1111/j.1365-2435.2008.01490.x

Moreno, E. & Møller, A.P. (1996). Morphological aspects of avian tail

movements: a functional approach in hirundines. – Auk 113: 647–654.

https://doi.org/10.2307/4088985

Møller, A.P. (1988). Female choice selects for male sexual tail ornaments in

the monogamous swallow. Nature 332: 640–642.

https://doi.org/10.1038/332640a0

Møller, A.P. (1994). Sexual selection and the barn swallow. – Oxford Univ.

Press, Oxford.

Møller, A.P. (1996). The cost of secondary sexual characters and the evolution

of cost-reducing traits. – Ibis 138: 112–119.

https://doi.org/10.1111/j.1474-919X.1996.tb04317.x

Møller, A.P., de Lope, F. & Saino, N. (1995). Sexual selection in the barn

swallow Hirundo rustica. VI. Aerodynamic adaptations. – J. Evol. Biol.

8: 671–687. https://doi.org/10.1046/j.1420-9101.1995.8060671.x

Norberg, R. (1994). Swallow tail streamer is a mechanical device for

selfdeflection of tail leading edge enhancing aerodynamic efficiency and

flight manoeuvrability. – Proc. R. Soc. Lond. B 257: 227–233.

https://doi.org/10.1098/rspb.1994.0119

Oufiero, C.E. & Garland, Jr T. (2007). Evaluating performance costs of

sexually selected traits. – Funct. Ecol. 21: 676–689.

https://doi.org/10.1111/j.1365-2435.2007.01259.x

R Core Team. (2021). R: A language and environment for statistical

computing. – R Foundation for Statistical Computing, Vienna, Austria.

Available at http://www.R-project.org/ (accessed 11 September 2021)

Romano, A., Constanzo, A., Rubolini, D., Saino, N. & Møller, A.P. (2017).

Geographical and seasonal variation in the intensity of sexual selection

in the barn swallow *Hirundo rustica*: A metaanalysis. – Biol. Rev. 92:

1582–1600. https://doi.org/10.1111/brv.12297

Rowe, L.V., Evans, M.R. & Buchanan, K.L. (2001). The function and

evolution of the tail streamer in hirundines. – Behav. Ecol. 12: 157–163.

https://doi.org/10.1093/beheco/12.2.157

Saino, N., Cuervo, J.J., Ninni, P., de Lope, F. & Møller, A.P. (1997).

Haematocrit correlates with tail ornament size in three populations of

barn swallow (*Hirundo rustica*). – Funct. Ecol. 11: 604–610.

https://doi.org/10.1046/j.1365-2435.1997.00131.x

Stinchcombe, J.R., Agrawal, A.F., Hohenlohe, P.A., Arnold, S.J. & Blows,

M.W. (2008). Estimating nonlinear selection gradients using quadratic

regression coefficients: double or nothing? – Evolution 62: 2435–2440.

https://doi.org/10.1111/j.1558-5646.2008.00449.x

Thavarajah, N.K., Tickle, P.G., Nudds, R.L. & Codd, J.R. (2016). The peacock

train does not handicap cursorial locomotor performance. – Sci. Rep. 6:

36512. https://doi.org/10.1038/srep36512

Thomas, A.L.R. & Rowe, L.V. (1997). Experimental tests on tail elongation

and sexual selection in swallows (*Hirundo rustica*) do not affect the tail

streamer and cannot test its function. – Behav. Ecol. 8: 580–581.

https://doi.org/10.1093/beheco/8.5.580

Trappett, A., Condon, C.H., White, C., Matthew, P. & Wilson, R.S. (2013).

Extravagant ornaments of male threadfin rainbowfish (*Iriatherina*

werneri) are not costly for swimming. – Funct. Ecol. 27: 1034–1041.

https://doi.org/10.1111/1365-2435.12097

Tubaro, P.L. (2003). A comparative study of aerodynamic function and

flexural stiffness of outer tail feathers in birds. – J. Avian Biol. 34:

243–250. https://doi.org/10.1034/j.1600-048X.2003.03084.x

Turner, A.K. (2006). The barn swallow.– T & AD Poyser, London.

Table 1.   Bayesian linear mixed-effects model of stability of chuonchuon

in relation to experimental manipulation of elongated tails (N_{ID} = 4, $Ntotal$ =

20).

Variable	Coefficient [95% CI]	P_{MCMC}
Intercept	**22.37 [18.95, 25.97]**	**<0.0001**
Linear term of manipulation	**0.96 [0.71, 1.23]**	**<0.0001**
Quadratic term of manipulation	**8.36 [6.21, 10.51]**	**<0.0001**

560 The dependent variable is the absolute angle of chuonchuon deviated from

561 the horizontal plane. I included chuonchuon ID as a random factor (variance

562 = 0.20) in addition to units (variance = 15.39).

563 Significant test results (i.e. $P < 0.05$) are indicated in bold.

565 Figure legends

Type of selection	A Viability selection	B Viability selection +Sexual selection	C Sexual selection
Predicted function			

Deviation of outermost tail feathers | Deviation of outermost tail feathers | Deviation of outermost tail feathers

Methods & Results

ca. -10mm

Authors tested prediction by shortening tail feather tips

The results matched the prediction of "B"

Conclusion — Sexual selection extended tail length by around 10mm beyond its aerodynamic optimum

Figure 1.   Details of manipulation experiment of Evans and colleagues (see Evans, 1998; Buchanan & Evans, 2000; Rowe et al., 2001 for original figures; also see Evans & Thomas, 1997; Thomas & Rowe, 1997 for similar predictions). The vertical line in each plot indicates current ornament expression, and the triangle indicates ornament expression providing peak performance (note that lower values indicate higher performance here). Although they originally called deviation of outermost tail feathers (or possibly, narrowed part of outermost tail feather length) as a "streamer," I avoided this wording, because it implies some kinds of aerodynamic function of the ornaments, and because the formal definition of the streamer is unclear (and not to confound size and shape: Matyjasiak et al., 2009). Likewise, I used viability selection rather than natural selection to be precise.

Figure 2.   Chuonchuon (upper panel), chuonchuon with elongated tails
and extended wings (middle panel), showing how manipulation was
conducted, and results of chuonchuon experiment (lower panel). A simple
quadratic regression line is denoted (see Table 1 for a formal analysis).
Individual identity was denoted by different coloration. Mean ± SE values of
stability, measured by absolute angle deviated from horizontal plane, in
ancestral state shown in upper panel was 4.87 ± 1.78 (range: 1.38–8.87). Not
surprisingly, tail elongation without extended wings (i.e., models without
compensatory traits) made models unbalanced in the range of the current
study (e.g., when elongated tails = 30 mm, which corresponds to the
manipulation = −80 mm, mean stability ± SE = 36.60 ± 2.30, range =
32.69–42.93).

Figure 3. An example of hypothetical performance surface in relation to ornament and phenotypic expression of compensatory traits (upper panel). Darker background coloration indicates lower performance (note that the best performance is attained in the lower left corner). Black arrow indicates hypothetical evolutionary pathway to the current state (filled circle). Light blue arrow indicates phenotypic change when researchers experimentally reduced ornamentation (in which compensatory traits remain unchanged). Lower panel shows performance cost measurement (which is here denoted as the best performance minus the performance of the focal coordinate) in relation to ornament expression along with black or light blue arrows. The evolutionary pathway to the current state is biased toward right side because of intense sexual selection favoring long tails (i.e., actual evolutionary pathway depends on sexual selection in addition to viability selection due to performance cost and other costs, such as production costs of ornament and compensatory traits; see text). Narrow contour width

indicates that small deviation from integrated, co-opted trait sets had a strong negative effect on performance, as predicted by Norberg (1994). Here, for the illustrative purpose, I assumed that performance is determined by $-4x^2 - 4y^2 + 7xy$ where x denotes ornament expression whereas y denotes the expression of compensatory traits. The current state is determined here as the point in which the sum of performance surface and sexual selection surface (here, $10x$) is maximized (see text for detailed explanation). Dotted line is shown here as a ridge line.

Figure S1. An example of performance surface with a saddle (left panel) and

an example of performance surface with a ridge (right panel), both of which

are given by performance surface = $0.5*\gamma_{11}*x_1^2 + \beta_1*x_1 + 0.5*\gamma_{22}*x_2^2 + \beta_2*x_2 +$

$\gamma_{12}*x_1*x_2 + \alpha$ with the condition, $\gamma_{12}^2 > \gamma_{11}*\gamma_{22}$ (former) and $\gamma_{12}^2 = \gamma_{11}*\gamma_{22}$ (latter).

Darker background coloration indicates lower performance.

Figure S2. The "corrected" predictions for manipulation experiment of Evans

and colleagues under the presence of compensatory traits. Note that two type

of scenarios (B and C) cannot be distinguished. See text and Fig. 1 for

detailed explanations (also see Figs. 3 & S3).

Figure S3. An example of hypothetical performance surface in relation to ornament and phenotypic expression of compensatory traits (upper panel). Darker background coloration indicates lower performance (note that the lightest background located near the current state, shown as a filled circle, in this example). Black arrow indicates hypothetical evolutionary pathway to the current state. Light blue arrow indicates phenotypic change when researchers experimentally reduced ornament expression (in which compensatory traits remain unchanged). Lower panel shows performance cost measurement (which is here denoted as the best performance minus the performance of the focal coordinate) in relation to ornament expression along with black or light blue arrows. The evolutionary pathway to the current state is biased toward right side because of intense sexual selection favoring long tails (i.e., actual evolutionary pathway depends on sexual selection in addition to viability selection due to flight cost and other costs, such as production costs of tail length and compensatory traits; see text). Narrow contour width indicates that small deviation from integrated, co-opted character sets had a strong negative effect on flight performance, as predicted by Norberg (1994). Here, for the illustrative purpose, I assumed that performance is determined by $-4(x-3)^2 - 4(y-3)^2 + 7(x-3)*(y-3)$ where x denotes ornament expression whereas y denotes the expression of compensatory traits. The current state is determined here as the point in which the sum of performance surface and sexual selection surface (here, $5x$) is maximized (see text for detailed explanation).

(See next page for Figure S3)

Figure S4. A phylogenetic comparative experiment. Rather than reducing a
large extent of ornamentation in a single species, which can result in
overcompensation (light blue arrow), researchers should reduce small
extents of ornamentation in multiple species with varying expressions of
ornamentation, without being bothered by overcompensation (dark green
arrow). **Black arrow indicates a hypothetical evolutionary pathway to the**
current state of the most ornamented species. Less-ornamented species are
assumed to be present somewhere near the evolutionary pathway (i.e., they
are here assumed to be hypothetical ancestral states of the most ornamented
species; sensu Matyjasiak et al. 2009). See Fig. 3 for detailed information
(note that this figure is derived from the upper panel in Fig. 3). For easy
understanding, dotted line is shown here as a ridge line, above which
overcompensation occurs.