Testosterone eliminates strategic prosocial behavior in healthy males

Hana H. Kutlikova*[a,b]
Lei Zhang[a,c,d]
Christoph Eisenegger [a] [†]
Jack van Honk[b,e]
Claus Lamm[a,f]

[a] Department of Cognition, Emotion, and Methods in Psychology, University of Vienna, Vienna, Austria
[b] Department of Experimental Psychology, Helmholtz Institute, Utrecht University, Utrecht, the Netherlands
[c] Centre for Human Brain Health, School of Psychology, University of Birmingham, Birmingham, UK
[d] Institute for Mental Health, School of Psychology, University of Birmingham, Birmingham, UK
[e] Department of Psychiatry and Mental Health, University of Cape Town, Cape Town, South Africa
[f] Vienna Cognitive Science Hub, University of Vienna, Vienna, Austria

[†] Deceased February 27, 2017

*Corresponding author: Hana H. Kutlikova, Liebiggasse 5, University of Vienna, Vienna 1010, Austria, hana.kutlikova@univie.ac.at

Author Contributions: H.H.K. and L.Z. share the first authorship of this article. H.H.K, C.E., and C.L. designed research; H.H.K. performed research; H.H.K. and L.Z. analyzed data; H.H.K., L.Z., C.L., and J.v.H. wrote the paper.

Competing Interest Statement: The authors declare no competing interest.

Abstract

Humans are strategically more prosocial when their actions are being watched by others than when

they act alone. Using a psychopharmacogenetic approach, we investigated the computational and

endocrinological mechanisms of such audience-driven prosociality. 187 male participants received

either a single dose of testosterone or a placebo and performed a prosocial and self-benefitting

reinforcement learning task. Crucially, the task was performed either in private or when being

watched. Rival theories suggest that the hormone might either diminish or strengthen audience-

dependent prosociality. We show that exogenous testosterone fully eliminated strategic, i.e., feigned,

prosociality and thus decreased submission to audience expectations. We next performed

reinforcement-learning drift-diffusion computational modeling to elucidate which latent aspects of

decision-making testosterone acted on. The modeling revealed that testosterone compared to placebo

did not deteriorate reinforcement learning per se. Rather, when being watched, the hormone

altered the degree to which the learned information on choice value translated to action selection.

These results indicate that when being watched, the hormone was influencing participants towards

choices that conform less with audience expectations, and thus lower prosociality. Exploratory

analysis of personal values related to conformity and tradition corroborates this view, as it suggests

that the shift from socially conforming to nonconforming decisions co-varies with individual

differences in these values. Together, these findings indicate that instead of deceptively increasing

socially desirable behavior to manage reputation and status, testosterone, by acting on implicit

decision-making processes, counteracts conformity and submission to audience expectations.

Keywords: testosterone, prosocial behavior, audience-effect, reinforcement learning, drift-diffusion

model, conformity

Introduction

Humans behave more prosocially when their actions are watched by others[1]. This phenomenon has been demonstrated across a variety of social behaviors, such as blood donations[2], church offerings[3], or monetary donations to charitable organizations[4], and is often referred to as strategic prosociality[5], or the audience effect[6]. From an evolutionary perspective, making one's generosity visible to others has an important signaling value, in that it advertises an individual's qualities as a potential partner or a valuable group member[7]. In the present study, we propose and investigate whether the steroid hormone testosterone plays a crucial role in shaping such audience effects.

Research in the past decade has demonstrated that testosterone is implicated in a wide spectrum of socially dominant behaviors[8,9]. Exogenous testosterone alleviates subordination to the dominance of others[10-12] and reduces the physiological response to being evaluated by others[13]. Given that enhanced submission to audience expectations has been associated with intense apprehension about social evaluation[14], one possible prediction is that testosterone administration will decrease audience effects.

Contrasting with this view, the hypothesis that testosterone drives status-seeking via reputation building rather than dominance[15,16] would predict that based on the social context, testosterone might conditionally promote prosocial and especially socially desirable behavior to build up a reputation and increase status.

The present paper is the first that aimed to distinguish between these two alternatives of boosting one's social status that testosterone may act on. One option is that, in line with the social dominance hypothesis[17], the hormone prioritizes dominant status-seeking and would hence diminish the submission to audience expectations. The other option is that testosterone primarily promotes reputable status-seeking[15,16]. If true, the hormone could increase strategic prosocial behavior.

Through what neurobiological pathways could testosterone modulate such complex social behaviors? Previously, exogenous testosterone was found to increase dopamine levels in the rat ventral striatum[18], suggesting that the hormone exerts its effects through the modulation of dopaminergic activity in reward-related neural circuits. Besides this insight from animal research, testosterone and reward processing have also been linked in humans[19,20]. It remains to be shown, though, which specific aspects of reward processing testosterone acts on. For one, during value learning, testosterone may influence the incorporation of so-called prediction errors (PE), which track the difference between predicted and actual outcome[21] and are encoded by the phasic activity of midbrain dopaminergic neurons projecting to the ventral striatum[22,23]. Alternatively, testosterone may impact the conversion of the learned values into action selection, or the temporal dynamics of the evidence accumulation.

The present study thus not only aimed to investigate if testosterone influences strategic prosociality,

but also whether this is achieved by impacting reward-related computations. We employed a novel

modeling approach, by combining reinforcement learning with diffusion decision models (RLDDM).

This provided a more comprehensive account of the latent processes involved in prosocial decision-

making than previous separate RL and DDM approaches[21,24]. Besides describing how subjective

values of the choice options are learned through PEs (*learning rate parameters*) and converted to

actions (*choice consistency parameter*), the new combination of reinforcement learning and diffusion

decision modeling also enabled us to explore the temporal dynamics of these latent processes

(*decision threshold and drift-scaling parameters,* see *SI Table S3* for parameter description)[24].

Male participants (n = 187) underwent a double-blind, between-subject, placebo-controlled,

topical testosterone (150 mg) administration and then performed a reinforcement learning task (Figure

1). On each trial, participants were required to choose between one of two symbols. One symbol was

associated with a high probability (75%), and one was associated with a low probability (25%) of a

reward. These contingencies were not instructed but had to be learned through trial and error. While

charitable donation tasks[16] and neuroeconomic games[8,9] classically measure participants' overall

prosociality using deliberated decisions, such as deciding how much money to share with another

person, the RL task allowed us to furthermore characterize the hidden individual steps in the process

of learning about the consequences actions have for oneself and others (see[25,26] for similar recent

approaches). To compare self- and other-oriented decision-making, participants completed the task for

themselves and for an NGO of their choice (within-subject condition). Critically, the task was

performed either in private or when being watched (between-subject condition, see *Materials and*

Methods).

Based on previous audience-effect research[2-6], we predicted that when the participants are

watched, they will be relatively more prosocial (i.e., make more correct choices for the other vs self)

than in private. Crucially, we expected that such an audience effect will be underpinned by relatively

faster incorporation of the PEs (captured by the learning rate parameter α in RL); higher consistency

in converting values to action probability (captured by the inverse temperature parameter tau in RL,

also known as value sensitivity, exploration parameter, or 1/β); and more integrated evidence

necessary for making a decision (captured by the threshold parameter in DDM). In other words,

participants would learn more efficiently, learned values would inform their behavior more

consistently, and their decisions would be more cautious.

Our main hypothesis was that the effects of being watched on other- vs. self-benefitting

behavior will be modulated by testosterone administration. Given that testosterone reduces

submission signals and stress responses to the social evaluation, allowing for dominant status-

seeking[10-13], we hypothesized that testosterone would reduce the audience effect expected in the

placebo group. As an alternative prediction, we reasoned that if testosterone does not primarily cause

dominant status-seeking, but instead, in non-threatening environments, promotes more agreeable and

reputable status-enhancing behaviors[16,17], participants in the testosterone (vs placebo) group should

show a larger audience effect. Irrespective of whether testosterone would increase or decrease

prosocial behavior under the audience effect, we also predicted that testosterone's effects will be

associated with changes in the efficiency of PE-based value updating (α in RL), choice consistency

(tau in RL), and evidence necessary for making a decision (threshold parameter in DDM).

Furthermore, considering that testosterone possibly modulates social behavior through both

androgenic and dopaminergic pathways[27], we explored whether a CAG repeat polymorphism of the

androgen receptor, as well as a DAT1 polymorphism of the dopamine transporter, interacts with

testosterone administration effects. Next, as previous research has shown that testosterone effects on

status-seeking and decision-making are influenced by endogenous cortisol levels[28,29], we examined

whether testosterone administration effects interact with salivary cortisol levels measured at baseline

and with cortisol reactivity to being watched. Moreover, as it has been suggested that sensitivity of

dopaminergic pathways is heightened among highly dominant individuals[30] and that these individuals

show more pronounced effects of testosterone administration[27,31], we tested whether testosterone

effects on strategic prosociality vary as a function of self-reported trait dominance[32]. Lastly, we

conducted an exploratory post-hoc analysis to shed light on self-reported motivational value

systems[33,34] putatively underlying testosterone's effects on strategic prosociality.

Figure 1. Experimental design and task. *(A)* Timeline of the experimental session. *(B)* Prosocial reinforcement
learning task. Participants performed the task either in private or watched by an observer introduced as an NGO
association representative. The observation was signaled by a red frame. Each participant completed three
blocks of 16 trials for themself and three blocks of 16 trials to benefit an NGO of his choice. *(C)* Schematic of
the reinforcement learning drift diffusion model (RLDDM). Left panel: trial-by-trial value updates in RL; right
panel: evidence accumulation in DDM. Importantly, the drift rate in DDM is calculated from the value
difference between choice options in RL.

Results

Manipulation check

In the testosterone, compared to the placebo group, we observed higher salivary testosterone levels
110 minutes after gel administration, and this difference remained stable until the end of the
experiment (drug treatment x time: $F(3, 522.59) = 47.82$, $p < .001$, $\eta^2 p = .423$, see *Figure S1* and *SI:*
Supplementary analysis of hormone data for analysis details). The observation condition did not
significantly influence testosterone levels (visibility x time: $F(3,520.88) = 1.528$, $p = .206$).

Analysis of the subjective ratings of being watched showed that participants in the observed condition
felt watched to a greater extent than in the private condition ($X^2 (2, 187) = 111.77$, $p < .001$), and that
testosterone administration did not influence the perception of being watched ($X^2 (2, 187) = 0.007$, p
$= .997$). The degree to which the participants felt watched, was positively associated $r (179) = .173$, p
$= .019$) with cortisol reactivity to the visibility manipulation (Δ cortisol levels 20 minutes after the end
of the task - cortisol levels immediately before the task). Note also that the feelings of being watched
were not related to the cortisol levels measured at the baseline $r (179) = -.082$, $p = .317$).

Testosterone eliminates the audience effect

The three-way interaction of the factors drug treatment (P/T), visibility (private/observed), and type of
recipient (self/other)predicted the number of correct choices (i.e., options that have higher reward
probability) the participants made ($OR = 1.06$, $CI = [1.00, 1.13]$, $p = .046$; Figure 2A). Follow-up
analysis using treatment contrasts showed that participants in the placebo group showed more
prosocial behavior, as indicated by relatively more correct prosocial choices when being watched
compared to the private setting in which they were not watched (recipient x visibility interaction in
the placebo group: $OR = 1.44$, $CI = [1.02, 2.02]$, $p = .039$). Supporting our prediction based on the
social dominance hypothesis, this audience effect was absent in the testosterone group (recipient x
visibility interaction in the testosterone group: $OR = 0.88$, $CI = [0.63, 1.23]$, $p = .461$). Specifically,
when participants were observed, testosterone, compared to placebo, reduced the number of correct
choices made for another ($OR = 0.67$, $CI = [0.49, 0.91]$, $p = .011$, Figure 2C). The number of correct
choices made for self, however, was not influenced by the drug treatment ($OR = 1.12$, $CI = [0.87$,
$1.45]$, $p = .384$), visibility ($OR = 0.98$, $CI = [0.75, 1.27]$, $p = .861$, or their interaction ($OR = 0.91$, CI
$= [0.63, 1.33]$, $p = .638$, (Figure 2B).

Figure 2. Differences in the number of correct choices. *(A)* Three-way interaction of the factors treatment x
recipient x visibility. Participants in the placebo group behaved more prosocially (as captured by the prosociality
index = correct choices for other – correct choices for self) when being observed than in privacy. Exogenous
testosterone eliminated this audience effect. Note that the analyses were performed with raw data in the 2 x 2 x 2
factorial design; the plotted difference score (other minus self) is to improve readability and interpretability.
Breaking down the three-way interaction by using treatment contrasts showed that there was no significant
effect of the treatment and visibility factors, or their interaction, on the number of correct choices made for
oneself *(B)*. On the other hand, testosterone, compared to placebo, decreased the number of correct choices
made for the NGO when being observed *(C)*. Dots represent the data of individual participants, lines represent
mean values per group, and boxes 95% confidence intervals.

Behavior is best explained by a reinforcement learning drift diffusion model with dual learning
rates

Next, we sought to uncover the computational mechanisms underlying the experiment-condition-
specific behavioral differences on a trial-by-trial basis. The winning model (winning over five other
candidate models; see *Materials and Methods* and *SI: Model selection and validation*) entailed
combined RL and DDM components, and thus simultaneously predicted individuals' choices and RTs
(see *SI: Table S3* for a complete list of parameters and their description). The RL component section
predicted participants' learning behavior via the value updates through the computation of PEs with
separate learning rates for positive and negative PEs (i.e., α^{posPE} and α^{negPE} Equation (1)). In other
words, the model that best accounted for the data assumed a differential speed of learning with and
without positive feedback:

242
$$V_{c,t} = \begin{cases} V_{c,t-1} + \alpha^{pos}(O_{t-1} - V_{c,t-1}), \text{if } O_{t-1} > 0 \\ V_{c,t-1} + \alpha^{neg}(O_{t-1} - V_{c,t-1}), \text{otherwise} \end{cases} \quad (1)$$

where O_{t-1} denotes the outcome, and $V_{c,t-1}$ the subjective value of choice c at trial $t-1$.

In addition, the DDM component predicted RTs by assuming an evidence accumulation process (as

quantified by the drift rate; decisions were made when the evidence reached a certain threshold[35]).

Importantly, the marriage between RL and DDM allowed a fine-grained investigation into how the

drift rate (v_t) was shaped by the value difference between two symbols at the trial-by-trial level

(Equation (2); *S*, a non-linear transformation function; $v_{scaling}$, a weight parameter that maps accuracy-

coded value difference into the drift rate[24]).

250
$$v_t = S\left[v_{\text{scaling}}\left(V_{\text{correct},t} - V_{\text{incorrect},t}\right)\right]_{,} \tag{2}$$

251 We fitted all candidate models (see *Materials and Methods* and *SI: Supplementary information on*

252 *computational modeling*) under the hierarchical Bayesian estimation scheme[36] to incorporate both

253 group-level commonality and individual differences, according to our task design (effects of drug

254 treatment (P/T), visibility (private/observed), and type of recipient (self/other)).

Testosterone's impact on strategic prosocial behavior is associated with choice consistency

Next, we investigated which RLDDM parameters of our validated winning model are associated with

the effects found in the behavioral analysis of the correct choice. As a first step, we tested the

parameters for the 3-way interaction effect of drug treatment, visibility, and type of recipient.

In the second step, we examined whether the parameters that showed a three-way interaction effect of

our experimental manipulation predict behavioral prosociality. Behavioral prosociality was measured

by the difference between correct choices made for others and self.

Out of the five parameters (learning rate for positive PE, learning rate for negative PE, choice

consistency, threshold, drift-scaling parameter), only choice consistency showed the requisite three-

way interaction of our experimental manipulations ($B = 0.98$, $CI = [0.97, 0.99]$, $p < .001$), and at the

same time significantly predicted behavioral prosociality (Bonferroni correction for multiple

comparisons, $B = 5.45$, $CI = [3.71, 7.19]$, $p < .001$; Figure 3D). Specifically, participants in the

placebo group displayed relatively higher consistency in choices made for the other (vs. self) when

being observed than in privacy (recipient x visibility interaction in the placebo group: $B = 1.08$, $CI =$

$[1.05, 1.10]$, $p < .001$). On the contrary, in the testosterone group, observation, compared to privacy,

decreased the consistency of choices made for the other (vs. self) (recipient x visibility interaction in

testosterone group: $B = 0.94$, $CI = [0.92, 0.97]$, $p < .001$; Figure 3B). When participants were

observed, testosterone, compared to placebo, diminished the relative consistency of prosocial choices

(recipient x treatment interaction in observed condition: $OR = 0.87$, $CI = [0.85, 0.89]$, $p = .001$). In the

private condition, there was no evidence for such an effect (recipient x treatment interaction in private

condition: $OR = 0.99$, $CI = [0.97, 1.02]$, $p = .605$; for analysis of all RLDDM parameters, see *SI:*

Analysis of the RLDDM parameters and their association with prosocial behavior).

Altogether, these results suggest that testosterone eliminates audience-dependent prosocial behavior

by affecting choice consistency.

Interaction of testosterone effects with trait dominance

In further support of the social dominance hypothesis, trait dominance interacted with testosterone's

effects on correct choice (recipient x drug treatment x visibility x trait dominance: $OR = 1.04$, $CI =$

[1.01, 1.09], $p = .026$). In a follow-up analysis aimed at decoding this interaction, the continuous

measure of dominance was replaced by a categorical variable with levels of high and low dominance,

based on the median split of dominance scores (Med = 3.875). Decomposition of the four-way

interaction revealed that testosterone reduced the number of correct choices made for others during

observation specifically among men with high trait dominance ($OR = 0.60$, $CI = [0.42, 0.87]$, p

$= .008$) and this effect was weaker and non-significant among those with low dominance ($OR = 0.74$,

$CI = [0.52, 1.04]$, $p = .084$). Trait dominance did not significantly interact with the RLDDM

parameters (all $ps > .220$ see *SI: Interaction of trait dominance with testosterone effects on RLDDM*

parameters).

Interaction of testosterone effects with genetic polymorphisms and cortisol levels

CAG-repeat and DAT1 polymorphisms did not interact with the effects of testosterone on correct

choice or RLDDM parameters (all $ps > .090$, see *SI: Supplementary information on the analysis of*

genetic data). Furthermore, neither baseline cortisol levels, nor cortisol reactivity to visibility

manipulation (Δ cortisol levels 20 minutes after the end of task - cortisol levels immediately before

the task) interacted with the effects of testosterone on correct choice or RLDDM parameters (all ps

$> .074$, see *SI: Supplementary analysis of hormone data*).

Interaction of testosterone effects with personal value orientations

We next explored whether four principal value orientations (self-enhancement, self-transcendence,

openness, conservation) [33,34] measured using a self-report questionnaire interacted with testosterone's

influence on choice consistency. This revealed a significant interaction with individual differences in

conservation value orientation, which reflects how someone identifies with values such as

conformity, security, tradition, and self-restriction (recipient x visibility x administration x

conservation: $B = 0.05$, $CI = [0.02, 0.09]$, $p = .001$). Concretely, testosterone (vs placebo)

administration significantly altered the link between the conservation value and audience effect on

choice consistency. Among placebo participants, the treatment contrast analysis showed a

nonsignificant positive association between conservation value and the prosocial audience effect

(recipient x visibility x conservation: $B = 0.17$, $CI = [-0.03, 0.36]$, $p = .088$). In the testosterone

group, the opposite pattern emerged, and higher conservation value was significantly associated with

lower audience effect (recipient x visibility x conservation: $B = -0.25$, $CI = [-0.42, -0.08]$, $p = .003$).

No other interaction between value orientations and testosterone effect on choice behavior or

RLDDM parameters was detected (all $ps > .101$ see *SI: Supplementary information on the*

questionnaire data).

Figure 3. Differences in the parameters estimated by the reinforcement learning drift diffusion model (RLDDM). *(A)* In the placebo group, observation compared to privacy relatively decreased the prosocial learning rate for negative PE (i.e., the difference between α^{negPE} in the other condition and α^{negPE} in the self condition). Testosterone administration reversed the observation effect. The results suggest that for better performance in the task, a lower learning rate from negative PE is more suitable. *(B)* In the placebo group, observation compared to privacy, relatively increased the consistency of the prosocial choices. Testosterone administration reversed this audience effect. *(C)* In the placebo group, observation compared to privacy, relatively increased the DDM threshold for prosocial choices. Testosterone administration reversed the audience effect. *(D)* Inverse temperature parameter tau that captures choice consistency significantly predicted prosociality. Dots represent the data of individual participants, lines represent mean values per group, and boxes 95% confidence intervals.

Learning parameters in relation to optimal learning rates

To gain a deeper understanding of how the learning parameters were related to the task performance in our experimental design, we performed a simulation study to identify optimal learning rates[37] (see *SI: Simulations of optimal learning rates*). In all conditions, both the posterior positive and negative

learning rates were smaller with respect to the optimal ones (see Figure 4A, 4C). Crucially, to validate

whether the choice accuracy corresponding to the posterior parameters in our winning model could

capture key patterns in our behavior findings (i.e., posterior predictive check), we let our winning

model generate synthetic data and analyzed the generated prosocial behavior (i.e., choice accuracy for

other minus choice accuracy for self) in the same way as we analyzed the observed data. We found

that results from the generated data (Figure 4B, 4D) greatly resembled the behavioral patterns

reported in Figure 2A.

Figure 4. Optimal learning rates and posterior predictive checks. Posterior learning rates in relation to the
optimal learning rates in the private *(A)* and observed *(C)* conditions. Orange dots represent the optimal
combination between learning rates for positive and negative PE identified via simulation; red crosses indicate
the posterior means of learning rates. The posterior learning rates were employed to perform posterior predictive
checks for the main behavioral findings for the private *(B)* and observed *(D)* conditions. Simulated data from
posteriors were analyzed in a similar fashion as the real data and the model prediction largely matched our main
behavioral effect (cf. Figure 2A).

Discussion

Using pharmacological manipulation and a novel computational model integrating reinforcement learning with the drift diffusion modeling framework (RLDDM), we tested and characterized testosterone's role in audience-dependent prosocial behavior. The results show that testosterone diminishes the typical audience effect present in the placebo condition. Computational modeling pinpoints this effect to a reduction in the extent to which the performance of prosocial (vs. selfish) choices is consistent with learned reward values. Moreover, the effects are more pronounced in participants with higher trait dominance. Taken together, these findings are in line with the social dominance hypothesis and are thus consistent with the notion that testosterone decreases submission to audience expectations, rather than promoting the strategic display of socially pleasing behavior[38,39].

A growing body of evidence suggests that testosterone exerts its behavioral effects through the modulation of reward-related processes[19,20]. However, to our knowledge, no study investigated the computational mechanisms underlying such effects. Using joint RLDDMs, we found that in the placebo group, observation (vs privacy) increased the relative consistency of prosocial choices. Testosterone administration eliminated this audience effect, making the performance of prosocial (vs. self-benefitting) choices less consistent with value computations. Low choice consistency means that individuals select options with non-maximal expected values, which is often referred to as exploratory behavior[40]. In environments with static reward probabilities, participants can maximize their reward by initially exploring which option tends to be more fruitful. Once learners discover the better option, exploration yields no benefit. One possible explanation of the present effect could therefore be that testosterone impaired individuals' ability to adapt and control the amount of exploration. However, our data do not indicate that testosterone affects exploration in general, as we did not find any testosterone influence on choice consistency in the private setting. Instead, the effects of testosterone appeared only in a situation where social status was at stake. Interestingly, research in monkeys has shown that socially challenging environments (vs socially isolated environments) increased the availability of D_2 receptors[41], which play a role in choice consistency[42]. This effect was present in particular in dominant monkeys and absent in the subordinate ones. Such findings are in line with our results showing enhanced testosterone effects in individuals with high trait dominance. We, therefore, suggest that future studies should examine the potentially mediating role of D2 receptors in testosterone's effect on reward learning in social situations. Alternatively, it could be speculated that the elimination of the audience effect by testosterone stems from the hormone's ability to reduce fear in social situations. Indeed, earlier research shows that exogenous testosterone diminishes the physiological stress response to the presence of an observer[13] and has anxiolytic-like properties in humans and across species[10,43,44]. However, we did not observe any interaction of the testosterone's effect with cortisol levels measured at baseline or with cortisol reactivity, thus our data do not provide

support for such an interpretation. To examine this topic further, studies on testosterone and audience

effect should include more explicit measures of subjectively perceived stress and anxiety.

l

The present data also shows that testosterone administration substantially alters the relation

between the audience effect and self-reported conservation value. Conservation value, as

measured here, reflects personal values with an emphasis on conformity, security, tradition, and

self-restriction[33]. The significant interaction between the degree to which one identifies with

conservation value and testosterone's influence on the audience effect was driven by a positive

association in the placebo group, and a reversal of this relationship in the testosterone group,

where higher conservation values went along with lower audience effects. In consideration of the

conformity aspect of the conservation value, it is of interest that in Western societies,

nonconforming people are perceived as having higher status and competence than those who

conform to the social expectations[45]. Thus, these findings are suggestive of the interpretation that

by inducing a more non-confirming attitude, testosterone reaches its principle goal - which is to

be observed as having high status and competence[17,39]. We note though that the analyses of value

orientation are exploratory and were performed post-hoc, and thus need independent

confirmation.

Variability in dominance, cultural differences in social status attaintment, and conservation

value can also account for the results of another recent study, which was conducted among Chinese

students and showed that testosterone enhanced audience effects[16]. Indeed, contrary to Western

society, in Eastern cultures, high social status is associated with increased self-restriction and other-

orientation, including generosity to those with lower status[46]. These cultural differences have been

linked to polymorphisms in the dopamine D4 receptor gene[47], implying a putative biological

mechanism that could explain cultural differences in testosterone effects. Contrasting results may also

stem from differences in the applied methods. While Wu et al.[16] used a modified dictator game, where

participants were explicitly asked whether they want to donate a certain monetary amount to charity,

in the present task, the donation to charity was determined indirectly by the participant's performance

in a reinforcement learning task. Our paradigm thus presents a more implicit measure of prosociality.

Eventually, as the bioavailability of testosterone after topical administration may be influenced by the

amount of subcutaneous fat at the application site[48], certain differences in the effects of an identical

testosterone dose might also result from ethnic differences in fat distribution[49].

Our results are, furthermore, in line with studies showing that testosterone decreases

deception[50-52]. Further research is, however, needed to determine whether testosterone reduces lying

per se, or only in situations where dishonest behavior may be considered "cheap", dishonorable and

lower the subject's feelings of pride and self-image[50]. In this respect, we note that our reinforcement

learning paradigm tapped into implicit processes and decisions that were likely less conducive to
deliberate or intended deception.

It is worth noting that our winning model with differential learning rates for positive and
negative PEs is in line with previous studies on learning and decision-making that report asymmetric
learning effect[53]. The theoretical interpretation of these differences, however, is mixed in the
literature. While some studies reported enhanced learning after positive PE compared to negative and
related this feature to optimism bias[54], others report a higher learning rate for negative than for
positive PEs, interpreted as possibly reflecting risk aversion[55]. Our findings in this regard may
contribute to this debate such that learning update is much quicker after receiving positive feedback,
especially when the feedback is associated with appetitive stimuli (e.g., monetary reward). This is
likely to be generalized to situations with relatively positive feedback, rather than the actual positive
feedback *per se*. For instance, in aversive learning, receiving no feedback (e.g., a neutral outcome) is,
on a relative scale, more positive than actual negative feedback (e.g., an electric shock). Learning
rates for such no-feedback events have been shown to be higher than those for stimuli with negative
feedback, including learning under social contexts[26].

There are also some limitations inherent to the methodology of our study. Due to the sex
differences in testosterone metabolism and unknown pharmacokinetics following the topical
administration of testosterone in women[56], the study included only male participants. Hence, the
generalization of these findings to females requires further investigation. Future research will also be
required to more systematically test whether testosterone's influence on the prosocial audience effect
is sensitive to the number and salience of the present observers. However, the data from the survey
suggest that the salience of the observers used in our study was representative of a wide range of
social contacts and that the subjective feeling of being watched even scaled with a well-established
stress biomarker.

In conclusion, we conducted a multifaceted examination of the computational,
endocrinological, and genetic mechanisms underlying audience effect and showed that testosterone
reduced strategic prosocial learning through impairment of choice consistency. These findings
provide the first evidence that in the Western student sample, testosterone abolishes audience effects,
and therefore does not foster the seeking of social leadership by reputational politics. Furthermore,
our research reveals that testosterone impacts status-seeking by influencing how the learned values are
expressed in behavior.

Materials and Methods

Participants

The study sample consisted of 192 healthy adult men aged between 18 and 40 years ($M = 24.89$, $SD = 4.08$). The sample size was determined based on previous testosterone administration studies[8,13,31] and our pilot study. Our sample size gave us 90% power to detect relatively small effects of size $f \geq 0.15$, for the main analyses of interest (i.e., interaction effects of the factors drug treatment, visibility, and type of recipient on the RLDDM parameters). Participants were recruited via flyers placed around university campuses and online advertisements. The volunteers, who replied to these advertisements, were screened via an online questionnaire and a telephone interview. The exclusion criteria comprised a history of neurological or psychiatric disorders, endocrine or other internal diseases, obesity, substance dependence, and the use of steroids. Only male participants were included as testosterone metabolism is subject to sex differences and the pharmacokinetics of topical administration of testosterone are unclear in women[56]. Three participants were excluded because of an unreliable change in hormone levels (see *SI: Supplementary analysis of hormone data)* and two participants were excluded because they continually clicked on the same response key irrespective of changing stimuli and reward probabilities for more than 80% of the block trials, and thus were classified as non-compliant. All participants gave written consent and received a financial reward for their participation consisting of a flat fee and a bonus based on their task performance. All procedures were approved by the local research ethics board and conducted following the latest revision of the Declaration of Helsinki[56].

Procedure and experimental conditions

Testing took place in groups of three to five participants, who were seated individually in small cubicles within the same testing room. All experimental sessions started between 01:00 and 02:30 p.m. First, a buccal smear sample for CAG repeat and DAT1 polymorphisms analysis was taken (see *SI*: *Supplementary information on the analysis of genetic data*). 20 min after arrival, participants were asked to drool 2 mL of saliva into a polyethylene collection tube. All salivary samples were frozen on-site and stored at - 30 °C until analysis. Afterward, participants were administered topical testosterone or placebo gel in a double-blind between-subjects design with random group allocation. Those allocated to the testosterone group received a single dose of testosterone gel, containing 150 mg testosterone [Androgel®]; participants in the placebo group received an equivalent amount of placebo gel. The only difference between the testosterone and placebo gel was that the placebo gel did not contain testosterone. Participants rubbed the gel onto their upper arms and shoulders using disposable latex gloves. Gel administration was followed by a 2-hour waiting period, during which participants remained in the laboratory premises, completed personality and demographic questionnaires, and were offered leisure-time reading materials. One hour and 50 minutes after the gel application, participants provided a second saliva sample and subsequently began the experimental

task (see Figure 1). Two more saliva samples were taken during the course of the study: 20 and 60

minutes after the end of the experimental task. After data collection was complete, saliva samples

were analyzed by liquid chromatography-tandem mass spectrometry in Dresden LabService GmbH,

Germany.

Participants performed the task under one of the two randomly assigned between-subject conditions,

either in private or when being observed. In the private condition, participants were informed that

their task performance is completely anonymous and no one (including the experimenter) would know

how much money they would earn for themselves and the charitable organization. In the observed

condition, two female observers, introduced as NGO association representatives entered the room and

watched participants perform the task. The observers were seated at a desk with a laptop and had an

equal view of all the participants. In addition, when participants were observed, a red frame was

shown on their computer´s screen and the displays of the participants' computers were transmitted

onto the observers' laptop screens.

Participants were thus randomly assigned into four experimental groups corresponding to the levels of

two between-subject factors: (1) treatment (testosterone/placebo) and (2) visibility (observed/private).

These groups did not differ in age, trait dominance, basal hormone levels, or distribution of AR CAG

and DAT1 genotype (see *SI: Table S1*).

Prosocial learning task

Participants performed a probabilistic reinforcement learning task[25], where they could earn rewards

either for themselves (self condition) or for an NGO of their choice (other condition). On each trial,

participants were presented with two abstract symbols, one associated with a high (75%) and the other

with a low (25%) reward probability. These contingencies were not instructed but had to be learned

through trial and error. Participants selected a symbol by a button press and then received feedback on

whether they obtained points or not. This way participants learned which symbol to choose to

maximize the rewards in the long run. The points were converted to monetary rewards at the end of

the experiment. Participants completed 6 blocks, 3 blocks in self and 3 blocks in the other condition.

Each block started with a new pair of symbols and consisted of 16 trials/choices. In the self condition,

the blocks started with "YOU" displayed and had the word "YOU" at the top of each screen. In the

other condition, the blocks started with "NGO" displayed and had the word "NGO" at the top of each

screen. The order of the blocks was pseudo-randomized so that the same recipient block did not occur

twice in a row, and that half of the participants' sample started the task with the *self* condition and the

other half with the *other* condition. At the end of the experimental task, participants could choose the

recipient of the money they earned in the other condition from a list of 6 different charities.

Immediately after completing the task, participants were asked to fill in a post-task questionnaire to

estimate their subjective perception of being watched. The participants were asked the question: "Did

you feel that you were being watched while performing the task?" The answers were classified into

three categories: 1 (Not at all), 2 (Moderately), and 3 (Strongly).

Statistical analysis of correct choices

Statistical analysis was performed using R statistical language[57]. We analyzed the treatment

(testosterone/placebo) x visibility (observed/private) x recipient (self/other) interaction effect on

correct choice using generalized linear mixed models (GzLMM) with binomial distribution and logit

link function[58]. The correct choice was defined as choosing the symbol with a higher reward

probability. The participant's identity was modeled as a random intercept effect and the within-subject

factor recipient (self/other) was entered as a random slope.

To examine whether the effects of testosterone on correct choice varied as a function of trait

dominance, CAG repeat, DAT1 polymorphism, baseline cortisol, cortisol reactivity, and personal

value orientations, we added these variables separately as predictors in interaction with the other

factors specified in the above GzLMM.

P-values were based on Type III Wald chi-square tests from the R car package, and post-hoc tests of

significant three-way interactions were conducted with the R sjPlot package that provided reported

odds ratios (ORs) together with 95% confidence intervals (95%CIs). Plots were created using yarrr

and ggplot2 R packages (See *SI* for references).

Reinforcement learning drift-diffusion modeling

To uncover the cognitive computational processes underlying our learning task, we performed

modeling analysis under the joint reinforcement learning drift diffusion model (RLDDM)

framework[24,35]. In essence, RLDDM bridges RL, which typically models choices, and DDM, which

commonly models response times (RT). This approach has been proven to provide more granularity

than using RL or DDM alone[24].

Previous studies have reported differences in learning following positive and negative PEs[55,60] and

these differences have been linked to the distinctive roles of striatal D1 and D2 dopamine receptors in

segregated cortico-striatal pathways[61]. Furthermore, analyses of genetic polymorphisms of DARPP-32

that predict choice behavior associated with a positive outcome, and the DRD2 gene predicting

avoidance of choices associated with negative outcomes, support the notion that independent

dopaminergic mechanisms contribute to learning from positive and negative feedback[62]. RL models

with dual learning rates have been successful in capturing this asymmetric learning effect[53,63],

including studies in social neuroscience examining prosocial behavior[26]. We therefore tested

candidate models with a single learning rate (Rescorla-Wagner models) as well as models with dual

learning rates. Together, we tested 6 candidate RLDDM models, and the winning model is described

below (see *SI: Supplementary information on computational modeling* for full model description,

estimation, and comparison procedures). The RL part of the winning RLDDM model was

implemented with a dual learning rates reinforcement learning model, where both the learning rate for

a positive prediction error and the learning rate for a negative prediction error were employed to

update values (i.e., V(A) and V(B) for two-choice options) [54] (Equation (1); see also *SI:*

Supplementary information on computational modeling).

The DDM part of the winning RLDDM model was implemented via a non-linear transformation of

the accuracy-codded value differences computed from the RL counterpart, to construct the trial-by-

trial drift rates[24] (Equation (2); see *also SI: Supplementary information on computational modeling*).

The winning model contained 14 parameters: 7 separate parameters for each between-subject

condition (i.e., placebo/testosterone, private/observed), and differential parameters for the within-

subject condition (i.e. other/self; see *SI: Table S3* for the parameter list and description).

In all models, we simultaneously modeled participants' choice and RT, separately for each between-

subject condition (i.e., placebo/testosterone; observed/private). Model estimations of all candidate

models were performed with hierarchical Bayesian analysis (HBA). HBA was particularly useful

when the number of trials was limited (here, 16 trials per block per participant), because in HBA,

group-level and individual-level parameters were mutually informing each other during model

estimation. All models reached convergence (see *SI: Supplementary information on computational*

modeling).

Statistical analysis of model parameters

The drug treatment (testosterone/placebo) x visibility (observed/private) x recipient (other/self) effect

on the extracted free parameters was analyzed using GzLMMs analogous to the analysis of the correct

choice. Due to the non-normal distribution of residuals, gamma distribution with a log link function

was used for the parameter analyses. Finally, we tested whether the RLDDM parameter estimates,

which were affected by the interaction of the drug treatment, visibility, and recipient could explain the

differences observed in the behavioral prosociality measure. To do so, we conducted multiple linear

regressions with the difference in the number of correct choices made for other and self (*prosociality*

index) as a dependent variable and the differences in the RLDDM parameter estimates (αneg_{other}-

αneg_{self}, τ_{other} – τ_{self}, $threshold_{other}$ -$threshold_{self}$, $drift\text{-}scaling_{other}$-$drift\text{-}scaling_{self}$) as separate predictors.

Bonferroni correction for multiple comparisons was used.

Simulation to identify optimal learning rates to aid interpretation

As it is recommended to interpret model parameters in the context of task structures (e.g., reward

probability, trial number[37]), we conducted simulations with grid approximation to identify the

parameter sets of "optimal learning rates" for each condition. The optimal learning rates here referred

to the combination of the two learning rates that could generate the highest choice frequency of

choosing the better option within each pair. We then compared the estimated learning rates for each

condition with respect to the optimal learning rates and draw further inferences (Figure 4; see also *SI: Supplementary information on the computational analysis*).

Acknowledgments

We thank Felix Dörflinger and Maximilian Kathofer for their dedicated assistance with data collection. We also thank Shawn Geniole for his helpful feedback. This work was supported by the Vienna Science and Technology Fund (WWTF VRG 13-007).

References

1. Bradley A, Lawrence C, Ferguson E. 2018. Does observability affect prosociality? *Proceedings of the Royal Society B: Biological Sciences* **285**:20180116. doi:10.1098/rspb.2018.0116

2. Lacetera N, Macis M. 2010. Social image concerns and prosocial behavior: Field evidence from a nonlinear incentive scheme. *Journal of Economic Behavior & Organization* **76**:225–237. doi:10.1016/j.jebo.2010.08.007

3. Soetevent AR. 2005. Anonymity in giving in a natural context—a field experiment in 30 churches. *Journal of Public Economics* **89**:2301–2323. doi:10.1016/j.jpubeco.2004.11.002

4. Li Y, Météreau E, Obeso I, Butera L, Villeval MC, Dreher J-C. 2020. Endogenous testosterone is associated with increased striatal response to audience effects during prosocial choices. *Psychoneuroendocrinology* **122**:104872. doi:10.1016/j.psyneuen.2020.104872

5. Böckler A, Tusche A, Singer T. 2016. The Structure of Human Prosociality: Differentiating Altruistically Motivated, Norm Motivated, Strategically Motivated, and Self-Reported Prosocial Behavior. *Social Psychological and Personality Science* **7**:530–541. doi:10.1177/1948550616639650

6. Hamilton AF de C, Lind F. 2016. Audience effects: what can they tell us about social neuroscience, theory of mind and autism? *Cult Brain* **4**:159–177. doi:10.1007/s40167-016-0044-5

7. Gintis H, Smith EA, Bowles S. 2001. Costly Signaling and Cooperation. *Journal of Theoretical Biology* **213**:103–119. doi:10.1006/jtbi.2001.2406

8. Dreher J-C, Dunne S, Pazderska A, Frodl T, Nolan JJ, O'Doherty JP. 2016. Testosterone causes both prosocial and antisocial status-enhancing behaviors in human males. *PNAS* **113**:11633–11638. doi:10.1073/pnas.1608085113

9. Ou J, Wu Y, Hu Y, Gao X, Li H, Tobler PN. 2021. Testosterone reduces generosity through cortical and subcortical mechanisms. *PNAS* **118**. doi:10.1073/pnas.2021745118

10. Terburg D, Syal S, Rosenberger LA, Heany SJ, Stein DJ, Honk J van. 2016. Testosterone abolishes implicit subordination in social anxiety. *Psychoneuroendocrinology* **72**:205–211. doi:10.1016/j.psyneuen.2016.07.203

11. Geniole SN, Proietti V, Bird BM, Ortiz TL, Bonin PL, Goldfarb B, Watson NV, Carré JM. 2019. Testosterone reduces the threat premium in competitive resource division. *Proc Biol Sci* **286**:20190720. doi:10.1098/rspb.2019.0720

12. Kutlikova HH, Geniole SN, Eisenegger C, Lamm C, Jocham G, Studer B. 2021. Not giving up: Testosterone promotes persistence against a stronger opponent. *Psychoneuroendocrinology* **128**. doi:10.1016/j.psyneuen.2021.105214

13. Kutlikova HH, Durdiaková JB, Wagner B, Vlček M, Eisenegger C, Lamm C, Riečanský I. 2020. The effects of testosterone on the physiological response to social and somatic stressors. *Psychoneuroendocrinology* **117**:104693. doi:10.1016/j.psyneuen.2020.104693

14. Cañigueral R, Hamilton AF de C. 2019. Being watched: Effects of an audience on eye gaze and prosocial behaviour. *Acta Psychol (Amst)* **195**:50–63. doi:10.1016/j.actpsy.2019.02.002

15. Eisenegger C, Haushofer J, Fehr E. 2011. The role of testosterone in social interaction. *Trends in Cognitive Sciences* **15**:263–271. doi:10.1016/j.tics.2011.04.008

16. Wu Y, Zhang Y, Ou J, Hu Y, Zilioli S. 2020. Exogenous testosterone increases the audience effect in healthy males: evidence for the social status hypothesis. *Proceedings of the Royal Society B: Biological Sciences* **287**:20200976. doi:10.1098/rspb.2020.0976

17. Terburg D, van Honk J. 2013. Approach–Avoidance versus Dominance–Submissiveness: A Multilevel Neural Framework on How Testosterone Promotes Social Status. *Emotion Review* **5**:296–302. doi:10.1177/1754073913477510

18. de Souza Silva MA, Mattern C, Topic B, Buddenberg TE, Huston JP. 2009. Dopaminergic and serotonergic activity in neostriatum and nucleus accumbens enhanced by intranasal administration of testosterone. *European Neuropsychopharmacology* **19**:53–63. doi:10.1016/j.euroneuro.2008.08.003

19. Hermans EJ, Bos PA, Ossewaarde L, Ramsey NF, Fernández G, van Honk J. 2010. Effects of exogenous testosterone on the ventral striatal BOLD response during reward anticipation in healthy women. *Neuroimage* **52**:277–283. doi:10.1016/j.neuroimage.2010.04.019

20. Op de Macks ZA, Moor BG, Overgaauw S, Güroğlu B, Dahl RE, Crone EA. 2011. Testosterone levels correspond with increased ventral striatum activation in response to monetary rewards in adolescents. *Developmental Cognitive Neuroscience*, Special Issue on Motivation **1**:506–516. doi:10.1016/j.dcn.2011.06.003

21. Sutton RS, Barto AG. 2018. Reinforcement Learning, second edition: An Introduction. MIT Press.

22. Cohen JY, Haesler S, Vong L, Lowell BB, Uchida N. 2012. Neuron-type-specific signals for reward and punishment in the ventral tegmental area. *Nature* **482**:85–88. doi:10.1038/nature10754

23. Diederen KMJ, Spencer T, Vestergaard MD, Fletcher PC, Schultz W. 2016. Adaptive Prediction Error Coding in the Human Midbrain and Striatum Facilitates Behavioral Adaptation and Learning Efficiency. *Neuron* **90**:1127–1138. doi:10.1016/j.neuron.2016.04.019

24. Fontanesi L, Gluth S, Spektor MS, Rieskamp J. 2019. A reinforcement learning diffusion decision model for value-based decisions. *Psychon Bull Rev* **26**:1099–1121. doi:10.3758/s13423-018-1554-2

25. Lockwood PL, Apps MAJ, Valton V, Viding E, Roiser JP. 2016. Neurocomputational mechanisms of prosocial learning and links to empathy. *PNAS* **113**:9763–9768. doi:10.1073/pnas.1603198113

26. Lengersdorff LL, Wagner IC, Lockwood PL, Lamm C. 2020. When Implicit Prosociality Trumps Selfishness: The Neural Valuation System Underpins More Optimal Choices When Learning to Avoid Harm to Others Than to Oneself. *J Neurosci* **40**:7286–7299. doi:10.1523/JNEUROSCI.0842-20.2020

27. Geniole SN, Procyshyn TL, Marley N, Ortiz TL, Bird BM, Marcellus AL, Welker KM, Bonin PL, Goldfarb B, Watson NV, Carré JM. 2019. Using a Psychopharmacogenetic Approach To Identify the Pathways Through Which—and the People for Whom—Testosterone Promotes Aggression. *Psychol Sci* **30**:481–494. doi:10.1177/0956797619826970

28. Knight EL, Sarkar A, Prasad S, Mehta PH. 2020. Beyond the challenge hypothesis: The
emergence of the dual-hormone hypothesis and recommendations for future research.
Hormones and Behavior, 30th Anniversary of the Challenge Hypothesis **123**:104657.
doi:10.1016/j.yhbeh.2019.104657

29. Prasad S, Knight EL, Mehta PH. 2019. Basal testosterone's relationship with dictator game
decision-making depends on cortisol reactivity to acute stress: A dual-hormone perspective on
dominant behavior during resource allocation. *Psychoneuroendocrinology* **101**:150–159.
doi:10.1016/j.psyneuen.2018.11.012

30. Qu C, Ligneul R, Van der Henst J-B, Dreher J-C. 2017. An Integrative Interdisciplinary
Perspective on Social Dominance Hierarchies. *Trends in Cognitive Sciences* **21**:893–908.
doi:10.1016/j.tics.2017.08.004

31. Carré JM, Geniole SN, Ortiz TL, Bird BM, Videto A, Bonin PL. 2017. Exogenous
Testosterone Rapidly Increases Aggressive Behavior in Dominant and Impulsive Men. *Biol
Psychiatry* **82**:249–256. doi:10.1016/j.biopsych.2016.06.009

32. Cheng JT, Tracy JL, Henrich J. 2010. Pride, personality, and the evolutionary foundations of
human social status. *Evolution and Human Behavior* **31**:334–347.
doi:10.1016/j.evolhumbehav.2010.02.004

33. Schwartz SH. 2007. Chapter 7 A Proposal for Measuring Value Orientations across Nations.

34. Schwartz, SH, Breyer, B, & Danner, D. 2015. Human Values Scale (ESS). *Zusammenstellung
sozialwissenschaftlicher Items und Skalen (ZIS)*. doi:10.6102/zis234

35. Ratcliff R, McKoon G. 2008. The Diffusion Decision Model: Theory and Data for Two-
Choice Decision Tasks. *Neural Comput* **20**:873–922. doi:10.1162/neco.2008.12-06-420

36. Ahn W-Y, Haines N, Zhang L. 2017. Revealing Neurocomputational Mechanisms of
Reinforcement Learning and Decision-Making With the hBayesDM Package. *Comput
Psychiatr* **1**:24–57. doi:10.1162/CPSY_a_00002

37. Zhang L, Lengersdorff L, Mikus N, Gläscher J, Lamm C. 2020. Using reinforcement learning
models in social neuroscience: frameworks, pitfalls and suggestions of best practices. *Social
Cognitive and Affective Neuroscience* **15**:695–707. doi:10.1093/scan/nsaa089

38. Dabbs JM. 1998. Testosterone and the concept of dominance. *Behavioral and Brain Sciences*
21:370–371. doi:10.1017/S0140525X98331222

39. Mazur A, Booth A. 1998. Testosterone and dominance in men. *Behav Brain Sci* **21**:353–363;
discussion 363-397.

40. Daw ND, O'Doherty JP, Dayan P, Seymour B, Dolan RJ. 2006. Cortical substrates for
exploratory decisions in humans. *Nature* **441**:876–879. doi:10.1038/nature04766

41. Morgan D, Grant KA, Gage HD, Mach RH, Kaplan JR, Prioleau O, Nader SH, Buchheimer
N,
Ehrenkaufer RL, Nader MA. 2002. Social dominance in monkeys: dopamine D2 receptors
and cocaine self-administration. *Nat Neurosci* **5**:169–174. doi:10.1038/nn798

42. Eisenegger C, Naef M, Linssen A, Clark L, Gandamaneni PK, Müller U, Robbins TW. 2014.
Role of Dopamine D2 Receptors in Human Reinforcement Learning.
Neuropsychopharmacology **39**:2366–2375. doi:10.1038/npp.2014.84

43. Aikey JL, Nyby JG, Anmuth DM, James PJ. 2002. Testosterone rapidly reduces anxiety in male house mice (Mus musculus). *Horm Behav* **42**:448–460. doi:10.1006/hbeh.2002.1838

44. Fernández-Guasti A, Martínez-Mota L. 2005. Anxiolytic-like actions of testosterone in the burying behavior test: role of androgen and GABA-benzodiazepine receptors. *Psychoneuroendocrinology* **30**:762–770. doi:10.1016/j.psyneuen.2005.03.006

45. Bellezza S, Gino F, Keinan A. 2014. The red sneakers effect: Inferring status and competence from signals of nonconformity. *Journal of Consumer Research* **41**:35–54. doi:10.1086/674870

46. Miyamoto Y, Yoo J, Levine CS, Park J, Boylan JM, Sims T, Markus HR, Kitayama S, Kawakami N, Karasawa M, Coe CL, Love GD, Ryff CD. 2018. Culture and Social Hierarchy: Self- and Other-Oriented Correlates of Socioeconomic Status across Cultures. *J Pers Soc Psychol* **115**:427–445. doi:10.1037/pspi0000133

47. Kitayama S, King A, Yoon C, Tompson S, Huff S, Liberzon I. 2014. The dopamine D4 receptor gene (DRD4) moderates cultural difference in independent versus interdependent social orientation. *Psychol Sci* **25**:1169–1177. doi:10.1177/0956797614528338

48. Miller J, Britto M, Fitzpatrick S, McWhirter C, Testino SA, Brennan JJ, ZumBrunnen TL. 2011. Pharmacokinetics And Relative Bioavailability Of Absorbed Testosterone After Administration Of A 1.62% Testosterone Gel To Different Application Sites In Men With Hypogonadism. *Endocrine Practice* **17**:574–583. doi:10.4158/EP10192.OR

49. Wu C-H, Heshka S, Wang J, Pierson RN, Heymsfield SB, Laferrère B, Wang Z, Albu JB, Pi-Sunyer X, Gallagher D. 2007. Truncal fat in relation to total body fat: influences of age, sex, ethnicity and fatness. *Int J Obes* **31**:1384–1391. doi:10.1038/sj.ijo.0803624

50. Wibral M, Dohmen T, Klingmüller D, Weber B, Falk A. 2012. Testosterone Administration Reduces Lying in Men. *PLOS ONE* **7**:e46774. doi:10.1371/journal.pone.0046774

51. van Honk J, Will G-J, Terburg D, Raub W, Eisenegger C, Buskens V. 2016. Effects of Testosterone Administration on Strategic Gambling in Poker Play. *Sci Rep* **6**:18096. doi:10.1038/srep18096

52. Henderson A, Thoelen G, Nadler A, Barraza J, Nave G. 2018. Testing the influence of testosterone administration on men's honesty in a large laboratory experiment. *Sci Rep* **8**:11556. doi:10.1038/s41598-018-29928-z

53. den Ouden HEM, Daw ND, Fernandez G, Elshout JA, Rijpkema M, Hoogman M, Franke B, Cools R. 2013. Dissociable effects of dopamine and serotonin on reversal learning. *Neuron* **80**:1090–1100. doi:10.1016/j.neuron.2013.08.030

54. Lefebvre G, Lebreton M, Meyniel F, Bourgeois-Gironde S, Palminteri S. 2017. Behavioural and neural characterization of optimistic reinforcement learning. *Nat Hum Behav* **1**:1–9. doi:10.1038/s41562-017-0067

55. Gershman SJ. 2015. Do learning rates adapt to the distribution of rewards? *Psychon Bull Rev* **22**:1320–1327. doi:10.3758/s13423-014-0790-3

56. Eisenegger C, von Eckardstein A, Fehr E, von Eckardstein S. 2013. Pharmacokinetics of testosterone and estradiol gel preparations in healthy young men. *Psychoneuroendocrinology* **38**:171–178. doi:10.1016/j.psyneuen.2012.05.018

57. WMA - The World Medical Association-WMA Declaration of Helsinki – Ethical Principles for Medical Research Involving Human Subjects (June 29, 2021).

58. R Core Team, *R: A Language and Environment for Statistical Computing* (R Foundation for Statistical Computing, 2020).

59. Bates D, Mächler M, Bolker B, Walker S. 2015. Fitting Linear Mixed-Effects Models Using lme4. *Journal of Statistical Software* **67**:1–48. doi:10.18637/jss.v067.i01

60. Pedersen ML, Frank MJ, Biele G. 2017. The drift diffusion model as the choice rule in reinforcement learning. *Psychon Bull Rev* **24**:1234–1251. doi:10.3758/s13423-016-1199-y

61. Cox SML, Frank MJ, Larcher K, Fellows LK, Clark CA, Leyton M, Dagher A. 2015. Striatal D1 and D2 signaling differentially predict learning from positive and negative outcomes. *Neuroimage* **109**:95–101. doi:10.1016/j.neuroimage.2014.12.070

62. Frank MJ, Moustafa AA, Haughey HM, Curran T, Hutchison KE. 2007. Genetic triple dissociation reveals multiple roles for dopamine in reinforcement learning. *Proc Natl Acad Sci U S A* **104**:16311–16316. doi:10.1073/pnas.0706111104

63. Crawley D, Zhang L, Jones EJH, Ahmad J, Oakley B, Cáceres ASJ, Charman T, Buitelaar JK, Murphy DGM, Chatham C, Ouden H den, Loth E, Group the E-AL. 2020. Modeling flexible behavior in childhood to adulthood shows age-dependent learning mechanisms and less optimal learning in autism in each age group. *PLOS Biology* **18**:e3000908. doi:10.1371/journal.pbio.3000908

951   **Supplementary information for:**
952
953   **Testosterone eliminates strategic prosocial behavior in healthy males**
954

955   Hana H. Kutlikova, Lei Zhang, Christoph Eisenegger, Jack van Honk, Claus Lamm
956

957   Corresponding author: Hana H. Kutlikova
958   Email: hana.kutlikova@univie.ac.at

959
960
961   This PDF file includes:
962
963   Supplementary text
964   Supplementary information on the use of statistical software and functions
965   Supplementary analysis of hormone data
966   Supplementary information on computational modeling
967   Supplementary information on the analysis of the genetic data
968   Supplementary information on the questionnaire data
969   Figure S1
970   Tables S1 to S3
971   SI References
972
973   Other supplementary materials for this manuscript include the following:
974   Data and codes can be accessed at: https://osf.io/qr4ve/

Supplementary information on the use of statistical software and functions

Statistical analysis was performed using the R statistical language[1] and the packages: lme4[2] for construction of general linear mixed models (GLMM) and generalized linear mixed model (GzLMM); car package[3] for construction of general linear models and computation of *p*-values based on Type III Wald chi-square tests; sjPlot package[4] for post-hoc tests of significant three-way interactions including odds ratios (ORs) and 95% confidence intervals (95%CIs); yarrr[5] and ggplot2[6] packages for construction of plots. Correlations were compared using cocor pachage[7]. Computational modeling was performed using Markov chain Monte Carlo with the statistical computing language Stan[8] while following the hBayesDM package[9]. Model comparison and evaluation were performed using the LOO package[10].

Supplementary analysis of hormone data

The effect of drug treatment on hormone levels.

After data collection was complete, saliva samples were shipped on dry ice to Dresden LabService GmbH led by Clemens Kirschbaum, Germany. Liquid chromatography-tandem mass spectrometry was used to determine the hormonal levels. To examine the change of the hormonal levels throughout the experimental session, testosterone, cortisol, and estradiol levels were analyzed using GLMMs with the fixed factors drug treatment (testosterone/placebo), visibility (observed/private), time (baseline/1 h 50 min after drug treatment/20 min after the end of the task/60 min after the end of the task), and participant's identity as a random intercept. Due to the non-normal distribution of residuals, hormonal data were log-transformed. Baseline hormonal levels did not significantly differ across experimental groups (all *p*s > .263, see Table S1). As expected, 1h 50 min after gel administration, we observed higher testosterone levels in the testosterone group ($M_{Sample2}$ = 5014.10 pg/mL, 95%CI [3866.10, 6502.88]) compared to the placebo group ($M_{Sample2}$ = 134.30, 95%CI [103.54, 175.92]; drug treatment x time: $F(3,522.59)$ = 47.82, $p < .001$, $\eta2$ =.423), a difference that remained stable until the end of the experiment (see Figure S1).

Testosterone levels of two participants in the testosterone group decreased after the drug administration (Sample 1/Sample 2: 629.29/308.95 pg/mL and 350.76/294.76 pg/mL), while testosterone levels of one participant in the placebo group increased considerably (Sample 1/Sample 2: 546.133/1099.08 pg/mL). These three participants were excluded from the analyses.

There were no effects of drug treatment on cortisol (drug treatment x time: $F(3,550.23)$ = 0.149, p = .930) or estradiol levels (drug treatment x time: $F(3,551.12)$ = 1.420, p = .236).

The observation condition did not significantly influence any hormonal levels (testosterone: visibility x time: $F(3,520.88)$ = 1.528, p = .206; cortisol: visibility x time: $F(3,520.6)$ = 0.9, p = 441; estradiol: visibility x time: $F(3,521.37)$ = 1.699, p = .166).

Contamination of salivary samples.

Out of a total of 192 baseline samples, we noted that 34 contained above-normal testosterone levels,

atypical for normal young men (> 2000 pg/mL). All other baseline values were hormonally typical.

The samples with abnormally high testosterone values appeared only in the participants, who later

received testosterone treatment, not in the placebo group. Previous research[11,12] described similar

abnormally high testosterone levels and attributed it to the testosterone contamination of the common

surfaces (e.g., doorknobs, keyboards), excluding the option of physiological contamination. Based on

their recommendation, we implemented a cleaning protocol that included the wearing of disposable

sterile gloves, and cleaning of keyboards, computer mice, tables, and doorknobs with an alcohol-

based solution after each session. Although these precautions successfully prevented between-session

contamination, we suspect that they still did not reliably impede within-session contamination of the

saliva containers. For future studies, we, therefore, recommend even stricter sanitizing protocols and

more careful handling of the saliva collection tubes and boxes, before, during, and also after sample

collection.

In our sample, the abnormally high values were present only in the sessions where testosterone was

administered, and this notwithstanding, the testosterone group showed a reliable testosterone increase

after the drug administration in comparison to the placebo group. We, therefore, decided to retain the

participants with contaminated baseline samples for the behavioral analyses, except for the analysis

that includes baseline testosterone levels.

Interaction of cortisol levels with testosterone effects on the correct choice and RLDDM

parameters.

To examine whether cortisol levels interacted with testosterone´s effect on correct choice and

reinforcement learning drift diffusion model (RLDDM) parameters, we separately added baseline

cortisol levels and cortisol reactivity as predictors to our main analysis. The cortisol reactivity was

defined as a difference between cortisol levels detected in the saliva sample taken 20 minutes after the

end of visibility manipulation and cortisol levels from the sample taken immediately before the start

of the paradigm. The values were mean-centered and entered as a predictor in interaction with the

other factors (recipient, drug treatment, visibility) to the GzLMM of correct choice and the GzLMMs

of the RLDDM parameters with log link function (see *Materials and Methods* in the main manuscript

text). The analysis revealed no significant interaction of baseline cortisol levels with testosterone

effect on correct choice (recipient x drug treatment x visibility baseline cortisol: $OR = 0.86$, $CI =$

$[0.73, 1.013]$, $p = .075$), α^{posPE} (recipient x drug treatment x visibility baseline cortisol: $B = 0.99$, $CI =$

$[0.92, 1.09]$, $p = .854$), α^{negPE} (recipient x drug treatment x visibility baseline cortisol: $B = 0.95$, $CI =$

$[0.857, 1.06]$, $p = .349$), choice consistency (recipient x drug treatment x visibility baseline cortisol: B

$= 1.00$, $CI = [0.980, 1.01]$, $p = .468$) or decision threshold (recipient x drug treatment x visibility

baseline cortisol: $OR = 1.00$, $CI = [0.99, 1.01]$, $p = .490$).

Similarly, the analysis revealed no significant interaction of cortisol reactivity with testosterone effect

on correct choice (recipient x drug treatment x visibility x post-task cortisol: $OR = 1.04$, $CI = [0.99,$

$1.09]$, $p = .122$), α^{posPE} (recipient x drug treatment x visibility x post-task cortisol: $B = 1.01$, $CI =$

$[0.99, 1.03]$, $p = .462$), α^{negPE} (recipient x drug treatment x visibility x post-task cortisol: $B = 1.01$, CI

$= [0.97, 1.04]$, $p = .738$), choice consistency (recipient x drug treatment x visibility x post-task

cortisol: $B = 1.00$, $CI = [1.00, 1.01]$, $p = .249$) or decision threshold (recipient x drug treatment x

visibility x post-task cortisol: $B = 1.00$, $CI = [1.00, 1.00]$, $p = .983$).

1073

Supplementary information on computational modeling

Rescorla-Wagner (RW) model.

We started with the simple Rescorla-Wagner[13] model as our baseline model. On each trial, the value

($V_{c,t}$) of the chosen option was updated with the reward prediction error (RPE):

$$V_{c,t} = V_{c,t-1} + \alpha \left(O_{t-1} - V_{c,t-1} \right),$$ (1)

where O_{t-1} was the received outcome, and α ($0 < \alpha < 1$) denoted the learning rate.

Dual learning rates (DLR) model.

Studies have suggested that individuals may have separate updates for positive and negative

feedback[14]. Hence, we tested a dual learning rates model on top of the RW model:

$$V_{c,t} = \begin{cases} V_{c,t-1} + \alpha^{pos} \left(O_{t-1} - V_{c,t-1} \right), \text{if } O_{t-1} > 0 \\ V_{c,t-1} + \alpha^{neg} \left(O_{t-1} - V_{c,t-1} \right), \text{otherwise} \end{cases},$$ (2)

where α^{posPE} and α^{negPE} were the learning rates for positive and negative RPEs, respectively.

In both RW and DLR models, action values were converted to action probabilities using the softmax

function. Let A and B be the choice symbols per trial, the probability of choosing A was computed via

the difference between $V(A)$ and $V(B)$:

$$p(A) = \frac{1}{1 + e^{\beta(-(V_A - V_B))}}, \; p(B) = 1 - p(A),$$ (3)

where β ($\beta > 0$) was the inverse temperature that represented choice consistency. Higher β indicated

that individuals' choices were more consistent with their value computation, where lower β indicated

that individuals behaved more randomly. The action probability was then used to model participants'

choice data with a categorical distribution:

$$choice_t \sim categorical\left([p_t(A), p_t(B)] \right)$$ (4)

Drift diffusion model (DDM).

The drift diffusion model (a.k.a., diffusion decision model[15]) was a widely used computational

framework to model individuals' response times (RTs). In its canonical expression, DDM contained

four parameters, namely, the drift rate (v; v > 0), the initial bias (z; z > 0), non-decision time (T; 0 < T

min(RT)), as well as the decision threshold (a). For simplicity in learning tasks with abstract symbols,

the initial bias z was fixed at 0.5. Trial-by-trial RTs were distributed according to the Wiener first

passage time (WFPT[16]):

$$RT_t \sim wfpt\left(a, T, z, v\right)$$

1101   (5)

1102

Reinforcement learning drift diffusion model (RLDDM).

In value-based decision-making, individuals' RTs may vary as the function of trial-by-trial valuation,

such that the larger the value difference between choice alternatives, the faster the RT. Therefore, a

joint reinforcement learning drift diffusion model (RLDDM) framework has been proposed[17,18],

bridging RL and DDM. This approach provides more granularity than using RL or DDM alone[18]. In

essence, the drift rate in DDM was characterized by the accuracy-coded value differences computed

from the RL counterpart. This way, the drift rate was no more a constant parameter throughout the

entire experiment, instead, it varied across trials (i.e., v_t, instead of v) according to the values

computed from RL updates (in the present study, RW or RP). In the simplest RLDDM, trial-by-trial

drift rates were constructed via a linear function of value difference:

$$v_t = v_{\text{scaling}}\left(V_{\text{correct},t} - V_{\text{incorrect},t}\right)$$

1113   (6)

1114   where v_{scaling} ($v_{\text{scaling}} > 0$) was the scaling parameter that quantified the impact of value difference. Note

1115   that we employed stimulus coding in our RLDDM, so that in Equation (6), the drift rate was always a

1116   function of the value difference between the correct (i.e., more rewarding, 75% reward probability)

1117   and the incorrect options (i.e., less rewarding, 25% reward probability), rather than between the

1118   chosen and unchosen options.

1119

1120   ***Reinforcement learning drift diffusion model with non-linear transformation (RLDDM-nonlin).***

1121   There is evidence that a non-linear mapping between value difference and the drift rate could better

1122   capture individuals' RTs as opposed to a linear transformation[18]. This is likely because non-linear

1123   functions may provide more sensitivity, akin to the softmax function in choice models. We thus

1124   implemented an RLDDM-nonlin following:

$$v_t = S\left[v_{\text{scaling}}\left(V_{\text{correct},t} - V_{\text{incorrect},t}\right)\right]$$

1125   (7)

1126   with

$$S(x) = 2 \cdot \frac{v_{\max}}{1 + e^{-x}} - v_{\max}$$

1127   (8)

where $S(x)$ was a non-linear sigmoid function centered at 0, that convert x to lie between $-v_{\max}$ and

$v_{\max}$ ($v_{\max} > 0$). It is worth noting that $v_{\max}$ only affected the maximum value of the drift rate, whereas

$v_{scaling}$, as in Equation 6, established the trial-by-trial mapping between the value difference and the

drift rate.

In both RLDDM and RLDDM-nonlin, all other DDM parameters (i.e., a, T, z) were identical to the

canonical DDM model, and RTs were distributed with *wfpt* using trial-by-trial drift rate (v_t):

1134
$$RT_i \sim wfpt\left(a, T, z, v_t\right)_{.} \tag{9}$$

1135 Note that, in all candidate models (Table S1), we introduced differential parameters for the within-

1136 subject condition of our experiment, namely, all parameters were separately modeled for the "self"

1137 and the "other" conditions.

1138

Model estimation.

The model estimation and model selection procedures were largely similar to[19]. Hence, below we

echoed these procedures from[19] to enhance reproducibility, with modifications that were specific to

the current study.

In all models, we simultaneously modeled participants' choice and RT, separately for each between-

subject condition (i.e., placebo vs. testosterone; observed vs. private). Model estimations of all

candidate models were performed with hierarchical Bayesian analysis (HBA)[20] using the statistical

computing language Stan[8] in R. Stan utilizes a Hamiltonian Monte Carlo (HMC; an efficient Markov

Chain Monte Carlo, MCMC) sampling scheme to perform full Bayesian inference and obtain the

actual posterior distribution. We performed HBA rather than maximum likelihood estimation (MLE)

because HBA provides much more stable and accurate estimates than MLE[18]. Following the approach

in the "hBayesDM" package[9] for using Stan in the field of reinforcement learning, we assumed, for

instance, that a generic individual-level parameter φ was drawn from a group-level normal

distribution, namely, $\varphi \sim$ Normal (μ_φ, σ_φ), with μ_φ and σ_φ. being the group-level mean and standard

deviation, respectively. Both these group-level parameters were specified with weakly-informative

priors[20]: $\mu_\varphi \sim$ Normal (0, 1) and $\sigma_\varphi \sim$ half-Cauchy (0, 1). This was to ensure that the MCMC sampler

traveled over a sufficiently wide range to sample the entire parameter space. Appropriate parameter

transformations were applied to double-bounded parameters (e.g., learning rate, [0, 1]) with the

inverse probit function (i.e., the cumulative distribution function of the standard normal distribution),

and single-bounded parameters (e.g., drift rate, (0, +∞)) with the soft-plus function (i.e., ln(1 + e^x)),

respectively.

1160

In HBA, all group-level parameters and individual-level parameters were simultaneously estimated

through the Bayes' rule by incorporating behavioral data. We fit each candidate model with four

independent MCMC chains using 1,000 iterations after 1,000 iterations for the initial algorithm

warmup per chain, which resulted in 4,000 valid posterior samples. The convergence of MCMC

chains was assessed both visually (from the trace plot) and through the Gelman-Rubin $\hat{R}$ Statistics[21].

R̂ values of all parameters were smaller than 1.05 in the current study), which indicated adequate

convergence.

1168

Model selection and validation.

For model comparison and model selection, we computed the Leave-One-Out information criterion

(LOOIC) score per candidate model[22]. The LOOIC score provides the point-wise estimate (using the

entire posterior distribution) of out-of-sample predictive accuracy in a fully Bayesian way, which is

more reliable compared to information criteria using point-estimate (e.g., Akaike information

criterion, AIC; deviance information criterion, DIC). By convention, a lower LOOIC score indicates

better out-of-sample prediction accuracy of the candidate model. We selected the model with the

lowest LOOIC as the winning model. We additionally performed Bayesian model averaging (BMA)

with Bayesian bootstrap[23] to compute the probability of each candidate model being the best model.

Conventionally, the BMA probability of 0.8 (or higher) is a decisive indication.

Moreover, given that model comparison provided merely relative performance among candidate

models[22], we then tested how well our winning model's posterior prediction was able to replicate the

key features of the observed data (a.k.a., posterior predictive checks, PPCs). Since we only found an

effect in choice data, we performed PPCs only for choices (excluding RTs). To this end, we applied a

one-step-ahead PPC[19,24] that factored in participants' actual action and outcome sequences to generate

predictions with the entire posterior MCMC samples. Specifically, we let the winning model generate

choices as many times as the number of MCMC samples (i.e., 4,000 times) per trial per participant,

and we analyzed the generated data the same way as we did for the observed data. We then assessed

whether these analyses could reproduce the behavioral pattern in our behavioral analyses (Figure 4B,

4D in the main text).

1189

Simulations of optimal learning rates.

To better understand the magnitude of the posterior learning rates, we performed simulations to obtain

"optimal learning rates", and then compared the posterior parameters in relation to these optimal

parameters (Figure 4A, 4C in the main text). Because there were two learning rates (α^{posPE} and α^{negPE}),

to reduce complexity, we fixed the inverse temperature parameter to be the corresponding group-level

posterior mean in each condition. For each simulation, we took a small grid per parameter (0:0.01:1)

and computed the choice accuracy across 16 trials (identical to the main experiment) for each

combination of the parameters. Each simulation was repeated 1000 times to obtain stable results. We

then considered the parameters that gave the highest choice accuracy as the optimal learning rates.

1199

1200

Analysis of the RLDDM parameters and their association with prosocial behavior.

Next, we examined whether the behavioral pattern found in the analysis of the correct choice would

be associated with differences in the individual model parameters.

As a first step, we tested the parameters of our validated winning model for the 3-way interaction

effect of drug treatment, visibility, and type of recipient. There was no significant 3-way interaction in

the learning rate for positive PE (B = 1.03, CI = [1.00, 1.06], p = .071). The analysis of the learning

rate for negative PE revealed a three-way interaction of drug treatment, visibility, and type of

recipient (B = 1.07, CI = [1.03, 1.11], p = .002) so that the participants in the placebo group had a

relatively lower negative learning rate for prosocial choices when being watched than in privacy

(recipient x visibility interaction in the placebo group: B = 0.77, CI = [0.61, 0.98], p = .034).

Conversely, in the testosterone group, observation (vs privacy) relatively increased the negative

learning rate for prosocial choices (recipient x visibility interaction in testosterone group: B = 1.31, CI

= [1.04, 1.66], p = .021; Figure 2A). Moreover, the analysis of the choice consistency (inverse

temperature parameter tau, described also in the main text) likewise showed a three-way interaction

(B = 0.98, CI = [0.97, 0.99], p < .001). Placebo group participants had relatively higher consistency in

choices made for the other (vs. self) when being observed than in privacy (recipient x visibility

interaction in the placebo group: B = 1.08, CI = [1.05, 1.10], p < .001). On the contrary, in the

testosterone group, observation, compared to privacy, decreased the consistency of choices made for

the other (vs. self) (recipient x visibility interaction in testosterone group: B = 0.94, CI = [0.92, 0.97],

p < .001). When participants were observed, testosterone, compared to placebo, diminished the

relative consistency of prosocial choices (recipient x treatment interaction in observed condition: OR

= 0.87, CI = [0.85, 0.89], p = .001). In the private condition, there was no evidence for such an effect

(recipient x treatment interaction in private condition: OR = 0.99, CI = [0.97, 1.02], p = .605).

The analysis of the DDM threshold parameter revealed a three-way interaction as well (B= 1.01, CI =

[1.00, 1.02], p < .001; Figure 2C). Placebo group participants had a relatively higher threshold for

choices made for another (vs. self) when being observed than in privacy (recipient x visibility

interaction in the placebo group: B = 1.03, CI = [1.01, 1.05], p < .001). Conversely, in the testosterone

group, observation, compared to privacy, decreased the amount of information required for choices

made for another (vs. self) (recipient x visibility interaction in testosterone group: B = 0.95, CI =

[0.92, 0.98], p < .001). When participants were observed, testosterone, compared to placebo,

decreased the relative threshold of prosocial choices (recipient x treatment interaction in observed

condition: B = 0.95, CI = [0.92, 0.98], p < . 001). The analysis of the DDM drift-scaling parameter

revealed a three-way interaction (B = 0.83, CI = [0.82, 0.84], p < .001). Participants in both placebo

(recipient x visibility interaction in placebo group B = 0.92, CI = [0.91, 0.93], p < .001) and

testosterone group (recipient x visibility interaction in placebo group B = 0.77, CI = [0.76, 0.78], p

< .001) showed relatively lower drift scaling for choices made for another (vs. self) when being

observed than in privacy (recipient x visibility interaction in placebo group B = 0.92, CI = [0.91,

0.93], p < .001). When participants were observed, testosterone, compared to placebo, decreased the

relative drift scaling of prosocial choices (recipient x treatment interaction in the observed group: $B =$

0.93, $CI = [0.92, 0.94]$, p < .001).

As a second step, we examined whether the RLDDM parameters that were impacted by testosterone

administration predict behavioral prosociality, measured by the difference between correct choices

made for other and self across the whole sample. Out of the five parameters, negative learning rate (B

$= -8.82$, $CI = [-15.89, -1.75]$, $p < .015$), choice consistency ($B = 5.45$, $CI = [3.71, 7.19]$, $p < .001$), and

DDM threshold ($B = 9.56$, $CI = [0.33, 18.18]$, $p < .043$) predicted prosociality, however, only choice

consistency survived the Bonferroni correction for multiple comparisons ($p < .01$). Altogether, as

reported in the main text, these results suggest that testosterone's impact on strategic prosocial

behavior (i.e., audience effect) is strongly linked to testosterone's effect on choice consistency

(inverse temperature parameter tau).

1250

Analysis of the drift-scaling parameter and response times.

As specified in equation (7), on each trial t, the drift rate v_t was defined with a drift-scaling parameter,

$v_{scaling}$ that scales the value difference between the correct and incorrect symbol. Drift-scaling

parameter affects the curvature of the function: smaller values lead to a more linear mapping between

the value difference and the drift rate, and therefore less sensitivity to value differences.

Drift scaling is conceptually linked to the speed of integration and response times[15], we, therefore,

tested whether drift-scaling parameter predicted response times and found a significant association (B

$= 0.96$, $CI = [0.95, 0.98]$, $p < .001$). However, contrary to correct choices, response times did not

differ across experimental groups (drug treatment: $B = 1.01$, $CI = [0.97, 1.04]$, p < .706; visibility: $B =$

1.02, $CI = [0.99, 1.06]$, $p < .211$; recipient: $B = 0.99$, $CI = [0.98, 1.01]$, $p < .315$; drug treatment x

visibility x recipient: $B = 1.01$, $CI = [0.99, 1.01]$, $p < .841$).

1262

1263   **Supplementary information on the analysis of genetic data**

1264

Previous research suggested that testosterone may influence behavior through dopaminergic

pathways[25]. In humans, testosterone administration enhanced activation of the ventral striatum to

monetary rewards[26] and the enhancing effects of exogenous testosterone on competitive status-

seeking were more pronounced among individuals with a 9/10R compared to 10/10R genotype of the

dopamine transporter (DAT)[27]. The expression of DAT, which regulates striatal dopamine, is linked to

a 40 base-pair variable number tandem repeat polymorphism of the DAT1 gene[28]. Homozygous

10/10-repeat carriers of this polymorphism have higher DAT expression (i.e., lower striatal dopamine)

than heterozygous, 9-repeat variant, individuals[29].

Testosterone's effects on status-seeking behavior have likewise been shown to be enhanced among

individuals with fewer CAG repeats in exon 1 of the androgen-receptor gene[27,30]. In-vitro

experimental work suggests that increasing the number of CAG repeats within the androgen receptor

(AR) gene reduces the receptor's transcriptional potential[31]. In other words, the efficiency of the

androgen receptors is negatively related to the CAG repeat[32].

We, therefore, tested whether testosterone's effects on strategic prosociality depended on individual

differences in striatal dopamine, assessed by DAT1 polymorphism, and the efficiency of ARs,

assessed by the CAG repeat polymorphism.

1281

Genotyping of AR CAG repeat and DAT1 polymorphisms.

DNA was extracted from buccal swabs and isolated using a resin-based method with Chelex®100

(Sigma Aldrich, USA). For amplification of the CAG repeat polymorphism in exon 1 of the AR gene

primers forward - 5' GCGCGAAGTGATCCAGAAC 3' tagged with 6–carboxyfluorescein and

reverse - 5' CTCATCCAGGACCAGGTAGC 3', and for amplification of the DAT1-3'UTR

VNTR polymorphism primers forward - 5' GTCCTTGTGGTGTAGGGAAC 3' tagged with 6–

carboxyfluorescein and reverse - 5' CTGGAGGTCACGGCTCAAG 3' were used in PCR with 20 μL

reaction using 250 nmol/L final primer molarity. As PCR masterrmix 5x Hot FIREPol Blend

Mastermix with 7.5 mM MgCl2 (Solis Biodyne, Tartu, Estonia) was used in all amplifications.

The following PCR program was used: initial denaturation step at 95°C for 15 min, followed by 30

cycles each consisting of denaturation at 95°C for 30 s, annealing at 60°C for 30 s and polymerization

at 72°C for 1 min. The number of repeats of AR CAG STR and DAT1-3'UTR VNTR was analyzed

by fragment analysis using Sanger sequencing on ABI 3500 Genetic Analyzer (Applied Biosystems,

USA).

1296

Interaction of DAT1 polymorphism with testosterone effects on the correct choice and RLDDM

parameters.

There were no significant differences in the distribution of the genotype among our experimental

groups (χ^2 (6, $N = 187$) = 6.95, $p = .326$). The 9/10R and the 10/10R genotypes accounted for most of

the observed DAT1 genotypes in our sample (36% (N=66) and 56% (N=103), respectively), and we

thus used these two genotypes in the analyses by adding DAT1 polymorphisms as a predictor in

interaction with the other factors (recipient, drug treatment, visibility) to the GzLMM of correct

choice and the GzLMMs of the RLDDM parameters (see *Materials and Methods* in the main

manuscript text). The analysis revealed no significant interaction of DAT1 polymorphism with

testosterone effect on correct choice (recipient x drug treatment x visibility x DAT1: *OR* = 0.97, *CI* =

[0.91, 1.03], *p* = .307), α^{posPE} (recipient x drug treatment x visibility x DAT1: *B* = 1.17, *CI* = [0.70,

1.94], *p* = .555), α^{negPE} (recipient x drug treatment x visibility x DAT1: *B* = 0.58, *CI* = [0.29, 1.18], *p*

= .133), choice consistency (recipient x drug treatment x visibility x DAT1: *B* = 0.96, *CI* = [0.89,

1.03], *p* = .266) or decision threshold (recipient x drug treatment x visibility x DAT1: *B* = 1.00, *CI* =

[1.00, 1.00], *p* = .947).

1312

Interaction of AR CAG repeat polymorphism with testosterone effects on the correct choice and RLDDM parameters.

Mean-centered CAG repeat lengths of the AR gene in exon 1 were included as a predictor in interaction with the other factors (recipient, drug treatment, visibility) to the GzLMM of correct choice and the GzLMMs of the RLDDM parameters (see *Materials and Methods* in the main text). The analysis revealed no significant interaction of CAG repeat polymorphism with testosterone effect on correct choice (recipient x drug treatment x visibility x CAG: $OR = 1.00$, $CI = [0.99, 1.02]$, $p = .769$), α^{posPE} (recipient x drug treatment x visibility x CAG: $B = 1.00$, $CI = [1.00, 1.01]$, $p = .329$), αneg^{PE} (recipient x drug treatment x visibility x CAG: $B = 1.01$, $CI = [1.00, 1.02]$, $p = .092$), choice consistency (recipient x drug treatment x visibility x CAG: $B = 1.00$, $CI = [0.99, 1.01]$, $p = .696$) or decision threshold (recipient x drug treatment x visibility x CAG: $B = 1.01$, $CI = [1.00, 1.02]$, $p = .122$).

Interaction of trait dominance with testosterone effects on RLDDM parameters.

Mean-centered dominance scores[33] were included as a predictor in interaction with the other factors (recipient, drug treatment, visibility) to the GzLMM of correct choice (reported in the main text) and the GzLMMs of the RLDDM parameters. Contrary to the former, the latter analysis revealed no significant interaction of dominance scores with testosterone effect RLDDM parameters: learning rate for positive PE (recipient x drug treatment x visibility x dominance: $B = 1.01$, $CI = [0.98, 1.04]$, $p = .546$), learning rate for negative PE (recipient x drug treatment x visibility x dominance: $B = 1.02$, $CI = [0.98, 1.07]$, $p = .341$), choice consistency (recipient x drug treatment x visibility x dominance: $B = 1.00$, $CI = [1.00, 1.01]$, $p = .229$), or decision threshold (recipient x drug treatment x visibility x dominance: $B = 1.00$, $CI = [1.00, 1.00]$, $p = .492$).

Supplementary information on the questionnaire data

Post-task questionnaire.

To estimate the subjective perception of being watched, a post-task questionnaire was administered immediately after the end of the reinforcement-learning paradigm. The participants were asked the question: "Did you feel that you were being watched while performing the task?" The answers were classified into three categories: 1 (Not at all), 2 (Moderately), and 3 (Strongly). First, we investigated whether the subjective feelings of being watched were related to the cortisol reactivity to visibility manipulation (see *Manipulation check* in the manuscript's main text). We next examined whether the subjective feelings of being watched interact with the testosterone administration effect on prosocial choice and RLDDM parameters. As the factors *visibility* and *subjective feeling of being watched* are

not orthogonal, we did not add the subjective feelings of being watched as a fourth factor to the

analysis. Instead, we replaced the factor *visibility* with the factors *subjective feeling of being watched*

and compared whether such a model explains more variance than the original one. We did not find a

significant interaction of the factors administration x recipient x subjective feeling of being watched

($OR = 0.96$, $CI = [0.89, 1.03]$, $p = .269$). This model did not explain more variance (*Conditional*

R2=.101) than the original administration x recipient x visibility model (*Conditional R2*=.102).

1354

Portrait Values Questionnaire.

We conducted exploratory analyses to examine what motivational constructs may have interacted

with testosterone's effects. To this end, we analyzed data from the Portrait Values Questionnaire[33,34]

which had been administered as part of the study's extensive questionnaire battery. Note that these are

exploratory analyses and that this specific questionnaire was originally intended to be used for

analyses related to other tasks that were part of the overall study. They should therefore be considered

with caution, and we have clearly labeled them as exploratory and post-hoc. In brief, the Portrait

Values Questionnaire is designed to capture the structure of human values explaining the motivational

bases of our attitudes and behavior. The questionnaire allowed us to distinguish four principal value

orientations: self-transcendence, self-enhancement, conservation, and openness. The four value

subscales were separately added to our analysis of recipient x administration x visibility interaction as

mean-centered interaction terms. Only conservation value significantly (Bonferroni adjusted α level

= .0125) interacted with testosterone effects on choice consistency (recipient x visibility x

administration x conservation: $B = 0.05$, $CI = [0.02, 0.09]$, $p = .001$), the follow-up analysis is

reported in the main manuscript text. Conservation did not significantly interact with testosterone

effects on correct choice (recipient x drug treatment x visibility x conservation: $OR = 1.03$, $CI =$

$[0.98, 1.07]$, $p = .220$). Self-transcendence did not significantly interact with testosterone effects on

correct choice (recipient x drug treatment x visibility x self-transcendence: $OR = 0.97$, $CI = [0.91,$

$1.03]$, $p = .265$) or choice consistency (recipient x drug treatment x visibility x self-transcendence: B

$= 0.04$, $CI = [-0.01, 0.09]$, $p = .102$). Similarly, self-enhancement value did not significantly interact

with testosterone effects on correct choice (recipient x drug treatment x visibility x self-enhancement:

$OR = 1.03$, $CI = [0.99, 1.07]$, $p = .126$) or choice consistency (recipient x drug treatment x visibility x

self-enhancement: $B = -0.01$, $CI = [-0.04, 0.02]$, $p = .598$). Lastly, openness value did not

significantly interact with testosterone effects on correct choice (recipient x drug treatment x visibility

x openness: $OR = 1.02$, $CI = [0.97, 1.06]$, $p = .453$) or choice consistency (recipient x drug treatment

x visibility x openness: $B = 0.06$, $CI = [-0.22, 0.33]$, $p = .693$).

1381

Observers' salience survey.

To estimate the salience of the observers introduced as NGO representatives in our paradigm, we

conducted an additional online survey. Using Amazon Mechanical Turk (MTruk) we interviewed

N = 73 volunteers (sample size determined by G*Power[34] to achieve 95 % power to detect a small

effect size f = .20) aged M = 34.55 (SD = 9.12) years. Participants were asked to perform the task

alone and rate how much they agree with the statements regarding different types of observers.

Together they rated 7 statements with the following wording, and each statement concerned a

different type of observer:

"As I perform this task on my computer/mobile device alone, I anticipate I would feel

differently if ……… would watch me perform this task."

The observer's types were: an unknown other, a friend, a parent, a shop assistant, an NGO

representative, a colleague from work, and a technical support worker. A repeated-measures

ANOVA comparing the ratings of the respective observers showed that there was a difference

across the levels of the within-subject factor $F(6, 396)$ = 2.69, p = .014, f =.204. A follow-up

analysis using treatment contrasts and the NGO as a reference level showed a significant

difference between the levels of NGO representative and parent (B = -0.57, p < 0.001), meaning

that participants would feel different to a greater extent if they were watched by an NGO

representative than a parent. Crucially, no other pairwise comparisons were significant, all ps

>.169)."

1401

1402

1403

1404

1405

1406

1407

1408

1409

1410

1411

1412

1413

1414

1435 **Figure S1.** Testosterone levels during the experimental session. Error bars = Mean ± 95%CI.
1436

1437   **Table S1.** Summary statistics across experimental groups: Mean (95%CI), ANOVA.

Drug treatment	Placebo		Testosterone			
Visibility	**Private**	**Observed**	**Private**	**Observed**	**F**	**p**
N	46	45	52	44		
Age	25.2 (24.0, 26.4)	24.5 (23.3, 25.7)	24.4 (23.3, 25.5)	25.2 (24.0, 26.4)	1.141	.236
Baseline testosterone [pg/mL]	143 (76.7, 209)	131 (64.5, 197)	221 (150.2, 293)	261 (172.1, 350)		
log	4.75 (4.52, 4.98)	4.71 (4.48, 4.94)	4.92 (4.67, 5.16)	4.97 (4.66, 5.28)	0.128	.721
Baseline cortisol [nmol/L]	2.84 (1.81, 3.87)	3.69 (2.66, 4.72)	3.92 (2.96, 4.88)	3.57 (2.52, 4.61)		
log	0.77 (0.52, 1.02)	1.00 (0.75, 1.25)	0.91 (0.68, 1.15)	0.86 (0.61, 1.11)	1.260	.263
Baseline estradiol [pg/mL]	3.68 (3.12, 4.23)	3.73 (3.17, 4.29)	3.77 (3.25, 4.30)	3.73 (3.16, 4.29)		
log	1.23 (1.09, 1.36)	1.19 (1.05, 1.33)	1.21 (1.08, 1.33)	1.22 (1.08, 1.36)	0.112	.739
CAG-repeat polymorphism	19.6 (18.5, 20.7)	19.9 (18.9, 21.0)	19.4 (18.4, 20.4)	19.4 (18.3, 20.4)	0.083	.773
Dominance	3.89 (3.62, 4.17)	4.02 (3.74, 4.30)	4.05 (3.79, 4.31)	3.84 (3.56, 4.12)	1.432	.232

1438
1439
1440

Table S2. Model space and model evidence.

RW, Rescorla-Wagner model; DLR dual learning rates model; DDM, drift diffusion model;

RLDDM, reinforcement learning drift diffusion model; RLDDM-nonlinear, RLDDM with a non-

linear transformation function; LOOIC leave-one-out information criterion (lower LOOIC value

indicates better out-of-sample predictive accuracy); weight, model weight calculated with Bayesian

model averaging using Bayesian bootstrap (higher model weight value indicates a higher probability

of the candidate model to have generated the observed data). The winning model is highlighted in

bold. Because LOOIC provides relative quantifications of model performance, and the data to be

modeled varied across conditions, the absolute magnitude of LOOIC should not be interpreted.

1450

Model space	Task condition	Placebo Observed		Placebo Private		Testosterone observed		Testosterone Private	
		LOOIC	Weight	LOOIC	Weight	LOOIC	Weight	LOOIC	Weight
RW	DDM	9688	0	10697	0	9991	0	12427	0
	RLDDM	8365	0	9522	0	8925	0	11170	0
	RLDDM-nonlinear	8235	0	9395	0	8782	0	10903	0
DLR	DDM	9538	0	10493	0	9813	0	12199	0
	RLDDM	7878	0.006	8868	0.053	8444	0.002	10425	0.001
	RLDDM-nonlinear	**7832**	**0.994**	**8837**	**0.947**	**8352**	**0.998**	**10261**	**0.999**

1451

1452

1453 **Table S3.** Summary of parameters in the winning model.

1454 Note that all parameters were further separated for "self" versus "other", hence for each between-

1455 subject condition, the winning model contained 14 parameters. The initial bias z in DDM was fixed at

1456 0.5. min(RT), lowest response time from observed data.

1457

Component	Parameter	Meaning	Interpretation
RL	$0 < \alpha^{posPE} < 1$	learning rate for updating from positive reward prediction error	the learning rate weighs the effect of the reward prediction error in the value update; a higher (lower) learning rate means a faster (slower) value update from the most recent outcome
	$0 < \alpha^{negPE} < 1$	learning rate for updating from negative reward prediction error	
	$\beta > 0$	inverse temperature in softmax action selection function	choice consistency parameter, captures how much choices rely on the value updates
DDM	$v_{max} > 0$	maximum value of the drift rate in the non-linear function	defines the upper/lower boundaries in the sigmoid non-linear function
	$v_{scaling} > 0$	drift scaling that maps value difference into the drift rate	scales the effect of value difference between the choice options on the drift rate
	$a > 0$	decision threshold, i.e., the distance between choice alternatives	"the endpoint" of the evidence accumulation process, captures the amount of information necessary to make a decision
	$0 < T < min(RT)$	non-decision time	considered to capture sensory delay and/or movement initiation

1458
1459

SI References

1. R Core Team. 2020. R: A Language and Environment for Statistical Computing. Vienna, Austria: R Foundation for Statistical Computing.

2. Bates D, Mächler M, Bolker B, Walker S. 2015. Fitting Linear Mixed-Effects Models Using lme4. *Journal of Statistical Software* **67**:1–48. doi:10.18637/jss.v067.i01

3. Fox J, Weisberg S. 2019. An R Companion to Applied Regression, Third. ed. Thousand Oaks CA: Sage.

4. Lüdecke D. 2020. sjPlot: Data Visualization for Statistics in Social Science.

5. Phillips N. 2017. yarrr: A Companion to the e-Book "YaRrr!: The Pirate's Guide to R."

6. Wickham H. 2016. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New York.

7. Diedenhofen B, Musch J. 2015. cocor: A Comprehensive Solution for the Statistical Comparison of Correlations. *PLoS One* **10**:e0121945. doi:10.1371/journal.pone.0121945

8. Carpenter B, Gelman A, Hoffman MD, Lee D, Goodrich B, Betancourt M, Brubaker M, Guo J, Li P, Riddell A. 2017. Stan: A Probabilistic Programming Language. *Journal of Statistical Software* **76**:1–32. doi:10.18637/jss.v076.i01

9. Ahn W-Y, Haines N, Zhang L. 2017. Revealing Neurocomputational Mechanisms of Reinforcement Learning and Decision-Making With the hBayesDM Package. *Comput Psychiatr* **1**:24–57. doi:10.1162/CPSY_a_00002

10. Vehtari A, Gelman A, Gabry J. 2017. Practical Bayesian model evaluation using leave-one-out cross-validation and WAIC. *Statistics and Computing*, **27**, 1413–1432. doi: 10.1007/s11222-016-9696-4

11. Knight EL, McShane BB, Kutlikova HH, Morales PJ, Christian CB, Harbaugh WT, Mayr U, Ortiz TL, Gilbert K, Ma-Kellams C, Riečanský I, Watson NV, Eisenegger C, Lamm C, Mehta PH, Carré JM. 2020. Weak and Variable Effects of Exogenous Testosterone on Cognitive Reflection Test Performance in Three Experiments: Commentary on Nave, Nadler, Zava, and Camerer (2017). *Psychol Sci* **31**:890–897. doi:10.1177/0956797619885607

12. Nave G, Nadler A, Zava D, Camerer C. 2017. Single-Dose Testosterone Administration Impairs Cognitive Reflection in Men. *Psychol Sci* **28**:1398–1407. doi:10.1177/0956797617709592

13. R. A. Rescorla, A. R. Wagner, "A theory of Pavlovian conditioning: Variations in the effectiveness of reinforcement and nonreinforcement" in *Classical Conditioning II: Current Research and Theory*, (Appleton-Century-Crofts, 1972), pp. 64–99.

14. den Ouden HEM, Daw ND, Fernandez G, Elshout JA, Rijpkema M, Hoogman M, Franke B, Cools R. 2013. Dissociable effects of dopamine and serotonin on reversal learning. *Neuron* **80**:1090–1100. doi:10.1016/j.neuron.2013.08.030

15. Ratcliff R, McKoon G. 2008. The Diffusion Decision Model: Theory and Data for Two-Choice Decision Tasks. *Neural Comput* **20**:873–922. doi:10.1162/neco.2008.12-06-420

16. Navarro DJ, Fuss IG. 2009. Fast and accurate calculations for first-passage times in Wiener diffusion models. *Journal of Mathematical Psychology* **53**:222–230. doi:10.1016/j.jmp.2009.02.003

17. Pedersen ML, Frank MJ, Biele G. 2017. The drift diffusion model as the choice rule in reinforcement learning. *Psychon Bull Rev* **24**:1234–1251. doi:10.3758/s13423-016-1199-y

18. Fontanesi L, Gluth S, Spektor MS, Rieskamp J. 2019. A reinforcement learning diffusion decision model for value-based decisions. *Psychon Bull Rev* **26**:1099–1121. doi:10.3758/s13423-018-1554-2

19. Zhang L, Gläscher J. 2020. A brain network supporting social influences in human decision-making. *Science Advances* **6**:eabb4159. doi:10.1126/sciadv.abb4159

20. Gelman A, Carlin JB, Stern HS, Dunson DB, Vehtari A, Rubin DB. 2015. Bayesian Data Analysis, 3rd ed. New York: Chapman and Hall/CRC. doi:10.1201/b16018

21. Gelman A, Rubin DB. 1992. Inference from Iterative Simulation Using Multiple Sequences. *Statistical Science* **7**:457–472.

22. Vehtari A, Gelman A, Gabry J. 2016. Practical Bayesian model evaluation using leave-one-out cross-validation and WAIC. *arXiv:150704544 [stat]*. doi:10.1007/s11222-016-9696-4

23. Yao Y, Vehtari A, Simpson D, Gelman A. 2018. Using Stacking to Average Bayesian Predictive Distributions (with Discussion). *Bayesian Analysis* **13**:917–1007. doi:10.1214/17-BA1091

24. Zhang L, Lengersdorff L, Mikus N, Gläscher J, Lamm C. 2020. Using reinforcement learning models in social neuroscience: frameworks, pitfalls and suggestions of best practices. *Social Cognitive and Affective Neuroscience* **15**:695–707. doi:10.1093/scan/nsaa089

25. Purves-Tyson TD, Owens SJ, Double KL, Desai R, Handelsman DJ, Weickert CS. 2014. Testosterone Induces Molecular Changes in Dopamine Signaling Pathway Molecules in the Adolescent Male Rat Nigrostriatal Pathway. *PLoS One* **9**:e91151. doi:10.1371/journal.pone.0091151

26. Hermans EJ, Bos PA, Ossewaarde L, Ramsey NF, Fernández G, van Honk J. 2010. Effects of exogenous testosterone on the ventral striatal BOLD response during reward anticipation in healthy women. *Neuroimage* **52**:277–283. doi:10.1016/j.neuroimage.2010.04.019

27. Losecaat Vermeer AB, Krol I, Gausterer C, Wagner B, Eisenegger C, Lamm C. 2020. Exogenous testosterone increases status-seeking motivation in men with unstable low social status. *Psychoneuroendocrinology* **113**:104552. doi:10.1016/j.psyneuen.2019.104552

28. Vandenbergh DJ, Persico AM, Hawkins AL, Griffin CA, Li X, Jabs EW, Uhl GR. 1992. Human dopamine transporter gene (DAT1) maps to chromosome 5p15.3 and displays a VNTR. *Genomics* **14**:1104–1106. doi:10.1016/s0888-7543(05)80138-7

29. Heinz A, Goldman D, Jones DW, Palmour R, Hommer D, Gorey JG, Lee KS, Linnoila M, Weinberger DR. 2000. Genotype Influences In Vivo Dopamine Transporter Availability in Human Striatum. *Neuropsychopharmacol* **22**:133–139. doi:10.1016/S0893-133X(99)00099-8

30. Geniole SN, Procyshyn TL, Marley N, Ortiz TL, Bird BM, Marcellus AL, Welker KM, Bonin PL, Goldfarb B, Watson NV, Carré JM. 2019. Using a Psychopharmacogenetic

1570 Approach To Identify the Pathways Through Which—and the People for Whom—
1571 Testosterone Promotes Aggression. *Psychol Sci* **30**:481–494.
1572 doi:10.1177/0956797619826970

1573

1574 31. Chamberlain NL, Driver ED, Miesfeld RL. 1994. The length and location of CAG
1575 trinucleotide repeats in the androgen receptor N-terminal domain affect transactivation
1576 function. *Nucleic Acids Res* **22**:3181–3186. doi:10.1093/nar/22.15.3181

1577

1578 32. Zitzmann M, Nieschlag E. 2003. The CAG repeat polymorphism within the androgen
1579 receptor gene and maleness. *Int J Androl* **26**:76–83. doi:10.1046/j.1365-2605.2003.00393.x

1580

1581 33. Schwartz SH. 2007. Chapter 7 A Proposal for Measuring Value Orientations across
1582 Nations.

1583

1584 34. Schwartz, SH, Breyer, B, & Danner, D. 2015. Human Values Scale
1585 (ESS).*Zusammenstellung sozialwissenschaftlicher Items und Skalen (ZIS)*.
1586 doi:10.6102/zis234

1587

1588 35. Faul F, Erdfelder E, Lang A-G, Buchner A. 2007. G*Power 3: a flexible statistical power
1589 analysis program for the social, behavioral, and biomedical sciences. *Behav Res Methods*
1590 **39**:175–191. doi:10.3758/bf03193146

1591

1592

1593

1594