The single-cell atlas of the murine reproductive tissues during preterm labor

Valeria Garcia-Flores[1,2], Roberto Romero[1,3-7,]*, Azam Peyvandipour[1-3], Jose Galaz[1,2],

Errile Pusod[1,2], Bogdan Panaitescu[1,2], Derek Miller[1,2], Yi Xu[1,2], Li Tao[1,2], Zhenjie Liu[1,2],

Adi L. Tarca[1,2,8], Roger Pique-Regi[1-3,]*, Nardhy Gomez-Lopez[1,2,9,10,]*

[1]Perinatology Research Branch, Division of Obstetrics and Maternal-Fetal Medicine, Division of

Intramural Research, *Eunice Kennedy Shriver* National Institute of Child Health and Human

Development, National Institutes of Health, U.S. Department of Health and Human Services

(NICHD/NIH/DHHS); Bethesda, Maryland, and Detroit, Michigan, USA.

[2]Department of Obstetrics and Gynecology, Wayne State University School of Medicine;

Detroit, Michigan, USA.

[3]Center for Molecular Medicine and Genetics, Wayne State University; Detroit, Michigan, USA.

[4]Department of Obstetrics and Gynecology, University of Michigan; Ann Arbor, Michigan,

USA.

[5]Department of Epidemiology and Biostatistics, Michigan State University; East Lansing,

Michigan, USA.

[6]Center for Molecular Medicine and Genetics, Wayne State University; Detroit, Michigan, USA.

[7]Detroit Medical Center; Detroit, Michigan, USA.

[8]Department of Computer Science, Wayne State University College of Engineering; Detroit,

Michigan, USA.

[9]Department of Biochemistry, Microbiology, and Immunology, Wayne State University School

of Medicine; Detroit, Michigan, USA.

[10]Lead contact

*Correspondence

nardhy.gomez-lopez@wayne.edu

prbchiefstaff@med.wayne.edu

rpique@wayne.edu

SUMMARY

Preterm birth, the leading cause of perinatal morbidity and mortality worldwide,

frequently results from the syndrome of preterm labor. Intra-amniotic infection is the best-

established causal link to preterm labor, and involves premature activation of the parturition

cascade in the reproductive tissues. Herein, we utilized single-cell RNA-sequencing (scRNA-

seq) to generate a single-cell atlas of the murine uterus, decidua, and cervix in a model of

infection-induced preterm labor. We show that preterm labor affects the transcriptomic profiles

of specific immune and non-immune cell subsets. Shared and tissue-specific gene expression

signatures were identified among affected cells. Importantly, determination of intercellular

communications implicates specific cell types preterm labor-associated signaling pathways

across tissues. Last, *in silico* comparison of murine and human uterine cell-cell interactions

reveals conserved signaling pathways implicated in labor. Thus, scRNA-seq provides new

insights into the preterm labor-driven cellular landscape and communications in the reproductive

tissues.

KEYWORDS: single cell, scRNA-seq, decidua, uterus, cervix, mouse, human

INTRODUCTION

Preterm birth is a devastating clinical condition that affects 15 million infants each year and is the leading cause of neonatal morbidity and mortality worldwide (Chawanpaiboon et al., 2019; Liu et al., 2015b). Spontaneous preterm birth often results from preterm labor, a syndrome for which multiple etiologies have been proposed (Goldenberg et al., 2008; Romero et al., 2014a). Among these, the best-established causal link to preterm birth is intra-amniotic infection, a clinical condition resulting from the invasion of microbes into the amniotic cavity (Agrawal and Hirsch, 2012; Bastek et al., 2011; Combs et al., 2014; Gibbs et al., 1992; Gomez et al., 1995; Gonçalves et al., 2002; Keelan et al., 2003; Romero et al., 2019; Theis et al., 2020). The most frequently detected bacteria in amniotic fluid of women diagnosed with intra-amniotic infection include genital mycoplasmas, *Streptococcus agalactiae*, *Gardnerella vaginalis*, and *Escherichia coli*, among others (Burnham et al., 2020; DiGiulio et al., 2010; Gibbs et al., 1982; Gravett et al., 1986; Oh et al., 2019; Romero et al., 2015a; Romero et al., 2014b; Romero et al., 2015b; Romero et al., 2015c; Romero et al., 1989; Yoon et al., 1998; Yoon et al., 2019). Human descriptive studies have consistently shown that such microbial invasion of the amniotic cavity is accompanied by a local acute inflammatory response that includes the infiltration of leukocytes into the amniotic cavity (inc. amniotic fluid (Galaz et al., 2020a; Galaz et al., 2020b; Gomez-Lopez et al., 2019a; Gomez-Lopez et al., 2019b; Gomez-Lopez et al., 2021b; Gomez-Lopez et al., 2017b; Gomez-Lopez et al., 2019c; Gomez-Lopez et al., 2018b; Gomez et al., 1994; Martinez-Varea et al., 2017; Romero et al., 1991; Romero et al., 1993a; Romero et al., 1993b; Yoon et al., 1996) and placental tissues (Gomez-Lopez et al., 2017a; Guzick and Winn, 1985; Hillier et al., 1988; Kim et al., 2015a; Kim et al., 2009; Kim et al., 2015b; Odibo et al., 1999; Oh et al., 2021; Oh et al., 2011; Olding, 1970; Pacora et al., 2002; Park et al., 2009; Redline, 2012;

Redline et al., 2003; Svensson et al., 1986; van Hoeven et al., 1996; Yoon et al., 2001; Zhang et al., 1985)) as well as the reproductive tissues (Keski-Nisula et al., 2003; Makieva et al., 2017). More recently, animal models coupled with omics technologies have been utilized to strengthen this concept and establish causality between intra-amniotic infection and the inflammatory milieu observed in the reproductive tissues (e.g., uterus, decidua, and cervix) that serve to orchestrate the premature activation of the common pathway of labor (Migale et al., 2016; Motomura et al., 2020a; Toothaker et al., 2020; Willcockson et al., 2018). However, the simultaneous investigation of the cellular landscape and interaction networks at single-cell resolution in the reproductive tissues implicated in preterm parturition has not been undertaken.

Single-cell technology has emerged as a useful tool for evaluating cellular composition, transcriptomic activity, and communication networks in gestational and reproductive tissues (Huang et al., 2021; Nelson et al., 2016; Pavličev et al., 2017; Pique-Regi et al., 2019; Suryawanshi et al., 2018; Tsang et al., 2017; Vento-Tormo et al., 2018). Indeed, we have applied single-cell RNA-sequencing (scRNA-seq) to investigate the physiological and pathological processes of labor in the placenta and extraplacental membranes (Pique-Regi et al., 2019). More recently, we utilized scRNA-seq to unravel the myometrial cell types that participate in the normal process of term parturition as well as key cell-cell interactions taking place in this compartment (Pique-Regi et al., 2022). Importantly, the discovery of single-cell signatures derived from the placental tissues and myometrium possesses translational value as these can serve as potential non-invasive biomarkers of labor progression and/or obstetrical disease (Gomez-Lopez et al., 2022; Pique-Regi et al., 2022; Pique-Regi et al., 2019; Tarca et al., 2021; Tarca et al., 2019; Tsang et al., 2017).

90   In the current study, we utilized scRNA-seq coupled with an allogeneic murine model of

91 intra-amniotic infection to investigate the cellular landscape and cell-cell communications in the

92 reproductive tissues (uterus, decidua, and cervix) during the process of preterm labor. We

93 utilized a murine model of preterm labor and birth induced by the intra-amniotic inoculation of

94 *E. coli*, and assessed cervical shortening to establish the timing of active preterm labor. Next,

95 using scRNA-seq and computational approaches, we generated a single-cell atlas of the uterus,

96 decidua, and cervix during preterm labor as well as their cell type-specific transcriptomic

97 activity. In addition, we established the cell-cell communication networks between cell types in

98 each tissue during preterm labor and identified key signaling pathways implicated in this process.

99 Last, we integrated cell-cell signaling pathways derived from the murine uterus with those from

100 the human myometrium during the processes of preterm and term labor, respectively, to

101 demonstrate conserved labor-associated signaling.

RESULTS

A single-cell atlas of the murine reproductive tissues during preterm labor induced by *E.*

coli

Intra-amniotic infection has been documented as inducing inflammatory changes in the

tissues surrounding the amniotic cavity (Faro et al., 2019; Motomura et al., 2021; Motomura et

al., 2020b; Senthamaraikannan et al., 2016). To investigate such changes at single-cell

resolution, we first established a murine model of preterm labor and birth induced by the intra-

amniotic inoculation with *E. coli*, one of the microorganisms commonly identified in the

amniotic fluid of women with intra-amniotic infection (Gibbs et al., 1982; Romero et al., 2015c;

Romero et al., 1989; Yoneda et al., 2016a). Mice with an allogeneic pregnancy underwent

ultrasound-guided intra-amniotic injection of *E. coli* or vehicle control on 16.5 days *post coitum*

(dpc) (Figure 1A). Intra-amniotic inoculation with *E. coli* reduced gestational length in the

majority of dams (Figure 1B), resulting in an 83.3% (5/6) rate of preterm birth (Figure 1C). We

then intra-amniotically injected a second cohort of mice with *E. coli* or PBS to perform tissue

collection for single-cell analyses. To ensure that the *E. coli*-injected mice were undergoing

preterm labor at the time of tissue collection, we utilized ultrasound to evaluate cervical length

just prior to intra-amniotic injection and again 24 h later as a readout of cervical effacement

(Figure 1D). Cervical shortening was observed in all dams that received intra-amniotic *E. coli* at

24 h post-injection, indicating that these dams were in active preterm labor at the time of tissue

collection, whereas no cervical shortening was observed in controls (Figure 1E). Therefore,

intra-amniotic inoculation with *E. coli* represents a translational model that resembles the clinical

scenario of intra-amniotic infection leading to preterm labor and birth.

Preterm parturition includes the activation of the common pathway of labor that

comprises increased uterine contractility, the triggering of local immune response in the decidual

tissues, and cervical dilatation (Lopez Bernal, 2003; Norwitz et al., 1999; Romero et al., 2014a;

Romero et al., 2006; Romero et al., 1994; Smith, 2007). Therefore, to establish a single-cell atlas

of the murine reproductive tissues in preterm labor, we utilized the uterus, decidua, and cervix of

dams that received intra-amniotic inoculation with *E. coli* in the active phase of parturition (17.5

dpc) for single-cell RNA-sequencing (scRNA-seq) (Figure 1F). We identified 31 cell clusters

across the uterus, decidua, and cervix that corresponded to multiple cell types: smooth muscle (2

clusters), epithelial (10 clusters), fibroblast (3 clusters), stromal (3 clusters), Endothelial,

Neutrophil, Monocyte, macrophage (2 clusters), Dendritic Cell, T cell, B cell, NK cell (2

clusters), Erythroid, Plasmocyte, and Trophoblast (Figure 1G). The heterogeneous and distinct

cellular composition of the uterus, decidua, and cervix was highlighted by assigning tissue

identity to each cell cluster (Figure 1H). In control dams, the uterus, decidua, and cervix each

displayed a distinct basal cellular repertoire: the uterus showed a predominance of fibroblast

(clusters 0 and 1) and non-decidual stromal (clusters 2 and 12) cell types, and the decidua also

included a unique subset of stromal cells (cluster 4) (likely corresponding to conventional

decidual stromal cells) (Figure S1A). The uterus and decidua of control mice also included

modest populations of innate immune cells such as Monocyte and macrophage subsets as well as

lymphocytes such as T cell, NK cell-1, NK cell-2, and B cell (Figure S1A), likely representing

the resident immune populations that have been characterized in human and murine tissues

(Bartmann et al., 2014; Li et al., 2018; Pique-Regi et al., 2022; Pique-Regi et al., 2019; Trundley

and Moffett, 2004; Vento-Tormo et al., 2018). By contrast with the uterus and decidua, the

cervix of control mice comprised a diverse compartment of epithelial subsets (clusters 5, 7, 8, 10,

11, 14) and other major cell types (Figure S1A), as previously shown (Chumduri et al., 2021;

Koh et al., 2019; Zhao et al., 2021). Immune cells were scarce in the cervix, although a modest

Macrophage-1 population was observed (Figure S1A) that is consistent with prior reports of

cervical cell composition in late gestation (Dobyns et al., 2015; Dubicke et al., 2016; Osman et

al., 2003; Timmons et al., 2009). Together, these data provide an overview of the single-cell

composition and diversity in the murine uterus, decidua, and cervix in late murine pregnancy.

Moreover, by dissecting the cellular repertoire of the cervix we demonstrate the

underappreciated heterogeneity of this compartment.

Preterm labor induced by *E. coli* dysregulates the repertoire of immune and non-immune

cell types in the reproductive tissues

We then examined the effects of preterm labor on the abundance of each cell type

identified across all tissues (Figure 2A) as well as within the uterus, decidua, and cervix (Figure

2B-D, Figure S1B, and Table S1). During preterm labor, a relative increase in innate immune

cell clusters such as Monocytes, macrophages, Dendritic cells, and Neutrophils (clusters 3, 6, 9,

and 18) was observed in the uterus, decidua, and, to a lesser extent, the cervix (Figure 2B-D and

Figure S1B). The NK cell-2 and Plasmocyte subsets in the uterus and decidua also showed

changes with preterm labor (Figure 2B-C and Figure S1B). Moreover, Dendritic cells were

increased in the decidua (Figure 2C and Figure S1B), with a similar tendency observed in the

uterus (Figure 2B and Figure S1B). Interestingly, the Macrophage-1 cell type was decreased in

the uterus with preterm labor (Figure 2B and Figure S1B). Notably, the T-cell population (cluster

16) also appeared to increase in the uterus and decidua with preterm labor (Figure 2B&C and

Figure S1B), which is consistent with prior studies implicating T-cell infiltration and activation

as a component of parturition (Arenas-Hernandez et al., 2019; Gomez-Lopez et al., 2009;

Gomez-Lopez et al., 2011a; Gomez-Lopez et al., 2011b; Gomez-Lopez et al., 2013; Sindram-

Trujillo et al., 2003; Slutsky et al., 2019; Vargas et al., 1993). Although not visually apparent

from the UMAP plots (Figure S1B), both the uterus and decidua showed a substantial decrease in

non-immune subsets such as Fibroblast-1, -2, and Stromal-3 with preterm labor (Figure 2B&C),

with Stromal-2 also showing modest changes in the uterus (Figure 2B). Interestingly, a subset of

epithelial cells (cluster 11, Epithelial-10) that was largely absent in the uterus and decidua of

controls became apparent in preterm labor (Figure S1B), suggesting labor-induced differentiation

or activation of these cells. By contrast with the uterus and decidua, the cervix only showed

changes in two cell types: Neutrophil and Epithelial-8 were both increased with preterm labor

(Figure 2D), indicating that a modest cellular response to intra-amniotic infection occurs in this

tissue. We also evaluated whether cells of fetal origin were represented among the populations of

the uterus, decidua, and cervix during preterm labor (Figure S1C-E). A small population of fetal

cells (Trophoblasts) was detected in the uterus and decidua, which is consistent with prior single-

cell studies of the human myometrium (Pique-Regi et al., 2022) and may represent residual

placental cells attached to the uterus and decidua.

To validate the leukocyte infiltration of the uterus, decidua, and cervix indicated by our

single-cell data, we undertook a series of histological and immunohistochemistry analyses

(Figure S2). We observed collagen degradation in the uterine and cervical tissues with preterm

labor, and mucin production by cervical cells appeared to increase compared to control tissues

(Figure S2A-C). Histological changes in preterm labor were accompanied by increased CD45+

leukocyte infiltration in the uterus and decidua (Figure S2D-F). Uterine leukocytes were more

evenly distributed among neutrophils, monocytes, and macrophages, whereas decidual

leukocytes were predominantly neutrophils and, to a lesser extent, monocytes (Figure S2G-I).

Similar to our scRNA-seq results, the leukocyte abundance in the cervical tissues was largely

comparable between the control and preterm labor groups (Figure S2I).

Taken together, our scRNA-seq indicate a shift in the cellular composition of the murine

uterus, decidua, and cervix that accompanies preterm labor.

Preterm labor induced by *E. coli* dysregulates the gene expression of immune and non-

immune cell types in the reproductive tissues

Given that preterm labor altered the cellular composition of the uterus, decidua, and

cervix, we next explored whether this inflammatory process also resulted in transcriptomic

changes to the identified cell types. Consistent with their altered abundance, multiple fibroblast,

stromal, and epithelial cell types in the uterus and decidua displayed upregulated gene expression

with preterm labor (Figure 2E&F and Table S2), while the cervical non-immune cells with

upregulated gene expression were exclusively epithelial (Figure 2G and Table S2). Interestingly,

innate immune cell types showed strong dysregulation of gene expression in both directions that

was inconsistent among tissues: while Monocyte showed more downregulated differentially

expressed genes (DEGs) in the uterus (Figure 2E) and cervix (Figure 2G), this cell type showed

more upregulated DEGs in the decidua (Figure 2F). Neutrophil showed stronger downregulation

of DEGs in the uterus and decidua (Figure 2E&F), whereas in the cervix DEGs were primarily

upregulated in this cell type (Figure 2G). Macrophage-1, Dendritic cell, and NK cell-1

consistently displayed predominantly upregulated DEGs in the uterus and decidua (Figure

2E&F), and were not represented in the cervix as previously noted. In addition, the uterine

Macrophage-2 and NK cell-2 populations displayed upregulated DEGs (Figure 2E), which was

not observed in other tissues (Figure 2F&G). Although not as abundant as innate immune cells,

the T cell population also displayed upregulated DEGs with preterm labor in the uterus and

decidua (Figure 2E&F). Moreover, quantile-quantile plots of DEGs from enriched cell types

indicate that the uterus is the tissue most affected by the process of labor (Figure 2H-J). Thus,

preterm labor primarily induces gene expression in the dominant cell types from each tissue; yet,

the substantial amount of downregulated gene expression in innate immune cells may indicate an

immunological switch from one transcriptomic program to another to combat infection.

Preterm labor induced by *E. coli* involves conserved cell types that display distinct

processes in the reproductive tissues

The transcriptomic profiling of cell types suggested that specific subsets show conserved

responses with preterm labor across the reproductive tissues. Therefore, we next focused on the

shared preterm labor-specific gene expression among the uterus, decidua, and cervix. The Venn

diagram displayed in Figure 3A highlights the overlap in DEGs across tissues, particularly the

uterus and decidua. Correlation analyses indicated stronger relationships between preterm labor

and gene expression changes in the uterus and decidua than in the cervix (Figure S3A), which

was reflective of the total preterm labor-associated DEGs in each tissue. This observation was

further confirmed by the correlation between the gene expression profiles of the uterus and

decidua, which was stronger than the correlations between the decidua and cervix or the uterus

and cervix (Figure S3B). Given that the uterus, decidua, and cervix all displayed some degree of

correlation for preterm labor-associated gene expression, we evaluated the cell type-specific

transcriptomic changes that were conserved across all three tissues. We found that innate

immune cell types (Monocyte and Neutrophil) as well as non-immune cell types (Epithelial-3, -

4, -6 and Endothelial) showed conserved gene expression changes associated with preterm labor

across the uterus, decidua, and cervix (Figure 3B). We reasoned that, although the transcriptome

profiles of specific cell types were affected across all tissues, such cells may display distinct

biological processes according to their location. Gene Ontology (GO) analysis of the Neutrophil

cell type in the uterus, decidua, and cervix revealed that, while these cells shared some processes

such as "response to bacterium" and "response to lipopolysaccharide", processes specific to

Neutrophils in each tissue were also observed (Figure 3C). Uterine Neutrophils showed

enrichment of processes related to cytokine signaling and anti-viral response, whereas decidual

Neutrophils showed enrichment of cellular activation-associated processes (Figure 3C). In the

cervix, unique enriched Neutrophil processes were primarily associated with response to external

stimuli and bacteria (Figure 3C). Uterine and decidual Monocyte and Macrophage-1 cells also

shared enriched processes related to cytokine production and response to

bacteria/lipopolysaccharide, with decidual Monocytes also showing enrichment of activation-

associated processes (Figure 3D). By contrast, cervical Monocytes displayed highly distinct

processes related to protein synthesis and humoral immune response (Figure 3D), suggesting that

such cells may functionally differ from their counterparts in the uterus and decidua. Epithelial-6,

which had sufficient DEGs to perform GO analysis in all three tissues, displayed largely

consistent processes across the uterus, decidua, and cervix that were related to inflammation,

anti-bacterial response, and cytokine production (Figure 3E). The uterine Epithelial-4 cell type

displayed enrichment of several chemotaxis-associated processes, suggesting involvement in

leukocyte recruitment to this tissue, whereas the cervical Epithelial-4 showed enrichment of

effector functions such as production of NO, IL-1, and IFNγ (Figure 3E). Epithelial-3, which

only displayed sufficient DEGs for GO analysis in the cervix, showed enrichment of multiple

processes related to promotion of B-cell and antibody responses (Figure 3E). Thus, the

conserved cell types affected by preterm labor in the uterus, decidua, and cervix each display

distinct enrichment of biological processes, suggesting that similar cell types display tissue-

specific functions in the context of intra-amniotic infection leading to preterm labor. Together

with the observed increase in cervical Epithelial-8 cell counts with preterm labor, it is possible

that the upregulation of inflammatory gene expression represents an infection-induced

differentiation of cervical epithelial cells to better participate in host defense mechanisms in this

compartment.

To further infer cellular functionality in preterm labor, we then utilized the Kyoto

Encyclopedia of Genes and Genomes database to evaluate the pathways enriched in labor-

associated DEGs in each cell type (Figure S3C). Strikingly, both immune and non-immune cell

types with altered gene expression in preterm labor showed enrichment of immunological

pathways such as "cytokine-cytokine receptor interaction", "NOD-like receptor signaling

pathway", and "viral protein interaction with cytokine and its receptor" across the three tissues

(Figure S3C). Such findings are consistent with previous studies showing the upregulation of

immune-related pathways in decidual endothelial (Huang et al., 2020) and stromal cells (Huang

et al., 2021) from women with labor. Additional inflammatory pathways such as "NF-kappa B

signaling pathway" and "Toll-like receptor signaling pathway" were also represented, to a lesser

extent, by immune cells (e.g., NK cells and Neutrophil) as well as non-immune cells such as

epithelial cells, including in the cervix (Figure S3C).

We also investigated the biological processes enriched in several non-immune cell

subsets that were conserved between the uterus and decidua with preterm labor (Figure S4).

Interestingly, Stromal-1 and Stromal-2 in the uterus showed largely similar enrichment of

biological processes, as did the Stromal-1 and Stromal-2 cell types in the decidua (Figure S4A).

Yet, these cell types differed between tissues, given that the decidual stromal cells were enriched

for leukocyte migration and chemotaxis whereas the uterine stromal cells showed enrichment for

response to immune signaling (Figure S4A). The Fibroblast-1 and Stromal-3 cell types showed

less diversity in their enriched processes compared between the uterus and decidua, with the

former being associated with host defense against infection and the latter associated with

immune activation, including adaptive immunity (Figure S4B). Fibroblast-2 and Fibroblast-3

were also comparable between the uterus and decidua; however, the decidual Fibroblast-3

showed more striking enrichment of responses to microorganisms and cytokine signaling (Figure

S4C). Finally, the uterine and decidual Endothelial cell types displayed similar enrichment of

processes related to host defense, innate immunity, and cytokine signaling, with the decidual

subset showing modestly higher enrichment for processes related to neutrophil migration (Figure

S4D).

Taken together, these data indicate that the uterus, decidua, and cervix contain cell types

that display distinct tissue-specific gene expression profiles in preterm labor, pointing to

differing functional roles for these cells in the host response to intra-amniotic infection. Yet,

there is an overall tendency for the enrichment of similar immunological pathways in both

immune and non-immune cell types across tissues, likely as part of the common host response to

intra-amniotic infection.

Preterm labor influences cell-cell communications in the reproductive tissues

Having established that preterm labor drives distinct transcriptomic changes in specific

cell types in the uterus, decidua, and cervix, we next leveraged our single-cell data to elucidate

cell-cell communication networks in these tissues.

Cell-cell communications in the uterus

The uterus is a highly heterogeneous organ with multiple described regions that differ in

cellular composition and function (Bukowski et al., 2006; Danforth and Ivy, 1949; Liu et al.,

2015a; Mosher et al., 2013; Patwardhan et al., 2015; Pollard et al., 2000; Schwalm and

Dubrauszky, 1966; Sooranna et al., 2006; Sparey et al., 1999; Wikland et al., 1982). To unravel

the intercellular communications taking place in the murine uterus with preterm labor, we

performed a correlation analysis across preterm labor-associated genes for each pair of identified

cell types (Figure S5A). The strongest correlations were observed for non-immune cell types,

such as stromal, epithelial, fibroblast, smooth muscle, and Endothelial cell types (Figure S5A),

suggesting that these cells exhibit similar changes in gene expression with preterm labor. Innate

and adaptive immune cell types also showed moderate correlations; namely, T cell, NK-cell-1,

NK-cell-2, Macrophage-1, Macrophage-2, Monocyte, and Dendritic cell (Figure S5A). The cell

types with the weakest correlations were primarily epithelial subsets as well as Neutrophil,

Erythroid, Stromal-3, and SMC-2 (Figure S5A), indicating that such cells show more distinct

gene expression changes with preterm labor.

We next used CellChat to infer cell-cell communications within the uterus using our

single-cell gene expression data and a database of established interactions between signaling

ligands, receptors, and their cofactors (Jin et al., 2021). Signaling pathways that were enriched or

diminished in preterm labor, as well as those that were unaffected by this process, are shown in

Figure S5B and Table S3. The alluvial plots shown in Figure 4A&B display the major cell-cell

communication processes taking place in the uterus as well as the cell types that participate as

senders or receivers in each process. Both innate and adaptive immune cell subsets (Neutrophil,

Macrophage-1, Monocyte, Dendritic cell, NK cell-1, T-cell) contribute to the top signaling

pathways implicated in preterm labor, such as CCL, CXCL, complement, IFN-I and IFN-II, IL-

1, and IL-6 (Figure 4A&B). Notably, multiple non-immune subsets also participate in these

processes; namely, fibroblast, stromal, epithelial, SMC, and Endothelial cell types (Figure

4A&B). While both immune and non-immune cell types served as receivers of preterm labor-

associated signaling, specific responders to each pathway could be distinguished (Figure 4B).

For example, the signaling pathway of IL-6, which is commonly utilized as a biomarker of intra-

amniotic inflammation (Yoon et al., 2001), was primarily driven by immune cell types (Figure

4A); yet, the receiver cells for this pathway were non-immune subsets (Figure 4B). Conversely,

the primary senders for the Annexin signaling pathway were non-immune cell types (Figure 4A),

with the downstream receivers being predominantly immune cells (Figure 4B). The changes in

cell-cell communication occurring as a result of preterm labor were visualized using the arrow

plot in Figure 4C, where the directionality of each cell type arrow reflects the propensity for

increased outgoing and/or incoming interaction strength. Cell types such as Macrophage-2,

Stromal-2, Stromal-3, and Fibroblast-3 showed primarily even increases in incoming and

outgoing signaling with preterm labor (Figure 4C). Other cell types were more biased towards

incoming interactions, such as Macrophage-1, Neutrophil, Dendritic cell, Plasmocyte, Monocyte,

and Epithelial-6, or outgoing interactions, such as Stromal-1, Fibroblast-1, and Fibroblast-2

(Figure 4C). Interestingly, several cell types showed a net decrease in signaling with preterm

labor; namely, T cell and SMC-1 (Figure 4C).

The top 25% of aggregated cellular interactions in the uterus were then contrasted

between the control and preterm labor groups, emphasizing the overall increase in cell-cell

signaling with preterm labor as well as the incorporation of new signaling pathways from cell

types that were rarely present in control tissues, such as Neutrophil (Figure 4D). Interestingly,

while Macrophage-1 signaling was increased, Macrophage-2 signaling decreased, which could

indicate a homeostatic role for the latter subset that is diminished in preterm labor, as previously

reported (Gomez-Lopez et al., 2021a). Next, we examined the top contributors within uterine

cell-cell signaling pathways enriched with preterm labor (Figure 4E). We found that macrophage

subsets and Dendritic cells were primary contributors to CCL signaling between control uterine

cell types, and such signaling was further strengthened in preterm labor (Figure 4E). By contrast,

the Galectin signaling pathway, already enriched in control uterine tissues, was upregulated in

new cell types in preterm labor (e.g., Epithelial-6) and diminished in others (e.g., Macrophage-2)

(Figure 4E).

We also explored the changes in cell type-specific expression of genes related to

progesterone and prostaglandin signaling in the uterus (Figure S5C and Figure S6A). As

expected, progesterone-related gene expression was consistently downregulated across uterine

cell types in preterm labor (Figure S5C). Prostaglandin-related gene expression showed more

activity in the uterus than in other tissues (Figure S6); yet, preterm labor-associated changes in

each gene were consistent across uterine cell types (Figure S6A), further supporting the

involvement of multiple immune and non-immune cell populations in labor-mediator signaling

pathways.

Taken together, these findings highlight the complex cell-cell communication network

taking place in the murine uterus and how such interactions are modulated by the inflammatory

process of preterm labor in both immune and non-immune cell types.

Cell-cell communication in the decidua

We next examined the correlations across preterm labor-associated changes in gene

expression for each pair of cell types identified in the decidua (Figure S7A). Similar to the

uterine tissues, the strongest correlations were observed for non-immune cell types (e.g., stromal,

epithelial, fibroblast, smooth muscle, and Endothelial), followed by innate and adaptive immune

cells, of which the macrophage and Monocyte clusters were best correlated (Figure S7A).

Similarly, the weakest correlations were observed for some epithelial cell types and Neutrophils

(Figure S7A). Thus, decidual cells display preterm labor-associated changes in gene expression

with varying magnitude of sharing among cell types, which resemble those observed in the

uterine tissues.

The inferred cell-cell signaling pathways that were enriched or diminished in the decidua

with preterm labor are shown in Figure S7B and Table S3. From among these, the top pathways

are displayed with their participating sender and receiver cell types in Figure 5A&B. Similar to

the uterine tissues, key cell-cell communication pathways were primarily related to immune

functions such as cytokine and chemokine signaling (Figure 5A&B). Unique among the three

compared tissues, the IL-17 pathway was most prominent in the decidua (Figure 5A&B), which

was consistent with a previous report of IL-17 signaling in endothelial cells derived from the

human peripartum decidua (Huang et al., 2020) and suggests that decidual T cells participate in

the local inflammatory response to intra-amniotic infection. Among other identified signaling

pathways, both immune and non-immune cell subsets contributed as senders or receivers,

including the NK-cell-2 subset that was not implicated in uterine cell-cell signaling (Figure

5A&B). The decidual Epithelial-5 cell type appeared to be primarily functioning as a receiver of

cell-cell signaling in this tissue (Figure 5A&B). We then visualized the preterm labor-driven

changes in incoming and outgoing signaling, and observed that subsets such as Monocyte,

Macrophage-1, Neutrophil, NK cells, and Dendritic cells showed predominantly incoming

interactions (Figure 5C). On the other hand, stromal and fibroblast subsets as well as T cells

tended towards increased outgoing signaling, while SMC-1 and Endothelial cells showed an

overall reduction in interaction strength (Figure 5C). Interestingly, outgoing interaction strength

was greater in decidual T cells compared to uterine T cells (Figure 5C vs. Figure 4C), which

further emphasizes a role for T cell-derived signals in the pathophysiology of preterm labor

associated with intra-amniotic infection (Gershater et al., 2022). Consistent with enhanced cell-

cell signaling in preterm labor, aggregated cellular interaction plots demonstrated an overall net

increase in decidual intercellular interactions compared to controls (Figure 5D). Similar to the

uterine tissues, enriched signaling pathways such as CCL and Galectin were primarily driven by

Macrophage-1, Monocyte, and Dendritic cell in preterm labor, with overall interactions among

cell types increasing compared to controls (Figure 5E).

Similar to the changes observed in the uterus, decidual expression of progesterone-related

genes was consistently downregulated across cell types with preterm labor (Figure S7C). The

patterns of change in prostaglandin-related gene expression were also similar between the

decidua and uterus; yet, some difference in the magnitude of change between compartments were

observed for multiple genes, potentially indicating a stronger upregulation of preterm labor-

associated prostaglandin signaling in the uterus relative to the decidua (Figure S6B).

These data provide insight into the distinct cellular interactions taking place in the

decidua during the process of preterm labor, including the involvement of cell types and

signaling pathways not observed in other tissues. Yet, the decidua and uterus also share cell type-

specific communications that are affected by preterm labor.

Cell-cell communication in the cervix

Our investigation of the cell type-specific changes taking place in the cervix with preterm

labor had indicated that the Neutrophil and Monocyte subsets were most affected, in tandem with

previous studies showing labor-associated infiltration of immune cells (Gonzalez et al., 2011;

Osman et al., 2003; Payne et al., 2012; Sakamoto et al., 2005; Timmons et al., 2009), as well as

multiple epithelial cell subsets (Figures 2 and 3). Consistently, correlation analysis of these cell

types showed that the strongest associations in gene expression changes driven by preterm labor

were observed among epithelial cell types (Figure S8A), while Neutrophils and Monocytes

showed modest correlation of genes affected by preterm labor (Figure S8A). Inferred cell-cell

signaling pathways were noticeably fewer compared to the other tissues and included multiple

processes unique to the cervix (Figure S8B and Table S3), which could be attributed to the less

diverse cell type composition observed in this tissue. Indeed, as shown by the participating

senders and receivers, signaling pathways that were strongly implicated in the uterus and decidua

with preterm labor were not as enriched in cervical cell types (Figure 6A&B). On the other hand,

cell-cell signaling pathways related to extracellular matrix were strongly represented (Figure

6A&B), which is consistent with the primarily connective tissue composition of the cervix

(Granstrom et al., 1989; Krantz and Phillips, 1962; Ludmir and Sehdev, 2000). As expected,

given their inferred roles as receivers, most cervical epithelial cell types showed strong incoming

interactions with preterm labor, whereas the SMC-1, Fibroblast-2, and Stromal-1 subsets showed

a tendency towards increased outgoing interactions (Figure 6B&C). This was further supported

by the aggregated cervical cell-cell interactions in the control and preterm labor groups showing

increased receipt of signaling by Epithelial-1 and Epithelial-8 as well as SMC-1 and Fibroblast-2

(Figure 6D). Notably, Fibroblast-2 and SMC-1 were top contributors to enriched signaling

pathways such as Collagen and Tenascin (Figure 6E). It is possible that the Fibroblast-2 and/or

SMC-1 cell clusters include cervical myofibroblasts, given that a previous histological

investigation indicated a pregnancy-specific accumulation of such cells, which could be

interacting with the extracellular matrix to aid in supporting the mechanical stresses present

during labor (Montes et al., 2002). In the last decade, a new paradigm for the role of SMCs in the

human cervix has emerged, suggesting a sphincter-like function of the internal os, in which the

SMCs express contractility-associated proteins that are responsive to oxytocin signaling (Vink et

al., 2016). Together with our current findings, these observations support the involvement of

SMC-1 and Fibroblast-2 subsets in preterm labor-associated signaling in the murine cervix, and

further emphasize the unique cell-cell signaling pathways taking place in the cervix during

preterm labor.

Shared cellular signaling pathways in the murine uterus and human myometrium during

the processes of preterm and term labor

Last, to examine the shared pathways implicated in the process of parturition in mice and

humans, we utilized the differential cellular interactions in the murine uterus with preterm labor

together with our previously generated single-cell atlas of the human myometrium with labor at

term (Pique-Regi et al., 2022) (Figure 7). We investigated the interaction strength between cell

types affected by labor as well as prominent signaling pathways, and contrasted these between

the murine and human tissues. Overall, we found that labor-associated cell-cell interactions were

primarily driven by SMC, stromal, fibroblast, and innate immune cell types in both the murine

uterus and human myometrium, independently of differences in sender/receiver status (Figure

7A&B).

Specifically, in the murine uterus, non-immune cell types such as Fibroblast-1, -2, and -3,

Stromal-2 and -3, and SMC-1 showed the strongest labor-associated increase in outgoing cell-

cell signaling, whereas outgoing signaling by Macrophage-2 was greatly diminished (Figure 7A).

The top receivers of labor-associated signaling were Fibroblast-3, Stromal-2 and -3, and

Epithelial-6 as well as innate immune cell types (Monocyte, Neutrophil, and Macrophage-1)

(Figure 7A). Endothelial, SMC-1, Plasmocyte, and Macrophage-2 showed diminished incoming

signaling with preterm labor (Figure 7A).

In the human myometrium, labor involved increased outgoing signaling almost

exclusively by smooth muscle cell and stromal subsets, with Myofibroblast and LED cells also

contributing to this process (Figure 7B). Receivers of such outgoing interactions included

macrophage subsets and Monocytes, which is consistent with previous investigations indicating

that myometrial cell contraction during labor is promoted by crosstalk with macrophages in a co-

culture model (Wendremaire et al., 2020). Multiple myometrial cell subsets displayed

substantially reduced outgoing signaling with labor, such as EVT, Macrophage-4, Plasmablast,

Unciliated Epithelial, and ILC (Figure 7B). For incoming signaling, the majority of human

myometrial cell types tended to have increased interaction only with those cells that displayed

greater outgoing signals, such as smooth muscle and stromal cells (Figure 7B). Some exceptions

seemed to be the Macrophage-1, SMC-3, and Stromal-1 subsets, whose incoming signaling from

the majority of other cell types was strengthened (Figure 7B).

We then examined the top 25% of aggregated cell-cell interactions in the human

myometrium at term without labor and at term with labor (Figure S8A). Consistent with the

correlation analysis above (Figure 7B), a clear shift in the cell types contributing to myometrial

cell-cell signaling was observed between groups, with the EVT, Macrophage-4, Plasmablast, and

Unciliated epithelial clusters showing greatly diminished interactions in labor, and smooth

muscle and stromal cell subsets acquiring increased interactions (Figure S9A). This shift in

interaction with labor was also reflected by the combined differential interaction analysis shown

in Figure S8B. When examining the specific cell-cell signaling pathways that were impacted by

labor, terms such as Collagen, IL-1, CCL, Complement, and CXCL were found to be shared

between the murine uterus and human myometrium, indicating shared labor-associated cellular

signaling between species (Figure 7C&D and Figure S9C-G); yet, the closer inspection of these

shared signaling pathways revealed differences in the cell types contributing to each (Figure

7C&D and Figure S9C-G). The relevance of immune pathways such as chemokine signaling in

the process of labor is supported by previous reports demonstrating that the use of a chemokine

inhibitor on human myometrial cells decreased contraction and gap junction formation, thereby

disrupting intercellular communication (Boros-Rausch et al., 2021; Coleman et al., 2020). Taken

together, these data indicate that labor-associated cell-cell interactions heavily involve SMC,

stromal, fibroblast, and innate immune cell types in the murine uterus and the human

myometrium, providing evidence of labor-specific signaling processes between immune and

non-immune cells that are shared between species. Yet, this interpretation should be taken with

caution, given that we compared the physiologic process of labor in the human myometrium with

the pathologic process of labor induced by bacteria in the murine uterus.

DISCUSSION

The current study provides a single-cell atlas of the murine uterus, decidua, and cervix

that highlights the cell type composition, transcriptional profiles, and cell-cell signaling taking

place in these tissues in normal late gestation or in the context of infection-induced preterm labor

and birth. scRNA-seq allowed for the deep characterization of immune and non-immune cells

that demonstrates the underappreciated heterogeneity of cell types conventionally considered to

be uniform in function, such as uterine smooth muscle cells and cervical epithelial cells. Our data

can thus serve as a reference for future studies seeking to target specific subsets of these cells,

which may have differing roles in pregnancy and labor as indicated by their distinct

transcriptional profiles. Shared modulation of gene expression was noted between uterine and

decidual cell types and was reflected by the similar enrichment of labor-associated signaling

pathways, which is consistent with the spatial proximity of these tissues. Yet, the comparison of

individual cell types across tissues indicated that the most represented biological processes can

vary according to location, and therefore the tissue of origin should be taken into consideration

when inferring cellular function. Importantly, here we provide scRNA-seq characterization of the

understudied cervical tissues, demonstrating a plethora of epithelial subsets with different

potential functions as well as smooth muscle and fibroblast cell types that indicate an unexpected

level of heterogeneity in the cervix. Recent evidence has suggested a sphincter-like function of

SMCs in the internal os of the human cervix (Vink et al., 2016), and our current findings support

this concept. Inferred cell-cell communications provided evidence of substantial crosstalk among

uterine, decidual, and cervical cell types during the process of preterm labor, highlighting key

signaling pathways that could potentially be targeted in future translational studies aimed at

preventing spontaneous preterm labor. In addition, such analysis demonstrated cell types with

elevated or diminished interactions driven by inflammation, which can serve to identify those

cell types that are most and least involved in such signaling. To demonstrate the application of

our single-cell dataset, we have leveraged prior single-cell analysis of the human myometrium in

term labor to evaluate cellular interactions compared to our murine model of preterm labor. This

analysis provided useful insight into shared signaling pathways associated with the inflammatory

process of labor, providing a practical demonstration of how our scRNA-seq dataset can be

leveraged for *in silico* discovery of specific cell types, pathways, or genes that can be

subsequently targeted *in vitro* and/or *in vivo*.

The current study has some limitations. First, it is important to consider that spontaneous

preterm labor is a syndrome, for which intra-amniotic infection represents only one known

etiology (Romero et al., 2014a). Here, we focused on preterm labor triggered by intra-amniotic

infection of *E. coli*, a Gram positive bacteria, using an established animal model that resembles

the clinical condition (Gomez-Lopez et al., 2018a). Other known or proposed etiologies for

spontaneous preterm labor may have subtle differences in the involved cell types and associated

signaling pathways, and thus further characterization of the cellular atlas in each preterm labor

subtype is necessary for their distinction. Second, intra-amniotic infection is often polymicrobial

and/or can be induced by a variety of bacterial species (Burnham et al., 2020; Romero et al.,

2015c), of which *Ureaplasma* species are the most commonly found in the amniotic cavity

(Motomura et al., 2020b; Romero et al., 1989; Sweeney et al., 2016; Viscardi, 2010; Yoneda et

al., 2016b; Yoon et al., 1998); therefore, the immune responses triggered by each bacterium or

cluster of bacteria may differentially affect cellular responses in the reproductive and gestational

tissues. However, the *in vivo* standardization of a polymicrobial infection model and the isolation

of clinically-relevant *Ureaplasma* species are challenging, and thus here we utilized intra-

amniotic infection with an easily-cultured bacterium, *E. coli*, to induce preterm birth in mice.

Future investigations may utilize other bacteria detected in the amniotic cavity of women

diagnosed with intra-amniotic infection (DiGiulio et al., 2010; Romero et al., 2015c). In addition,

it should be noted that single-cell RNA-seq is a discovery-based approach that we have utilized

to generate an atlas of the murine reproductive tissues, and thus careful interpretation is required

when extrapolating specific findings to the human context. Last, it is worth mentioning that RNA

techniques using single-cell suspensions lose information about the spatial relationships among

cell types in target tissues; therefore, such data may be complemented by using spatial

transcriptomics and/or proteomics. Nonetheless, our data can serve as a resource for targeted

studies that can validate such findings using human samples.

ACKNOWLEDGEMENTS

The authors would like to acknowledge Yesong Liu for his assistance in performing some of the immunofluorescence imaging.

This research was supported, in part, by the Perinatology Research Branch, Division of Obstetrics and Maternal-Fetal Medicine, Division of Intramural Research, *Eunice Kennedy Shriver* National Institute of Child Health and Human Development, National Institutes of Health, U.S. Department of Health and Human Services (NICHD/NIH/DHHS); and, in part, with Federal funds from NICHD/NIH/DHHS under Contract No. HHSN275201300006C. Dr. Romero has contributed to this work as part of his official duties as an employee of the United States Federal Government. This research was also supported by the Wayne State University Perinatal Initiative in Maternal, Perinatal and Child Health. The funders had no role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript. Figures include art created with BioRender.com

AUTHOR CONTRIBUTIONS

Conceptualization, R.R., R.P-R., and N.G-L.; Methodology, V.G-F., J.G., Y.X., and N.G-L.; Validation, A.P., R.P-R., and A.L.T.; Formal Analysis, V.G-F., A.P., R.P-R., and N.G-L.; Investigation, V.G-F., J.G., E.P., B.P., D.M., Y.X., L.T., Z.L., and A.L.T.; Resources, R.R., A.L.T., R.P-R., and N.G-L.; Data Curation, V.G-F., A.P., and R.P-R.; Writing - Original Draft, V.G-F., R.R., A.P., J.G., B.P., D.M.; Writing - Review & Editing, V.G-F., R.R., A.P., J.G., E.P., B.P., D.M., Y.X., L.T., Z.L., A.L.T, R.P-R., and N.G-L.; Visualization, V.G-F., A.P., J.G., E.P.; Supervision, R.R., R.P-R., and N.G-L.; Project Administration, N.G-L.; Funding Acquisition, R.R., and N.G-L.

DECLARATION OF INTERESTS

The authors declare that no competing interests exist.

FIGURE TITLES AND LEGENDS

Figure 1. Single-cell atlas of the murine reproductive tissues during preterm labor induced

by intra-amniotic infection. **(A)** Experimental design for the ultrasound-guided intra-amniotic

injection of *E. coli* or PBS (vehicle control) into pregnant mice on 16.5 days *post coitum* (dpc) (n

= 3-6 per group). Mice were monitored to determine pregnancy outcomes **(B-C)**. Gestational age

and preterm birth rate of dams intra-amniotically injected with *E. coli or* PBS. Gestational age

was compared using a 2-sided Mann-Whitney U-test, and preterm birth rates were compared

using a 2-sided Fisher's test. $P < 0.05$ was considered significant. **(D)** Experimental design for

the determination of cervical length on 16.5 dpc, prior to intra-amniotic injection of *E. coli* or

PBS, and 24 h later (17.5 dpc) (n = 6-9 per group). **(E)** Cervical length of dams intra-

amniotically injected with *E. coli* or PBS at 16.5 and 17.5 dpc. Cervical length was compared

between time points using a 2-sided Mann-Whitney U-tests. $P < 0.05$ was considered significant.

The change in cervical shortening was calculated by considering the measurement at 16.5 dpc as

100%. **(F)** Diagram illustrating the generation of single-cell suspensions from the uterus,

decidua, and cervix collected for single-cell RNA-sequencing experiments (scRNA-seq) (n = 4

per group). **(G)** Uniform manifold approximation and projection (UMAP) plot showing all cell

types present in the uterus, decidua, and cervix. **(H)** UMAP color-coded plot showing tissue-

specific predominance of distinct cell types in the uterus (blue), decidua (pink), and cervix

(orange). Blue dotted lines distinguish major cell types: myeloid, endothelial, stromal, smooth

muscle, epithelial, and lymphoid. Abbreviations used: SMC, smooth muscle cell; NK cell,

natural killer cell.

Figure 2. **Preterm labor induced by *E. coli* dysregulates the repertoire and gene expression**

of immune and non-immune cell types in the reproductive tissues. **(A)** Color-coded uniform

manifold approximation and projection (UMAP) plot showing the effects of preterm labor on the

abundance of specific cell types (shown in red) compared to control (shown in blue). **(B-D)** Bar

plots showing the numbers of each cell type in the uterus, decidua, and cervix. The comparison

of cell numbers between the two study groups for each cell type was performed using a 2-sided t-

test. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$. **(E-G)** Bar plots showing the

numbers of differentially expressed genes (DEGs) induced by preterm labor in ~~each~~ cell type in

the uterus, decidua, and cervix. Red and pink bars indicate upregulated and downregulated

DEGs, respectively (derived from DESeq2, $q < 0.1$). The comparisons of the fraction of

downregulated and upregulated DEGs in each cell type between the study groups were

calculated using two-sided binomial tests. * $q < 0.05$, ** $q < 0.01$, *** $q < 0.001$, **** $q <$

0.0001. **(H-J)** Quantile-quantile plot showing differential expression of genes analyzed for

selected enriched cell types from the uterus, decidua and cervix. Deviation above the 1:1 line

(solid black line) indicates enrichment. Abbreviations used: SMC, smooth muscle cell; NK cell,

natural killer cell.

Figure 3. **Preterm labor induced by *E. coli* involves conserved cell types that display**

distinct processes in the reproductive tissues. **(A)** Venn diagrams showing the numbers of

differential expressed genes (DEGs, $q < 0.1$) that are unique to or shared among the uterus,

decidua, and cervix. **(B)** Box plots showing the correlation of specific cell types affected by

preterm labor and conserved across the uterus, decidua, and cervix using the Spearman's method.

Cluster profiler dot plots showing the preterm labor-associated Gene Ontology (GO) biological

processes that are unique or shared among **(C)** Neutrophil, **(D)** Monocyte and Macrophage, and

(E) Epithelial cell types from the uterus, decidua, and cervix. The size and color of each dot

represent gene ratio and significance level, respectively. 1-sided Fisher's exact tests were used.

Abbreviations used: SMC, smooth muscle cell; NK cell, natural killer cell.

Figure 4. Cellular interactions in the uterus during preterm labor. (A-B) Alluvial plots

showing the roles of specific cell types as senders or receivers of preterm labor-associated

signaling in the uterus based on selected enriched signaling pathways comparing the overall

information flow within networks between preterm labor and control derived from CellChat (full

list of pathways in Figure S5B). Connecting lines are color-coded and represent the participation

of each cell type as senders or receivers of the indicated signaling pathway. Line thickness is

proportional to the strength of interaction. **(C)** Arrow plot showing changes in outgoing and

incoming interactions strength between preterm labor (point of the arrow) and control condition

(base of the arrow) for specific cell types in the uterus. **(D)** Circle plots showing the top 25% of

aggregated interactions among cell types in the uterus for control and preterm labor groups. Each

node represents a cell type and the interaction is shown by lines color-coded based on the sender

cell. **(E)** Circle plots showing the top 25% increased (red) or decreased (blue) signaling

interactions in the uterus for specific pathways in preterm labor compared to controls.

Figure 5. Cellular interactions in the decidua during preterm labor. **(A-B)** Alluvial plots

showing the roles of specific cell types as senders or receivers of preterm labor-associated

signaling in the decidua based on 11 enriched signaling pathways comparing the overall

information flow within networks between preterm labor and control derived from CellChat (full

list of pathways in Figure S6B). Connecting lines are color-coded and represent the participation

of each cell type as senders or receivers of the indicating signaling pathway. Line thickness is

proportional to the strength of interaction. **(C)** Arrow plot showing changes in the outgoing and

incoming interaction strength between preterm labor (point of the arrow) and control condition

(base of the arrow) for specific cell types in the decidua. **(D)** Circle plots showing the top 25% of

aggregated interactions among cell types in the decidua for control and preterm labor groups.

Each node represents a cell type and the interaction is shown by color-coded lines. **(E)** Circle

plots showing the top 25% increased (red) or decreased (blue) signaling interactions in the

decidua for specific pathways in preterm labor compared to controls.

Figure 6. Cellular interactions in the cervix during preterm labor. **(A-B)** Alluvial plots

showing the roles of specific cell types as senders or receivers of preterm labor-associated

signaling in the cervix based on selected enriched signaling pathways (full list of pathways in

Figure S7B). Connecting lines are color-coded and represent the participation of each cell type as

senders or receivers of the indicating signaling pathway. Line thickness is proportional to the

strength of interaction. **(C)** Arrow plot showing changes in the strength of outgoing and

incoming interactions between preterm labor (point of the arrow) and control (base of the arrow)

for specific cell types in the cervix. **(D)** Circle plots showing the top 25% of aggregated

interactions among cell types in the cervix for control and preterm labor groups. Each node

represents a cell type and the interaction is shown by color-coded lines. **(E)** Circle plots showing

the top 25% increased (red) or decreased (blue) signaling interactions in the cervix for specific

pathways in preterm labor compared to controls.

Figure 7. Shared cellular signaling pathways in the murine uterus and human myometrium during the processes of preterm and term labor. (A) Heatmap showing the differential interaction strength among cell types in the murine uterus with preterm labor. Red and blue shading indicate increased or decreased signaling, respectively, in preterm labor compared to control. **(B)** Heatmap showing the differential interaction strength among cell types in the human myometrium with term labor. Red and blue shading indicate increased or decreased signaling, respectively, in term labor compared to term without labor. **(C)** Circle plots representing the top 25% murine uterine cell-cell communications inferred for the Collagen, IL-1, CCL pathways for the control and preterm labor groups. **(D)** Circle plots representing the top 25% human myometrial cell-cell communications inferred for the Collagen, IL-1, and CCL pathways for the control and preterm labor groups. Abbreviations used: SMC, smooth muscle cell; NK cell, natural killer cell; EVT, extravillous trophoblast; ILC, innate lymphoid cell; LED, lymphoid endothelial decidual cell.

STAR METHODS

KEY RESOURCES TABLE

(Submitted separately)

RESOURCE AVAILABILITY

Lead contact

Further information and requests for resources and reagents should be directed to and will

be fulfilled by the lead contact, Nardhy Gomez-Lopez (nardhy.gomez-lopez@wayne.edu).

Material Availability

This study did not generate new unique reagents.

Data and Code Availability

The scRNA-seq data reported in this study has been submitted to the Gene Expression

Omnibus repository (GEO accession number GSE200289). All software and R packages used

herein are detailed in materials and methods section. Scripts detailing the single-cell analysis are

available at GitHub (https://github.com/piquelab/murine_scRNA)

EXPERIMENTAL MODEL DETAILS

Mice

Mice were purchased from The Jackson Laboratory (Bar Harbor, ME, USA) and bred in

the animal care facility at the C.S. Mott Center for Human Growth and Development, Wayne

State University (Detroit, MI, USA). Mice were under a circadian cycle of light:dark = 12:12 h.

Eight- to twelve-week-old C57BL/6 (RRID:IMSR_JAX:000664) female mice were bred with

BALB/cByJ male mice (RRID:IMSR_JAX:001026), and females were examined daily between

0800 and 0900 to check for the presence of a vaginal plug, which was considered as 0.5 days

post coitum (dpc). Upon observation of a vaginal plug, females were removed from mating cages

and housed separately, and their weights were monitored daily. At 12.5 dpc, a weight gain of $\geq$2g

was considered confirmation of pregnancy. Mice were randomized to receive intra-amniotic

injection of *E. coli* or vehicle control, and investigators were not blinded to control or treatment

assignment. No experimental mice were excluded from analysis. All procedures and experiments

were approved by the Institutional Animal Care and Use Committee (IACUC) at Wayne State

University under Protocol nos. 18-03-0584 and 21-04-3506.

METHOD DETAILS

Preparation of E. coli for intra-amniotic injection

Escherichia coli was purchased from the American Type Culture Collection (ATCC, Manassas, VA, USA) (ATCC 12014) and was grown in Luria-Bertani (LB) broth (cat. no. L8050, Teknova, Hollister, CA, USA) at 37°C. From an overnight culture, a sub-culture was placed with fresh LB broth and grown to the logarithmic phase (OD_{600} 0.9 - 1). Additional dilution was performed using sterile 1X phosphate-buffered saline (PBS, Thermo Fisher Scientific/Gibco, Grand Island, NY, USA) to reach a working concentration of 10 CFU/25µL.

Measurement of cervical length by ultrasound

Dams were anesthetized on 16.5 dpc by inhalation of 2% isoflurane (Aerrane; Baxter Healthcare Corporation, Deerfield, IL, USA) and 1 to 2 liters/min of oxygen in an induction chamber. Anesthesia was maintained with a mixture of 1.5 to 2% isoflurane and 1.5 to 2 liters/min of oxygen. Mice were positioned on a heating pad and stabilized with adhesive tape. Fur removal from the abdomen was performed using Nair cream (Church & Dwight Co., Inc., Ewing, NJ, USA). Sterile forceps were utilized to expose the vulva and 200 µL of Sterile Aquasonic® 100 ultrasound transmission gel (Parker laboratories, Fairfield, NJ, USA) was used to fill the vagina to create contrast and allow for clear visualization of the external limit of the uterine cervix (i.e., the external os). A trans-abdominal ultrasound approach was utilized to evaluate the cervix. The transducer was slowly moved toward the lower part of the abdomen and the cervix was positioned in a longitudinal view. The cervical length was measured from the internal to the external os at least three times per mouse, and its average was utilized as the final value for cervical length. This procedure was performed prior to ultrasound-guided injection

with either *E. coli* or PBS and repeated 24 h later (on 17.5 dpc) (i.e., prior to tissue collection).

The change in cervical length was determined as a percentage by considering the cervical length

on 16.5 dpc as 100% and then calculating the percentage of the cervical length on 17.5 dpc.

Intra-amniotic inoculation with E. coli

Dams that underwent cervical measurement were maintained on the heating pad under

anesthesia as described above. The ultrasound transducer was slowly moved toward the abdomen

to localize the amniotic sacs. The syringe with *E. coli* suspension (10 CFU/25μL) was stabilized

by a mechanical holder (VisualSonics). Ultrasound-guided intra-amniotic inoculation with *E.*

coli was performed in each amniotic sac using a 30G needle (BD PrecisionGlide needle; Becton

Dickinson, Franklin Lakes, NJ, USA). Controls were injected with 25 μL of sterile 1X PBS into

each amniotic sac. After the ultrasound injection, the dams were placed under a heat lamp for

recovery (defined as when the mouse resumed normal activities such as walking and

responding), which typically occurred 10 min after removal from anesthesia. After recovery,

mice were video monitored to observe pregnancy outcomes.

Video Monitoring

Pregnancy outcomes were recorded via video camera (Sony Corporation, Tokyo, Japan)

to determine gestational length, and therefore rate of preterm birth. Preterm birth was defined as

delivery occurring before 18.5 dpc, based on the earliest delivery of PBS-injected control dams,

and its rate was represented by the percentage of females delivering preterm among the total

number of mice injected.

Tissue collection prior to preterm birth

Dams were euthanized on 17.5 dpc and the reproductive tissues (uterus, decidua, and

cervix) were collected. Tissues collected for the preparation of single-cell suspensions were

placed in sterile 1X PBS, while tissues for histological analyses were fixed in 10% Neutral

Buffered Formalin (Surgipath, Leyca Biosystems, Wetzlar, Germany) and embedded in paraffin.

Five-µm-thick sections were cut and mounted on Superfrost® Plus microscope slides (Cat. No.

48311-703, VWR International, LLC. Radnor, PA, USA).

Histological characterization of murine reproductive tissues

Leukocyte detection using DAB immunohistochemistry

Five-µm-thick tissue sections from mice injected with PBS or *E. coli* were deparaffinized

and rehydrated using xylene and a series of decreasing ethanol concentrations, respectively.

Immunohistochemistry staining using the Monoclonal Rabbit Anti-Mouse CD45 (AB_2799780;

clone D3F8Q, cat. no. 70257S, Cell Signaling Technology, Danvers, MA, USA) was performed

using the Leica Bond Max Automatic Staining System in a peroxidase-mediated oxidation of

3,30-diaminobenzidine (DAB) from the Bond™ Polymer Refine Detection Kit (both from Leica

Microsystems, Wetzlar, Germany). The negative control used was the Rabbit FLEX Universal

Negative Control (cat. no. IR60066-2, Agilent, Santa Clara, CA, USA). Images were scanned

using the Brightfield setting of the Vectra Polaris Multispectral Imaging System.

Movat's pentachrome staining

Five-µm-thick tissue sections from mice injected with PBS or *E. coli* were histologically

characterized for the presence of collagen, elastin, muscle, and mucin using the Movat

Pentachrome Stain Kit (Modified Russell-Movat; ScyTek Laboratories, Inc. Logan, UT, USA),

following manufacturer's instructions with modifications. Briefly, tissue sections were

deparaffinized, stained with working Elastic Stain solution for 20 min, and rinsed in running tap

water for 1 min followed by rinsing with deionized water. Then, the following reagents from the

kit were sequentially applied to the entire tissue section with distilled water rinsing in between

each application: 2% Ferric Chloride for 5-8 s, 5% Sodium Thiosulfate Solution for 1 min,

Alcian Blue Solution (pH 2.5) for 20 min, Biebrich Scarlet-Acid Fuchsin Solution for 2 min, 5%

Phosphotungstic Acid Solution for 7 min, and 1% Acetic Acid Solution for 3 min. Excess Acetic

Acid Solution was drained from the slides and Yellow Stain Solution was immediately applied

for 20 min. The slides were then rinsed in 100% ethanol followed by rinsing with xylene. Images

were scanned using the Brightfield setting of the Vectra Polaris Multispectral Imaging System

(Akoya Biosciences, Marlborough, MA, USA).

OPAL multiplex immunofluorescence

OPAL multiplex immunofluorescence staining was performed using the OPAL Multiplex

7-color IHC kit (Cat. no. NEL811001KT; Akoya Biosciences), according to the manufacturer's

instructions. Prior to multiplex staining, the order of antibody staining was optimized using

single-plex staining paired with tyramide signal amplification (TSA)-conjugated OPAL

fluorophores. The optimized detection panel includes antibody-OPAL fluorophore pairs in the

following order: Monoclonal Rabbit Anti-Mouse F4/80 (AB_2799771; clone D2S9R; cat. no.

70076S, Cell Signaling Technology) with OPAL 520, Monoclonal Rabbit Anti-Mouse CD3ε

(AB_2889902; clone E4T1B; cat. no. 78588S, Cell Signaling Technology) with OPAL 570,

Monoclonal Rabbit Anti-Mouse Klrb1c/CD161c (AB_2892989; clone E6Y9G; cat. no. 39197S,

Cell Signaling Technology) with OPAL 620, Polyclonal Rabbit Anti-Mouse Ly6C (cat. no.

HA500088, HuaBio, Boston, MA, USA) with OPAL 650, and Monoclonal Rabbit Anti-Mouse

Ly6G (AB_2909808; clone E6Z1T; cat. no. 87048S, Cell Signaling Technology) with OPAL

690. The Rabbit FLEX Universal Negative Control (Agilent) was used as isotype. Briefly, 5-µm-

thick tissue sections from mice injected with PBS or *E. coli* were deparaffinized and rehydrated

using xylene and a series of decreasing ethanol concentrations, respectively. The slides were

rinsed in deionized water and epitope retrieval was performed by submerging the slides in

appropriate antigen retrieval (AR) buffer and boiling in a microwave oven. Non-specific binding

was prevented by incubating slides in OPAL antibody diluent/blocking solution prior to

incubating with each primary antibody at room temperature. Next, the slides were rinsed in

TBST prior to incubation with anti-mouse secondary antibody-horse radish peroxidase (HRP)

conjugate followed by the selected TSA-conjugated OPAL fluorophore. Cycles of sequential

epitope retrieval, target detection, and signal amplification were repeated using the optimized

antibody-OPAL fluorophore pair. Once all targets were detected, the slides were incubated with

DAPI (4′,6-diamidino-2-phenylindole) as a nuclear counterstain and mounted using AquaSlip™

Aqueous Permanent Mounting Medium (American MasterTech). Fluorescence image acquisition

was performed using the Vectra Polaris Multispectral Imaging System at 20x magnification.

Multispectral images were analyzed using the inForm software version 2.4 (Akoya Biosciences).

Tissue dissociation of murine uterus and decidua

Immediately following tissue collection, the uterus and decidua were dissociated to

prepare single-cell suspensions. The tissues were mechanically dissociated and enzymatically

digested by incubating at 37ºC using enzymes from the Umbilical Cord Dissociation Kit

(Miltenyi Biotec). A second round of mechanical dissociation was performed using the

gentleMACS Dissociator (Miltenyi Biotec), and dissociated cells were rinsed with 1X PBS

(Thermo Fisher Scientific) prior to filtration using a 100µm cell strainer (Miltenyi Biotec).

Filtered cells were pelleted by centrifugation at 300 x g for 5 min, erythrocytes were eliminated

using ACK Lysing Buffer (Life Technologies), and the cells were rinsed in 0.04% BSA (Sigma

Aldrich) and 0.5 mM EDTA (Sigma Aldrich) diluted in 1X PBS. Finally, the cells were filtered

using a 30µm cell strainer (Miltenyi Biotec), and the Dead Cell Removal Kit was used to remove

dead cells to obtain a cell viability of ≥80%.

Tissue dissociation of the murine cervix

Immediately following the collection of the cervix, the tissue was mechanically

dissociated and enzymatically digested using Collagenase A (160 mg/mL) (Sigma Aldrich, St.

Louis, MO, USA) and incubated at 37ºC. Then, the dissociated cells were pelleted by

centrifugation at 16,000 x g for 10 min at 20ºC and resuspended with 0.05% trypsin-EDTA

(Thermo Fisher Scientific, Waltham, MA) prior to a second round of mincing and incubation in

0.05% trypsin-EDTA at 37ºC. The enzymatic reaction was stopped by the addition of FBS (Fetal

Bovine Serum, Thermo Fisher). Cells were then filtered using a 70µm cell strainer (Miltenyi

Biotec, San Diego, CA, USA) and pelleted by centrifugation at 300 x g for 10 min. Erythrocytes

were removed using ACK Lysing Buffer (Life Technologies, Grand Island, NY, USA). Finally,

the cells were resuspended in 0.04% Bovine Serum Albumin (BSA) (Sigma Aldrich) diluted in

1X PBS (Thermo Fisher Scientific, NY, USA) and filtered through a 30µm cell strainer

(Miltenyi Biotec). The cell concentration and viability were determined using an automatic cell

counter (Cellometer Auto 2000, Nexcelom Bioscience, Lawrence, MA, USA) and the Dead Cell

Removal Kit (Miltenyi Biotec) was used to remove dead cells to obtain a cell viability of ≥80%.

Generation of gel beads-in-emulsion (GEMs) and library preparation

Generation of gel beads-in-emulsion (GEMs) and preparation of library constructs was

performed on viable single-cell suspensions using the 10x Genomics Chromium Single Cell 3'

Gene Expression Version 3.1 Kit (10x Genomics, Pleasanton, CA, USA), according to the

manufacturer's instructions. Briefly, viable single cells were encapsulated in partitioning oil

together with a single Gel Bead with barcoded oligonucleotides within the Chromium Controller.

Reverse transcription of mRNA into complementary (c)DNA was performed using the Veriti 96-

well Thermal Cycler (Thermo Fisher Scientific, Wilmington, DE, USA). Dynabeads MyOne

SILANE (10x Genomics) and the SPRIselect Reagent (Beckman Coulter, Indianapolis, IN,

USA) were used to purify resulting cDNA, which was optimized by enzymatic fragmentation,

end-repair, and A-tailing. Next, adaptors and sample index were incorporated by ligation. The

sample index PCR product was then amplified using the Veriti 96-well Thermal Cycler and

double-sided size selection was performed using the SPRIselect Reagent. Following the

formation of cDNA and final library construct, the Agilent Bioanalyzer High Sensitivity DNA

Chip (Agilent Technologies, Wilmington, DE, USA) was used determine sample quality and

concentration.

Sequencing

Prior to sequencing of post-library constructs, samples were quantified using the Kapa

DNA Quantification Kit for Illumina platforms (Kapa Biosystems, Wilmington, MA, USA),

889 following the manufacturer's instructions. The sequencing of 10x scRNA-seq libraries was

890 performed on the Illumina NextSeq 500 at the Genomics Services Center (GSC) of the Center

891 for Molecular Medicine and Genetics (Wayne State University School of Medicine, Detroit, MI,

892 USA). The Illumina 75 Cycle Sequencing Kit (Illumina, San Diego, CA, USA) was used with 58

893 cycles for R2, 26 for R1, and 8 for I1.

895 **QUANTIFICATION AND STATISTICAL ANALYSIS**

896 ***scRNA-seq data normalization and pre-processing***

897 Sequencing data were processed using Cell Ranger version 4.0.0 (10x Genomics). The

898 "cellranger counts" was also used to align the scRNA-seq reads by using the STAR

899 aligner(Dobin et al., 2013) to produce the bam files necessary for demultiplexing the individual

900 of origin based on genotype information using demuxlet(Kang et al., 2018) and a custom vcf file.

901 The genotype data were downloaded from ftp://ftp-

902 [mouse.sanger.ac.uk/current_snps/mgp.v5.merged.snps_all.dbSNP142.vcf.gz](ftp://ftp-mouse.sanger.ac.uk/current_snps/mgp.v5.merged.snps_all.dbSNP142.vcf.gz), the strains

903 C57BL_6NJ and BALB_cJ were extracted, and a new synthetic vcf file was generated consisting

904 of all the genetic variants where these two strains diverge, and containing a maternal genotype

905 column identical to the C57BL_6NJ strain and a fetal genotype column with a "0/1"

906 heterozygote genotype. Ambient RNA contamination and doublets were removed using SoupX

907 version 1.5.2(Young and Behjati, 2020) and DoubletFinder 2.0.3(McGinnis et al., 2019).

908 Additionally, any cell with < 200 genes or > 20,000 genes detected, or that had > 10%

909 mitochondrial reads, was excluded (Table S4). All count data matrices were then normalized and

910 combined using the Seurat package in R (Seurat version 4.0.3)(Hafemeister and Satija, 2019;

911 Stuart et al., 2019). The first 100 principal components were obtained, and the different libraries

were integrated and harmonized using the Harmony package in R version 1.0.0(Korsunsky et al.,

2019). The top 30 harmony components were then processed to embed and visualize the cells in

a two-dimensional map via the Uniform Manifold Approximation and Projection for Dimension

Reduction (UMAP) algorithm(Becht et al., 2018; McInnes et al., 2018). A resolution of 0.5 was

used to cluster the single cells.

Annotation of cell types

The SingleR(Aran et al., 2019) package in R version 1.6.1 was used to annotate cell types

based on their similarities to reference datasets with known labels(Buechler et al., 2021; Tabula

Muris et al., 2018). SingleR annotates single cells from query data by computing the Spearman's

correlation coefficient between the single-cell gene expression data and samples from the

reference dataset. The correlation is measured only based on the variable genes in the reference

dataset. The multiple correlation coefficients per cell type are combined according to the cell

type labels of the reference dataset to assign a score per cell type. Additionally, we confirmed the

cell type identities by identifying the top DEGs (see below) and the gene-cell type mapping data

provided by the Mouse Cell Atlas and single-cell MCA (scMCA) package(Han et al., 2018) in R

version 0.2.0. Using different annotations obtained from the reference mapping workflows, the

final cell type labels were assigned based on a majority vote. If multiple clusters were assigned

to the same consensus cell type, we added a sub-index to that cell type for each different original

Seurat cluster: Clusters 0, 1, and 21 were annotated as Fibroblast-1, Fibroblast-2, and Fibroblast-

3; clusters 2, 4, and 12 were annotated as Stromal-1, Stromal-2 (Decidua), and Stromal-3;

clusters 5, 7, 8, 10, 11, 13, 14, 20, 23, and 28 were annotated as Epithelial-1 (basal), Epithelial-2

(squamous), Epithelial-3 (squamous), Epithelial-4 (glandular), Epithelial-10 (proliferative),

Epithelial-5 (luminal), Epithelial-6 (secretory), Epithelial-7 (glandular), Epithelial-8

(Enterocyte), and Epithelial-9 (Secretory); clusters 9 and 24 were annotated as Macrophage-1

and Macrophage-2 (progenitor); clusters 15 and 26 were annotated as SMC-1 and SMC-2; and

clusters 19 and 27 were annotated as NK-cell-1 and NK-cell-2. All remaining clusters were

assigned a unique cell type identifier (Table S5).

Differential gene expression for cell type analysis

For this analysis, the differential expression of selected marker genes for each cell

type/cluster was identified using the Wilcoxon Rank Sum test and the FindAllMarkers function

from Seurat (Table S5). For this analysis, we first compared each cluster to all cell types. We

further used the top cell markers [ranked based on $\log_2$(Fold change) and requiring q < 0.1]

assigned to each sub-cluster to annotate the clusters using the Mouse Cell Atlas and scMCA

package(Han et al., 2018).

Differential gene expression in preterm labor

The identification of preterm labor-associated DEGs between study groups was

performed using the DESeq2 R package version 1.32.0(Love et al., 2014). A term for each

library was added to the DESeq2 model to correct for technical batch effects (library identifier).

For each cell type/replicate combination, we only used combinations with more than 20 cells;

otherwise, it was treated as non-observed. Cell types found in < 3 combinations per study group

were dropped from the differential gene expression analysis (Table S2 contains all genes

determined as differentially expressed). Note that these thresholds imply that clusters with < 120

cells are not analyzed to ensure robust gene expression estimation. Quantile-quantile plots were

used to show that p-values are well calibrated under the null hypothesis of no effect of preterm

labor, and also to show which tissues and cell types are more enriched for preterm labor-

associated gene expression changes (Figure 2H-J). Multiple comparison correction was

performed by controlling for false discovery rate using Benjamini-Hochberg's method and genes

with $q < 0.1$ were reported in Figure 2E-G and Table S2. Statistical difference between the

fraction of genes that were upregulated versus downregulated by preterm labor in each cell type

was assessed with a binomial test and corrected for multiple comparisons using Benjamini-

Hochberg's method. To compare the effects of preterm labor on gene expression across different

tissues and cell types, we performed Spearman's correlation between the $\log_2$FC obtained in each

DESeq2 analysis performed using genes that had been detected as differentially expressed in at

least one cell-type/tissue, $q < 0.1$. These correlations were visualized as a heatmap in Figure

S5A, S6A, and S7A and in boxplots for relevant tissue and cell-type combinations in Figure 3B-

D.

Gene ontology and pathway enrichment analysis of genes affected by preterm labor

The clusterProfiler in R version 4.0.4(Yu et al., 2012) was used to perform the Over-

Representation Analysis (ORA) separately for each list of genes obtained as differentially

expressed for each cell type based on the Gene Ontology (GO), Kyoto Encyclopedia of Gene and

Genomes (KEGG), and Reactome databases. The functions "enrichPathway", "enrichKEGG",

and "enrichGO", from "clusterProfiler" were used. In ORA analyses, the universe of genes for

each cell type was the subset that was expressed at a level sufficient to be tested in differential

gene expression analysis. When results are combined across cell types, any genes tested (with a

calculated p-value) in any of the cell types are used for the universe. Only ORA results that were

significant after correction were reported with $q < 0.05$ being considered statistically significant.

Cell-cell communication analysis

CellChat (Jin et al., 2021) was used to infer the cell-cell communications using the

single-cell gene expression data from preterm labor and control conditions and a database of

prior knowledge of the interactions between signaling ligands, receptors, and their cofactors. The

top 25% of significant cell-cell communications ($p < 0.05$) across different pathways were shown

for the two conditions of preterm labor and control. Next, the aggregated cell-cell

communication between different cell groups was calculated for the two study groups, and the

interaction strength was compared among different cell types from the two study groups. The

differential interaction strength was represented with circle plots with red (or blue) edges

showing the increased (or decreased) signaling in preterm labor compared to controls.

Additionally, the detailed differential interaction strengths were shown using heatmap

representations. Major signaling sender and receiver cells were displayed using scatter plots

where the changes in signaling strength from control to preterm labor were represented by

arrows. The R packages CellChat version 1.1.2, ggalluvial version 0.12.3, and ggplot2 version

3.3.5 were used to visualize cell-cell communication analyses. The major sending and receiving

signaling roles based on context-specific pathways across different cell groups were identified

using a cut-off of 0.5 when visualizing the connection. The overall information flow [sum of the

significant communication probability ($p < 0.05$) in the inferred cell-cell network] for each

signaling network was represented using a bar plot. The comparison between the overall

information flow from the two study groups (preterm labor and control) was performed using the

paired Wilcoxon test with the function "rankNet" from CellChat.

1004

1005   ***Comparison between cell-cell communication in human and murine uterine tissues***

1006   We inferred cell-cell communications using the human myometrial single-cell gene

1007   expression data from term in labor (TIL) and term without labor (TNL) study groups(Pique-Regi

1008   et al., 2022), and compared the inferred interactions between mouse (uterus) and human

1009   (myometrium) across the top common signaling pathways with highest numbers of DEGs.

1010

Statistical analysis

Observational mouse data were analyzed by using SPSS v19.0 and GraphPad Prism

version 8. For comparing the rates of preterm birth, the Fisher's exact test was used. For

gestational length and cervical shortening, the statistical significance of group comparisons was

assessed using the Mann-Whitney U-test or paired t-test, respectively.

SUPLEMENTAL INFORMATION

Supplementary Figure Legends

Figure S1. Preterm labor induced by intra-amniotic infection alters the cellular

composition of the murine reproductive tissues, related to Figure 1. (A) UMAP plots

showing the cell types present in the uterus, decidua, and cervix of control mice and **(B)** mice

with preterm labor. **(C-E)** UMAP plots showing the distribution of cells according to fetal

(purple) or maternal (grey) origin in the uterus, decidua, and cervix. Abbreviations used: SMC,

smooth muscle cell; NK cell, natural killer cell.

1024

Figure S2. Leukocyte infiltration in the uterus, decidua, and cervix, related to Figure 2. (A-

C) Representative Movat pentachrome staining images of the uterus, decidua, and cervix from

control (top row) and preterm labor (bottom row) mice. Red staining indicates muscle/fibrin,

dark purple staining indicates elastic fibers, blue staining indicates mucin, and yellow indicates

collagen/reticular fibers. Nuclei appear as dark blue/black. Whole-slide images taken at (A)

control: 3.7X, preterm labor 4X; (B) control: 2.9X, preterm labor: 4.9X; (C) control: 3.2 X,

preterm labor: 3.1X (scale bars = 300µm). Zoomed images were all taken at 20X magnification

(scale bars = 50µm). **(D-F)** Representative images showing 3, 3'diaminobenzidine (DAB)

immunohistochemistry to detect the pan-leukocyte marker CD45 in the uterus, decidua, and

cervix of control mice (top row) and preterm labor (bottom row) mice (n = 3 per group). Whole-

slide brightfield images taken at (E) control: 3.5X, preterm labor: 3.3X; (F) control: 2.9X,

preterm labor 3.9X; (G) control: 3.4X, preterm labor: 3.2X (scale bars = 300µm). Zoomed

images were taken at 20X magnification (scale bars = 50µm). **(G-I)** Representative merged

image showing the co-localized immunofluorescence detection of neutrophils (Ly6G+ cells,

pink), monocytes (Ly6C+ cells, cyan), macrophages (F4/80+ cells, red), T cells (CD3+ cells,

yellow) and NK cells (CD161c+ cells, green) in the uterus, decidua, and cervix of control and

preterm labor mice only (n = 3). Nuclear staining is shown in blue (4',6-diamidino-2-

phenylindole; DAPI). Images were taken at 20X magnification. Scale bar = 100μm.

1043

Figure S3. Shared and unique biological processes in specific cell types impacted by labor

across tissues, related to Figure 3. (**A**) Box plots showing the Spearman's correlation of the

preterm labor-associated $\log_2$(Fold change, FC) among cell types within each tissue. (**B**) Box

plots showing the Spearman's correlations between tissue pairs for the preterm labor-associated

logFC in cell types present in the two tissues. (**C**) ClusterProfiler dot plot showing preterm labor-

associated Kyoto Encyclopedia of Genes and Genomes (KEGG) pathways enriched in specific

cell 588 types in the uterus, decidua, and cervix. The size and color of each dot represents

enrichment score and significance level, respectively. Significant KEGG pathways (q < 0.05)

were identified based on over-representation analysis using one-sided Fisher's exact tests.

Abbreviations used: SMC, smooth muscle cell; NK cell, natural killer cell.

1054

Figure S4. The uterus and decidua share enrichment of biological processes in preterm

labor, related to Figure 3. Cluster profiler dot plots showing the Gene Ontology (GO)

biological processes enriched with preterm labor in (**A**) Stromal-1 and Stromal-2, (**B**) Stromal-3

and Fibroblast-1, (**C**) Fibroblast-2 and Fibroblast-3, and (**D**) Endothelial cells in the uterus and

decidua. The size and color of each dot represent gene ratio and significance level, respectively.

A 1-sided Fisher's exact test was used.

1061

Figure S5. Preterm labor-associated enrichment of signaling pathways in uterine cell types,

related to Figure 4. (A) Heatmap showing correlations among uterine cell types where red and

white blocks signify increased and decreased correlation, respectively. Pearson correlation tests

was used. **(B)** Bar plots showing the high expression of specific signaling pathways in the uterus

of control mice (blue bars/pathway names) or mice with preterm labor (red bars/pathway names).

(C) Forest plot showing the $\log_2$(FC, fold change) and 95% confidence intervals of differentially

expressed (DEGs) across cell types in murine uterus. DEGs shown are significant with FDR (q <

0.1). Abbreviations used: SMC, smooth muscle cell; NK cell, natural killer cell.

1070

1071 **Figure S6. Preterm labor-induced induced changes in the expression of prostaglandin-**

1072 **associated genes in different cell types across the reproductive tissues, related to Figures 4-**

1073 **6.** Forest plot showing the $\log_2$(FC, fold change) and 95% confidence intervals of differentially

1074 expressed (DEGs) in selected cell types across the murine cervix, decidua and uterus. DEGs

1075 shown are significant with FDR (q < 0.01).

1076

Figure S7. Preterm labor-induced changes in interaction between decidual cell types,

related to Figure 5. (A) Heatmap showing correlations among decidual cell types where red and

white blocks signify increased and decreased correlation, respectively. Pearson correlation tests

was used. **(B)** Bar plots showing the high expression of specific signaling pathways in the

decidua of control mice (blue bars/pathway names) or mice with preterm labor (red bars/pathway

names). **(C)** Forest plot showing the $\log_2$(FC, fold change) and 95% confidence intervals of

differentially expressed (DEGs) across cell types in murine decidua. DEGs shown are significant

with FDR (q < 0.1). Abbreviations used: SMC, smooth muscle cell; NK cell, natural killer cell.

1085

1086 **Figure S8. Preterm labor-induced changes in interaction between cervical cell types,**

1087 **related to Figure 6. (A)** Heatmap showing correlations among cervical cell types where red and

1088 white blocks signify increased and decreased correlation, respectively. Pearson correlation tests

1089 was used. **(B)** Bar plots showing the high expression of specific signaling pathways in the cervix

1090 of control mice (blue bars/pathway names) or mice with preterm labor (red bars/pathway names).

1091

Figure S9. Labor-associated signaling in the human myometrium partially overlaps with

preterm labor-associated changes in the murine uterus, related to Figure 7. (A) Circle plots

showing the top aggregated interactions among cell types in the myometrium from humans

without (left) or with labor at term (right). Each node represents a cell type and the interaction is

shown by lines color-coded based on the sender cell. Representation of aggregated interactions

with p < 0.05 using cell chat. **(B)** Circle plot showing the increased (red) or decreased (blue)

signaling interactions in the human myometrium in labor compared to controls without labor.

Representation of top 25% differential interaction strength. **(C)** Venn diagram showing the

overlap in upregulated signaling pathways between the murine uterus in preterm labor (left

circle, pink) and the human myometrium in term labor (right circle, orange). Shared labor- and

inflammation-associated pathways are highlighted in red. **(D-E)** Circle plots representing the top

25% human myometrial cell-cell communications inferred for the Complement and CXCL

pathways for the Term not in labor and Term in labor groups. **(F-G)** Circle plots representing the

top 25% murine uterine cell-cell communications inferred for the Complement and CXCL

pathways for the control and preterm labor groups. Abbreviations used: SMC, smooth muscle

cell; NK cell, natural killer cell; EVT, extravillous trophoblast; ILC, innate lymphoid cell; LED,

lymphoid endothelial decidual cell.

SUPPLEMENTAL EXCEL TABLE TITLES

Table S1. Summary of the numbers of uterine, decidual and cervical cells encapsulated within

the 10x Genomics Gel-bead-in-emulsion (GEM), related to Figure 2.

1112

1113 **Table S2.** Preterm labor-associated differentially expressed genes in uterine, decidual and

1114 cervical cell types, as well as the total differentially expressed genes across tissue, related to

1115 Figures 2 and 3.

1116

1117 **Table S3.** Summary of genes and pathways used in the CellChat intercellular communication

1118 analysis for the uterus, decidua and cervix, related to Figures 4, 5, and 6.

1119

1120 **Table S4.** Summary of quality control metrics calculated with the 10x Genomics Cell Ranger

1121 pipeline for each library, related to methods section ***"scRNA-seq data normalization and pre-***

1122 ***processing"***.

1123

1124 **Table S5.** List of marker genes for the identification of cell types in the murine uterus, decidua

1125 and cervix, related to methods section ***"Annotation of cell types"***.

REFERENCES

Agrawal, V., and Hirsch, E. (2012). Intrauterine infection and preterm labor. Semin Fetal Neonatal Med *17*, 12-19. 10.1016/j.siny.2011.09.001.

Aran, D., Looney, A.P., Liu, L., Wu, E., Fong, V., Hsu, A., Chak, S., Naikawadi, R.P., Wolters, P.J., Abate, A.R., et al. (2019). Reference-based analysis of lung single-cell sequencing reveals a transitional profibrotic macrophage. Nat Immunol *20*, 163-172. 10.1038/s41590-018-0276-y.

Arenas-Hernandez, M., Romero, R., Xu, Y., Panaitescu, B., Garcia-Flores, V., Miller, D., Ahn, H., Done, B., Hassan, S.S., Hsu, C.D., et al. (2019). Effector and Activated T Cells Induce Preterm Labor and Birth That Is Prevented by Treatment with Progesterone. J Immunol *202*, 2585-2608. 10.4049/jimmunol.1801350.

Bartmann, C., Segerer, S.E., Rieger, L., Kapp, M., Sütterlin, M., and Kämmerer, U. (2014). Quantification of the predominant immune cell populations in decidua throughout human pregnancy. Am J Reprod Immunol *71*, 109-119. 10.1111/aji.12185.

Bastek, J.A., Gómez, L.M., and Elovitz, M.A. (2011). The role of inflammation and infection in preterm birth. Clin Perinatol *38*, 385-406. 10.1016/j.clp.2011.06.003.

Becht, E., McInnes, L., Healy, J., Dutertre, C.A., Kwok, I.W.H., Ng, L.G., Ginhoux, F., and Newell, E.W. (2018). Dimensionality reduction for visualizing single-cell data using UMAP. Nat Biotechnol. 10.1038/nbt.4314.

Boros-Rausch, A., Shynlova, O., and Lye, S.J. (2021). A Broad-Spectrum Chemokine Inhibitor Blocks Inflammation-Induced Myometrial Myocyte-Macrophage Crosstalk and Myometrial Contraction. Cells *11*. 10.3390/cells11010128.

Buechler, M.B., Pradhan, R.N., Krishnamurty, A.T., Cox, C., Calviello, A.K., Wang, A.W., Yang, Y.A., Tam, L., Caothien, R., Roose-Girma, M., et al. (2021). Cross-tissue organization of the fibroblast lineage. Nature *593*, 575-579. 10.1038/s41586-021-03549-5.

Bukowski, R., Hankins, G.D., Saade, G.R., Anderson, G.D., and Thornton, S. (2006). Labor-associated gene expression in the human uterine fundus, lower segment, and cervix. PLoS Med *3*, e169. 10.1371/journal.pmed.0030169.

Burnham, P., Gomez-Lopez, N., Heyang, M., Cheng, A.P., Lenz, J.S., Dadhania, D.M., Lee, J.R., Suthanthiran, M., Romero, R., and De Vlaminck, I. (2020). Separating the signal from the noise in metagenomic cell-free DNA sequencing. Microbiome *8*, 18. 10.1186/s40168-020-0793-4.

Chawanpaiboon, S., Vogel, J.P., Moller, A.B., Lumbiganon, P., Petzold, M., Hogan, D., Landoulsi, S., Jampathong, N., Kongwattanakul, K., Laopaiboon, M., et al. (2019). Global, regional, and national estimates of levels of preterm birth in 2014: a systematic review and modelling analysis. Lancet Glob Health *7*, e37-e46. 10.1016/s2214-109x(18)30451-0.

Chumduri, C., Gurumurthy, R.K., Berger, H., Dietrich, O., Kumar, N., Koster, S., Brinkmann, V., Hoffmann, K., Drabkina, M., Arampatzi, P., et al. (2021). Opposing Wnt signals regulate cervical squamocolumnar homeostasis and emergence of metaplasia. Nat Cell Biol *23*, 184-197. 10.1038/s41556-020-00619-0.

Coleman, M., Orvis, A., Wu, T.Y., Dacanay, M., Merillat, S., Ogle, J., Baldessari, A., Kretzer, N.M., Munson, J., Boros-Rausch, A.J., et al. (2020). A Broad Spectrum Chemokine Inhibitor Prevents Preterm Labor but Not Microbial Invasion of the Amniotic Cavity or Neonatal Morbidity in a Non-human Primate Model. Front Immunol *11*, 770. 10.3389/fimmu.2020.00770.

Combs, C.A., Gravett, M., Garite, T.J., Hickok, D.E., Lapidus, J., Porreco, R., Rael, J., Grove, T., Morgan, T.K., Clewell, W., et al. (2014). Amniotic fluid infection, inflammation, and

colonization in preterm labor with intact membranes. Am J Obstet Gynecol *210*, 125 e121-125 e115. 10.1016/j.ajog.2013.11.032.

Danforth, D.N., and Ivy, A.C. (1949). The lower uterine segment; its derivation and physiologic behavior. Am J Obstet Gynecol *57*, 831-841. 10.1016/0002-9378(49)90639-0.

DiGiulio, D.B., Romero, R., Kusanovic, J.P., Gómez, R., Kim, C.J., Seok, K.S., Gotsch, F., Mazaki-Tovi, S., Vaisbuch, E., Sanders, K., et al. (2010). Prevalence and diversity of microbes in the amniotic fluid, the fetal inflammatory response, and pregnancy outcome in women with preterm pre-labor rupture of membranes. Am J Reprod Immunol *64*, 38-57. 10.1111/j.1600-0897.2010.00830.x.

Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M., and Gingeras, T.R. (2013). STAR: ultrafast universal RNA-seq aligner. Bioinformatics *29*, 15-21. 10.1093/bioinformatics/bts635.

Dobyns, A.E., Goyal, R., Carpenter, L.G., Freeman, T.C., Longo, L.D., and Yellon, S.M. (2015). Macrophage gene expression associated with remodeling of the prepartum rat cervix: microarray and pathway analyses. PLoS One *10*, e0119782. 10.1371/journal.pone.0119782.

Dubicke, A., Ekman-Ordeberg, G., Mazurek, P., Miller, L., and Yellon, S.M. (2016). Density of Stromal Cells and Macrophages Associated With Collagen Remodeling in the Human Cervix in Preterm and Term Birth. Reprod Sci *23*, 595-603. 10.1177/1933719115616497.

Faro, J., Romero, R., Schwenkel, G., Garcia-Flores, V., Arenas-Hernandez, M., Leng, Y., Xu, Y., Miller, D., Hassan, S.S., and Gomez-Lopez, N. (2019). Intra-amniotic inflammation induces preterm birth by activating the NLRP3 inflammasome. Biol Reprod *100*, 1290-1305. 10.1093/biolre/ioy261.

Galaz, J., Romero, R., Slutsky, R., Xu, Y., Motomura, K., Para, R., Pacora, P., Panaitescu, B., Hsu, C.D., Kacerovsky, M., and Gomez-Lopez, N. (2020a). Cellular immune responses in amniotic fluid of women with preterm prelabor rupture of membranes. J Perinat Med *48*, 222-233. 10.1515/jpm-2019-0395.

Galaz, J., Romero, R., Xu, Y., Miller, D., Slutsky, R., Levenson, D., Hsu, C.D., and Gomez-Lopez, N. (2020b). Cellular immune responses in amniotic fluid of women with preterm clinical chorioamnionitis. Inflamm Res *69*, 203-216. 10.1007/s00011-019-01308-x.

Gershater, M., Romero, R., Arenas-Hernandez, M., Galaz, J., Motomura, K., Tao, L., Xu, Y., Miller, D., Pique-Regi, R., Martinez, G., 3rd, et al. (2022). IL-22 Plays a Dual Role in the Amniotic Cavity: Tissue Injury and Host Defense against Microbes in Preterm Labor. J Immunol. 10.4049/jimmunol.2100439.

Gibbs, R.S., Blanco, J.D., St Clair, P.J., and Castaneda, Y.S. (1982). Quantitative bacteriology of amniotic fluid from women with clinical intraamniotic infection at term. J Infect Dis *145*, 1-8. 10.1093/infdis/145.1.1.

Gibbs, R.S., Romero, R., Hillier, S.L., Eschenbach, D.A., and Sweet, R.L. (1992). A review of premature birth and subclinical infection. Am J Obstet Gynecol *166*, 1515-1528. 10.1016/0002-9378(92)91628-n.

Goldenberg, R.L., Culhane, J.F., Iams, J.D., and Romero, R. (2008). Epidemiology and causes of preterm birth. Lancet *371*, 75-84. 10.1016/s0140-6736(08)60074-4.

Gomez-Lopez, N., Estrada-Gutierrez, G., Jimenez-Zamudio, L., Vega-Sanchez, R., and Vadillo-Ortega, F. (2009). Fetal membranes exhibit selective leukocyte chemotaxic activity during human labor. J Reprod Immunol *80*, 122-131. 10.1016/j.jri.2009.01.002.

Gomez-Lopez, N., Garcia-Flores, V., Chin, P.Y., Groome, H.M., Bijland, M.T., Diener, K.R., Romero, R., and Robertson, S.A. (2021a). Macrophages exert homeostatic actions in pregnancy

to protect against preterm birth and fetal inflammatory injury. JCI Insight *6*. 10.1172/jci.insight.146089.

Gomez-Lopez, N., Romero, R., Arenas-Hernandez, M., Panaitescu, B., Garcia-Flores, V., Mial, T.N., Sahi, A., and Hassan, S.S. (2018a). Intra-amniotic administration of lipopolysaccharide induces spontaneous preterm labor and birth in the absence of a body temperature change. J Matern Fetal Neonatal Med *31*, 439-446. 10.1080/14767058.2017.1287894.

Gomez-Lopez, N., Romero, R., Galaz, J., Bhatti, G., Done, B., Miller, D., Ghita, C., Motomura, K., Farias-Jofre, M., Jung, E., et al. (2022). Transcriptome changes in maternal peripheral blood during term parturition mimic perturbations preceding spontaneous preterm birth. Biol Reprod *106*, 185-199. 10.1093/biolre/ioab197.

Gomez-Lopez, N., Romero, R., Galaz, J., Xu, Y., Panaitescu, B., Slutsky, R., Motomura, K., Gill, N., Para, R., Pacora, P., et al. (2019a). Cellular immune responses in amniotic fluid of women with preterm labor and intra-amniotic infection or intra-amniotic inflammation. Am J Reprod Immunol *82*, e13171. 10.1111/aji.13171.

Gomez-Lopez, N., Romero, R., Leng, Y., Garcia-Flores, V., Xu, Y., Miller, D., and Hassan, S.S. (2017a). Neutrophil extracellular traps in acute chorioamnionitis: A mechanism of host defense. Am J Reprod Immunol *77*. 10.1111/aji.12617.

Gomez-Lopez, N., Romero, R., Leng, Y., Xu, Y., Slutsky, R., Levenson, D., Pacora, P., Jung, E., Panaitescu, B., and Hsu, C.D. (2019b). The origin of amniotic fluid monocytes/macrophages in women with intra-amniotic inflammation or infection. J Perinat Med *47*, 822-840. 10.1515/jpm-2019-0262.

Gomez-Lopez, N., Romero, R., Varrey, A., Leng, Y., Miller, D., Done, B., Xu, Y., Bhatti, G., Motomura, K., Gershater, M., et al. (2021b). RNA Sequencing Reveals Diverse Functions of Amniotic Fluid Neutrophils and Monocytes/Macrophages in Intra-Amniotic Infection. J Innate Immun *13*, 63-82. 10.1159/000509718.

Gomez-Lopez, N., Romero, R., Xu, Y., Leng, Y., Garcia-Flores, V., Miller, D., Jacques, S.M., Hassan, S.S., Faro, J., Alsamsam, A., et al. (2017b). Are amniotic fluid neutrophils in women with intraamniotic infection and/or inflammation of fetal or maternal origin? Am J Obstet Gynecol *217*, 693.e691-693.e616. 10.1016/j.ajog.2017.09.013.

Gomez-Lopez, N., Romero, R., Xu, Y., Miller, D., Arenas-Hernandez, M., Garcia-Flores, V., Panaitescu, B., Galaz, J., Hsu, C.D., Para, R., and Berry, S.M. (2019c). Fetal T Cell Activation in the Amniotic Cavity during Preterm Labor: A Potential Mechanism for a Subset of Idiopathic Preterm Birth. J Immunol *203*, 1793-1807. 10.4049/jimmunol.1900621.

Gomez-Lopez, N., Romero, R., Xu, Y., Miller, D., Leng, Y., Panaitescu, B., Silva, P., Faro, J., Alhousseini, A., Gill, N., et al. (2018b). The immunophenotype of amniotic fluid leukocytes in normal and complicated pregnancies. Am J Reprod Immunol *79*, e12827. 10.1111/aji.12827.

Gomez-Lopez, N., Vadillo-Perez, L., Hernandez-Carbajal, A., Godines-Enriquez, M., Olson, D.M., and Vadillo-Ortega, F. (2011a). Specific inflammatory microenvironments in the zones of the fetal membranes at term delivery. Am J Obstet Gynecol *205*, 235.e215-224. 10.1016/j.ajog.2011.04.019.

Gomez-Lopez, N., Vadillo-Perez, L., Nessim, S., Olson, D.M., and Vadillo-Ortega, F. (2011b). Choriodecidua and amnion exhibit selective leukocyte chemotaxis during term human labor. Am J Obstet Gynecol *204*, 364.e369-316. 10.1016/j.ajog.2010.11.010.

Gomez-Lopez, N., Vega-Sanchez, R., Castillo-Castrejon, M., Romero, R., Cubeiro-Arreola, K., and Vadillo-Ortega, F. (2013). Evidence for a role for the adaptive immune response in human term parturition. Am J Reprod Immunol *69*, 212-230. 10.1111/aji.12074.

Gomez, R., Ghezzi, F., Romero, R., Muñoz, H., Tolosa, J.E., and Rojas, I. (1995). Premature
labor and intra-amniotic infection. Clinical aspects and role of the cytokines in diagnosis and
pathophysiology. Clin Perinatol *22*, 281-342.

Gomez, R., Romero, R., Galasso, M., Behnke, E., Insunza, A., and Cotton, D.B. (1994). The
value of amniotic fluid interleukin-6, white blood cell count, and gram stain in the diagnosis of
microbial invasion of the amniotic cavity in patients at term. Am J Reprod Immunol *32*, 200-210.
10.1111/j.1600-0897.1994.tb01115.x.

Gonçalves, L.F., Chaiworapongsa, T., and Romero, R. (2002). Intrauterine infection and
prematurity. Ment Retard Dev Disabil Res Rev *8*, 3-13. 10.1002/mrdd.10008.

Gonzalez, J.M., Dong, Z., Romero, R., and Girardi, G. (2011). Cervical remodeling/ripening at
term and preterm delivery: the same mechanism initiated by different mediators and different
effector cells. PLoS One *6*, e26877. 10.1371/journal.pone.0026877.

Granstrom, L., Ekman, G., Ulmsten, U., and Malmstrom, A. (1989). Changes in the connective
tissue of corpus and cervix uteri during ripening and labour in term pregnancy. Br J Obstet
Gynaecol *96*, 1198-1202. 10.1111/j.1471-0528.1989.tb03196.x.

Gravett, M.G., Hummel, D., Eschenbach, D.A., and Holmes, K.K. (1986). Preterm labor
associated with subclinical amniotic fluid infection and with bacterial vaginosis. Obstet Gynecol
67, 229-237. 10.1097/00006250-198602000-00013.

Guzick, D.S., and Winn, K. (1985). The association of chorioamnionitis with preterm delivery.
Obstet Gynecol *65*, 11-16.

Hafemeister, C., and Satija, R. (2019). Normalization and variance stabilization of single-cell
RNA-seq data using regularized negative binomial regression. Genome Biol *20*, 296.
10.1186/s13059-019-1874-1.

Han, X., Wang, R., Zhou, Y., Fei, L., Sun, H., Lai, S., Saadatpour, A., Zhou, Z., Chen, H., Ye,
F., et al. (2018). Mapping the Mouse Cell Atlas by Microwell-Seq. Cell *172*, 1091-1107 e1017.
10.1016/j.cell.2018.02.001.

Hillier, S.L., Martius, J., Krohn, M., Kiviat, N., Holmes, K.K., and Eschenbach, D.A. (1988). A
case-control study of chorioamnionic infection and histologic chorioamnionitis in prematurity. N
Engl J Med *319*, 972-978. 10.1056/nejm198810133191503.

Huang, J., Li, Q., Peng, Q., Xie, Y., Wang, W., Pei, C., Zhao, Y., Liu, R., Huang, L., Li, T., et al.
(2021). Single-cell RNA sequencing reveals heterogeneity and differential expression of
decidual tissues during the peripartum period. Cell Prolif *54*, e12967. 10.1111/cpr.12967.

Huang, J.R., Nie, J., Liu, L.J., Zhang, X.W., Xie, Y.M., Peng, Q.Z., Wang, W.N., Pei, C.L.,
Zhao, Y.H., Liu, R., et al. (2020). Single-cell transcriptomics reveals the heterogeneity of the
decidual endothelial cells that participate in labor onset. Eur Rev Med Pharmacol Sci *24*, 10359-
10365. 10.26355/eurrev_202010_23385.

Jin, S., Guerrero-Juarez, C.F., Zhang, L., Chang, I., Ramos, R., Kuan, C.H., Myung, P., Plikus,
M.V., and Nie, Q. (2021). Inference and analysis of cell-cell communication using CellChat. Nat
Commun *12*, 1088. 10.1038/s41467-021-21246-9.

Kang, H.M., Subramaniam, M., Targ, S., Nguyen, M., Maliskova, L., McCarthy, E., Wan, E.,
Wong, S., Byrnes, L., Lanata, C.M., et al. (2018). Multiplexed droplet single-cell RNA-
sequencing using natural genetic variation. Nat Biotechnol *36*, 89-94. 10.1038/nbt.4042.

Keelan, J.A., Blumenstein, M., Helliwell, R.J., Sato, T.A., Marvin, K.W., and Mitchell, M.D.
(2003). Cytokines, prostaglandins and parturition--a review. Placenta *24 Suppl A*, S33-46.
10.1053/plac.2002.0948.

Keski-Nisula, L.T., Aalto, M.L., Kirkinen, P.P., Kosma, V.M., and Heinonen, S.T. (2003). Myometrial inflammation in human delivery and its association with labor and infection. Am J Clin Pathol *120*, 217-224. 10.1309/kc6k-dtx9-8lfy-b3j7.

Kim, C.J., Romero, R., Chaemsaithong, P., Chaiyasit, N., Yoon, B.H., and Kim, Y.M. (2015a). Acute chorioamnionitis and funisitis: definition, pathologic features, and clinical significance. Am J Obstet Gynecol *213*, S29-52. 10.1016/j.ajog.2015.08.040.

Kim, M.J., Romero, R., Gervasi, M.T., Kim, J.S., Yoo, W., Lee, D.C., Mittal, P., Erez, O., Kusanovic, J.P., Hassan, S.S., and Kim, C.J. (2009). Widespread microbial invasion of the chorioamniotic membranes is a consequence and not a cause of intra-amniotic infection. Lab Invest *89*, 924-936. 10.1038/labinvest.2009.49.

Kim, S.M., Romero, R., Park, J.W., Oh, K.J., Jun, J.K., and Yoon, B.H. (2015b). The relationship between the intensity of intra-amniotic inflammation and the presence and severity of acute histologic chorioamnionitis in preterm gestation. J Matern Fetal Neonatal Med *28*, 1500-1509. 10.3109/14767058.2014.961009.

Koh, W., Wu, A., Penland, L., Treutlein, B., Neff, N.F., Mantalas, G.L., Blumenfeld, Y.J., El-Sayed, Y.Y., Stevenson, D.K., Shaw, G.M., and Quake, S.R. (2019). Single Cell Transcriptomes Derived from Human Cervical and Uterine Tissue during Pregnancy. Adv Biosyst *3*, e1800336. 10.1002/adbi.201800336.

Korsunsky, I., Millard, N., Fan, J., Slowikowski, K., Zhang, F., Wei, K., Baglaenko, Y., Brenner, M., Loh, P.R., and Raychaudhuri, S. (2019). Fast, sensitive and accurate integration of single-cell data with Harmony. Nat Methods *16*, 1289-1296. 10.1038/s41592-019-0619-0.

Krantz, K.E., and Phillips, W.P. (1962). Anatomy of the human uterine cervix, gross and microscopic. Ann N Y Acad Sci *97*, 551-563. 10.1111/j.1749-6632.1962.tb34666.x.

Li, Y., Lopez, G.E., Vazquez, J., Sun, Y., Chavarria, M., Lindner, P.N., Fredrickson, S., Karst, N., and Stanic, A.K. (2018). Decidual-Placental Immune Landscape During Syngeneic Murine Pregnancy. Front Immunol *9*, 2087. 10.3389/fimmu.2018.02087.

Liu, L., Li, H., Dargahi, D., Shynlova, O., Slater, D., Jones, S.J., Lye, S.J., and Dong, X. (2015a). HoxA13 Regulates Phenotype Regionalization of Human Pregnant Myometrium. J Clin Endocrinol Metab *100*, E1512-1522. 10.1210/jc.2015-2815.

Liu, L., Oza, S., Hogan, D., Perin, J., Rudan, I., Lawn, J.E., Cousens, S., Mathers, C., and Black, R.E. (2015b). Global, regional, and national causes of child mortality in 2000-13, with projections to inform post-2015 priorities: an updated systematic analysis. Lancet *385*, 430-440. 10.1016/s0140-6736(14)61698-6.

Lopez Bernal, A. (2003). Mechanisms of labour--biochemical aspects. BJOG *110 Suppl 20*, 39-45. 10.1046/j.1471-0528.2003.00023.x.

Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biol *15*, 550. 10.1186/s13059-014-0550-8.

Ludmir, J., and Sehdev, H.M. (2000). Anatomy and physiology of the uterine cervix. Clin Obstet Gynecol *43*, 433-439. 10.1097/00003081-200009000-00003.

Makieva, S., Dubicke, A., Rinaldi, S.F., Fransson, E., Ekman-Ordeberg, G., and Norman, J.E. (2017). The preterm cervix reveals a transcriptomic signature in the presence of premature prelabor rupture of membranes. Am J Obstet Gynecol *216*, 602 e601-602 e621. 10.1016/j.ajog.2017.02.009.

Martinez-Varea, A., Romero, R., Xu, Y., Miller, D., Ahmed, A.I., Chaemsaithong, P., Chaiyasit, N., Yeo, L., Shaman, M., Lannaman, K., et al. (2017). Clinical chorioamnionitis at term VII: the amniotic fluid cellular immune response. J Perinat Med *45*, 523-538. 10.1515/jpm-2016-0225.

McGinnis, C.S., Murrow, L.M., and Gartner, Z.J. (2019). DoubletFinder: Doublet Detection in Single-Cell RNA Sequencing Data Using Artificial Nearest Neighbors. Cell Syst *8*, 329-337 e324. 10.1016/j.cels.2019.03.003.

McInnes, L., Healy, J., and Melville, J. (2018). UMAP: Uniform Manifold Approximation and Projection for Dimension Reduction. arXiv:1802.03426.

Migale, R., MacIntyre, D.A., Cacciatore, S., Lee, Y.S., Hagberg, H., Herbert, B.R., Johnson, M.R., Peebles, D., Waddington, S.N., and Bennett, P.R. (2016). Modeling hormonal and inflammatory contributions to preterm and term labor using uterine temporal transcriptomics. BMC Med *14*, 86. 10.1186/s12916-016-0632-4.

Montes, G.S., Zugaib, M., Joazeiro, P.P., Varayoud, J., Ramos, J.G., Munoz-de-Toro, M., and Luque, E.H. (2002). Phenotypic modulation of fibroblastic cells in the mucous layer of the human uterine cervix at term. Reproduction *124*, 783-790. 10.1530/rep.0.1240783.

Mosher, A.A., Rainey, K.J., Bolstad, S.S., Lye, S.J., Mitchell, B.F., Olson, D.M., Wood, S.L., and Slater, D.M. (2013). Development and validation of primary human myometrial cell culture models to study pregnancy and labour. BMC Pregnancy Childbirth *13 Suppl 1*, S7. 10.1186/1471-2393-13-s1-s7.

Motomura, K., Romero, R., Galaz, J., Tarca, A.L., Done, B., Xu, Y., Leng, Y., Garcia-Flores, V., Arenas-Hernandez, M., Theis, K.R., et al. (2021). RNA Sequencing Reveals Distinct Immune Responses in the Chorioamniotic Membranes of Women with Preterm Labor and Microbial or Sterile Intra-amniotic Inflammation. Infect Immun *89*. 10.1128/IAI.00819-20.

Motomura, K., Romero, R., Tarca, A.L., Galaz, J., Bhatti, G., Done, B., Arenas-Hernandez, M., Levenson, D., Slutsky, R., Hsu, C.D., and Gomez-Lopez, N. (2020a). Pregnancy-specific transcriptional changes upon endotoxin exposure in mice. J Perinat Med *48*, 700-722. 10.1515/jpm-2020-0159.

Motomura, K., Romero, R., Xu, Y., Theis, K.R., Galaz, J., Winters, A.D., Slutsky, R., Garcia-Flores, V., Zou, C., Levenson, D., et al. (2020b). Intra-Amniotic Infection with Ureaplasma parvum Causes Preterm Birth and Neonatal Mortality That Are Prevented by Treatment with Clarithromycin. mBio *11*. 10.1128/mBio.00797-20.

Nelson, A.C., Mould, A.W., Bikoff, E.K., and Robertson, E.J. (2016). Single-cell RNA-seq reveals cell type-specific transcriptional signatures at the maternal-foetal interface during pregnancy. Nat Commun *7*, 11414. 10.1038/ncomms11414.

Norwitz, E.R., Robinson, J.N., and Challis, J.R. (1999). The control of labor. N Engl J Med *341*, 660-666. 10.1056/NEJM199908263410906.

Odibo, A.O., Rodis, J.F., Sanders, M.M., Borgida, A.F., Wilson, M., Egan, J.F., and Campbell, W.A. (1999). Relationship of amniotic fluid markers of intra-amniotic infection with histopathology in cases of preterm labor with intact membranes. J Perinatol *19*, 407-412. 10.1038/sj.jp.7200210.

Oh, J.W., Moon, K.C., Park, C.W., Park, J.S., and Jun, J.K. (2021). Acute chorioamnionitis and intra-amniotic inflammation are more severe according to outside-in neutrophil migration within the same chorio-decidua. Taiwan J Obstet Gynecol *60*, 639-652. 10.1016/j.tjog.2021.05.011.

Oh, K.J., Park, K.H., Kim, S.N., Jeong, E.H., Lee, S.Y., and Yoon, H.Y. (2011). Predictive value of intra-amniotic and serum markers for inflammatory lesions of preterm placenta. Placenta *32*, 732-736. 10.1016/j.placenta.2011.07.080.

Oh, K.J., Romero, R., Park, J.Y., Lee, J., Conde-Agudelo, A., Hong, J.S., and Yoon, B.H. (2019). Evidence that antibiotic administration is effective in the treatment of a subset of patients

with intra-amniotic infection/inflammation presenting with cervical insufficiency. Am J Obstet
Gynecol *221*, 140.e141-140.e118. 10.1016/j.ajog.2019.03.017.
Olding, L. (1970). Value of placentitis as a sign of intrauterine infection in human subjects. Acta
Pathol Microbiol Scand A *78*, 256-264. 10.1111/j.1699-0463.1970.tb03300.x.
Osman, I., Young, A., Ledingham, M.A., Thomson, A.J., Jordan, F., Greer, I.A., and Norman,
J.E. (2003). Leukocyte density and pro-inflammatory cytokine expression in human fetal
membranes, decidua, cervix and myometrium before and during labour at term. Mol Hum
Reprod *9*, 41-45. 10.1093/molehr/gag001.
Pacora, P., Chaiworapongsa, T., Maymon, E., Kim, Y.M., Gomez, R., Yoon, B.H., Ghezzi, F.,
Berry, S.M., Qureshi, F., Jacques, S.M., et al. (2002). Funisitis and chorionic vasculitis: the
histological counterpart of the fetal inflammatory response syndrome. J Matern Fetal Neonatal
Med *11*, 18-25. 10.1080/jmf.11.1.18.25.
Park, C.W., Moon, K.C., Park, J.S., Jun, J.K., Romero, R., and Yoon, B.H. (2009). The
involvement of human amnion in histologic chorioamnionitis is an indicator that a fetal and an
intra-amniotic inflammatory response is more likely and severe: clinical implications. Placenta
30, 56-61. 10.1016/j.placenta.2008.09.017.
Patwardhan, M., Hernandez-Andrade, E., Ahn, H., Korzeniewski, S.J., Schwartz, A., Hassan,
S.S., and Romero, R. (2015). Dynamic Changes in the Myometrium during the Third Stage of
Labor, Evaluated Using Two-Dimensional Ultrasound, in Women with Normal and Abnormal
Third Stage of Labor and in Women with Obstetric Complications. Gynecol Obstet Invest *80*,
26-37. 10.1159/000370001.
Pavličev, M., Wagner, G.P., Chavan, A.R., Owens, K., Maziarz, J., Dunn-Fletcher, C., Kallapur,
S.G., Muglia, L., and Jones, H. (2017). Single-cell transcriptomics of the human placenta:
inferring the cell communication network of the maternal-fetal interface. Genome Res *27*, 349-
361. 10.1101/gr.207597.116.
Payne, K.J., Clyde, L.A., Weldon, A.J., Milford, T.A., and Yellon, S.M. (2012). Residency and
activation of myeloid cells during remodeling of the prepartum murine cervix. Biol Reprod *87*,
106. 10.1095/biolreprod.112.101840.
Pique-Regi, R., Romero, R., Garcia-Flores, V., Peyvandipour, A., Tarca, A.L., Pusod, E., Galaz,
J., Miller, D., Bhatti, G., Para, R., et al. (2022). A single-cell atlas of the myometrium in human
parturition. JCI Insight *7*. 10.1172/jci.insight.153921.
Pique-Regi, R., Romero, R., Tarca, A.L., Sendler, E.D., Xu, Y., Garcia-Flores, V., Leng, Y.,
Luca, F., Hassan, S.S., and Gomez-Lopez, N. (2019). Single cell transcriptional signatures of the
human placenta in term and preterm parturition. Elife *8*. 10.7554/eLife.52004.
Pollard, A.J., Sparey, C., Robson, S.C., Krainer, A.R., and Europe-Finner, G.N. (2000). Spatio-
temporal expression of the trans-acting splicing factors SF2/ASF and heterogeneous ribonuclear
proteins A1/A1B in the myometrium of the pregnant human uterus: a molecular mechanism for
regulating regional protein isoform expression in vivo. J Clin Endocrinol Metab *85*, 1928-1936.
10.1210/jcem.85.5.6537.
Redline, R.W. (2012). Inflammatory response in acute chorioamnionitis. Semin Fetal Neonatal
Med *17*, 20-25. 10.1016/j.siny.2011.08.003.
Redline, R.W., Faye-Petersen, O., Heller, D., Qureshi, F., Savell, V., and Vogler, C. (2003).
Amniotic infection syndrome: nosology and reproducibility of placental reaction patterns.
Pediatr Dev Pathol *6*, 435-448. 10.1007/s10024-003-7070-y.
Romero, R., Dey, S.K., and Fisher, S.J. (2014a). Preterm labor: one syndrome, many causes.
Science *345*, 760-765. 10.1126/science.1251816.

Romero, R., Espinoza, J., Kusanovic, J.P., Gotsch, F., Hassan, S., Erez, O., Chaiworapongsa, T., and Mazor, M. (2006). The preterm parturition syndrome. BJOG *113 Suppl 3*, 17-42. 10.1111/j.1471-0528.2006.01120.x.

Romero, R., Gomez-Lopez, N., Winters, A.D., Jung, E., Shaman, M., Bieda, J., Panaitescu, B., Pacora, P., Erez, O., Greenberg, J.M., et al. (2019). Evidence that intra-amniotic infections are often the result of an ascending invasion - a molecular microbiological study. J Perinat Med *47*, 915-931. 10.1515/jpm-2019-0297.

Romero, R., Mazor, M., Munoz, H., Gomez, R., Galasso, M., and Sherer, D.M. (1994). The preterm labor syndrome. Ann N Y Acad Sci *734*, 414-429. 10.1111/j.1749-6632.1994.tb21771.x.

Romero, R., Miranda, J., Chaemsaithong, P., Chaiworapongsa, T., Kusanovic, J.P., Dong, Z., Ahmed, A.I., Shaman, M., Lannaman, K., Yoon, B.H., et al. (2015a). Sterile and microbial-associated intra-amniotic inflammation in preterm prelabor rupture of membranes. J Matern Fetal Neonatal Med *28*, 1394-1409. 10.3109/14767058.2014.958463.

Romero, R., Miranda, J., Chaiworapongsa, T., Chaemsaithong, P., Gotsch, F., Dong, Z., Ahmed, A.I., Yoon, B.H., Hassan, S.S., Kim, C.J., et al. (2014b). A novel molecular microbiologic technique for the rapid diagnosis of microbial invasion of the amniotic cavity and intra-amniotic infection in preterm labor with intact membranes. Am J Reprod Immunol *71*, 330-358. 10.1111/aji.12189.

Romero, R., Miranda, J., Chaiworapongsa, T., Chaemsaithong, P., Gotsch, F., Dong, Z., Ahmed, A.I., Yoon, B.H., Hassan, S.S., Kim, C.J., et al. (2015b). Sterile intra-amniotic inflammation in asymptomatic patients with a sonographic short cervix: prevalence and clinical significance. J Matern Fetal Neonatal Med *28*, 1343-1359. 10.3109/14767058.2014.954243.

Romero, R., Miranda, J., Kusanovic, J.P., Chaiworapongsa, T., Chaemsaithong, P., Martinez, A., Gotsch, F., Dong, Z., Ahmed, A.I., Shaman, M., et al. (2015c). Clinical chorioamnionitis at term I: microbiology of the amniotic cavity using cultivation and molecular techniques. J Perinat Med *43*, 19-36. 10.1515/jpm-2014-0249.

Romero, R., Quintero, R., Nores, J., Avila, C., Mazor, M., Hanaoka, S., Hagay, Z., Merchant, L., and Hobbins, J.C. (1991). Amniotic fluid white blood cell count: a rapid and simple test to diagnose microbial invasion of the amniotic cavity and predict preterm delivery. Am J Obstet Gynecol *165*, 821-830. 10.1016/0002-9378(91)90423-o.

Romero, R., Sirtori, M., Oyarzun, E., Avila, C., Mazor, M., Callahan, R., Sabo, V., Athanassiadis, A.P., and Hobbins, J.C. (1989). Infection and labor. V. Prevalence, microbiology, and clinical significance of intraamniotic infection in women with preterm labor and intact membranes. Am J Obstet Gynecol *161*, 817-824. 10.1016/0002-9378(89)90409-2.

Romero, R., Yoon, B.H., Mazor, M., Gomez, R., Diamond, M.P., Kenney, J.S., Ramirez, M., Fidel, P.L., Sorokin, Y., Cotton, D., and et al. (1993a). The diagnostic and prognostic value of amniotic fluid white blood cell count, glucose, interleukin-6, and gram stain in patients with preterm labor and intact membranes. Am J Obstet Gynecol *169*, 805-816. 10.1016/0002-9378(93)90009-8.

Romero, R., Yoon, B.H., Mazor, M., Gomez, R., Gonzalez, R., Diamond, M.P., Baumann, P., Araneda, H., Kenney, J.S., Cotton, D.B., and et al. (1993b). A comparative study of the diagnostic performance of amniotic fluid glucose, white blood cell count, interleukin-6, and gram stain in the detection of microbial invasion in patients with preterm premature rupture of membranes. Am J Obstet Gynecol *169*, 839-851. 10.1016/0002-9378(93)90014-a.

Sakamoto, Y., Moran, P., Bulmer, J.N., Searle, R.F., and Robson, S.C. (2005). Macrophages and not granulocytes are involved in cervical ripening. J Reprod Immunol *66*, 161-173. 10.1016/j.jri.2005.04.005.

Schwalm, H., and Dubrauszky, V. (1966). The structure of the musculature of the human uterus--muscles and connective tissue. Am J Obstet Gynecol *94*, 391-404. 10.1016/0002-9378(66)90661-2.

Senthamaraikannan, P., Presicce, P., Rueda, C.M., Maneenil, G., Schmidt, A.F., Miller, L.A., Waites, K.B., Jobe, A.H., Kallapur, S.G., and Chougnet, C.A. (2016). Intra-amniotic Ureaplasma parvum-Induced Maternal and Fetal Inflammation and Immune Responses in Rhesus Macaques. J Infect Dis *214*, 1597-1604. 10.1093/infdis/jiw408.

Sindram-Trujillo, A., Scherjon, S., Kanhai, H., Roelen, D., and Claas, F. (2003). Increased T-cell activation in decidua parietalis compared to decidua basalis in uncomplicated human term pregnancy. Am J Reprod Immunol *49*, 261-268. 10.1034/j.1600-0897.2003.00041.x.

Slutsky, R., Romero, R., Xu, Y., Galaz, J., Miller, D., Done, B., Tarca, A.L., Gregor, S., Hassan, S.S., Leng, Y., and Gomez-Lopez, N. (2019). Exhausted and Senescent T Cells at the Maternal-Fetal Interface in Preterm and Term Labor. J Immunol Res *2019*, 3128010. 10.1155/2019/3128010.

Smith, R. (2007). Parturition. N Engl J Med *356*, 271-283. 10.1056/NEJMra061360.

Sooranna, S.R., Grigsby, P.L., Engineer, N., Liang, Z., Sun, K., Myatt, L., and Johnson, M.R. (2006). Myometrial prostaglandin E2 synthetic enzyme mRNA expression: spatial and temporal variations with pregnancy and labour. Mol Hum Reprod *12*, 625-631. 10.1093/molehr/gal061.

Sparey, C., Robson, S.C., Bailey, J., Lyall, F., and Europe-Finner, G.N. (1999). The differential expression of myometrial connexin-43, cyclooxygenase-1 and -2, and Gs alpha proteins in the upper and lower segments of the human uterus during pregnancy and labor. J Clin Endocrinol Metab *84*, 1705-1710. 10.1210/jcem.84.5.5644.

Stuart, T., Butler, A., Hoffman, P., Hafemeister, C., Papalexi, E., Mauck, W.M., 3rd, Hao, Y., Stoeckius, M., Smibert, P., and Satija, R. (2019). Comprehensive Integration of Single-Cell Data. Cell *177*, 1888-1902 e1821. 10.1016/j.cell.2019.05.031.

Suryawanshi, H., Morozov, P., Straus, A., Sahasrabudhe, N., Max, K.E.A., Garzia, A., Kustagi, M., Tuschl, T., and Williams, Z. (2018). A single-cell survey of the human first-trimester placenta and decidua. Sci Adv *4*, eaau4788. 10.1126/sciadv.aau4788.

Svensson, L., Ingemarsson, I., and Mårdh, P.A. (1986). Chorioamnionitis and the isolation of microorganisms from the placenta. Obstet Gynecol *67*, 403-409.

Sweeney, E.L., Kallapur, S.G., Gisslen, T., Lambers, D.S., Chougnet, C.A., Stephenson, S.A., Jobe, A.H., and Knox, C.L. (2016). Placental Infection With Ureaplasma species Is Associated With Histologic Chorioamnionitis and Adverse Outcomes in Moderately Preterm and Late-Preterm Infants. J Infect Dis *213*, 1340-1347. 10.1093/infdis/jiv587.

Tabula Muris, C., Overall, c., Logistical, c., Organ, c., processing, Library, p., sequencing, Computational data, a., Cell type, a., Writing, g., et al. (2018). Single-cell transcriptomics of 20 mouse organs creates a Tabula Muris. Nature *562*, 367-372. 10.1038/s41586-018-0590-4.

Tarca, A.L., Romero, R., Erez, O., Gudicha, D.W., Than, N.G., Benshalom-Tirosh, N., Pacora, P., Hsu, C.D., Chaiworapongsa, T., Hassan, S.S., and Gomez-Lopez, N. (2021). Maternal whole blood mRNA signatures identify women at risk of early preeclampsia: a longitudinal study. J Matern Fetal Neonatal Med *34*, 3463-3474. 10.1080/14767058.2019.1685964.

Tarca, A.L., Romero, R., Xu, Z., Gomez-Lopez, N., Erez, O., Hsu, C.D., Hassan, S.S., and Carey, V.J. (2019). Targeted expression profiling by RNA-Seq improves detection of cellular

dynamics during pregnancy and identifies a role for T cells in term parturition. Sci Rep *9*, 848. 10.1038/s41598-018-36649-w.

Theis, K.R., Romero, R., Motomura, K., Galaz, J., Winters, A.D., Pacora, P., Miller, D., Slutsky, R., Florova, V., Levenson, D., et al. (2020). Microbial burden and inflammasome activation in amniotic fluid of patients with preterm prelabor rupture of membranes. J Perinat Med *48*, 115-131. 10.1515/jpm-2019-0398.

Timmons, B.C., Fairhurst, A.M., and Mahendroo, M.S. (2009). Temporal changes in myeloid cells in the cervix during pregnancy and parturition. J Immunol *182*, 2700-2707. 10.4049/jimmunol.0803138.

Toothaker, J.M., Presicce, P., Cappelletti, M., Stras, S.F., McCourt, C.C., Chougnet, C.A., Kallapur, S.G., and Konnikova, L. (2020). Immune Cells in the Placental Villi Contribute to Intra-amniotic Inflammation. Front Immunol *11*, 866. 10.3389/fimmu.2020.00866.

Trundley, A., and Moffett, A. (2004). Human uterine leukocytes and pregnancy. Tissue Antigens *63*, 1-12. 10.1111/j.1399-0039.2004.00170.x.

Tsang, J.C.H., Vong, J.S.L., Ji, L., Poon, L.C.Y., Jiang, P., Lui, K.O., Ni, Y.B., To, K.F., Cheng, Y.K.Y., Chiu, R.W.K., and Lo, Y.M.D. (2017). Integrative single-cell and cell-free plasma RNA transcriptomics elucidates placental cellular dynamics. Proc Natl Acad Sci U S A *114*, E7786-e7795. 10.1073/pnas.1710470114.

van Hoeven, K.H., Anyaegbunam, A., Hochster, H., Whitty, J.E., Distant, J., Crawford, C., and Factor, S.M. (1996). Clinical significance of increasing histologic severity of acute inflammation in the fetal membranes and umbilical cord. Pediatr Pathol Lab Med *16*, 731-744.

Vargas, M.L., Sántos, J.L., Ruiz, C., Montes, M.J., Alemán, P., García-Tortosa, C., and García-Olivares, E. (1993). Comparison of the proportions of leukocytes in early and term human decidua. Am J Reprod Immunol *29*, 135-140. 10.1111/j.1600-0897.1993.tb00578.x.

Vento-Tormo, R., Efremova, M., Botting, R.A., Turco, M.Y., Vento-Tormo, M., Meyer, K.B., Park, J.E., Stephenson, E., Polanski, K., Goncalves, A., et al. (2018). Single-cell reconstruction of the early maternal-fetal interface in humans. Nature *563*, 347-353. 10.1038/s41586-018-0698-6.

Vink, J.Y., Qin, S., Brock, C.O., Zork, N.M., Feltovich, H.M., Chen, X., Urie, P., Myers, K.M., Hall, T.J., Wapner, R., et al. (2016). A new paradigm for the role of smooth muscle cells in the human cervix. Am J Obstet Gynecol *215*, 478 e471-478 e411. 10.1016/j.ajog.2016.04.053.

Viscardi, R.M. (2010). Ureaplasma species: role in diseases of prematurity. Clin Perinatol *37*, 393-409. 10.1016/j.clp.2009.12.003.

Wendremaire, M., Hadi, T., Pezze, M., Barrichon, M., Lopez, T., Neiers, F., Sagot, P., Garrido, C., and Lirussi, F. (2020). Macrophage-induced reactive oxygen species promote myometrial contraction and labor-associated mechanisms. Biol Reprod *102*, 1326-1339. 10.1093/biolre/ioaa032.

Wikland, M., Lindblom, B., Wilhelmsson, L., and Wiqvist, N. (1982). Oxytocin, prostaglandins, and contractility of the human uterus at term pregnancy. Acta Obstet Gynecol Scand *61*, 467-472. 10.3109/00016348209156592.

Willcockson, A.R., Nandu, T., Liu, C.L., Nallasamy, S., Kraus, W.L., and Mahendroo, M. (2018). Transcriptome signature identifies distinct cervical pathways induced in lipopolysaccharide-mediated preterm birth. Biol Reprod *98*, 408-421. 10.1093/biolre/iox180.

Yoneda, N., Yoneda, S., Niimi, H., Ueno, T., Hayashi, S., Ito, M., Shiozaki, A., Urushiyama, D., Hata, K., Suda, W., et al. (2016a). Polymicrobial Amniotic Fluid Infection with Mycoplasma/Ureaplasma and Other Bacteria Induces Severe Intra-Amniotic Inflammation

Associated with Poor Perinatal Prognosis in Preterm Labor. Am J Reprod Immunol *75*, 112-125. 10.1111/aji.12456.

Yoneda, S., Shiozaki, A., Yoneda, N., Ito, M., Shima, T., Fukuda, K., Ueno, T., Niimi, H., Kitajima, I., Kigawa, M., and Saito, S. (2016b). Antibiotic Therapy Increases the Risk of Preterm Birth in Preterm Labor without Intra-Amniotic Microbes, but may Prolong the Gestation Period in Preterm Labor with Microbes, Evaluated by Rapid and High-Sensitive PCR System. Am J Reprod Immunol *75*, 440-450. 10.1111/aji.12484.

Yoon, B.H., Chang, J.W., and Romero, R. (1998). Isolation of Ureaplasma urealyticum from the amniotic cavity and adverse outcome in preterm labor. Obstet Gynecol *92*, 77-82. 10.1016/s0029-7844(98)00122-7.

Yoon, B.H., Romero, R., Moon, J.B., Shim, S.S., Kim, M., Kim, G., and Jun, J.K. (2001). Clinical significance of intra-amniotic inflammation in patients with preterm labor and intact membranes. Am J Obstet Gynecol *185*, 1130-1136. 10.1067/mob.2001.117680.

Yoon, B.H., Romero, R., Park, J.Y., Oh, K.J., Lee, J., Conde-Agudelo, A., and Hong, J.S. (2019). Antibiotic administration can eradicate intra-amniotic infection or intra-amniotic inflammation in a subset of patients with preterm labor and intact membranes. Am J Obstet Gynecol *221*, 142.e141-142.e122. 10.1016/j.ajog.2019.03.018.

Yoon, B.H., Yang, S.H., Jun, J.K., Park, K.H., Kim, C.J., and Romero, R. (1996). Maternal blood C-reactive protein, white blood cell count, and temperature in preterm labor: a comparison with amniotic fluid white blood cell count. Obstet Gynecol *87*, 231-237. 10.1016/0029-7844(95)00380-0.

Young, M.D., and Behjati, S. (2020). SoupX removes ambient RNA contamination from droplet-based single-cell RNA sequencing data. Gigascience *9*. 10.1093/gigascience/giaa151.

Yu, G., Wang, L.G., Han, Y., and He, Q.Y. (2012). clusterProfiler: an R package for comparing biological themes among gene clusters. OMICS *16*, 284-287. 10.1089/omi.2011.0118.

Zhang, J.M., Kraus, F.T., and Aquino, T.I. (1985). Chorioamnionitis: a comparative histologic, bacteriologic, and clinical study. Int J Gynecol Pathol *4*, 1-10.

Zhao, Z., Wang, Y., Wu, Y., Li, D., Zhang, T., Ma, Y., Teng, X., and Zuo, W. (2021). Single-cell analysis defines the lineage plasticity of stem cells in cervix epithelium. Cell Regen *10*, 36. 10.1186/s13619-021-00096-2.

A

Intra-amniotic injection

Allogeneic pregnancy

Ultrasound system

16.5 *dpc*

Needle

Amniotic cavity

Fetus

B Gestational age

p = 0.03

Days *post coitum*

19.5

18.5

18.0

17.5

PBS   *E.coli*

C Rate of preterm birth

p = 0.04

Percentage %

PBS   *E.coli*

D

2nd Cohort

Measurement of the cervical length

Measurement of the cervical length

16.5 *dpc*

Bladder

Vagina

Cervix

17.5 *dpc*

Intra-amniotic injection

E Shortening of cervical length

p < 0.001

Percentage %

PBS   *E.coli*

F

Uterus   **Decidua**   **Cervix**   **Isolation of single-cells**   **Single-cell technology**

G

UMAP_2

-5

-10

-15   -10   -5   0   5   10

UMAP_1

H

UMAP_2

-5

-10

Myeloid

Stromal

Endothelial

Lymphoid

Smooth muscle cells

Epithelial

-15   -10   -5   0   5   10

UMAP_1

Uterus
Decidua
Cervix

Cell types

0	Fibroblast-1	6	Neutrophil	12	Stromal-3	18	Dendritic cell	24	Macrophage-2 (Progenitor)
1	Fibroblast-2	7	Epithelial-2 (Squamous)	13	Epithelial-5 (Luminal)	19	NK cell-1	25	Plasmocyte
2	Stromal-1	8	Epithelial-3 (Squamous)	14	Epithelial-6 (Secretory)	20	Epithelial-7 (Glandular)	26	SMC-2
3	Monocyte	9	Macrophage-1	15	SMC-1	21	Fibroblast-3	27	NK cell-2
4	Stromal-2 (Decidual)	10	Epithelial-4 (Glandular)	16	T cell	22	Erythroid	28	Epithelial-9 (Secretory)
5	Epithelial-1 (Basal)	11	Epithelial-10 (Glandular)	17	Endothelial	23	Epithelial-8 (Enterocyte)	29	Trophoblast
								30	B cell

Decidua

Cervix

A Sender cell / Signaling pathway

B Signaling pathway / Receiver cell

C Outgoing interaction strength (Control → PTL) vs Incoming interaction strength (Control → PTL)

D Differential interactions

Sender cell → Receiver cell, Self signaling

—— Increased signaling —— Decreased signaling

1 Fibroblast-2 2 Stromal-1 5 Epithelial-1 (Basal) 7 Epithelial-2 (Squamous) 8 Epithelial-3 (Squamous) 10 Epithelial-4 (Glandular) 11 Epithelial-10 (Glandular) 14 Epithelial-6 (Secretory) 15 SMC-1 20 Epithelial-7 (Glandular) 22 Erythroid 23 Epithelial-8 (Enterocyte)

E Collagen — Control / Preterm labor ; Tenascin — Control / Preterm labor

A Murine uterus

Differential interaction strength

Sources (Sender): Fibroblast-1, Fibroblast-2, Stromal-3, Epithelial-5, Epithelial-6, SMC-1, T cell, Endothelial, Dendritic cell, NK cell-1, Stromal-1, Fibroblast-3, Monocyte, Stromal-2, Neutrophil, Macrophage-1, Macrophage-2, Plasmocyte

Receivers: Fibroblast-1, Fibroblast-2, Stromal-3, Epithelial-5, Epithelial-6, SMC-1, T cell, Endothelial, Dendritic cell, NK cell-1, Stromal-1, Fibroblast-3, Monocyte, Stromal-2, Neutrophil, Macrophage-1, Macrophage-2, Plasmocyte

Relative values −4 −2 0 2 4

B Human myometrium

Differential interaction strength

Sources (Sender): B cell, CD4 T cell, CD8 T cell, Dendritic cell, Decidua, Endothelial-1, Endothelial-2, ILC, LED, Macrophage-1, Macrophage-2, Macrophage-3, Monocyte, Myofibroblast, NK cell, SMC-1, SMC-2, SMC-3, Stromal-1, Stromal-2, EVT, Macrophage-4, Plasmablast, Uncil. Epithelial

Receivers: B cell, CD4 T cell, CD8 T cell, Dendritic cell, Decidua, Endothelial-1, Endothelial-2, ILC, LED, Macrophage-1, Macrophage-2, Macrophage-3, Monocyte, Myofibroblast, NK cell, SMC-1, SMC-2, SMC-3, Stromal-1, Stromal-2, EVT, Macrophage-4, Plasmablast, Uncil. Epithelial

Relative values −4 −2 0 2 4

C COLLAGEN | IL-1 | CCL

Control

Preterm labor

Murine uterus

0 Fibroblast-1		15 SMC-1	
1 Fibroblast-2		16 T cell	
2 Stromal-1		17 Endothelial	
3 Monocyte		18 Dendritic cell	
4 Stromal-2		19 NK cell-1	
6 Neutrophil		21 Fibroblast-3	
9 Macrophage-1		22 Erythroid	
13 Epithelial-5		24 Macrophage-2	
14 Epithelial-6		25 Plasmocyte	

D

Term not in labor

Term in labor

Human myometrium

a Endothelial-1	m	B cell
b Endothelial-2	n	Uncilliated epithelial
c Decidual	o	EVT
d Dendritic cell	p	Stromal-1
e CD4 T cell	q	Stromal-2
f CD8 T cell	r	SMC-1
g SMC-2	s	Macrophage-1
h SMC-3	t	Macrophage-2
i NK cell	u	Macrophage-3
j Myofibroblast	v	Macrophage-4
k Monocyte	w	ILC
l Plasmablast	x	LED