Research Article

EGFR inhibition leads to enhanced desmosome assembly and cohesion in

cardiomyocytes

Condensed title: EGFR inhibition-induced positive adhesiotropy

Maria Shoykhet[1], Orsela Dervishi[1], Philipp Menauer[1], Matthias Hiermaier[1], Jens Waschke[1]

and Sunil Yeruva[1*]

[1]Faculty of Medicine, Ludwig-Maximilian-University (LMU) Munich, Munich, Germany

Corresponding author:

*Sunil Yeruva, Anatomical Institute, Department I – Vegetative Anatomy, Ludwig-

Maximilian-University, Pettenkoferstraße 11, 80336 Munich, Germany; Phone: +49-89-2180-

72703; sunil.yeruva@med.uni-muenchen.de

Summary: Shoykhet *et al*. show EGFR inhibition-enhanced cardiomyocyte cohesion. This

positive adhesiotropy is achieved via an increase in desmosomal assembly and DP/DSG2

localization at cell borders and is the first step towards a novel therapeutic approach for

arrhythmogenic cardiomyopathy.

Key Words:

Epidermal growth factor receptor, erlotinib, desmoglein 2, desmoplakin,

arrhythmogenic cardiomyopathy

Conflict of Interest

The authors declare that no conflict of interest exists.

Abstract

The epidermal growth factor receptor (EGFR) is a ubiquitously expressed receptor tyrosine

kinase, which is dysregulated in several kinds of cancer and heart diseases. Arrhythmogenic

cardiomyopathy (AC) is a familial heart disease in part caused by impaired desmosome

turnover. Thus, stabilization of desmosome integrity, may provide potential new treatment

options. Desmosomes, apart from cellular cohesion, provide the structural framework of a

signaling hub. Here, we investigated the role of EGFR inhibition on cardiomyocyte cohesion

under physiological and pathophysiological conditions using the murine plakoglobin

knockout AC model, in which EGFR was upregulated. EGFR inhibition led to enhanced

cardiomyocyte cohesion. Immunoprecipitation showed an interaction of EGFR and

desmoglein 2 (DSG2). Immunostaining and AFM revealed enhanced DSG2 localization and

binding at cell borders upon EGFR inhibition. Enhanced area composita length and

desmosome assembly were observed upon EGFR inhibition, confirmed by enhanced DSG2

and desmoplakin (DP) recruitment to cell borders. Thus, inhibiting EGFR, thereby stabilizing

desmosome integrity, might provide new treatment options for AC.

Abbreviation	full form
AC	arrhythmogenic cardiomyopathy
AFM	atomic force microscopy
BSA/NGS	bovine serum albumin and normal goat serum
DES	desmin
DP	desmoplakin
DSC2	desmocollin 2
DSG2	desmoglein 2
EGFR	epidermal growth factor receptor
Erlo	erlotinib
FRAP	fluorescence recovery after photobleach
IgG hc	IgG heavy chain
Jup	junctional plakoglobin
N-CAD	N-Cadherin
NT	non-target
PFA	paraformaldehyde
PG	plakoglobin
PKP2	plakophilin 2
SRC	proto-oncogene kinase Src
STED	stimulated emission depletion
WGA	wheat germ agglutinin
τ	tau; halftime of recovery

Introduction

The epidermal growth factor receptor (EGFR) is a ubiquitously expressed transmembrane receptor tyrosine kinase. Dysregulation of EGFR signaling often results in pathologies, such as different kinds of cancer, but also inflammatory skin and bowel disease or other skin, inflammatory or renal disorders (Bhargava et al., 2005, Kleinschmidt-DeMasters et al., 2005, Shia et al., 2005, da Cunha Santos et al., 2011, Campbell et al., 2014, Harskamp et al., 2016, Iwamoto et al., 2019). Today, EGFR inhibitors, either small inhibitory molecules (e.g., erlotinib, gefitinib or lapatinib) or inhibitory antibodies (e.g., cetuximab or panitumumab) are approved as medication to treat non-small cell lung cancer, pancreatic cancer, breast cancer, colorectal carcinoma or squamous cell carcinoma of head and neck (Westover et al., 2018, Sabbah et al., 2020). EGFR can be transactivated through G protein-coupled receptors via second messengers, several protein kinases or matrix metalloproteases (Noma et al., 2007, Makki et al., 2013). In addition, EGFR signaling can be activated, enhanced and mediated by the proto-oncogene tyrosine-protein kinase Src (SRC) (Chen et al., 2018). Vice versa, SRC can be activated by EGFR (Wilde et al., 1999). In the heart, EGFR was found to play a crucial role in cell metabolism, proliferation, cell survival, as well as hypertrophy and remodeling (Stroedecke et al., 2021). EGFR inhibition can lead to heart failure in the healthy heart, and a total knockout of the receptor can lead to severe hypertrophy and hypotension (Schreier et al., 2013, Kenigsberg et al., 2017, Guo et al., 2021). However, under pathological conditions, EGFR inhibition can be cardioprotective, as EGFR has been shown to be upregulated in a diabetic setting (Matrougui, 2010), while EGFR inhibition protected vasculature and heart from diabetes-induced damages (Belmadani et al., 2008, Benter et al., 2009, Akhtar et al., 2012, Choi et al., 2012, El-Hashim et al., 2017). Furthermore, recent studies indicate that EGFR inhibition has a protective effect on cardiomyocytes in hypoxic conditions, i.e., after myocardial infarctions (Heliste et al., 2021). It seems that like in other tissues, EGFR signaling is tightly controlled in the heart, and due to its double role and the multitude of its

downstream effectors, any alterations in EGFR signaling can have severe effects on heart

function (Shraim et al., 2021, Forrester et al., 2016).

Arrhythmogenic cardiomyopathy (AC) is a disease most commonly caused by mutations in

genes coding for desmosomal proteins, such as desmoglein 2 (DSG2), plakoglobin (PG),

desmoplakin (DP), desmocollin 2 (DSC2), N-Cadherin (N-CAD) and plakophilin 2 (PKP2)

(Gerull and Brodehl, 2020, Corrado et al., 2017), hence considered as a disease of the

desmosome. Causative mutations of AC often lead to alterations in desmosomal contacts

between cells, thereby weakening cellular cohesion. Several pathways regulating

cardiomyocyte cohesion have been identified so far (Genet et al., 2012, Schlipp et al., 2014,

Schinner et al., 2017, Zhao et al., 2019, Shoykhet et al., 2020). Furthermore, there is

increasing evidence that desmosomes not only provide cellular cohesion, but also act as

signaling hubs (Waschke, 2019, Muller et al., 2021), meaning that desmosomes can influence,

but are also influenced by cellular signaling processes. Recent advancement in research

suggests that similar molecular mechanisms are present in AC and pemphigus, an

autoimmune skin disease, as well as inflammatory bowel diseases (Waschke, 2019, Shoykhet

et al., 2020, Schmitt and Waschke, 2021, Yeruva and Waschke, 2022).

EGFR inhibition led to increased cellular cohesion in keratinocytes (Walter et al., 2019),

whereas in squamous carcinoma cell lines, EGFR inhibition increased protein levels of DSG2

and DSC2 (Lorch et al., 2004). However, EGFR inhibition by erlotinib in intestinal epithelial

cells caused a decrease in cellular cohesion (Ungewiss et al., 2018), revealing that the role of

EGFR can vary between different cells or tissues.

EGFR seems to play a major role in the heart function. Furthermore, links between EGFR and

DSG2 and other desmosomal proteins have been established. Thus, in this study, we

investigated whether inhibition of the EGFR signaling pathway can be used as a therapeutic

option for AC, via regulating cardiomyocyte cohesion, using a $Jup^{-/-}$ mouse model, which we

characterized previously (Schinner et al., 2017, Schinner et al., 2020, Shoykhet et al., 2020).

To this end, we used the well-known and clinically approved first-generation EGFR inhibitor

erlotinib and an EGFR signaling molecule SRC inhibitor, PP2. Erlotinib was used to directly

inhibit EGFR and PP2 to indirectly inhibit EGFR, by inhibiting the EGFR signaling pathway,

since SRC acts both up- and downstream of EGFR. This approach was used as a proof of

concept that EGFR inhibition, either using drug or a signaling inhibitor, rescues impaired

cardiomyocyte cohesion.

Results

Inhibition of EGFR or SRC lead to positive adhesiotropy in HL-1 cardiomyocytes and murine

cardiac slice cultures

In keratinocytes and intestinal epithelial cells, as well as in cancer cells, it is known that

EGFR can modulate the desmosomal structure, assembly and signaling and vice versa (Lorch

et al., 2004, Klessner et al., 2009, Bektas et al., 2013, Kamekura et al., 2014, Ungewiss et al.,

2018). We used the cardiomyocyte-specific plakoglobin knockout ($Jup^{-/-}$) mouse, representing

a pathogenic AC model, which shows fibrotic replacement of myocardial tissue as well as

arrhythmias (Schinner et al., 2020, Schinner et al., 2017, Shoykhet et al., 2020) and found an

upregulation of EGFR protein in the $Jup^{-/-}$ mice as compared to wild type ($Jup^{+/+}$) mice

(Figure 1A and Supplementary Figure 1A). Thus, we investigated the role of EGFR in

cardiomyocyte cohesion. To this end, we inhibited EGFR directly using erlotinib or SRC,

thereby inhibiting the EGFR pathway, using PP2 and performed dispase-based dissociation

assays with HL-1 cardiomyocytes and murine cardiac slice cultures. Therefore, treated HL-1

cell monolayers or murine cardiac slices were incubated with a mixture of dispase II and

collagenase and mechanical stress was applied to challenge cell-cell contacts, upon which

fragments of the monolayer (for HL-1 cells) or single cardiomyocytes (for murine cardiac

slices) resulted. The results from this assay shall be interpreted as the lesser the number of

fragments or single cardiomyocytes, the higher the cardiomyocyte cohesion or vice versa.

We observed that both EGFR and SRC inhibition enhanced cardiomyocyte cohesion, which

we term as positive adhesiotropy, in HL-1 cardiomyocytes and murine cardiac slice cultures

(Figure 1B and 1C). Inhibition of EGFR or SRC by erlotinib or PP2, respectively, was

confirmed by assessing the phosphorylation levels of their downstream targets: ERK and

EGFR (Supplementary Figure 1B-D). We also performed immunostaining of N-CAD and

DSG2 in HL-1 cardiomyocytes and found an increase of DSG2 at the cell borders after EGFR

or SRC inhibition, compared to controls (Figures 1D and 1E). Next, to assess whether EGFR

is in a complex with desmosomal proteins, we immunoprecipitated DSG2 from HL-1 lysates

and performed immunoblots for EGFR, PG, DP and PKP2. Indeed, EGFR was found in

complex with DSG2 in addition to PG, DP and PKP2 (Figure 1F). Nevertheless, treatment

with erlotinib did not alter the composition of this complex.

These experiments showed that inhibition of either EGFR or SRC leads to positive

adhesiotropy in HL-1 cardiomyocytes and murine cardiac slice cultures, paralleled by

enhanced DSG2 localization along cell junctions.

Inhibition of EGFR increases the DSG2 binding event frequency at cell borders

Since we observed an increase in DSG2 localization at the membrane after EGFR or SRC

inhibition, we assessed the role of DSG2 in strengthening cellular cohesion upon EGFR

inhibition in cardiomyocytes. For this, we performed atomic force microscopy (AFM)

measurements using HL-1 cardiomyocytes. Here, a cantilever with a protein-coated tip

approached the cells, where the protein at the tip could eventually interact with proteins at the

cell surface. We measured the binding frequency of DSG2, DSC2, or N-CAD-coated tips, as

well as the unbinding forces at cell borders and cell surfaces. The areas were chosen using a

topography image of the cells, acquired in the force-based quantitative imaging mode. We

observed an increase in binding frequency at cell borders for DSG2-coated tips only (Figure

2A). In the case of N-CAD coated tips no change in binding frequency was observed (Figure

2B). However, in the case of DSC2-coated tips, reduced binding frequency at the cell surface,

but not the cell border, was found (Figure 2C). On the other hand, no changes in unbinding

forces for any of the proteins analyzed were observed (Figure 2D-F). These data indicate that

enhanced binding frequency of DSG2 correlates with the strengthening of intercellular

adhesion in HL-1 cardiomyocytes upon EGFR inhibition.

Apart from DSG2, positive adhesiotropy induced by EGFR or SRC inhibition is also

dependent on DP

To further understand the mechanism of erlotinib or PP2-induced positive adhesiotropy, we

performed siRNA-mediated knockdown in HL-1 cardiomyocytes. Firstly, we knocked down

Egfr, which did not alter cellular cohesion (Figure 3A and Supplementary Figure 2A). We

then knocked down *Egfr* and treated the cells with erlotinib or PP2. Upon *Egfr* knockdown,

erlotinib-induced positive adhesiotropy was abolished, confirming that the effect observed

after erlotinib treatment is mediated through EGFR. Interestingly, SRC inhibition was still

effective after *Egfr* knockdown, indicating that the PP2 effect is either independent of EGFR

or that SRC is downstream of EGFR (Figure 3B and Supplementary Figure 2A). Next, we

knocked down *Dsp* or *Dsg2*. Western blots confirmed all knockdowns (Figure 3C). Knocking

down *Dsp* or *Dsg2* did not alter cellular cohesion (Figure 3D and Supplementary Figure 2B

and 2C), as observed before (Shoykhet et al., 2020, Schinner et al., 2020). Both, erlotinib and

PP2 enhanced cellular cohesion after *Dsg2* knockdown. This suggests that the remaining

DSG2 protein might be sufficient enough to mediate the positive adhesiotropic effect of

EGFR or SRC inhibition. In previous reports inhibition of EGFR enhanced DP expression in

breast cancer cells and led to enhanced DP at the cell border in oral squamous cell carcinoma

cells (Lorch et al., 2004, Pang et al., 2004), therefore, we knocked down *Dsp*. EGFR or SRC

inhibition-mediated positive adhesiotropy was abrogated upon *Dsp* knockdown, indicating the

crucial role of DP in erlotinib or PP2-mediated positive adhesiotropy (Figure 3E and

Supplementary Figure 2C). These data highlight the role of DP, along with DSG2, in positive

adhesiotropy induced by EGFR or SRC inhibition.

EGFR or SRC inhibition leads to increased area composita length in HL-1 cardiomyocytes

Since both, DSG2 and DP seem to be involved in the positive adhesiotropy induced by EGFR

or SRC inhibition, we next performed immunostaining of these proteins together with

Phalloidin to stain the cortical actin cytoskeleton. Confocal microscopy images revealed an

increase in the localization of DP and DSG2 at cell borders after inhibition of EGFR or SRC

(Figure 4A-C). At the same time DSG2 and DP staining looked enlarged in erlotinib and PP2-

treated cells compared to vehicle controls, which indicated changes at the area composita, that

were shown to be present in intercellular junctions of HL-1 cells (Schlipp et al., 2014).

Therefore, we performed stimulated emission depletion (STED) imaging for DSG2 and DP,

which showed that area composita number remained unchanged, whereas the area composita

length increased after EGFR or SRC inhibition (Figure 4D-F). Apart from that, we observed

an increase in desmin (DES) insertion into areae compositae upon treatment with erlotinib or

PP2 (Supplementary Figure 3).

Enhanced DP and DSG2 staining in murine cardiac slices after EGFR or SRC inhibition

To confirm the findings from HL-1 cardiomyocytes in a mouse model, we performed

immunostaining of murine cardiac slice cultures, both in $Jup^{+/+}$ and in $Jup^{-/-}$ mice. As

observed before (Yeruva et al., 2020), DSG2 was less expressed in the ICDs of $Jup^{-/-}$ mice

compared to $Jup^{+/+}$ mice (Figure 5A and D). No changes in staining intensity or ICD lengths

between mediator-treated and untreated slice cultures were found in murine cardiac slice

cultures from $Jup^{+/+}$ or $Jup^{-/-}$ mice. Nonetheless, both DP and DSG2 staining were broader

after EGFR or SRC inhibition (Figure 5A-C). Interestingly, in cardiac slices obtained from

$Jup^{-/-}$ mice, only DSG2 staining width was increased after EGFR or SRC inhibition, whereas

changes in DP staining width were not observed (Figure 5D-F). Taken together,

immunostaining showed increased DSG2 and DP localization at cellular junctions after EGFR

or SRC inhibition.

EGFR or SRC inhibition promotes desmosomal assembly in HL-1 cardiomyocytes

Since the localization of DSG2 at cell borders was increased after EGFR or SRC inhibition,

we wondered whether EGFR inhibition alters DSG2 mobility. Therefore, we transfected HL-1

cardiomyocytes with a DSG2-GFP construct and performed fluorescence recovery after

photobleach (FRAP) experiments. However, we found no changes neither in the halftime of

recovery (τ), nor in the immobile fraction after EGFR inhibition (Figures 6A and 6B).

Furthermore, we did not observe significant protein localization changes between cytosolic

and membrane-bound fractions in Triton-X-100 assays (Supplementary Figure 4A and 4B).

Therefore, we next studied desmosomal assembly and performed a Ca^{2+}-switch assay by

depleting cells of Ca^{2+} for 90 minutes, using the Ca^{2+}-chelator EGTA to disrupt desmosomes.

We performed dispase-based dissociation assays after changing the medium to Ca^{2+}-

containing medium together with erlotinib or PP2. EGFR or SRC inhibition led to increased

cellular cohesion after a Ca^{2+}-switch, indicating increased desmosomal assembly (Figure 6C

and Supplementary Figure 4C). Indeed, immunostaining for DP and DSG2 revealed that both

EGFR and SRC inhibition led to enhanced recruitment of DP and DSG2 to the cell borders

after the Ca^{2+}-switch in HL-1 cardiomyocytes (Figure 6D-F). Immunoprecipitation of DP

after EGFR and SRC inhibition revealed more association of PG and DP in HL-1

cardiomyocytes, while PKP2 and DSG2 association with DP was only increased upon SRC

inhibition (Figure 6G). These data confirmed that EGFR and SRC inhibition led to enhanced

desmosomal assembly.

Discussion

AC is a disease with complex pathogenicity, which at least in part is caused by disturbed

desmosome turnover (Liang et al., 2021). The current therapeutic options for AC patients

include changes in lifestyle, treatment with antiarrhythmic drugs, catheter ablation,

implantable cardiac defibrillator, and end-stage heart transplantation (Corrado et al., 2017).

However, in patients carrying mutations in genes coding for proteins of the desmosomal

complex, lifestyle changes, such as restraining from physical endurance activities and β-

blocker therapy are most commonly used (Corrado et al., 2017). In our previous work, we

discovered adrenergic signaling via PKA-mediated PG-phosphorylation at serine 665 as a

driving force for DSG2-mediated enhanced cardiomyocyte cohesion, which we termed

positive adhesiotropy (Schlipp et al., 2014, Schinner et al., 2017, Shoykhet et al., 2020). This

indicated that cardiomyocyte adhesion at ICDs is precisely regulated and can be enhanced via

drugs. Because EGFR inhibitors are used in clinics for the treatment of different kinds of

cancers (Westover et al., 2018, Sabbah et al., 2020), in this work, we characterized the role of

EGFR on cardiomyocyte cohesion and the efficacy of erlotinib to modulate adhesion. Indeed,

we show that inhibition of EGFR and its effector molecule, SRC, in HL-1 cardiomyocytes

and murine cardiac slice cultures from $Jup^{+/+}$ or $Jup^{-/-}$ mice enhanced cardiomyocyte

cohesion, paralleled by increased desmosome assembly into the area composita (Figure 7).

As a first step, we analyzed the expression of EGFR in $Jup^{+/+}$ or $Jup^{-/-}$ mice hearts and found

an increase in EGFR expression in the hearts of $Jup^{-/-}$ mice. To our knowledge, this is the first

instant of finding EGFR overexpression in murine models of AC. In fact, EGFR activation

and signaling was shown to promote TGF-β dependent renal fibrosis (Chen et al., 2012,

Samarakoon et al., 2013). Since in AC, replacement of myocardium with fibrotic tissue is a

common phenomenon (Corrado et al., 2017), we believe that the observed EGFR

overexpression might be related to fibrosis observed in $Jup^{-/-}$ mice (Shoykhet et al., 2020,

Yeruva et al., 2020, Schinner et al., 2017). It has been shown that upon treatment with the

EGFR inhibitor gefitinib, inflammation and hypertrophy caused by chronic treatment with the

adrenergic mediator isoprenaline, was ameliorated in mice (Grisanti et al., 2015, Talarico et

al., 2014). Moreover, increased EGFR expression was positively correlated with the size of

atherosclerotic plaques (Lamb et al., 2004). However, mice with a strong downregulation of

EGFR in the heart, have severe hypertrophy and hypotension (Schreier et al., 2013).

We used erlotinib or PP2 to inhibit EGFR and SRC kinases, respectively, and focused more

on erlotinib, since it is a clinically approved drug. Dispase-based dissociation assays using

erlotinib and PP2 in HL-1 cardiomyocytes and murine cardiac slice cultures from $Jup^{+/+}$ or

$Jup^{-/-}$ mice revealed that both the inhibitors enhance cardiomyocyte cohesion. Similar to our

findings in this study, EGFR inhibition has been shown to increase cellular cohesion in

squamous cell carcinomas (Lorch et al., 2004), whereas in keratinocytes EGFR inhibition

protected against DSG3 autoantibody-induced loss of keratinocyte adhesion (Walter et al.,

2019). Immunoprecipitation revealed EGFR along with PKP2, DP and PG being in a complex

with DSG2. In intestinal epithelial cells, as well as in a cell-free AFM setting, an interaction

of DSG2 and EGFR has been observed (Ungewiss et al., 2018). However, to our knowledge

in cardiomyocytes this complex was not known before. It has been very well demonstrated

that desmosomes in the epithelia or ICDs in the cardiomyocytes not only provide cellular

cohesion but can also be part of a signaling hub (Broussard et al., 2015, Najor et al., 2017,

Muller et al., 2021, Yeruva and Waschke, 2022), thus disrupting EGFR signaling in this

context might lead to changes at the desmosome.

On a single molecule level in an AFM setting, we found that binding events mediated by

DSG2, but not DSC2 and N-CAD were increased after erlotinib treatment in HL-1

cardiomyocytes. Increased binding frequency at the cell borders indicates an increased

binding partner localization towards the cell borders. However, it might also be that active

EGFR is disturbing the interaction possibility of the binding partner with the DSG2-coated

AFM tip, which is restored upon EGFR inhibition. In a cell-free AFM setting DSG2 does not

only interact homophilically with DSG2, but also heterophilically with DSC2 (Fuchs et al.,

2022). In the AFM setting, only interactions with "free" proteins can be measured. Since we

observed an increase of DSG2 at the cell borders upon EGFR inhibition, we suppose that the

increased binding frequency at cell borders, in line with our results from the Ca^{2+}-switch

assays indicate increased desmosomal assembly, where freshly assembled desmosomal

precursors are available for interaction with the opposite cell DSG2 or DSC2. At the same

time, the decrease of DSC2 binding frequency might indicate that either DSC2 or its binding

partners are not available for interactions with the DSC2-coated tips, especially at the cell

surface. Since we did not observe any significant changes in the DSC2 binding frequency at

the cell border, we assumed that the observed decrease in DSC2 binding frequency at cell

surface was not off significant concern.

When we knocked down *Egfr* by siRNA, SRC inhibition was still effective in inducing

positive adhesiotropy, indicating that the SRC inhibition-mediated effect on cardiomyocyte

cohesion is either downstream or at least in part independent of EGFR. However, Western

blot analyses revealed that PP2 also caused a significant decrease in EGFR phosphorylation at

Y845 and Y1068, indicating SRC acts upstream of EGFR. We have previously shown in

keratinocytes that SRC via its target cytoskeletal protein cortactin contributes to the loss of

cell cohesion mediated by DSG3 autoantibodies (Kugelmann et al., 2019). Similarly, in

cardiomyocytes SRC might also have a direct interaction with desmosomal proteins,

independent of the EGFR molecule. Infact, in intestinal epithelial cells DSG2, EGFR and

SRC were found to be in one complex (Ungewiss et al., 2018). Upon knockdown of *DSG2*,

DSC2 has been shown to be upregulated in colonic adenocarcinoma cell lines (Kamekura et

al., 2014), which might explain why we can still observe a positive adhesiotropic effect of

EGFR or SRC inhibition upon *Dsg2* knockdown. On the other hand, the remaining DSG2

upon *Dsg2* knockdown could be sufficient to induce positive adhesiotropy after EGFR

inhibition. Indeed, we never observed a decrease in cell cohesion upon *Dsg2* knockdown

(Schinner et al., 2020, Shoykhet et al., 2020) which further confirms that the partial presence

of DSG2 can provide mechanical stability to the cells.

Further, we observed an increased recruitment of DSG2 and DP to the membranes of HL-1

cardiomyocytes, increasing area composita length. The longer areae compositae in HL-1

cardiomyocytes could be the cellular correlate of increased thickness of DSG2 and DP

staining in murine cardiac slices. A similar reorganization of ICDs has already been observed

in electron microscopy upon adrenergic signaling, which also induces positive adhesiotropy in

cardiomyocytes (Yeruva et al., 2020). We found that the recruitment of DSG2 and DP into the

cellular junctions is not caused by increased mobility, rather by an enhanced desmosomal

assembly, which is supported by increased DES insertions into DP. Interestingly, in $Jup^{-/-}$

mice, we observed changes in DSG2 but not DP localization at the ICD, indicating that the

effect of EGFR and SRC inhibition in intact cardiac tissue is at least in part mediated by PG.

Indeed, it was shown previously that in $Jup^{-/-}$ keratinocytes, steady-state protein levels and

recruitment of desmosome components to the membrane were reduced (Yin et al., 2005).

Mechanistically, PG-phosphorylation at tyrosine residues was shown to be involved in EGFR-

induced loss of cell adhesion, which decreased association of PG to DP (Gaudry et al., 2001),

and triggered translocation of PG and DP from the membrane towards the cytoplasm (Gaudry

et al., 2001, Bektas et al., 2013). Furthermore, EGF-induced phosphorylation of PG could be

prevented by EGFR inhibition (Lorch et al., 2004). In consensus with the above studies, an

increase in co-immunoprecipitation of PG and PKP2 proteins with DP in HL-1

cardiomyocytes after inhibition of EGFR or SRC was observed in this study. These results

further strengthen our hypothesis that desmosomal assembly is increased upon EGFR or SRC

inhibition.

Based on our results, we anticipate that short-term EGFR inhibition can be beneficial in the

setting of AC. Furthermore, unlike other EGFR inhibitors, reports of adverse cardiac effects

after erlotinib treatment are very rare (Kenigsberg et al., 2017, Stuhlmiller et al., 2017, Barish

et al., 2019, Nagashio et al., 2021). Taken together, we conclude that inhibiting EGFR,

thereby enhancing desmosome assembly stabilizes desmosome integrity. However, further

studies, investigating the *in vivo* effect of EGFR inhibition utilizing AC mouse models as well

as the effects on electromechanical coupling are needed, which are beyond the scope of

current study. Nevertheless, we believe that our study is a first step that paves the way for

future studies targeting EGFR inhibition by erlotinib as a treatment option for AC.

Materials and Methods

1.1.Cell culture

HL-1 cells, which are immortalized murine atrial myocyte cells, provided by William

Claycomb (LSU Health Sciences Center, New Orleans, USA), were cultured in supplemented

medium as described before (Shoykhet et al., 2020). Cells were cultured in medium

containing norepinephrine, while for experiments, medium without norepinephrine was used.

1.2.Murine cardiac slice culture

Animal handling was performed in accordance with the guidelines from Directive

2010/63/EU of the European Parliament. Previously characterized 12 week old C57BL6/J

wildtype and cardiomyocyte specific $Jup^{-/-}$ mice were used for experiments (Schinner et al.,

2017, Schinner et al., 2020). In brief, mice with loxP sites flanking exon 1 of the junctional

plakoglobin gene (*Jup*) on both alleles (Juptm1.1Glr/J mice, the Jackson Laboratory, Bar

Harbor, USA) were bred with mice heterozygous for loxP sites in *Jup* with an expression of

the recombinase Cre under the control of cardiac-specific alpha myosin heavy chain promoter

(Myh6), which were generated by crossing with B6.FVB-Tg(Myh6-cre)2182Mds/J mice (The

Jackson Laboratory). Murine cardiac slice cultures were obtained as described before

(Shoykhet et al., 2020) with minor adjustments. In brief, after anesthetizing mice with

isoflurane, mice were sacrificed by cervical dislocation, the hearts were collected, put into

pre-cooled slicing buffer, embedded in low-melt agarose and sliced with a vibratome

(LeicaVT1200S vibratome) into 200 µm sections. For treatments, consecutive slices were

used to minimize slice-to-slice variability. After respective treatments, slices were processed

for lysate preparation, dispase-based dissociation assays and immunostainings. For

immunostainings, slices were embedded in NEG-50™ frozen section medium and stored at -

80°C until further processing.

1.3.Mediators and reagents

Samples were treated with mediators for 90 minutes. Erlotinib (Santa Cruz, #sc-202154) was

used as an EGFR inhibitor at a final concentration of 2.5 µM. PP2 (Merck, #529573) was

used to inhibit the EGFR signaling pathway by inhibiting SRC at a final concentration of 10

μM. When a Ca^{2+}-switch was performed, cells were treated with 5 mM EGTA (VWR, #0732)

for 90 minutes. Then, medium was changed back to Ca^{2+}-containing medium for 90 minutes

and mediators were added at the same time.

1.4.Lysate preparation

For Western blot analyses, lysates were prepared in SDS lysis buffer with protease and

phosphatase inhibitors (cOmplete™ Protease Inhibitor Cocktail, Roche, #CO-RO and

PhosStop™, Roche, #PHOSS-RO). In the case of HL-1 cardiomyocytes, the medium was

removed after treatments and cells were kept on ice, washed with PBS, scraped into SDS lysis

buffer and sonicated. When working with murine cardiac slice cultures, slices were treated,

washed with TBS and snap-frozen in liquid nitrogen. To prepare the tissue lysates, SDS lysis

buffer was added, the samples were transferred into gentleMACS™ M-tubes (Miltenyi

Biotec, #130-093-236) and dissociated using the protein_01_01 program of the

gentleMACS™ OctoDissociator (Miltenyi Biotec, #130-095-937).

For co-immunoprecipitation experiments, cells were cultured in T75 flasks. After treatments,

RIPA lysis buffer (10 mM Na_2HPO_4, 150 mM NaCl, 1% Triton X-100, 0.25% SDS, 1%

sodium deoxycholate, pH=7.2) containing protease and phosphatase inhibitors was added to

the cells and incubated for 30 minutes on ice on a rocking platform. The lysate was then

transferred to gentleMACS™ M-tubes, and dissociated using the protein_01_01 program of

the gentleMACS™ OctoDissociator.

For Triton assays, cells were washed with ice cold PBS and cells were kept on ice on a

rocking platform in Triton buffer (0.5% Triton X-100, 50 mM MES, 25 mM EGTA, 5 mM

$MgCl_2$, pH=6.8) with protease and phosphatase inhibitors for 20 minutes. After that, the

lysate was transferred to Eppendorf tubes and centrifuged at 4°C for 10 minutes at 15000 g in

an Eppendorf 5430R centrifuge. The supernatant containing the Triton-soluble proteins was

transferred to a new reaction tube, while the pellets were washed with Triton buffer, dissolved

in SDS lysis buffer and sonicated.

For all assays, protein concentration was measured using the Pierce™ Protein Assay Kit

(Thermo Fisher, #23225) according to the manufacturer's protocol.

1.5.Western blot analyses

Lysates were denatured in Laemmli buffer and boiled for 10 minutes at 95°C, before being

loaded on SDS-PAGE gels. The PageRuler™ Plus Prestained Protein Ladder (Thermo Fisher,

#26620) was used as a marker. After electrophoresis, proteins were transferred to a

nitrocellulose membrane (Thermo Fisher, #LC2006) using the wet-blot method. Membranes

were then blocked in 5% milk/TBST, in case of anti-phospho-EGFR antibodies in 5%

BSA/TBST, or in case of anti-DP antibodies in ROTI®Block (Carl Roth, #A151.2). For

Western blots with lysates from murine tissue samples, as well as Triton blots, the No-Stain™

Protein Labeling Reagent (Thermo Fisher, #A44449) was used as loading control according to

the manufacturer's protocol. Following primary antibodies were added overnight at 4°C on a

rocking platform: anti-DP (Progen, #61003), anti-DSG1/2 (Progen #61002), anti-phospho-

EGFR Y845 (Cell Signaling #2231), anti-phospho-EGFR Y1068 (Cell Signaling, #2234),

anti-EGFR (sc-373746 (for HL-1 cardiomyocytes) and ab52894 (for murine tissue)), anti-

phospho-ERK1/2 ((E-4) Santa Cruz, #sc-7383), anti-ERK (Cell Signaling, #9102), anti-

phospho-PG (S665, self-made as described before (Schinner et al., 2020)), anti-PG ((PG5.1)

Progen #61005), anti-PKP2 (Progen, #651167) and anti-α-tubulin ((DM1A), abcam,

#ab7291). After washing, membranes were incubated with species-matched, HRP-coupled

secondary antibodies (Dianova, #111-035-045 or #115-035-068) diluted in TBST or 5%

milk/TBST in case of phospho-EGFR antibodies, and developed using an iBright™ FL1500

(Thermo Fisher) developer with SuperSignal™ West Pico PLUS Chemiluminescent Substrate

(Thermo Fisher, #34577). All primary antibodies were used at 1:1000 dilutions, apart from

phospho-EGFR and EGFR antibodies (1:500), phospho-PG (1:20), anti-PKP2 (1:25) and anti-

α-tubulin (1:4000). Quantification was performed using the Image Studio Lite v.5.2 or the

iBright™ Analysis Software (Invitrogen).

1.6.Dissociation Assays

Dissociation assays in HL-1 cardiomyocytes were performed as described before (Shoykhet et

al., 2020). In brief: after treatment, medium was removed and Liberase-DH (Sigma-Aldrich,

#LIBDH-RO) was added, followed by dispase II (Sigma-Aldrich, #D4693). Cells were

incubated at 37°C, 5% CO_2, until the cell monolayers detached from the well bottoms. Then,

the enzyme mix was carefully replaced by HBSS and mechanical stress was applied by

shaking at 1.31 g for 5 minutes on an orbital shaker (Stuart SSM5 orbital shaker). For better

visibility, MTT (Sigma-Aldrich, #M5655) was added to the cells and images were taken to

count the number of fragments. The amount of fragmentation serves as an indirect

measurement of intercellular cohesion, as fewer fragments suggest a stronger cellular

cohesion.

Dissociation assays in murine cardiac slices were performed similarly to dissociation assays

in HL-1 cardiomyocytes. However, in this case, Liberase-DH and dispase II were added

simultaneously for 30 minutes. MTT was added and equal mechanical stress was applied to

all samples of an experiment using an electrical pipette. Then, the content of each well was

filtered using a 70 µm nylon membrane (PluriSelect, #43-10070-60), and transferred to a

gridded 96-well plate. Pictures of the well were taken and stitched together using the

AutoStich software (Brown, 2018). For quantification, only the number of viable dissociated

cardiomyocytes was counted, distinguishable by their rod shape and the MTT staining. To

control variations due to different slice size or location in the ventricle, consecutive slices for

control and treatments were used, and treatments were normalized to the respective control

slice.

1.7.Immunostainings

Cells were seeded on coverslips, and fixed with paraformaldehyde (PFA) after respective

treatments. 7 µm thick sections of murine cardiac slices were cut with a cryostat

(HM500OMV, Microm, Walldorf, Germany), sections were transferred to glass slides and

adhered by warming up to 38°C. After PFA fixation, the samples were permeabilized with

0.1% Triton-X 100 and blocked with bovine serum albumin and normal goat serum

(BSA/NGS). Primary antibodies diluted in PBS were added and incubated overnight at 4°C in

a wet chamber. Following primary antibodies were used: anti-DES (abcam, #ab32362), anti-

DP (Progen, #61003), anti-DSG2 (Progen, #610121) and anti-N-CAD (BD Transduction

Laboratories™, #610921). When working with HL-1 cardiomyocytes, PBS was used for

washing steps, while for murine tissue samples, 50 mM NH_4Cl in PBS was used. After

washing, species-matched, fluorophore-coupled secondary antibodies ((anti-mouse Alexa

Fluor® 488 (Dianova #115-545-003), anti-mouse Cy3 (Dianova 115-165-164) or anti-rabbit

Cy5 (Dianova, 111-175-144)) were added for 60 minutes. In the case of wheat germ

agglutinin (WGA) or Phalloidin staining (Thermo Fisher, #W11261 and Thermo Fisher,

#A12379), the dye was added together with the secondary antibodies. WGA was used at 2

µg/ml and Phalloidin at 130 nM concentrations. During the last 10 minutes of secondary

antibody incubation, DAPI (Sigma Aldrich, #D9542, 0.5 µg/ml) was added. After washing,

coverslips were mounted on microscope slides using NPG (Sigma-Aldrich, #P3130). Slides

were analyzed using a Leica SP5 II confocal microscope (Leica, Mannheim, Germany) with a

63X oil objective with a numerical aperture of 1.4 using Immersol 518F (Zeiss). Images were

acquired at room temperature with the LAS-AF software. Z-scans were performed at 0.25 µm

thickness spanning the whole cell volume. Colocalization analyses were performed as

described before (Shoykhet et al., 2020), in brief: the membrane region was marked as a

region of interest (Supplementary Figure 5A). In this region, the amount of stained pixels in

the other channel, where colocalization was to be quantified, was measured and a ratio of

colocalization at the membrane was calculated. Measurement of staining width in

immunostainings performed in murine cardiac slice cultures, is exemplified in Supplementary

Figure 5B.

For stimulated emission depletion (STED) immunostainings, a slightly modified staining

protocol was used: instead of washing with PBS, cells were washed with 50 mM NH_4Cl in

PBS. Primary antibodies were diluted in BSA/NGS and the following secondary antibodies

were used: STAR RED anti-rabbit (abberior, #STRED-1002) and Alexa Fluor® 594 anti-

mouse (abcam #ab150120), both 1:200 dilution. Slides were mounted with ProLong™

Diamond Antifade (Thermo Scientific, #P36961). Samples were analyzed at room

temperature using a STED expert line microscope from Abberior with a 100X oil objective

with a numerical aperture of 1.4 using IMMOIL-F30CC (Olympus). Images were acquired

with a DMK33G274 camera and the Imspector software. For quantification, areas with a

"railroad-like" staining of DP were considered as area composita. To quantify DES insertions

into the area composita, the number of spots where DES filaments were inserting into the area

composita were counted.

1.8.Immunoprecipitation

For pulldowns, 1 µg of respective antibodies were added to 1.5 mg of protein lysate

overnight, and kept at 4°C on a spinning wheel. To rule out unspecific antibody or bead

binding, IgG control samples were prepared, in parallel. Therefore, the same amount of

protein lysate was used as for pulldowns, however, instead of primary antibody, 1 µg of

normal mouse IgG (Merck, #12-371) was added to the samples. Pulldowns were performed

with Protein G Dynabeads™ (Thermo Fisher, #10004D), by adding the lysate-antibody mix

for 1 hour at 4°C on a spinning wheel to beads prewashed in RIPA buffer with protease and

phosphatase inhibitors. Washing steps were performed using a magnetic rack. Then, beads

were washed twice with wash buffer 1 (50 mM Tris-HCl, 150 mM NaCl, 0.1 mM EDTA,

0.5% Tween20, pH=7.5), thrice with wash buffer 2 (100 mM Tris-HCl, 200 mM NaCl, 2 M

Urea, 0.5% Tween20, pH=7.5), and twice with 1% Triton X-100 in PBS. All wash buffers

were supplemented with protease and phosphatase inhibitors. After the last wash step, beads

were resuspended in Laemmli buffer, denatured for 10 minutes at 95°C, the lysate was

separated from the beads using a magnet and loaded on a SDS-PAGE gel for Western blot

analyses. 20 µg of total lysate was used as input control for the Western blots.

1.9.Atomic Force Microscopy

Atomic Force Microscopy (AFM) measurements were performed as described before

(Schinner et al., 2017, Shoykhet et al., 2020) using a Nanowizard® III AFM (JPK

instruments, Berlin, Germany) with an optical fluorescence microscope (Axio Observer D1,

Carl Zeiss) at 37°C. Cantilevers (MLCT AFM Probes, Bruker, Calle Tecate, CA) were

functionalized with recombinant proteins as described elsewhere (Ebner et al., 2007). The

following recombinant proteins were used: DSC2-Fc, DSG2-Fc (both self-made by Protein A

purification of cell culture supernatants from overexpressing CHO cells, as described in

(Waschke et al., 2005)) or recombinant N-CAD-Fc (R&D Systems, #6626-NC-50). The

canonical pyramidal-shaped D-tip was used. The thermal noise method (Hutter and

Bechhoefer, 1993) was used to determine the spring constants of the tips in order to obtain

correct values for unbinding forces. For experiments, HL-1 cardiomyocytes were seeded on

24 mm glass coverslips. A bright field image of the cells of interest was acquired with a 63X

objective. Afterwards an overview picture of an area of interest was taken in the force-based

quantitative imaging (QI) mode with the following settings: pulling speed: 50 µm/s, setpoint:

0.2 nN, z-length 2.5 µm. Based on the QI image, areas for adhesion measurements were

chosen (Supplementary Figure 5C). Adhesion measurements were performed on cell borders

as well as cell surfaces on an area of 1.5 µm x 5.0 µm using the force mapping mode with the

following settings: relative setpoint: 0.2 nN, z-length: 2.5 µm, extension speed: 5.0 µm/s,

extend and retract delay: 0.1 s. Information of approach and retraction forces was recorded in

force-distance curves (FDCs). The resulting FDCs were analyzed using the JPKSPM Data

processing software. Binding events are visible by a step in the FDCs, representing the force

needed to rupture the binding between the functionalized tip and the binding partner on the

cell. Hence, this force is the unbinding force. Unbinding forces were analyzed using the

extreme fit distribution in Origin 2017 software.

1.10. siRNA-mediated knockdown

For siRNA-mediated knockdown, cells were transfected with a transfection reagent,

comprised of OptiMEM (Thermo Scientific, #31985070), RNAiMAX (Thermo Scientific

#13778150) and the ON-Target siRNA (Dharmacon, siNT #D-001810-10, si*Egfr* L-040411-

00-0005, si*Dsg2* #L-042514-01-005, si*Dsp* #L-040653-01-005) according to manufacturer's

protocol. Cells were transfected for 24 hours. 72 hours post transfection, medium was

changed again. Experiments were performed 84 hours post-transfection. Each knockdown

was confirmed by Western blot.

1.11. Fluorescence Recovery After Photobleach

For Fluorescence Recovery After Photobleach (FRAP) measurements, HL-1 cardiomyocytes

were seeded in ibidi 15 μ-Slide 8 Well Glass Bottom chambers (ibidi, #80827). 24 hours after

seeding, cells were transfected with 1 μg DSG2-GFP construct as described before (Schinner

et al., 2017), using TurboFect (Thermo, #R0531). Images were captured with a Leica SP5

confocal microscope (Leica, Mannheim, Germany) with the LAS-AF software starting 60

minutes after treatment with the vehicle or erlotinib. Using the Leica FRAP Wizard, 5

pictures were taken before bleaching, then an area of interest was bleached with 20 pulses at

100% laser intensity for 0.523 s. After that, 60 pictures were taken every second, 20 pictures

were taken every 3 seconds and 10 pictures were taken every 10 seconds. Halftime of

recovery, as well as immobile fractions, were calculated using the Leica FRAP Wizard.

1.12. Imaging and Statistics

Images were processed with Adobe Photoshop CS5, Image Studio Lite v.5.2 and ImageJ

software. The cartoon was created using BioRender. Statistical analyses were performed using

GraphPad Prism 8. Student's *t*-tests or 1-Way ANOVA with Post-hoc tests were applied after

outlier removal and are explained in figure legends. Data are represented as mean ± standard

deviation. For dissociation assays as well as for quantification of Western blots in HL-1

cardiomyocytes, values were normalized to the average control value of all experiments. For

dissociation assays in murine cardiac slices, the number of dissociated cells was normalized to

the respective control slice of the same mouse. Significance was assumed for p ≤ 0.05.

Online Supplementary Material

Figure S1 shows Western blot confirmations of the erlotinib and PP2 effects on HL-1

cardiomyocytes. Figure S2 shows representative images of the dispase-based dissociation

assay in HL-1 cells after erlotinib or PP2 treatments. Figure S3 shows an increased DES

insertion into DP upon erlotinib or PP2 treatments. Figure S4 shows that EGFR or SRC

inhibition do not cause changes in protein localization between detergent soluble and

insoluble fractions. Figure S5 shows exemplified immunostaining quantifications and the area

choice in the AFM measurements.

Acknowledgements

This work was supported by the Ludwig-Maximilian-University Munich through the FöFoLe

program and the Deutsche Forschungsgemeinschaft (grant WA2474/11-1 to JW). The authors

declare no competing financial interests. We thank Martina Hitzenbichler, Daniela

Kugelmann, Kathleen Plietz, Kilian Skowranek and the AFM tip coating team for their help

and technical assistance.

Author contributions

MS, OD, PM and MH acquired the data. MS and SY analyzed the data and drafted the

manuscript. All authors proof-read the manuscript. JW and SY made critical revision of the

manuscript for important intellectual content, handled supervision and designed the research.

Statements and declarations

Conflict of Interest

The authors declare that no conflict of interest exists.

Ethical Standards

The manuscript does not contain clinical studies or patient data. Animal handling was in

accordance with the guidelines from the Directive 2010/63/EU of the European Parliament

and approved by the regional government of Upper Bavaria (Gz. ROB-55.2-2532.Vet_02-19-

172).

Figure 1 EGFR or SRC inhibition induces positive adhesiotropy in cardiomyocytes

A: Representative Western blot showing protein expression of EGFR and PG in $Jup^{+/+}$ and

$Jup^{-/-}$ mice. **B**: Dispase-based dissociation assay in HL-1 cardiomyocytes upon inhibition of

EGFR or SRC by erlotinib or PP2, respectively, with representative pictures of the wells, * p

$\leq$ 0.05 (p=0.0003 for erlotinib and p<0.0001 for PP2), 1-way ANOVA with Holm-Sidak

correction, N=7 biological replicates. **C**: Dispase-based dissociation assay in murine cardiac

slice cultures obtained from $Jup^{+/+}$ and $Jup^{-/-}$ mice upon inhibition of EGFR or SRC, * p $\leq$

0.05 ($Jup^{+/+}$ mice: p=0.0012 for DMSO vs erlotinib and p=0.0024 for DMSO vs PP2. $Jup^{-/-}$

mice: p=0.0322 for DMSO vs erlotinib and PP2), 1-way ANOVA with Holm-Sidak

correction, N=7 for Jup$^{+/+}$ mice and N=6 for $Jup^{-/-}$ mice. **D**: Maximum projections of

immunostainings in HL-1 cardiomyocytes for N-CAD and DSG2 showing an increase of

DSG2 at the cell borders after erlotinib or PP2 treatments. Z-scans spanning the whole cell

volume, z-steps = 0.25 µm. White arrows indicate areas of increased DSG2 localization at the

cell membrane. Scale bar: 10 µm. **E:** Quantification of colocalization of N-CAD and DSG2 in

HL-1 cardiomyocytes, * p $\leq$ 0.05 (p<0.0001 for DMSO vs erlotinib and p=0.0042 for DMSO

vs PP2), 1-way ANOVA with Holm-Sidak correction. Each data point represents one

individual cell border, N=5 biological replicates. **F:** Representative Western blots for

immunoprecipitation of DSG2, co-immunoprecipitation of DP, EGFR, PKP2 and PG, IgG

heavy chain (IgG hc) served as loading control for immunoprecipitated samples, N=3

biological replicates.

Figure 2 EGFR inhibition leads to increased DSG2 binding frequency at cell borders

A-C: Binding frequency and topography during atomic force microscopy (AFM)

measurements on HL-1 cardiomyocytes before and after erlotinib treatment at or close to cell

borders and cell surfaces with tips coated with **A**: DSG2, **B**: N-CAD, **C**: DSC2.

Measurements were first performed in 90 minutes DMSO-treated samples, then medium was

changed, erlotinib was added and experiment was performed after 90 minutes. Green dots in topography images indicate binding events; cyan dots indicate cell borders. Topography images are 1.5 µm x 5 µm. In the graphs, per data-point, 1500 curves were analysed across two areas (1.5 µm x 5 µm). * p ≤ 0.05. Statistical significance was calculated between DMSO vs erlotinib. (DSG2: cell borders p=0.0426, cell surface: p=0.0785; N-CAD: cell borders p=0.433, cell surface: p=0.1357; DSC2: cell borders p=0.1151, cell surface: p=0.041), unpaired Student's *t*-test, N=3 biological replicates. **D-F**: Unbinding forces measured during AFM measurements on HL-1 cardiomyocytes at cell borders and cell surfaces with tips coated with **D**: DSG2, **E**: N-CAD, **f**: DSC2. Per data-point, 1500 curves were analysed across two areas (1.5 µm x 5 µm). * p ≤ 0.05. Statistical significance was calculated between DMSO vs erlotinib. (DSG2 cell borders 0.9808, cell surface p=0.9722; N-CAD: p=0.5380, cell surface p=0.958; DSC2: p=0.9258, cell surface p=0.4421), unpaired, Student's *t*-test, N=3 biological replicates.

Figure 3 Positive adhesiotropy induced by EGFR or SRC inhibition is dependent on DP

A: Dispase-based dissociation assay in HL-1 cardiomyocytes after siRNA-mediated knockdown of EGFR and non-target (NT) control knockdown. Unpaired Student's *t*-test (p=0.9858), N=6 biological replicates. **B**: Dispase-based dissociation assay in HL-1 cardiomyocytes after siRNA-mediated knockdown of EGFR and treatments with erlotinib or PP2. *p ≤ 0.05 (siNT: p=0.0264 for DMSO vs erlotinib and PP2; siEGFR: p=0.8257 for DMSO vs erlotinib and p=0.0494 for DMSO vs PP2), 1-way ANOVA with Holm-Sidak correction, N=6. **C**: Representative Western blot confirmation of siRNA-mediated knockdown efficiency for EGFR, DP and DSG2, α-tubulin served as a loading control, N=6 biological replicates. **D**: Dispase-based dissociation assay in HL-1 cardiomyocytes after siRNA-mediated knockdown of DSG2 or DP *p ≤ 0.05 (p=0.982 for siNT vs siDSG2 and p=0.2204 for siNT vs siDSP), unpaired Student's *t*-test. **E**: Dispase-based dissociation assay

in HL-1 cardiomyocytes after siRNA-mediated knockdown of DSG2 or DP and treatments with erlotinib or PP2. *p ≤ 0.05 (siNT: p=0.0002 for DMSO vs erlotinib and p= 0.001 for DMSO vs PP2; siDSG2: p=0.0081 for DMSO vs erlotinib and p=0.0097 for DMSO vs PP2; siDSP: p=0.9512 for DMSO vs erlotinib and PP2), 1-way ANOVA with Holm-Sidak correction, N=6-8 biological replicates.

Figure 4 Increased DP and DSG2 recruitment to the cell membranes leads to longer areae compositae upon EGFR or SRC inhibition

A: Immunostaining of DP and DSG2 in HL-1 cardiomyocytes with Phalloidin as membrane marker. Scale bar: 10 µm. White arrows represent an increase in DSG2 and DP localization to the membrane compared to DMSO. **B**: Quantification of colocalization of DP and Phalloidin. *p ≤ 0.05 (p=0.0001 for DMSO vs erlotinib and p<0.0001 for DMSO vs PP2), 1-way ANOVA with Holm-Sidak correction. **C**: Quantification of colocalization of DSG2 and Phalloidin. *p ≤ 0.05 (p=0.0002 for DMSO vs erlotinib and p=0.0122 for DMSO vs PP2), 1-way ANOVA with Holm-Sidak correction. Each data point represents one individual cell border, N=6 biological replicates. **D**: STED images of HL-1 cardiomyocytes stained for DP and DSG2. Scale bar in overviews: 2 µm, scale bar in zooms: 500 nm. **E**: Number of areae compositae over six areas (15 µm x 15 µm) from N=5 biological replicates, *p ≤ 0.05 (p=0.2959 for DMSO vs erlotinib and p=0.6994 for DMSO vs PP2), 1-way ANOVA with Holm-Sidak correction. **F**: Quantification of area composita length. *p ≤ 0.05 (p=0.0078 for DMSO vs erlotinib and p=0.0001 for DMSO vs PP2), 1-way ANOVA with Holm-Sidak correction, N=4 biological replicates.

Figure 5 EGFR or SRC inhibition leads to increased recruitment of DP and DSG2 into the ICDs in $Jup^{+/+}$ and $Jup^{-/-}$ mice

A: Immunostaining of DP and DSG2 in *Jup*^(+/+) murine cardiac slices with WGA as membrane

marker. Scale bar: 8 μm. **B**: DP staining width. **C**: DSG2 staining width. * p ≤ 0.05, (DP

staining width: p<0.0001 for DMSO vs erlotinib and PP2; DSG2 staining width: p<0.0001 for

DMSO vs erlotinib and p=0.0004 for DMSO vs PP2), 1-way ANOVA with Holm-Sidak

correction. Each data point represents one ICD, N=4 mice. **D**: Immunostainings of *Jup*^(-/-)

murine cardiac slices for DP and DSG2 with WGA as membrane marker. Scale bar: 8 μm. **E**:

DP staining width. **F**: DSG2 staining width. * p ≤ 0.05 (DP staining width: p=0.7121 for

DMSO vs erlotinib and p=0.5042 for DMSO vs PP2; DSG2 staining width: p=0.0189 for

DMSO vs erlotinib and p=0.0147 for DMSO vs PP2), 1-way ANOVA with Holm-Sidak

correction. Each data point represents one ICD, N=4 mice. White arrows represent an increase

staining width in DSG2 or DP staining compared to DMSO.

Figure 6 Increased recruitment of DP and DSG2 towards the cell membrane is achieved by

enhanced desmosome assembly

A-B: Fluorescence recovery after photobleach (FRAP) measurements in HL-1

cardiomyocytes transfected with DSG2-GFP and treated with DMSO or erlotinib.

Measurements were performed 60-120 minutes post treatment **A**: Immobile fraction **B**:

Halftime of recovery (τ). * p ≤ 0.05 (immobile fraction p=0.6933; τ p=0.5252 for DMSO vs

erlotinib), unpaired Student's *t*-test, N=4 biological replicates. **C:** Dispase-based dissociation

assay in HL-1 cardiomyocytes after 90 minutes of Ca^{2+}-depletion and subsequent Ca^{2+}-

repletion together with erlotinib or PP2. *p ≤ 0.05 (p=0.0201 for DMSO vs erlotinib and

PP2), 1-way ANOVA with Holm-Sidak correction, N=6 biological replicates. **D:**

Immunostaining of DP and DSG2 in HL-1 cardiomyocytes with Phalloidin as membrane

marker after 90 minutes of Ca^{2+}-depletion and subsequent Ca^{2+}-repletion together with

erlotinib or PP2. White arrows indicate areas of increased DP or DSG2 recruitment to the cell

membrane. Scale bar: 10 μm. **E-F:** Quantification of colocalization of **E**: DP or **F**: DSG2 and

Phalloidin, * p ≤ 0.05 (DP: p=0.0216 for DMSO vs erlotinib and p=0.0036 for DMSO vs

PP2); DSG2: p=0.0073 for DMSO vs erlotinib and p=0.0004 for DMSO vs PP2), 1-way

ANOVA with Holm-Sidak correction, N=4 biological replicates. Each data point represents

one individual cell border. **G:** Immunoprecipitation of DP from HL-1 lysates, co-

immunoprecipitation of EGFR, DSG2, PKP2 and PG, N=3 biological replicates.

Fig. 7 Schematic overview of EGFR or SRC inhibition-mediated positive adhesiotropy

EGFR exists in complex with DSG2, along with DP, PG and PKP2, either at the cell borders

or in the cytoplasm. EGFR or SRC inhibition, via erlotinib or PP2, respectively, led to

positive adhesiotropy in cardiomyocytes by enhancing the desmosome assembly, which

increased area composita length.

References

AKHTAR, S., YOUSIF, M. H., DHAUNSI, G. S., CHANDRASEKHAR, B., AL-FARSI, O. & BENTER, I. F. 2012.
Angiotensin-(1-7) inhibits epidermal growth factor receptor transactivation via a Mas
receptor-dependent pathway. *Br J Pharmacol,* 165, 1390-400.
BARISH, R., GATES, E. & BARAC, A. 2019. Trastuzumab-Induced Cardiomyopathy. *Cardiol Clin,* 37,
407-418.
BEKTAS, M., JOLLY, P. S., BERKOWITZ, P., AMAGAI, M. & RUBENSTEIN, D. S. 2013. A pathophysiologic
role for epidermal growth factor receptor in pemphigus acantholysis. *J Biol Chem,* 288, 9447-
56.
BELMADANI, S., PALEN, D. I., GONZALEZ-VILLALOBOS, R. A., BOULARES, H. A. & MATROUGUI, K. 2008.
Elevated epidermal growth factor receptor phosphorylation induces resistance artery
dysfunction in diabetic db/db mice. *Diabetes,* 57, 1629-37.
BENTER, I. F., BENBOUBETRA, M., HOLLINS, A. J., YOUSIF, M. H., CANATAN, H. & AKHTAR, S. 2009.
Early inhibition of EGFR signaling prevents diabetes-induced up-regulation of multiple gene
pathways in the mesenteric vasculature. *Vascul Pharmacol,* 51, 236-45.
BHARGAVA, R., GERALD, W. L., LI, A. R., PAN, Q., LAL, P., LADANYI, M. & CHEN, B. 2005. EGFR gene
amplification in breast cancer: correlation with epidermal growth factor receptor mRNA and
protein expression and HER-2 status and absence of EGFR-activating mutations. *Mod Pathol,*
18, 1027-33.
BROUSSARD, J. A., GETSIOS, S. & GREEN, K. J. 2015. Desmosome regulation and signaling in disease.
Cell Tissue Res, 360, 501-12.
BROWN, M. 2018. *Autostitch Software* [Online]. Available:
http://matthewalunbrown.com/autostitch/autostitch.html [Accessed].
CAMPBELL, P., MORTON, P. E., TAKEICHI, T., SALAM, A., ROBERTS, N., PROUDFOOT, L. E., MELLERIO,
J. E., AMINU, K., WELLINGTON, C., PATIL, S. N., AKIYAMA, M., LIU, L., MCMILLAN, J. R.,
ARISTODEMOU, S., ISHIDA-YAMAMOTO, A., ABDUL-WAHAB, A., PETROF, G., FONG, K.,
HARNCHOOWONG, S., STONE, K. L., HARPER, J. I., IRWIN MCLEAN, W. H., SIMPSON, M. A.,
PARSONS, M. & MCGRATH, J. A. 2014. Epithelial inflammation resulting from an inherited
loss-of-function mutation in EGFR. *J Invest Dermatol,* 134, 2570-2578.
CHEN, J., CHEN, J. K., NAGAI, K., PLIETH, D., TAN, M., LEE, T. C., THREADGILL, D. W., NEILSON, E. G. &
HARRIS, R. C. 2012. EGFR signaling promotes TGFbeta-dependent renal fibrosis. *J Am Soc*
Nephrol, 23, 215-24.
CHEN, Z., OH, D., DUBEY, A. K., YAO, M., YANG, B., GROVES, J. T. & SHEETZ, M. 2018. EGFR family and
Src family kinase interactions: mechanics matters? *Curr Opin Cell Biol,* 51, 97-102.
CHOI, S. K., GALAN, M., PARTYKA, M., TREBAK, M., BELMADANI, S. & MATROUGUI, K. 2012. Chronic
inhibition of epidermal growth factor receptor tyrosine kinase and extracellular signal-
regulated kinases 1 and 2 (ERK1/2) augments vascular response to limb ischemia in type 2
diabetic mice. *Am J Pathol,* 180, 410-8.
CORRADO, D., BASSO, C. & JUDGE, D. P. 2017. Arrhythmogenic Cardiomyopathy. *Circ Res,* 121, 784-
802.
DA CUNHA SANTOS, G., SHEPHERD, F. A. & TSAO, M. S. 2011. EGFR mutations and lung cancer. *Annu*
Rev Pathol, 6, 49-69.
EBNER, A., WILDLING, L., KAMRUZZAHAN, A. S., RANKL, C., WRUSS, J., HAHN, C. D., HOLZL, M., ZHU,
R., KIENBERGER, F., BLAAS, D., HINTERDORFER, P. & GRUBER, H. J. 2007. A new, simple
method for linking of antibodies to atomic force microscopy tips. *Bioconjug Chem,* 18, 1176-
84.
EL-HASHIM, A. Z., KHAJAH, M. A., RENNO, W. M., BABYSON, R. S., UDDIN, M., BENTER, I. F.,
EZEAMUZIE, C. & AKHTAR, S. 2017. Src-dependent EGFR transactivation regulates lung
inflammation via downstream signaling involving ERK1/2, PI3Kdelta/Akt and NFkappaB
induction in a murine asthma model. *Sci Rep,* 7, 9919.

FORRESTER, S. J., KAWAI, T., O'BRIEN, S., THOMAS, W., HARRIS, R. C. & EGUCHI, S. 2016. Epidermal
Growth Factor Receptor Transactivation: Mechanisms, Pathophysiology, and Potential
Therapies in the Cardiovascular System. *Annu Rev Pharmacol Toxicol,* 56**,** 627-53.
FUCHS, M., KUGELMANN, D., SCHLEGEL, N., VIELMUTH, F. & WASCHKE, J. 2022. Desmoglein 2 can
undergo Ca(2+)-dependent interactions with both desmosomal and classical cadherins
including E-cadherin and N-cadherin. *Biophys J,* 121**,** 1322-1335.
GAUDRY, C. A., PALKA, H. L., DUSEK, R. L., HUEN, A. C., KHANDEKAR, M. J., HUDSON, L. G. & GREEN,
K. J. 2001. Tyrosine-phosphorylated plakoglobin is associated with desmogleins but not
desmoplakin after epidermal growth factor receptor activation. *J Biol Chem,* 276**,** 24871-80.
GENET, G., GUILBEAU-FRUGIER, C., HONTON, B., DAGUE, E., SCHNEIDER, M. D., COATRIEUX, C.,
CALISE, D., CARDIN, C., NIETO, C., PAYRE, B., DUBROCA, C., MARCK, P., HEYMES, C., DUBRAC,
A., ARVANITIS, D., DESPAS, F., ALTIE, M. F., SEGUELAS, M. H., DELISLE, M. B., DAVY, A.,
SENARD, J. M., PATHAK, A. & GALES, C. 2012. Ephrin-B1 is a novel specific component of the
lateral membrane of the cardiomyocyte and is essential for the stability of cardiac tissue
architecture cohesion. *Circ Res,* 110**,** 688-700.
GERULL, B. & BRODEHL, A. 2020. Genetic Animal Models for Arrhythmogenic Cardiomyopathy. *Front*
Physiol, 11**,** 624.
GRISANTI, L. A., REPAS, A. A., TALARICO, J. A., GOLD, J. I., CARTER, R. L., KOCH, W. J. & TILLEY, D. G.
2015. Temporal and gefitinib-sensitive regulation of cardiac cytokine expression via chronic
beta-adrenergic receptor stimulation. *Am J Physiol Heart Circ Physiol,* 308**,** H316-30.
GUO, S., OKYERE, A. D., MCEACHERN, E., STRONG, J. L., CARTER, R. L., PATWA, V. C., THOMAS, T. P.,
LANDY, M., SONG, J., LUCCHESE, A. M., MARTIN, T. G., GAO, E., RAJAN, S., KIRK, J. A., KOCH,
W. J., CHEUNG, J. Y. & TILLEY, D. G. 2021. Epidermal growth factor receptor-dependent
maintenance of cardiac contractility. *Cardiovasc Res.*
HARSKAMP, L. R., GANSEVOORT, R. T., VAN GOOR, H. & MEIJER, E. 2016. The epidermal growth
factor receptor pathway in chronic kidney diseases. *Nat Rev Nephrol,* 12**,** 496-506.
HELISTE, J., JOKILAMMI, A., VAPARANTA, K., PAATERO, I. & ELENIUS, K. 2021. Combined genetic and
chemical screens indicate protective potential for EGFR inhibition to cardiomyocytes under
hypoxia. *Sci Rep,* 11**,** 16661.
HUTTER, J. L. & BECHHOEFER, J. 1993. Calibration of atomic-force microscope tips. *Review of*
Scientific Instruments, 64**,** 1868-1873.
IWAMOTO, M., SASO, W., SUGIYAMA, R., ISHII, K., OHKI, M., NAGAMORI, S., SUZUKI, R., AIZAKI, H.,
RYO, A., YUN, J. H., PARK, S. Y., OHTANI, N., MURAMATSU, M., IWAMI, S., TANAKA, Y.,
SUREAU, C., WAKITA, T. & WATASHI, K. 2019. Epidermal growth factor receptor is a host-
entry cofactor triggering hepatitis B virus internalization. *Proc Natl Acad Sci U S A,* 116**,** 8487-
8492.
KAMEKURA, R., KOLEGRAFF, K. N., NAVA, P., HILGARTH, R. S., FENG, M., PARKOS, C. A. & NUSRAT, A.
2014. Loss of the desmosomal cadherin desmoglein-2 suppresses colon cancer cell
proliferation through EGFR signaling. *Oncogene,* 33**,** 4531-6.
KENIGSBERG, B., JAIN, V. & BARAC, A. 2017. Cardio-oncology Related to Heart Failure: Epidermal
Growth Factor Receptor Target-Based Therapy. *Heart Fail Clin,* 13**,** 297-309.
KLEINSCHMIDT-DEMASTERS, B. K., LILLEHEI, K. O. & VARELLA-GARCIA, M. 2005. Glioblastomas in the
older old. *Arch Pathol Lab Med,* 129**,** 624-31.
KLESSNER, J. L., DESAI, B. V., AMARGO, E. V., GETSIOS, S. & GREEN, K. J. 2009. EGFR and ADAMs
cooperate to regulate shedding and endocytic trafficking of the desmosomal cadherin
desmoglein 2. *Mol Biol Cell,* 20**,** 328-37.
KUGELMANN, D., ROTZER, V., WALTER, E., EGU, D. T., FUCHS, M. T., VIELMUTH, F., VARGAS-ROBLES,
H., SCHNOOR, M., HERTL, M., EMING, R., ROTTNER, K., SCHMIDT, A., SPINDLER, V. &
WASCHKE, J. 2019. Role of Src and Cortactin in Pemphigus Skin Blistering. *Front Immunol,* 10**,**
626.
LAMB, D. J., MODJTAHEDI, H., PLANT, N. J. & FERNS, G. A. 2004. EGF mediates monocyte chemotaxis
and macrophage proliferation and EGF receptor is expressed in atherosclerotic plaques.
Atherosclerosis, 176**,** 21-6.

LIANG, Y., LYON, R. C., PELLMAN, J., BRADFORD, W. H., LANGE, S., BOGOMOLOVAS, J., DALTON, N. D.,
GU, Y., BOBAR, M., LEE, M. H., IWAKUMA, T., NIGAM, V., ASIMAKI, A., SCHEINMAN, M.,
PETERSON, K. L. & SHEIKH, F. 2021. Desmosomal COP9 regulates proteome degradation in
arrhythmogenic right ventricular dysplasia/cardiomyopathy. *J Clin Invest,* 131.
LORCH, J. H., KLESSNER, J., PARK, J. K., GETSIOS, S., WU, Y. L., STACK, M. S. & GREEN, K. J. 2004.
Epidermal growth factor receptor inhibition promotes desmosome assembly and strengthens
intercellular adhesion in squamous cell carcinoma cells. *J Biol Chem,* 279**,** 37191-200.
MAKKI, N., THIEL, K. W. & MILLER, F. J., JR. 2013. The epidermal growth factor receptor and its
ligands in cardiovascular disease. *Int J Mol Sci,* 14**,** 20597-613.
MATROUGUI, K. 2010. Diabetes and microvascular pathophysiology: role of epidermal growth factor
receptor tyrosine kinase. *Diabetes Metab Res Rev,* 26**,** 13-6.
MULLER, L., HATZFELD, M. & KEIL, R. 2021. Desmosomes as Signaling Hubs in the Regulation of Cell
Behavior. *Front Cell Dev Biol,* 9**,** 745670.
NAGASHIO, K., TAJIRI, K., SATO, K. & IEDA, M. 2021. Erlotinib-Induced Cardiomyopathy in a Patient
with Metastatic Non-Small Cell Lung Cancer. *Int Heart J,* 62**,** 1171-1175.
NAJOR, N. A., FITZ, G. N., KOETSIER, J. L., GODSEL, L. M., ALBRECHT, L. V., HARMON, R. & GREEN, K. J.
2017. Epidermal Growth Factor Receptor neddylation is regulated by a desmosomal-COP9
(Constitutive Photomorphogenesis 9) signalosome complex. *Elife,* 6.
NOMA, T., LEMAIRE, A., NAGA PRASAD, S. V., BARKI-HARRINGTON, L., TILLEY, D. G., CHEN, J., LE
CORVOISIER, P., VIOLIN, J. D., WEI, H., LEFKOWITZ, R. J. & ROCKMAN, H. A. 2007. Beta-
arrestin-mediated beta1-adrenergic receptor transactivation of the EGFR confers
cardioprotection. *J Clin Invest,* 117**,** 2445-58.
PANG, H., ROWAN, B. G., AL-DHAHERI, M. & FABER, L. E. 2004. Epidermal growth factor suppresses
induction by progestin of the adhesion protein desmoplakin in T47D breast cancer cells.
Breast Cancer Res, 6**,** R239-45.
SABBAH, D. A., HAJJO, R. & SWEIDAN, K. 2020. Review on Epidermal Growth Factor Receptor (EGFR)
Structure, Signaling Pathways, Interactions, and Recent Updates of EGFR Inhibitors. *Curr Top*
Med Chem, 20**,** 815-834.
SAMARAKOON, R., DOBBERFUHL, A. D., COOLEY, C., OVERSTREET, J. M., PATEL, S., GOLDSCHMEDING,
R., MELDRUM, K. K. & HIGGINS, P. J. 2013. Induction of renal fibrotic genes by TGF-beta1
requires EGFR activation, p53 and reactive oxygen species. *Cell Signal,* 25**,** 2198-209.
SCHINNER, C., OLIVARES-FLOREZ, S., SCHLIPP, A., TRENZ, S., FEINENDEGEN, M., FLASWINKEL, H.,
KEMPF, E., EGU, D. T., YERUVA, S. & WASCHKE, J. 2020. The inotropic agent digitoxin
strengthens desmosomal adhesion in cardiac myocytes in an ERK1/2-dependent manner.
Basic Res Cardiol, 115**,** 46.
SCHINNER, C., VIELMUTH, F., ROTZER, V., HIERMAIER, M., RADEVA, M. Y., CO, T. K., HARTLIEB, E.,
SCHMIDT, A., IMHOF, A., MESSOUDI, A., HORN, A., SCHLIPP, A., SPINDLER, V. & WASCHKE, J.
2017. Adrenergic Signaling Strengthens Cardiac Myocyte Cohesion. *Circ Res,* 120**,** 1305-1317.
SCHLIPP, A., SCHINNER, C., SPINDLER, V., VIELMUTH, F., GEHMLICH, K., SYRRIS, P., MCKENNA, W. J.,
DENDORFER, A., HARTLIEB, E. & WASCHKE, J. 2014. Desmoglein-2 interaction is crucial for
cardiomyocyte cohesion and function. *Cardiovasc Res,* 104**,** 245-57.
SCHMITT, T. & WASCHKE, J. 2021. Autoantibody-Specific Signalling in Pemphigus. *Front Med*
(Lausanne), 8**,** 701809.
SCHREIER, B., RABE, S., SCHNEIDER, B., BRETSCHNEIDER, M., RUPP, S., RUHS, S., NEUMANN, J.,
RUECKSCHLOSS, U., SIBILIA, M., GOTTHARDT, M., GROSSMANN, C. & GEKLE, M. 2013. Loss of
epidermal growth factor receptor in vascular smooth muscle cells and cardiomyocytes causes
arterial hypotension and cardiac hypertrophy. *Hypertension,* 61**,** 333-40.
SHIA, J., KLIMSTRA, D. S., LI, A. R., QIN, J., SALTZ, L., TERUYA-FELDSTEIN, J., AKRAM, M., CHUNG, K. Y.,
YAO, D., PATY, P. B., GERALD, W. & CHEN, B. 2005. Epidermal growth factor receptor
expression and gene amplification in colorectal carcinoma: an immunohistochemical and
chromogenic in situ hybridization study. *Mod Pathol,* 18**,** 1350-6.

SHOYKHET, M., TRENZ, S., KEMPF, E., WILLIAMS, T., GERULL, B., SCHINNER, C., YERUVA, S. &
841   WASCHKE, J. 2020. Cardiomyocyte adhesion and hyperadhesion differentially require ERK1/2
842   and plakoglobin. *JCI Insight,* 5.
843 SHRAIM, B. A., MOURSI, M. O., BENTER, I. F., HABIB, A. M. & AKHTAR, S. 2021. The Role of Epidermal
844   Growth Factor Receptor Family of Receptor Tyrosine Kinases in Mediating Diabetes-Induced
845   Cardiovascular Complications. *Front Pharmacol,* 12, 701390.
846 STROEDECKE, K., MEINEL, S., MARKWARDT, F., KLOECKNER, U., STRAETZ, N., QUARCH, K., SCHREIER,
847   B., KOPF, M., GEKLE, M. & GROSSMANN, C. 2021. The mineralocorticoid receptor leads to
848   increased expression of EGFR and T-type calcium channels that support HL-1 cell
849   hypertrophy. *Sci Rep,* 11, 13229.
850 STUHLMILLER, T. J., ZAWISTOWSKI, J. S., CHEN, X., SCIAKY, N., ANGUS, S. P., HICKS, S. T., PARRY, T. L.,
851   HUANG, W., BEAK, J. Y., WILLIS, M. S., JOHNSON, G. L. & JENSEN, B. C. 2017. Kinome and
852   Transcriptome Profiling Reveal Broad and Distinct Activities of Erlotinib, Sunitinib, and
853   Sorafenib in the Mouse Heart and Suggest Cardiotoxicity From Combined Signal Transducer
854   and Activator of Transcription and Epidermal Growth Factor Receptor Inhibition. *J Am Heart*
855   *Assoc,* 6.
856 TALARICO, J. A., CARTER, R. L., GRISANTI, L. A., YU, J. E., REPAS, A. A. & TILLEY, D. G. 2014. beta-
857   adrenergic receptor-dependent alterations in murine cardiac transcript expression are
858   differentially regulated by gefitinib in vivo. *PLoS One,* 9, e99195.
859 UNGEWISS, H., ROTZER, V., MEIR, M., FEY, C., DIEFENBACHER, M., SCHLEGEL, N. & WASCHKE, J. 2018.
860   Dsg2 via Src-mediated transactivation shapes EGFR signaling towards cell adhesion. *Cell Mol*
861   *Life Sci,* 75**,** 4251-4268.
862 WALTER, E., VIELMUTH, F., WANUSKE, M. T., SEIFERT, M., POLLMANN, R., EMING, R. & WASCHKE, J.
863   2019. Role of Dsg1- and Dsg3-Mediated Signaling in Pemphigus Autoantibody-Induced Loss
864   of Keratinocyte Cohesion. *Front Immunol,* 10, 1128.
865 WASCHKE, J. 2019. Desmogleins as signaling hubs regulating cell cohesion and tissue/organ function
866   in skin and heart - EFEM lecture 2018. *Ann Anat,* 226**,** 96-100.
867 WASCHKE, J., BRUGGEMAN, P., BAUMGARTNER, W., ZILLIKENS, D. & DRENCKHAHN, D. 2005.
868   Pemphigus foliaceus IgG causes dissociation of desmoglein 1-containing junctions without
869   blocking desmoglein 1 transinteraction. *J Clin Invest,* 115, 3157-65.
870 WESTOVER, D., ZUGAZAGOITIA, J., CHO, B. C., LOVLY, C. M. & PAZ-ARES, L. 2018. Mechanisms of
871   acquired resistance to first- and second-generation EGFR tyrosine kinase inhibitors. *Ann*
872   *Oncol,* 29**,** i10-i19.
873 WILDE, A., BEATTIE, E. C., LEM, L., RIETHOF, D. A., LIU, S. H., MOBLEY, W. C., SORIANO, P. & BRODSKY,
874   F. M. 1999. EGF receptor signaling stimulates SRC kinase phosphorylation of clathrin,
875   influencing clathrin redistribution and EGF uptake. *Cell,* 96**,** 677-87.
876 YERUVA, S., KEMPF, E., EGU, D. T., FLASWINKEL, H., KUGELMANN, D. & WASCHKE, J. 2020. Adrenergic
877   Signaling-Induced Ultrastructural Strengthening of Intercalated Discs via Plakoglobin Is
878   Crucial for Positive Adhesiotropy in Murine Cardiomyocytes. *Front Physiol,* 11**,** 430.
879 YERUVA, S. P. & WASCHKE, J. 2022. Structure and regulation of desmosomes in intercalated discs:
880   Lessons from epithelia. *J Anat.*
881 YIN, T., GETSIOS, S., CALDELARI, R., GODSEL, L. M., KOWALCZYK, A. P., MULLER, E. J. & GREEN, K. J.
882   2005. Mechanisms of plakoglobin-dependent adhesion: desmosome-specific functions in
883   assembly and regulation by epidermal growth factor receptor. *J Biol Chem,* 280**,** 40355-63.
884 ZHAO, G., QIU, Y., ZHANG, H. M. & YANG, D. 2019. Intercalated discs: cellular adhesion and signaling
885   in heart health and diseases. *Heart Fail Rev,* 24**,** 115-132.